KREIN'S STRINGS WHOSE SPECTRAL FUNCTIONS ARE OF POLYNOMIAL GROWTH

S.KOTANI

ABSTRACT. In the case of Krein's strings with spectral functions of polynomial growth a necessary and sufficient condition for the Krein's correspondence to be continuous is given.

1. INTRODUCTION

Let \mathcal{M} be the totality of non-decreasing, right continuous functions on $[0,\infty)$ satisfying

$$m(0-) = 0, \quad m(x) \le \infty,$$

and set

$$\left\{ \begin{array}{l} l = \inf \left\{ x \ge 0; \ m(x) = \infty \right\}, \\ a = \inf \left\{ x \ge 0; \ m(x) > 0 \right\}. \end{array} \right.$$

For $m \in \mathcal{M}$ denote $\varphi_{\lambda}(x), \psi_{\lambda}(x)$ the solutions to

$$\begin{cases} \varphi_{\lambda}(x) = 1 - \lambda \int_{0}^{x} (x - y) \varphi_{\lambda}(y) dm(y), \\ \psi_{\lambda}(x) = x - \lambda \int_{0}^{x} (x - y) \psi_{\lambda}(y) dm(y), \end{cases}$$

and define

$$h(\lambda) = \lim_{x \to l} \frac{\psi_{\lambda}(x)}{\varphi_{\lambda}(x)} = \int_{0}^{l} \varphi_{\lambda}(x)^{-2} dx.$$

Then it is known that there exists a unique measure σ on $[0,\infty)$ satisfying

$$h(\lambda) = a + \int_0^\infty \frac{1}{\xi - \lambda} d\sigma(\xi)$$

and conversely, h determines m uniquely. Conventionally it is understood that for $m \in \mathcal{M}$ taking ∞ identically on $[0, \infty)$ the h vanishing identically corresponds, and for $m \in \mathcal{M}$ vanishing identically on $[0, \infty)$ the h taking identically ∞ corresponds. This is the theorem obtained by Krein[8] and m is called Krein's (regular) string. Later Kasahara[1] established the continuity for the correspondence and applied it to show limit theorems for 1D diffusion processes with m their speed measures. Recently Kotani[7] extended Kasahara's result to a certain kind of singular strings m, namely to m which is a non-decreasing and right continuous function on $(-\infty, \infty)$ satisfying

$$m(-\infty) = 0, \quad m(x) \le \infty,$$

and

(1.1)
$$\int_{-\infty}^{a} x^2 dm(x) < \infty$$

²⁰⁰⁰ Mathematics Subject Classification. Primary 34L05, 34B20; Secondary 60J60, 60J55.

Key words and phrases. generalized diffusion process, Krein's correspondence, inverse spectral problem.

for some a. When the condition (1.1) is satisfied, the boundary $-\infty$ is called as the limit circle type for the associated generalized second order differential operator $d^2/dmdx$. In this case he introduced a new h by

$$h(\lambda) = \lim_{x \to -\infty} \left(x + \varphi_{\lambda}(x) \int_{x}^{l} \frac{dy}{\varphi_{\lambda}(y)^{2}} \right) = a + \int_{0}^{\infty} \left(\frac{1}{\xi - \lambda} - \frac{\xi}{\xi^{2} + 1} \right) d\sigma\left(\xi\right),$$

which satisfies

$$h'(\lambda) = \int_{-\infty}^{l} \frac{\partial}{\partial \lambda} \varphi_{\lambda}(x)^{-2} dx,$$

and proved the continuity of the correspondence between m and h. Probabilistic applications of this result were given by Kasahara-Watanabe[2, 3] and it was interpreted from the point of view of the excursion theory by Yano[9]. In this article we consider m satisfying a milder condition than (1.1), namely

$$\int_{-\infty}^{a} |x| \, dm(x) < \infty,$$

and obtain the continuity result under additional conditions on m, which allows any power growth of the spectral measures at ∞ .

2. Preliminaries

Let m(x) be a non-decreasing and right continuous function on $(-\infty, \infty)$ satisfying

$$m(-\infty) = 0, \quad m(\infty) \le \infty.$$

Set

$$l = \sup \{x > -\infty, m(x) < +\infty\}, \quad l_+ = \sup \operatorname{supp} dm, \quad l_- = \inf \operatorname{supp} dm.$$

Note $m(l) = \infty$ if $l < \infty$. Assume

(2.1)
$$\int_{-\infty}^{a} |x| \, dm(x) < \infty$$

with some $a \in (l_-, l_+)$. Let \mathcal{E} be the totality of non-decreasing functions m satisfying (2.1). We exclude m vanishing identically on $(-\infty, \infty)$ from \mathcal{E} . One can regard dm as a distribution of weight and in this case m works as a string. On the other hand, one can associate a generalized diffusion process with generator L

$$L = \frac{d}{dm} \frac{d}{dx}$$

if we impose a suitable boundary condition if necessary. The condition (2.1) is called as entrance condition in 1D diffusion theory developed by W.Feller, so we call *m* satisfying (2.1) a string of entrance type. For an entrance type *m*, it is easy to show that for $\lambda \in \mathbf{C}$ an integral equation

$$\varphi(x) = 1 - \lambda \int_{-\infty}^{x} (x - y)\varphi(y)dm(y)$$

has a unique solution, which is denoted by $\varphi_{\lambda}(x)$. Introduce a subspace

$$L_0^2(dm) = \left\{ f \in L^2(dm); \text{ supp} f \subset (-\infty, l) \right\},\$$

and for $f \in L^2_0(dm)$ define a generalized Fourier transform by

$$\widehat{f}(\lambda) = \int_{-\infty}^{l} f(x)\varphi_{\lambda}(x)dm(x).$$

Krein's spectral theory implies there exists a measure σ on $[0,\infty)$ satisfying

(2.2)
$$\int_{-\infty}^{l} |f(x)|^2 dm(x) = \int_{0}^{\infty} \left| \widehat{f}(\xi) \right|^2 d\sigma(\xi) \text{ for any } f \in L^2_0(dm)$$

 σ is called a spectral measure for the string m. The non-uniqueness of such σ occurs if and only if

$$(2.3) l_+ + m(l_+) < \infty$$

The number $l (\geq l_+)$ possesses its meaning only when (2.3) is satisfied, and in this case there exists a σ satisfying (2.2) with the boundary condition

$$f(l_{+}) + (l - l_{+}) f^{+}(l_{+}) = 0$$

at l_+ . Here f^+ is the derivative from the right hand side. If $l = \infty$, this should be interpreted as

$$f^+\left(l_+\right) = 0.$$

At the left boundary l_{-} no boundary condition is necessary if $l_{-} = -\infty$, and if $l_{-} > -\infty$ we impose the reflective boundary condition, namely

$$f^{-}(l_{-}) = 0$$
 the derivative from left.

Generally, for a string m of entrance type it is known that for $\lambda < 0$ there exists uniquely f such that

$$\begin{cases} -Lf = \lambda f, \quad f > 0, \quad f^+ \le 0, \quad f(l-) = 0\\ f(x)\varphi_{\lambda}^+(x) - f^+(x)\varphi_{\lambda}(x) = 1 \end{cases}$$

This unique f is denoted by f_{λ} and contains information of the boundary condition we are imposing on -L at the right boundary l_+ , and f_{λ} can be represented by φ_{λ} as

(2.4)
$$f_{\lambda}(x) = \varphi_{\lambda}(x) \int_{x}^{l} \frac{dy}{\varphi_{\lambda}(y)^{2}}$$

The right side integral is always convergent for $\lambda < 0$, because if $\operatorname{supp} dm \neq \phi$, then choosing $a \in \operatorname{supp} dm$, we see for x > a

$$\varphi_{\lambda}(x) \ge 1 - \lambda \int_{-\infty}^{x} (x - y) \, dm(y) \ge 1 - \lambda \int_{-\infty}^{a} (x - y) \, dm(y) \ge 1 - \lambda \, (x - a) \, m(a),$$

hence

$$\int_{x}^{l} \frac{dy}{\varphi_{\lambda}(y)^{2}} \leq \int_{x}^{l} \frac{dy}{\left(1 - \lambda \left(y - a\right) m(a)\right)^{2}} < \infty$$
 for $x > a$. If $\operatorname{supp} dm = \phi$, then $l < \infty$ and

(2.5)
$$m(x) = \begin{cases} 0 & \text{for } x < l \\ \infty & \text{for } x > l \end{cases}$$

which implies

$$\varphi_{\lambda}(x) = \begin{cases} 1 & \text{for } x < l \\ \infty & \text{for } x > l \end{cases}$$

and

$$\int_{x}^{l} \frac{dy}{\varphi_{\lambda}(y)^{2}} = l - x < \infty.$$

Here note that we have excluded m = 0 identically on $(-\infty, \infty)$, hence $l < \infty$. If m is a non-decreasing function of (2.5) the spectral measure vanishes identically on $[0, \infty)$. If m is ∞ identically on $(-\infty, \infty)$, the spectral function σ is defined to be 0 identically on $[0, \infty)$. Conversely if a spectral measure vanishes identically on $[0, \infty)$, then the associated string m should be of (2.5). $\varphi_{\lambda}(x)$ is an entire function of minimal exponential type as a function of λ and the zeroes of $\varphi_{\lambda}(x)$ coincide

S.KOTANI

with the eigenvalues of -L defined as a self-adjoint operator on $L^2(dm, (-\infty, x])$ with the Dirichlet boundary condition at x, which means that $\varphi_{\lambda}(x)$ has simple zeroes on $(0, \infty)$. The Green function g_{λ} for -L on $L^2(dm)$ is given by

$$g_{\lambda}(x,y) = g_{\lambda}(y,x) = f_{\lambda}(y)\varphi_{\lambda}(x)$$

for $x \leq y$. The relationship between σ and g_{λ} is described by an identity

$$\int_{-\infty}^{l} \int_{-\infty}^{l} g_{\lambda}(x,y) f(x) \overline{f(y)} dm(x) dm(y) = \int_{0}^{\infty} \frac{\left|\widehat{f}(\xi)\right|^{2}}{\xi - \lambda} \sigma(d\xi)$$

for any $f \in L^2(dm)$, and

$$g_{\lambda}(x,y) = \int_{0}^{\infty} \frac{\varphi_{\xi}(x)\varphi_{\xi}(y)}{\xi - \lambda} d\sigma\left(\xi\right),$$

through which σ is determined uniquely from the string *m*. Distinct *m*s may give an equal σ , namely for $a \in \mathbf{R}$ a new string

$$m_a\left(x\right) = m\left(x+a\right)$$

defines the same σ , because

$$\varphi_{\lambda}^{a}(x) = \varphi_{\lambda}(x+a), \quad f_{\lambda}^{a}(x) = f_{\lambda}(x+a),$$

hence

$$g_{\lambda}^{a}(x,x) = \varphi_{\lambda}^{a}(x)f_{\lambda}^{a}(x) = g_{\lambda}(x+a,x+a) = \int_{0}^{\infty} \frac{\varphi_{\xi}(x+a)^{2}}{\xi-\lambda} d\sigma\left(\xi\right).$$

On the other hand

$$g_{\lambda}^{a}(x,x) = \int_{0}^{\infty} \frac{\varphi_{\xi}^{a}(x)\varphi_{\xi}^{a}(x)}{\xi - \lambda} d\sigma_{a}\left(\xi\right) = \int_{0}^{\infty} \frac{\varphi_{\xi}(x+a)^{2}}{\xi - \lambda} d\sigma_{a}\left(\xi\right),$$

hence an identity

$$_{a}\left(\xi\right) = \sigma\left(\xi\right)$$

 σ

should be held. Conversely we have

Theorem 1. (Kotani[5, 6]) If two strings m_1 and m_2 of \mathcal{E} have the same spectral measure σ , then $m_1(x+c) = m_2(x)$ for a $c \in \mathbf{R}$.

If we hope to obtain the continuity of the correspondence between m and σ , we have to keep the non-uniqueness in mind. Namely, for m of \mathcal{E} a sequence $\{m_n\}_{n\geq 1}$ of \mathcal{E} defined by

$$m_n\left(x\right) = m\left(x-n\right)$$

converges to the trivial function 0 as $n \to \infty$. However the associated σ s are independent of n. Therefore we shall give several alternative definitions of convergence by imposing certain extra conditions (related to tightness) in addition to pointwise convergence. Set

$$M(x) = \int_{-\infty}^{x} (x - y) dm(y) = \int_{-\infty}^{x} m(y) dy.$$

Then, the condition (2.1) is equivalent to

 $M(x) < \infty$

for x < l. Using a convention

$$[-\infty, a) = (-\infty, a), (a, \infty] = (a, \infty)$$
 and so on,

we can see M is a non-decreasing convex function on $(-\infty, \infty)$ satisfying

$$\begin{cases} M(x) = 0 \text{ on } (-\infty, l_{-}],\\ \text{continuous and strictly increasing on } [l_{-}, l),\\ M(x) = \infty \text{ on } (l, \infty). \end{cases}$$

For a fixed positive number c, we assume

(2.6)
$$0 \in (l_{-}, l] \text{ and } M(l) \ge c,$$

and normalize such an m by

$$(2.7) M(0) = c$$

Denote by $\mathcal{E}^{(c)}$ the set of all elements of \mathcal{E} satisfying (2.6), (2.7) and set

$$\mathcal{E}_+ = \bigcup_{c>0} \mathcal{E}^{(c)}.$$

In this definition of \mathcal{E}_+ among functions satisfying (2.1) any function m defined by (2.5) for some $l \leq \infty$ is excluded from \mathcal{E}_+ . Therefore, $\mathcal{E} \setminus \mathcal{E}_+$ consists of m satisfying (2.5) for some $l < \infty$. The uniqueness of the correspondence between m and σ holds under this normalization. Set

 \mathcal{S} = the set of all spectral measures for strings of \mathcal{E} .

Any suitable characterization of S is not known yet, however, any measure on $[0, \infty)$ with polynomial growth at ∞ belongs to S.

We prepare a basic estimate for φ_{λ} . φ_{λ} can be represented as

(2.8)
$$\varphi_{\lambda}(x) = \sum_{n=0}^{\infty} (-\lambda)^n \phi_n(x),$$

where $\{\phi_n\}_{n>0}$ are

$$\phi_n(x) = \int_{-\infty}^x (x - y) \phi_{n-1}(y) dm(y), \quad \phi_0(x) = 1.$$

Then, the convergence of the above series can be seen by

Lemma 1. φ_{λ} is given by an absolute convergent series (2.8) and satisfies

$$\varphi_{\lambda}(x) \leq \exp\left(\left|\lambda\right| M(x)\right).$$

Proof. First we show for any $k \ge 0$

(2.9)
$$\phi_k(x) \le \frac{M(x)^k}{k!}$$

holds. Observe

$$\phi_1(x) = \int_{-\infty}^x (x - y) \, dm(y) = M(x).$$

Assuming (2.9) for some k, we have

$$\begin{split} \phi_{k+1}(x) &\leq \frac{1}{k!} \int_{-\infty}^{x} (x-y) \, M(y)^{k} dm(y) \\ &= \frac{1}{k!} \int_{-\infty}^{x} \left(M(y) - k \, (x-y) \, M'(y) \right) M(y)^{k-1} M'(y) dy \\ &\leq \frac{1}{k!} \int_{-\infty}^{x} M'(y) M(y)^{k} dy = \frac{M(x)^{k+1}}{(k+1)!}, \end{split}$$

which proves (2.9) for general k. Then the estimate of φ_{λ} is clear.

Here we clarify the convergence of a sequence of monotone functions taking value ∞ . For non-negative and non-decreasing function m which may take ∞ , set

$$\widehat{m}(x) = \frac{2}{\pi} \tan^{-1} m(x), \quad x \in \mathbf{R}.$$

Then

$$\widehat{m}\left(x\right)\in\left[0,1\right]$$

and right continuous non-decreasing function satisfying

$$0 \le \widehat{m} \left(-\infty \right) \le \widehat{m} \left(x \right) \le \widehat{m} \left(l - \right) \le \widehat{m} \left(l \right) = 1,$$

if $l < \infty$. A sequence of non-negative and non-decreasing functions m_n is defined to converge to m as $n \to \infty$ if

$$(2.10)\qquad \qquad \widehat{m}_n\left(x\right) \to \widehat{m}\left(x\right)$$

holds at any point of continuity of $\widehat{m}(x)$.

Lemma 2. Suppose $m_n \in \mathcal{E}$ converges to $m \in \mathcal{E}$ as $n \to \infty$. Then it holds that

$$\lim_{n \to \infty} l_n \ge l.$$

Proof. Let x < l be a point of continuity for \hat{m} . Then

$$\widehat{m}_n\left(x\right) \to \widehat{m}\left(x\right) < 1,$$

hence

$$\widehat{m}_n\left(x\right) < 1$$

for every sufficiently large n, which implies $x < l_n$ and completes the proof. \Box

The continuity of the correspondence from \mathcal{E} to \mathcal{S} is not hard to show. Let m_n, m be strings of \mathcal{E} and define the convergence of m_n to m by

(A) $m_n(x) \to m(x)$ for every point of continuity of m. (B) $\lim_{x \to -\infty} \sup_{n \ge 1} M_n(x) = 0$

Theorem 2. Suppose $m_n \in \mathcal{E}$ converge to $m \in \mathcal{E}$. Then, for every $\lambda < 0$ the Green functions $g_{\lambda}^{(n)}(x, y)$ of the string m_n converge to the Green function $g_{\lambda}(x, y)$ of m for any x, y < l. In particular the spectral functions $\sigma_n(\xi)$ converge to $\sigma(\xi)$ at every point of continuity of σ .

Proof. Under the conditions it is easy to see that the φ -functions $\varphi_{\lambda}^{(n)}(x)$ of m_n converge to the φ -function $\varphi_{\lambda}(x)$ of m compact uniformly with respect to $(x, \lambda) \in (-\infty, l) \times C$ from the uniform bound for $\varphi_{\lambda}^{(n)}$ due to Lemma 1.. Moreover, if m(a) > 0 at some a, a point of continuity of m, then there exists a positive constant C such that

$$\varphi_{\lambda}^{(n)}(y) \ge 1 - \lambda M_n(y) \ge 1 + C(y - a)$$

holds for any y > a, hence

$$f_{\lambda}^{(n)}(x) = \varphi_{\lambda}^{(n)}(x) \int_{x}^{l_{n}} \frac{1}{\varphi_{\lambda}^{(n)}(y)^{2}} dy$$

also converge to $f_{\lambda}(x)$. If supp $m = \phi$, namely m(x) = 0 identically on $(-\infty, l)$, then

$$f_{\lambda}^{(n)}(x) \to l - x$$

if $l < \infty$. The case $l = \infty$ is excluded. Consequently, we have

$$g_{\lambda}^{(n)}(x,y) = \varphi_{\lambda}^{(n)}(y) f_{\lambda}^{(n)}(x) \to \varphi_{\lambda}(y) f_{\lambda}(x) = g_{\lambda}(x,y)$$

for any $y \leq x < l$. The identity

$$g_{\lambda}^{(n)}(x,y) = \int_{0}^{\infty} \frac{\varphi_{\xi}^{(n)}(x)\varphi_{\xi}^{(n)}(y)}{\xi - \lambda} d\sigma_{n}\left(\xi\right)$$

shows the last statement of the theorem.

3. Scales and estimates by trace

The straight converse statement of the theorem 2 is hopeless to be true, because there is no characterization for a measure σ on $[0, \infty)$ to be a spectral measure of a string $m \in \mathcal{E}$. Therefore we prove the converse continuity of the correspondence by imposing a condition on $\{\sigma_n\}$. In the process of the proof we have to estimate $\varphi_{\lambda}(x)^{-2}$ in terms of m. A better way to investigate $\varphi_{\lambda}(x)^{-2}$ is to use probabilistic methods. Recall that for each fixed a, $\varphi_{\lambda}(a)$ has simple zeroes $\{\mu_n\}_{n\geq 1}$ which are eigenvalues of -L on $(-\infty, a]$ with Dirichlet boundary condition at x = a. Since the Green function for this operator is

$$a - (x \lor y)$$

we see

(3.1)
$$\sum_{n=1}^{\infty} \mu_n^{-1} = \operatorname{tr} (-L)^{-1} = \int_{-\infty}^a (a-x) \, dm(x) = M(a) < \infty.$$

Choosing a b < l, we denote by $\phi_{\lambda}^{b}(\psi_{\lambda}^{b})$ the solutions of

4

$$-\frac{d}{dm}\frac{d}{dx}f = \lambda f, \text{ with } f(b) = 1, \ f'(b) = 0 \ (f(b) = 0, \ f'(b) = 1) \text{ respectively.}$$

Then we see an identity

$$\varphi_{\lambda}(x) = \varphi_{\lambda}(b)\phi_{\lambda}^{b}(x) + \varphi_{\lambda}'(b)\psi_{\lambda}^{b}(x)$$

holds. Lemma 1 implies $\varphi_{\lambda}(b)$ and $\varphi'_{\lambda}(b)$ are entire functions of at most exponential type M(b) as functions of λ . On the other hand, $\phi^b_{\lambda}(x)$ and $\psi^b_{\lambda}(x)$ are entire functions of order at most 1/2 as functions of λ . Therefore we know that $\varphi_{\lambda}(x)$ is an entire function of minimal exponential type, which combined with (3.1) shows

$$\varphi_{\lambda}(a) = \prod_{n=1}^{\infty} \left(1 - \frac{\lambda}{\mu_n}\right)$$

For the detail refer to page 441 of [5]. Now let $\{X_n\}_{n\geq 1}$ be independent random variables each of which has an exponential distribution of mean 1. Then, an identity

$$E \exp\left(\lambda \sum_{n=1}^{\infty} \mu_n^{-1} X_n\right) = \prod_{n=1}^{\infty} \left(1 - \frac{\lambda}{\mu_n}\right)^{-1} = \varphi_{\lambda}(a)^{-1}$$

holds. Therefore, letting $\left\{\widetilde{X}_n\right\}_{n\geq 1}$ be independent copies of $\{X_n\}_{n\geq 1}$ and setting

$$Y_n = X_n + \widetilde{X}_n, \quad X = \sum_{n=1}^{\infty} \mu_n^{-1} Y_n,$$

we have

(3.2)
$$\varphi_{\lambda}(a)^{-2} = E \exp(\lambda X).$$

We denote X = X(a) if necessary, because the eigenvalues $\{\mu_n\}_{n \ge 1}$ depends on the boundary a.

Lemma 3. Suppose the spectral measure σ of an $m \in \mathcal{E}$ satisfies

$$p(t) = \int_0^\infty e^{-t\xi} d\sigma\left(\xi\right) < \infty \quad \text{for any } t > 0$$

Then, for any non-negative Borel measurable function f on $[0,\infty)$

(3.3)
$$\int_{-\infty}^{l} Ef(X(x)) dx = \int_{0}^{\infty} p(t)f(t) dt$$

holds by permitting for the integrals to take the value ∞ simultaneously.

Proof. From (2.4) it follows that for any x < l and $\lambda < 0$

$$\int_{x}^{l} Ee^{\lambda X(y)} dy = Ee^{\lambda X(x)} \int_{0}^{\infty} \frac{\varphi_{\xi}(x)^{2}}{\xi - \lambda} d\sigma\left(\xi\right) = \int_{0}^{\infty} Ee^{\lambda(X(x)+t)} p(t, x, x) dt,$$

holds, where p(t, x, y) is the transition probability density defined by

$$p(t, x, y) = \int_0^\infty e^{-t\xi} \varphi_{\xi}(x) \varphi_{\xi}(y) \, d\sigma(\xi) \,,$$

hence a functional monotone class theorem shows that the identity below holds for any non-negative bounded continuous function f on $[0, \infty)$.

(3.4)
$$\int_{x}^{l} E\left(f\left(X\left(y\right)\right)e^{\lambda X\left(y\right)}\right) dy = \int_{0}^{\infty} E\left(f\left(X\left(x\right)+t\right)e^{\lambda\left(X\left(x\right)+t\right)}\right)p(t,x,x)dt$$
Since for $t > 2M(x)$

Since, for t > 2M(x)

$$p(t, x, x) = \int_0^\infty e^{-\xi t} \varphi_{\xi}(x)^2 \, d\sigma(\xi) \le \int_0^\infty e^{-\xi t} e^{2\xi M(x)} \, d\sigma(\xi) = p(t - 2M(x))$$

holds, assuming f(t) = 0 for $t < \epsilon$, we see

$$\int_{-\infty}^{l} E\left(f\left(X\left(y\right)\right)e^{\lambda X\left(y\right)}\right) dy = \int_{0}^{\infty} f\left(t\right)e^{\lambda t}p(t)dt$$

by letting $x \to -\infty$. Here we have used the fact

$$X(x) \to 0$$
 as $x \to -\infty$.

The rest of the proof is a routine.

Now we define a scale function ϕ on [0,1] as a function satisfying the following properties.

(S.1) ϕ is strictly increasing, convex and $\phi(0) = 0, \phi'(1-) < \infty$.

(S.2) For each x > 0

$$\overline{\lim_{y \downarrow 0}} \frac{\phi(xy)}{\phi(y)} < \infty.$$

(S.3) For each $x \in (0, 1]$

$$\underline{\lim}_{y \downarrow 0} \frac{\phi\left(xy\right)}{\phi\left(y\right)} > 0$$

The property (S.1) enables us to extend ϕ linearly to $[1, \infty)$, namely

$$\phi(x) = \phi(1) + \phi'(1-)(x-1)$$

for x > 1. Then ϕ becomes non-negative, convex and non-decreasing function on $[0, \infty)$. Throughout the paper ϕ is always extended to $[1, \infty)$ linearly in this way. A regularly varying function at 0 satisfies the condition (S.2), (S.3). Set

$$C_{+}(x) = \sup_{y>0} \frac{\phi(xy)}{\phi(y)} < \infty, \quad C_{-}(x) = \inf_{y \in (0,1]} \frac{\phi(xy)}{\phi(y)} > 0.$$

Then C_+ becomes non-negative, convex and non-decreasing on $[0, \infty)$. It satisfies the submultiplicative property

$$C_+(xy) \le C_+(x)C_+(y)$$

for any x, y > 0, hence, setting

$$\alpha_{+} = \sup_{x>1} \frac{\log C_{+}\left(e^{x}\right)}{x} \in [0,\infty)$$

we see

$$C_+(x) \le x^{\alpha_+}$$

8

holds for any $x \ge e$. α_+ should be not less than 1 due to the convexity of C_+ . Since

$$\phi\left(xy\right) \le C_{+}(x)\phi\left(y\right)$$

holds for any x, y > 0, we have

$$\phi(x) \ge \frac{\phi(1)}{C_+(1/x)} \ge \phi(1) x^{\alpha_+}$$

for any $x \in [0, 1/e]$. Therefore the property (S.2) restricts ϕ not to decay faster than with a power order.

(3.5)
$$\phi(xy) \le C_+(x)\phi(y)$$

 C_{-} satisfies

$$C_{-}(xy) \ge C_{-}(x)C_{-}(y)$$

for any $x, y \in [0, 1]$. Typical examples for functions satisfying (S.1)~(S.3) are

$$\phi(x) = x^{\alpha}, \quad x^{\alpha} \left(c - \log x \right),$$

where $\alpha \geq 1$ and c is a sufficiently large positive constant.

Lemma 4. Let $\{Y_n\}_{n\geq 1}$ be a sequence of identically distributed non-negative random variables with mean μ . and $\{\lambda_n\}_{n\geq 1}$ be a non-negative sequence satisfying

$$\sum_{n=1}^{\infty} \lambda_n < \infty,$$

and set

$$X = \sum_{n=1}^{\infty} \lambda_n Y_n$$

Then, we have (1) If ϕ satisfies (S.1), then

$$\phi\left(EX\right) \leq E\phi\left(X\right).$$

(2) If ϕ satisfies (S.1), (S.2), then

$$E\phi(X) \le \left(EC_+\left(\frac{Y_1}{\mu}\right)\right)\phi(EX)$$

Proof. Jensen's inequality implies the inequality in (1). To show the second inequality we set

$$m_n = m^{-1}\lambda_n, \quad m = \sum_{k=1}^{\infty}\lambda_k.$$

Then (3.5) implies

$$\phi(X) = \phi\left(m\mu\sum_{n=1}^{\infty}m_n\frac{Y_n}{\mu}\right) \le \phi(m\mu)C_+\left(\sum_{n=1}^{\infty}m_n\frac{Y_n}{\mu}\right)$$

Since the function C_+ is convex, we have

$$C_+\left(\sum_{n=1}^{\infty} m_n \frac{Y_n}{\mu}\right) \le \sum_{n=1}^{\infty} m_n C_+\left(\frac{Y_n}{\mu}\right),$$

and

$$E\phi(X) \le \phi(m\mu) \sum_{n=1}^{\infty} m_n EC_+\left(\frac{Y_n}{\mu}\right) = \phi(EX) EC_+\left(\frac{Y_1}{\mu}\right).$$

Let X be the same as was defined in (3.2) and set

$$C_{\phi} = EC_{+}\left(\frac{Y_{1}}{\mu}\right) = \int_{0}^{\infty} te^{-t}C_{+}(t/2) dt < \infty.$$

Lemma 5. We have

(1) If ϕ satisfies (S.1), then

$$E\left(\phi\left(X\right)e^{\lambda X}\right) \geq \varphi_{\lambda}\left(a\right)^{-2}\phi\left(\int_{-\infty}^{a}\varphi_{\lambda}\left(x\right)^{2}dm\left(x\right)\int_{x}^{a}\varphi_{\lambda}\left(y\right)^{-2}dy\right).$$

(2) If ϕ satisfies (S.1), (S.2), then

$$E\left(\phi\left(X\right)e^{\lambda X}\right) \leq C_{\phi}\varphi_{\lambda}\left(a\right)^{-2}\phi\left(\int_{-\infty}^{a}\varphi_{\lambda}\left(x\right)^{2}dm\left(x\right)\int_{x}^{a}\varphi_{\lambda}\left(y\right)^{-2}dy\right).$$

Proof. For a fixed $\lambda < 0$ let Z be a non-negative random variable satisfying

$$Ee^{\mu Z} = \varphi_{\lambda+\mu} \left(a\right)^{-2} \varphi_{\lambda} \left(a\right)^{2} = \prod_{n=1}^{\infty} \left(1 - \frac{\mu}{\mu_{n} - \lambda}\right)^{-2}$$

Then, note an identity

(3.6)
$$E\left(\phi\left(X\right)e^{\lambda X}\right) = \varphi_{\lambda}\left(a\right)^{-2}E\left(\phi\left(Z\right)\right),$$

which can be shown from the observation

$$\frac{\partial^{k}}{\partial\lambda^{k}}\varphi_{\lambda}\left(a\right)^{-2} = \varphi_{\lambda}\left(a\right)^{-2} \left.\frac{\partial^{k}}{\partial\mu^{k}}\left(\varphi_{\lambda+\mu}\left(a\right)^{-2}\varphi_{\lambda}\left(a\right)^{2}\right)\right|_{\mu=0}$$

for any $k \ge 0$, because this implies the identity when $\phi(x) = x^k$. To apply Lemma 4 to Z we need to compute EZ. If we denote the Green operator for L on $(-\infty, a]$ with Dirichlet boundary condition at a by G_{λ} , then

$$G_{\lambda}(x,y) = \varphi_{\lambda}(y) \varphi_{\lambda}(x) \int_{x}^{a} \varphi_{\lambda}(z)^{-2} dz \quad \text{for } x \ge y$$

hence

$$EZ = \sum_{j=1}^{\infty} \frac{1}{\mu_j - \lambda} = \operatorname{tr} G_{\lambda} = \int_{-\infty}^{a} \varphi_{\lambda} (x)^2 dm (x) \int_{x}^{a} \varphi_{\lambda} (y)^{-2} dy$$

holds, and we have the inequalities in the statement.

The right side of the inequalities in Lemma 5 can be estimated further.

Lemma 6. For $\lambda < 0$ the following inequalities are valid. (1) $\int_{-\infty}^{a} \varphi_{\lambda}(x)^{2} dm(x) \int_{x}^{a} \varphi_{\lambda}(y)^{-2} dy \ge M(a)\varphi_{\lambda}(a)^{-2}$ (2) $\int_{-\infty}^{a} \varphi_{\lambda}(x)^{2} dm(x) \int_{x}^{a} \varphi_{\lambda}(y)^{-2} dy \le M(a) \wedge \left(\frac{\log \varphi_{\lambda}(a)}{-\lambda}\right)$

Proof. The inequality (1) and the first inequality of (2) follow from the monotonicity of $\varphi_{\lambda}(z)$, namely we have

$$\int_{x}^{a} \varphi_{\lambda}(y)^{-2} dy \ge \varphi_{\lambda}(a)^{-2} (a-x), \quad \int_{x}^{a} \varphi_{\lambda}(y)^{-2} dy \le \varphi_{\lambda}(x)^{-2} (a-x),$$

which implies

$$\int_{-\infty}^{a} \varphi_{\lambda}(x)^{2} dm(x) \int_{x}^{a} \varphi_{\lambda}(y)^{-2} dy \ge \varphi_{\lambda}(a)^{-2} \int_{-\infty}^{a} (a-x) dm(x) = \varphi_{\lambda}(a)^{-2} M(a),$$

and
$$\int_{-\infty}^{a} \varphi_{\lambda}(x)^{2} dm(x) \int_{x}^{a} \varphi_{\lambda}(y)^{-2} dy \le \int_{-\infty}^{a} (a-x) dm(x) = M(a).$$

The second inequality of (2) follows by using the equation satisfied by $\varphi_{\lambda}(x)$

$$d\varphi_{\lambda}'(y) = -\lambda\varphi_{\lambda}(y) \, dm(y) \, ,$$

which yields

$$-\lambda \int_{-\infty}^{a} \varphi_{\lambda}(y)^{2} dm(y) \int_{y}^{a} \varphi_{\lambda}(z)^{-2} dz$$

$$= \int_{-\infty}^{a} \varphi_{\lambda}(y) d\varphi_{\lambda}'(y) \int_{y}^{a} \varphi_{\lambda}(z)^{-2} dz$$

$$= \varphi_{\lambda}(y) \varphi_{\lambda}'(y) \int_{y}^{a} \varphi_{\lambda}(z)^{-2} dz \Big|_{-\infty}^{a} - \int_{-\infty}^{a} \varphi_{\lambda}'(y)^{2} dy \int_{y}^{a} \varphi_{\lambda}(z)^{-2} dz + \int_{-\infty}^{a} \frac{\varphi_{\lambda}'(y)}{\varphi_{\lambda}(y)} dy.$$

Noting

$$\varphi_{\lambda}(y) \varphi_{\lambda}'(y) \int_{y}^{a} \varphi_{\lambda}(z)^{-2} dz \underset{y \to -\infty}{\sim} -\lambda m(y) (a-y) \underset{y \to -\infty}{\to} 0,$$

we see

$$-\lambda \int_{-\infty}^{a} \varphi_{\lambda}(y)^{2} dm(y) \int_{y}^{a} \varphi_{\lambda}(z)^{-2} dz = \log \varphi_{\lambda}(a) - \int_{-\infty}^{a} \varphi_{\lambda}'(y)^{2} dy \int_{y}^{a} \varphi_{\lambda}(z)^{-2} dz$$
$$\leq \log \varphi_{\lambda}(a),$$

which completes the proof.

As the last lemma in this section we have

Lemma 7. The following two estimates hold. (1) For any function ϕ satisfying (S.1) it holds that

$$\int_{0}^{\infty} p(t)\phi(t) e^{\lambda t} dt \ge \int_{-\infty}^{l} \phi\left(M(x)\varphi_{\lambda}(x)^{-2}\right)\varphi_{\lambda}(x)^{-2} dx$$

(2) For any function ϕ satisfying (S.1), (S.2) it holds that

$$\int_{0}^{\infty} p(t)\phi(t) e^{\lambda t} dt \leq C_{\phi} \int_{-\infty}^{l} \phi\left(M(x) \wedge \left(\frac{\log \varphi_{\lambda}(x)}{-\lambda}\right)\right) \varphi_{\lambda}(x)^{-2} dx$$
$$\leq C_{\phi} \int_{-\infty}^{a} \phi(M(x)) \varphi_{\lambda}(x)^{-2} dx + C_{\phi} \frac{-\lambda}{\varphi_{\lambda}'(a)} \int_{0}^{\infty} \phi(t) e^{\lambda t} dt,$$

Proof. All we have to show is an estimate of the integral

$$\int_{a}^{l} \phi\left(\frac{\log \varphi_{\lambda}\left(x\right)}{-\lambda}\right) \varphi_{\lambda}\left(x\right)^{-2} dx.$$

Noting the monotonicity of $\varphi_{\lambda}(x)$, $\varphi'_{\lambda}(x)$ and $\varphi_{\lambda}(x) \ge 1$, we see

$$\begin{split} &\int_{a}^{l} \phi\left(\frac{\log\varphi_{\lambda}\left(x\right)}{-\lambda}\right)\varphi_{\lambda}\left(x\right)^{-2} dx \\ &= \int_{\varphi_{\lambda}\left(a\right)}^{\varphi_{\lambda}\left(l\right)} \phi\left(\frac{\log z}{-\lambda}\right) \frac{1}{z^{2}\varphi_{\lambda}'\left(\varphi_{\lambda}^{-1}\left(z\right)\right)} dz \\ &\leq \frac{1}{\varphi_{\lambda}'\left(\varphi_{\lambda}^{-1}\left(\varphi_{\lambda}\left(a\right)\right)\right)} \int_{\varphi_{\lambda}\left(a\right)}^{\varphi_{\lambda}\left(l\right)} \phi\left(\frac{\log z}{-\lambda}\right) \frac{dz}{z^{2}} \leq \frac{-\lambda}{\varphi_{\lambda}'\left(a\right)} \int_{0}^{\infty} \phi\left(t\right) e^{\lambda t} dt. \end{split}$$

S.KOTANI

4. Continuity of the correspondence from ${\mathcal S}$ to ${\mathcal E}$

In this section we give a partial converse of Theorem 2. The lemma below will be useful later.

Lemma 8. Let
$$m_n \in \mathcal{E}$$
 and σ_n be its spectral function. Suppose

(4.1)
$$\lim_{n \to \infty} M_n(l_n) = 0$$

holds. Then $\sigma_n(\xi) \to 0$ for any $\xi > 0$.

Proof. Since
$$M_n(l_n) < \infty$$
, we have $l_n < \infty$. Set $\widetilde{m}_n(x) = m_n(x+l_n)$. Then
(4.2) $\widetilde{M}_n(0) \to 0$

and its spectral measure coincides with σ_n . Since the condition (4.2) implies

$$\widetilde{m}_n(x) \to \begin{cases} 0 & \text{for } x < 0 \\ \infty & \text{for } x > 0 \end{cases}$$

in \mathcal{E} , Theorem 2 shows $\sigma_n \to 0$.

Theorem 3. Let $m_n \in \mathcal{E}$ and σ_n be its spectral function satisfying

$$p_n(t) = \int_0^\infty e^{-t\xi} d\sigma_n\left(\xi\right) < \infty \quad \text{for any } t > 0$$

and

(4.3)
$$\sup_{n \ge 1} \int_0^1 p_n(t)\phi(t)dt < \infty$$

for a function ϕ satisfying (S.1). Assume there exists a non-trivial measure σ on $[0,\infty)$ satisfying

$$\sigma_{n}\left(\xi\right) \to \sigma\left(\xi\right)$$

at every point of continuity of σ . Then

$$\lim_{n \to \infty} p_n(t) = p(t), \quad \underline{\lim}_{n \to \infty} M_n(l_n) > 0$$

hold. Choose c such that

$$0 < c < \lim_{n \to \infty} M_n\left(l_n\right)$$

and define a_n by the solution $M_n(a_n) = c$. Then there exists a unique $m \in \mathcal{E}^{(c)}$ with spectral measure σ and it holds that $m_n(\cdot + a_n) \to m$ in \mathcal{E} , hence $\sigma \in \mathcal{S}$.

Proof. For any $\epsilon > 0$ and any N > 0

$$\int_0^{\epsilon} p_n(t)\phi(t)dt = \int_0^{\epsilon} \phi(t)dt \int_0^{\infty} e^{-t\xi} d\sigma_n\left(\xi\right) \ge e^{\epsilon N} \int_N^{\infty} e^{-2\epsilon\xi} d\sigma_n\left(\xi\right) \int_0^{\epsilon} \phi(t)dt$$

holds, and the condition (4.3) implies that there exists a constant C_{ϵ} such that

$$\int_{N}^{\infty} e^{-2\epsilon\xi} d\sigma_n\left(\xi\right) \le e^{-\epsilon N} \frac{\int_{0}^{\epsilon} p_n(t)\phi(t)dt}{\int_{0}^{\epsilon} \phi(t)dt} \le e^{-\epsilon N} C_{\epsilon}$$

is valid for any n, N, which yields

(4.4)
$$p_n(t) \to p(t) = \int_0^\infty e^{-t\xi} d\sigma\left(\xi\right)$$

as $n \to \infty$. Applying (3.3) to $\phi(t) e^{\lambda t}$ for $\lambda < 0$ shows

$$\int_{-\infty}^{l_n} E\left(\phi\left(X_n\left(x\right)\right)e^{\lambda X_n\left(x\right)}\right) dx = \int_0^\infty p_n(t)e^{\lambda t}\phi(t)dt$$
$$\leq \int_0^1 p_n(t)e^{\lambda t}\phi(t)dt + p_n(1)\int_1^\infty e^{\lambda t}\phi(t)dt \leq C$$

with a constant C, where $X_n(x)$ is defined by the eigenvalues $\{\mu_j(x)\}_{j\geq 1}$ corresponding to m_n . Since we assume the limiting spectral measure σ is non-trivial, Lemma 8 shows

$$\underline{\lim_{n \to \infty}} M_n \left(l_n \right) > 0.$$

For c such that

$$0 < c < \underline{\lim}_{n \to \infty} M_n\left(l_n\right)$$

define a_n by the solution $M_n(a_n) = c$ and set

$$\widetilde{m}_n\left(x\right) = m_n\left(x + a_n\right).$$

Then $\widetilde{m}_n \in \mathcal{E}^{(c)}$ and an inequality (1) of Lemma 7 implies

$$\int_{-\infty}^{l_n} E\left(\phi\left(X_n\left(x\right)\right)e^{\lambda X_n\left(x\right)}\right) dx \ge \int_{-\infty}^{\widetilde{l}_n} \widetilde{\varphi}_{\lambda}^{(n)}\left(x\right)^{-2} \phi\left(\widetilde{M}_n\left(x\right)\widetilde{\varphi}_{\lambda}^{(n)}\left(x\right)^{-2}\right) dx,$$

and from Lemma 1 we have

$$\int_{-\infty}^{\widetilde{l}_n} \widetilde{\varphi}_{\lambda}^{(n)}(x)^{-2} \phi\left(\widetilde{M}_n(x) \, \widetilde{\varphi}_{\lambda}^{(n)}(x)^{-2}\right) dx \ge \int_{-\infty}^{l_n} e^{2\lambda \widetilde{M}_n(x)} \phi\left(\widetilde{M}_n(x) \, e^{2\lambda \widetilde{M}_n(x)}\right) dx$$
$$\ge \int_{-\infty}^0 e^{2\lambda c} \phi\left(\widetilde{M}_n(x) \, e^{2\lambda c}\right) dx.$$

Thus

$$\int_{-\infty}^{0} e^{2\lambda c} \phi\left(\widetilde{M}_{n}\left(x\right) e^{2\lambda c}\right) dx \leq C$$

is valid for any $n \ge 1$. Therefore, for any x < 0

$$e^{2\lambda c}\left(-x\right)\phi\left(\widetilde{M}_{n}\left(x\right)e^{2\lambda c}\right) \leq \int_{x}^{0}e^{2\lambda c}\phi\left(\widetilde{M}_{n}\left(y\right)e^{2\lambda c}\right)dy \leq C$$

which implies that $\{\widetilde{m}_n\}_{n\geq 1}$ has a convergent subsequence in the sense of the convergence in \mathcal{E} , namely the convergence under the conditions (A), (B). Since we have proved (4.4), the uniqueness of the spectral measure in $\mathcal{E}^{(c)}$ and Theorem 2 complete the proof.

Corollary 1. Suppose a measure σ on $[0,\infty)$ satisfies

$$\int_0^1 p(t)\phi(t)dt < \infty$$

with a function ϕ on [0,1] satisfying (S.1). Then, there exists an $m \in \mathcal{E}$ with spectral measure σ in \mathcal{S} .

Proof. Define σ_n by

$$\sigma_n\left(\xi\right) = \begin{cases} \sigma\left(\xi\right) & \text{for } \xi < n\\ \sigma\left(n\right) & \text{for } \xi \ge n \end{cases}$$

Then this σ_n satisfies all the conditions of Theorem 3. Applying the theorem we easily obtain the corollary.

This corollary provides a plenty of spectral measures in S growing faster than any power order at ∞ .

S.KOTANI

5. Continuity of the correspondence between \mathcal{E}_{ϕ} and \mathcal{S}_{ϕ}

In this section we give a necessary and sufficient condition for the continuity of the correspondence by restricting the order of growth of spectral measures at ∞ . We call ϕ to be a scale function if it satisfies the conditions (S.1), (S.2), (S.3). For a scale function ϕ set

$$\mathcal{E}_{\phi} = \left\{ m \in \mathcal{E}; \ \int_{-\infty}^{a} \phi\left(M\left(x\right)\right) dx < \infty \ \text{ for } \exists a \in (l_{-}, l_{+}) \right\},$$

and

$$\mathcal{S}_{\phi} = \left\{ \sigma; \int_{1}^{\infty} \widetilde{\phi}\left(\xi\right) \sigma\left(d\xi\right) < \infty \right\},$$

where

$$\widetilde{\phi}\left(\xi\right) = \int_{0}^{\infty} e^{-t\xi} \phi\left(t\right) dt.$$

It is easy to see that

$$\int_{1}^{\infty} \widetilde{\phi}\left(\xi\right) \sigma\left(d\xi\right) < \infty \Longleftrightarrow \int_{0}^{1} p(t)\phi\left(t\right) dt < \infty.$$

Moreover, from the properties of scales, it is always valid that for $\sigma \in \mathcal{S}_{\phi}$

$$\int_{1}^{\infty} \xi^{-\alpha - 1} \sigma\left(d\xi\right) < \infty$$

for an $\alpha \geq 1$. On the other hand, if $m \in \mathcal{E}_{\phi}$, (2) of Lemma 7 yields

$$\int_{0}^{\infty} p(t)\phi(t) e^{\lambda t} dt \leq C_{\phi} \int_{-\infty}^{a} \phi(M(x)) \varphi_{\lambda}(x)^{-2} dx + C_{\phi} \frac{-\lambda}{\varphi_{\lambda}'(a)} \int_{0}^{\infty} \phi(t) e^{\lambda t} dt$$
$$\leq C_{\phi} \int_{-\infty}^{a} \phi(M(x)) dx + C_{\phi} \frac{-\lambda}{\varphi_{\lambda}'(a)} \int_{0}^{\infty} \phi(t) e^{\lambda t} dt.$$

Therefore it holds that

$$\int_0^\infty p(t)\phi(t)\,e^{\lambda t}dt < \infty,$$

which shows $\sigma \in S_{\phi}$. Conversely assume $\sigma \in S_{\phi}$. Then

$$\int_0^\infty p(t)\phi(t) e^{\lambda t} dt = \int_0^1 p(t)\phi(t) e^{\lambda t} dt + \int_1^\infty p(t)\phi(t) e^{\lambda t} dt$$
$$\leq \int_0^1 p(t)\phi(t) dt + p(1) \int_1^\infty \phi(t) e^{\lambda t} dt < \infty$$

for any $\lambda < 0$. Therefore

$$\int_{-\infty}^{l} \phi\left(M(x)\varphi_{\lambda}(x)^{-2}\right)\varphi_{\lambda}(x)^{-2} dx < \infty$$

holds. Since ϕ satisfies the condition (S.3)

$$\phi\left(M(x)\varphi_{\lambda}(x)^{-2}\right) \ge C_{-}\left(\varphi_{\lambda}(x)^{-2}\right)\phi\left(M(x)\right)$$

holds. Due to

$$\varphi_{\lambda}(x)^{-2} \ge e^{2\lambda M(x)}$$

and $M(x) \to 0$ as $x \to -\infty$, we easily see

$$\int_{-\infty}^{a} \phi\left(M(x)\right) dx < \infty,$$

which implies $m \in \mathcal{E}_{\phi}$. Therefore, a string belongs to \mathcal{E}_{ϕ} if and only if its spectral measure is an element of \mathcal{S}_{ϕ} .

For ϕ let m_n, m be strings of \mathcal{E}_{ϕ} and define the convergence of m_n to m in \mathcal{E}_{ϕ} by

(C) $\lim_{x \to -\infty} \sup_{n \ge 1} \int_{-\infty}^{x} \phi(M_n(y)) \, dy = 0,$

in addition to the condition (A). The convergence of spectral measures in \mathcal{S}_{ϕ} is defined by

(A') $\sigma_n(\xi) \to \sigma(\xi)$ at every point of continuity of σ .

(C')
$$\lim_{N \to \infty} \sup_{n \ge 1} \int_{N}^{\infty} \widetilde{\phi}(\xi) \sigma_n(d\xi) = 0.$$

An equivalent statement is possible by p(t).

 $(A'') \quad p_n(t) \to p(t) \text{ for any } t > 0.$ $(C'') \quad \lim_{\epsilon \downarrow 0} \sup_{n \ge 1} \int_0^\epsilon p_n(t) \phi(t) \, dt = 0.$

Set

$$\mathcal{E}_{\phi}^{(c)} = \mathcal{E}_{\phi} \cap \mathcal{E}^{(c)}.$$

Theorem 4. Let $\{\sigma_n\}_{n\geq 1}, \sigma$ be elements of \mathcal{S}_{ϕ} and $m_n, m \in \mathcal{E}_{\phi}^{(c)}$ be the strings corresponding to σ_n, σ respectively. Then, $m_n \to m$ in $\mathcal{E}_{\phi}^{(c)}$ if and only if $\sigma_n \to \sigma$ in \mathcal{S}_{ϕ} .

Proof. Suppose $m_n \to m$ in $\mathcal{E}_{\phi}^{(c)}$. Then, Theorem 2 shows the validity of the condition (A'). Therefore we have only to check the condition (C''). From (2) of Lemma 7

$$\int_0^\infty p_n(t)\phi(t)\,e^{\lambda t}dt \le C_\phi \int_{-\infty}^0 \phi\left(M_n(x)\right)\varphi_\lambda^{(n)}\left(x\right)^{-2}\,dx + C_\phi \frac{-\lambda}{\varphi_\lambda^{(n)\prime}\left(0\right)} \int_0^\infty \phi\left(t\right)e^{\lambda t}dt$$

is valid. Fix $\epsilon > 0$ and choose a < 0 such that

$$C_{\phi} \int_{-\infty}^{a} \phi\left(M_n(x)\right) dx < \epsilon$$

for any $n \ge 1$. Since $M_n(x)$, $\varphi_{\lambda}^{(n)}(x)$ converge to M(x), $\varphi_{\lambda}(x)$ uniformly on $(-\infty, 0]$ and the estimate

$$\varphi_{\lambda}^{(n)}(0) \ge 1 - \lambda M_n(0) = 1 - \lambda d$$

show that if $-\lambda$ is sufficiently large, then

$$C_{\phi} \int_{a}^{0} \phi\left(M_{n}(x)\right) \varphi_{\lambda}^{(n)}\left(x\right)^{-2} dx < \epsilon$$

is valid for any $n \ge 1$. Moreover, due to c > 0

$$\underline{\lim}_{n \to \infty} m_n \left(0 \right) \ge m \left(0 - \right) > 0$$

holds, hence

$$C_{\phi}\frac{-\lambda}{\varphi_{\lambda}^{(n)\prime}\left(0\right)}\int_{0}^{\infty}\phi\left(t\right)e^{\lambda t}dt \leq C_{\phi}\frac{1}{m_{n}\left(0\right)}\int_{0}^{\infty}\phi\left(t\right)e^{\lambda t}dt < \epsilon$$

also holds for any $n \ge 1$ if we choose sufficiently large $-\lambda$, which implies

$$\sup_{n\geq 1} \int_0^\infty p_n(t)\phi(t) e^{\lambda t} dt \le 3\epsilon$$

From

$$\int_{0}^{-1/\lambda} p_n(t)\phi(t) \, dt \le e \int_{0}^{\infty} p_n(t)\phi(t) \, e^{\lambda t} dt \le 3e\epsilon$$

the condition (C'') is confirmed. Conversely assume $\sigma_n \to \sigma$ in S_{ϕ} . Then Theorem 3 shows the condition (A) holds. Hence we have only to check the condition (C). From (1) of Lemma 7

$$\int_0^\infty p_n(t)\phi(t)\,e^{\lambda t}dt \ge \int_{-\infty}^l \phi\left(M_n(x)\varphi_\lambda^{(n)}(x)^{-2}\right)\varphi_\lambda^{(n)}(x)^{-2}\,dx$$

follows. The property (S.3) implies

$$\phi\left(M_n(x)\varphi_{\lambda}^{(n)}(x)^{-2}\right) \ge C_{-}\left(\varphi_{\lambda}^{(n)}(x)^{-2}\right)\phi\left(M_n(x)\right)$$

and, as was pointed out in the proof of Theorem 3, $\varphi_{\lambda}^{(n)}(x) \to 1$ as $x \to -\infty$ uniformly with respect to *n*. Therefore, the condition (C'') guarantees the condition (C).

The last theorem can be restated as the convergence in \mathcal{E}_{ϕ} .

Theorem 5. Let $\{\sigma_n\}_{n\geq 1}, \sigma$ be elements of S_{ϕ} and $m_n, m \in \mathcal{E}_{\phi}$ be the strings corresponding to σ_n, σ respectively. Assume $\sigma_n \to \sigma$ in S_{ϕ} and σ is non-trivial. Then, there exist a sequence $\{a_n\}_{n\geq 1}$ in \mathbf{R} and c > 0 with $m_n (\cdot + a_n) \in \mathcal{E}_{\phi}^{(c)}$, and

$$m_n (\cdot + a_n) \to m$$

holds in \mathcal{E}_{ϕ} .

Proof. Since σ is non-trivial, we can apply Lemma 8. The rest of the proof is clear from Theorem 4.

For applications it will be helpful to rewrite the condition (C) as Kasahara-Watanabe did in [4].

Lemma 9. Assume ϕ satisfies (S.1). Then a sequence $\{m_n\}_{n\geq 1}$ converges to m in \mathcal{E}_{ϕ} , if and only if $\{m_n\}_{n\geq 1}$ and m satisfy the condition below. (D) For any $x \in \mathbf{R}$

$$\int_{-\infty}^{x} \phi\left(M_{n}(y)\right) dy \to \int_{-\infty}^{x} \phi\left(M(y)\right) dy.$$

Similarly the set of conditions (A') and (C') is equivalent to (D'), and that of (A'') and (C'') is equivalent to (D'').

(D') For any $\lambda < 0$

$$\int_0^\infty \widetilde{\phi}\left(\xi - \lambda\right) \sigma_n\left(d\xi\right) \to \int_0^\infty \widetilde{\phi}\left(\xi - \lambda\right) \sigma\left(d\xi\right).$$

(D'') For any $\lambda < 0$

$$\int_{0}^{\infty} p_{n}\left(t\right)\phi\left(t\right)e^{t\lambda}dt \rightarrow \int_{0}^{\infty} p\left(t\right)\phi\left(t\right)e^{t\lambda}dt$$

Proof. Assume $m_n \to m$ in \mathcal{E}_{ϕ} . Then, (C) implies that there exists c < l such that

$$\int_{-\infty}^{c} \phi\left(M_n(y)\right) dy \le 1,$$

hence for any x < c

$$\phi\left(M_n(x)\right)\left(c-x\right) \le \int_x^c \phi\left(M_n(y)\right) dy \le \int_{-\infty}^c \phi\left(M_n(y)\right) dy \le 1,$$

which shows

$$M_n(x) \le \phi^{-1}\left(\frac{1}{c-x}\right)$$

for any $n \ge 1$ and x < c. Then it is easy to see that $M_n(x) \to M(x)$ at every point x, and this together with (C) implies (D). Conversely, for any $\epsilon > 0$, choose c < l such that

$$\int_{-\infty}^{c} \phi\left(M(y)\right) dy < \epsilon.$$

Then, clearly (C) follows from (D). The condition (A) can be derived from (D) by the monotonicity of ϕ and M_n . We omit the proof for (D') and (D'').

6. Application

Typical examples of m belonging to \mathcal{E} are

$$m_{\alpha}(x) = \begin{cases} C_{\alpha} x^{-\beta} & x > 0 & \text{if } 0 < \alpha < 1 \\ e^{x} & x \in \mathbf{R} & \text{if } \alpha = 1 \\ C_{\alpha} (-x)^{-\beta} & x < 0 & \text{if } \alpha > 1. \end{cases} \quad \text{with } \beta = \frac{\alpha}{\alpha - 1}$$

and the spectral measures and p(t) are

$$\sigma_{\alpha}(d\xi) = \frac{\alpha^{2\alpha}}{\Gamma(1+\alpha)^2} d\xi^{\alpha}, \quad p_{\alpha}(t) = \frac{\alpha^{2\alpha}}{\Gamma(1+\alpha)} t^{-\alpha},$$

where

$$C_{\alpha} = \begin{cases} \left(\frac{1-\alpha}{\alpha}\right)^{\frac{\alpha}{1-\alpha}} &, \quad 0 < \alpha < 1\\ \left(\frac{\alpha-1}{\alpha}\right)^{-\frac{\alpha}{\alpha-1}} &, \quad 1 < \alpha \end{cases}$$

.

In this section we consider the asymptotic behavior of the spectral measures and the transition probability densities when strings are close to the above typical ones. If $\alpha \in (0, 2)$, the following results are already known. Here we denote

$$f(x) \sim g(x)$$
 as $x \uparrow 0, (x \to \infty)$

if

$$\lim_{x \to 0} \frac{f(x)}{g(x)} = 1, \quad \left(\lim_{x \to \infty} \frac{f(x)}{g(x)} = 1\right)$$

hold respectively. Let φ be a function regularly varying at 0 with exponent $\alpha - 1$.

Theorem 6. (Kasahara[1], Kasahara-Watanabe[4]) The following asymptotic relationship between m and p is valid.

(1) If $\alpha \in (0,1)$, then

$$m(x) \sim \frac{(-\beta)^{\beta}}{x\varphi^{-1}(x)} \quad as \ x \uparrow \infty$$

holds if and only if

$$p(t) \sim \frac{\alpha^{2\alpha}}{\Gamma(1+\alpha)} \frac{1}{t} \varphi\left(\frac{1}{t}\right) \quad as \ t \to \infty$$

(2) If $\alpha \in (1, 2)$, then

$$m(x) \sim \frac{\beta^{\beta}}{-x\varphi^{-1}\left(-x\right)} \quad as \ x \uparrow 0$$

holds if and only if

$$p(t) \sim \frac{\alpha^{2\alpha}}{\Gamma(1+\alpha)} \frac{1}{t} \varphi\left(\frac{1}{t}\right) \quad as \ t \to \infty$$

They showed an analogous result in case $\alpha = 1$ in Kasahara-Watanabe[4]. In this section we extend their results to the case $\alpha \ge 2$ by applying Theorem 4. The basic idea, which was first employed by Kasahara[1], is to use the continuity between m and p and the scaling relationship

(6.1)
$$abm(ax) \leftrightarrow \frac{1}{ab}p\left(b^{-1}t\right)$$

for any a, b > 0. The proof proceeds just like Kasahara-Watanabe[4], especially the case $\alpha = 1$.

Let $m \in \mathcal{E}$ be a non-decreasing function with l = 0, namely

 $m(x) < \infty$ on $(-\infty, 0)$ and $m(x) = \infty$ on $(0, \infty)$.

Let φ be a regularly varying function at 0 with exponent $\alpha - 1$ and set

$$m_{\nu}(x) = \nu \varphi^{-1}(\nu) m(\nu x).$$

Then from (6.1) we have

(6.2)
$$\begin{cases} M_{\nu}(x) = \varphi^{-1}(\nu) M(\nu x) \\ p_{\nu}(t) = \nu^{-1} \varphi^{-1}(\nu)^{-1} p\left(\varphi^{-1}(\nu)^{-1} t\right) \\ \sigma_{\nu}(\xi) = \nu^{-1} \varphi^{-1}(\nu)^{-1} \sigma\left(\varphi^{-1}(\nu)\xi\right) \end{cases}$$

To consider an extension of Theorem 6 we introduce conditions on m and σ ;

(6.3)
$$m(x) \sim \frac{\beta^{\beta}}{-x\varphi^{-1}(-x)} \quad \text{as} \quad x \uparrow 0,$$

which means

(6.4)
$$M(x) \sim \frac{\beta^{\beta}}{(\beta - 1)\varphi^{-1}(-x)} \quad \text{as} \ x \uparrow 0,$$

and

(6.5)
$$p(t) = \int_0^\infty e^{-t\xi} d\sigma(\xi) < \infty \quad \text{for any } t > 0.$$

Proposition 1. If $m \in \mathcal{E}$ satisfies (6.3), then it holds that

(6.6)
$$\sigma\left(\xi\right) \sim \frac{\alpha^{2\alpha}}{\Gamma\left(1+\alpha\right)^2} \xi^{\alpha} \quad as \ \xi \downarrow 0.$$

Moreover, if m satisfies (6.5) as well, then (6.7) below holds.

(6.7)
$$p(t) \sim \frac{\alpha^{2\alpha}}{\Gamma(1+\alpha)} \frac{1}{t} \varphi\left(\frac{1}{t}\right) \quad as \quad t \to \infty.$$

Proof. Since

$$M_{\nu}(x) = \int_{-\infty}^{x} m_{\nu}(y) \, dy = \varphi^{-1}(\nu) \, M(\nu x)$$

holds, from (6.4) we know

$$M_{\nu}(x) \to \frac{\beta^{\beta}}{\beta - 1} (-x)^{1 - \beta} \text{ as } \nu \to 0$$

for any x < 0, which means that $\{M_{\nu}\}$ satisfies the condition (B). Applying Theorem 2 yields

$$\sigma_{\nu}(\xi) \to \frac{\alpha^{2\alpha}}{\Gamma(1+\alpha)^2} \xi^{\alpha} \text{ for any } \xi > 0$$

as $\nu \to 0$, which is equivalent to (6.6) due to (6.2). If we assume the condition (6.5) as well on σ , the Abelian theorem for Laplace transform shows the property (6.7).

To obtain a converse statement to the above proposition we need

Lemma 10. Assume $\sigma \in S$ satisfies the condition (6.5) and a condition

(6.8)
$$\int_0^1 p(t)\phi(t)\,dt < \infty$$

for a positive function ϕ on [0,1] satisfying

(6.9)
$$\phi(st) \le Ct^k \phi(s) \text{ for any } s, t \le 1 \text{ for some } k > \alpha - 1.$$

Then $\{p_{\nu}(t)\}$ satisfies the condition (28), namely

(6.10)
$$\sup_{\nu>0}\int_0^1 p_{\nu}(t)\,\phi(t)\,dt < \infty.$$

Proof. Since φ is a regularly varying function at 0 with exponent $\alpha - 1$, $t^{-1}\varphi(t^{-1})$ is regularly varying at ∞ with exponent $-\alpha$, and there exists a slowly varying function l(t) such that

$$\frac{1}{t}\varphi\left(\frac{1}{t}\right) = t^{-\alpha}l(t).$$

Generally a slowly varying function l(t) has an expression

(6.11)
$$l(t) = c(t) \exp\left(\int_{a}^{t} \frac{\epsilon(u)}{u} du\right)$$

with a positive constant a and functions c(t), $\epsilon(t)$ behaving as

 $c(t) \rightarrow c > 0, \ \ \epsilon \left(t \right) \rightarrow 0 \ \ \text{as} \ \ t \rightarrow \infty.$

Now we decompose the integral in (6.10) into two parts:

$$\int_{0}^{1} p_{\nu}(t) \phi(t) dt = \nu^{-1} \varphi^{-1}(\nu)^{-1} \int_{0}^{1} p\left(\varphi^{-1}(\nu)^{-1} t\right) \phi(t) dt = I_{1} + I_{2}$$

with

$$\begin{cases} I_1 = \nu^{-1} \varphi^{-1} (\nu)^{-1} \int_{N\varphi^{-1}(\nu)}^{1} p\left(\varphi^{-1} (\nu)^{-1} t\right) \phi(t) dt, \\ I_2 = \nu^{-1} \varphi^{-1} (\nu)^{-1} \int_0^{N\varphi^{-1}(\nu)} p\left(\varphi^{-1} (\nu)^{-1} t\right) \phi(t) dt, \end{cases}$$

where N is chosen so that

$$|\epsilon(u)| \leq \delta$$
 for any $u \geq N$

holds with a positive δ satisfying $\delta < k - (\alpha - 1)$. Since the condition (6.7) implies

$$0 < \frac{p\left(\varphi^{-1}\left(\nu\right)^{-1}t\right)}{\varphi^{-1}\left(\nu\right)^{\alpha}t^{-\alpha}l\left(\varphi^{-1}\left(\nu\right)^{-1}t\right)} \le C'$$

for any $t \ge N\varphi^{-1}(\nu)$ with some constant C', we have

$$I_{1} \leq C' \nu^{-1} \varphi^{-1} (\nu)^{-1+\alpha} l \left(\varphi^{-1} (\nu)^{-1}\right) \int_{N \varphi^{-1}(\nu)}^{1} t^{-\alpha} \frac{l \left(\varphi^{-1} (\nu)^{-1} t\right)}{l \left(\varphi^{-1} (\nu)^{-1}\right)} \phi(t) dt.$$

First note

$$\nu^{-1}\varphi^{-1}(\nu)^{-1+\alpha}l\left(\varphi^{-1}(\nu)^{-1}\right) = \nu^{-1}\varphi^{-1}(\nu)^{-1+\alpha}\varphi^{-1}(\nu)^{1-\alpha}\varphi\left(\varphi^{-1}(\nu)\right) = 1,$$

and the (6.11) shows for $t \ge N\varphi^{-1}(\nu)$

$$\frac{l\left(\varphi^{-1}\left(\nu\right)^{-1}t\right)}{l\left(\varphi^{-1}\left(\nu\right)^{-1}\right)} = \exp\left(\int_{a}^{\varphi^{-1}\left(\nu\right)^{-1}t} \frac{\epsilon\left(u\right)}{u} du - \int_{a}^{\varphi^{-1}\left(\nu\right)^{-1}} \frac{\epsilon\left(u\right)}{u} du\right)$$
$$= \exp\left(-\int_{\varphi^{-1}\left(\nu\right)^{-1}t}^{\varphi^{-1}\left(\nu\right)^{-1}t} \frac{\epsilon\left(u\right)}{u} du\right)$$
$$\leq \exp\left(\delta \log t^{-1}\right) = t^{-\delta}.$$

In (6.9) setting s = 1, we have $\phi(t) \leq C\phi(1)t^k$, hence

$$I_1 \le C \int_{N\varphi^{-1}(\nu)}^{1} t^{-\alpha} t^{-\delta} \phi(t) \, dt \le CC' \phi(1) \int_0^1 t^{-\alpha-\delta+k} dt$$

is valid. Due to (6.9) I_2 can be estimated as

$$I_{2} = \nu^{-1} \int_{0}^{N} p(s) \phi\left(\varphi^{-1}(\nu)s\right) ds$$

$$\leq C\nu^{-1} \left(\varphi^{-1}(\nu)\right)^{k} \int_{0}^{N} p(s) \phi(s) ds \leq C'' \nu^{-1 + \frac{k}{\alpha - 1} - \delta'} \int_{0}^{N} p(s) \phi(s) ds,$$

where $\delta' > 0$ can be chosen so that

$$-1 + \frac{k}{\alpha - 1} - \delta' > 0$$

holds. Consequently we have

$$\int_{0}^{1} p_{\nu}(t) \phi(t) dt \leq CC' \phi(1) \int_{0}^{1} t^{-\alpha-\delta+k} dt + C'' \nu^{-1+\frac{k}{\alpha-1}-\delta'} \int_{0}^{N} p(s) \phi(s) ds,$$

and (6.8) implies the second assertion of (6.10).

and (6.8) implies the second assertion of (6.10).

Remark 1. The property (6.9) are satisfied not only by $\phi(t) = t^k$ with $k > \alpha - 1$ but also by subexponential functions: for p > 1, c > 0

$$\phi(t) = \exp\left(-c\left(-\log t\right)^p\right).$$

Within the knowledge of the previous sections the best converse statement to Proposition 1 is as follows.

Proposition 2. Let $m \in \mathcal{E}$ be a non-decreasing function with l = 0 and $M(0) = \infty$. Assume $\sigma \in S$ satisfies the conditions (6.5) and (6.8) with a positive function ϕ on [0,1] satisfying (S.1) and (6.9). Then the property (6.6) (equivalently (6.7)) implies (6.3).

Proof. First note (6.6) is equivalent to

$$\sigma_{\nu}\left(\xi\right) \to \frac{\alpha^{2\alpha}}{\Gamma\left(1+\alpha\right)^{2}}\xi^{\alpha} \quad \text{for any } \xi > 0,$$

as $\nu \to 0$. Since we are assuming $M(0) = \infty$,

$$M_{\nu}\left(0\right) = \varphi^{-1}\left(\nu\right) M\left(0\right) = \infty$$

holds for any $\nu > 0$, and there exists uniquely $a_{\nu} < 0$ such that

$$M_{\nu}\left(a_{\nu}\right) = \frac{\beta^{\beta}}{\beta - 1} \equiv c.$$

Set

$$\widetilde{M}_{\nu}(x) = M_{\nu}(x + a_{\nu} + 1).$$

Then, taking -1 instead of 0 as a normalization point, Lemma 10 makes it possible to apply Theorem 3 and we have

$$\widetilde{M}_{\nu}(x) \to \begin{cases} c(-x)^{1-\beta} & \text{for } x < 0\\ \infty & \text{for } x > 0 \end{cases}$$

holds in \mathcal{E} as $\nu \to 0$, from which

(6.12)
$$\varphi^{-1}(\nu) M(\nu(x+a_{\nu}+1)) \to \begin{cases} c(-x)^{1-\beta} & \text{for } x < 0 \\ \infty & \text{for } x > 0 \end{cases}$$

follows. To simplify the involved formula (6.12) we take their inverse. Set

$$u = M (\nu (x + a_{\nu} + 1)), \quad \lambda = c \varphi^{-1} (\nu)^{-1}$$

Since $\varphi^{-1}(\nu) M(\nu a_{\nu}) = c$, we easily see

$$\varphi\left(c\lambda^{-1}\right)\left(x+1\right) + M^{-1}\left(\lambda\right) = M^{-1}\left(u\right)$$

Denoting $y = \lambda^{-1} u$, (6.12) is equivalent to

$$y \to (-x)^{1-\beta} \,,$$

from which

$$\frac{M^{-1}(\lambda y) - M^{-1}(\lambda)}{\varphi(c\lambda^{-1})} = x + 1 \to 1 - y^{-(\beta - 1)}$$

follows for any y > 0 as $\lambda \to \infty$. Since φ is regularly varying at 0 with exponent $\alpha - 1$,

$$\frac{\varphi(c\lambda^{-1})}{\varphi(\lambda^{-1})} \to c^{\alpha-1} = (\alpha-1)^{-1} \alpha^{\alpha} \quad \text{as } \lambda \to \infty.$$

and

(6.13)
$$\lim_{\lambda \to \infty} \frac{M^{-1}(\lambda x) - M^{-1}(\lambda)}{\varphi(\lambda^{-1})} = (\alpha - 1)^{-1} \alpha^{\alpha} \left(1 - x^{-(\alpha - 1)}\right)$$

follow. Then Lemma 11 below shows (6.3).

Lemma 11. (6.13) *implies* (6.3).

Proof. Assume (6.13). Since $M^{-1}(x)$ has a monotone density

$$(M^{-1}(x))' = \frac{1}{m(M^{-1}(x))},$$

the monotone density theorem implies

$$\lim_{\lambda \to \infty} \left(\frac{M^{-1} \left(\lambda x \right) - M^{-1} \left(\lambda \right)}{\varphi \left(\lambda^{-1} \right)} \right)' = \left(\left(\alpha - 1 \right)^{-1} \alpha^{\alpha} \left(1 - x^{-(\alpha - 1)} \right) \right)',$$

which is

$$\lim_{\lambda \to \infty} \frac{\lambda}{\varphi(\lambda^{-1}) m(M^{-1}(\lambda x))} = \alpha^{\alpha} x^{-\alpha}.$$

Setting x = 1 and $u = M^{-1}(\lambda)$, we have

$$\lim_{u \to 0} \frac{M(u)}{\varphi(M(u)^{-1}) m(u)} = \alpha^{\alpha}.$$

For any $\epsilon > 0$ there exists $\delta > 0$ such that for any $u \in (-\delta, 0)$

$$\alpha^{-\alpha} - \epsilon \le \frac{\varphi\left(M(u)^{-1}\right)m(u)}{M(u)} \le \alpha^{-\alpha} + \epsilon$$

are valid. Noting m(u) = M(u)', we see

$$\left(\alpha^{-\alpha} - \epsilon\right)(-x) \le \int_{x}^{0} \frac{\varphi\left(M(u)^{-1}\right)}{M(u)} dM\left(u\right) \le \left(\alpha^{-\alpha} + \epsilon\right)(-x)$$

for any $x \in (-\delta, 0)$, hence

$$\left(\alpha^{-\alpha} - \epsilon\right)(-x) \le \int_{M(x)}^{\infty} \frac{\varphi\left(y^{-1}\right)}{y} dy = \int_{0}^{M(x)^{-1}} \frac{\varphi\left(z\right)}{z} dz \le \left(\alpha^{-\alpha} + \epsilon\right)(-x).$$

Since $\varphi(z)/z$ is a regularly varying function at 0 with exponent $\alpha - 2$,

$$\int_0^y \frac{\varphi(z)}{z} dz \sim \frac{\varphi(y)}{\alpha - 1} \text{ as } y \downarrow 0$$

is valid, which implies

$$\frac{\varphi\left(M\left(x\right)^{-1}\right)}{\alpha-1} \sim \alpha^{-\alpha}\left(-x\right) \quad \text{as } x \uparrow 0,$$

hence

$$M(x) \sim \frac{\beta^{\beta}}{\beta - 1} \varphi^{-1} (-x)^{-1} \quad \text{as } x \uparrow 0$$

This is equivalent to (6.3).

In the above two propositions we stated the conditions which should be satisfied by $m \in \mathcal{E}$ in terms of its spectral function σ . It may be preferable to describe the result by m itself directly. To do so, unfortunately we have to impose a more restrictive condition on m, and combining Proposition 1 and Proposition 2 we have

Theorem 7. Let $\alpha \geq 2$, $k > \alpha - 1$ and φ is a regularly varying function at 0 with exponent $\alpha - 1$. Let $m \in \mathcal{E}$ be a non-decreasing function with l = 0 and $M(0) = \infty$. Assume m satisfies

$$\int_{-\infty}^{-1} M(x)^k dx < \infty.$$

Then, the property

$$p\left(t\right)\sim\frac{\alpha^{2\alpha}}{\Gamma\left(1+\alpha\right)}\frac{1}{t}\varphi\left(\frac{1}{t}\right)\quad as \ t\rightarrow\infty$$

holds if and only if the asymptotics below is valid.

$$m(x) \sim \frac{\beta^{\beta}}{-x\varphi^{-1}(-x)} \quad as \ x \uparrow 0.$$

Proof. The proof is immediate from the above two propositions if we observe $\phi(t) = t^k$ satisfies all the requirements needed in Proposition 2.

Acknowledgement. The author would like to express his hearty thanks to Professor Y.Kasahara who allowed him to read a preprint, which was very helpful in the course of proving Proposition 2 and Lemma 11.

References

- Y.Kasahara. Spectral theory of generalized second order differential operators and its application to Markov processes, Japan. J. Math. 1 (1975), 67-84
- [2] Y.Kasahara and S.Watanabe. Brownian representation of a class of Lévy processes and its application to occupation times of diffusion processes, Illinois J. Math., 50 (2006), 515-539
- [3] Y.Kasahara and S.Watanabe. Remarks on Krein-Kotani's correspondence between strings and Herglotz functions, Proc. Japan. Acad. 85 Ser.A (2009), 22-26
- [4] Y.Kasahara and S.Watanabe. Asymptotic behavior of spectral measures and Krein's and Kotani's strings, Kyoto Journal of Math. 50 (2010), 623-644
- [5] S.Kotani. On a generalized Sturm-Liouville operator with a singular boundary, J. Math. Kyoto Univ. 15 (1975), 423-454
- [6] S.Kotani. A remark to the ordering theorem of L.de Branges, J. Math. Kyoto Univ. 16 (1976), 665-674
- [7] S.Kotani. Krein's strings with singular left boundary, Rep. Math. Phys. 59 (2007), 305-316

22

KREIN'S STRINGS WHOSE SPECTRAL FUNCTIONS ARE OF POLYNOMIAL GROWTH 23

- [8] M.G.Krein. On a generalization of investigation of Stieltjes, Dokl. Akad. Nauk SSSR 87 (1952), 881-884 (Russian)
- K.Yano. Excursion measure away from an exit boundary of one-dimensional diffusion processes, Publ. RIMS 42, (2006), 837-878

KWANSEI GAKUIN UNIVERSITY E-mail address: kotani@kwansei.ac.jp