

# GMRT discovery of PSR J1544+4937, an eclipsing black-widow pulsar identified with a *Fermi* LAT source

B. Bhattacharyya<sup>1</sup>, J. Roy<sup>1</sup>, P. S. Ray<sup>2</sup>, Y. Gupta<sup>1</sup>, D. Bhattacharya<sup>3</sup>, R. W. Romani<sup>4</sup>, S. M. Ransom<sup>5</sup>, E. C. Ferrara<sup>6</sup>, M. T. Wolff<sup>2</sup>, F. Camilo<sup>7,8</sup>, I. Cognard<sup>9</sup>, A. K. Harding<sup>6</sup>, P. R. den Hartog<sup>4</sup>, S. Johnston<sup>10</sup>, M. Keith<sup>10</sup>, M. Kerr<sup>4</sup>, P. F. Michelson<sup>4</sup>, P. M. Saz Parkinson<sup>11</sup>, D. L. Wood<sup>12</sup>, K. S. Wood<sup>2</sup>

## ABSTRACT

Using the Giant Metrewave Radio Telescope (GMRT) we performed deep observations to search for radio pulsations in the directions of unidentified *Fermi* Large Area Telescope (LAT)  $\gamma$ -ray sources. We report the discovery of an eclipsing black-widow millisecond pulsar, PSR J1544+4937, identified with the uncataloged  $\gamma$ -ray source FERMI J1544.2+4941. This 2.16 ms pulsar is in a 2.9

---

<sup>1</sup>National Centre for Radio Astrophysics, Tata Institute of Fundamental Research, Pune 411 007, India

<sup>2</sup>Space Science Division, Naval Research Laboratory, Washington, DC 20375-5352, USA

<sup>3</sup>Inter-University Centre for Astronomy and Astrophysics, Pune 411 007, India

<sup>4</sup>W. W. Hansen Experimental Physics Laboratory, Kavli Institute for Particle Astrophysics and Cosmology, Department of Physics and SLAC National Accelerator Laboratory, Stanford University, Stanford, CA 94305, USA

<sup>5</sup>National Radio Astronomy Observatory (NRAO), Charlottesville, VA 22903, USA

<sup>6</sup>NASA Goddard Space Flight Center, Greenbelt, MD 20771, USA

<sup>7</sup>Columbia Astrophysics Laboratory, Columbia University, New York, NY 10027, USA

<sup>8</sup>Arecibo Observatory, Arecibo, Puerto Rico 00612, USA

<sup>9</sup>Laboratoire de Physique et Chimie de l'Environnement, LPCE UMR 6115 CNRS, F-45071 Orléans Cedex 02, and Station de radioastronomie de Nançay, Observatoire de Paris, CNRS/INSU, F-18330 Nançay, France

<sup>10</sup>CSIRO Astronomy and Space Science, Australia Telescope National Facility, Epping NSW 1710, Australia

<sup>11</sup>Santa Cruz Institute for Particle Physics, Department of Physics and Department of Astronomy and Astrophysics, University of California at Santa Cruz, Santa Cruz, CA 95064, USA

<sup>12</sup>Praxis Inc., Alexandria, VA 22303, resident at Naval Research Laboratory, Washington, DC 20375, USA

hours compact circular orbit with a very low-mass companion ( $M_c > 0.017M_\odot$ ). At 322 MHz this pulsar is found to be eclipsing for 13% of its orbit, whereas at 607 MHz the pulsar is detected throughout the low-frequency eclipse phase. Variations in the eclipse ingress phase are observed, indicating a clumpy and variable eclipsing medium. Moreover, additional short-duration absorption events are observed around the eclipse boundaries. Using the radio timing ephemeris we were able to detect  $\gamma$ -ray pulsations from this pulsar, confirming it as the source powering the  $\gamma$ -ray emission.

*Subject headings:* pulsars: general; binaries: eclipsing, pulsars: individual (PSR J1544+4937)

## 1. Introduction

The Large Area Telescope (LAT; Atwood et al. 2009) aboard the *Fermi Gamma-ray Space Telescope* has discovered a large number of  $\gamma$ -ray point sources, of which many are unidentified or even unassociated with any known potential counterpart (Ackermann et al. 2012). The LAT can localize most of these sources well enough that they can be covered in a single pointing with the large primary beams of radio telescopes at low-frequencies, allowing them to be searched efficiently. Targeted radio searches of unassociated LAT point sources by the *Fermi* Pulsar Search Consortium (PSC) have resulted in the discovery of 43 radio millisecond pulsars (MSPs; Ray et al. 2012). MSPs are thought to evolve from normal pulsars in binary systems via transfer of angular momentum from companions. Thus, the majority of MSPs are naturally expected to be in binaries ( $\sim 83\%$  being the binary fraction for MSPs in the Galactic field<sup>1</sup>). Binary systems where the pulsar wind evaporates the companion are one way to form isolated MSPs. Such systems where the interaction is ongoing are called black-widow (BW) pulsars. Many exhibit long eclipses ( $\sim 10\%$  of the orbital period, apparently larger than the companion’s Roche lobe) that are believed to be caused by the material blown from the very low mass companion ( $M_c \ll 0.1M_\odot$ ) by the pulsar wind. There were two such eclipsing BW systems in the Galactic field known before the launch of *Fermi* – PSR B1957+20 (Fruchter et al. 1988) and PSR J2051–0827 (Stappers et al. 1996). The BW pulsars are found to have higher values of spin-down energy-loss rate ( $\dot{E} \sim 10^{34}$  erg s<sup>−1</sup>) compared to other MSPs, making these systems good candidates for pulsed  $\gamma$ -ray emission (Roberts 2011). Among 43 new MSPs found in *Fermi*-directed searches there are at least

---

<sup>1</sup><http://astro.phys.wvu.edu/GalacticMSPs/>

10 BWs (Ray et al. 2012). This Letter describes the discovery and follow-up study of an eclipsing BW MSP, J1544+4937, with the GMRT.

## 2. Observations and search analysis

As a part of the PSC search effort, we observed mid- and high-Galactic-latitude unassociated *Fermi* point sources with the GMRT at 607 MHz. The GMRT Software Back-end (GSB; Roy et al. 2010) produces simultaneous incoherent and coherent beam filter-bank outputs of  $512 \times 0.0651$  MHz sampled every  $61.44 \mu\text{s}$ . The wider incoherent beam of the GMRT (beam-width  $40'$  at 607 MHz) can easily cover error circles associated with the *Fermi* sources. In addition a coherent beam that is  $3 \times$  more sensitive and narrower ( $1.5'$  at 607 MHz) can be useful if the pulsar happens to be near the pointing center.

One of the targets was FERMI J1544.2+4941, a  $\gamma$ -ray source from an unpublished internal source list created by the LAT Collaboration using 18 months of data in preparation for the 2FGL catalog (Nolan et al. 2012). The source location (J2000) from that analysis (and used for our telescope pointing) was R.A. =  $236^\circ.074$ , Decl. =  $49^\circ.695$ , with a 95% confidence error circle of radius  $9.5'$ . This source is very weak, with a test statistic (TS) of 26.2 (Mattox et al. 1996) in the 18 month analysis, and did not make the significance cut to be included in the 2FGL catalog itself.

We processed the data on an IUCAA HPC cluster with Fourier-based acceleration search methods using search techniques implemented in PRESTO (Ransom et al. 2002). We investigated trial dispersion measures (DMs) ranging from  $0 \text{ pc cm}^{-3}$  up to  $350 \text{ pc cm}^{-3}$ . A linear drift of up to 200 Fourier-frequency bins for the highest summed harmonic was allowed. The powerline, 50 Hz, and its subsequent harmonics were excised. Using parameters of 32 MHz bandwidth, 10% duty-cycle, incoherent array gain of 2.3 K/Jy, for 30 minutes of observing, we estimate the search sensitivity as  $(92\text{K} + T_{\text{sky}})/(335\text{K}) \text{ mJy}$  for a  $5\sigma$  detection at 607 MHz. Considering  $|b| > 5^\circ$ , where  $T_{\text{sky}} \sim 10\text{--}45 \text{ K}$ , our search sensitivity is 0.3–0.4 mJy.

In a 30-minute pointing on 2011 February 1, towards FERMI J1544.2+4941 we discovered a binary MSP of period 2.16 ms with significant acceleration of  $2.25 \text{ m s}^{-2}$  at a DM of  $23.2 \text{ pc cm}^{-3}$ .

## 3. Follow-up timing

We localized J1544+4937 with an accuracy of  $5''$  (positions listed in Table 1) using continuum imaging followed by multi-pixel beamforming (Roy et al. 2012), which allowed us

to have sensitive follow-up studies using the GMRT coherent array. We estimate a flux of 5.4 mJy at 322 MHz, and a spectral index of  $-2.3$ . We started the regular timing campaign for J1544+4937 in April 2011 at 322 MHz. With the derived position from the multi-pixel search and an *a priori* binary model predicted by Bhattacharyya & Nityananda (2008), we obtained phase-connected time-of-arrivals (TOAs) from TEMPO<sup>2</sup>, using the JPL DE405 solar system ephemeris (Standish 2004). The binary timing model used is ELL1 (Lange et al. 2001), since J1544+4937 is in a very low eccentricity system. This MSP is in a very compact binary with an orbital period of 2.9 hours. We derive a minimum companion mass (for  $90^\circ$  orbital inclination) of  $0.017 M_\odot$  using the Keplerian mass function, assuming a pulsar mass of  $1.4 M_\odot$ . J1544+4937 is eclipsed for about 13% of the orbit at 322 MHz (Sec.6). The best-fit timing model (MJD 55680.927–56332.90) is obtained excluding the TOAs around the eclipse phase. We achieved a post-fit rms timing residual of  $6.9 \mu\text{s}$  from 652 days of timing (Fig. 1). We estimate a precise DM equal to  $23.2258(1) \text{ pc cm}^{-3}$  by a timing fit using 322 and 607 MHz TOAs from non-eclipsing binary phases (excluding TOAs between phases 0.05–0.35). The resulting ephemeris, position from timing fit and derived parameters are listed in Table 1.




Fig. 1.— Post-fit timing residuals of J1544+4937 considering non-eclipsing binary phases. The best-fit timing solution is given in Table 1.

Table 1. Parameters of J1544+4937

Parameter	Value <sup>a</sup>
Interferometric position <sup>b</sup>	
Right ascension (J2000) .....	15 <sup>h</sup> 44 <sup>m</sup> 04 <sup>s</sup> .166±0 <sup>s</sup> .3
Declination (J2000) .....	+49°37′57″.45±4″.7
Offset from survey beam center .....	4.3′
Parameters from radio timing	
Right ascension (J2000) .....	15 <sup>h</sup> 44 <sup>m</sup> 04 <sup>s</sup> .48722 (2)
Declination (J2000) .....	+49°37′55″.2545 (2)
Position Epoch (MJD) .....	51544.0
Pulsar period $P$ (ms) .....	2.15928839043289 (5)
Pulsar frequency $f$ (Hz) .....	463.11553585462 (1)
Frequency derivative $\dot{f}$ (Hz s <sup>-1</sup> ) .....	-6.29 (1)×10 <sup>-16</sup>
Period Epoch (MJD) .....	56007.0
Dispersion measure $DM$ (cm <sup>-3</sup> pc) .....	23.2258 (1)
Binary model .....	ELL1
Orbital period $P_b$ (days) .....	0.1207729895 (1)
Projected semi-major axis $x$ (lt-s) .....	0.0328680 (4)
Epoch of ascending node passage $T_{ASC}$ (MJD) .....	56124.7701121 (2)
Span of timing data (MJD) .....	652
Number of TOAs .....	280
Post-fit residual rms ( $\mu$ s) .....	6.9
Reduced chi-square .....	2.7
Derived parameters	
Mass function $f$ ( $M_\odot$ ) .....	0.0000026132
Min companion Mass $m_c$ ( $M_\odot$ ) .....	0.017
DM distance <sup>c</sup> (kpc) .....	1.2
Flux density at 322 MHz (mJy) .....	5.4
Flux density at 607 MHz (mJy) .....	1.2
Spectral index .....	-2.3
Surface magnetic field $B_s$ (10 <sup>8</sup> G) .....	0.805 (1)
Spin down luminosity $\dot{E}$ (10 <sup>34</sup> erg s <sup>-1</sup> ) .....	1.150 (8)
Characteristic age $\tau$ (Gyr) .....	11.65 (3)
$\gamma$ -ray parameters	
Photon flux (> 0.1 GeV, cm <sup>-2</sup> s <sup>-1</sup> ) .....	1.6(8) ×10 <sup>-9</sup>
Energy flux (> 0.1 GeV, erg cm <sup>-2</sup> s <sup>-1</sup> ) .....	2.1(6) ×10 <sup>-12</sup>
Luminosity, $L_\gamma/f_\Omega$ (> 0.1 GeV, erg cm <sup>-2</sup> s <sup>-1</sup> ) .....	3.6 ×10 <sup>32</sup>
Efficiency, $\eta_\gamma/f_\Omega$ (> 0.1 GeV) .....	0.03

<sup>a</sup>Errors in the last digit are in parentheses.

<sup>b</sup>Roy et al. (2012)

<sup>c</sup>Cordes & Lazio (2002)




Fig. 2.— Phase-aligned radio and LAT  $\gamma$ -ray light curves of J1544+4937. The horizontal dashed lines are an estimate of the background level from sources other than the pulsar.

#### 4. $\gamma$ -ray pulsations

The radio timing position of J1544+4937 is  $4.3'$  from the LAT localization of FERMI J1544.2+4941. This is well within the radius of the 95% confidence error circle, suggesting an association between the pulsar and the LAT source, which we sought to confirm with detection of pulsations. In addition, the  $\gamma$ -ray detectability metric  $\dot{E}^{1/2}/d^2$  for J1544+4937 is  $7.4 \times 10^{16} \text{ erg}^{1/2} \text{ kpc}^{-2} \text{ s}^{-1/2}$ , which is comparable to other  $\gamma$ -ray detected MSPs (Fig. 12 of Abdo et al. 2010).

Because of the low significance of the LAT source, a spectral analysis of the region

---

<sup>2</sup><http://www.atnf.csiro.au/research/pulsar/tempo>

is required to optimize the sensitivity of the H-test for pulsed significance using photon probability weighting (Kerr 2011). We performed a binned likelihood spectral analysis of the region using the *Fermi* Science Tools<sup>3</sup>. We selected LAT data with reconstructed energies in the range 100 MeV to 20 GeV collected between 2008 August 4 and 2013 February 6. We used the P7SOURCE\_V6 instrument response functions and excluded events at a zenith angle  $> 100^\circ$  and times when the LAT was in the SAA or the rocking angle was greater than  $52^\circ$ . We modeled<sup>4</sup> a region of radius  $15^\circ$  using a source list from the 2FGL catalog with an additional source added at the location of the pulsar. We modeled the pulsar with an exponentially cutoff power-law spectrum. Because it is so faint, we fitted only for the power-law index and normalization, keeping the cutoff energy fixed at 2.5 GeV, a value typical of other MSPs (Abdo et al. 2010). The spectral model included both the isotropic and Galactic diffuse contributions with normalizations free in the fit. We detected the source with a TS of 38.2, corresponding to  $\sim 6\sigma$ . The spectral index and cutoff energy are very poorly constrained for this weak source, so to estimate the uncertainty on the integrated flux we repeated the spectral fit with the cutoff frozen at 0.7, 2.5, and 5.0 GeV, and with the index frozen at 1.3 (the average value for LAT-detected MSPs) and the cutoff free. We find that the uncertainty due to the spectral shape is about  $3 \times 10^{-10} \text{ cm}^{-2} \text{ s}^{-1}$  for the photon flux and  $2 \times 10^{-13} \text{ erg cm}^{-2} \text{ s}^{-1}$  for the energy flux. These are smaller than the statistical errors for this faint source and are added in quadrature to compute the errors on the fluxes reported in Table 1. Using the fitted energy flux ( $G$ ), and the DM distance ( $d$ ), we compute the  $\gamma$ -ray luminosity  $L_\gamma = 4\pi f_\Omega G d^2$  and efficiency  $\eta = L_\gamma / \dot{E}$ , where the beaming factor  $f_\Omega$  is assumed to be 1 (Waters et al. 2009).

Using this model, we selected events within  $2^\circ$  of the source and computed the probability that they originated from the source with the *Fermi* Science Tool `gtsrcprob`. Using these probabilities as weights and pulse phases computed using the radio timing model, we computed a weighted H-test of 37.1, corresponding to a detection significance of  $5.1 \sigma$  and confirming that J1544+4937 is indeed a  $\gamma$ -ray pulsar and identifying the LAT source as this MSP. The LAT weighted lightcurves in two energy bands, phase aligned to the radio profiles, are shown in Fig. 2. For this faint source, it is difficult to determine the peak multiplicity, but the  $\gamma$ -ray emission seems to be mainly between phases 0.4 and 0.6, relative to the single radio peak. Fitting a single gaussian peak to the light curve gives a peak at phase  $0.54 \pm 0.02$  with a width of  $0.25 \pm 0.05$ .

---

<sup>3</sup><http://fermi.gsfc.nasa.gov/ssc/data/analysis/software/>

<sup>4</sup><http://fermi.gsfc.nasa.gov/ssc/data/access/lat/BackgroundModels.html>

## 5. X-ray and optical observations

We analyzed 5 *Swift*/XRT observations (totaling 9.3 ks) of this field from 2013 March 19–24. No X-ray counterpart is detected, providing a flux upper limit of  $\sim 1.7 \times 10^{-14}$  ergs  $\text{cm}^{-2} \text{s}^{-1}$  for 0.3–10 keV.

We have checked for an optical counterpart for J1544+4937 using archival SDSS data. This field was imaged (Adelman-McCarthy et al. 2008) on MJD 52046.294, at binary phase 0.57. No optical counterpart was detected. The strongest upper limits were  $g' > 22.5$ ,  $r' > 22.0$  and  $i' > 21.5$  (90% CL). We also obtained a 300s H $\alpha$  image with the MiniMo camera on the 3.5m WIYN telescope on MJD 55975.521 (Brownsberger & Romani 2013). No stellar counterpart or extended emission was seen with an H $\alpha$  flux upper limit of  $2.5 \times 10^{-16}$  erg  $\text{cm}^{-2} \text{s}^{-1}$  (90% CL) for a compact  $< 5''$  nebula.

We compare these limits to fluxes measured for the archetypal, and best studied, BW B1957+20 (Reynolds et al. 2007; van Kerkwijk et al. 2011) which has a DM distance  $\sim 2.5$  kpc (with an estimated extinction  $A_V = 1.2$ ),  $P_b = 9.17$  hours and  $\dot{E} = 7.7 \times 10^{34}$  erg  $\text{s}^{-1}$ . At the orbital phase corresponding to J1544+4937’s upper limits (0.32 past eclipse) the studies above give extinction corrected magnitudes  $g' \sim 20.2$ ,  $r' \sim 19.7$  and  $i' \sim 19.4$ . Assuming a Roche lobe filling factor similar to that of the B1957+20 (since we do not know the true size of the companion), the ratio of reprocessed heating flux is just the ratio of spin-down luminosities, i.e. J1544+4937 is  $6.4 \times$  lower. Whereas, if we bring B1957+20 to 1.2 kpc, it would be  $4.3 \times$  brighter. The extinction correction for J1544+4937 is very small ( $A_V = 0.05$ ) in this direction. Corrected magnitudes of B1957+20, scaled to J1544+4937, would be 0.4mag fainter. Observed limits on J1544+4937 are  $\Delta g' = +1.8\text{mag}$ ,  $\Delta r' = +1.9\text{mag}$  and  $\Delta i' = +1.7\text{mag}$ , fainter than the scaled B1957+20 values. Thus the optical data imply  $\sim 2.3 \times$  larger distance of J1544+4937. Alternatively, the companion radius may be smaller by a similar factor or the companion temperature may be lower than the heating comparison above.

Our H $\alpha$  limit for J1544+4937 corresponds to  $< 0.15$  of the highest surface brightness  $\sim 5''$  patch of the B1957+20 bow shock. The average flux ratio is even more constraining, with an upper limit of  $\sim 0.03$  of the total bow-shock flux. However, given the small fraction of MSPs that show optical bow shocks (likely due to the small filling factor of the neutral interstellar medium), the non-detection at such a large distance from the Galactic plane ( $b \sim 50^\circ$ ) is not particularly surprising.



## 6. Eclipse characteristics

Fig. 3 presents the timing residuals (and electron column densities) around the eclipse phase at 322 and 607 MHz, with 90s time resolution. The effect of the eclipses is generally seen from 0.18 to 0.31 orbital phase (eclipse-zone hereafter) at 322 MHz. The eclipses are centered at binary phase 0.24 with a duration of around 22 minutes. We estimate the radius of the companion’s Roche lobe,  $R_L$  (Eggleton 1983),

$$R_L = \frac{0.49aq^{2/3}}{0.6q^{2/3} + \ln(1 + q^{1/3})} \sim 0.13R_\odot \quad (1)$$

where  $q = m_c/m_p$  is the mass ratio of the companion and the pulsar, and  $a$  is the separation of the companion from the pulsar ( $a \sim 1.2R_\odot$  for J1544+4937, indicating an extremely compact binary). The opaque portion of the companion’s orbit is  $0.98 R_\odot$ , much larger than  $R_L$  of  $0.13 R_\odot$ , so the volume occupied by the eclipsing material is well outside the companion’s Roche lobe, and thus is not gravitationally bound to the companion. This confirms that this binary is a BW, where the pulsar is ablating its companion, creating a significant amount of intrabinary material that obscures the pulsar’s emission.

Our sample consisted of six eclipses at 322 MHz where the pulsar emission was fully obscured by companion and its wind and two 607 MHz observing sessions covering the full orbit (Fig. 3). At 607 MHz we detect the MSP throughout the 322 MHz eclipse-zone. However, we observe a flux fading at 607 MHz near the superior conjunction (orbital phase 0.24). Significant delays in pulse arrival times are observed at 607 MHz during the eclipse-zone at 322 MHz. Maximum delay in pulse arrival time at 607 MHz near the eclipse superior conjunction is around  $300 \mu\text{s}$ , which corresponds to an increase in DM of  $0.027 \text{ pc cm}^{-3}$  and an added electron density,  $N_e$  of  $8 \times 10^{16} \text{ cm}^{-2}$ .

Thompson et al. (1994) (T94 hereafter) elucidate a collection of eclipse mechanisms. According to them, eclipsing due to refraction of the radio beam demands an order of magnitude higher group delay ( $\sim$  few tens of milliseconds) than we observe for J1544+4937 ( $250 \mu\text{s}$  at 322 MHz egress). Using  $N_e \sim 8 \times 10^{16} \text{ cm}^{-2}$  observed during superior conjunction at 607 MHz and absorption length  $L$  about twice the size of the eclipse-zone, according to equation 11 of T94 we find that free-free absorption is possible (absorption optical depth  $\tau_{ff} > 1$ ) if the plasma temperature  $T \leq 4 \times f_{cl}^{2/3} \text{ K}$ , where  $f_{cl}^{2/3}$  is the clumping factor. This demands either a very low temperature or a very high value of the clumping factor, both of which are not physically achievable. Eclipsing by pulse smearing (due to the increase of  $N_e$  along the line of sight) can be ruled out, as the excess electron column density inferred from 607 MHz predicts  $374 \mu\text{s}$  smearing of pulses at 322 MHz near superior conjunction (considering incoherent dedispersion), which is less than one fifth of pulse period. Since

J1544+4937 has a narrow main-pulse, and no significant profile evolution is apparent at the eclipse boundary, pulse broadening due to scattering can reduce the detectability but cannot explain the eclipse. In addition, since J1544+4937 is relatively weak, nearby and has a shallower spectrum than B1957+20, the expected induced Compton scattering optical depth is much less than one (equation 26 of T94). Another eclipse mechanism considered by T94 is cyclotron-synchrotron absorption of the radio waves by non-relativistic/relativistic electrons, which requires a magnetic field in the vicinity of the companion. We calculate a magnetic field  $B \sim 11$  G and corresponding cyclotron absorption frequency  $\sim 31$  MHz (equations 35, 37 of T94, assuming a moment-of-inertia of  $10^{45}$  g cm<sup>2</sup>). Thus 322 and 607 MHz will correspond to 10th and 20th harmonics of the cyclotron resonance. For a fixed temperature, the optical depth for cyclotron absorption drops with harmonics, which may explain the lack of absorption seen at 607 MHz. A modified model is proposed by Khechinashvili et al. (2000) based on kinematic treatment of cyclotron damping, assuming white-dwarf companions with reasonably strong surface magnetic fields. Further observations over a wider radio spectrum may help to investigate the frequency-dependent degree of damping predicted by this model.

We observe a temporal variation of the ingress phase at 322 MHz. For two eclipses, the ingress phase is shifted to 0.16 (from 0.18 for the other four eclipses), whereas there are no apparent shifts in the egress phase. This corresponds to an increase of the opaque portion by  $0.15 R_{\odot}$  and  $N_e > 1.2 \times 10^{16}$  cm<sup>-2</sup> at the ingress boundary. Such asymmetric increase of eclipse duration is unique to this BW and may indicate that our line of sight is probing a wind zone where there is systematic outflow of eclipse material.

We also observe strong phase modulations and additional short-duration absorptions at ingress and egress, in time-series data at higher resolution. The durations of these features are in general around 10–20 s, and hence they are not seen in Fig. 3. However, in one of the observing epochs these modulations lasted longer – phase modulation of duration 100 s, followed by a short-duration absorption of  $\sim 180$  s, then regular emission resumes for 500 s, after which the eclipse starts (Fig. 4). Such intriguing modulations, indicating fluctuation of plasma density around eclipsing zone, are not reported for other eclipsing BWs. Fragmented blobs of plasma randomly oriented around eclipsing zone, obscuring radiation from the pulsar, can explain these short-duration absorptions.

## 7. Discussion

We report the GMRT discovery of an eclipsing BW pulsar, at the position of an unassociated LAT source, FERMI J1544.2+4941. This is the first Galactic field MSP discovered at the GMRT. The detection of pulsed  $\gamma$ -rays from this pulsar demonstrates it as the source

powering FERMI J1544.2+4941. Due to the limited significance of the source in  $\gamma$ -rays additional data are needed before conclusions on peak multiplicity and system geometry can be drawn. The implied efficiency ( $\sim 3\%$ ) of converting spin-down energy into  $\gamma$ -rays is typical of LAT-detected MSPs. This is the first discovery of a radio MSP in a LAT source fainter than the 2FGL catalog limit (Ray et al. 2012). Since the radio pulsar is relatively bright, this provides strong justification to continue these searches as new LAT sources are revealed in analyses of longer datasets. The radio flux is uncorrelated with the  $\gamma$ -ray flux (Ackermann et al. 2012), so even faint new LAT sources can harbor bright radio MSPs.

By assuming a comparable companion Roche lobe filling factor and orbital inclination, we find that the irradiated flux/unit area should be  $0.73\times$  that of B1957+20, leading to an effective temperature of the heated side of  $T_{max} \sim 7700$  K. The flux limits from section 5 imply a distance  $> 2.7$  kpc. This would imply a larger  $\gamma$ -ray efficiency of  $> 15\%$ , which is still typical of LAT-detected MSPs. Deeper optical observations with multicolor light curves of the companion can be used to test these estimates.

Eclipsing BW pulsars have the potential of providing information on the evolutionary connection between the low-mass X-ray binaries and isolated MSPs. With long term monitoring of this pulsar we aim to estimate  $\dot{P}_b$  and its higher derivatives, which can provide an estimate of the life span of the system. Bates et al. (2011) noted that for BW systems the measured value of  $\dot{E}/a^2$  is an order of magnitude higher than for other MSP binaries, indicative of greater energy flux needed to ablate the companion. For J1544+4937 we calculate  $\dot{E}/a^2 \sim 1.5 \times 10^{33}$  erg lt-s $^{-2}$ s $^{-1}$ , which is similar to other BW systems in the Galactic field. Dual frequency observations presented in this paper suggest that cyclotron absorption by the plasma formed via interaction of the pulsar wind with ablated material can obscure the pulsed emission. However, exploring the radio spectrum on either side to probe the reduced/increased opaqueness of the stellar wind during the eclipse phase may provide better insight into the plausible eclipse mechanism.

The *Fermi* LAT Collaboration acknowledges support from a number of agencies and institutes for both development and the operation of the LAT as well as scientific data analysis. These include NASA and DOE in the United States, CEA/Irfu and IN2P3/CNRS in France, ASI and INFN in Italy, MEXT, KEK, and JAXA in Japan, and the K. A. Wallenberg Foundation, the Swedish Research Council and the National Space Board in Sweden. Additional support from INAF in Italy and CNES in France for science analysis during the operations phase is also gratefully acknowledged. We acknowledge support of telescope operators of the GMRT, which is run by the National Centre for Radio Astrophysics of the Tata Institute of Fundamental Research. We thank the *Swift* team at Pennsylvania State University, especially Abe Falcone. We acknowledge help of C. Cheung in interpreting the

XRT data.

## REFERENCES

- Abdo, A. A. et al. 2010, *ApJS*, 187, 460.
- Ackermann, M. et al. 2012, *ApJ*, 753, 83
- Adelman-McCarthy, J. et al. 2008, *ApJS*, 175, 297.
- Atwood, W. B. et al. 2009, *ApJ*, 697, 1071.
- Bates, S. D. et al. 2011, *MNRAS*, 416, 2455.
- Bhattacharyya, B., Nityananda, R. 2008, *MNRAS*, 387, 273.
- Brownsberger, S., Romani, R. W. 2013, *ApJ* in prep.
- Cordes, J. M., Lazio, T. J. W. 2002, *arXiv:astro-ph/0207156*.
- Eggleton, P. P. 1983, *ApJ*, 268, 368.
- Fruchter, A. S., Stinebring, D. R., Taylor, J. H. 1988, *Nature*, 333, 237.
- Kerr, M. 2011, *ApJ*, 732, 38
- Khechinashvili, D. G., Melikidze, G. I., Gil, J. I. 2000, *ApJ*, 541, 335.
- Lange, C. et al. 2001, *MNRAS*, 326, 274.
- Mattox, J. R. et al. 1996, *ApJ*, 461, 396.
- Roberts, M. S. E. 2011, *AIP Conf. Proc.* 1357, 40.
- Nolan, P. L. et al. 2012, *ApJS*, 199, 31
- Ransom, S. M., Eikenberry, S. S., Middleditch, J. 2002, *AJ*, 124, 1788.
- Ray, P. S. et al., 2012, *2011 Fermi Symposium Proceedings (eConf C110509)*, *arXiv:1205.3089*.
- Reynolds, M. P. J. et al. 2007, *MNRAS*, 379, 1117.
- Roy, J. et al. 2010, *Experimental Astronomy*, 28, 55.

- Roy, J., Bhattacharyya, B., Gupta, Y., 2012, MNRAS Letters, 27, 90.
- Stappers, B. W. et al. 1996, ApJ, 465, 119.
- Stappers, B. W. et al. 2001, MNRAS, 321, 576.
- Standish, E. M. 2004, A&A, 417, 1165.
- Thompson, C., Blandford, R. D., Evans, C. R., Phinney, E. S. 1994, ApJ, 422, 304.
- van Kerkwijk, M. H., Breton, R. P., Kulkarni, S. R. 2011, ApJ, 728, 95.
- Watters, K. P., Romani, R. W., Weltevrede, P., Johnston, S. 2009, ApJ, 695, 1289.




Fig. 3.— Variation of timing residuals and electron column density (TOAs of 90 s time resolution) around eclipse phase at 322 MHz (top) and 607 MHz (bottom).




Fig. 4.— Example of additional short-duration absorption beyond the eclipse phase at 322 MHz on 2012 June 3. Left: variation of timing residuals and electron column density at 42 s time resolution. Right: phaseogram showing short-duration absorption.