

PROVING THE CEP WITH COMPACT STARS?*

D.E. Alvarez-Castillo^{1,2†} and D. Blaschke^{1,3,4‡}

¹ Bogoliubov Laboratory of Theoretical Physics, JINR Dubna, 141980 Dubna, Russia

² Instituto de Física, Universidad Autónoma de San Luis Potosí, S.L.P. 78290, México

³ Institute for Theoretical Physics, University of Wrocław, 50-204 Wrocław, Poland

⁴ Fakultät für Physik, Universität Bielefeld, 33501 Bielefeld, Germany

ANNOTATION

We present a model for hybrid compact stars composed of a quark core and a hadronic mantle with an abrupt first order phase transition at the interface and in accordance with the latest astrophysical measurements of two $2 M_{\odot}$ pulsars. We demonstrate the possibility of a disconnected mass-radius sequence (third family) of high-mass pulsars as a distinct feature due to a large jump $\Delta\epsilon$ in the energy density of the first order phase transition setting in at $\epsilon_{\text{crit}} \approx 500 \text{ MeV}/\text{fm}^3$ and fulfilling $\Delta\epsilon/\epsilon_{\text{crit}} > 0.6$. We conclude that the measurement of so called *twin* compact stars at high mass ($\sim 2 M_{\odot}$) would support the existence of a first order phase transition in symmetric matter at zero temperature entailing the existence of a critical end point (CEP) in the QCD phase diagram.

INTRODUCTION

Recently the study of neutron stars has become an ever more active field of research since it complements the investigations of matter under extreme conditions in heavy-ion collisions and in ab-initio lattice QCD simulations. The interiors of compact stars are unique places to investigate the otherwise inaccessible cold, dense equation of state (EoS) which determines the shape of of stable compact star sequence(s) in the mass - radius diagram as well as other properties like the composition, the moment of inertia etc. In this regard, recently the masses of two heavy compact stars have been precisely measured: $1.97 \pm 0.04 M_{\odot}$ for PSR J1614-2230 [1] and $2.01 \pm 0.04 M_{\odot}$ for PSR J0348+0432 [2], presenting a stringent constraint on models for the high density EoS. Would it be possible to gain

*Work supported by CompStar, a research networking programme of the European Science foundation and by CompStar-POL a project of the Polish MNiSW supporting it as well as by NCN under grant number DEC-2011/02/A/ST2/00306 and by RFBR under grant number 11-02-01538a.

†alvarez@theor.jinr.ru

‡blaschke@ift.uni.wroc.pl

evidence for a strong first order deconfinement transition in compact stars this would prove the existence of at least one CEP in the QCD phase diagram since in the high temperature region explored by lattice QCD simulations [3] the transition is crossover.

MASSIVE HYBRID STARS & TWINS

A first order phase transition in neutron star matter can take place just as in symmetric matter where it is searched for in heavy ion collisions. Adopting the setting of [4, 6], we construct hybrid stars with a hybrid EoS composed of a given hadronic EoS, here DD2 [5], and a quark matter EoS parametrized by its squared speed of sound c_{QM}^2 which pretty well describes [6] results of a color superconducting NJL model [7]

$$P(\varepsilon) = P_{\text{DD2}}(\varepsilon)\Theta(\varepsilon_{\text{crit}} - \varepsilon) + c_{\text{QM}}^2 \varepsilon \Theta(\varepsilon - \varepsilon_{\text{crit}} - \Delta\varepsilon). \quad (1)$$

The critical energy density $\varepsilon_{\text{crit}}$ and the discontinuity $\Delta\varepsilon$ complete the EoS model which is capable of describing compact star sequences with a third family of stars in the mass-radius diagram [8, 9, 10, 11]. Searching for sequences obeying the mass constraint [1, 2] we obtain a quasi-horizontal branch disconnected from the almost vertical neutron star branch, as a consequence of a strong phase transition. Fig. 1 shows the EoS (1) for the parameters: $c_{\text{QM}}^2 = 0.94$, $\Delta\varepsilon = 0.67 \varepsilon_{\text{crit}}$ and $\varepsilon_{\text{crit}} = 485 \text{ MeV/fm}^3$. The latter corresponds to $P(\varepsilon_{\text{crit}}) = 100 \text{ MeV/fm}^3$ and a baryon density at the quark matter onset of $n_{\text{crit}} = 2.9 n_0$ with $n_0 = 0.16 \text{ fm}^{-3}$. Fig. 2 shows the mass-radius relation for this hybrid EoS and the energy density profile for the example of a twin pair of $2 M_{\odot}$ stars is given in Fig. 1 (right).

CONCLUSIONS

In this work we have demonstrated for a simple hybrid star EoS model that the characteristic mass-radius relationship of nonrotating compact star configurations obtained by solving the TOV equations has a third family of stable hybrid stars at high masses that is in accordance with astrophysical observations. This includes in particular the constraint for the minimal radius from RXJ 1856-3754 and the new pulsar mass measurements for PSR J1614-2230 and PSR J0348+0432 at about $2 M_{\odot}$. Our result shows that stars of this high mass which are the best candidates for containing exotic matter (like quark matter) in their interiors could actually be approximate mass twins: although having about the same mass their interiors may show a vastly different composition: while the lower density twin has a larger radius ($\sim 13 \text{ km}$) and is made of nuclear matter only, the high-density twin




Figure 1. Left panel: Hadronic EoS (DD2, green dashed line) vs. hybrid one (Eq. 1, indigo solid line). Right panel: Energy density profile of $2 M_{\odot}$ twins: a hadronic (green dashed line) and a hybrid (indigo solid line) compact star.

is more compact ($\sim 10.5 - 12$ km) with a phase transition at about half the stars radius. The precise determination of compact star radii would allow to prove or disprove whether the sequences of stable stars in the mass-radius diagram form a pattern like the one discussed here: with an almost vertical branch of hadronic stars and an almost horizontal branch of hybrid stars, possibly disconnected from the former thus representing a “third family” of compact stars and allowing for high-mass star twins. Identifying this pattern by observation would indicate that compact star matter undergoes a phase transformation with a large jump in energy density that could correspond to a strong first order transition in symmetric matter. Given the knowledge from lattice QCD that at zero baryon density the QCD phase transition proceeds as a crossover, twins would then support the existence of a CEP in the QCD phase diagram, see also [13] for a microphysical study.

ACKNOWLEDGEMENTS

We gratefully acknowledge numerous discussions and collaboration work on the topic addressed in this contribution with our colleagues, in particular with S. Benic, G. Contrera, H. Grigorian, O. Kaczmarek, T. Klähn, E. Laermann, R. Łastowiecki, M.C. Miller, G. Poghosyan, S.B. Popov, J. Trümper, D.N. Voskresensky and F. Weber.




Figure 2. Mass-radius diagram for hybrid star EoS (1). The almost vertical branch of pure hadronic stars (green solid line) is separated from the almost horizontal hybrid star branch (indigo solid line) by an unstable branch (black dashed line). Recently measured masses for two $2 M_{\odot}$ pulsars shown by red and blue hatched regions. Also shown: causality constraint (upper left forbidden area) and the minimal radius constraint from RXJ 1856-3754 [12].

REFERENCES

- [1] P. Demorest *et al.*, *Nature* **467**, 1081 (2010).
- [2] J. Antoniadis *et al.*, *Science* **340**, 6131 (2013).
- [3] O. Kaczmarek *et al.*, *Phys. Rev. D* **83**, 014504 (2011).
- [4] M. G. Alford, S. Han and M. Prakash, arXiv:1302.4732 [astro-ph.SR].
- [5] S. Typel and H. H. Wolter, *Nucl. Phys. A* **656**, 331 (1999).
- [6] J. L. Zdunik and P. Haensel, arXiv:1211.1231 [astro-ph.SR].
- [7] R. Lastowiecki *et al.*, *Acta Phys. Polon. Supp.* **5**, 535 (2012) [arXiv:1112.6430 [nucl-th]].
- [8] U. H. Gerlach, *Phys. Rev.* **172**, 1325 (1968).
- [9] B. Kämpfer, *J. Phys. A* **14**, L471 (1981).
- [10] K. Schertler *et al.*, *Nucl. Phys. A* **677**, 463 (2000).
- [11] N. K. Glendenning and C. Kettner, *Astron. Astrophys.* **353**, 795 (2000).
- [12] J. E. Trümper *et al.*, *Nucl. Phys. Proc. Suppl.* **132**, 560 (2004).
- [13] D. Blaschke *et al.*, arXiv:1302.6275 [hep-ph].