

Exact Green's function of the reversible ABCD reaction in two space dimensions

Thorsten Prüstel and Martin Meier-Schellersheim

Laboratory of Systems Biology
National Institute of Allergy and Infectious Diseases
National Institutes of Health


June 10, 2021

Abstract

We derive an exact expression for the Green's functions in the time domain of the reversible diffusion-influenced ABCD reaction $A + B \leftrightarrow C + D$ in two space dimensions. Furthermore, we calculate the corresponding survival and reaction probabilities. The obtained expressions should prove useful for the study of reversible membrane-bound reactions in cell biology and can serve as a useful ingredient of enhanced stochastic particle-based simulation algorithms.

1 Introduction

The reversible so-called ABCD reaction refers to the following reaction schemes [7]



In particular, it represents the enzymatic reaction



*Email: prustelt@niaid.nih.gov, mms@niaid.nih.gov

where the enzyme E converts a substrate S into a product. Note that the reaction of Eq. (1.4) is the limiting case of



under the assumption that the concentration of the intermediate complex does not change on the timescale of the association/dissociation reactions.

Exact analytical expressions for the Green's functions (GF) of the reversible ABCD reaction for an isolated pair have been derived for 1, 2 and 3 dimensions (3D) in the Laplace domain and for 1 and 3 dimensions in the time domain [7]. However, an exact solution valid for all times in 2D is still lacking despite both its broad applicability to membrane phenomena and its conceptual importance. In fact, in the context of cell biological applications, the 2D case provides the basis for a better understanding of processes such as signal-induced inhomogeneities and receptor clustering on cell membranes [2], where diffusion constants that govern the lateral motion of receptors are typically small. Furthermore, from a conceptual point of view, diffusion in two dimensions is special, because 2D is the critical dimension with regard to recurrence and transience of random walks [10].

Moreover, GFs for an isolated pair play an important role in a number of particle-based stochastic simulation algorithms, because any biochemical reaction network may be thought of as composed of unimolecular and bimolecular reactions. In this context, analytical expressions for GFs describing an isolated pair can be employed to enhance the efficiency and accuracy of Brownian dynamics simulations [4, 6, 8, 11].

2 Theory

In this section we will closely follow Ref. [7]. The two different possible states of the ABCD reaction require considering two different probability density functions: $p_1(r_1, t|\xi_1)$ yields the probability to find an A, B pair at a distance equal to r_1 at time t , given that it was initially separated by a distance equal to ξ_1 , and $p_2(r_2, t|\xi_1)$ provides the corresponding probability for the C, D pair, given that initially an A, B pair was separated by a distance equal to ξ_1 . The relative diffusion constants in the two states are denoted by D_1 and D_2 , respectively. When the distance equals the reaction distance $a_1(a_2)$, the pair A, B (C, D) may undergo a reaction and be converted to C, D (A, B). The transitions from one state to the other are characterized by the intrinsic rate constants κ_1 and κ_2 , respectively. The time evolution of the

probability densities is governed by the two equations [7]

$$\begin{aligned} \frac{\partial p_i(r_i, t|\xi_1)}{\partial t} &= D_i \left(\frac{\partial^2}{\partial r_i^2} + \frac{1}{r_i} \frac{\partial}{\partial r_i} \right) p_i(r_i, t|\xi_1) - \kappa_i \frac{\delta(r_i - a_i)}{2\pi r_i} p_i(r_i, t|\xi_1) \\ &+ \kappa_j \frac{\delta(r_i - a_i)}{2\pi r_i} p_j(a_j, t|\xi_1), \quad \text{where } i, j = 1, 2 \text{ and } i \neq j. \end{aligned} \quad (2.1)$$

The Eqs. (2.1) have to be supplemented by an appropriate initial condition. As already mentioned, we assume, without loss of generality, that the initial state at time $t = 0$ is given by an A, B pair separated by a distance equal to ξ_1 . Hence, the initial conditions read

$$p_1(r_1, t = 0|\xi_1) = \frac{\delta(r_1 - \xi_1)}{2\pi r_1}, \quad (2.2)$$

$$p_2(r_2, t = 0|\xi_1) = 0. \quad (2.3)$$

In the following we are interested in the GF $p_1(r_1, t|\xi_1)$ and $p_2(r_2, t|\xi_1)$ that satisfy the initial value problem posed by Eqs. (2.1), (2.2), (2.3).

Employing the Laplace transform

$$\tilde{p}_i(r_i, s|\xi_1) = \int_0^\infty e^{-st} p_i(r_i, t|\xi_1) dt, \quad i = 1, 2, \quad (2.4)$$

Popov and Agmon [7] obtained explicit expressions for the GF in the Laplace domain

$$\begin{aligned} \tilde{p}_1(r_1, s|\xi_1) &= \tilde{G}_{1,\text{ref}}(r_1, s|\xi_1) \\ &- \frac{\kappa_1 \tilde{G}_{1,\text{ref}}(r_1, s|a_1) \tilde{G}_{1,\text{ref}}(a_1, s|\xi_1)}{1 + \kappa_1 \tilde{G}_{1,\text{ref}}(a_1, s|a_1) + \kappa_2 \tilde{G}_{2,\text{ref}}(a_2, s|a_2)}, \end{aligned} \quad (2.5)$$

$$\tilde{p}_2(r_2, s|\xi_1) = \frac{\kappa_1 \tilde{G}_{2,\text{ref}}(r_2, s|a_2) \tilde{G}_{1,\text{ref}}(a_1, s|\xi_1)}{1 + \kappa_1 \tilde{G}_{1,\text{ref}}(a_1, s|a_1) + \kappa_2 \tilde{G}_{2,\text{ref}}(a_2, s|a_2)}, \quad (2.6)$$

where $\tilde{G}_{i,\text{ref}}(r, s|\xi)$ denotes the Laplace transform of the non-reactive GF that satisfies at the encounter distance a reflective boundary condition

$$\left. \frac{\partial G_{i,\text{ref}}(r, s|\xi)}{\partial r} \right|_{r=a_i} = 0, \quad i = 1, 2. \quad (2.7)$$

Clearly, the probability density of a non-reactive isolated pair has to satisfy

$$2\pi \int_{a_i}^\infty \tilde{G}_{i,\text{ref}}(r_i, s|a_i) r_i dr_i = \frac{1}{s}, \quad i = 1, 2. \quad (2.8)$$

Due to this property, the survival or reaction probabilities

$$\tilde{S}_i(s|\xi_1) = 2\pi \int_{a_i}^\infty \tilde{p}_i(r_i, s|\xi_1) r_i dr_i \quad (2.9)$$

can easily be obtained from Eqs. (2.5), (2.6) [7]

$$s\tilde{S}_1(s|\xi_1) = 1 - \frac{\kappa_1\tilde{G}_{1,\text{ref}}(a_1, s|\xi_1)}{1 + \kappa_1\tilde{G}_{1,\text{ref}}(a_1, s|a_1) + \kappa_2\tilde{G}_{2,\text{ref}}(a_2, s|a_2)}, \quad (2.10)$$

$$s\tilde{S}_2(s|\xi_1) = \frac{\kappa_1\tilde{G}_{1,\text{ref}}(a_1, s|\xi_1)}{1 + \kappa_1\tilde{G}_{1,\text{ref}}(a_1, s|a_1) + \kappa_2\tilde{G}_{2,\text{ref}}(a_2, s|a_2)}. \quad (2.11)$$

It follows immediately from Eqs. (2.10), (2.11) that $\tilde{S}_1(s|\xi_1) + \tilde{S}_2(s|\xi_1) = 1/s$, which translates to

$$S_1(t|\xi_1) + S_2(t|\xi_1) = 1, \quad (2.12)$$

in the time domain, as it should be.

3 Explicit form of the Green's functions and survival probabilities

Eqs. (2.5), (2.6) permit obtaining $\tilde{p}_i(r_i, s|\xi_1)$ once the explicit form of the non-reactive GF in the Laplace domain is known. In 2D, it is given by [7]

$$\tilde{G}_{i,\text{ref}}(r_i, s|\xi_i) = \tilde{G}_{i,\text{free}}(r_i, s|\xi_i) + \frac{1}{2\pi D_i} K_0(q_i r_i) K_0(q_i \xi_i) \frac{I_1(q_i a_i)}{K_1(q_i a_i)}. \quad (3.1)$$

Here, I_n, K_n refer to the modified Bessel function of first and second kind, respectively [1, Sect. 9.6] and $\tilde{G}_{i,\text{free}}(r, s|\xi)$ denotes the free-space GF [3, Ch. 14.8, Eq. (2)]

$$\tilde{G}_{i,\text{free}}(r_i, s|\xi_i) = \frac{1}{2\pi D_i} \begin{cases} I_0(q_i \xi_i) K_0(q_i r_i), & r_i > \xi_i \\ I_0(q_i r_i) K_0(q_i \xi_i), & r_i < \xi_i \end{cases} \quad (3.2)$$

and we define $q_i = \sqrt{s/D_i}$ for $i = 1, 2$. Combining Eqs. (2.6), (2.11) with Eqs. (3.1), (3.2) one arrives at

$$\tilde{p}_2(r_2, t|\xi_1) = \frac{h_1}{2\pi D_2 a_2} \frac{K_0(q_2 r_2) K_0(q_1 \xi_1)}{q_2 K_1(q_2 r_2)} \frac{1}{\mathcal{D}}, \quad (3.3)$$

$$\tilde{S}_2(t|\xi_1) = \frac{h_1}{s} \frac{K_0(q_1 \xi_1)}{\mathcal{D}}, \quad (3.4)$$

where the denominator \mathcal{D} is given by

$$\mathcal{D} = q_1 K_1(q_1 a_1) \left[1 + h_2 \frac{K_0(q_2 a_2)}{q_2 K_1(q_2 a_2)} \right] + h_1 K_0(q_1 a_1). \quad (3.5)$$

Here, we introduced

$$h_i = \kappa_i / (2\pi D_i a_i), \quad i = 1, 2. \quad (3.6)$$

In the following we employ the inversion theorem for the Laplace transformation to obtain the corresponding GF and survival probabilities in the time domain according to

$$p_i(r_i, t|\xi_1) = \frac{1}{2\pi i} \int_{\gamma-i\infty}^{\gamma+i\infty} e^{st} \tilde{p}_i(r_i, s|\xi_1) ds, \quad (3.7)$$

$$S_i(t|\xi_1) = \frac{1}{2\pi i} \int_{\gamma-i\infty}^{\gamma+i\infty} e^{st} \tilde{S}_i(s|\xi_1) ds. \quad (3.8)$$

It is convenient to consider $p_2(r_2, t|\xi_1)$ first. We proceed analogously to the calculations presented in Ref. [9], where the GF of the reversible diffusion-influenced reaction $A + B \rightleftharpoons C$ for an isolated pair in two space dimensions was obtained. In particular, to calculate the Bromwich contour integral, we note that $\tilde{p}_2(r_2, t|\xi_1)$ is multi-valued and has a branch point at $s = 0$. Therefore, we consider the contour of Fig. 1 with a branch cut along the negative real axis, cp. [3, Ch. 12.3, Fig. 40]. In addition, we note that the integrand has no poles within and on the contour [5] and the contribution from the small circle around the origin vanishes. Thus, the only contributions come from the integration above and below the branch cut

$$\begin{aligned} & \int_{\gamma-i\infty}^{\gamma+i\infty} e^{st} \tilde{p}_2(r_2, s|\xi_1) ds = \\ & - \int_{\mathcal{C}_2} e^{st} \tilde{p}_2(r_2, s|\xi_1) ds - \int_{\mathcal{C}_4} e^{st} \tilde{p}_2(r_2, s|\xi_1) ds. \end{aligned} \quad (3.9)$$

To calculate the integrals $\int_{\mathcal{C}_2}, \int_{\mathcal{C}_4}$, we choose, without loss of generality, $s = D_1 x^2 e^{i\pi}$ and use [3, Append. 3, Eqs. (25), (26)]

$$I_n(xe^{\pm\pi i/2}) = e^{\pm n\pi i/2} J_n(x), \quad (3.10)$$

$$K_n(xe^{\pm\pi i/2}) = \pm \frac{1}{2} \pi i e^{\mp n\pi i/2} [-J_n(x) \pm iY_n(x)]. \quad (3.11)$$

$J_n(x), Y_n(x)$ denote the Bessel functions of first and second kind, respectively [1, Sect. 9.1]. It follows that

$$\begin{aligned} \int_{\mathcal{C}_2} e^{st} \tilde{p}_2(r_2, s|\xi_1) ds &= \frac{h_1}{\pi a_2} \sqrt{\frac{D_1}{D_2}} \int_0^\infty e^{-D_1 x^2 t} \times \\ & \frac{[\Omega(r_2 \sqrt{D_1/D_2}, \xi_1) - i\Pi(r_2 \sqrt{D_1/D_2}, \xi_1)] [-Y_1(x\phi) + iJ_1(x\phi)] [\beta - i\alpha]}{[\alpha^2 + \beta^2] [J_1^2(x\phi) + Y_1^2(x\phi)]} dx. \end{aligned} \quad (3.12)$$

Here, we introduced a fair amount of new notation. First, we have

$$\alpha = \alpha_{\text{rad}} + \alpha_b, \quad (3.13)$$

$$\beta = \beta_{\text{rad}} + \beta_b, \quad (3.14)$$

where $\alpha_{\text{rad}}, \beta_{\text{rad}}$ refer to the functions that appear in the expression of the GF that describes irreversible association of an isolated pair in 2D, i.e. a GF that satisfies a radiation boundary condition at contact a_1 [3, Chap. 14.8, Eqs. (12), (13)]

$$p_{\text{rad}}(r, t|\xi) = \frac{1}{2\pi} \int_0^\infty e^{-D_1 t x^2} T_{\text{rad}}(x, r) T_{\text{rad}}(x, \xi) x dx, \quad (3.15)$$

where

$$T_{\text{rad}}(x, r) = \frac{J_0(rx)\beta_{\text{rad}} - Y_0(rx)\alpha_{\text{rad}}}{[\alpha_{\text{rad}}^2 + \beta_{\text{rad}}^2]^{1/2}}, \quad (3.16)$$

and

$$\alpha_{\text{rad}} = xJ_1(xa_1) + h_1J_0(xa_1), \quad (3.17)$$

$$\beta_{\text{rad}} = xY_1(xa_1) + h_1Y_0(xa_1). \quad (3.18)$$

Furthermore, we defined

$$\alpha_b = \sigma Y_1(xa_1) + \rho J_1(xa_1), \quad (3.19)$$

$$\beta_b = \rho Y_1(xa_1) - \sigma J_1(xa_1), \quad (3.20)$$

where

$$\rho = \psi(Y_0(x\phi)Y_1(x\phi) + J_0(x\phi)J_1(x\phi)), \quad (3.21)$$

$$\sigma = -\psi \frac{2}{\pi x \phi}, \quad (3.22)$$

and

$$\psi = h_2 \sqrt{\frac{D_2}{D_1}} \frac{1}{[J_1(x\phi)^2 + Y_1(x\phi)^2]}, \quad \phi = \sqrt{\frac{D_1}{D_2}} a_2. \quad (3.23)$$

Finally, we introduced

$$\Omega(r, \xi) = J_0(xr)J_0(x\xi) - Y_0(xr)Y_0(x\xi), \quad (3.24)$$

$$\Pi(r, \xi) = Y_0(xr)J_0(x\xi) + J_0(xr)Y_0(x\xi). \quad (3.25)$$

Next, we consider the integral along the contour \mathcal{C}_4 below the branch cut. It can be evaluated by choosing $s = D_1 x^2 e^{-i\pi}$ and after a calculation analogous to the evaluation of $\int_{\mathcal{C}_2}$ we obtain

$$\int_{\mathcal{C}_2} e^{ps} \tilde{p}_2(r_2, s|\xi_1) ds = - \left(\int_{\mathcal{C}_4} e^{st} \tilde{p}_2(r_2, s|\xi_1) ds \right)^*, \quad (3.26)$$

where $*$ means complex conjugation. Thus, the exact Green's function in the time domain is given by

$$\begin{aligned}
p_2(r_2, t|\xi_1) &= -\frac{1}{\pi} \Im \left(\int_{\mathcal{C}_2} e^{st} \tilde{p}_2(r_2, s|\xi_1) ds \right) \\
&= -\frac{h_1}{\pi^2 a_2} \sqrt{\frac{D_1}{D_2}} \int_0^\infty e^{-D_1 x^2 t} \times \\
&\quad \left\{ \frac{\alpha [Y_1(x\phi)\Omega(r_2\sqrt{D_1/D_2}, \xi_1) - J_1(x\phi)\Pi(r_2\sqrt{D_1/D_2}, \xi_1)]}{[\alpha^2 + \beta^2][J_1^2(x\phi) + Y_1^2(x\phi)]} \right. \\
&\quad \left. + \frac{\beta [J_1(x\phi)\Omega(r_2\sqrt{D_1/D_2}, \xi_1) + Y_1(x\phi)\Pi(r_2\sqrt{D_1/D_2}, \xi_1)]}{[\alpha^2 + \beta^2][J_1^2(x\phi) + Y_1^2(x\phi)]} \right\} dx. \quad (3.27)
\end{aligned}$$

To obtain the corresponding reaction probability $S_2(t|\xi_1)$, we can proceed analogously to the calculation of the GF $p_2(r_2, t|\xi_1)$, in particular, we can employ the same integration contour. There is one important difference, though. In the case of the survival probability, the contribution from the integration along the small circle around the origin does not vanish. To calculate this contribution we use $s = D_1 x^2 e^{i\varphi}$ and integrate from $-\pi$ to π . Using Eq. (2.11), (3.5) and the small argument expansions of the modified Bessel functions [1, Eqs. (9.6.7)-(9.6.9)], we arrive at

$$\begin{aligned}
&2\pi i \text{ contribution from the integral around the origin} \\
&= -i \lim_{x \rightarrow 0} \int_\pi^{-\pi} \frac{h_1 K_0(q_1 \xi_1)}{\mathcal{D}} d\varphi = -i \lim_{x \rightarrow 0} \int_\pi^{-\pi} \frac{h_1 \ln x}{h_1 \ln x + h_2 a_2 / a_1 \ln x} d\varphi \\
&= 2\pi i \frac{h_1 a_1}{h_1 a_1 + h_2 a_2}. \quad (3.28)
\end{aligned}$$

Taking into account the contributions from the integral above and below the branch cut and around the origin, we finally obtain

$$\begin{aligned}
S_2(t|\xi_1) &= \frac{h_1 a_1}{h_1 a_1 + h_2 a_2} \\
&\quad - \frac{2h_1}{\pi} \int_0^\infty e^{-D_1 t x^2} \frac{\alpha Y_0(x\xi_1) - \beta J_0(x\xi_1)}{\alpha^2 + \beta^2} \frac{dx}{x}. \quad (3.29)
\end{aligned}$$

In particular, we get for the steady-state values of $S_2(t|\xi_1)$ and of $S_1(t|\xi_1)$ (due to Eq. (2.12))

$$\lim_{t \rightarrow \infty} S_1(t|\xi_1) = \frac{h_2 a_2}{h_1 a_1 + h_2 a_2}, \quad (3.30)$$

$$\lim_{t \rightarrow \infty} S_2(t|\xi_1) = \frac{h_1 a_1}{h_1 a_1 + h_2 a_2}, \quad (3.31)$$

in agreement with Ref. [7]. Also, the explicit form of $S_1(t|\xi_1)$ follows immediately from Eqs. (3.29), (2.12).

It remains to calculate the GF $p_1(r_1, t|\xi_1)$. We know from Eq. (2.5) that it can be written as

$$p_1(r_1, t|\xi_1) = G_{1,\text{ref}}(r_1, t|\xi_1) + p_{1bc}(r_1, t|\xi_1), \quad (3.32)$$

where the explicit form of $G_{1,\text{ref}}(r_1, t|\xi_1)$ is well-known and can be recovered from Eq. (3.15) in the limit $h_1 \rightarrow 0$. $p_{1bc}(r_1, t|\xi_1)$ can easily be obtained from Eq. (3.27), exploiting the similarity of $\tilde{p}_{1bc}(r_1, s|\xi_1)$ and $\tilde{p}_2(r_2, s|\xi_1)$, cp. Eqs. (2.5), (2.6). Thus, we obtain

$$p_{1bc}(r_1, t|\xi_1) = \frac{h_1}{\pi^2 a_1} \int_0^\infty e^{-D_1 x^2 t} \times \left\{ \frac{\alpha[Y_1(xa_1)\Omega(r_1, \xi_1) - J_1(xa_1)\Pi(r_1, \xi_1)]}{[\alpha^2 + \beta^2][J_1^2(xa_1) + Y_1^2(xa_1)]} + \frac{\beta[J_1(xa_1)\Omega(r_1, \xi_1) + Y_1(xa_1)\Pi(r_1, \xi_1)]}{[\alpha^2 + \beta^2][J_1^2(xa_1) + Y_1^2(xa_1)]} \right\} dx. \quad (3.33)$$

Note that in the limit $\kappa_2 \rightarrow 0$ one recovers from Eqs. (3.32),(3.33) the known GF that satisfies a radiation BC Eq. (3.15) and describes the irreversible association $A + B \rightarrow C$.

Acknowledgments

This research was supported by the Intramural Research Program of the NIH, National Institute of Allergy and Infectious Diseases.




Figure 1: Integration contour used in Eq. (3.9).

References

- [1] M. Abramowitz and I.A. Stegun. *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*. Dover, New York, 1965.
- [2] I. Bethani, S.S. Skanland, I. Dikic, and A. Acker-Palmer. *EMBO J.*, 29:2677, 2010.
- [3] H.S. Carslaw and J.C. Jaeger. *Conduction of Heat in Solids*. Clarendon Press, New York, 1986.
- [4] A.L. Edelstein and N. Agmon. *J. Chem. Phys.*, 99:5396, 1993.
- [5] A. Erdelyi and W.O. Kermack. *Proc. Camb. Phil. Soc.*, 41:74, 1945.
- [6] H. Kim, M. Yang, and K.J. Shin. *J. Chem. Phys.*, 111:1068, 1999.
- [7] A.V. Popov and N. Agmon. *J. Chem. Phys.*, 117:5770, 2002.
- [8] A.V. Popov and N. Agmon. *J. Chem. Phys.*, 118:11057, 2003.
- [9] T. Prüstel and M. Meier-Schellersheim. *J. Chem. Phys.*, 137:054104, 2012.
- [10] D. Toussaint and F. Wilczek. *J. Chem. Phys.*, 78:2642, 1983.
- [11] J.S. van Zon and P.R. ten Wolde. *Phys. Rev. Lett.*, 94:128103, 2005.