

# Bayesian Test for Colocalisation Between Pairs of Genetic Association Studies Using Summary Statistics

Claudia Giambartolomei<sup>1,\*</sup>, Damjan Vukcevic<sup>2</sup>, Eric E. Schadt<sup>3</sup>, Aroon D. Hingorani<sup>4</sup>, Chris Wallace<sup>5</sup>, Vincent Plagnol<sup>1</sup>

**1 UCL Genetics Institute, University College London (UCL), Darwin Building, Gower Street, London WC1E 6BT, UK**

**2 Murdoch Childrens Research Institute, Royal Children’s Hospital, Melbourne, Australia**

**3 Department of Genetics and Genomics Sciences, Mount Sinai School of Medicine, New York, New York, United States of America**

**4 Institute of Cardiovascular Science, University College London, London WC1E 6BT, UK**

**5 Juvenile Diabetes Research Foundation/Wellcome Trust Diabetes and Inflammation Laboratory, Cambridge Institute for Medical Research, Department of Medical Genetics, NIHR Biomedical Research Centre, University of Cambridge, Addenbrookes Hospital, Hills Road, Cambridge, CB2 0XY, UK**

**\* E-mail: claudia.giambartolomei.10@ucl.ac.uk**

## Abstract

Genetic association studies, in particular the genome-wide association study (GWAS) design, have provided a wealth of novel insights into the aetiology of a wide range of human diseases and traits, in particular cardiovascular diseases and lipid biomarkers. The next challenge consists of understanding the molecular basis of these associations. The integration of multiple association datasets, including gene expression datasets, can contribute to this goal. We have developed a novel statistical methodology to assess whether two association signals are consistent with a shared causal variant. An application is the integration of disease scans with expression quantitative trait locus (eQTL) studies, but any pair of GWAS datasets can be integrated in this framework. A key feature of the method is the ability to derive the output statistics from single SNP summary statistics, hence making it possible to perform systematic meta-analysis type comparisons across multiple GWAS datasets (implemented online at <http://coloc.cs.ucl.ac.uk/coloc/>). We demonstrate the value of the approach by re-analysing a gene expression dataset in 966 liver samples with a published meta-analysis of lipid traits including > 100,000 individuals of European ancestry [1]. Combining all lipid biomarkers, our re-analysis supported 29 out of 38 reported colocalisation results with eQTLs. Two clearly discordant findings (*IFT172*, *CPNE1*), as well as multiple new colocalisation results, highlight the value of a formal systematic statistical test. Our findings provide information about the causal gene in associated intervals and have direct implications for the understanding of complex diseases as well as the design of drugs to target disease pathways.

## Author Summary

Genome-wide association studies (GWAS) have found a large number of genetic regions (“loci”) affecting clinical end-points and phenotypes, many outside coding intervals. One approach to understanding the biological basis of these associations has been to explore whether GWAS signals from intermediate cellular phenotypes, in particular gene expression, are located in the same loci (“colocalise”) and are potentially mediating the disease signals. However it is not clear how to assess whether the same variants are responsible for the two GWAS signals or whether it is distinct causal variants close to each other. In this paper, we describe a statistical method which can use only single variant summary statistics to test for colocalisation of GWAS signals. We describe one application of our method to a meta-analysis of blood lipids and liver expression, although any two datasets resulting from association studies can be used. Our method is able to detect the subset of GWAS signals explained by regulatory effects and

identify candidate genes affected by the same GWAS variants. As summary GWAS data are increasingly available, applications of colocalisation methods to integrate the findings will be essential for functional follow-up, and will also be particularly useful to identify tissue specific signals in eQTL datasets.

## Introduction

In the last decade, hundreds of genomic loci affecting complex diseases and disease relevant intermediate phenotypes have been found and robustly replicated using genome-wide association studies (GWAS, [2]). At the same time, gene expression measurements derived from microarray [3] or RNA sequencing [4] studies have been used extensively as an outcome trait for the GWAS design. Such studies are usually referred to as expression quantitative trait locus (eQTL) analysis. While GWAS datasets have provided a steady flow of positive and replicable results, the interpretation of these findings, and in particular the identification of underlying molecular mechanisms, has proven to be challenging. Integrating molecular level data and other disease relevant intermediate phenotypes with GWAS results is the natural step forward in order to understand the biological relevance of these results.

In this context, a natural question to ask is whether two independent association signals at the same locus, typically generated by two GWAS studies, are consistent with a shared causal variant. If the answer is positive, we refer to this situation as colocalised traits, and the likelihood that both traits are causally linked increases greatly. A typical example involves an eQTL study and a disease association result, which points to the causal gene and the tissue in which the effect is mediated [5–7]. The same questions can also be considered between pairs of eQTLs [8, 13], or pairs of diseases [9].

However, identifying the traits that share a common association signal is not a trivial statistical task. Visual comparison of overlaps of association signals with an expression dataset is a step in this direction (using for example Sanger tool Genevar <http://www.sanger.ac.uk/resources/software/genevar/>), but the abundance of eQTLs in the human genome and across different tissues makes an accidental overlap between these signals very likely [3]. Therefore visual comparison is not enough to make inferences about causality and formal statistical tests must be used to address this question.

Nica et al. [5] proposed a methodology to rank the SNPs with an influence on two traits based on the residual association conditional on the most associated SNP. By comparing the GWAS SNP score with all other SNPs in the associated region, this method accounts for the local LD structure. However this is not a formal test of a null hypothesis for, or against, colocalisation at the locus of interest. A formal test of colocalisation has been developed in a regression framework, either frequentist [10] or Bayesian [11]. In this context, the question of colocalisation amounts to testing whether the estimated regression coefficients against a chosen set of SNPs are proportional between two datasets. Both of these approaches have been shown to be closely linked, at least when the sample size is large [11]. They also share the drawback of having to specify a subset of SNPs to base the test on, and Wallace [12] shows that this step can generate significant biases. The main source of bias is the fact that the causal variant may not always be the most strongly associated one, which can over-estimate the differences between two datasets and reject colocalisation in situations where the causal SNP is in fact shared.

Although approaches exist to overcome this [12], perhaps the greatest limitation is the requirement for individual level genotype data, which are rarely available for large scale eQTL datasets.

The success of GWAS meta-analyses has shown that there is considerable benefit in being able to derive association tests on the basis of summary statistics. With these advantages in mind, He et al. [7] developed a statistical test to match the pattern of gene expression with a GWAS dataset. This approach, coded in the software Sherlock, can accommodate  $p$ -values as input. However, their hypothesis of interest differs from the question of colocalisation, with the focus of the method being on genomewide convergence of signals, assuming an abundance of trans eQTLs. In particular, SNPs that are not associated with gene expression do not contribute to the test statistic. Such variants can provide strong evidence against colocalisation if they are strongly associated with the GWAS outcome.

These limitations motivate the development of novel methodologies to test for colocalisation between pairs of traits. Here, we derive a novel Bayesian statistical test for colocalisation that addresses many of the shortcomings of existing tools. Our underlying model is closely related to the approach developed by Flutre et al. [13], which considers the different but related problem of maximising the power to discover eQTLs in expression datasets of multiple tissues. A key feature of our approach is that it only requires single SNP  $p$ -values and their minor allele frequencies (MAFs), or estimated allelic effect and standard error, combined with closed form analytical results that enable quick comparisons, even at the genome-wide scale. Our procedure provides intuitive posterior probabilities that can be easily interpreted. A key application is the systematic comparison between a new GWAS dataset and a large catalogue of association studies in order to identify novel shared mechanisms. We demonstrate the value of the method by re-analysing a large scale meta-analysis of blood lipids [1] in combination with a gene expression study in 966 liver samples [14].

## Results

### Overview of the method

We consider a situation where two traits (denoted by  $Y_1, Y_2$ ) have been measured in two distinct datasets of unrelated individuals. We assume that samples are drawn from the same ethnic group, i.e. allele frequencies and pattern of linkage disequilibrium (LD) are identical in both populations. We consider, for each variant, a linear trend model between the outcome phenotype  $Y$  and the genotypes  $X$  (or a log-odds generalized linear model if one of the two outcome phenotypes  $Y$  is binary):

$$Y = \mu + \beta X$$

We are interested in a situation where single variant association  $p$ -values and MAFs, or estimated regression coefficients  $\hat{\beta}$  and their estimated precisions  $var(\hat{\beta})$ , are available for both datasets at  $Q$  variants, typically SNPs but also indels. We make two additional assumptions and discuss later in this paper how these can be relaxed. Firstly, that the causal variant is included in the genotyping panel, either directly typed or well imputed [15–17]. Secondly, that at most one association is present for each trait. We are interested in exploring whether the data support a shared causal variant for both traits. While the method is fully applicable to case-control outcome, we consider two quantitative traits in this initial description.

SNP causality in a region of  $Q$  variants can be summarised for each trait using a vector of length  $Q$  of  $(0, 1)$  values, where 1 means that the variant is causally associated with the trait of interest and at most one entry is non-zero. A schematic illustration of this framework is provided in Figure 1 in a region that contains 8 SNPs. Each possible pair of vectors (for traits 1 and 2, which we refer to as “configuration”) can be assigned to one of five hypothesis:

- $\mathbb{H}_0$ : No association with either trait
- $\mathbb{H}_1$ : Association with trait 1, not with trait 2
- $\mathbb{H}_2$ : Association with trait 2, not with trait 1
- $\mathbb{H}_3$ : Association with trait 1 and trait 2, two independent SNPs
- $\mathbb{H}_4$ : Association with trait 1 and trait 2, one shared SNP

In this framework, the colocalisation problem can be re-formulated as assessing the support for all configurations (i.e. pairs of binary vectors) in hypothesis  $\mathbb{H}_4$ .

Our method is Bayesian in the sense that it integrates over all possible configurations. This process requires the definition of prior probabilities which are defined at the SNP level (see Methods). A likelihood can be assigned to each configuration, and these likelihoods can be summed over all configurations and combined with the prior to assess the support for each hypotheses  $(\mathbb{H})_1^5$ . The result of this procedure is five posterior probabilities (PP0, PP1, PP2, PP3 and PP4). A large posterior probability for hypothesis 3, PP3, indicates support for two independent causal SNPs associated with each trait. In contrast, if PP4 is large, the data support a shared causal variant. While the method uses Approximate Bayes Factor computations (ABF, [18], and Methods), no iterative computation scheme (such as Monte Carlo Markov Chain) is required. Therefore, computations are quick and do not require any specific computing infrastructure. Importantly, the use of ABF enable the computation of posterior probabilities from single variant association  $p$ -values and MAFs, although the estimated single SNP regression coefficients  $\hat{\beta}$  and their variances or standard errors are preferred for imputed data.

### Sample size required for colocalisation analysis

Given the well understood requirements for large sample size for GWAS data, we used simulations to investigate the power of our approach in a situation where the causal variant is shared between both traits in the two datasets, the “eQTL dataset” and the “biomarker dataset”. Using the original sample size of the liver eQTL dataset described herein [14], and a variance explained of 10% for expression [19, 20], we show the results of simulations for different sample sizes for the biomarker dataset, as well as different levels of variance explained by the causal variant for the biomarker data (Figure 2). We find that given sample size of 2,000 for the biomarker dataset, the causal variant needs to explain close to 2% of the variance of the biomarker to provide reliable evidence in favour of a colocalised signal (lower 10<sup>th</sup> percentile for PP4 > 80%).

### Consequence of limited variant density

Owing to the assumption that the causal variant is included in the genotype panel, we hypothesised that the use of non-imputed GWAS data (such as traditional SNP arrays) would have an impact on the computed metrics. To investigate this possibility, we used densely imputed data in regions covered by the MetaboChip Illumina array and only selected the subset of variants that were present in the Illumina 660K genotyping array. We simulated data under the assumption of a shared causal variant, with 4,000 individuals in the biomarker dataset. We then computed the PP4 statistic with and without restricting the SNP set to the Illumina 660K Chip SNPs (Figure 3). We also considered the different settings whereby the causal SNP is or is not directly genotyped by the Illumina array (see Figure S1 and Figure S2 for more exhaustive simulations). Our results show that when the causal variant is directly genotyped by the low density array, the use of imputed data is not essential provided that the sample size is sufficiently large (Figure 3). However, in cases where the causal variant is not typed or imputed, the variance of PP4 is much higher (Figure 3B). In this situation, the resulting PP4 statistic tends to decrease even though considerable variability is observed.

### Comparison with existing colocalisation tests

We compared the behaviour of our proposed test with that of proportional colocalisation testing [10, 12] in the specific case of a biomarker dataset with 10,000 samples (Figure 4). Broadly, in the case of either a single common causal variant or two distinct causal variants, our proposed method could infer the simulated hypotheses correctly (PP4 or PP3 > 0.9) with good confidence, and PP3 > 0.9 slightly more often than the proportional testing  $p$ -value < 0.05, indicating greater power. However, our proposed method is relatively more affected by using only tagging SNPs compared to proportional testing. We also

examined the behaviour when the assumption of a single causal variant was violated. In the case where there were more than two or more causal variants, none common to both traits, H3 consistently had the greatest support.

It has been proposed that gene expression may be subject to both global regulatory variation which acts across multiple tissues and secondary tissue specific regulators [21]. Neither approach covers this case explicitly in its construction, but it is instructive to examine their expected behaviour. The proportional approach tends to reject a null of colocalisation, suggesting that a single distinct causal variant can be sufficient to violate the null hypothesis of proportional regression coefficients. In contrast, the Bayesian approach tends to favour the shared variant in the cases covered by our simulations (median PP4 > median PP3), and either hypotheses H3 or H4 can potentially have strong support (PP4 > 0.9 in close to 50% of simulations, and PP3 > 0.9 in around 25% of simulations). Of course, the ultimate goal should be to extend these tests to cover multiple causal variants, but in the meantime, it can be useful to know that a high PP4 in our proposed Bayesian analysis indicates strong support for “at least one causal variant” and that rejection of the null of proportionality of regression coefficients indicates that the two traits do not share all causal variants, not that they cannot share one.

## Application to a meta-analysis of blood lipids combined with a liver expression dataset

Teslovich et al. [1] reported common variants associated with plasma concentrations of low-density lipoprotein cholesterol (LDL), high-density lipoprotein cholesterol (HDL) and triglyceride (TG) levels in more than 100,000 individuals of European ancestry. They then reported the correlations between the lead SNPs at the loci they found and the expression levels of transcripts in liver. They defined a region within 500 kilobases of the lead SNP, and the threshold for significance is  $10^{-8}$ . At this threshold, they found 38 SNP-to-gene eQTLs in liver ([1] Supplementary Table 8). The liver expression dataset used in this analyses is the same as the one used in Teslovich et al.

Table 1 lists the results of our re-analysis of the associations that were reported correlated with both lipid levels and expression in human liver in the published dataset but that were not supported by our method. To assess the role of the prior, we varied the critical parameter  $p_{12}$ , which codes for the prior probability that a variant is associated with both traits. Here we report the results using  $p_{12} = 10^{-5}$ . The complete list of results is provided in Table S1.

For 7 of the reported genes, our method does not support a shared association between primary signal for liver expression and one of the four lipid traits (PP4 < 80%). In addition, we found strong evidence of distinct signals between *HLA-DQ/HLA-DR* and TC (Table S1) but these results must be interpreted with caution owing to the extensive polymorphism in the major histocompatibility complex region.

The data support distinct associations (PP3 > 90% and PP4 < 10%) for two loci (*IFT172* and TG or TC, *CPNE1* and TC), suggesting that expression in liver is not causally associated with the listed traits (table 1 and Figure S5).

*FRK* expression and TC (PP3 = 61%), and *TBKBP1* and TC PP3 = 78% or LDL PP3 = 47%), do not exclude a shared causal variant, but in these regions with extensive level of LD, it is difficult to obtain firm support for a shared causal variant, which leads to lower PP4 estimates. In one case, *CEP250*, our re-analysis of the expression data did not identify an eQTL for this gene. In such a situation, both PP3 and PP4 are low and PP0, PP1 and PP2 concentrate most of the posterior distribution.

We also found several loci not previously reported with a high probability of influencing the lipid traits in liver (PP4 > 80%, Tables S2,S3,S4,S5).

To explore the possibility that secondary signals may colocalise, we performed the test using conditional analysis results for the eQTL dataset. Two of the loci (*SYPL2*/LDL or TC, *APOC4* and TG) show evidence of colocalisation with expression after conditional analysis. The interpretation of these cases is discussed in the section below.

Table 2 lists all the eQTLs with strong colocalisation support for more than one of the three biomarkers LDL, HDL and TG, possibly suggesting a common pathway between the biomarkers and liver expression.

## Dealing with several independent associations for the same trait

We have so far assumed that each trait is associated with at most one causal variant per locus. However, recent studies have shown that it is not unusual to observe two or more independent associations at a locus for a trait of interest [22]. In the presence of multiple associations for either of the two datasets, the method will test for colocalisation between the most significant of these multiple associations within each dataset.

We accommodate for multiple associations by conditioning on the most associated variant, computing conditional  $p$ -values and repeating the colocalisation test. Two of the six genes listed in table 1 were found to have a high probability of a shared signal using the conditional association signal for expression. An example of this procedure for the gene *SYPL2* and LDL is provided in Figure 5. We find that the top liver eQTL signal is clearly discordant with LDL association (Table 1 and Figure 5). However, conditioning on the top eQTL signal reveals a second independent association for *SYPL2* expression in liver. This secondary *SYPL2* eQTL colocalises with the LDL association (PP4 > 90%, Figure 5).

## Web based resource

We developed a web site designed for integration of GWAS results using only  $p$ -values and the sample size of the datasets (<http://coloc.cs.ucl.ac.uk/coloc/>). The website was developed using RWUI [23]. Results include a list of potentially causal genes with the associated PP4 with their respective plots and ABF, and can be viewed either interactively or returned by email.

Researchers can request a genome-wide scan of results from a genetic association analysis, and obtain a list of genes with a high probability of mediating the GWAS signals in a particular tissue. The tools also allows visualisation of the signals within a genetic region of interest (see Figure S5 for examples of output of regional association plots).

The database and browser currently include the possibility of investigating colocalisation only with liver expression data, however the resource will soon be extended to include expression in different tissues (including brain, [24, 25]), a multiplicity of molecular phenotypes and clinical variables, and different diseases. This method, as well as alternative approaches for colocalisation testing [10, 12], are also available with additional input options in an R package, *coloc*, from the Comprehensive R Archive Network (<http://cran.rproject.org/web/packages/coloc>).

## Discussion

We have developed a novel Bayesian statistical procedure to assess whether two association signals are colocalised. The strength of this approach lies in its speed and analytical forms, combined with the fact that it can use single variant  $p$ -values when only these are available.

Our method differs from a typical fine-mapping exercise in the sense that we are not interested in knowing which variant, potentially shared between datasets, is likely to be causal but only whether a shared causal variant is plausible. We currently assume that each variant is equally likely a priori to affect gene expression or trait. A straightforward addition to our methodology would consider location specific priors for each variant, which would depend for example on the distance to the gene of interest, or the presence of functional elements in this chromosome region [26, 27]. Our computation of the BF also assumes that, under  $H_4$ , the effect sizes for both traits are independent. This assumption may not be appropriate for example if one compares eQTLs across different tissue types, or the same trait in two different studies. A more general BF computation which includes cases where the effects are correlated

has been proposed in [28] and could be used to extend our approach to these situations. Our results also show that high genotyping density, and/or accurate use of imputation techniques, are essential to provide an accurate answer to the colocalisation problem.

Good imputation quality is another important parameter. Indeed, while the variance of the regression coefficient can be estimated solely on the basis of the minor allele frequency for typed SNPs and sample size (and the case control ratio in the case of a binary outcome) [16, 29], this ignores the uncertainty due to the imputation procedure. Filtering out poorly imputed SNPs partially addresses this problem. However, filtering may lead to the exclusion of the causal variant, which can make our colocalisation test overly conservative (Figure 3B). Hence, providing estimates of the variance of the MLE, together with the effect estimates, will result in greater accuracy. This additional option is available on the coloc package in R (<http://cran.rproject.org/web/packages/coloc>).

Another issue is the choice of prior probabilities. For the eQTL analysis, we used a  $10^{-4}$  prior probability for a cis-eQTL. A more stringent threshold may be better suited for variants further away from the gene under genetic control. We also used a prior probability of  $10^{-4}$  for the lipid associations. Although our knowledge about this is still lacking, this estimate has been suggested in the literature in the context of GWAS [18, 35, 36]. We assigned a prior probability of  $1 \times 10^{-5}$  for  $p_{12}$ , which encodes the probability that a variant affects both traits. It has been shown that SNPs associated with complex traits are more likely to be eQTLs compared to other SNPs chosen at random from GWAS platforms [31], and a higher weighting for these SNPs has been proposed when performing Bayesian association analyses [37, 38]. Also, eQTLs have been shown to be enriched for disease-associated SNPs when a disease-relevant tissue is used [8, 39]. Our sensitivity analysis for the  $p_{12}$  parameter showed broadly consistent results (Table S1). In cases where GWAS data are available for both traits, [13] show that it is possible to estimate these parameters from the data using a hierarchical model. This addition is a possible extension of our approach.

Our comparison with the proportional colocalisation testing approach demonstrates a further advantage of our proposed method. When proportional testing fails to reject its null hypothesis of no colocalisation, it cannot be distinguished whether there truly is a shared causal variant or whether the effects or sample sizes are simply too small to provide sufficient power to reject the null. In contrast, our method clearly provides support for either a shared or distinct causal variants in its partitioning of the hypothesis space.

The interpretation of our posterior probabilities requires caution. For example, a low PP4 probability (which supports a shared causal variant) may not imply that the association signals are distinct: if the sample size is not sufficient, PP3 will also be low and it will not be possible to draw meaningful conclusions. Even if PP3 (i.e. the posterior probability supporting different causal variants) is high, one cannot exclude that this might be a consequence of the causal variant not being typed or imputed (Figure 3). Lastly, a high PP4 is a measure of correlation, not causality. To illustrate this point, one can consider the relatively common situation where a single variant appears to affect the expression of several genes in the same chromosome region. All three eQTLs will be colocalised and potentially also colocalised with the biomarker of interest. In this situation one would typically expect that a single gene is causally involved in the biomarker pathway but the colocalisation test with the biomarker will generate high PP4 values for all genes in the interval.

We show that we can use conditional  $p$ -values to deal with multiple independent associations with the same trait at one locus. The identification of loci with several independent liver eQTLs, with only some of them associated with a biomarker of interest (as in Figure 5), points to intriguing biological mechanisms. Different effects on several isoforms may explain these discrepancies but additional follow-up will be required to fully understand such loci. In situations where only single SNP summary statistics are available, the approximate conditional meta-analysis framework proposed by Visscher et al [30] can be used to obtain conditional  $p$ -values. As pointed out by Wallace [12], this top SNP selection for the conditional analysis can create biases, although the bias is small in the case of large samples and/or strong

effects. Alternatively, if individual genotype data are available, one can estimate Bayes factors for pairs of variants rather than single variants. This modification would allow up to two associations per trait while avoiding the issue of SNP selection for the conditional analysis. At the expense of computational speed, this solution can be extended to all set of  $n$  variants hence enabling up to  $n$  independent associations per locus and per outcome trait.

Importantly, GWAS signals can be explained by eQTLs only when the causal variant affects the phenotype by altering the amount of protein production, but not when the phenotype is affected by changing the type of protein produced, although the former seems to be the most common [31]. Furthermore, since many diseases manifest their phenotype in certain tissues exclusively [3, 21, 32, 33], colocalisation results will be dependent on the expression dataset used. In addition to identifying the causal genes, the identification of tissue specificity for the molecular effects underlying GWAS signals is a key outcome of our method. We anticipate that building a reference set of eQTL studies in multiple tissues will provide a useful check for every new GWAS dataset, pointing directly to potential candidate genes/tissue types where these effects are mediated.

While this report focuses on finding shared signals between a biomarker dataset and a liver expression dataset, we plan to utilise summary results of multiple GWAS and eQTL studies, for a variety of cell types and traits. Disease/disease, eQTL/disease or disease/biomarkers comparisons are all of biological interest. We expect that the fact that the test can be based on single SNP summary statistics will be key to overcome data sharing concerns, hence enabling a large scale implementation of this tool. The increasing availability of RNA-Seq eQTL studies will further increase the opportunity to detect isoform specific eQTLs and their relevance to disease studies. Owing to the increasing availability of GWAS datasets, the systematic application of this approach will potentially provide clues into the molecular mechanisms underlying GWAS signals and the aetiology of the disorders.

## Materials and Methods

### Expression dataset

We used in our analysis gene expression and genotype data from 966 human liver samples. The samples were collected post-mortem or during surgical resection from unrelated European-American subjects from two different non-overlapping studies, which have been described in [14]. The cohorts were both genotyped using Illumina 650Y BeadChip array, and 39,000 expression probes were profiled using Agilent human gene expression arrays. All of the expression data has been normalised as one unit even though they were part of different studies, since high concordance between data generated using the same array platforms has been previously reported. Probe sequences were searched against the human reference genome GRCh37 from 1000genomes using BLASTN. Multiple probes mapping to one gene were kept in order to examine possible splicing. The probes were kept and annotated to a specific gene if they were entirely included in genes defined by Ensembl ID or by HGNC symbol using the package biomaRt in R [34]. After mapping and annotating the probes, we were left with 40,548 mapped probes covering 24,927 genes.

### Imputation of genetic data

Quality control filters were applied both before and after imputation. Before imputation, individuals with more than 10% missing genotypes were removed, and SNPs showing a missing rate greater than 10%, a deviation for HWE at a  $p$ -value less than 0.001 were dropped. After imputation, monomorphic SNPs were excluded from analyses.

To speed up the imputation process, the genome was broken into small chunks that were phased and imputed separately and then re-assembled. This was achieved using the ChunkChromosome tool



(<http://genome.sph.umich.edu/wiki/ChunkChromosome>), and specifying chunks of 1000 SNPs, with an overlap window of 200 SNPs on each side, which improves accuracy near the edges during the phasing step. Each chunk was phased using the program MACH1 (Versions 1.0.18, downloaded from: <http://www.sph.umich.edu/csg/abecasis/MACH/download/>) with the number of states set to 300 and the number of rounds of MCMC set to 20 for all chunks. Phased haplotypes were used as a basis for imputation of untyped SNPs using the program Minimac (version 2011.10.27, downloaded from: <http://www.sph.umich.edu/csg/afuchs/minimac-beta-2012.8.15.tgz>) with 1000 Genomes EUR reference haplotypes (phase1 version 3, March 2012) to impute SNPs not genotyped on the Illumina array. The data was then collated in probability format that can be used by the R Package `snpStats` [34].

## eQTL analysis

eQTL  $p$ -values, effect sizes, and standard errors were obtained by fitting a linear trend test regression between the expression of each gene and all variants 200 kilobases upstream and downstream from each probe. After filtering out the variants with  $MAF < 0.001$ , monomorphic SNPs, multi-allelic SNPs (as reported in 1000 Genomes or in the Ensembl database) and variants not sufficiently well imputed ( $Rsq < 0.3$ , as defined by minimac <http://genome.sph.umich.edu/wiki/minimac>) between both datasets, we applied our colocalisation procedure. We conducted conditional analysis on SNPs with  $p$ -values  $< 10^{-4}$  for the expression associations, and repeated the colocalisation test using expression data conditioned on the most significant SNP. The aim of this analysis is to explore whether additional signals for expression other than the main one are shared with the biomarker signal.

## Biomarker dataset

The biomarker  $p$ -values from the meta-analyses (with genomic control correction) were obtained from a publicly available repository (<http://www.sph.umich.edu/csg/abecasis/public/lipids2010/>).

## Posterior Computation

We call a ‘‘configuration’’ one possible combination of pairs of binary vectors indicating whether the variant is associated with the selected trait. We can group the configurations into five sets,  $S_0, S_1, S_2, S_3, S_4$ , containing assignments of all SNPs  $Q$  to the functional role corresponding to the five hypothesis  $\mathbb{H}_0, \mathbb{H}_1, \mathbb{H}_2, \mathbb{H}_3, \mathbb{H}_4$ . We can compute the posterior probabilities for each of these 5 hypothesis using a Bayesian approach. Firstly, we calculate the likelihood of each hypothesis,

$$L(H_h) = L(H_h | D) = \sum_{S \in S_h} P(D | S)P(S) \quad (1)$$

where  $P(S)$  is the prior probability of a configuration,  $P(D | S)$  is the probability of the observed data  $D$  under a given configuration (which we will refer to as the likelihood of this configuration), and the sum is over all configurations  $S$  which are consistent with a given hypothesis  $H_h$ ,  $h=(1,2,3,4)$ . Thus, each configuration likelihood is weighted by the prior probability of that configuration.

Next, we compute the posterior probability by taking the ratio of the hypothesis likelihoods under the different hypotheses. For example, the posterior probability under hypothesis 4 is:

$$PP4 = \frac{L(H_4)}{L(H_0) + L(H_1) + L(H_2) + L(H_3) + L(H_4)} \quad (2)$$

If we also divide each probability by the baseline  $L(H_0)$  we get:

$$PP4 = \frac{\frac{L(H_4)}{L(H_0)}}{1 + \frac{L(H_1)}{L(H_0)} + \frac{L(H_2)}{L(H_0)} + \frac{L(H_3)}{L(H_0)} + \frac{L(H_4)}{L(H_0)}} \quad (3)$$

The ratios in the numerator and denominator of equation 3 are:

$$\frac{L(H_h)}{L(H_0)} = \sum_{S \in S_h} \frac{P(D | S)}{P(D | S_0)} \times \frac{P(S)}{P(S_0)} \quad (4)$$

The first ratio in this equation is a Bayes Factor (BF) for each configuration, and the second ratio is the prior odds of a configuration compared with the baseline configuration  $S_0$ . The BF can be computed for each variant from the  $p$ -value, or estimated regression coefficient  $\hat{\beta}$  and variance of  $\hat{\beta}$ , using Wakefield's method. By summing over all configurations in  $S_h$  we are effectively comparing the support in the data for one alternative hypothesis versus the null hypothesis.

## Bayes factor computation

A Bayes Factor for each SNP and each trait 1 and 2 was computed using the Approximate Bayes Factor (ABF, [18]). Wakefield's method yields a Bayes factor that measures relative support for a model in which the SNP is associated with the trait compared to the null model of no association.

The equation used is the following:

$$ABF = \frac{1}{\sqrt{1-r}} \times \exp \left[ -\frac{Z^2}{2} \times r \right] \quad (5)$$

where  $Z = \hat{\beta}/\sqrt{V}$  is the usual Z statistic and the shrinkage factor  $r$  is the ratio of the variance of the prior and total variance ( $r = W/(V + W)$ ).

Assuming a normal distribution, the  $p$ -value of each SNP can be converted to standard one-tailed Z-score by using inverse normal cumulative distribution function. So for a SNP, all that is needed are the  $p$ -values from a standard regression output, and  $\sqrt{W}$ , the standard deviation of the normal prior  $N(0, W)$  on  $\beta$ .

The variance of the effect estimate,  $V$ , can be approximated using the MAF and sample size. However for imputed data it is preferable to use the variance outputted in standard regression analysis directly in the ABF equation. For the expression dataset used here, the variance and effect estimates from the regression analysis were used for computation of ABFs. More details can be found in the Supplementary materials.

## Choice of priors

Prior probabilities are assigned at the SNP level and correspond to mutually exclusive events. We assigned a prior of  $1 \times 10^{-4}$  for  $p_1$  and  $p_2$ , the probability that a SNP is associated with either of the two traits. Since all SNPs are assumed to have the same prior probability of association, this prior can be interpreted as an estimate for the proportion of SNPs that we expect to be associated with the trait in question.

We also assigned a prior probability of  $1 \times 10^{-5}$  for  $p_{12}$ , the probability that one SNP is associated with both traits. This probability can be better understood when it is re-expressed as the conditional probability of a SNP being associated with trait 2, given that it is associated with trait 1. So assigning a probability of  $1 \times 10^{-5}$  means that 1 in 10 SNPs that are associated with trait 1 is also associated with the other. As a sensitivity analysis, we ran the comparison with Teslovich et al. using two other prior probabilities for  $p_{12}$ ,  $p_{12} = 2 \times 10^{-6}$  which means 1 in 50 SNPs that are associated with one trait is also associated with the other; and  $p_{12} = 10^{-6}$  which means 1 in 100 SNPs.

To compute the ABF, we also needed to specify the standard deviation for the prior, and we set this to 0.20 for binary traits and 0.15 for quantitative traits (more details in Supplementary).

## Acknowledgments

Claudia Giambartolomei is supported by a grant from the British Heart Foundation (CD6H).  
Chris Wallace is supported by the Wellcome Trust (WT089989).


## References

1. Teslovich TM, Musunuru K, Smith AV, Edmondson AC, Stylianou IM, et al. (2010) Biological, clinical and population relevance of 95 loci for blood lipids. *Nature* 466: 707–713.
2. Feero WG, Guttmacher AE, Manolio TA (2010) Genomewide association studies and assessment of the risk of disease. *New England Journal of Medicine* 363: 166–176.
3. Nica AC, Dermitzakis ET (2008) Using gene expression to investigate the genetic basis of complex disorders. *Human molecular genetics* 17: R129–R134.
4. Pickrell JK, Marioni JC, Pai AA, Degner JF, Engelhardt BE, et al. (2010) Understanding mechanisms underlying human gene expression variation with rna sequencing. *Nature* 464: 768–772.
5. Nica AC, Montgomery SB, Dimas AS, Stranger BE, Beazley C, et al. (2010) Candidate causal regulatory effects by integration of expression qtls with complex trait genetic associations. *PLoS genetics* 6: e1000895.
6. Hunt KA, Zhernakova A, Turner G, Heap GA, Franke L, et al. (2008) Newly identified genetic risk variants for celiac disease related to the immune response. *Nature genetics* 40: 395–402.
7. He X, Fuller CK, Song Y, Meng Q, Zhang B, et al. (2013) Sherlock: Detecting gene-disease associations by matching patterns of expression qtl and gwas. *The American Journal of Human Genetics* 92: 667–680.
8. Ding J, Gudjonsson JE, Liang L, Stuart PE, Li Y, et al. (2010) Gene expression in skin and lymphoblastoid cells: Refined statistical method reveals extensive overlap in cis-eqtl signals. *The American Journal of Human Genetics* 87: 779–789.
9. Cotsapas C, Voight BF, Rossin E, Lage K, Neale BM, et al. (2011) Pervasive sharing of genetic effects in autoimmune disease. *PLoS genetics* 7: e1002254.
10. Plagnol V, Smyth DJ, Todd JA, Clayton DG (2009) Statistical independence of the colocalized association signals for type 1 diabetes and rps26 gene expression on chromosome 12q13. *Biostatistics* 10: 327–334.
11. Wallace C, Rotival M, Cooper JD, Rice CM, Yang JH, et al. (2012) Statistical colocalization of monocyte gene expression and genetic risk variants for type 1 diabetes. *Human molecular genetics* 21: 2815–2824.
12. Wallace C (2013) Unbiased statistical testing of shared genetic control for potentially related traits. arXiv preprint arXiv:13015510 .
13. Flutre T, Wen X, Pritchard J, Stephens M (2013) A statistical framework for joint eqtl analysis in multiple tissues. *PLoS Genet* 9: e1003486.


14. Schadt EE, Woo S, Hao K (2012) Bayesian method to predict individual snp genotypes from gene expression data. *Nature genetics* 44: 603–608.
15. Howie B, Marchini J, Stephens M (2011) Genotype imputation with thousands of genomes. *G3: Genes, Genomes, Genetics* 1: 457–470.
16. Marchini J, Howie B (2010) Genotype imputation for genome-wide association studies. *Nature Reviews Genetics* 11: 499–511.
17. Howie B, Fuchsberger C, Stephens M, Marchini J, Abecasis GR (2012) Fast and accurate genotype imputation in genome-wide association studies through pre-phasing. *Nature genetics* .
18. Wakefield J (2009) Bayes factors for genome-wide association studies: comparison with p-values. *Genetic Epidemiology* 33: 79–86.
19. Schadt EE, Molony C, Chudin E, Hao K, Yang X, et al. (2008) Mapping the genetic architecture of gene expression in human liver. *PLoS biology* 6: e107.
20. Innocenti F, Cooper GM, Stanaway IB, Gamazon ER, Smith JD, et al. (2011) Identification, replication, and functional fine-mapping of expression quantitative trait loci in primary human liver tissue. *PLoS genetics* 7: e1002078.
21. Brown CD, Mangravite LM, Engelhardt BE (2012) Integrative modeling of eqtls and cis-regulatory elements suggest mechanisms underlying cell type specificity of eqtls. *arXiv preprint arXiv:12103294* .
22. Trynka G, Hunt KA, Bockett NA, Romanos J, Mistry V, et al. (2011) Dense genotyping identifies and localizes multiple common and rare variant association signals in celiac disease. *Nature genetics* 43: 1193–1201.
23. Newton R, Wernisch L (2007) *Rwui: A web application to create user friendly web interfaces for r scripts*. *New Functions for Multivariate Analysis* : 32.
24. Trabzuni D, Ryten M, Walker R, Smith C, Imran S, et al. (2012) Quality control parameters on a large dataset of regionally dissected human control brains for whole genome expression studies. *Journal of Neurochemistry* 120: 473–473.
25. Ramasamy A, Trabzuni D, Gibbs JR, Dillman A, Hernandez DG, et al. (2013) Resolving the polymorphism-in-probe problem is critical for correct interpretation of expression qtl studies. *Nucleic Acids Research* .
26. Khatun J (2012) An integrated encyclopedia of dna elements in the human genome. *Nature* .
27. Gerstein MB, Kundaje A, Hariharan M, Landt SG, Yan KK, et al. (2012) Architecture of the human regulatory network derived from encode data. *Nature* 489: 91–100.
28. Wen X, Stephens M (2011) Bayesian methods for genetic association analysis with heterogeneous subgroups: from meta-analyses to gene-environment interactions. *arXiv preprint arXiv:11111210* .
29. Guan Y, Stephens M (2008) Practical issues in imputation-based association mapping. *PLoS Genet* 4: e1000279.
30. Yang J, Ferreira T, Morris AP, Medland SE, Madden PA, et al. (2012) Conditional and joint multiple-snp analysis of gwas summary statistics identifies additional variants influencing complex traits. *Nature genetics* 44: 369–375.

31. Nicolae DL, Gamazon E, Zhang W, Duan S, Dolan ME, et al. (2010) Trait-associated snps are more likely to be eqtls: annotation to enhance discovery from gwas. *PLoS genetics* 6: e1000888.
32. Dimas AS, Deutsch S, Stranger BE, Montgomery SB, Borel C, et al. (2009) Common regulatory variation impacts gene expression in a cell type-dependent manner. *Science* 325: 1246–1250.
33. Hernandez DG, Nalls MA, Moore M, Chong S, Dillman A, et al. (2012) Integration of gwas snps and tissue specific expression profiling reveal discrete eqtls for human traits in blood and brain. *Neurobiology of disease* .
34. Team RDC (2013) R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria. URL <http://www.R-project.org>.
35. Stephens M, Balding DJ (2009) Bayesian statistical methods for genetic association studies. *Nature Reviews Genetics* 10: 681–690.
36. Burton PR, Clayton DG, Cardon LR, Craddock N, Deloukas P, et al. (2007) Genome-wide association study of 14,000 cases of seven common diseases and 3,000 shared controls. *Nature* 447: 661–678.
37. Knight J, Barnes MR, Breen G, Weale ME (2011) Using functional annotation for the empirical determination of bayes factors for genome-wide association study analysis. *PloS one* 6: e14808.
38. Johansson M, Roberts A, Chen D, Li Y, Delahaye-Sourdeix M, et al. (2012) Using prior information from the medical literature in gwas of oral cancer identifies novel susceptibility variant on chromosome 4-the adapt method. *PloS one* 7: e36888.
39. Richards A, Jones L, Moskvina V, Kirov G, Gejman P, et al. (2011) Schizophrenia susceptibility alleles are enriched for alleles that affect gene expression in adult human brain. *Molecular psychiatry* 17: 193–201.


## Figures



**Figure 1.** Example of one configuration under different hypotheses represented by one binary vector for each trait of  $(0, 1)$  values of length  $n = 8$ , the number of shared variants in a region. The value of 1 means that the variant is causally involved in disease, 0 that it is not. The first plot shows the case where only one dataset shows an association. The second plot shows that the causal SNP is different for the biomarker dataset compared to the expression dataset. The third plot shows the configuration where the single causal variant is the fourth one.




**Figure 2. Simulation analysis with a shared causal variant between two studies: one eQTL (sample size 966 samples, 10% of the variance explained by the variant) and one biomarker (such as LDL).** The variance explained by the biomarker is colour coded and the x-axis shows the sample size of the biomarker study. The y axis shows the median, 10% and 90% quantile of the distribution of PP4 values (which supports a shared common variant).




**Figure 3. Simulation analysis with a shared causal variant between two studies: one eQTL (sample size 966 samples) and one biomarker (sample size of 4,000 samples).** The variance explained by the biomarker and the expression is the same and is colour coded. The x-axis shows the estimated PP4 for 1,000 simulations using data imputed from metaboChip Illumina array. The y-axis uses the same dataset restricted to variants present on the Illumina 660W genotyping array to assess the impact of a lower variant density. **A.** The causal variant is included in the Illumina 660W panel. **B.** The causal SNP not included in Illumina 660W panel.





**Figure 4. Summary of proportional and Bayesian colocalisation analysis of simulated data under a variety of scenarios.** Each plot shows a different scenario, the total number of causal variants in a region is indicated by number of circles in the plot titles with causal variants affecting both traits, the eQTL trait only, or the biomarker trait only, indicated by full circles, top-shaded circles and bottom-shaded circles respectively. For proportional testing, under either a PC or BMA approach, we show the proportion of simulations with  $p$  or  $ppp < 0.05$  (black horizontal line), while for the Bayesian analysis we plot the proportion of simulations with the posterior probability of the indicated hypothesis  $> 0.9$ . Error bars show 95% confidence intervals. In all cases, for the eQTL trait, there are 1,000 samples and genetic variants explain a total of 10% of trait variance; for the biomarker trait, there are 10,000 samples.



**Figure 5. LDL association and eQTL association plots at the *SYPL2* locus.** The x-axis shows the physical position on the chromosome (Mb) **A:**  $-\log_{10}(p)$  association  $p$ -values for LDL. The  $p$ -values are from the Teslovich et al published meta-analysis of > 100,000 individuals. **B:**  $-\log_{10}(p)$  association  $p$ -values for *SYPL2* expression in 966 liver samples. **C:**  $-\log_{10}(p)$  association  $p$ -values for *SYPL2* expression conditional on the top eQTL associated SNP at this locus (rs2359653).

## Tables

**Table 1. Loci discordant with published results**

Chr	Region	Gene	Trait	Primary signal		Secondary signal*	
				PP3 (%)	PP4 (%)	PP4 (%)	conditional SNP
1	109824678:110224737	<i>SYPL2</i>	LDL	> 99	< 1	99	rs2359653
			TC	> 99	< 1		
2	27467244:27867303	<i>IFT172</i>	TG	> 99	< 1		
			TC	> 99	< 1		
6	116062804:116462863	<i>FRK</i>	LDL	90	10		
			TC	61	39		
17	45589357:45989416	<i>TBKBP1</i>	LDL	47	51		
			TC	78	17		
19	45248464:45648523	<i>APOC4</i>	TG	> 99	< 1	96	19:45452692:A_AG
20	33899702:34299761	<i>CEP250</i>	TC	< 1	< 1		
20	34013995:34414054	<i>CPNE1</i>	TC	95	5		

Gene/eQTL associations previously reported as having a probable shared variant but not supported by our method based on PP4 (posterior probability for a shared signal values) < 80%. \*Secondary signals are reported only when there is a secondary eQTL at a  $p$ -value greater than  $10^{-4}$ . colocalisation tests are computed using the expression data conditioned on the listed SNP.

**Table 2. Loci associated with multiple biomarkers**

Chr	Region	Gene	Traits
1	109618271:110018330	<i>CELSR2</i>	LDL, HDL
1	109622208:110022267	<i>PSRC1</i>	LDL, HDL
1	109652373:110052432	<i>SORT1</i>	LDL, HDL
1	109744528:110144587	<i>PSMA5</i>	LDL, HDL
1	62849818:63249877	<i>DOCK7</i>	LDL, TG
1	62870388:63270447	<i>ANGPTL3</i>	LDL, TG
8	8795514:9195573	<i>PPP1R3B</i>	LDL, HDL
8	8992223:9392282	ENSG00000254235	LDL, HDL
11	61367291:61767350	<i>FADS1</i>	LDL, HDL, TG
11	61441012:61841071	<i>FADS3</i>	LDL, HDL, TG
12	111508189:111908248	<i>CUX2</i>	LDL, HDL
15	58334099:58734158	<i>ALDH1A2</i>	HDL, TG
15	58653103:59053162	<i>LIPC</i>	HDL, TG
16	71910841:72310900	<i>HPR</i>	LDL, TG
16	71910841:72310900	<i>HP</i>	LDL, TG
20	44327404:44727463	<i>PLTP</i>	HDL, TG

eQTLs that co-localise with two or more of the three biomarkers LDL, HDL, TG, defined by PP4 greater than 90% using Teslovich et al. biomarker data and the 966 liver eQTL study.

## Supplemental figures legends

**Figure S1. Simulation analysis with a shared causal variant between two studies, one eQTL (sample size 966 samples) and one biomarker, comparing results using imputed versus not imputed data where the causal SNP is included in both the cases.** Each plot shows different sample sizes for one dataset. The variance explained by the causal variant for both the traits is colour coded. The x-axis shows the estimated PP4 for 1,000 simulations using data imputed from metaboChip Illumina array (Methods). The y-axis uses the same dataset restricted to variants present on the Illumina 660W genotyping array to assess the impact of a lower variant density. The causal variant is included in the Illumina 660W panel.

**Figure S2. Simulation analysis with a shared causal variant between two studies, one eQTL (sample size 966 samples) and one biomarker, comparing results using imputed versus not imputed data where the causal SNP is not included in one of the datasets.** Each plot shows different sample sizes for one dataset. The variance explained by the causal variant for both the traits is colour coded. Column and row headings are the same as in previous figure. The causal SNP is not included in Illumina 660W panel.

**Figure S3. The relationship between PP4 and the posterior predictive  $p$ -value (on a  $-\log_{10}$  scale) from proportional testing.** Proportional testing uses the BMA approach, integrating over all possible two SNP hypotheses. Each row shows a different scenario, the total number of causal variants in a region is indicated by number of symbols in the plot titles with the type of causal variant indicated by the symbol: full circle - affects both traits; top only - affects one trait; bottom only- affects other trait. For proportional testing, the grey vertical line indicates the threshold ppp of 0.05. Each column shows the total proportion of trait variance for the biomarker explained by all variants in a region, with variance explained spread equally over all variants. In all cases, for the eQTL trait,  $n=1,000$ , 10% of the variance explained by the variant; for the biomarker trait,  $n=10,000$ .

**Figure S4. The relationship between PP4 and the posterior predictive  $p$ -value (on a  $-\log_{10}$  scale) from proportional testing, using subset of SNPs which appear on the Illumina HumanOmniExpress genotyping array.** For the eQTL trait,  $n=1,000$ , 10% of the variance explained by the variant; for the biomarker trait,  $n=10,000$ , 1% or 2% of the variance explained by the variant. Column and row headings are the same as in previous figure.

**Figure S5.** Graphical output using the coloc server (<http://coloc.cs.ucl.ac.uk/coloc/>) for testing genes reported in Teslovich et al. [1] as having a probable shared variant but not supported by our method based on PP4 (posterior probability for a shared signal values)  $< 80\%$ . The Manhattan plots focus on a specific region of the genome with a range of  $\sim 400$  kilobases around the expression probe of the gene specified below each plot. The top plots use the  $-\log_{10}(p\text{-value})$  from the published meta-analysis with one of the four lipid biomarkers; the bottom plots show the  $-\log_{10}(p\text{-value})$  computed by fitting a generalized linear model with expression as dependent variable and SNP genotypes as independent variable. Each cross represents one SNP, imputed or directly typed. The value on the top of each plot shows the PP4 from the colocalisation test between the two top SNP of the expression and biomarker associations.

## Supplemental tables legends

**Table S1. Results using reported loci that colocalise with liver eQTL:** Published results of loci correlating with both liver expression and one of the four lipid traits (Teslovich et al. Supplementary Table 8) and posterior probability of different signal (PP3) and common signal (PP4) after applying colocalisation test. Each row lists the results for one probe, and the multiple entries for the same locus and trait represent multiple probes mapping to the same locus.

**Table S2. eQTL/LDL colocalisation:** Positive (PP4 > 80%) eQTL/LDL colocalisation results between the liver eQTL dataset and the Teslovich meta-analysis. In the column “New”, “Y” indicates that this genes has not been reported to be an intermediate for LDL association by Teslovich et al.

**Table S3. eQTL/HDL colocalisation:** Positive (PP4 > 80%) eQTL/HDL colocalisation results between the liver eQTL dataset and the Teslovich meta-analysis. In the column “New”, “Y” indicates that this genes has not been reported to be an intermediate for LDL association by Teslovich et al.

**Table S4. eQTL/TG colocalisation:** Positive (PP4 > 80%) eQTL/TG (logarithm-transformed) colocalisation results between the liver eQTL dataset and the Teslovich meta-analysis. In the column “New”, “Y” indicates that this genes has not been reported to be an intermediate for LDL association by Teslovich et al.

**Table S5. eQTL/TC colocalisation:** Positive (PP4 > 80%) eQTL/TC colocalisation results between the liver eQTL dataset and the Teslovich meta-analysis. In the column “New”, “Y” indicates that this genes has not been reported to be an intermediate for LDL association by Teslovich et al.