

Environmental and fishing effects on the dynamic of brown tiger prawn (*Penaeus esculentus*) in Moreton Bay (Australia)

Marco Kienzle*, Anthony J. Courtney† and Michael F. O’Neill†

August 20, 2018

Abstract

This analysis of the variations of brown tiger prawn (*Penaeus esculentus*) catch in the Moreton Bay multispecies trawl fishery estimated catchability using a delay difference model. It integrated several factors responsible for variations in catchability: targeting of fishing effort, increasing fishing power and changing availability. An analysis of covariance was used to define fishing events targeted at brown tiger prawns. A general linear model estimated inter-annual variations of fishing power. Temperature induced changes in prawn behaviour played an important role in the dynamic of this fishery. Maximum likelihood estimates of targeted catchability ($3.92 \pm 0.40 \cdot 10^{-4}$ boat-days⁻¹) were twice as large as non-targeted catchability ($1.91 \pm 0.24 \cdot 10^{-4}$ boat-days⁻¹). The causes of recent decline in fishing effort in this fishery were discussed.

1 Introduction

Moreton Bay covers a 1500 km² area on the east coast of Australia (Fig. 1). Its shallow water (< 36 m, average 6.8 m) provide habitats suitable for at least 12 species of prawn (Hyland, 1987), five of which (greasy-back (*Metapenaeus bennettiae*), eastern king (*Melicertus plebejus*), brown tiger (*Penaeus esculentus*), endeavour (*Metapenaeus endeavouri*) and banana prawns (*Fenneropenaeus merguensis*)) are caught by a commercial otter trawl fishery. This industry grew rapidly after 1952 prompting the government to regulate the expansion of the fleet (Parke, 2013). Today, the fishery is managed by input controls in the form of limited entry, vessel and gear restrictions. Spatial closures were introduced in 1993 by the creation of a Marine Park which coverage was extended from 0.5% to 16% of the Bay in 2009. The fishery has provided predominantly small prawns (less than 20 g) to the local market for human consumption and bait. Total catch in 1952–53 was 136 tonnes (t) and reached up to a 1000 t in 1990. In recent years, the total number of vessels in the fishery has declined by 70%, from a peak at 206 vessels to 57 (Fig. 2(a), (Courtney et al., 2012)) in response to falling demand, decreasing prawn prices and increasing fuel costs (Pascoe et al., 2013). Landings’s species composition shifted from being dominated by small size prawns, mostly greasybacks and juveniles eastern king, to an increasing proportion of larger, more valuable, brown tiger prawn (Fig. 2(b)).

*DAFF biometry, Brisbane, Queensland, Australia

†DAFF agri-science, Brisbane, Queensland, Australia.

17 The brown tiger prawn is endemic to Australia and distributed across tropical and subtropical coastal waters
18 (Fig. 1) in depths to 200 m. A characteristic of penaeid coastal shrimps is their short life span, of the order
19 of two years, and their presence in the fishery in significant quantities for a period generally little more than
20 a year (García and Reste, 1981). In Moreton Bay, the population of brown tiger prawns has been assumed
21 to have largely non overlapping generations (Ovenden et al., 2007) because previous estimates of natural and
22 fishing mortality (Wang, 1999; Somers and Wang, 1997) suggested that less than 1% of a cohort would be alive
23 after 1 year of exploitation. Most eggs are produced in a single, clearly defined peak in October–November
24 although spawning continues to May each year (Courtney and Masel, 1997). Larval survival depends on wa-
25 ter temperature and salinity. Nursery habitats for *P. esculentus* are shallow inshore areas prone to estuarine
26 fluctuations of temperature and salinity (Keys, 2003). Peak settlement of juveniles in sea grass in the southern
27 part of the Bay occurs between September and November and late January and April (O’Brien, 1994). Brown
28 tiger prawns recruit to the fishery at a large size (20 g and 27 mm carapace length) compared to the other
29 species (Courtney et al., 1995). Adult brown tiger prawns are benthic and nocturnal, remaining buried during
30 the day and emerging in the evening to feed and mate (Keys, 2003). The duration of nightly emergence from
31 the substrate and the rate of activities such as swimming and foraging are dependent on diurnal rhythms and
32 water temperature (Hill, 1985), light intensity and moulting events (Keys, 2003). Catchability of *P. esculentus*
33 in wild fisheries has been linked with temperatures (White, 1975) and was postulated to play an important role
34 in determining the magnitude of brown tiger catch in Moreton Bay (Hill, 1985).

35
36 The Moreton Bay fishery is a small component of Queensland’s East Coast Otter Trawl fishery (ECOTF,
37 (Pascoe et al., 2013)) which fishing capacity increased significantly since the introduction of otter trawling in
38 this region. New technologies effective at improving fishing efficiency were quickly adopted by this fleet (Robins
39 et al., 1998). Since the mid-1980s, fishing power grew between 0.5 and 4.7% yr⁻¹ depending on the sub-fishery
40 considered (Bishop et al., 2008; O’Neill and Turnbull, 2006; O’Neill and Leigh, 2007). Technology creep was
41 always perceived as a concealed threat to their sustainability. As a consequence they have been closely moni-
42 tored and subject to quantitative stock assessments for many years in order to manage the risk of recruitment
43 overfishing characteristic of tiger prawns (Dichmont et al., 2006). Recent changes in the Moreton Bay trawl
44 fishery have prompted the industry to investigate alternative fishing strategies but a lack of quantitative stock
45 assessment for this area precluded such evaluation. Therefore a delay difference model (Schnute, 1985) was ap-
46 plied to fill this gap. The present stock assessment was implemented to estimate catchability in order to quantify
47 the impact of fishing on the survival of this population of brown tiger prawn. The model was developed to
48 take into account the interactions between environment, technology and fishing effort. A major challenge arose
49 from applying this single species stock assessment model to data from this multispecies fishery and required to
50 define the portion of effort targeted at the brown tiger prawns. Model uncertainties on several aspects of the
51 fishery generated a large amount of plausible stock assessment models that were fitted to the data by maximum
52 likelihood (Burnham and Anderson, 2003) to identify which hypotheses best described historical variations of
53 brown tiger catch in Moreton Bay .

54 2 Materials and methods

55 2.1 Data sources

56 The present assessment of the Moreton Bay stock of brown tiger prawns is based on (a) compulsory commercial
57 logbook data collected since 1988 that provided total catch by species, vessel and day and (b) skipper inter-
58 views conducted in 2000 and 2010 used to construct the history of fishing gear used by each vessel. The fishing

power analysis used this information at the highest temporal resolution (catch by vessel per day). The stock assessment model was fitted to weekly catches grouped into 22 biological years (1989–2010). Each biological year was made of 52 weeks, started around the first of July when brown tiger prawn activity in Moreton Bay was at its lowest and finishing a year later at the end of June.

Logbook records from 1988 to 2010 were combined with vessel and gear descriptions collected during the skipper interviews. The number of nets, total head-rope length, mesh size, type and size of the otter-boards, steaming speed, engine power, propeller diameter, presence/absence of kort nozzle, maximum trawling speed and engine revolution speed were combined using the Prawn Trawl Prediction Model (Sterling, 2005b; Bishop et al., 2008) into an estimate of swept area rate (SAR, in hectares per hour) for each vessel/net configuration available (134 in total). Ten fishing technologies (colour echo-sounder, satellite navigation, global positioning system (GPS), plotter, auto-pilot, GPS coupled with auto-pilot, by-catch-reduction device (BRD) and turtle excluding devices (TED)) were coded as binary variables to indicate presence or absence on-board a vessel during each fishing event. Finally, a continuous variable describing the moon phase was also associated with each logbook record.

Sea surface temperatures in Moreton Bay vary from 16°C in winter to 29°C in summer whereas its range at Cape Moreton is attenuated to 18.5–25.5°C by the ocean (Davie et al., 2011). Seasonal average sea surface temperatures collected within a 60 nautical miles radius around Moreton Island (Anon. (2012b), Fig. 3(c)) were combined with experimental duration of emergence of tiger prawns from Hill (1985) (Fig. 3(b)) to create a seasonal index of brown tiger prawn availability (γ , Fig. 3(d)). This index was made to vary weekly from high availability in summer to low availability in winter. Its amplitude represented a decline in availability of about 50% between summer and winter. This variable was kept constant between years. It was included in several versions of the delay difference model as a multiplier of catchability to determine if the hypothesis that variations of temperature influence the magnitude of brown tiger prawn catch in Moreton Bay (Hill, 1985) was supported by logbooks data.

2.2 Fishing power analysis

In fisheries, catch is often found to increase linearly as a function of effort on the log-scale (Hilborn and Walters, 1992). Multispecies fisheries such as the Moreton Bay fishery exploit different species opportunistically throughout the year as they become available. An un-discriminated analysis of the data showed no relationship between brown tiger catch and fishing effort. This was the result of including records with very low catch rates across the range of effort because non-target species were caught at random or were present on the fishing ground at a lower abundance than the target species (Fig. 4). A rule to classify each unit of fishing effort (in boat-days) into fishing targeted or not targeted at brown tiger prawn was required to analyze these data. A large number of targeting rules were proposed and assessed against the data using an analysis of covariance (ANCOVA, Tab. 3). The data were partitioned according to each rule and both targeted and non-targeted groups of data were fitted with a separate linear regression between catch and effort on the log-scale (Fig. 4). The minimum residual sum of square of an ANCOVA was used to choose the targeting rule that explained the largest variability in the data.

Catch and effort data from fishing records classified as targeted at brown tiger prawns were standardized using a Generalized Linear Model (GLM) to estimate yearly variation of fishing power (Maunder and Punt, 2004). Logarithm of daily brown tiger catch for each vessel (C_i in kg) was expressed as a linear combination of (a) the logarithm of area swept (SA) by the otter trawl estimated as the product of SAR and number of hours

2.3 Stock assessment model

103 fished; (b) several binary variables coding for presence or absence of particular technologies j (represented by
104 matrix $X_{i,j}$); (c) an index of abundance for years, months (taken as factors) and their interaction (represented
105 by matrix $Y_{i,y,m,l}$) and (d) lunar phase (L_i)

$$\log(C_i) = a + b \log(\text{SA}_i) + X_{i,j} c_j + Y_{i,y,m,l} d_{k,l,m} + e L_i \quad (1)$$

106 The parameters of the linear model were estimated in R (R Core Team, 2013) using a Generalized Linear
107 Model (GLM) with quasi family, log-link and variance proportional to the square of mean. An alternative fit
108 using the Gaussian family with log-link provided a poorer fit to the data and was abandoned.

109
110 Multicollinearity in the data set was identified and treated regressing all pairs of explanatory variables
111 against each other (Draper and Smith, 1998) to calculate the variability (R^2) of one variable that was explained
112 by the other (Tab. 4); one variable in each pair was discarded from the pool of possible GLM co-variates when
113 $R^2 > 20\%$. The same procedure using a threshold value of 5% eliminated many more variables from the GLM
114 and was abandoned because it was deemed too stringent. This approach to treat multicollinearity provided
115 stability to the parameter estimates, in particular to the estimates of fishing power.

116
117 An estimate of fishing power variations between 1988 and 2010 was obtained using the GLM to calculate
118 the average catch per hour trawled using a fixed level of abundance (Bishop et al., 2008). These estimates were
119 expressed relative to the beginning of the time-series (1988) were insensitive to the fixed level of abundance
120 chosen. Standard errors for the relative fishing power time-series were obtained by propagating uncertainties
121 from the GLM predictions (Bevington and Robinson, 2003).

2.3 Stock assessment model

2.3.1 Population dynamics

122
123
124
125 A Schnute-Deriso delay-difference model (Schnute, 1985; Deriso, 1980; Hilborn and Walters, 1992) was used to
126 estimate weekly variations in biomass (B_t) of brown tiger prawns in Moreton Bay between 1989 and 2010

$$B_t = s_{t-1} B_{t-1} + \rho s_{t-1} B_{t-1} - \rho s_{t-1} s_{t-2} B_{t-2} - s_{t-1} \rho w_{k-1} R_{t-1} + w_k R_t, \quad 3 \leq t \leq 264 \quad (2)$$

127 Sex-combined growth parameters (ρ , w_{k-1} and w_k), derived from von Bertalanffy estimates (Gribble and
128 Dredge, 1994), were fixed in the model (Tab. 1). This model assumed all prawns were fully recruited to the
129 fishery (knife-edged selectivity) at an age of 22 weeks ($k = 22$), weighing 19.5 grams. w_{k-1} , the pre-recruitment
130 weight was interpreted as a parameter rather than the actual weight at age $k - 1$ and was estimated according
131 to Schnute (1985). Survival (s_t) varied as a function of a fixed natural mortality rate (M , Tab. 1) and fishing
132 mortality (F_t) proportional to effort ($F_t = qE_t$)

$$s_t = \exp[-(M + qE_t)] \quad (3)$$

133 where catchability (q) was estimated.

134
135 This model estimated the magnitude of recruitment in each week (R_t) using 1 parameter to describe total
136 recruitment in each year (R_y) between 1989 and 2010 and 2 parameters (μ , κ) from the von Mises probability
137 density function (Mardia and Jupp, 1999) to allocate a proportion of the total recruitment within each year to
138 each 52 weeks

$$f(x|\mu, \kappa) = \frac{\exp[\kappa \cos(x - \mu)]}{2\pi I_0(\kappa)} \quad (4)$$

139 where $I_0(x)$ is the modified Bessel function of order 0.

140

141 2.3.2 Alternative fishing mortality models

142 Effort was used as a co-variate in the delay difference model to compare the capacity of different fishing mortality
 143 models to explain the weekly variation in brown tiger prawn catch. A total of 7 models were evaluated: the first
 144 model used all effort reported catching tiger prawns (model 1 in Tab. 2); model 2 used effort split into targeted
 145 and non-targeted effort to estimate two coefficients of catchability; the third and fifth models corrected time
 146 series of effort by yearly variations of fishing power (Fig. 6(a)); model 4 and 6 allowed for availability to vary
 147 within years as a function of temperature (Fig. 3). The seventh model was similar to model 4 but non-targeted
 148 effort was not corrected by fishing power.

149 2.3.3 Fitting method

150 A total of 28–29 parameters (1 or 2 catchability parameters, two initial biomass (B_1 and B_2), two von Mises
 151 parameters and 22 annual recruitment parameters and σ , the standard deviation of observation errors) were
 152 estimated by maximum likelihood assuming the square-root of predicted catch (\hat{C}_t , (Quinn and Deriso, 1999))

$$\hat{C}_t = \frac{F_t}{M + F_t} B_t (1 - \exp[-(M + F_t)]) \quad (5)$$

153 provided the mean of a Normal distribution of the square-root of observed catches (C_t) (Dichmont et al., 2003)
 154 with residual standard deviation (σ). The negative log-likelihood function used to fit the model was (Haddon,
 155 2010)

$$-\log(L) = n \log(\sqrt{2\pi}\sigma) + \frac{1}{2\sigma^2} \sum_{t=1}^{n=1144} (\sqrt{\hat{C}_t} - \sqrt{C_t})^2 \quad (6)$$

156 The logarithm was tested as an alternative transformation to the catch data but was abandoned due to
 157 non-normal errors. The model was implemented in C++ and used MINUIT minimization library (James and
 158 Winkler, 2004) available through ROOT (Brun and Rademakers, 1997). Requests regarding the availability of
 159 the code should be directed to the first author of this publication, M. Kienzle.

160 2.3.4 Projections, equilibrium and reference points

161 The proportion of mature female biomass estimated from scientific surveys (Fig 3(a), Courtney and Masel
 162 (1997)) were combined with the estimated stock biomass to calculate spawning stock biomass (SSB) assuming
 163 an even sex-ratio. A Ricker model was fitted to SSB and recruitment estimates lagged by 1 year using linear
 164 regression on transformed data (Hilborn and Walters, 1992). The parameters of this stock-recruitment relation-
 165 ship and uncertainty were used to close the biological cycle by simulating recruitment on the log-scale using
 166 a Gaussian random number generator. The dynamic of the stock at several constant level of exploitation was
 167 projected over a 150-year period to calculate maximum sustainable yield (MSY) and associated effort (E_{MSY}).
 168 In these simulations, effort was distributed within each year according to the average intra-annual pattern
 169 observed between 2006 and 2010.

3 Results

3.1 Fishery statistics

Since the introduction of compulsory logbooks in this fishery in 1988, total catch of all prawn species nearly halved from a 5-year average of 743 ± 107 t at the beginning of the time series to 392 ± 66 t in the last 5 years (Fig. 2(a)). Over the same period, the total number of boats operating in Moreton Bay declined by 2/3 from a median of 198 to 66. Meanwhile, brown tiger prawn catches almost doubled, from an average of 100 tonnes per year before 2000 to an average of 180 tonnes afterwards (Fig. 5). Nominal CPUE fluctuated around 16.0 ± 4.5 kg/boat-day until 2002 before increasing sharply to an average 40.2 ± 8.5 kg/boat-day between 2006 and 2010.

3.2 Effort targeted at brown tiger prawn

Several rules were applied to classify daily individual vessel logbook records into targeted and non-targeted effort at brown tiger prawns. The residual sum of square of an ANCOVA reached a minimum when effort was assumed targeted at brown tiger if this species represented more than 20% of the catch (Tab. 3 and Fig. 4). According to this rule, 90% of tiger catch was associated with targeted effort. Total effort reported catching brown tiger prawn increased from below 6000 boat-days in 1988 up to above 8000 boat-days in 1998–2000 and then declined to around 3500 boat-days (Fig. 5). The proportion of non-targeted effort that was frequently above 30% prior to 2000, declined to around 15% in recent years.

3.3 Fishing power analysis

Vessel identifier (*i.e.* Boat Mark) was confounded with most other co-variates ($R^2 \geq 0.34$) and was eliminated from subsequent analysis (Tab. 4). Auto-pilot was eliminated because it explained 42% of the variability of colour echo-sounder. As a result of this selection process, catch data were fitted with a GLM using the following co-variates: year, month and their interaction; the logarithm of swept area (SA); lunar phase and the presence/absence of satellite navigation system (satnav); differential GPS (dGPS); plotter; GPS coupled with autopilot; computer mapping devices; by-catch reduction devices (BRD); turtle excluding devices (TED) and colour echo-sounder.

Single term deletion from the full model (performed using the *drop1* function in R Core Team (2013)) indicated that abundance terms (year, month and interaction between year and month) explained the largest portion of catch variability (Tab. 5), followed by, in decreasing order of importance, swept area (SA), computer mapping, satnav, plotter, BRD, lunar phase, dGPS, colour echo-sounder, GPS coupled with autopilot and TED. The satellite navigation systems were found to have a positive effect on fishing efficiency: satnav was estimated to improve catch by 25%; dGPS by 7% but effect of GPS was not found to be significant (Tab. 6). Electronic mapping systems were also found to improve catch: computer mapping improved catch by 16% and plotter by 12% respectively. Bycatch reduction devices were found to improve catch of brown tiger by around 10%. On the other hand, turtle excluding devices reduced catch by around 9%. Fishing at full moon was less effective than at new moon. Autopilot coupled with GPS was found to reduce catch rates.

The combined effect of all fishing variables accounted in the GLM estimated fishing power increased by 40–50% from 1988 to 2010 (Fig. 6(a)). Little fishing power variation could be identified between 1988 and 1994. Two large improvements occurred starting in 1994 and 2001, each enhancing the fishing efficiency of this fleet by 20%. Uncertainties associated with this time-series declined through time as the number of skippers providing

210 relevant information for this analysis increased.

211

212 3.4 Stock assessment

213 Catch of brown tiger prawns was characterised by a seasonal variation from a minimum in June–August to a
214 maximum in January–March (Fig. 7(a)). The delay difference model captured both the intra-annual and inter-
215 annual variability but systematically under-estimated large catches. Residuals were normally distributed with
216 a slight tendency to increase over time and an intra-annual pattern was evident (Fig. 7). Both discrepancies
217 were small and did not influence the ranking of hypotheses presented below. Allowing catchability to vary
218 seasonally as a function of temperature provided the largest improvement to the delay difference model (model
219 5 to 6 and model 3 to 4 in Tab. 7). The second biggest improvement was achieved by differentiating catchability
220 associated to targeted effort against non-targeted fishing (model 1 to 2 and model 6 to 4). Relative fishing
221 power corrections to all effort time series improved the fit to a lesser extent (model 2 to 3 and model 1 to
222 5) and reduced estimates of targeted catchability (q_1) by 12–17%. The asymmetric correction of targeted and
223 non-targeted effort time series improved the fit slightly more (model 4 to 7).

224

225 The best description of tiger prawn catch was achieved by combining environmental and fishing effects into
226 model 7. This model estimated targeted catchability equal to $q_1 = 3.92 \pm 0.40 \cdot 10^{-4}$ boat-days⁻¹. A unit
227 of non-targeted fishing effort was estimated to inflict around half the fishing mortality of a unit of targeted
228 effort ($q_2 = 1.91 \pm 0.24 \cdot 10^{-4}$ boat-days⁻¹). 90% of recruitment to the fishery was estimated to occur between
229 mid-November to the end of April and peak at the beginning of February (week 32, Fig. 7(c)). Magnitude
230 of recruitment increased, in average, by a factor of 1.8 before and after 2001 (Fig. 7(d)). A linear regression
231 between recruitment and spawning stock biomass (SSB) was not significant ($P = 0.10$). A fit of the Ricker
232 stock-recruitment relationship showed that this aspect of the dynamic of the stock was the most uncertain
233 (Fig. 7(e)).

234

235 Projections of this stock model indicated that a maximum sustainable yield of 153 ± 50 tonnes can be
236 achieved by applying 5600 targeted-fishing boat-days (in 1989 units), equivalent to 4000 boat-days in 2010 units
237 (Fig. 8(a)). During most its recorded history, the stock of brown tiger prawn in Moreton has been overfished:
238 its spawning stock biomass stayed around 60% of SSB_{MSY} until 2001 and then increased beyond this refer-
239 ence point (Fig. 8(b)). Data from 1989 were not alike those from the same period. Fishing effort was well
240 below E_{MSY} at the beginning of the time series but increased up to 2000 without influencing much the SSB.
241 Between 2000 and 2006, both SSB and fishing effort increased. After 2006, fishing mortality decreased while
242 SSB increased to levels that were below (respectively above) those required to maintain maximum brown tiger
243 prawn production in Moreton Bay. In 2010, the stock was not overfished nor was overfishing occurring.

244

245 4 Discussion

246 The delay difference model was implemented to quantify the impact of fishing on survival of brown tiger prawn
247 in Moreton Bay. Defining fishing effort targeted at brown tiger prawn provided an important improvement to
248 the fit, consistent with Zhou et al. (2011) recommending to weight these time-series differently in stock assess-
249 ments. This model of the fishery can address the effect of shifting effort on and off tiger prawn but further
250 research is needed to address shifts in effort between species in this multispecies fishery. Maximum likelihood

251 estimate of targeted catchability ($3.92 \cdot 10^{-4}$ boat-days⁻¹) was about 5 times larger than $8.1\text{--}8.8 \cdot 10^{-5}$ estimated
252 by Zhou et al. (2011) and Wang (1999) in the NPF. This difference reflects to some extent the difference in stock
253 size between these areas, Moreton Bay being 200 times smaller than the NPF. The model for stock-recruitment
254 relationship was by far the most uncertain aspect of this fishery. The lack of significant linear relationship
255 between recruitment and SSB is not un-expected given (a) the small number of observations and (b) the large
256 variability in recruitment. The projections clearly encompassed the range of catch observed in this fishery over
257 the past 20 years.

258
259 This stock assessment indicated that recent levels of exploitation of brown tiger prawn in Moreton Bay
260 were sustainable. Increases in brown tiger prawn catch and catch rates were associated with declining fleet size
261 and effort. The delay difference model estimated that recruitment increased simultaneously suggesting that
262 brown tiger prawn has recovered from recruitment overfishing. Economics is the most likely driver of effort
263 decline in this fishery: Australia almost doubled its import of prawn in the past 10 years, which account today
264 for over 60% of the total consumption of prawn in this country (Anon., 2011). Imports of larger volumes of
265 aquaculture production, in particular the white leg shrimp (*Penaeus vannamei*), have increased consumption
266 and commercialization of species that were once primarily caught by local fishermen. The rapid decline of
267 crustacean prices, 30–40% (Anon., 2010; Adams et al., 2005; Curtotti et al., 2011), was strongly correlated with
268 total effort in the fishery ($\rho = 0.79$). This accentuated competition in the seafood market affected the revenue of
269 many fishing operations whose profits had already been eroded by years of increasing fuel prices (Anon., 2012a;
270 Sterling, 2005a). These two economic factors have certainly affected the fishery bionomic equilibrium (Clark,
271 1990) especially because trawling is an energy intensive fishing method. The changes observed in Moreton Bay
272 trawl fishery in recent years are probably the result of aquaculture reducing pressure on wild fish stocks. The
273 likelihood of fishing effort becoming a threat to the sustainability of brown tiger prawn harvest in Moreton Bay
274 is low given that the fundamental economic drivers of the fishery are not likely to improve in a foreseeable future.

275
276 The simultaneous increase in SSB and effort targeted at tiger prawn between 2000 and 2006 was peculiar
277 and suggested that factors other than those accounted for in this analysis might have affected the dynamic
278 of the stock. Between those, a reduction of growth overfishing could explain a reduction in mortality and in-
279 crease in brown tiger catch. The relative importance of this hypothesis was difficult to assess given the lack
280 of information regarding the variation of catch's size-composition throughout the entire time-series. On the
281 other hand, improvements in habitat, such as increased seagrass area or recent Marine Park closures in Moreton
282 Bay, are unlikely to have contributed to the increase in *P. esculentus* population size. Hyland et al. (1989)
283 mapped the seagrass densities and distributions in Moreton Bay in the late 1980s and to our knowledge there
284 are no evidence to suggest that these habitats have extended significantly since. In fact, large reclamations
285 of intertidal areas associated with expansion of the Port of Brisbane have probably contributed to a decline in
286 such habitats for tiger prawns. The Queensland Government closed some area in the Bay to trawling in 2009
287 under the Moreton Bay Marine Park Plan. However, while these areas may have had a positive impact on the
288 brown tiger population size, they occurred after the dramatic decline in effort and after the population showed
289 signs of recovery.

290
291 The estimated average rate of increase in fishing power in Moreton Bay of $1.7\% \text{ yr}^{-1}$ is at the higher end of
292 the range estimated for the fleet operating on the east coast of Australia ($0.5\text{--}1.8\% \text{ yr}^{-1}$) for the same period
293 (O'Neill et al., 2003; O'Neill and Leigh, 2007)) and at the lower end of envelope of the lower possible cases (1.8%
294 and 2.8%) estimated by (Bishop et al., 2008; Zhou et al., 2011) for 1980–2007. This difference might result from
295 (a) an intrinsic difference in fishing improvements between smaller vessels (< 14 m) operating in the sheltered
296 waters of Moreton Bay compared to vessels operating in the Gulf of Carpentaria capable of long fishing trip

297 requiring to withstand bad weather at sea; or (b) statistical treatment to allow for the possibility of greater
298 impacts of technologies, referred to as the "high" treatment (Bishop et al., 2008). The steep increase in fishing
299 power starting in 1995 corresponds to the adoption of GPS in the fleet O'Neill et al. (2003). The increase in
300 fishing power starting in 2003 is not clearly associated with any technological improvement and might be the
301 result of less efficient boat leaving the fishery. The year term in catch rates standardizations (Maunder and
302 Punt, 2004) corresponds to an indice of abundance, providing the size of areas fished, and the spatial pattern
303 of effort, have remained constant over the years (Bishop et al., 2008). The present analysis did not account for
304 spatial information which might result in interpreting wrongly variations of catch rates as a result of variations
305 of abundance and fishing gear rather than a change in fishing locations (Campbell, 2004; Walters, 2003).

306
307 A range of hypotheses were compared to determine which influenced most the dynamic of this fishery.
308 Temperature was found to determine the magnitude of catch by changing the duration of emergence of brown
309 tiger prawns (Hill, 1985). This behavioural change is related to the frequency of feeding that dependent on
310 metabolic and digestive rates regulated by ambient temperatures in aquatic poikilotherms (Fonds et al., 1992).
311 The time-series of temperatures used in this analysis under-estimates the amplitude of variations in the Bay
312 because shallow waters warm and cool faster in response air temperature than larger bodies of seawater. Such
313 difference could explain the discrepancies between the model and the data evidenced by the residues's weekly
314 pattern. Future research should include time-series of temperature collected *in-situ* and assess how they im-
315 prove the fit of the delay difference model. Moreover this work provides a framework to evaluate the effect of
316 climate change on the dynamic of tiger prawn fisheries. Given the present results, a rise in water temperature
317 is expected to benefit fishermen in Moreton Bay by increasing brown tiger prawn catchability. The importance
318 of this effect relative to, for example, fishing power increases could be quantified. Nevertheless, other effects of
319 temperature such as its influence on timing and length of spawning season will need to be included to provide
320 a comprehensive model of the effects of climate change on this tiger prawn fishery.

322 Acknowledgements

323 This work was developed as part of the "Harvest strategy evaluations and co-management for the Moreton
324 Bay Trawl Fishery" Cooperative Research Center project 2009/774. We are grateful for the many discussions
325 with industry participants and representatives, in particular Dr. D. Sterling for estimating trawl swept area
326 rates using his prawn trawl performance model and Mr M. Wood for promoting a better understanding of this
327 fishery. We are grateful to Dr. G. Leigh, Dr. A. Campbell, Prof. Y.G. Wang and Dr. J. Robins for the many
328 discussions on this topic. We thank Dr N. Wade, Dr. A. Norman and Dr W. N. Venables from the CSIRO for
329 pointing relevant literature.

330 References

- 331 Adams, C., Keithly, W., Versaggi, S., 2005. The shrimp import controversy, in: Schmitz, A., Moss, C.B.,
332 Schmitz, T.G., Koo, W.W. (Eds.), International Agriculture Trade Disputes: Case Studies In North America.
333 Calgary: University of Calgary Press. chapter 15, pp. 225–244.
- 334 Anon., 2010. Economic Indicators for the Spencer Gulf and West Coast Prawn Fisheries. Technical Report.
335 Econsearch.

- 336 Anon., 2011. Australian fisheries statistics 2011. Dept. of Agriculture, Fisheries and Forestry, Australian
337 Government.
- 338 Anon., 2012a. Agricultural commodity statistics 2012. Dept. of Agriculture, Fisheries and Forestry, Australian
339 Government.
- 340 Anon., 2012b. Hydrography, Meteorology and Oceanography services. Australian Government – Department
341 of Defence. URL: www.metoc.gov.au. last accessed on 9th January 2012.
- 342 Bevington, P.R., Robinson, D.K., 2003. Data reduction and error analysis. 3rd ed., Mac Graw Hill.
- 343 Bishop, J., Venables, W.N., Dichmont, C.M., J., S.D., 2008. Standardizing catch rates: is logbook information
344 by itself enough ? ICES Journal of Marine Science 65, 255–266.
- 345 Brun, R., Rademakers, F., 1997. Root - an object oriented data analysis framework, in: Proceedings AI-
346 HENP'96 Workshop, Lausanne, Sep. 1996, Nucl. Inst. & Meth. in Phys. Res. A 389, pp. 81–86. See also
347 <http://root.cern.ch/>.
- 348 Burnham, K.P., Anderson, D., 2003. Model Selection and Multi-Model Inference. 2nd ed., Springer-Verlag.
- 349 Campbell, R.A., 2004. CPUE standardisation and the construction of indices of stock abundance in a spatially
350 varying fishery using general linear models. Fisheries Research 70, 2009–227.
- 351 Clark, C., 1990. Mathematical bioeconomics: the optimal management of renewable resources. Pure and applied
352 mathematics, Wiley.
- 353 Courtney, A., Kienzle, M., Pascoe, S., O'Neill, M., Leigh, G., Wang, Y.G., Innes, J., Landers, M., Braccini, M.,
354 Prosser, A., Baxter, P., Sterling, D., Larkin, J., 2012. Harvest strategy evaluations and co-management for
355 the Moreton Bay Trawl Fishery. Technical Report Project 2009/774. Australian Seafood CRC.
- 356 Courtney, A., Masel, J., 1997. Spawning stock dynamics of two penaeid prawns, *Metapenaeus bennettiae* and
357 *Penaeus esculentus*, in Moreton Bay, Queensland, Australia. Marine Ecology Progress Series 148, 37–47.
- 358 Courtney, A., Masel, J., Die, D., 1995. Temporal and spatial patterns in recruitment of three penaeid prawns
359 in Moreton Bay, Queensland, Australia. Estuarine, Coastal and Shelf Science 41.
- 360 Curtotti, R., Hormis, M., McGill, K., Pham, T., Vieira, S., Perks, C., George, D., 2011. Australian fisheries
361 - outlook and economic indicators, in: Outlook conference, Australian Bureau of Agricultural and Resource
362 Economics and Sciences (ABARES).
- 363 Davie, P., Cranitch, G., Wright, J., Cowell, B., 2011. Wild guide to Moreton Bay and adjacent coasts. 2nd ed.,
364 Queensland Museum, Brisbane, Australia.
- 365 Deriso, R., 1980. Harvesting strategies and parameter estimation for an age-structured model. Canadian Journal
366 of Fisheries and Aquatic Sciences 37, 268–282.
- 367 Dichmont, C., Deng, A., Punt, A., Venables, W., Haddon, M., 2006. Management strategies for short-lived
368 species: The case of Australia's Northern Prawn Fishery: 1. Accounting for multiple species, spatial structure
369 and implementation uncertainty when evaluating risk. Fisheries Research 82, 204–220.

- 370 Dichmont, C., Punt, A., Deng, A., Dell, Q., Venables, W., 2003. Application of a weekly delay-difference model
371 to commercial catch and effort data for tiger prawns in australia's northern prawn fishery. *Fisheries Research*
372 65, 335–350.
- 373 Draper, N., Smith, H., 1998. *Applied Regression Analysis*. 3rd ed., Wiley & Sons.
- 374 Fonds, M., Cronie, R., Vethaak, A., Puyl, P.V.D., 1992. Metabolism, food consumption and growth of plaice
375 (*pleuronectes platessa*) and flounder (*platichthys flesus*) in relation to fish size and temperature. *Netherlands*
376 *Journal of Sea Research* 29, 127–143.
- 377 García, S., Reste, L., 1981. Life cycles, dynamics, exploitation and management of coastal penaeid shrimp
378 stocks. FAO fisheries technical paper, FAO.
- 379 Grey, D., Dall, W., Baker, A., 1983. A guide to the Australian Penaeid Prawns. Technical Report 2005/239.
380 Department of Primary Production.
- 381 Gribble, N., Dredge, M., 1994. Mixed-species yield-per-recruit simulations of the effect of seasonal closure on a
382 central queensland coastal prawn trawling ground. *Canadian Journal of Fisheries and Aquatic Sciences* 51,
383 998–1010.
- 384 Haddon, M., 2010. *Modelling and quantitative methods in fisheries*. Chapman & Hall/CRC.
- 385 Hilborn, R., Walters, C., 1992. *Quantitative fisheries stock assessment: choice, dynamics and uncertainty*.
386 Chapman and Hall.
- 387 Hill, B., 1985. Effect of temperature on duration of emergence, speed of movement, and catchability of the
388 prawn *Penaeus esculentus*, in: Rothlisberg PC, Hill BJ, S.D. (Ed.), *Proceedings of the Second Australian*
389 *National Prawn Seminar*, Simpson Halligan, Brisbane, Australia. pp. 77–83.
- 390 Hyland, S., 1987. An investigation of the nekto-benthic organisms in Logan River and Moreton Bay (Queensland)
391 with an emphasis on penaeid prawns. Ph.D. thesis. University of Queensland.
- 392 Hyland, S.J., Courtney, A.J., Butler, C.T., 1989. Distribution of seagrass in the Moreton Region from Coolan-
393 gatta to Noosa. Information Series QI89010. Queensland Dept of Primary Industry.
- 394 James, F., Winkler, M., 2004. *Minuit user's guide*. <http://www.cern.ch/minuit>.
- 395 Keys, S., 2003. Aspects of the biology and ecology of the brown tiger prawn, *penaeus esculentus*, relevant to
396 aquaculture. *Aquaculture* 217, 325 – 334.
- 397 Mardia, K., Jupp, P., 1999. *Directional statistics*. Wiley.
- 398 Maunder, M.N., Punt, A.E., 2004. Standardizing catch and effort data: a review of recent approaches. *Fisheries*
399 *Research* 70, 141–179.
- 400 O'Brien, C., 1994. Population dynamics of juvenile tiger prawns *Penaeus esculentus* in south Queensland,
401 Australia. *Marine Ecology Progress Series* 104, 247–256.
- 402 O'Neill, M., Courtney, A., Turnbull, C., Good, N., Yeomans, K., Staunton Smith, J., Shootingstar, C., 2003.
403 Comparison of relative fishing power between different sectors of the queensland trawl fishery, australia.
404 *Fisheries Research* , 309–321.

- 405 O'Neill, M., Leigh, G., 2007. Fishing power increases continues in Queensland's east coast trawl fishery. *Fisheries*
406 *Research* 85, 84–92.
- 407 O'Neill, M., Turnbull, C., 2006. Stock assessment of the Torres Strait tiger prawn fishery (*Penaeus esculentus*).
408 Information Series QI05132. Queensland Dept of Primary Industry and Fisheries.
- 409 Ovenden, J., Peel, D., Street, R., Courtney, A., Hoyle, S., Peel, S., Podlich, H., 2007. The genetic effective and
410 adult census size of an Australian population of tiger prawns (*Penaeus esculentus*). *Molecular Ecology* 16,
411 127–138.
- 412 Parke, J., 2013. Against the tide: Queensland's Moreton Bay fishing industry since 1824. 5Word Productions.
- 413 Pascoe, S., Dichmont, C., Brooks, K., Pears, R., Jebreen, E., 2013. Management objectives of queensland
414 fisheries: Putting the horse before the cart. *Marine Policy* 37, 115–122.
- 415 Quinn, T.J., Deriso, R.B., 1999. Quantitative fish dynamics. Oxford University Press.
- 416 R Core Team, 2013. R: A Language and Environment for Statistical Computing. R Foundation for Statistical
417 Computing. Vienna, Austria. URL: <http://www.R-project.org/>.
- 418 Robins, C.M., Wang, Y., Die, D., 1998. The impact of global positioning systems and plotters on fishing power
419 in the Northern Prawn Fishery. *Canadian Journal of Fisheries and Aquatic Sciences* 55.
- 420 Schnute, J., 1985. A general theory for analysis of catch and effort data. *Canadian Journal of Fisheries and*
421 *Aquatic Sciences* 42, 414–429.
- 422 Somers, I., Wang, Y.G., 1997. A simulation model for evaluating seasonal closures in australia's multispecies
423 northern prawn fishery. *North American Journal of Fisheries Management* 17, 114–130.
- 424 Sterling, D., 2005a. Fishing energy efficiency review for the fisheries Research and development corporation.
425 Technical Report 2005/239. Fisheries Research and Development Corporation.
- 426 Sterling, D., 2005b. Modelling the physics of prawn trawling for fisheries management. Ph.D. thesis. School of
427 applied physics. Curtin University of Technology, Perth, Australia.
- 428 Walters, C., 2003. Folly and fantasy in the analysis of spatial catch rate data. *Canadian Journal of Fisheries*
429 *and Aquatic Sciences* 60, 1433–1436.
- 430 Wang, Y., 1999. A maximum-likelihood method for estimating natural mortality and catchability coefficient
431 from catch and effort data. *Marine & Freshwater Research* 50, 307–11.
- 432 White, T., 1975. Factors affecting the catchability of a penaeid shrimp *penaeus esculentus*., in: Young, P. (Ed.),
433 Proceedings of the First Australian National Prawn Seminar, AGPS, Canberra, Australia. pp. 115–137.
- 434 Zhou, S., Punt, A.E., Deng, R., Bishop, J., 2011. Estimating multifleet catchability coefficients and natural
435 mortality from fishery catch and effort data: comparison of bayesian statespace and observation error models.
436 *Canadian Journal of Fisheries and Aquatic Sciences* 68, 1171–1181.

Figures caption

Fig. 1: Top map: the spatial distribution of the brown tiger prawn (*Penaeus esculentus*) in Australia (from Grey et al. (1983)) with the location of Moreton Bay indicated by a black square. Bottom map: the location of trawling ground covering an area of about 800 km² (striped area). The dots represent areas closed to trawling including habitat protection zones, conservation and marine national parks.

Fig. 2: Left: Recent trends in catch of all species of prawn and number of vessels fishing in Moreton Bay. Right: Species catch composition by year.

Fig. 3: Some relevant biological information: (a) proportion of mature female biomass; (b) duration of emergence determined experimentally by Hill (1985); (c) monthly average seawater temperature in Moreton Bay and (d) estimated indices of availability of brown tiger prawn in Moreton Bay.

Fig. 4: Relationship between total brown tiger prawn catch and effort (on the log-scale) by boat and year for fishing events targeted at brown tiger prawn or not.

Fig. 5: Time series of brown tiger prawn (*Penaeus esculentus*) catch and catch per unit of effort in Moreton Bay (left panel); time series of effort catching tiger prawn in Moreton Bay (right panel).

Fig. 6: (a) Estimated changes in fishing power relative to 1988 based on a GLM of brown tiger prawn catch. The vertical bars indicate 2 standard errors from the mean. (b) Variations of fishing effort corrected by fishing power.

Fig. 7: (a) Time series of residuals of the delay difference model fit (b) Proportion of recruitment in each month estimated using the von Mises distribution. (c) Time series of recruitment estimated from model 3. (d) Estimated relationship between spawning stock biomass and recruitment fitted with a Ricker function. The dotted lines represent 95% confidence interval of predictions from this model.

Fig. 8: (a) Simulated long-term yield of brown tiger prawn at fixed level of effort to determine maximum sustainable yield (MSY). (b) Trajectory of the fishery through time in relation to spawning stock biomass at MSY (x-axis) and effort at MSY (y-axis). Note that effort (E) on this graph correspond to targeted effort corrected by both fishing power and availability.

470 **Figures**




Figure 1:




Figure 2:




Figure 3:




Figure 4:




Figure 5:







Figure 6:




(a)




(b)



(c)



(d)



(e)

Figure 7:





Figure 8:

471 **Tables**

Parameter	Value	Reference
ρ	0.963	Gribble and Dredge (1994)
w_{k-1}	17.8 grams	based on the method of Schnute (1985)
w_k	19.5 grams	based on the method of Schnute (1985)
M	0.045 week ⁻¹	Dichmont et al. (2003)

Table 1: Values of parameters fixed in the delay difference model.

Model	Fishing mortality
1	$F_t = q_1 E_t$
2	$F_t = q_1 E_t(\text{targeted}) + q_2 E_t(\text{non targeted})$
3	$F_t = \beta (q_1 E_t(\text{targeted}) + q_2 E_t(\text{non targeted}))$
4	$F_t = \gamma \beta (q_1 E_t(\text{targeted}) + q_2 E_t(\text{non targeted}))$
5	$F_t = \beta q_1 E_t$
6	$F_t = \gamma \beta q_1 E_t$
7	$F_t = \gamma \beta q_1 E_t(\text{targeted}) + \gamma q_2 E_t(\text{non targeted})$

Table 2: Fishing mortality equations used in each delay difference model.

Definition of targeting	Res. SSQ
tiger / total > 0.1	18470
tiger / total > 0.2	17203
tiger / total > 0.3	17281
tiger / total > 0.4	17721
tiger / total > 0.5	18416
tiger / total > 0.6	19248
tiger / total > 0.7	20021
tiger / total > 0.8	21335
tiger / total > 0.9	22644
tiger > banana & tiger > greasyback & tiger > king	18029
banana = 0 & king = 0	22583

Table 3: Comparison of residual sum of squares (SSQ) of ANCOVAs of brown tiger prawn catch and effort on the log-scale for a variety of targeting definitions. The term "total" refers to the sum of tiger, banana, greasyback and eastern king prawns catch.

	Boat Mark	record number	Swept area	Lunar Quarters	colour echo-sounder	satnav	GPS	dGPS	plotter	autopilot	GPS coupled autopilot	GPS coupled radar	computer mapping	BRD	TED
Boat Mark	1.00														
record number		1.00													
Swept area	0.34	0.35	1.00	0.00	0.01	0.04	0.02	0.00	0.02	0.02	0.09	0.00	0.02	0.00	0.00
Lunar Quarters				1.00											
colour echo-sounder	0.95	0.95	0.01	0.00	1.00	0.01	0.00	0.01	0.01	0.42	0.00	0.01	0.10	0.01	0.00
satnav	0.74	1.00	0.04	0.00	0.01	1.00	0.00	0.01	0.01	0.00	0.11	0.00	0.07	0.03	0.00
GPS	0.69	0.79	0.02	0.00	0.00	0.00	1.00	0.00	0.19	0.00	0.05	0.01	0.10	0.04	0.00
dGPS	0.89	0.93	0.00	0.00	0.01	0.01	0.00	1.00	0.02	0.01	0.00	0.00	0.06	0.01	0.00
plotter	0.82	0.92	0.02	0.00	0.01	0.01	0.19	0.02	1.00	0.03	0.00	0.01	0.00	0.00	0.00
autopilot	0.95	0.95	0.02	0.00	0.42	0.00	0.00	0.01	0.03	1.00	0.03	0.01	0.05	0.00	0.00
GPS coupled autopilot	0.93	0.93	0.09	0.00	0.00	0.11	0.05	0.00	0.00	0.03	1.00	0.09	0.18	0.01	0.01
GPS coupled radar	0.62	0.62	0.00	0.00	0.01	0.00	0.01	0.00	0.01	0.01	0.09	1.00	0.02	0.00	0.00
computer.mapping	0.83	0.93	0.02	0.00	0.10	0.07	0.10	0.06	0.00	0.05	0.18	0.02	1.00	0.07	0.02
BRD	0.47	0.55	0.00	0.00	0.01	0.03	0.04	0.01	0.00	0.00	0.01	0.00	0.07	1.00	0.02
TED	0.40	0.54	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.02	0.02	1.00

Table 4: Proportion of variability explained (R^2) by a pair-wise linear regression using a single variable (in rows) as the dependent variable and a single variable (in column) as the explanatory variable.

	Df	Deviance	scaled dev.	Pr(> χ^2)
<none>		5458.53		
log(SA)	1	5908.35	1170.22	0.0000
BRD	1	5462.25	9.69	0.0019
TED	1	5460.71	5.69	0.0171
colour echo sounder	1	5461.75	8.38	0.0038
dGPS	1	5461.95	8.91	0.0028
satnav	1	5499.25	105.94	0.0000
plotter	1	5482.69	62.87	0.0000
GPSCoupledautopilot	1	5460.90	6.18	0.0129
computer mapping	1	5507.21	126.65	0.0000
lunar	1	5462.11	9.32	0.0023
Year:Month	244	10562.27	13277.33	0.0000

Table 5: Assessment of the effect of removing a single variable from the full GLM

	Estimate	Std. Error	t value	Pr(> t)
log(SA)	0.6384	0.0175	36.55	0.0000
BRD	0.0955	0.0308	3.10	0.0019
TED	-0.0907	0.0378	-2.40	0.0165
colour echo sounder	0.0579	0.0193	3.00	0.0027
dGPS	0.0688	0.0230	2.99	0.0028
satnav	0.2254	0.0217	10.37	0.0000
plotter	0.1104	0.0136	8.13	0.0000
GPSCoupledautopilot	-0.0332	0.0130	-2.56	0.0106
computer mapping	0.1521	0.0133	11.41	0.0000
lunar	-0.0450	0.0147	-3.07	0.0021
Year1990:Month01	-0.8020	0.6268	-1.28	0.2008
...

Table 6: Parameter estimates for the GLM co-variates.

Model	$-\log(L)$	Catchability		Recruitment distribution		Biomass		
		$q_1 (\times 10^{-4})$	$q_2 (\times 10^{-4})$	μ	κ	B(1)	B(2)	σ
7	3659.05	3.92 ± 0.4	1.91 ± 0.24	0.64 ± 0.03	2.02 ± 0.08	0.22 ± 0.01	0.3 ± 0.01	5.93 ± 0.13
4	3669.41	4.02 ± 0.42	1.7 ± 0.22	0.65 ± 0.03	2.04 ± 0.09	0.22 ± 0.02	0.3 ± 0.02	5.98 ± 0.13
6	3707.42	1.51 ± 0.19		0.51 ± 0.02	5.32 ± 0.58	0.5 ± 0.29	0.63 ± 0.26	6.18 ± 0.13
3	3759.51	3.96 ± 0.31	1.76 ± 0.2	0.49 ± 0.03	2.38 ± 0.12	0.06 ± 0.01	0.13 ± 0.01	6.47 ± 0.14
2	3764.42	4.9 ± 0.38	1.87 ± 0.22	0.46 ± 0.03	2.27 ± 0.12	0.05 ± 0.05	0.12 ± 0.05	6.5 ± 0.14
5	3801.99	3 ± 0.27		0.5 ± 0.02	3.46 ± 0.27	0.06 ± 0.06	0.13 ± 0.05	6.72 ± 0.14
1	3819.91	3.38 ± 0.27		0.48 ± 0.02	3.65 ± 0.29	0.06 ± 0.06	0.12 ± 0.05	6.82 ± 0.15

Table 7: Comparison of the negative log-likelihood and parameters estimates of different models. The results are ordered by increasing value of negative log-likelihood from top to bottom.