

Drake's Rule as a Consequence of Approaching Channel Capacity

Alexey A. Shadrin^{1,2}, Dmitri V. Parkhomchuk^{3*}

¹Fachbereich Mathematik und Informatik, Freie Universität Berlin, Berlin, Germany

²Department of Vertebrate Genomics, Max Planck Institute for Molecular Genetics, Berlin, Germany

³Institut für Medizinische Genetik und Humangenetik der Charité – Universitätsmedizin Berlin, Berlin, Germany

Email: *pdmitri@hotmail.com

Abstract

How mutations accumulate in genomes is the central question of molecular evolution theories. However, our understanding of this process is far from complete. Drake's rule is a notoriously universal property of genomes from microbes to mammals – the number of (functional) mutations per-genome per-generation is approximately constant within a phylum, despite orders of magnitude differences in genome sizes and diverse populations' properties. So far there is no concise explanation for this phenomenon. A formal model of storage of genetic information suggests that a genome of any species operates near its maximum informational storage capacity, and the mutation rate is near its upper limit, providing a simple explanation for the rule with minimal assumptions.

Keywords: Drake's rule, information theory, neutral theory

Introduction

For about a hundred years the key parameter in modeling of Darwinian selection is “fitness” – it defines which organisms are left to live and reproduce in a population and which have to be eliminated. Alleles or mutations (“variants”) are presumed to affect the fitness and a variant destiny (its frequency dynamics in a population) is traced with some mathematical models. There is a vast amount of models with different assumptions about how to model real populations “correctly”. For example the “Moran process” [1] defines an elementary time step as either death or reproduction of a random individual – “overlapping generation”, deriving the analytical solutions for some simple scenarios. Alternatively, the “Wright-Fisher model” [2] presumes the non-overlapping generations, such as annual plants. Then there is a question of how to calculate the cumulative fitness for a few independent variants, account for the effects of newly appearing variants, and many other subtleties. In the traditional models the fitness is “relative”, without any fixed “baseline” – an individual cannot be assigned a fitness value ignoring the rest of the population. It is impossible to compare a fitness of an elephant to a fitness of yeast. Such fitness keeps no population history – a gain (or loss) of fitness for a whole population is untraceable, because after the gain the organisms are competing with each other in formally the same way. So the evolution is presented as an opportunistic non-directional “Brownian” motion – fixation of accidental “positive” mutations. However, it would be desirable to have a measure, which is “absolute” – having a baseline and reflecting the organismal complexity – the total “genetic information” or “evolutionary progress”. Then, not only we can compare different species but also use it as a fitness function within a population, for modeling.

Despite the numerosity of models their explanatory power remains arguably limited, so that in 1996 Ohta and Gillespie admitted “a looming crisis” – “all current theoretical models suffer either from assumptions that are not quite realistic or from an inability to account readily for all phenomena.” [3]. Probably the limits of the current models are rooted in the basic fitness definition, because if it is similar in all models, the reshuffling of other

parameters will not drastically change the behavior and predictions on a fundamental level. Recently we proposed an information-theoretical model [4], which can provide a measure, which is “absolute”, estimates the total genomic information and can be used for the fitness calculations, sensibly accounting for interactions of any number of variants in a genome. Such fitness function is the most essential difference of our model from the traditional approaches, while the modes of reproduction and other parameters are of secondary importance.

Random mutations deteriorate the genomic information and must be compensated by selection and here we evaluate by simulations some simple scenarios of such process under the equilibrium condition. With some arguably plausible assumptions such process readily explains Drake's rule [5, 6, 7].

Methods

Information in sequence patterns

The measure of genetic information (GI) was proposed by Schneider et al. [8]. During the last 25 years it became a popular tool for investigation of binding sites patterns [9, 10]. It adapts the information theory (IT) concept of Shannon's entropy [11] in the context of biological sequences.

The acceptable variability in each position (P) is defined by the frequencies of four nucleotides in an equilibrium population and quantified by Shannon's entropy:

$$H(P) = - \sum_{N \in \{A, G, C, T\}} f_N \log_2 f_N \quad (1)$$

Where f_B , $B \in \{A, G, C, T\}$ is a frequency of nucleotide B at the position P . The genetic information for a single position is defined as: $GI(P) = 2 - H(P)$. One possible interpretation is that such function conveniently (additively and linearly) quantifies the amount of biases in alleles' equilibrium distribution.

For simplicity, we (after Schneider et al.) assume independent positions in patterns. However there are no indications that assuming some positional dependencies in patterns would drastically influence the main conclusions. Inevitably any model contains a set of assumptions representing a simplification of reality, though the simplicity cannot be immediately considered as model's shortcomings – more often it is the advantage. Recently we showed [4] that the sum of GIs can serve as a measure of positional information. This “additivity” should not be confused with a simple additivity of Shannon's entropy – the problem is to prove that the sum of GIs for a functional site (or a genome) is linearly linked to the “positional information” (specificity of molecular interactions). One could use some other measure of frequencies biases – why is the defined one “fundamental”? In order for the sum of site's GIs to have the informational meaning, the number of possible functional sequences for the site (the size of its “typical set”) must depend exponentially on the defined site's variability (the value reciprocal to the sum of GIs). This exponential dependence is the non-trivial result of IT (Shannon-McMillan-Breiman theorem). The corresponding “natural choice” of the logarithmic function for information measure is discussed in details in the classical Shannon's paper [11]. With such well-defined positional information measure it is possible to build a formal (“mechanistic”) model of “molecular machines” evolution.

Then we can use the position-specific GIs to calculate the total amount of information contained in a genome as a simple sum over all positions:

$$GI_{total} = 2L + \sum_{j=1}^L \sum_{B \in \{A, G, C, T\}} f_{jB} \log_2 f_{jB} \quad (2)$$

Where f_{jB} , $1 \leq j \leq L$, $B \in \{A, G, C, T\}$ is the frequency of nucleotide B at the position j . Let's also define an average density of genetic information in a population as $GI_p = GI_{total}/L$ bits per site. It is obvious that $0 \leq GI_p \leq 2$.

Hence a functional site (or a genome) is represented by the corresponding pattern – “ GI profile”, so that GIs and the corresponding 4-vectors of the acceptable equilibrium frequencies are defined in each position. As we discussed in [4] the equilibrium condition is important for the correct GI definition and measurement, while real populations are far from the equilibrium, in general. Importantly, the GI profile is the “prior”, inherent property of molecular functionality, for example a protein domain can be functional only within a certain set of sequences, e.g. in GI terms, a conserved domain has high GI value and a small “typical set”. Then the average density GI_p cannot be significantly different in close species – functional genes are conserved similarly, unless some novel mechanisms of molecular functioning are introduced. The actual variability in a population depends on this predefined GI and population history. The equilibrium population which we simulated here (effectively canceling out, “erasing” the history influence, revealing the unobscured “pure functionality” profile), is necessary for the correct GI measurement and the determination of an “error threshold”, however, a slice (small subset) of such population will have the same mutational properties as the whole equilibrium population, but smaller variability. Such subset represents a realistic population, which recently (relatively to mutation rate) underwent a bottleneck and experienced a “founder effect” - all individuals are closely related through a few recent population founders. The variability in this subset is not reflecting correctly the GI profile, however, this profile still “exists”, though more in a platonic sense. It could be revealed if this subset was allowed to diverge for sufficiently long time without disturbances.

Such regimes of mutagenesis and maintenance of variability are similar to those in quasispecies theories: “The quasispecies concept becomes important whenever mutation rates are high. This is often the case in viral and bacterial populations.” [12]. In these theories a population is represented by a “cloud” of diverged genotypes. However, the distinction between “normal” species and quasispecies is blurred, and nothing can prevent us from viewing a “normal” population as the aforementioned subset of quasispecies (in the process of divergence). Here we assume that this regime of high mutation rate is exactly the one, which deserves careful examination in large genomes of higher organisms as well – what matters is the mutation rate per-genome per-generation, and as we now know, this parameter is quite large in mammals also - about few hundreds mutations, with few in coding regions – actually that is the main point of Drake's rule. For simplicity we presume, that the selection has an opportunity to act compensatory (to increase GI) only in-between generations, ignoring possible germ-line selection issues - that is the reason for focusing on the per-generation mutation rates. In comparison, the quasispecies theory is used to address the evolution of HIV with 1-10 mutations per division and “it seems to operate very close to its error threshold”[12]. The existence of this “threshold” is our main postulate here. However we apply it to all species, and with the provided IT framework, such threshold seems to be well-defined and ready for modeling. The main difference between virus and mammals populations seems to be the generation time – the virus genotype “cloud” can be readily observed empirically. The equilibrium regime of maintaining variability is considered in quasispecies theories too, and after we introduce the pattern definition and the measure of genetic information with fitness function we arrive to our model. However, in the quasispecies theory the fitness is defined for the whole population of mutants, not for individuals [12].

Then (for $GI < 2$ bit) a large number of allowable sequences (constituting a “typical set”) are nearly “synonymous” and can coexist in a population in the case of equilibrium maintenance evolution. However, they are not completely synonymous so that the selection can maintain a pattern by discarding the most deviant (“atypical”) sequences. The advantage of the model is that it allows quantifying meaningfully the information contained in a pattern (or a genome) and given the defined weights matrices of a desired conservation profile, the model provides selective values of individuals accounting for all mutations, present and denovo. We showed [4] that the substitution rate in functional sequences can be arbitrary close to the neutral rate and the fraction of positive mutation can be high in general. About 50% of the retained mutations must be “positive” - a trivial requirement for the balance of GI . Our model, analogously to studying of evolution with Turing machines [13], can be described as a population of machines operating on symbol sequences (of limited length), reading out positional information recognizing corresponding patterns (via typical sets, technically, for a general typical set the assumption of positional independence is not necessary) of molecular interactions and calculating high-level phenotype. However, it seems that our machine is closer to describing the molecular computations through patterns recognition in comparison with the sequential algorithmic Turing machine. For the purposes of this investigation we don’t have to specify the phenotype calculations per se – once we define the patterns and typical sets in a “genome” we can address the problem of their maintenance or evolution (e.g. the cost or speed of patterns preservation or change). Here we focus on the maintenance properties, treating such machines as genetic information storage devices, which must resist random noise of mutagenesis. The only computation is done for selection actions – a genome “typicality” is used as fitness, accounting for all variants in a genome (Equation 3). As could be expected our fitness function is similar to the traditional one in its basic “common sense” features – for example a mutation in highly conserved site (high GI) will drop the fitness significantly. Notably, in this model all sites and variants are functional - there is no need to postulate “neutral” [14] or “near-neutral” [15] variants (to explain the high rates of sequence evolution) – in our case, the equilibrium can be interpreted as the cumulative neutrality of all mutations (remained in a population), while assuming the individual neutrality of all or most mutations would be throwing the baby out with the bathwater.

Simulation terms

An organism is represented by the nucleotide sequence of given length (L), $O = [B_1, B_2, \dots, B_L]$, where $\forall i \in [1, L], B_i \in \{A, G, C, T\}$. A population is a set of organisms of the same length. Parameters, which govern the process of simulation, are shown in Table 1. The mutational bias is included in the code for universality, but has no effect on the trends we investigate here.

Then, each organism (O) in a population can be associated with a weight specified by the weight matrix W :

$$W(O) = W([B_1, B_2, \dots, B_L]) = \sum_{i=1}^L W_i(B_i) \quad (3)$$

A “typical” probability is the expected probability of a sequence for a given GI -profile. It can be calculated through multiplication of the expected frequencies for corresponding positions. Here, for computational convenience, we define the fitness through summation of position-specific weights, which, for our purposes, is equivalent to the multiplication if we had used logarithms of frequencies. However, we do not know the resulting GI -profile before the

simulation is performed, hence the weight matrix defines a general direction of pattern conservation by selection, not the actual *GI*-profile per se.

Table 1. Simulation parameters.

Notation	Description
N	Number of organisms in the population (population size).
L	Length (number of bases) of genome of each organism in the population.
nd	Number of descendants each organism produces in a single round of reproduction.
P_m	Probability of mutation per base.
P_{ti}	Probability that an occurred mutation will be a transition mutation.
$W = (W_j \mid j \in [1,L])$	Selection weights of nucleotides in each position. Where $W_j = (w_{jA}, w_{jC}, w_{jG}, w_{jT})$, $W_j(B) = w_{jB}$, $B \in \{A,G,C,T\}$ – selection weight of the corresponding nucleotide B in j -th position.

This weight is used to determine preferences of selection, which tries to maintain a pattern. In our experience, the particular recipes for selection actions (e.g. probabilistic/deterministic) and reproduction modes (overlapping/non-overlapping generations) play little role for the described trends, as long as the main purpose of these actions is to maintain a pattern – a biased frequencies distribution, while the opposing force – random mutagenesis tries to flatten the bias. As we mentioned *GI* can be viewed as a convenient measure of functionally acceptable (we have to stress - not “observable”, in general) variants frequencies biases; such fitness definition, in our opinion, is the key departure from traditional models. For example it seems to be inherently difficult to approach Drake's rule explanation with a fitness function which is relative – it has no information on organisms' degree of complexity, hence, taken alone, it is “blind” to a genome size. In our case the total *GI* - organismal complexity is measured by the amount of pattern's (functionally acceptable) biases. It seems to be intuitively appealing quantification - the larger the total amount of biases (further from the equilibrium flat distribution) - the higher the information content. However such approach is a necessary simplification - it works under the assumption that the rest (“higher order”) information unfolding processes are approximately the same, which should work, at least for the similar species.

Presumably, the sophisticated error corrections mechanisms such as DNA repair constitute a biological burden. So we ask what is the maximum mutation rate, which is compatible with a given total *GI*. The differences of GI_p of functional sequences are assumed to be small for close species. Formally, for our phenotype-calculating machines, the conservation of *GI* is equivalent to the whole phenotype conservation, because as we reasoned in [4] the *GI* conservation preserves positional information of molecular interactions, so that a phenotype is mechanistically derived from the whole genome pattern.

Simulation Process

The entire simulation process can be divided in three successive stages: initialization, spawning and selection. The initialization occurs only in the very beginning and then the spawning and selection are repeated in a loop until the simulation process is stopped.

Initialization: the initial population, consisting of N organisms of length L is generated. All organisms in the initial population are identical and have maximum possible weight

according to matrix W , i.e. at each position j of each organism stands nucleotide B_j :

$$B_j = \left[B \mid w_{jB} = \max_B (w_{jB}, B \in \{A, G, C, T\}) \right].$$

Spawning: the progeny is spawned. Each organism in the population produces n_d descendants (here we consider in detail only the case of binary fission, i.e. when $n_d = 2$). A descendant organism has the same length as its parent and is obtained by the copying of the parental sequence with a certain probability of mutation (P_m) and with a bias of mutational spectrum (P_{ti}). The parental organism is excluded from the population after the reproduction, so the generations are non-overlapping and after this step the population consists of $n_d N$ organisms.

Selection: the selection reduces the number of organisms in the population back to the initial size. It acts deterministically, leaving N organisms, whose weight $W(O)$ is larger.

The choice of procedure of the initial population generation does not affect the steady state of the simulation process, so we can simply generate a random initial population. However, generating the initial population as described above will provide the faster convergence to the steady state – the equilibrium condition, which reveals the “error threshold” – the goal of our experiments. The above mode of reproduction describes the non-overlapping generations, for the simplicity of defining and counting mutations; however, we experimented with other regimes, including the overlapping generations similarly to Moran model, and found the trends invariant.

Results

GI Behavior in the Course of Simulation

At first GI_ρ of the initial population is equal to 2 bits, because all organisms are identical, but as we discussed earlier that is not the “correct” functional GI , but a formally computed value in the course of simulation. If we start the simulation process as described above with the probability of mutation P_m high enough to allow occurring mutations to propagate in the population, then the diversity will emerge and GI_ρ will start to decrease. While reducing, GI_ρ will finally reach the level, when the mutagenesis is balanced by the force of selection, and in the consequent iterations it will fluctuate in a vicinity of some value. This equilibrium is analogous to the equilibrium introduced in quasispecies theories, but again - it is not observed in a realistic population of higher organisms. The existence of the balance (mean GI_ρ) is clear because the capacity (the averaged effect) of random mutagenesis to decrease GI monotonically drops from some value at $GI_\rho = 2$, to zero at $GI_\rho = 0$, while the corresponding selection capacity to increase GI behaves reciprocally – having non-zero value at $GI_\rho = 0$ and zero at $GI_\rho = 2$, thus these two functions intersect at some equilibrium point. In our numerical experiments we consider that the population is already in the equilibrium state if during the last T generations ($T = 100$ in our tests) two conditions are met: the sum of all GI_ρ changes between consequent generations is less than a specified threshold ($1e-3$ in our tests), the maximum number of consequent generations increasing/decreasing GI_ρ is less than $0.1 * T$. The fluctuations around the equilibrium depend on particular features of modeling, however, in our experience, with higher population sizes the fluctuations are smaller, but the mean value of GI_ρ does not vary significantly - for the equilibrium there is no dependence on the population size, which is natural to expect for the population maintaining constant variant frequencies. We will denote this mean value of GI_ρ in equilibrium population as GI_{steady} . A biological interpretation is that it is a given species maintainable GI value. It should be, however, clearly understood, that the word “steady” here concerns only the genetic

information (and hence the phenotype), the genomes remain variable, because new mutations still appear with the steady rate. The “molecular clock” is ticking - and its empirical steadiness on the evolutionary scale is another indirect hint that the average GI density is a slowly varying parameter. For example, mutations are more frequent in a position with the lower GI value, so if the density would fluctuate strongly on the evolutionary scale, the clock would behave erratically. As we argued [4], GI increasing (positive) mutations constitute a significant fraction of random mutations (especially when GI in a position is low), thus allowing the same fraction (in the GI equivalent) of negative mutations to remain in the population. The monotonous molecular clock is a simple prediction of the provided model, alternatively it can be explained by the neutrality assumption, which seems to be an oversimplification of reality. Also the provided model shows that the steadiness of the clock is intimately connected with Drake's rule and the “error threshold”, while the neutral theory is inherently unable to make such connections. The convergence of GI_ρ for different parameters is presented in Figure 1.




Figure 1. Convergence of GI_ρ for different parameters.

Common parameters for all demonstrated cases are: $N = 1000$; $n_d = 2$; $P_{ii} = 2/3$; $W = (W_j = (0.8, 0.2, 0, 0)$ if j is even, else $W_j = (0.5, 0.3, 0.1, 0.1)$). Color determines organism length (L): green corresponds to $L = 100$, blue to $L = 200$ and red to $L = 400$. Line style determines probability of mutation per base (P_m): solid – $P_m = 0.01$, dashed – $P_m = 0.04$.

Counting Mutations

In simulation the number of fixated mutations, i.e. the observed mutations per generation can be counted directly. Following the common notation we denote the number of mutations per generation per base as u_b and the mutation rate per generation per genome as u_g . Despite the fact that the values u_b and P_m are closely related, u_b is always less or equal than P_m , since the organisms with more mutations are more likely to be eliminated at the selection stage.

Now let's look at somewhat inverse experiment: we can fix the value of GI_{steady} and all parameters from Table 1 except P_m , and then numerically find the value of P_m which corresponds to the fixed parameters. This procedure can be performed for the different lengths of organism (L) while maintaining the same values of all other parameters (N , n_d , P_{ii} , W , GI_{steady}). The fixed parameters in our experiment were set to: $N = 1000$; $n_d = 2$; $P_{ii} = 2/3$; $W = (W_j = (0.8, 0.2, 0, 0)$ if j is even, else $W_j = (0.5, 0.3, 0.1, 0.1)$); $GI_{steady} = 1.6$. The experiment was

performed for L values 64, 128, 256, 512 and 1024. Then we estimated the number of mutations observed in the GI -steady state and compared u_b and u_g parameters for different genome lengths. The results are summarized in Figure 2.




Figure 2. Relationship between the mutation rate per site per generation (u_b) and the genome size (L) observed in the simulation.

Red points – $GI_{steady} = 1.6$ bit per site, blue line – the corresponding regression line, based on the equation: $\log_2 u_b = -2.337 - 0.929 \log_2 L$ ($r^2 = 0.9993$).

Yellow points – $GI_{steady} = 1.4$ bit per site, green line – the corresponding regression, based on the equation: $\log_2 u_b = -1.251 - 0.982 \log_2 L$ ($r^2 = 0.9998$).

We also found the dependence of GI_{total} vs. GI_ρ when the mutations rate (P_m) is fixed while the genome size (L) varies (Figure 3). We considered the populations with genome sizes from 100 to 1100 with the step of 20. All other parameters of these populations were identical, namely: $N = 1000$; $n_d = 2$; $P_{ti} = 2/3$; $W = (W_j = (0.8, 0.2, 0, 0)$ if j is even, else $W_j = (0.5, 0.3, 0.1, 0.1)$) and mutation rate P_m was set to 0.007. For each population its GI density (GI_ρ) and the total GI (GI_{total}) were averaged over 1000 generations after the population reached GI -steady state.




Figure 3. Dependence of total GI (GI_{total}) on equilibrium GI density (GI_ρ) for a fixed mutation rate and different genome sizes (L).

Each point represents a population with organisms having genome of size $L \in [100, 120, \dots, 1080, 1100]$. For convenience of orientation some points are colored in red and the genome size of corresponding population is labeled.

Through defining different weights matrices (Equation 3) we tested different scenarios of the density distribution: with homogeneous *GI* distribution in a genome, and bimodal – one half of a genome consisting of highly conserved (“lethal”) sites – to model the regions such as conserved protein domains and the other half consisting of weakly conserved sites, to model the variable parts of proteins and weakly conserved non-coding regulatory DNA. The meaning of the obtained results for the average *GI* density versus mutation rates is the same, so that the actual distribution is not affecting the described trends.

Discussion and Conclusion

Naturally the model's applicability to the evolution of real molecular machines should be thoroughly investigated since any formal model has its limits. For example particular alleles might interact in more complex ways than described by the model. However, since it provides simple explanations for observable phenomena, with the minimal number of parameters and assumptions, and, in principle, is realizable in a hardware, which operates similarly to our understanding of molecular interactions, we believe that it fairly captures the general properties of real genetic systems. Interestingly, the model can be considered as a simple generalization of Hardy–Weinberg equilibrium (HWE) [16], explicitly including functional sites and their maintenance selection. This may explain the persistent (about half-century) illusion of neutrality - in usual tests (e.g. Tajima's), the mutations in an equilibrium population will pretend to be neutral, so such tests are actually testing for the equilibrium condition rather than for the individual mutations neutrality. Below we discuss some possible consequences for our understanding of real genetic systems, assuming that the model is sufficiently valid. To prevent unnecessary criticisms, we have to admit that we are discussing some possible features of an idealized population described by the model; the correspondence to real genetic systems reflects the practical value of the model (such assessment is often influenced by subjective personal “tastes”), which is of secondary importance. For example the “heated” debates of neutralists vs. selectionists - it seems that our view reconciles both camps - the evolution is mostly neutral (in stasis), though this neutrality is maintained by the selection of the most typical individuals. Technically, it is a normal epistemological practice to explore abstract models regardless of their immediate relevance to “reality”.

According to Drake [6] the genomic mutation rate “is likely to be determined by deep general forces, perhaps by a balance between the usually deleterious effects of mutation and the physiological costs of further reducing mutation rates”. As can be seen, Drake correctly did not include considerations for adaptive properties of evolution, practically solving the problem, hinting that it is rather the maintenance-related phenomenon, and once we interpret the maintenance as the equilibrium in alleles frequencies – the main property of our model – the population size is obviously out of the equation (as in the case of HWE).

The key assumption for Drake's rule explanation is that the total genomic information is saturated to its maximum maintainable value, or reciprocally and equivalently, the mutation rate is near its upper limit for a given species total *GI*. The mutation rates and thus the total *GI* are assumed to change slowly on evolutionary scale. We hypothesize that the rate decrease is a basic event required for progressive evolution, and it is promptly followed by the gain in total *GI*, restoring the equilibrium. The equilibrium can be regained “quickly” (~100 generations) judging by the speed of convergence to the steady state (Fig. 1). One difficult question is how to motivate the stability of the mutation rate for a given species – for the rate decrease, we can assume that it might happen due to the large difference of the

timescale of two phenomena. One is a merely long-term advantage of the lowered mutation rate – some generations must pass to fill newly accessible GI (if a niche requires it, which does not have to be the case, in general). And the other is the immediate disadvantage – “physiological costs” – since the lower rate, in principle, must be associated with a slower replication rate and/or additional energy expenditures. On the other hand, why the rates are not degrading, if increasing the rate might bring a fast advantage, but only a long-term disadvantage? At this point we can only speculate that for higher organisms, an increased somatic mutagenesis might cause also the short-term disadvantage, preventing the rate degradation (e.g. somatic mutations theories of aging or carcinogenesis). Beside the somatic mutagenesis, we could imagine any other selectively important phenotype, which is somehow linked to the changes in mutation rate. The other idea is that while the rate decrease must come at some “physiological costs”, the way back is not that easy – a mutation, which degrades the rate will not necessarily reduce the “physiological costs” back to the previous values. Such mutation must be rather specific “back-mutation” or, more likely, a number of them, making it improbable to achieve both – the rate increase and the corresponding costs reduction. Hence the rates can only go down, locked from above by both – short- and long-term disadvantages. Alternatively, the rate maintenance might require a regular population renewal, described below. Naturally there are examples of regressive evolution, which are not “interesting” for us here – such evolution can be easily caused, for example, by moving to a simpler niche (habitat) – “use it or lose it”.

Hypothetically a change of mutation rate would remold a species phenotype (suggesting an explanation for “punctuated equilibrium” phenomenon), since small relative change of the mutation rate can provide significant absolute change in total accessible GI and correspondingly a significant change in the phenotype. In principle it is assessable experimentally – if we were able to select flies (for example) for the lowered mutation rate, then the model predicts that such population has the capacity to produce more advanced species of flies. The problem is, however, that the population must be challenged with the proper external conditions, which could cause the evolutionary progress, promoting an increase of complexity.

It is natural to expect that the rates occupy discreet values, due to the discreet nature of corresponding modifying mutations and their (presumably) limited number. We could also hypothesize about speciation scenarios: suppose that in a large population the rates are heterogeneous and mixed, so that the population has some averaged rate. Then after a “founder” splits off, he produces a new population, which can have the rate different from the main population, leading to the fast phenotype changes.

The evolution of the (functional) genome size is presumed to occur through gene duplications, so that “gene families” grow in size. That also motivates our postulate of slow changes of GI_p for functional sequences – new sequences perform molecular functions similar to the original. The provided theory readily predicts that whole or partial genome duplications would lead to an increased rate of sequence evolution and to a subsequent shrinking back of the (functional) genome size, losing extra gene copies, due to inability to maintain higher total GI without changes of the mutation rates. The evolutionary progress (an increase in complexity and GI) is happening not due to duplications per se (which are relatively frequent events), but due to the mutation rate decrease and/or the adoption of the lower GI_p functionality (Fig. 3) – these changes are assumed to be “slow”. However the progressive evolution is naturally intervened with the external conditions – niche or habitat – it must be sufficiently complex to support the increase of species complexity. Duplications might also cause a reproductive isolation; hence together with the founder-specific mutation

rate hypothesis this might be a path to speciation and progressive evolution (when the founder retains the lowered mutation rate).

Hence we suggest that the “channel capacity” notion of IT is sufficiently deep and general principle to provide the desired understanding of Drake's rule; the notion also allows quantitative modeling of the process. Channel capacity is the upper bound for the information transmission rate for a given noise level. Practical solutions for information transmission are somewhat below this theoretical limit, and considerable engineering efforts are dedicated to approach the limit, simply because being closer to the limit saves energy. Hence another basic consideration is that if the nature would not use the genomic informational capacity to its full extent, it would not be “thrifty” - why waste resources on the unused capacity - the thriftiness should be favored by selection (though there are some opposing ideas of “selfish” or “parasitic” sequences). Intriguingly, in IT the problem of approaching the channel capacity limit has no general solution applicable to all practical situations, as it relates to the problem of achieving best compression rate, and, in practice, is limited by the memory and computational costs. Analogously to Chaitin proposals [17], we can speculate that molecular machines have an infinite field for exercising the mathematical creativity in attempts to approach the limit, explaining the drive to complexity in living systems [4]. Naturally the simple model captures only crude properties of the genetic information processing as there are many features not included – epigenetics, rearrangements, roles of transposable and repetitive elements, recombination, multiple ploidy, etc.

In comparison with the other recently proposed explanation of Drake's rule [7], our model does not call for additional difficult-to-define entities like “molecular refinements”, “drift barrier” or “effective population size” - the estimates of the later are admitted by the authors to be “fraught with difficulties”. It is not clear how to simulate that evolutionary model *in-silico*, to perform its validation, because genome-wide functionality and conservation is not defined. Hence there is no specific model for selection actions, and there are many arbitrary parameters. However a desirable feature of a “mechanistic” evolutionary model is the ability to simulate it, to evaluate its robustness in parameters space. Comparing Fig. 1A of [7] with Fig. 2, we can hypothesize that eukaryotes have lower *GI* density, in average, which is consistent with other observations – weaker genomic conservation, and Fig. 3 demonstrates that it can be advantageous to utilize the lower density *GI* storage. The *GI* storage strategy can be affected by particular demands for optimization: e.g. viruses or bacteria might prefer compact genomes with high *GI*, for faster replication or smaller physical size, utilizing the double stranded and overlapping coding and avoiding weakly conserved regulatory non-coding DNA.

The important consequence of our reasoning is that molecular evolution in average is not about a continuous increase of total *GI*. This suggest an explanation to a naive, but still valid question of why we see “living fossils” or don't see contemporary monkeys evolving into humans continuously, (anthropocentrically) assuming the later have higher *GI*; while on the other hand we can observe an amazing phenotypic plasticity (e.g. dogs pedigrees or Cetacean evolution). Despite being “adaptive”, for a given change of environment, the evolution is not “progressive” in terms of total *GI*, since we posit that each species already have the maximum *GI*, allowed by the mutation rate, which is assumed to vary slowly. That also calls to revisit the popular evolutionary concept that genes are near their best functional performance – the performance is as good as allowed by the corresponding channel capacity – balancing at the brink of “chaos and order”, so that a random mutation has high chances of being positive, in general. The dependence of the “evolvability” on the population size is

also practically a “dogma” in traditional theories, which, might be a consequence of the unconstrained (“open-ended”) opportunistic “Brownian” views on evolution. However if a population is at the *GI* limit, so that an advance in one function must be associated with the “costs” to others, the role of the population size might be diminished, at least, as we showed, for the maintenance regime. It seems that strong dependencies on population size in traditional models lead to some contradictions with observations, such as Lewontin’s “Paradox of Variation” [18], not to mention the general trend that more evolved forms have smaller population sizes, in average. Ironically, we can draw the opposite scenario for evolvability vs. population size: without the immediate negative effects, random mutations will degrade the rate in average. Hence the large population size in a long run can lead to the accumulation of variants, which increase the average mutation rate, leading to a degradation; then the way out is through the bottlenecks: the population must be regularly refreshed by the founding of subpopulations with decreased (below the average) rates. Such subpopulations will quickly gain an advantage and overcome the main population. In a sense it is the population genetic “ageing”, analogously to a somatic ageing. In that case, the reproductive barriers, bottlenecks and speciation events are the necessities of evolution, required for the renewal and progress, rather than peculiar accidental features. We would like to remind that in this model positive mutations are abundant, so there is no need in a large population size or waiting time.

The adaptation to new selection demands then happens at the price of decrease of adaptation to other demands, the phenomenon well known to breeders (which now may attempt to select for the lower mutation rates also). For our model this can be imagined as a reshaping of genomic *GI* profile (and correspondingly a phenotype) while keeping the total *GI* constant. In biological interpretation it is the directional decrease of variability (reflected in the increase of corresponding *GI*) of one phenotypic feature (which is in demand), while increasing variability (“loosening up”) of others. The traditional relative fitness function alone is unable to distinguish between such “reshaping” and “progressive” evolution modes, because the channel capacity notion is absent in traditional models (except for the somewhat analogous “error threshold” considerations, which are presumed to be narrowly applicable in some special cases). The general properties of such “reshaping selection” can be easily modeled with the suggested IT framework, to evaluate its basic features and the influences of diverse evolutionary strategies. In the case of eukaryotes we can expect that such evolutionary plasticity is residing mostly in non-coding regions with low *GI* density. Another straightforward application is the modeling of multifactorial and polygenic disorders, where a number of paradoxes are still waiting their resolution, such as the high heritability on the one hand, and the low predictive power derived from GWA studies, on the other. Probably some paradoxes can be approached by the abandoning the assumption of neutrality of common variants.

References

- [1] Moran PAP (1962) The statistical processes of evolutionary theory. Clarendon Press, Oxford.
- [2] Durrett R (2008) Probability Models for DNA Sequence Evolution. Springer, New York.
- [3] Ohta T, Gillespie JH (1996) Development of Neutral and Nearly Neutral Theories. *Theor Popul Biol* 49(2):128-42.

- [4] Shadrin AA, Grigoriev A, Parkhomchuk DV (2013) Positional information storage in sequence patterns. *Comput Mol Biosci.* 3(2):18-26.
- [5] Drake JW (1991) A constant rate of spontaneous mutation in DNA based microbes. *Proc Natl Acad Sci USA* 88: 7160–7164.
- [6] Drake JW, Charlesworth B, Charlesworth D, Crow JF (1998) Rates of Spontaneous Mutation. *Genetics* 148: 1667–1686.
- [7] Sung W, Ackerman MS, Miller SF, Doak TG, Lynch M (2012) Drift-barrier hypothesis and mutation-rate evolution. *Proc Natl Acad Sci USA* 109(45):18488-92.
- [8] Schneider TD, Stormo GD, Gold L, Ehrenfeucht A (1986) Information content of binding sites on nucleotide sequences. *J Mol Biol* 188:415-431.
- [9] Schneider TD, Stephens RM (1990) Sequence Logos: a New Way to Display Consensus Sequences. *Nucleic Acids Res* 18:6097-6100.
- [10] Hertz GZ, Stormo GD (1999) Identifying DNA and protein patterns with statistically significant alignments of multiple sequences. *Bioinformatics* 15(7-8):563-77.
- [11] Shannon CE (1948) A Mathematical Theory of Communication. *Bell System Technical Journal* 27:379-423, 623-656.
- [12] Nowak MA (1992) What is a quasispecies? *Trends Ecol Evol* 7: 118-121.
- [13] Feverati G, Musso F (2008) Evolutionary model with Turing machines. *Phys Rev E Stat Nonlin Soft Matter Phys* 77(6 Pt 1):061901.
- [14] Kimura M (1983) *The neutral theory of molecular evolution*. New York: Cambridge University Press.
- [15] Ohta T (1973) Slightly deleterious mutant substitutions in evolution. *Nature* 246(5428):96-8.
- [16] Hardy GH (2003) Mendelian proportions in a mixed population. 1908. *Yale J Biol Med* 76: 79-80.
- [17] Chaitin G (2012) *Proving Darwin: Making Biology Mathematical*. New York: Pantheon Books.
- [18] Lewontin RC (1974) *The genetic basis of evolutionary change*. Columbia University Press.