

# Fuzzy games with a countable space of actions and applications to systems of generalized quasi-variational inequalities

Monica Patriche

*University of Bucharest Faculty of Mathematics and Computer Science  
14 Academiei Street  
010014 Bucharest, Romania  
monica.patriche@yahoo.com*

---

## Abstract

In this paper, we introduce an abstract fuzzy economy (generalized fuzzy game) model with a countable space of actions and we study the existence of the fuzzy equilibrium. As applications, two types of results are obtained. The first ones concern the existence of the solutions for systems of generalized quasi-variational inequalities with random fuzzy mappings which we define. The last ones are new random fixed point theorems for correspondences with values in complete countable metric spaces.

*Keywords:* Abstract fuzzy economy, fuzzy equilibrium, incomplete information, random fixed point, random quasi-variational inequalities, random fuzzy mapping.

---

## 1. INTRODUCTION

In equilibrium theory, there are two directions for the study of the models with differential information. The first one is due to Harsanyi [9], whose first approach refers to Nash type models, and the second one is due to Radner [23], who introduced differential information into the Arrow-Debreu model [2]. Since then, an entire literature has developed on this topic.

A basic problem concerning the noncooperative games with differential information was the equilibrium existence. The classical deterministic models

of abstract economies due to Borglin and Keiding [3], Shafer and Sonnenschein [24], or Yannelis and Prabhakar [28] have had several generalizations. We can refer the reader to Yannelis's results, who introduced new information concepts based on measurability requirements or Patriche's models [17], [19], [21], [22].

A new approach of an abstract economy with private information and a countable set of actions [22] follows the ideas of Yu and Zhang in [29], who worked with the distributions of the atomless correspondences.

This paper has two aims. Firstly, we extend the study of the model defined in [22] in the fuzzy framework, by taking into account those situations in which agents may have partial control over the actions they choose. Since we are interested in a fuzzy setting, special attention is paid to the fuzzy equilibrium existence of the fuzzy games emphasizing that this approach can be used to treat more complicated economic systems. The uncertainties derived by the individual character of agents in election situations can be interpreted using fuzzy random mappings, which can be used for the purpose of modelling and analyzing the economic systems. We continue the way opened by Kim and Lee [11], who proved that the theory of fuzzy sets, initiated by Zadeh [31], has become a good framework for obtaining results concerning fuzzy equilibrium existence for abstract fuzzy economies. For other results in this domain, the reader is referred to [4], [10], [12], [13], [25], [27].

Finally, we search for applications of our results concerning the existence of the fuzzy equilibrium for the fuzzy games with countable action sets, in connection with the systems of generalized quasi-equilibrium inequalities with random fuzzy mappings. The variational inequalities were introduced in 1960s by Fichera and Stampacchia, who studied equilibrium problems arising from mechanics. Since then, this domain has been extensively developed and has been found very useful in many diverse fields of pure and applied sciences, such as mechanics, physics, optimization and control theory, operations research and several branches of engineering sciences. Recently, the random variational inequalities have been introduced and studied in [7], [8], [15], [26], [30]. We prove the existence of the solutions for systems of random generalized quasi-variational inequalities with random fuzzy mappings which we define. As consequences, we obtain new random fixed point theorems for correspondences with values in complete countable metric spaces.

The paper is organized as follows. In the next section, some notational and terminological conventions are given. In Section 3, the model of an abstract fuzzy economy with private information and a countable space of

actions is introduced and the main result is stated. Section 4 contains existence theorems for solutions of random quasi-variational inequalities with random fuzzy mappings and random fixed point theorems. In the last section, classical results are obtained as consequences.

## 2. NOTATION AND DEFINITION

Throughout this paper, we shall use the following notation:

$\mathbb{R}_{++}$  denotes the set of strictly positive reals.  $\text{co}D$  denotes the convex hull of the set  $D$ .  $\overline{\text{co}}D$  denotes the closed convex hull of the set  $D$ .  $2^D$  denotes the set of all non-empty subsets of the set  $D$ . If  $D \subset Y$ , where  $Y$  is a topological space,  $\text{cl}D$  denotes the closure of  $D$ .

For the reader's convenience, we review a few basic definitions and results concerning continuity and measurability of correspondences.

Let  $X, Y$  be topological spaces and  $F : X \rightarrow 2^Y$  be a correspondence.  $F$  is said to be *upper semicontinuous* if for each  $x \in X$  and each open set  $V$  in  $Y$  with  $F(x) \subset V$ , there exists an open neighbourhood  $U$  of  $x$  in  $X$  such that  $F(y) \subset V$  for each  $y \in U$ .  $F$  is said to be *lower semicontinuous (l.s.c)* if for each  $x \in X$  and each open set  $V$  in  $Y$  with  $F(x) \cap V \neq \emptyset$ , there exists an open neighbourhood  $U$  of  $x$  in  $X$  such that  $F(y) \cap V \neq \emptyset$  for each  $y \in U$ .

Each correspondence  $F : X \rightarrow 2^Y$  has two natural inverses. The upper inverse  $F^u$  (also called the *strong inverse*) of a subset  $A$  of  $Y$  is defined by  $F^u(A) = \{x \in X : F(x) \subset A\}$ . The lower inverse  $F^l$  (also called the *weak inverse*) of a subset  $A$  of  $Y$  is defined by  $F^l(A) = \{x \in X : F(x) \cap A \neq \emptyset\}$ .

An important result concerning upper (lower) semicontinuous correspondences is the following one.

**Lemma 1.** [30] *Let  $X$  and  $Y$  be two topological spaces and let  $A$  be a closed (resp. open) subset of  $X$ . Suppose  $F_1 : X \rightarrow 2^Y$ ,  $F_2 : X \rightarrow 2^Y$  are lower semicontinuous (resp. upper semicontinuous) such that  $F_2(x) \subset F_1(x)$  for all  $x \in A$ . Then the correspondence  $F : X \rightarrow 2^Y$  defined by*

$$F(x) = \begin{cases} F_1(x), & \text{if } x \notin A, \\ F_2(x), & \text{if } x \in A \end{cases}$$

*is also lower semicontinuous (resp. upper semicontinuous).*

Let  $(T, \mathcal{T})$  be a measurable space,  $Y$  a topological space and  $F : T \rightarrow 2^Y$  a correspondence.  $F$  is *weakly measurable* if  $F^l(A) \in \mathcal{T}$  for each open subset

$A$  of  $Y$ .  $F$  is *measurable* if  $F^l(A) \in \mathcal{T}$  for each closed subset  $A$  of  $Y$ . If  $(T, \mathcal{T})$  is a measurable space,  $Y$  a countable set and  $F : T \rightarrow 2^Y$  is a correspondence, then  $F$  is measurable if for each  $y \in Y$ ,  $F^{-1}(y) = \{t \in T : y \in F(t)\}$  is  $\mathcal{T}$ -measurable.

**Lemma 2.** [1]. *For a correspondence  $F : T \rightarrow 2^Y$  from a measurable space into a metrizable space we have the following:*

1. If  $F$  is measurable, then it is also weakly measurable;
2. If  $F$  is compact valued and weakly measurable, it is measurable.

The following properties are essential tools used to prove the existence of equilibria for abstract economies in Section 3. We follow Yu and Zhang [29]. Let  $Y$  be a countable complete metric space,  $(T, \mathcal{T}, \lambda)$  an atomless probability space and  $F : T \rightarrow 2^Y$  a measurable correspondence. The function  $f : T \rightarrow Y$  is said to be a *selection of  $F$*  if  $f(t) \in F(t)$  for  $\lambda$ -almost  $t \in T$ . Let us denote  $\mathcal{D}_F = \{\lambda f^{-1} : f \text{ is a measurable selection of } F\}$ .

We will present some regular properties of  $\mathcal{D}_F$ , also obtained by Yu and Zhang [29].

The next lemma states the convexity of  $\mathcal{D}_F$  for any correspondence  $F$ .

**Lemma 3.** [29]. *Let  $Y$  be a countable complete metric space,  $(T, \mathcal{T}, \lambda)$  an atomless probability space and  $F : T \rightarrow 2^Y$  a measurable correspondence. Then,  $\mathcal{D}_F$  is non-empty and convex in the space  $\mathcal{M}(Y)$  - the space of probability measure on  $Y$ , equipped with the topology of weak convergence.*

The compactness of  $\mathcal{D}_F$  is stated in the next result.

**Lemma 4.** [29]. *Let  $Y$  be a countable complete metric space, and  $(T, \mathcal{T}, \lambda)$  be an atomless probability space and  $F : T \rightarrow 2^Y$  be a measurable correspondence. If  $F$  is compact valued, then,  $\mathcal{D}_F$  is compact in  $\mathcal{M}(Y)$ .*

Lemma 5 concerns the upper semicontinuity property of  $\mathcal{D}_{F_x}$ .

**Lemma 5.** [29]. *Let  $X$  be a metric space,  $(T, \mathcal{T}, \lambda)$  an atomless probability space,  $Y$  a countable complete metric space and  $F : T \times X \rightarrow 2^Y$  a correspondence. Let us assume that, for any fixed  $x$  in  $X$ ,  $F(\cdot, x)$  (also denoted by*

$F_x$ ) is a compact-valued measurable correspondence, and for each fixed  $t \in T$ ,  $F(t, \cdot)$  is upper semicontinuous on  $X$ . Also, let us assume that there exists a compact valued correspondence  $H : T \times X \rightarrow 2^Y$  such that  $F(t, x) \subset H(t)$  for all  $t$  and  $x$ . Then  $\mathcal{D}_{F_x}$  is upper semicontinuous on  $X$ .

We will also need Kuratowski-Ryll-Nardzewski Selection Theorem in order to prove our main results in the next section.

**Theorem 6.** (*Kuratowski-Ryll-Nardzewski Selection Theorem*) [1]. *A weakly measurable correspondence with non-empty closed values from a measurable space into a Polish space admits a measurable selector.*

The fuzzy mappings are presented below.

Let  $F(Y)$  be a collection of all fuzzy sets over  $Y$ . A mapping  $F : \Omega \rightarrow F(Y)$  is called a *fuzzy mapping*. If  $F$  is a fuzzy mapping from  $\Omega$ ,  $F(\omega)$  is a fuzzy set on  $Y$  and  $F(\omega)(y)$  is the membership function of  $y$  in  $F(\omega)$ .

Let  $A \in F(Y)$ ,  $a \in [0, 1]$ , then the set  $(A)_a = \{y \in Y : A(y) \geq a\}$  is called an  $a$ -cut set of the fuzzy set  $A$ .

A fuzzy mapping  $F : \Omega \rightarrow F(Y)$  is said to be *measurable* if, for any given  $a \in [0, 1]$ ,  $(F(\cdot))_a : \Omega \rightarrow 2^Y$  is a measurable correspondence. A fuzzy mapping  $F : \Omega \rightarrow F(Y)$  is said to *have a measurable graph* if, for any given  $a \in [0, 1]$ , the correspondence  $(F(\cdot))_a : \Omega \rightarrow 2^Y$  has a measurable graph. A fuzzy mapping  $F : \Omega \times X \rightarrow F(Y)$  is called a *random fuzzy mapping* if, for any given  $x \in X$ ,  $F(\cdot, x) : \Omega \rightarrow F(Y)$  is a measurable fuzzy mapping.

### 3. FUZZY EQUILIBRIUM EXISTENCE FOR ABSTRACT FUZZY ECONOMIES WITH PRIVATE INFORMATION AND A COUNTABLE SET OF ACTIONS

#### 3.1. THE MODEL OF AN ABSTRACT FUZZY ECONOMY WITH PRIVATE INFORMATION

In this section we define a model of abstract fuzzy economy with private information and a countable set of actions. This model is a generalization of the one defined in [22]

Let  $I$  be a non-empty finite set (the set of agents). For each  $i \in I$ , the space of actions,  $S_i$  is a countable complete metric space and  $(\Omega_i, \mathcal{Z}_i)$  is a measurable space. Let  $(\Omega, \mathcal{F})$  be the product measurable space  $(\prod_{i \in I} \Omega_i, \otimes_{i \in I} \mathcal{Z}_i)$ ,

and let  $\mu$  be a probability measure on  $(\Omega, \mathcal{F})$ . For a point  $\omega = (\omega_1, \dots, \omega_n) \in \Omega$ , define the coordinate projections  $\tau_i(\omega) = \omega_i$ . The random mapping  $\tau_i(\omega)$  is interpreted as player  $i$ 's private information related to his action.

We also denote, for each  $i \in I$ ,  $\text{Meas}(\Omega_i, S_i)$  the set of measurable fuzzy mappings  $f$  from  $(\Omega_i, \mathcal{Z}_i)$  to  $S_i$ . An element  $g_i$  of  $\text{Meas}(\Omega_i, S_i)$  is called a *pure strategy* for player  $i$ . A *pure strategy profile*  $g$  is an  $n$ -vector function  $(g_1, g_2, \dots, g_n)$  that specifies a pure strategy for each player.

We suppose that there exists a fuzzy mapping  $X_i : \Omega_i \rightarrow \mathcal{F}(S_i)$  such that each agent  $i$  can choose an action from  $(X_i(\omega_i))_{z_i} \subset S_i$  for each  $\omega_i \in \Omega_i$ .

Let  $\mathcal{D}_{(X_i(\cdot))_{z_i}}$  be the set  $\{(\mu\tau_i^{-1})g_i^{-1} : g_i \text{ is a measurable selection of } (X_i(\cdot))_{z_i}\}$  and  $\mathcal{D}_{(X(\cdot))_z} := \prod_{i \in I} \mathcal{D}_{(X_i(\cdot))_{z_i}}$ .

For each  $i \in I$ , let us denote  $h_{g_i} = (\mu\tau_i^{-1})g_i^{-1}$ , where  $g_i$  is a measurable selection of  $(X_i(\cdot))_{z_i}$  and  $h_g = (h_{g_1}, h_{g_2}, \dots, h_{g_n})$ .

**Definition 1.** *An abstract fuzzy economy (or a generalized fuzzy game) with private information and a countable space of actions is defined as*

$\Gamma = (I, ((\Omega_i, \mathcal{Z}_i), (S_i, A_i, P_i, a_i, b_i, z_i)))_{i \in I}, \mu)$ , where:

- (a)  $X_i : \Omega_i \rightarrow \mathcal{F}(S_i)$  is the *action (strategy) fuzzy mapping* of agent  $i$ ,
- (b) for each  $\omega_i \in \Omega_i$ ,  $A_i(\omega_i, \cdot) : \mathcal{D}_{(X(\cdot))_z} \rightarrow \mathcal{F}(S_i)$  is the *random fuzzy constraint mapping* of agent  $i$ ;
- (c) for each  $\omega_i \in \Omega_i$ ,  $P_i(\omega_i, \cdot) : \mathcal{D}_{(X(\cdot))_z} \rightarrow \mathcal{F}(S_i)$  is the *random fuzzy preference mapping* of agent  $i$ ;
- (d)  $a_i : \mathcal{D}_{(X(\cdot))_z} \rightarrow (0, 1]$  is a *random fuzzy constraint function* and  $p_i : \mathcal{D}_{(X(\cdot))_z} \rightarrow (0, 1]$  is a *random fuzzy preference function*;
- (e)  $z_i \in (0, 1]$  is such that for all  $(\omega_i, h_g) \in \Omega_i \times \mathcal{D}_{(X(\cdot))_z}$ ,  $(A_i(\omega_i, h_g))_{a_i(h_g)} \subset (X_i(\omega_i))_{z_i}$  and  $(P_i(\omega_i, h_g))_{p_i(h_g)} \subset (X_i(\omega_i))_{z_i}$ .

**Definition 2.** *A fuzzy equilibrium for  $\Gamma$  is defined as a strategy profile  $g^* = (g_1^*, g_2^*, \dots, g_n^*) \in \prod_{i \in I} \text{Meas}(\Omega_i, S_i)$  such that for each  $i \in I$ :*

- 1)  $g_i^*(\omega_i) \in (A_i(\omega_i, h_{g^*}))_{a_i(h_{g^*})}$  for each  $\omega_i \in \Omega_i$ ;
- 2)  $(A_i(\omega_i, h_{g^*}))_{a_i(h_{g^*})} \cap (P_i(\omega_i, h_{g^*}))_{p_i(h_{g^*})} = \phi$  for each  $\omega_i \in \Omega_i$ .

### 3.2. EXISTENCE OF THE FUZZY EQUILIBRIUM FOR ABSTRACT FUZZY ECONOMIES WITH A COUNTABLE SET OF ACTIONS

In this subsection we prove the existence of fuzzy equilibrium of abstract fuzzy economies.

Theorem 7 is our main result.

Let us denote  $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_n) \in \mathcal{D}_{(X(\cdot))_z}$ .

**Theorem 7.** *Let  $\Gamma = (I, ((\Omega_i, \mathcal{Z}_i), (S_i, A_i, P_i, a_i, b_i, z_i))_{i \in I}, \mu)$  be an abstract economy with private information and a countable space of actions, where  $I$  is a finite index set such that for each  $i \in I$  :*

a)  $S_i$  is a countable complete metric space and  $(\Omega_i, \mathcal{Z}_i)$  is a measurable space;  $(\Omega, \mathcal{Z})$  is the product measurable space  $(\prod_{i \in I} (\Omega_i, \mathcal{Z}_i))$  and  $\mu$  an atomless probability measure on  $(\Omega, \mathcal{Z})$ ;

b) the fuzzy mapping  $X_i : \Omega_i \rightarrow \mathcal{F}(S_i)$  is measurable and  $(X_i(\cdot))_{z_i} : \Omega_i \rightarrow 2^{S_i}$  has compact values;

c) the fuzzy mapping  $A_i$  is such that for each  $\lambda \in \mathcal{D}_{(X(\cdot))_z}$ ,  $(A_i(\cdot, \lambda))_{a_i(\lambda)} : \Omega_i \rightarrow 2^{S_i}$  is measurable and, for all  $\omega_i \in \Omega_i$ ,  $(A_i(\omega_i, \cdot))_{a_i(\cdot)} : \mathcal{D}_{(X(\cdot))_z} \rightarrow 2^{S_i}$  is upper semicontinuous with non-empty compact values;

d) the fuzzy mapping  $P_i$  is such that for each  $\lambda \in \mathcal{D}_{(X(\cdot))_z}$ ,  $(P_i(\cdot, \lambda))_{p_i(\lambda)} : \Omega_i \rightarrow 2^{S_i}$  is measurable and, for all  $\omega_i \in \Omega_i$ ,  $(P_i(\omega_i, \cdot))_{p_i(\cdot)} : \mathcal{D}_{(X(\cdot))_z} \rightarrow 2^{S_i}$  is upper semicontinuous with non-empty compact values;

e) for each  $\omega_i \in \Omega_i$  and each  $g \in \prod_{i \in I} \text{Meas}(\Omega_i, S_i)$ ,  $g_i(\omega_i) \notin (P_i(\omega_i, h_g))_{p_i(h_g)}$ ;

f) the set  $U_i^{\omega_i} := \{\lambda \in \mathcal{D}_{(X(\cdot))_z} : (A_i(\omega_i, \lambda))_{a_i(\lambda)} \cap P_i((\omega_i, \lambda))_{p_i(\lambda)} = \emptyset\}$  is open in  $\mathcal{D}_{(X(\cdot))_z}$  in for each  $\omega_i \in \Omega_i$ .

Then, there exists  $g^* \in \prod_{i \in I} \text{Meas}(\Omega_i, S_i)$  an equilibrium for  $\Gamma$ .

*Proof.* According to Lemma 3,  $D_{(X_i(\cdot))_{z_i}}$  is non-empty and convex. According to Lemma 4,  $D_{(X_i(\cdot))_{z_i}}$  is compact. For each  $i \in I$  let us denote

$$U_i := \{(\omega_i, \lambda) \in \Omega_i \times \mathcal{D}_{(X(\cdot))_z} : (A_i(\omega_i, \lambda))_{a_i(\lambda)} \cap P_i((\omega_i, \lambda))_{p_i(\lambda)} = \emptyset\} \text{ and}$$

$$U_i^{\omega_i} := \{(\omega_i, \lambda) \in \Omega_i \times \mathcal{D}_{(X(\cdot))_z} : (A_i(\omega_i, \lambda))_{a_i(\lambda)} \cap P_i((\omega_i, \lambda))_{p_i(\lambda)} = \emptyset\}.$$

According to the assumption f), the set  $U_i^{\omega_i}$  is open in  $\mathcal{D}_{(X(\cdot))_z}$ .

Let us define  $F_i : \Omega_i \times \mathcal{D}_{(X(\cdot))_z} \rightarrow 2^{S_i}$  by

$$F_i(\omega_i, \lambda) = \begin{cases} (A_i(\omega_i, \lambda))_{a_i(\lambda)} \cap (P_i(\omega_i, \lambda))_{p_i(\lambda)} & \text{if } (\omega_i, \lambda) \notin U_i, \\ (A_i(\omega_i, \lambda))_{a_i(\lambda)} & \text{if } (\omega_i, \lambda) \in U_i. \end{cases}$$

Then, the correspondence  $F_i$  has non-empty compact values and is measurable with respect to  $\Omega_i$  and upper semicontinuous with respect to  $\lambda \in \mathcal{D}_{(X(\cdot))_z}$ .

We denote  $\mathcal{D}_{F_i}(\lambda) =$

$= \{h_{g_i} = (\mu\tau_i^{-1})g_i^{-1} : g_i \text{ is a measurable selection of } F_i(\cdot, \lambda)\}$ . Then:

i)  $\mathcal{D}_{F_i}(\lambda)$  is non-empty because there exists a measurable selection from the correspondence  $F_i$  according to Kuratowski-Ryll-Nardewski Selection Theorem.

ii)  $\mathcal{D}_{F_i}(\lambda)$  is convex and compact according to Lemma 3 and Lemma 4.

We define  $\Phi : \mathcal{D}_{(X(\cdot))_z} \rightarrow 2^{\mathcal{D}_{(X(\cdot))_z}}$ ,  $\Phi(\lambda) = \prod_{i \in I} \mathcal{D}_{F_i}(\lambda)$ .

The set  $\mathcal{D}_{(X(\cdot))_z}$  is non-empty compact and convex. According to Lemma 5, the correspondence  $\mathcal{D}_{F_i}$  is upper semicontinuous. Then, the correspondence  $\Phi$  is upper semicontinuous and has non-empty compact and convex values. According to Ky Fan fixed point Theorem [6], there exists a fixed point  $\lambda^* \in \Phi(\lambda^*)$ . In particular, for each player  $i$ ,  $\lambda_i^* \in \mathcal{D}_{F_i}(\lambda^*)$ . Therefore, for each player  $i$ , there exists  $g_i^* \in \text{Meas}(\Omega_i, S_i)$  such that  $g_i^*$  is a selection of  $F_i(\cdot, \lambda^*)$  and  $h_{g_i^*} = (\mu\tau_i^{-1})(g_i^*)^{-1} = \lambda_i^*$ . Let us denote  $h_{g^*} = (h_{g_1^*}, \dots, h_{g_n^*})$ .

We prove that  $g^*$  is an equilibrium for  $\Gamma$ . For each  $i \in I$ , because  $g_i^*$  is a selection of  $F_i(\cdot, h_{g_1^*}, \dots, h_{g_n^*})$ , it follows that  $g_i^*(\omega_i) \in (A_i(\omega_i, h_{g^*}))_{a_i(h_{g^*})} \cap (P_i(\omega_i, h_{g^*}))_{p_i(h_{g^*})}$  if  $(\omega_i, h_{g^*}) \notin U_i$  or  $g_i^*(\omega_i) \in (A_i(\omega_i, h_{g^*}))_{a_i(h_{g^*})}$  if  $(\omega_i, h_{g^*}) \in U_i$ .

According to the assumption d), it follows that  $g_i^*(\omega_i) \notin (P_i(\omega_i, h_{g^*}))_{p_i(h_{g^*})}$  for each  $\omega_i \in \Omega_i$ . Then  $g_i^*(\omega_i) \in (A_i(\omega_i, h_{g^*}))_{a_i(h_{g^*})}$  and  $(\omega_i, h_{g^*}) \in U_i$ . This is equivalent with the fact that  $g_i^*(\omega_i) \in (A_i(\omega_i, h_{g^*}))_{a_i(h_{g^*})}$  and  $(A_i(\omega_i, h_{g^*}))_{a_i(h_{g^*})} \cap (P_i(\omega_i, h_{g^*}))_{p_i(h_{g^*})} = \emptyset$  for each  $\omega_i \in \Omega_i$ . Consequently,  $g^* = (g_1^*, g_2^*, \dots, g_n^*)$  is an equilibrium for  $\Gamma$ .

## 4. RANDOM QUASI-VARIATIONAL INEQUALITIES WITH RANDOM FUZZY MAPPINGS

### 4.1. New types of systems of generalized quasi-variational inequalities

Noor and Elsanousi [15] introduced the notion of a random variational inequality. We also propose the next systems of generalized quasi-variational inequalities.

Let  $I$  be a non-empty and finite set. For each  $i \in I$ , let  $S_i$  be a countable complete metric space and  $(\Omega_i, \mathcal{Z}_i)$  a measurable space. Let  $(\Omega, \mathcal{F})$  be



the product measurable space  $(\prod_{i \in I} \Omega_i, \otimes_{i \in I} \mathcal{Z}_i)$  and  $\mu$  a probability measure on  $(\Omega, \mathcal{F})$ . For a point  $\omega = (\omega_1, \dots, \omega_n) \in \Omega$ , define the coordinate projections  $\tau_i(\omega) = \omega_i$ . We also denote, for each  $i \in I$ ,  $\text{Meas}(\Omega_i, S_i)$  the set of measurable fuzzy mappings  $f$  from  $(\Omega_i, \mathcal{Z}_i)$  to  $S_i$ . We suppose that there exists a correspondence  $X_i : \Omega_i \rightarrow S_i$  and let  $\mathcal{D}_{X_i}$  be the set  $\{(\mu\tau_i^{-1})g_i^{-1} : g_i \text{ is a measurable selection of } X_i\}$  and  $\mathcal{D}_X := \prod_{i \in I} \mathcal{D}_{X_i}$ .

Let  $A_i : \Omega_i \times \mathcal{D}_X \rightarrow 2^{S_i}$  be a correspondence and  $\psi_i : \Omega_i \times \mathcal{D}_X \times S_i \rightarrow \mathbb{R} \cup \{-\infty, +\infty\}$ .

We associate the next generalized quasi-variational problem with  $A_i$  and  $\psi_i$ :

- (1) Find  $\lambda^* \in \mathcal{D}_X$  such that:
  - i)  $\lambda_i^*(\omega_i) \in A_i(\omega_i, \lambda^*)$ ;
  - ii)  $\sup_{y_i \in A_i(\omega_i, \lambda^*)} \psi_i(\omega_i, \lambda^*, y_i) \leq 0$  for all  $\omega_i \in \Omega_i$ .

If  $S_i$  is a countable completely metrizable topological vector space, we introduce the following definition.

Let  $S'_i$  be the dual space of  $S_i$  and  $G_i : \Omega_i \times S_i \rightarrow 2^{S'_i}$ ,  $A_i : \Omega_i \times \mathcal{D}_X \rightarrow 2^{S_i}$  be correspondences.

- (2) Find  $\lambda^* \in \mathcal{D}_X$  such that:
  - i)  $\lambda^*(\omega_i) \in A_i(\omega_i, \lambda^*)$
  - ii)  $\sup_{y_i \in A_i(\omega_i, \lambda^*)} \sup_{v \in G_i(\omega_i, y_i)} \text{Re}\langle v, \lambda_i^*(\omega_i) - y_i \rangle \leq 0$  for all  $\omega_i \in \Omega_i$ , where the real part of pairing between  $S'_i$  and  $S_i$  is denoted by  $\text{Re}\langle v, x \rangle$  for each  $v \in S'_i$  and  $x \in S_i$ .

We will work on the next fuzzy model:

For each  $i \in I$ , let  $S_i$  be a countable complete metric space and  $(\Omega_i, \mathcal{Z}_i)$  is a measurable space. Let  $A_i : \Omega_i \times \mathcal{D}_X \rightarrow \mathcal{F}(S_i)$  be a fuzzy mapping and let  $a_i : \mathcal{D}_X \rightarrow (0, 1]$  be a fuzzy function. Let  $\psi_i : \Omega_i \times \mathcal{D}_X \times S_i \rightarrow \mathbb{R} \cup \{-\infty, +\infty\}$ .

Now, we are introducing the next type of variational inequality:

- (3) Find  $\lambda^* \in \mathcal{D}_X$  such that for every  $i \in I$ ,
  - i)  $\lambda_i^*(\omega_i) \in (A_i(\omega_i, \lambda^*))_{a_i(\lambda^*)}$ ;
  - ii)  $\sup_{y_i \in (A_i((\omega_i, \lambda^*)))_{a_i(\lambda^*)}} \psi_i(\omega_i, \lambda^*, y_i) \leq 0$  for all  $\omega_i \in \Omega_i$ . where  $(A_{i_{x^*}})_{a_i(x^*)} = \{z \in Y_i : A_{i_{x^*}}(z) \geq a_i(x^*)\}$ .

If  $A_i : X \rightarrow 2^{S_i}$  is a classical correspondence, then we get the variational inequality defined in (1).

Finally, we introduce the following system of generalized quasi-variational inequalities, in case that, for each  $i \in I$ ,  $S_i$  is a countable completely metrizable topological vector space,  $(\Omega_i, \mathcal{Z}_i)$  is measurable space,  $A_i : \Omega_i \times \mathcal{D}_X \rightarrow \mathcal{F}(S_i)$  and  $G_i : \Omega_i \times S_i \rightarrow \mathcal{F}(S'_i)$  are fuzzy mappings and  $a_i : \mathcal{D}_X \rightarrow (0, 1]$ ,  $g_i : S_i \rightarrow (0, 1]$  are fuzzy functions.

(4) Find  $\lambda^* \in \mathcal{D}_X$  such that for every  $i \in I$ ,

$$\begin{cases} \lambda_i^*(\omega_i) \in (A_i(\omega_i, \lambda^*))_{a_i(\lambda^*)}; \\ \sup_{y_i \in (A_i(\omega_i, \lambda^*))_{a_i(\lambda^*)}} \sup_{v \in (G_i(\omega_i, y_i))_{g_i(y_i)}} \operatorname{Re}\langle v, \lambda_i^*(\omega_i) - y_i \rangle \leq 0 \end{cases} \quad \text{for all } \omega_i \in \Omega_i.$$

#### 4.2. The existence of the solutions of the systems of generalized quasi-variational inequalities with random fuzzy mappings

In this section, we are establishing new results concerning the existence of the systems of generalized random quasi-variational inequalities with random fuzzy mappings and we also are stating random fixed point theorems. The proofs rely on the theorem of fuzzy equilibrium existence for the abstract fuzzy economy.

This is our first theorem.

**Theorem 8.** *Let  $I$  be a non-empty and finite set. For each  $i \in I$ ,  $S_i$  is a countable complete metric space and  $(\Omega_i, \mathcal{Z}_i)$  is a measurable space. Let  $(\Omega, \mathcal{Z})$  be the product measurable space  $(\prod_{i \in I} \Omega_i, \otimes_{i \in I} \mathcal{Z}_i)$ , and let  $\mu$  be a probability measure on  $(\Omega, \mathcal{Z})$ . Suppose that the following conditions are satisfied for each  $i \in I$ :*

a) *the correspondence  $X_i : \Omega_i \rightarrow \mathcal{F}(S_i)$  is measurable such that  $(X_i(\cdot))_{z_i} : \Omega_i \rightarrow 2^{S_i}$  has compact values;*

b) *the fuzzy mapping  $A_i$  is such that for each  $\lambda \in \mathcal{D}_{(X(\cdot))_z}$ ,  $(A_i(\cdot, \lambda))_{a_i(\lambda)} : \Omega_i \rightarrow 2^{S_i}$  is measurable and, for all  $\omega_i \in \Omega_i$ ,  $(A_i(\omega_i, \cdot))_{a_i(\cdot)} : \mathcal{D}_{(X(\cdot))_z} \rightarrow 2^{S_i}$  is upper semicontinuous with non-empty compact values;*

Let us assume that the mapping  $\psi_i : \Omega_i \times \mathcal{D}_{(X(\cdot))_z} \times S_i \rightarrow \mathbb{R} \cup \{-\infty, +\infty\}$  is such that:

(c)  *$\lambda \rightarrow \{y \in S_i : \psi_i(\omega, \lambda, y) > 0\} : \mathcal{D}_{(X(\cdot))_z} \rightarrow 2^{S_i}$  is upper semicontinuous with compact values on  $\mathcal{D}_{(X(\cdot))_z}$  for each fixed  $\omega_i \in \Omega_i$ ;*

(d)  *$\lambda_i(\omega_i) \notin \{y \in S_i : \psi_i(\omega_i, \lambda, y) > 0\}$  for each fixed  $(\omega_i, \lambda) \in \Omega_i \times \mathcal{D}_{(X(\cdot))_z}$ ;*

(e) for each  $\omega_i \in \Omega_i$ ,  $\{\lambda \in \mathcal{D}_{(X(\cdot))_z} : \alpha_i(\omega_i, \lambda) > 0\}$  is weakly open in  $\mathcal{D}_{(X(\cdot))_z}$ , where  $\alpha_i : \Omega_i \times \mathcal{D}_{(X(\cdot))_z} \rightarrow \mathbb{R}$  is defined by

$$\alpha_i(\omega_i, \lambda) = \sup_{y \in (A_i(\omega_i, \lambda))_{a_i(\lambda)}} \psi_i(\omega_i, \lambda, y) \text{ for each } (\omega_i, \lambda) \in \Omega_i \times \mathcal{D}_{(X(\cdot))_z};$$

(f)  $\{\omega_i : \alpha_i(\omega_i, \lambda) > 0\} \in \mathcal{Z}_i$  for each  $\lambda \in \mathcal{D}_{(X(\cdot))_z}$ .

Then, there exists  $\lambda^* \in \mathcal{D}_{(X(\cdot))_z}$  such that for every  $i \in I$ ,

i)  $\lambda_i^*(\omega_i) \in (A_i(\omega_i, \lambda^*))_{a_i(\lambda^*)}$ ;

ii)  $\sup_{y \in (A_i(\omega_i, \lambda^*))_{a_i(\lambda^*)}} \psi_i(\omega_i, \lambda^*, y) \leq 0$ .

*Proof.* For every  $i \in I$ , let  $P_i : \Omega_i \times \mathcal{D}_{(X(\cdot))_z} \rightarrow \mathcal{F}(S_i)$  and  $p_i : \mathcal{D}_{(X(\cdot))_z} \rightarrow (0, 1]$  such that  $(P_i(\omega, \lambda))_{p_i(\lambda)} = \{y \in S_i : \psi_i(\omega_i, \lambda, y) > 0\}$  for each  $(\omega_i, \lambda) \in \Omega_i \times \mathcal{D}_{(X(\cdot))_z}$ .

We shall show that the abstract economy  $\Gamma = (I, ((\Omega_i, \mathcal{F}_i), (S_i, A_i, P_i, a_i, b_i, z_i)))_{i \in I}, \mu)$  satisfies all the hypotheses of Theorem 7.

Suppose  $\omega_i \in \Omega_i$ .

According to c), we have that  $\lambda \rightarrow (P_i(\omega_i, \lambda))_{p_i(\lambda)} : \mathcal{D}_{(X(\cdot))_z} \rightarrow 2^{S_i}$  is upper semicontinuous with non-empty values and according to d),  $\lambda_i(\omega_i) \notin (P_i(\omega_i, \lambda))_{p_i(\lambda)}$  for each  $\lambda \in \mathcal{D}_{(X(\cdot))_z}$ .

According to the definition of  $\alpha_i$ , we note that, for each  $\omega_i \in \Omega_i$ ,  $\{\lambda \in \mathcal{D}_{(X(\cdot))_z} : (A_i(\omega_i, \lambda))_{a_i(\lambda)} \cap (P_i(\omega_i, \lambda))_{p_i(\lambda)} \neq \emptyset\} = \{\lambda \in \mathcal{D}_{(X(\cdot))_z} : \alpha_i(\omega_i, \lambda) > 0\}$  so that  $\{\lambda \in \mathcal{D}_{(X(\cdot))_z} : (A_i(\omega_i, \lambda))_{a_i(\lambda)} \cap (P_i(\omega_i, \lambda))_{p_i(\lambda)} \neq \emptyset\}$  is weakly open in  $\mathcal{D}_{(X(\cdot))_z}$  by e).

According to b) and f), it follows that for each  $\lambda \in \mathcal{D}_{(X(\cdot))_z}$ , the correspondences  $(A_i(\cdot, \lambda))_{a_i(\lambda)} : \Omega_i \rightarrow 2^{S_i}$  and  $(P_i(\omega_i, \lambda))_{p_i(\lambda)} : \Omega_i \rightarrow 2^{S_i}$  are measurable.

Thus, the abstract fuzzy economy  $\Gamma = (I, ((\Omega_i, \mathcal{Z}_i), (S_i, A_i, P_i, a_i, b_i, z_i)))_{i \in I}, \mu)$  satisfies all the hypotheses of Theorem 7. Therefore, there exists  $\lambda^* \in \mathcal{D}_{(X(\cdot))_z}$  such that for every  $i \in I$ :

$\lambda_i^*(\omega_i) \in (A_i(\omega_i, \lambda^*))_{a_i(\lambda^*)}$  and

$(A_i(\omega_i, \lambda^*))_{a_i(\lambda^*)} \cap (P_i(\omega_i, \lambda^*))_{p_i(\lambda^*)} = \emptyset$ ;

that is, there exists  $\lambda^* \in \mathcal{D}_{(X(\cdot))_z}$  such that for every  $i \in I$ :

i)  $\lambda_i^*(\omega_i) \in (A_i(\omega_i, \lambda^*))_{a_i(\lambda^*)}$ ;

ii)  $\sup_{y \in (A_i(\omega_i, \lambda^*))_{a_i(\lambda^*)}} \psi_i(\omega_i, \lambda^*, y) \leq 0$ .

If  $|I|=1$ , we obtain the following corollary.

**Corollary 9.** *Let  $S$  be a countable complete metric space,  $(\Omega, \mathcal{Z}, \mu)$  be a measure space. Suppose that the following conditions are satisfied:*

a) the fuzzy mapping  $X : \Omega \rightarrow \mathcal{F}(S)$  is measurable such that  $(X(\cdot))_z : \Omega_i \rightarrow 2^S$  has compact values;

b) the fuzzy mapping  $A$  is such that for each  $\lambda \in \mathcal{D}_{(X(\cdot))_z}$ ,  $(A(\cdot, \lambda))_{a(\lambda)} : \Omega \rightarrow 2^S$  is measurable and, for all  $\omega \in \Omega$ ,  $(A(\omega, \cdot))_{a(\cdot)} : \mathcal{D}_{(X(\cdot))_z} \rightarrow 2^S$  is upper semicontinuous with non-empty compact values;

The mapping  $\psi : \Omega \times \mathcal{D}_{(X(\cdot))_z} \times S \rightarrow \mathbb{R} \cup \{-\infty, +\infty\}$  is such that:

(c)  $\lambda \rightarrow \{y \in Y : \psi(\omega, \lambda, y) > 0\} : \mathcal{D}_{(X(\cdot))_z} \rightarrow 2^S$  is upper semicontinuous with compact values on  $\mathcal{D}_{(X(\cdot))_z}$  for each fixed  $\omega \in \Omega$ ;

(d)  $\lambda(\omega) \notin \{y \in S : \psi(\omega, \lambda, y) > 0\}$  for each fixed  $(\omega, \lambda) \in \Omega \times \mathcal{D}_{(X(\cdot))_z}$ ;

(e) for each  $\omega \in \Omega$ ,  $\{\lambda \in \mathcal{D}_{(X(\cdot))_z} : \alpha(\omega, \lambda) > 0\}$  is weakly open in  $\mathcal{D}_{(X(\cdot))_z}$ , where  $\alpha : Z \times \mathcal{D}_{(X(\cdot))_z} \rightarrow \mathbb{R}$  is defined by

$$\alpha(\omega, \lambda) = \sup_{y \in (A(\omega, \lambda))_{a(\lambda)}} \psi(\omega, \lambda, y) \text{ for each } (\omega, \lambda) \in Z \times \mathcal{D}_{(X(\cdot))_z};$$

(f)  $\{\omega : \alpha(\omega, \lambda) > 0\} \in \mathcal{Z}$  for each  $\lambda \in \mathcal{D}_{(X(\cdot))_z}$ .

Then, there exists  $\lambda^* \in \mathcal{D}_{(X(\cdot))_z}$  such that,

i)  $\lambda^*(\omega) \in (A(\omega, \lambda^*))_{a_i(\lambda^*)}$ ;

ii)  $\sup_{y \in (A(\omega, \lambda^*))_{a(\lambda^*)}} \psi(\omega, \lambda^*, y) \leq 0$ .

As a consequence of Theorem 8, we prove the following Tan and Yuan's type (1995) random quasi-variational inequality with random fuzzy mappings.

**Theorem 10.** Let  $I$  be a non-empty and finite set. For each  $i \in I$ ,  $S_i$  is a countable completely metrizable topological vector space and  $(\Omega_i, \mathcal{Z}_i)$  is a measurable space. Let  $(\Omega, \mathcal{Z})$  be the product measurable space  $(\prod_{i \in I} \Omega_i, \otimes_{i \in I} \mathcal{Z}_i)$ , and  $\mu$  a probability measure on  $(\Omega, \mathcal{Z})$ . Suppose that the following conditions are satisfied for each  $i \in I$  :

a) the correspondence  $X_i : \Omega_i \rightarrow \mathcal{F}(S_i)$  is measurable such that  $(X_i(\cdot))_{z_i} : \Omega_i \rightarrow 2^{S_i}$  has compact values;

b) the mapping  $A_i$  is such that for each  $\lambda \in \mathcal{D}_{(X(\cdot))_z}$ ,  $(A_i(\cdot, \lambda))_{a_i(\lambda)} : \Omega_i \rightarrow 2^{S_i}$  is measurable and, for all  $\omega_i \in \Omega_i$ ,  $(A_i(\omega_i, \cdot))_{a_i(\cdot)} : \mathcal{D}_{(X(\cdot))_z} \rightarrow 2^{S_i}$  is upper semicontinuous with non-empty compact values;

$G_i : \Omega_i \times S \rightarrow \mathcal{F}(S')$  and  $g_i : S \rightarrow (0, 1]$  are such that:

(c) For each fixed  $(\omega_i, y) \in \Omega_i \times S$ ,  $\lambda \rightarrow \{y \in S : \sup_{u \in (G_i(\omega_i, y))_{g_i(y)}} \operatorname{Re}\langle u, \lambda_i(\omega_i) - y \rangle > 0\} : \mathcal{D}_{(X(\cdot))_z} \rightarrow 2^S$  is upper semicontinuous with compact values;

(d) for each fixed  $\omega_i \in \Omega_i$ , the set

$\{\lambda \in \mathcal{D}_{(X(\cdot))_z} : \sup_{y \in (A_i(\omega_i, \lambda))_{a_i(\lambda)}} \sup_{u \in (G_i(\omega_i, y))_{g_i(y)}} \operatorname{Re}\langle u, \lambda_i(\omega_i) - y \rangle > 0\}$  is weakly open in  $\mathcal{D}_{(X(\cdot))_z}$

(e)  $\{\omega_i \in \Omega_i : \sup_{u \in (G_i(\omega, y))_{g_i(y)}} \operatorname{Re}\langle u, \lambda_i(\omega_i) - y \rangle > 0\} \in \mathcal{Z}_i$  for each  $\lambda \in \mathcal{D}_{(X(\cdot))_z}$ .

Then, there exists  $\lambda^* \in \mathcal{D}_{(X(\cdot))_z}$  such that for every  $i \in I$ :

i)  $\lambda_i^*(\omega_i) \in (A_i(\omega_i, \lambda^*))_{a_i(\lambda^*)}$ ;

ii)  $\sup_{u \in (G_i(\omega_i, y))_{g_i(y)}} \operatorname{Re}\langle u, \lambda_i^*(\omega_i) - y \rangle \leq 0$  for all  $y \in (A_i(\omega_i, \lambda^*))_{a_i(\lambda^*)}$ .

*Proof.* Let us define  $\psi_i : \Omega_i \times \mathcal{D}_{(X(\cdot))_z} \times S \rightarrow \mathbb{R} \cup \{-\infty, +\infty\}$  by

$\psi_i(\omega_i, \lambda, y) = \sup_{u \in (G_i(\omega_i, y))_{g_i(y)}} \operatorname{Re}\langle u, \lambda_i(\omega_i) - y \rangle$  for each  $(\omega_i, \lambda, y) \in \Omega_i \times \mathcal{D}_{(X(\cdot))_z} \times S$ .

We have that  $\lambda_i(\omega_i) \notin \{y \in S : \psi_i(\omega_i, \lambda, y) > 0\}$  for each fixed  $(\omega_i, \lambda) \in \Omega_i \times \mathcal{D}_{(X(\cdot))_z}$ .

All the hypotheses of Theorem 8 are satisfied. According to Theorem 8, there exists  $\lambda^* \in \mathcal{D}_{(X(\cdot))_z}$  such that  $\lambda_i^*(\omega_i) \in (A_i(\omega_i, \lambda^*))_{a_i(\lambda^*)}$  for every  $i \in I$  and

$\sup_{y \in (A_i(\omega_i, \lambda^*))_{a_i(\lambda^*)}} \sup_{u \in (G_i(\omega_i, y))_{g_i(y)}} \operatorname{Re}\langle u, \lambda_i^*(\omega_i) - y \rangle \leq 0$  for every  $i \in I$ .

If  $|I| = 1$ , we obtain the following corollary.

**Corollary 11.** *Let  $S$  be a countable completely metrizable topological vector space and let  $(\Omega, \mathcal{Z}, \mu)$  be a measure space. Suppose that the following conditions are satisfied:*

a) the fuzzy mapping  $X : \Omega \rightarrow \mathcal{F}(S)$  is measurable such that  $(X(\cdot))_z : \Omega \rightarrow 2^S$  has compact values;

b) the fuzzy mapping  $A$  is such that for each  $\lambda \in \mathcal{D}_{(X(\cdot))_z}$ ,  $(A(\cdot, \lambda))_{a(\lambda)} : \Omega \rightarrow 2^S$  is measurable and, for all  $\omega \in \Omega$ ,  $(A(\omega, \cdot))_{a(\cdot)} : \mathcal{D}_{(X(\cdot))_z} \rightarrow 2^S$  is upper semicontinuous with non-empty compact values;

$G : \Omega \times S \rightarrow \mathcal{F}(S')$  and  $g : S \rightarrow (0, 1]$  are such that:

(c) For each fixed  $(\omega, y) \in \Omega \times S$ ,  $\lambda \rightarrow \{y \in S : \sup_{u \in (G(\omega, y))_{g(y)}} \operatorname{Re}\langle u, \lambda(\omega) - y \rangle > 0\} : \mathcal{D}_{(X(\cdot))_z} \rightarrow 2^S$  is upper semicontinuous with compact values;

(d) for each fixed  $\omega \in \Omega$ , the set

$\{\lambda \in \mathcal{D}_{(X(\cdot))_z} : \sup_{y \in (A(\omega, \lambda))_{a(\lambda)}} \sup_{u \in (G(\omega, y))_{g(y)}} \operatorname{Re}\langle u, \lambda(\omega) - y \rangle > 0\}$  is weakly open in  $\mathcal{D}_{(X(\cdot))_z}$

(e)  $\{(\omega, \lambda) : \sup_{u \in (G(\omega, y))_{g(y)}} \operatorname{Re}\langle u, \lambda(\omega) - y \rangle > 0\} \in \mathcal{Z}$  for each  $\lambda \in \mathcal{D}_{(X(\cdot))_z}$ .

Then, there exists  $\lambda^* \in \mathcal{D}_{(X(\cdot))_z}$  such that:

i)  $\lambda^*(\omega) \in (A(\omega, \lambda^*))_{a(\lambda^*)}$ ;

ii)  $\sup_{u \in (G(\omega, y))_{g(y)}} \operatorname{Re}\langle u, \lambda^*(\omega) - y \rangle \leq 0$  for all  $y \in (A(\omega, \lambda^*))_{a(\lambda^*)}$ .

We obtain the following random fixed point theorem as a particular case of Theorem 8.

**Theorem 12.** *Let  $I$  be a non-empty finite set. For each  $i \in I$ ,  $S_i$  is a countable complete metric space and  $(\Omega_i, \mathcal{Z}_i)$  is a measurable space. Let  $(\Omega, \mathcal{Z})$  be the product measurable space  $(\prod_{i \in I} \Omega_i, \otimes_{i \in I} \mathcal{Z}_i)$ , and  $\mu$  a probability measure on  $(\Omega, \mathcal{Z})$ . Suppose that the following conditions are satisfied for each  $i \in I$ :*

a) *the fuzzy mapping  $X_i : \Omega_i \rightarrow \mathcal{F}(S_i)$  is measurable and  $(X_i(\cdot))_{z_i} : \Omega_i \rightarrow 2^{S_i}$  has compact values;*

b) *the fuzzy mapping  $A_i$  is such that for each  $\lambda \in \mathcal{D}_{(X(\cdot))_z}$ ,  $(A_i(\cdot, \lambda))_{a_i(\lambda)} : \Omega_i \rightarrow 2^{S_i}$  is measurable and, for all  $\omega_i \in \Omega_i$ ,  $(A_i(\omega_i, \cdot))_{a_i(\cdot)} : \mathcal{D}_{(X(\cdot))_z} \rightarrow 2^{S_i}$  is upper semicontinuous with non-empty compact values.*

*Then, there exists  $\lambda^* \in \mathcal{D}_{(X(\cdot))_z}$  such that for every  $i \in I$ ,  $\lambda_i^*(\omega_i) \in (A_i(\omega_i, \lambda^*))_{a_i(\lambda^*)}$ .*

If  $|I| = 1$ , we obtain the following result.

**Theorem 13.** *Let  $S$  be a countable complete metric space and let  $(\Omega, \mathcal{Z}, \mu)$  be a measure space. Suppose that the following conditions are satisfied:*

a) *the fuzzy mapping  $X : \Omega \rightarrow \mathcal{F}(S)$  is measurable and  $(X(\cdot))_z : \Omega \rightarrow 2^S$  has compact values;*

b) *the mapping  $A$  is such that for each  $\lambda \in \mathcal{D}_{(X(\cdot))_z}$ ,  $(A(\cdot, \lambda))_{a(\lambda)} : \Omega \rightarrow 2^S$  is measurable and, for all  $\omega \in \Omega$ ,  $(A(\omega, \cdot))_{a(\cdot)} : \mathcal{D}_{(X(\cdot))_z} \rightarrow 2^S$  is upper semicontinuous with non-empty compact values.*

*Then, there exists  $\lambda^* \in \mathcal{D}_{(X(\cdot))_z}$  such that  $\lambda^*(\omega) \in (A(\omega, \lambda^*))_{a(\lambda^*)}$ .*

## 5. Classical systems of generalized quasi-variational inequalities and random fixed point theorems

If we consider classical correspondences in the last section, we obtain several results concerning systems of generalized random quasi-variational inequalities and random fixed points, which are new in the literature. We present Theorem 14 as a consequence of Theorem 8.

**Theorem 14.** *Let  $I$  be a non-empty finite set. For each  $i \in I$ ,  $S_i$  is a countable complete metric space and  $(\Omega_i, \mathcal{Z}_i)$  is a measurable space. Let  $(\Omega, \mathcal{Z})$  be the product measurable space  $(\prod_{i \in I} \Omega_i, \otimes_{i \in I} \mathcal{Z}_i)$ , and let  $\mu$  be a probability measure on  $(\Omega, \mathcal{Z})$ . Suppose that the following conditions are satisfied for each  $i \in I$  :*

a) *the correspondence  $X_i : \Omega_i \rightarrow 2^{S_i}$  is measurable and  $(X_i(\cdot))_{z_i} : \Omega_i \rightarrow 2^{S_i}$  has compact values;*

b) *the correspondence  $A_i$  is such that for each  $\lambda \in \mathcal{D}_X$ ,  $A_i(\cdot, \lambda) : \Omega_i \rightarrow 2^{S_i}$  is measurable and, for all  $\omega_i \in \Omega_i$ ,  $A_i(\omega_i, \cdot) : \mathcal{D}_X \rightarrow 2^{S_i}$  is upper semicontinuous with non-empty compact values;*

Let us assume that the mapping  $\psi_i : \Omega_i \times \mathcal{D}_X \times S \rightarrow \mathbb{R} \cup \{-\infty, +\infty\}$  is such that:

(c)  *$\lambda \rightarrow \{y \in Y : \psi_i(\omega, \lambda, y) > 0\} : \mathcal{D}_X \rightarrow 2^S$  is upper semicontinuous with compact values on  $\mathcal{D}_X$  for each fixed  $\omega_i \in \Omega_i$ ;*

(d)  *$\lambda_i(\omega_i) \notin \{y \in S : \psi_i(\omega_i, \lambda, y) > 0\}$  for each fixed  $(\omega_i, \lambda) \in \Omega_i \times \mathcal{D}_X$ ;*

(e) *for each  $\omega_i \in \Omega_i$ ,  $\{\lambda \in \mathcal{D}_X : \alpha_i(\omega_i, \lambda) > 0\}$  is weakly open in  $\mathcal{D}_X$ , where  $\alpha_i : \Omega_i \times \mathcal{D}_X \rightarrow \mathbb{R}$  is defined by  $\alpha_i(\omega_i, \lambda) = \sup_{y \in A_i(\omega_i, \lambda)} \psi_i(\omega_i, \lambda, y)$  for each  $(\omega_i, \lambda) \in \Omega_i \times \mathcal{D}_X$ ;*

(f)  *$\{\omega_i : \alpha_i(\omega_i, \lambda) > 0\} \in \mathcal{Z}_i$  for each  $\lambda \in \mathcal{D}_X$ .*

*Then, there exists  $\lambda^* \in \mathcal{D}_X$  such that for every  $i \in I$ ,*

i)  *$\lambda_i^*(\omega_i) \in A_i(\omega_i, \lambda^*)$ ;*

ii)  *$\sup_{y \in A_i(\omega_i, \lambda^*)} \psi_i(\omega_i, \lambda^*, y) \leq 0$ .*

The next theorems concern the existence of the random fixed point for correspondences with values in complete countable metric spaces.

**Theorem 15.** *Let  $I$  be a non-empty finite set. For each  $i \in I$ ,  $S_i$  is a countable complete metric space and  $(\Omega_i, \mathcal{Z}_i)$  is a measurable space. Let  $(\Omega, \mathcal{Z})$  be the product measurable space  $(\prod_{i \in I} \Omega_i, \otimes_{i \in I} \mathcal{Z}_i)$ , and let  $\mu$  be a probability measure on  $(\Omega, \mathcal{Z})$ . Suppose that the following conditions are satisfied for each  $i \in I$  :*

a) *the correspondence  $X_i : \Omega_i \rightarrow 2^{S_i}$  is measurable with compact values;*

b) *for each  $\lambda \in \mathcal{D}_X$ ,  $A_i(\cdot, \lambda) : \Omega_i \rightarrow 2^{S_i}$  is measurable and, for all  $\omega_i \in \Omega_i$ ,  $A_i(\omega_i, \cdot) : \mathcal{D}_X \rightarrow 2^{S_i}$  is upper semicontinuous with non-empty compact values.*

*Then, there exists  $\lambda^* \in \mathcal{D}_X$  such that for every  $i \in I$ ,  $\lambda_i^*(\omega_i) \in A_i(\omega_i, \lambda^*)$ .*

If  $|I| = 1$ , we obtain the following result.

**Theorem 16.** *Let  $S$  be a countable complete metric space and let  $(\Omega, \mathcal{Z}, \mu)$  be a measure space. Suppose that the following conditions are satisfied:*

- a) the correspondence  $X : \Omega \rightarrow 2^S$  is measurable with compact values;*
  - b) for each  $\lambda \in \mathcal{D}_X$ ,  $A(\cdot, \lambda) : \Omega \rightarrow 2^S$  is measurable and, for all  $\omega \in \Omega$ ,  $A(\omega, \cdot) : \mathcal{D}_X \rightarrow 2^S$  is upper semicontinuous with non-empty compact values.*
- Then, there exists  $\lambda^* \in \mathcal{D}_X$  such that  $\lambda^*(\omega) \in A(\omega, \lambda^*)$ .*

## References

- [1] C. D. Aliprantis, K. C. Border, *Infinite Dimensional Analysis: A Hitchhiker's Guide*. Springer-Verlag, Berlin, 1994.
- [2] Arrow K. J., Debreu G. (1954), *Existence of an Equilibrium for a Competitive Economy*. *Econometrica*, **22**, 265-290.
- [3] Borglin A. and Keiding H. (1976), *Existence of equilibrium actions and of equilibrium: A note on the 'new' existence theorem*. *J. Math. Econom.* **3**, 313-316.
- [4] S.-S. Chang and K. K. Tan, *Equilibria and maximal elements of abstract fuzzy economies and qualitative fuzzy games*, *Fuzzy Sets and Systems* **125** (2002), 389-399.
- [5] Debreu G. (1952), *A social equilibrium existence theorem*, *Proc. Natl. Acad. Sci. USA* **38**, 886-903.
- [6] Fan K. (1952), *Fixed-point and minimax theorems in locally convex topological linear spaces*, *Proc. Nat. Acad. Sci. U.S.A.* **38**, 121-126.
- [7] Gwinner J. and Raciti F. (2006), *On a class of random variational inequalities on random sets*, *Numerical Functional Analysis and Optimization*, 27(5-6):619-636.
- [8] Gwinner J. and Raciti F. (2012), *Some equilibrium problems under uncertainty and random variational inequalities*, *Ann. Oper. Res.*, 200: 299-319.



- [9] Harsanyi, J.C.: Games with incomplete information played by “Bayesian” players. *Management Science* 14, 159–189 (1967).
- [10] Huang N.-J. (2001), *Existence of equilibrium for generalized abstract fuzzy economies*, *Fuzzy Sets and Systems* **117**, 151-156.
- [11] Kim W. K. and Lee K. H. (1998), *Fuzzy fixed point and existence of equilibria of fuzzy games*, *J. Fuzzy Math.* **6**, 193-202.
- [12] Kim W. K. and Lee K. H.(1999), *Existence of equilibria in generalized fuzzy games*, *J. Chungcheong Math. Soc.* **12**, 53-61.
- [13] Kim W. K. and Lee K. H.(2001), *Generalized fuzzy games and fuzzy equilibria*, *Fuzzy Sets and Systems* **122**, 293-301.
- [14] Nash J. F. (1950), *Equilibrium points in  $n$ -person games*. *Proc. Nat. Acad. Sci. U. S. A.*, **36**, 1, 48-49.
- [15] Noor M.A. and Elsanousi S.A.(1993), *Iterative algorithms for random variational inequalities*, *Pan Amer. Math. Journal* 3, 39-50.
- [16] Patriche M. (2009), *Equilibria for free abstract fuzzy economies*, *An. St. Ovidius University of Constanta*, Volume 17(2), 143-154.
- [17] Patriche M., *Bayesian abstract economy with a measure space of agents*, *Abstract and Applied Analysis*, Volume 2009 (2009), Article ID 523619.
- [18] Patriche M.(2010), *Abstract fuzzy economies and fuzzy equilibrium pairs*, *Mathematical Reports*, 12(62), 4, 373 – 382.
- [19] Patriche M., *Bayesian abstract economies*, *Optimization*, Vol. 60, Nr 4, 2011, 451-461.
- [20] Patriche M. (2011a), *Existence of fuzzy equilibria for fuzzy abstract economies with  $Q^1$ -majorized correspondences*, *Recent Researches in Neural Networks, Fuzzy Systems, Evolutionary Computing and Automation*, *International Conference on Fuzzy Systems*, 11 – 13 April, Brasov.
- [21] Patriche M. (2011b), *Equilibrium in games and competitive economies*, *The Publishing House of the Romanian Academy*.

- [22] Patriche M., Existence of equilibrium for an abstract economy with private information and a countable space of actions, Mathematical reports, to appear.
- [23] R. Radner, 1968. Competitive Equilibrium Under Uncertainty, *Econometrica*, 36 1, 31-58.
- [24] Shafer W. and Sonnenschein H. (1975), *Equilibrium in abstract economies without ordered preferences*, *Journal of Mathematical Economics* **2**, 345-348.
- [25] Wang L., ChoY Je, Huang N. (2011), *The robustness of generalized abstract fuzzy economies in generalized convex spaces*, *Fuzzy Sets and Systems*, 176 (1), 56-63.
- [26] Tan K.-K. and Yuan G. X.-Z. (1995), Random equilibria of random generalized games with applications to non-compact random quasi-variational inequalities, *Topological Methods in Nonlinear Analysis*, *Journal of the Juliusz Schauder Center*, Volume 5, 59–82.
- [27] L Wang, Y Je Cho, N Huang, *The robustness of generalized abstract fuzzy economies in generalized convex spaces*, *Fuzzy Sets and Systems*, 176 (1) (2011), 56-63.
- [28] Yannelis N. C. and Prabhakar N. D. (1983), Existence of maximal elements and equilibrium in linear topological spaces, *J. Math. Econom.* **12**, 233-245.
- [29] H. Yu and Z. Zhang, *Pure strategy equilibria in games with countable actions*, *Journal of Mathematical Economics* 43 (2007), 192-200.
- [30] Yuan X. Z. (1988), *The Study of Minimax inequalities and Applications to Economies and Variational inequalities*, *Memoirs of the American Society*, vol **132**, Nr 625.
- [31] L. A. Zadeh, *Fuzzy sets*, *Inform and Control* **8** (1965), 338-353.