Stationary modes and integrals of motion in nonlinear lattices with \mathcal{PT} -symmetric linear part

D. A. Zezyulin¹ and V. V. Konotop^{1,2}

¹Centro de Física Teórica e Computacional, Faculdade de Ciências, Universidade de Lisboa, Avenida Professor Gama Pinto 2, Lisboa 1649-003, Portugal
² Departamento de Física, Faculdade de Ciências, Universidade de Lisboa, Campo Grande, Ed. C8, Piso 6, Lisboa 1749-016, Portugal

Abstract. We consider finite-dimensional nonlinear systems with linear part described by a parity-time (\mathcal{PT} -) symmetric operator. We investigate bifurcations of stationary nonlinear modes from the eigenstates of the linear operator and consider a class of \mathcal{PT} -symmetric nonlinearities allowing for existence of the families of nonlinear modes. We pay particular attention to the situations when the underlying linear \mathcal{PT} -symmetric operator is characterized by the presence of degenerate eigenvalues or exceptional-point singularity. In each of the cases we construct formal expansions for small-amplitude nonlinear modes. We also report a class of nonlinearities allowing for the system to admit one or several integrals of motion, which turn out to be determined by the pseudo-Hermiticity of the nonlinear operator.

PACS numbers: 63.20.Pw, 05.45.Yv, 11.30.Er, 42.65.Wi

Submitted to: J. Phys. A: Math. Gen.

1. Introduction

Richness of solutions is a typical feature of nonlinear problems. Since a complete characterization of a vast set of all solutions usually is not possible, a simpler problem of classification of admissible *types* of solutions of a nonlinear system becomes a relevant and still nontrivial task. In this respect, there is an important distinction between conservative and non-conservative (i.e. dissipative) systems. Conservative nonlinear systems typically possess continuous *families* of nonlinear stationary solutions (modes) which exist for fixed values of the system parameters. A family of nonlinear modes can be parametrized by an "internal" quantity (or few quantities), like, for instance, L^2 -norm of the mode (which can correspond to energy, number of particles, or total power, depending on the particular physical statement) or its frequency (i.e. propagation constant or chemical potential). The situation becomes different when one considers a dissipative system characterized by the presence of gain and losses. In order to admit a stationary solution, such a system requires the absorbed energy to be compensated exactly by the gain. The requirement of energy balance imposes an additional constraint on the shape of the solution which has to have a nontrivial energy flow allowing for the energy transfer from the gain regions to the lossy ones. As a result, for given values of the system

parameters, a typical dissipative nonlinear system does not admit continuous families of nonlinear modes. Instead, isolated nonlinear modes appear. From the dynamical point of view, these isolated modes typically behave as attractors (being stable) or repellers (being unstable). In order to obtain a continuous set of physically distinct dissipative nonlinear modes, one has to vary parameters of the system, i.e. change the system itself. Then, instead of continuous families, characteristic for conservative systems with fixed parameters, the dissipative system admits continuous *branches* of nonlinear modes which are obtained through variation of the system parameters. We thus distinguish between parametric families and branches. The described dichotomy is particularly well-known in the context of the discussion of dissipative solitons vs. conventional solitons in the nonlinear Schrödinger-like and complex Ginzburg–Landau equations [1].

In this context, the physical concept of parity-time (\mathcal{PT}) symmetry [2] deserves a particular attention. This is because the nonlinear systems with \mathcal{PT} symmetry appear to occupy an "intermediate position" between the conventional conservative and dissipative systems. Whereas the systems governed by a \mathcal{PT} -symmetric operators are of dissipative nature and require a solution to generate a nontrivial energy flow in order to maintain itself, the exact gain and loss balance inherent to the \mathcal{PT} symmetry allows for the system to possess in some cases continuous families of nonlinear modes apart from the branches typical to dissipative systems. There are numerous studies where both branches and families of the nonlinear modes in \mathcal{PT} -symmetric systems have been reported. In particular, exact solutions were reported for the nonlinear \mathcal{PT} symmetric dimer (a system of two coupled nonlinear oscillators) in [3]. Families of nonlinear discrete modes received particular emphasis in [4, 5, 6, 7], while branches of the solutions were obtained for finite lattices like \mathcal{PT} -symmetric trimers and oligomers [5, 8, 9], as well as for infinite \mathcal{PT} -symmetric chains [6, 7, 10, 11].

Localized modes (or solitons) are also known to exist in extended nonlinear systems, where they first were found in the presence of a periodic \mathcal{PT} -symmetric potential and cubic nonlinearity [12]. Families of nonlinear modes in continuous systems were intensively investigated for gap solitons in Kerr [13] and χ^2 [14] media, in \mathcal{PT} -symmetric parabolic potential [15], in defect \mathcal{PT} -symmetric lattices [16], in systems with $\chi^{(2)}$ nonlinearity with embedded \mathcal{PT} -symmetric defect [17], and in the presence of a \mathcal{PT} -symmetric superlattice [18]. Lattices with \mathcal{PT} -symmetric nonlinear potentials were proposed in [19, 20]. Combined effect of linear and nonlinear \mathcal{PT} symmetric lattices was explored in discrete [21, 22] and continuous [23] statements. Models of two coupled waveguides (so-called \mathcal{PT} -symmetric nonlinear couplers) are also known to support solitons [24] and breathers [25]. Solitons in a system consisting of a necklace of optical \mathcal{PT} -symmetric waveguides were recently addressed in [26].

In the most of the mentioned studies, the nonlinearity inherent to the system was fixed by the physical statement of the problem. In optical applications one usually considers the cubic (namely, Kerr type) or quadratic (χ^2 -type) nonlinearities, while realization of \mathcal{PT} symmetry Bose-Einstein condensates [27, 28, 29] also implies the Kerr type nonlinearity. In many situations, however, the nonlinearity can be changed, which makes it relevant to study physical systems obeying the same linear properties but having nonlinearities of different types. This leads us to the *first goal* of the present paper, which is the effect of the type of the nonlinearity on the existence and dynamics of nonlinear modes. In particular, we will study bifurcations of families of nonlinear modes from the eigenstates of the underlying linear operator (i.e. the operator describing the linear part of the system) and argue that the bifurcations are possible only if the nonlinear operator admits a certain symmetry which can be conveniently referred to as weak \mathcal{PT} symmetry. This terminology is to emphasize that the class of weakly \mathcal{PT} -symmetric nonlinear operators contains as a subset a class of \mathcal{PT} -symmetric nonlinear operators which (by analogy with the definition of \mathcal{PT} symmetry for linear operators) commute with the \mathcal{PT} operator. The second goal of our paper is to study relevance of \mathcal{PT} -symmetric nonlinearities for dynamical properties of the system. In particular, we will study possibilities for the nonlinear system (1) to admit integrals of motion and show that if the nonlinearity is \mathcal{PT} symmetric (and has a certain additional simple property), then the nonlinear system does admit an integral of motion. On the other hand, we will also argue that integrals of motion can exist if the nonlinear operator is pseudo-Hermitian [30].

Apart from the type of nonlinearity, properties of the nonlinear modes may strongly depend on the character of the spectrum of the underlying linear operator. Most of the above mentioned studies dealt with the situations when the linear spectrum consists of simple real eigenvalues. Recently, nonlinear modes bifurcating from the doubly degenerate linear eigenstates were reported in [5]. The *third goal* of the paper is to perform the analysis of bifurcations of stationary nonlinear modes in a situation when the linear operator has a degenerate eigenstate of finite multiplicity or *exceptional point* singularities [31]. For both those cases we develop formal asymptotical expansions which describe bifurcations of the nonlinear modes from degenerate linear eigenstates. In order to construct the expansions, we explore the structure of the invariant subspace associated with the multiple eigenvalue. For the case of a semi-simple eigenvalue, we show that its invariant subspace can be spanned by a basis consisting of \mathcal{PT} -invariant linearly independent eigenvectors (see Proposition 1). For the situation when the geometric multiplicity of the eigenvector is less than its algebraic multiplicity, we show that there exists a \mathcal{PT} -invariant generalized eigenvector (Proposition 2).

The organization of the paper is as follows. In Sec. 2 we specify the chosen model and make some general remarks related to the subject. Next, we address the effect of the nonlinearity on bifurcation of nonlinear modes from the eigenstates of the linear lattice. We consider the essentially different situations of bifurcations from simple (Sec. 3) and semi-simple (Sec. 4) eigenvalues, as well as the nonlinear modes in the presence of the exceptional point singularity (Sec. 5). In Sec. 6 we address the question about the nonlinearities allowing nonlinear \mathcal{PT} -symmetric systems to have integrals of motion. The outcomes are summarized in the Conclusion.

2. The model and general remarks

To proceed with our studies, we specify the chosen model. In the present work, we study a nonlinear system of the form

$$i\dot{\mathbf{q}} = -H(\gamma)\mathbf{q} - F(\mathbf{q})\mathbf{q},\tag{1}$$

where $\mathbf{q} = \mathbf{q}(t)$ is a column-vector of N elements, an overdot stands for the derivative with respect to time $\dot{\mathbf{q}} = d\mathbf{q}/dt$ (or with respect to the propagation distance in optical terminology). The linear part of the finite lattice (1) is described by a $N \times N$ symmetric matrix $H(\gamma)$. We consider \mathcal{PT} -symmetric Hamiltonians $H(\gamma)$ [2], meaning that there exist parity, \mathcal{P} , and time-reversal, \mathcal{T} , operators such that $\mathcal{P}^2 = \mathcal{T}^2 = I$, $[\mathcal{P}, \mathcal{T}] = 0$ and $[\mathcal{PT}, H] = 0$ (hereafter I is the identity operator). The mentioned properties also imply that $(\mathcal{PT})^2 = I$. As it is customary, we define the time reversal operator \mathcal{T} by the complex conjugation, i.e. $\mathcal{T}\mathbf{q} = \mathbf{q}^*$ (hereafter an asterisk stands for the complex conjugation), and consider \mathcal{P} to be a matrix representation of the linear parity reversal operator. The above definition of \mathcal{T} ensures that entries of the matrix \mathcal{P} are real [32]. Moreover, \mathcal{PT} symmetry implies $H\mathcal{P} - \mathcal{P}H^* = 0$, and hence

$$H^{\dagger} = H^* = \mathcal{P}H\mathcal{P},\tag{2}$$

i.e. the pseudo-Hermiticity of $H(\gamma)$ [30]. We also require the matrix \mathcal{P} to be symmetric (for the discussion of the relevance of this requirement see [32]):

$$\mathcal{P} = \mathcal{P}^T = \mathcal{P}^\dagger. \tag{3}$$

The nonlinear operator $F(\mathbf{q})$ is a $N \times N$ matrix whose elements depend on the field \mathbf{q} . We focus on the case of the cubic nonlinearity when the entries $F_{pj}(\mathbf{q})$ of the matrix $F(\mathbf{q})$ are given as

$$F_{pj}(\mathbf{q}) = \mathbf{q}^{\dagger} \mathcal{F}_{pj} \mathbf{q} = \sum_{l,m=1}^{N} f_{pj}^{lm} q_{l}^{*} q_{m}, \quad p, j = 1, 2, \dots, N,$$
(4)

i.e. \mathcal{F}_{pj} are $N \times N$ matrices with time-independent entries f_{pj}^{lm} , l, m = 1, 2, ..., N. In other words, $F_{ij}(\mathbf{q})$ is a linear combination of pair-wise products of the elements of the vectors $\mathbf{q}(t)$ and $\mathbf{q}^*(t)$ (some of the coefficients f_{pj}^{lm} can be equal to zero). Notice that if $f_{pj}^{lm} = (f_{jp}^{ml})^* = f_{lj}^{pm} = f_{pm}^{lj}$ then in the dissipationless limit [i.e. at values γ for which $H(\gamma)$ becomes Hermitian] the system (1) is Hamiltonian i.e.

$$(F(\mathbf{q})\mathbf{q})_n = \frac{1}{2} \frac{\partial}{\partial q_n^*} \sum_{j,l,m,p=1}^N q_j^* q_l^* f_{jm}^{lp} q_m q_p, \quad n = 1, 2, \dots, N.$$

The particular examples of nonlinearities $F(\mathbf{q})$ considered below in this paper are of this type. Nevertheless, the analysis we develop is applicable for a general case, which in particular includes \mathcal{PT} -symmetric *dissipative* nonlinearities [19, 20, 21, 22], when the nonlinear operator $F(\mathbf{q})$ results in losses and gain.

Stationary nonlinear modes of system (1) correspond to solutions of the form $\mathbf{q}(t) = e^{ibt}\mathbf{w}$, where \mathbf{w} is a time-independent column vector solving the stationary nonlinear problem

$$b\mathbf{w} = H(\gamma)\mathbf{w} + F(\mathbf{w})\mathbf{w}, \qquad \text{Im}\,b = 0.$$
 (5)

Reality of b is required for the existence of stationary nonlinear modes, and equality $F(\mathbf{w}) = F(\mathbf{q})$ readily follows from Eq. (4). In the linear limit, which formally corresponds to $F(\mathbf{w}) = 0$, Eq. (5) is reduced to a linear eigenvalue problem for the operator $H(\gamma)$:

$$b\tilde{\mathbf{w}} = H(\gamma)\tilde{\mathbf{w}} \tag{6}$$

(hereafter tildes stand to indicate the eigenvectors and eigenvalues belonging to the linear spectrum). Let us now recall some relevant properties of the linear problem (6) (more details can be found e.g. in [2]). In a certain range of values of the parameter γ , a non-Hermitian \mathcal{PT} -symmetric operator $H = H(\gamma)$ may possess purely real spectrum. However, for a certain value (or values) of γ the system undergoes a spontaneous \mathcal{PT} symmetry breaking, corresponding to the transition from the real spectrum to complex one (in the latter situation the system is said to be in the phase of broken \mathcal{PT} symmetry). The transition to the broken \mathcal{PT} symmetry phase can be described in terms of exceptional-point spectral singularity [33, 34].

If an eigenvalue \tilde{b} of $H(\gamma)$ is real and possesses exactly one linearly independent eigenvector $\tilde{\mathbf{w}}$, then the latter can be chosen to be \mathcal{PT} -invariant [35], i.e.

$$\mathcal{PT}\tilde{\mathbf{w}} = \tilde{\mathbf{w}},\tag{7}$$

independently on whether the \mathcal{PT} symmetry of $H(\gamma)$ is unbroken or broken. Notice that condition (7) fixes the phase of the vector $\tilde{\mathbf{w}}$, while the model (1) is phase invariant thanks to the choice of the cubic nonlinearity of the form (4). Instead of vector $\tilde{\mathbf{w}}$ fixed by (7), one can consider any eigenvector of the form $e^{i\varphi}\tilde{\mathbf{w}}$, where φ is real. However, due to the phase invariance, this generalization does not lead to physically distinct solutions. Therefore, to simplify the algebra below we hold the definition of the \mathcal{PT} -invariant mode (7), bearing in mind that modes (either linear or nonlinear) \mathbf{w} and $e^{i\varphi}\mathbf{w}$ are physically equivalent.

Equation (7) trivially leads to the properties $\tilde{\mathbf{w}}^* = \mathcal{T}\tilde{\mathbf{w}} = \mathcal{P}\tilde{\mathbf{w}}$. Introducing the inner product as $\langle \mathbf{a}, \mathbf{b} \rangle = \mathbf{a}^{\dagger}\mathbf{b} = \sum_{j=1}^{N} a_j^* b_j$ we arrive at the conclusion that $\langle \tilde{\mathbf{w}}^*, \tilde{\mathbf{w}} \rangle$ is real. Indeed, for any two \mathcal{PT} -invariant vectors $\mathbf{a} = \mathcal{PT}\mathbf{a}$ and $\mathbf{b} = \mathcal{PT}\mathbf{b}$ one verifies that

$$\langle \mathbf{a}^*, \mathbf{b} \rangle = \langle \mathcal{P}\mathbf{a}, \mathbf{b} \rangle = \langle \mathcal{P}\mathbf{a}, \mathcal{P}\mathcal{T}\mathbf{b} \rangle = \langle \mathbf{a}, \mathcal{T}\mathbf{b} \rangle = \langle \mathbf{a}, \mathbf{b}^* \rangle = \langle \mathbf{a}^*, \mathbf{b} \rangle^*.$$
 (8)

In the situation when the eigenvalue \tilde{b} is multiple, one should distinguish between two different cases. In the first case, the multiple eigenvalue \tilde{b} (with algebraic multiplicity equal to n) possesses an invariant subspace spanned by n linearly independent eigenvectors. In other words, the eigenvalue \tilde{b} is *semi-simple*, and the Hamiltonian $H(\gamma)$ is diagonalizable (provided that all other eigenvalues are also simple or semi-simple). This situation arises, in particular, when two eigenvalues of a parameter-dependent Hamiltonian $H(\gamma)$ coalesce at some value of the control parameter γ but the two corresponding eigenvalues remain linearly independent. Physically such situation appears, in particular, when two (or more) identical systems are linearly coupled: the degeneracy occurs when the coupling becomes zero.

The second case corresponds to the situation when collision of the eigenvalues is accompanied by collisions of eigenvectors. Then the geometrical multiplicity of the eigenvalue \tilde{b} if less than the algebraic multiplicity, i.e. the dimension of the invariant subspace of the multiple eigenvalue \tilde{b} is less than n. This corresponds to so-called *exceptional* points [31]. As it was mentioned above, the relevance of such points is in particular related to the transition to the phase of broken \mathcal{PT} symmetry (see also the discussion in [36]).

3. Bifurcations of the families of nonlinear modes. Simple eigenvalue

We start by considering families of nonlinear modes bifurcating from a linear eigenstate of the \mathcal{PT} -symmetric Hamiltonian H. Let \tilde{b} be a simple real eigenvalue and $\tilde{\mathbf{w}}$ is the corresponding eigenvector solving linear problem (6). According to the discussion in Sec. 2, without loss of generality we can assume that $\tilde{\mathbf{w}}$ is \mathcal{PT} invariant, i.e. satisfies the condition (7).

In the vicinity of the *linear limit* the nonlinear modes bifurcating from the eigenstate $\tilde{\mathbf{w}}$ can be described using formal expansions [4]

$$\mathbf{w} = \varepsilon \tilde{\mathbf{w}} + \varepsilon^3 \mathbf{w}^{(3)} + o(\varepsilon^3) \quad \text{and} \quad b = \tilde{b} + \varepsilon^2 b^{(2)} + o(\varepsilon^2), \tag{9}$$

where ε is a real small parameter, $\varepsilon \ll 1$, and without loss of generality we impose the normalization condition $\langle \tilde{\mathbf{w}}, \tilde{\mathbf{w}} \rangle = 1$. Coefficients $\mathbf{w}^{(3)}$ and $b^{(2)}$ of the expansions are to be determined. Substituting expansions (9) into Eq. (5) and noticing from Eq. (4) that $F(\mathbf{w}) = \varepsilon^2 F(\tilde{\mathbf{w}}) + O(\varepsilon^3)$, one arrives at the following equation for the shift of the eigenvalue $b^{(2)}$:

$$b^{(2)}\tilde{\mathbf{w}} = (H - \tilde{b})\mathbf{w}^{(3)} + F(\tilde{\mathbf{w}})\tilde{\mathbf{w}}.$$
(10)

Multiplying this equation by $\tilde{\mathbf{w}}^*$ and using Eqs. (2) and (6), one readily obtains

$$b^{(2)} = \frac{\langle \tilde{\mathbf{w}}^*, F(\tilde{\mathbf{w}}) \tilde{\mathbf{w}} \rangle}{\langle \tilde{\mathbf{w}}^*, \tilde{\mathbf{w}} \rangle}.$$
(11)

Taking into account reality of the product $\langle \tilde{\mathbf{w}}^*, \tilde{\mathbf{w}} \rangle$ [see Eqs. (8)], one notices that bifurcation of a family nonlinear modes is possible only if

$$\operatorname{Im}\left\langle \tilde{\mathbf{w}}^{*}, F(\tilde{\mathbf{w}})\tilde{\mathbf{w}} \right\rangle = 0, \tag{12}$$

which is necessary for $b^{(2)}$ to be real. Thus (12) is a necessary condition for a family of nonlinear modes to bifurcate from the nondegenerate eigenstate of H corresponding to \tilde{b} .

By analogy with the condition (7) which guarantees reality of the denominator of the right hand side of Eq. (11), in order to ensure that the numerator is also real, we introduce the following condition for the nonlinear operator $F(\mathbf{w})$ [4]:

$$\mathcal{PTF}(\mathbf{w})\mathbf{w} = F(\mathbf{w})\mathbf{w}$$
 for all \mathbf{w} such that $\mathcal{PTw} = \mathbf{w}$. (13)

Obviously, this condition is equivalent to $[\mathcal{PT}, F(\mathbf{w})] = 0$, where the latter commutator is only considered on the set of vectors satisfying $\mathcal{PT}\mathbf{w} = \mathbf{w}$. In what follows the nonlinearities obeying (13) will be said to be *weakly* \mathcal{PT} symmetric. The set of weakly \mathcal{PT} -symmetric nonlinearities corresponding to the given parity operator \mathcal{P} will be denoted by $NL_{w\mathcal{PT}}(\mathcal{P})$. Thus the weak \mathcal{PT} symmetry of the nonlinear operator $F(\mathbf{w})$ appears to be a necessary condition for the system (1) to admit continuous families of nonlinear modes.

The choice of the term weak \mathcal{PT} symmetry can be understood if one considers a more restrictive condition for the nonlinear operator $F(\mathbf{w})$ to commute with the \mathcal{PT} operator for any vector \mathbf{w} :

$$[\mathcal{PT}, F(\mathbf{w})] = 0. \tag{14}$$

Nonlinear operators $F(\mathbf{w})$ of the latter type will be said to be \mathcal{PT} symmetric. The set of \mathcal{PT} -symmetric nonlinearities corresponding to the given parity operator \mathcal{P} will be denoted as $NL_{\mathcal{PT}}(\mathcal{P})$. Obviously, for a given parity operator \mathcal{P} , the class of weakly \mathcal{PT} -symmetric nonlinearities contains the class of \mathcal{PT} symmetric nonlinearities as a subset: $NL_{\mathcal{PT}}(\mathcal{P}) \subset NL_{w\mathcal{PT}}(\mathcal{P})$. The relevance of the $NL_{\mathcal{PT}}$ -type nonlinearities will be discussed in Sec. 6.

As an example, let us consider a finite nonlinear system with N = 4 (i.e. a quadrimer) and a Kerr nonlinearity, $F_K(\mathbf{w})$, whose elements are defined by $F_{K,pj}(\mathbf{w}) = \delta_{pj}|w_j|^2$ where δ_{pj} is the Kronecker delta, and p, j = 1, ..., 4. This nonlinearity satisfies the requirement (13) provided that the parity operator is chosen as

$$\mathcal{P}_{11} = \sigma_1 \otimes \sigma_1 = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}$$
(15)

(hereafter σ_j are the Pauli matrices). However, the nonlinearity $F_K(\mathbf{w})$ does not satisfy the condition (14). Therefore $F_K \in NL_{w\mathcal{PT}}(\mathcal{P}_{11}) \setminus NL_{\mathcal{PT}}(\mathcal{P}_{11})$.

Nonlinear lattices with \mathcal{PT} -symmetric linear part

Notice that expansions (9) suggest that the nonlinear modes \mathbf{w} bifurcating from the \mathcal{PT} -invariant eigenvector $\tilde{\mathbf{w}}$ will also be \mathcal{PT} invariant: $\mathcal{PT}\mathbf{w} = \mathbf{w}$. This fact is confirmed in numerous previous studies, see e.g. [4, 6, 7, 9, 11, 13, 27], where a large library of results dedicated to nonlinear modes bifurcating from the non-degenerate (simple) linear eigenstates and supported by the nonlinearities of the $NL_{w\mathcal{PT}}$ type can be found. Existence of the families of nonlinear modes was also established through the analytical continuation from the the anticontinuum limit [11] (the technique well known in the theory of conservative lattices [37]). Bifurcations of the nonlinear modes from the simple eigenvalues of the underlying linear lattices, as well as from the anticontinuum limit, were recently addressed in a rigorous mathematical framework [6].

4. Bifurcation from the semi-simple eigenvalue

4.1. Structure of the invariant subspace

In the previous section we have considered bifurcations of nonlinear modes from a nondegenerate (simple) eigenvalue of the underlying linear problem. Now we turn to the case of a semi-simple eigenvalue, i.e. a multiple eigenvalue whose geometric and algebraic multiplicities are equal. Let \tilde{b} be a real semi-simple eigenvalue of multiplicity equal to n. This means that there exist n linearly independent eigenvectors $\tilde{\mathbf{w}}_j$, $j = 1, \ldots, n$, such that $H\tilde{\mathbf{w}}_j = \tilde{b}\tilde{\mathbf{w}}_j$, and there is a n-dimensional H-invariant subspace spanned by $\tilde{\mathbf{w}}_j$ such that any vector from this subspace is an eigenvector of the operator H.

Adopting the approach of formal expansions used in Sec. 3 for the case of a simple eigenvalue, we are going to construct a family of nonlinear modes which in the vicinity of the bifurcation from the linear limit behave as $\mathbf{w} = \varepsilon \tilde{\mathbf{w}} + \ldots$, where $\tilde{\mathbf{w}}$ is some eigenvector of H corresponding to the eigenvalue \tilde{b} . Looking for \mathcal{PT} -invariant nonlinear modes $\mathcal{PT}\mathbf{w} = \mathbf{w}$, one has to find a \mathcal{PT} -invariant eigenvector $\tilde{\mathbf{w}}$ for the expansion to be valid. However, while in the case of a simple eigenvalue a \mathcal{PT} -invariant eigenvector $\tilde{\mathbf{w}}$ necessarily exists (and is unique up to irrelevant multiplier), in the case of a semi-simple eigenvalue \tilde{b} it is not readily obvious how many \mathcal{PT} -invariant eigenvectors exist (if any). Therefore, in the situation at hand it is necessary to explore structure of the invariant subspace associated with the multiple eigenvalue \tilde{b} . Let us show that in the invariant subspace of \tilde{b} one can always find a basis of n linearly independent \mathcal{PT} -invariant eigenvectors (independently on whether the \mathcal{PT} symmetry of H is broken or not).

Proposition 1. Let \hat{b} be a real semi-simple eigenvalue of multiplicity n. Then the invariant subspace of \tilde{b} has a complete basis $(\mathbf{u}_1, \mathbf{u}_2, \ldots, \mathbf{u}_n)$ of \mathcal{PT} -invariant eigenvectors: $\mathcal{PT}\mathbf{u}_j = \mathbf{u}_j, j = 1, 2, \ldots, n$.

Proof of Proposition 1. The condition of the proposition implies that there exist n linearly independent eigenvectors $\tilde{\mathbf{w}}_j$, j = 1, 2, ..., n, such that $H\tilde{\mathbf{w}}_j = \tilde{b}\tilde{\mathbf{w}}_j$. If the choice of the linearly independent eigenvectors $\tilde{\mathbf{w}}_j$ is arbitrary, then generally speaking $\mathcal{PT}\tilde{\mathbf{w}}_j \neq \tilde{\mathbf{w}}_j$, i.e. eigenvectors $\tilde{\mathbf{w}}_j$ (or some of them) are not \mathcal{PT} invariant. Thus, in order to prove the proposition we must find n linearly independent \mathcal{PT} -invariant eigenvectors.

To this end we apply \mathcal{PT} operator to each $\tilde{\mathbf{w}}_j$ and obtain a new set of vectors $\mathbf{v}_j = \mathcal{PT}\tilde{\mathbf{w}}_j, j = 1, \dots, n$. \mathcal{PT} symmetry of H and reality of \tilde{b} imply that $H\mathbf{v}_j = \tilde{b}\mathbf{v}_j$,

i.e. each \mathbf{v}_j belongs to the invariant subspace of the eigenvalue b. Notice also that $\mathcal{PT}\mathbf{v}_j = \mathbf{w}_j$ [thanks to $(\mathcal{PT})^2 = I$]. Then linear independence of vectors $\tilde{\mathbf{w}}_j$ implies that the vectors \mathbf{v}_j are also linearly independent and therefore constitute a basis in the invariant subspace of \tilde{b} . Therefore, there exists a $n \times n$ nonsingular matrix D such that $\mathbf{v}_j = \sum_{k=1}^n D_{jk} \tilde{\mathbf{w}}_k$.

Let us now introduce a new set of vectors \mathbf{u}_i , j = 1, 2, ..., n, given as

$$\mathbf{u}_j = e^{i\phi}\tilde{\mathbf{w}}_j + e^{-i\phi}\mathbf{v}_j = e^{-i\phi}\left(\sum_{k=0}^n D_{jk}\tilde{\mathbf{w}}_k + e^{2i\phi}\tilde{\mathbf{w}}_j\right) = e^{-i\phi}\sum_{k=0}^n M_{jk}\tilde{\mathbf{w}}_k,$$

where ϕ is an arbitrary (so far) real parameter, and M_{jk} are the entries of the $n \times n$ matrix M defined as $M = D + e^{2i\phi}I$ (here I is the $n \times n$ identity matrix). Then each \mathbf{u}_j belongs to the invariant subspace of \tilde{b} and is obviously \mathcal{PT} invariant. Besides, one can always find such ϕ that matrix M is nonsingular (it is necessary and sufficient to choose ϕ such that $\rho = -e^{2i\phi}$ does not belong to the spectrum of D). Once M is nonsingular, the eigenvectors \mathbf{u}_j are linearly independent and therefore constitute a complete \mathcal{PT} -invariant basis in the invariant subspace associated with the eigenvalue \tilde{b} . Thus the proposition is proven.

4.2. Expansions for nonlinear modes

Turning now to bifurcations of nonlinear modes, we employ the same expansions as in the case of a simple eigenvalue:

$$\mathbf{w} = \varepsilon \tilde{\mathbf{w}} + \varepsilon^3 \mathbf{w}^{(3)} + o(\varepsilon^3), \quad \text{and} \quad b = \tilde{b} + \varepsilon^2 b^{(2)} + o(\varepsilon^2). \tag{16}$$

However, now the vector $\tilde{\mathbf{w}}$ is a linear combination of the \mathcal{PT} -invariant eigenvectors \mathbf{u}_j constructed in Proposition 1: $\tilde{\mathbf{w}} = \sum_{j=1}^n c_j \mathbf{u}_j$. In order to assure that the linear solution $\tilde{\mathbf{w}}$ is \mathcal{PT} -invariant, i.e. satisfies (7), we require all the coefficients c_j to be real. However, an arbitrary set of coefficients c_j generally speaking does not represent a linear mode $\tilde{\mathbf{w}}$ allowing for a bifurcation of a family of nonlinear modes \mathbf{w} . The relevant relations among the coefficients can be found by multiplying Eq. (10) by \mathbf{u}_j^* leading to conditions as follows

for each
$$j = 1, 2, \dots n$$
: $b^{(2)} = \frac{\langle \mathbf{u}_j^*, F(\tilde{\mathbf{w}})\tilde{\mathbf{w}} \rangle}{\langle \mathbf{u}_j^*, \tilde{\mathbf{w}} \rangle},$ (17)

where $b^{(2)}$ is required to be real. One can readily ensure that if the nonlinearity $F(\mathbf{w})$ satisfies to requirement (13), i.e. $F(\mathbf{w}) \in NL_{w\mathcal{PT}}(\mathcal{P})$, then $b^{(2)}$ is real. Indeed:

$$\frac{\langle \mathbf{u}_{j}^{*}, F(\tilde{\mathbf{w}})\tilde{\mathbf{w}}\rangle^{*}}{\langle \mathbf{u}_{j}^{*}, \tilde{\mathbf{w}}\rangle^{*}} = \frac{\langle \mathcal{PT}\mathbf{u}_{j}, \mathcal{T}(F(\tilde{\mathbf{w}})\tilde{\mathbf{w}})\rangle}{\langle \mathcal{PT}\mathbf{u}_{j}, \mathcal{T}\tilde{\mathbf{w}}\rangle} = \frac{\langle \mathbf{u}_{j}^{*}, F(\tilde{\mathbf{w}})\tilde{\mathbf{w}}\rangle}{\langle \mathbf{u}_{j}^{*}, \tilde{\mathbf{w}}\rangle}$$

Bearing in mind the normalization condition $\langle \tilde{\mathbf{w}}, \tilde{\mathbf{w}} \rangle = 1$, the system (17) can be viewed as *n* algebraic equations with respect to n-1 independent real coefficients c_j ; the requirement of compatibility of these equation determines $b^{(2)}$. The existence and a number of the solutions obviously depends on the specific form of the nonlinearity $F(\mathbf{w})$, the latter determining the diversity of physically distinct families of nonlinear modes bifurcating from the eigenstates of the linear spectrum.

Finally, we mention a peculiar but physically relevant situation when for a certain set of the coefficients c_j the eigenstate $\tilde{\mathbf{w}}$ becomes an eigenvector of the nonlinear eigenvalue problem $F(\tilde{\mathbf{w}})\tilde{\mathbf{w}} = \lambda \tilde{\mathbf{w}}$, and the eigenvalue λ is real. Then Eqs. (17) are automatically satisfied provided that $b^{(2)} = \lambda$, i.e. $b^{(2)} = \langle \tilde{\mathbf{w}}, F(\tilde{\mathbf{w}})\tilde{\mathbf{w}} \rangle$. In this case the number of free parameters exceeds the number of the imposed constraints. Such a situation has been encountered for example in the model of a \mathcal{PT} -symmetric birefringent coupler [5] and occurs in the example considered in the next subsection.

4.3. Example: \mathcal{PT} -symmetric coupler of bi-chromatic light

As we already mentioned, the presence of multiply degenerated eigenvalues in the linear spectrum typically occurs when two identical systems are decoupled linearly, and coupling exists only due to the nonlinearity. Let us consider the following example:

$$i\dot{q}_1 = i\gamma q_1 + q_2 + (|q_1|^2 + |q_3|^2)q_1, \tag{18a}$$

$$i\dot{q}_2 = -i\gamma q_2 + q_1 + \varkappa (|q_2|^2 + |q_4|^2)q_2, \tag{18b}$$

$$i\dot{q}_3 = -i\gamma q_3 + q_4 + (|q_3|^2 + |q_1|^2)q_3, \tag{18c}$$

$$i\dot{q}_4 = i\gamma q_4 + q_3 + \varkappa (|q_4|^2 + |q_2|^2)q_4.$$
(18d)

It can be viewed as a model for propagation of a bi-chromatic light in a \mathcal{PT} -symmetric coupler. Then q_j and q_{j+2} (j = 1, 2) are the fields having the propagation constants k_1 and k_2 in *j*-th waveguide, where self-phase and cross-phase modulation of the two modes propagating in one arm are normalized to one. The modes k_1 and k_2 are subject to gain and absorption in the first waveguide (and *vice versa* in the second waveguide). The real parameter \varkappa describes the difference in the Kerr nonlinearities in the waveguides.

The linear Hamiltonian of (18a)-(18d) is described by the matrix $H_1(\gamma, -\gamma)$ where

$$H_{bc}(\gamma_1, \gamma_2) = \begin{pmatrix} i\gamma_1 & 1 & 0 & 0\\ 1 & i\gamma_2 & 0 & 0\\ 0 & 0 & -i\gamma_1 & 1\\ 0 & 0 & 1 & -i\gamma_2 \end{pmatrix}$$
(19)

and the nonlinearity is given by

$$F_{bc}(\mathbf{q}) = \begin{pmatrix} |q_1|^2 + |q_3|^2 & 0 & 0 & 0\\ 0 & \varkappa(|q_2|^2 + |q_4|^2) & 0 & 0\\ 0 & 0 & |q_1|^2 + |q_3|^2 & 0\\ 0 & 0 & 0 & \varkappa(|q_2|^2 + |q_4|^2) \end{pmatrix}.$$
(20)

Subscripts "bc" in Eqs. (19) and (20) stand for "bi-chromatic".

One readily verifies that $H_{bc}(\gamma_1, \gamma_2)$ is $\mathcal{P}_{10}\mathcal{T}$ -symmetric with respect to

$$\mathcal{P}_{10} = \sigma_1 \otimes \sigma_0 = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}.$$
 (21)

In the particular case $\gamma_1 = -\gamma_2 = \gamma$ an additional symmetry appears. Indeed, $H_{bc}(\gamma, -\gamma)$ is also $\mathcal{P}_{01}\mathcal{T}$ -symmetric with

$$\mathcal{P}_{01} = \sigma_0 \otimes \sigma_1 = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix}.$$




Figure 1. The graphs of the Hamiltonians $H_{bc}(\gamma_1, \gamma_2)$ (left graph) and $H_{bc}(\gamma, -\gamma)$ (right graph) and the symmetries corresponding to the respective parity operators.

The physical meaning of the operators \mathcal{P}_{10} and \mathcal{P}_{01} becomes particularly clear in terms of the graph representation [4], as this is illustrated in Fig. 1.

One also verifies that $F_{bc}(\mathbf{q}) \in NL_{\mathcal{PT}}(\mathcal{P}_{10}) \subset NL_{w\mathcal{PT}}(\mathcal{P}_{10})$. At $\varkappa = 1$ one additionally has $F_{bc}(\mathbf{q}) \in NL_{w\mathcal{PT}}(\mathcal{P}_{01})$.

As it is clear, the operator $H_{bc}(\gamma, -\gamma)$ represents two uncoupled linear \mathcal{PT} symmetric dimers. Spectrum of $H_{bc}(\gamma, -\gamma)$ consists of two double semi-simple
eigenvalues given by $\tilde{b}_{\pm} = \pm \sqrt{1 - \gamma^2}$. The $\mathcal{P}_{10}\mathcal{T}$ -invariant eigenvectors of $H_{bc}(\gamma, -\gamma)$ (which exist due to Proposition 1) can be written does as follows:

$$\begin{split} \mathbf{u}_{+,1} &= \begin{pmatrix} e^{i\varphi} \\ e^{-i\varphi} \\ e^{-i\varphi} \\ e^{i\varphi} \end{pmatrix}, \quad \mathbf{u}_{+,2} &= i \begin{pmatrix} e^{i\varphi} \\ e^{-i\varphi} \\ -e^{-i\varphi} \\ -e^{i\varphi} \\ e^{i\varphi} \\ e^{-i\varphi} \end{pmatrix}, \\ \mathbf{u}_{-,1} &= \begin{pmatrix} -e^{-i\varphi} \\ e^{i\varphi} \\ e^{i\varphi} \\ e^{-i\varphi} \\ e^{-i\varphi} \end{pmatrix}, \quad \mathbf{u}_{-,2} &= i \begin{pmatrix} -e^{-i\varphi} \\ e^{i\varphi} \\ e^{i\varphi} \\ -e^{-i\varphi} \end{pmatrix}, \end{split}$$

where φ is defined by requirements $\sin(2\varphi) = \gamma$ and $\cos(2\varphi) = \sqrt{1 - \gamma^2}$.

Let us search for nonlinear modes bifurcating from the eigenvalue \tilde{b}_+ (analysis for the eigenvalue \tilde{b}_- yields similar results). Following to the above approach, we search for the linear eigenvector in the form as a linear combination $\tilde{\mathbf{w}} = c_1 \mathbf{u}_{+,1} + c_2 \mathbf{u}_{+,2}$ where $c_{1,2}$ are real coefficients. Subject to the normalization $\langle \tilde{\mathbf{w}}, \tilde{\mathbf{w}} \rangle = 1$, we obtain $c_1^2 + c_2^2 = 1/4$. Straightforward algebra gives

$$\langle \mathbf{u}_{+,1}^*, F_{bc}(\tilde{\mathbf{w}})\tilde{\mathbf{w}} \rangle = c_1 \cos(2\varphi)(\varkappa + 1) + c_2 \sin(2\varphi)(\varkappa - 1), \qquad (22a)$$

$$\langle \mathbf{u}_{+,2}^*, F_{bc}(\tilde{\mathbf{w}})\tilde{\mathbf{w}} \rangle = c_1 \sin(2\varphi)(\varkappa - 1) - c_2 \cos(2\varphi)(\varkappa + 1), \qquad (22b)$$

$$\langle \mathbf{u}_{+,1}^*, \tilde{\mathbf{w}} \rangle = 4c_1 \cos(2\varphi), \quad \langle \mathbf{u}_{+,2}^*, \tilde{\mathbf{w}} \rangle = -4c_2 \cos(2\varphi).$$
 (22*c*)

Next, following Eqs. (17), we require

$$b^{(2)}\langle \mathbf{u}_{+,1}^*, \tilde{\mathbf{w}} \rangle = \langle \mathbf{u}_{+,1}^*, F_{bc}(\tilde{\mathbf{w}})\tilde{\mathbf{w}} \rangle, \qquad (23a)$$

$$b^{(2)}\langle \mathbf{u}_{+,2}^*, \tilde{\mathbf{w}} \rangle = \langle \mathbf{u}_{+,2}^*, F_{bc}(\tilde{\mathbf{w}})\tilde{\mathbf{w}} \rangle.$$
(23b)

Considering first the generic case $\varkappa \neq 1$, we substitute Eqs. (22a)-(22c) into Eqs. (23a)-(23b) and find that Eqs. (23a) and (23b) are compatible only if $\sin(4\varphi) = 0$, which means that bifurcations of nonlinear modes are possible only for $\gamma = 0$ (i.e. when the dissipation vanishes) or for $\gamma = \pm 1$. The latter case corresponds to the point of the phase transition to the broken \mathcal{PT} symmetry, which is described by the exceptional point singularity and requires a particular analysis (see Sec. 5).

The case $\varkappa = 1$ corresponds to the peculiar situation when for any choice of c_1 and c_2 one has $F_{bc}(\tilde{\mathbf{w}})\tilde{\mathbf{w}} = \tilde{\mathbf{w}}/2$, i.e. Eqs. (23*a*)–(23*b*) are automatically satisfied with $b^{(2)} = 1/2$. This is however a strongly degenerate case. Looking for solutions in the form $w_3^* = \chi w_1$, $w_4^* = \chi w_2$ (χ is arbitrary real) the system is reduced to the \mathcal{PT} -symmetric nonlinear dimer [3]

$$bw_1 = i\gamma w_1 + w_2 + (\chi^2 + 1)|w_1|^2 w_1,$$

$$bw_2 = -i\gamma w_2 + w_1 + (\chi^2 + 1)|w_2|^2 w_2,$$

which supports analytically computable families of nonlinear modes given by the substitution $w_1 = w_2^*$. Since χ is arbitrary, we can obtain a continuous set of solutions even for fixed γ and b.

5. Bifurcations of nonlinear modes in the presence of exceptional-point singularity

5.1. Expansions for nonlinear modes

Let us turn to a situation when a multiple eigenvalue b corresponds to the so-called exceptional point singularity which appears when coalescence of two (or more) simple eigenvalues of the Hamiltonian $H(\gamma)$ occurring at specific values of the parameter γ is accompanied by collision of the corresponding eigenvectors. Then the geometric multiplicity of the eigenvalue \tilde{b} is less than the algebraic multiplicity, and the Hamiltonian $H(\gamma)$ becomes nondiagonalizable. Presence of exceptional points is a typical feature of \mathcal{PT} -symmetric systems, since such points naturally appear at the "boundary" between phases of broken and unbroken \mathcal{PT} symmetries [33, 34]. Nonlinear behavior of \mathcal{PT} -symmetric optical lattices near the phase-transition point was recently considered in [38, 39].

We illustrate our ideas considering an exceptional point where two linear eigenstates coalesce, forming a real multiple eigenvalue \tilde{b} with total multiplicity equal to two and having exactly one linearly independent eigenvector $\tilde{\mathbf{w}}$. We thus have

$$\left(H - \tilde{b}\right)\tilde{\mathbf{w}} = 0, \quad \left(H - \tilde{b}\right)\mathbf{v} = \tilde{\mathbf{w}}.$$
 (24)

where we have introduced a generalized eigenvector \mathbf{v} . Since H is \mathcal{PT} symmetric and $\tilde{\mathbf{w}}$ is the only linearly independent eigenvector corresponding to \tilde{b} , we can assume that $\mathcal{PT}\tilde{\mathbf{w}} = \tilde{\mathbf{w}}$. We can also assume the the generalized eigenvector \mathbf{v} is also \mathcal{PT} invariant, i.e. $\mathcal{PT}\mathbf{v} = \mathbf{v}$. This assumption is valid thanks to the following proposition.

Proposition 2. There exists a \mathcal{PT} -invariant generalized eigenvector \mathbf{v} : $\mathcal{PT}\mathbf{v} = \mathbf{v}$.

Proof of Proposition 2. Let \mathbf{u} is an arbitrarily chosen generalized eigenvector, i.e. $(H - \tilde{b})\mathbf{u} = \tilde{\mathbf{w}}$. Applying \mathcal{PT} operator to the both sides of the latter equality and using that operator $H - \tilde{b}$ commutes with \mathcal{PT} and that $\mathcal{PT}\tilde{\mathbf{w}} = \tilde{\mathbf{w}}$, we find that $\mathcal{PT}\mathbf{u}$ is also a generalized eigenvector. Then the \mathcal{PT} -invariant generalized eigenvector \mathbf{v} can be found as $\mathbf{v} = \frac{1}{2}(\mathbf{u} + \mathcal{PT}\mathbf{u})$.

Multiplying the second of equations (24) by $\mathcal{P}\tilde{\mathbf{w}} = \tilde{\mathbf{w}}^*$ and using that $(H - \tilde{b})^{\dagger} = \mathcal{P}(H - \tilde{b})\mathcal{P}$, we find out that the eigenvector $\tilde{\mathbf{w}}$ is self-orthogonal [40] in the sense of the indefinite \mathcal{PT} inner product [35], i.e. $\langle \mathcal{P}\tilde{\mathbf{w}}, \tilde{\mathbf{w}} \rangle = 0$. Therefore, now Eq. (11) as well as the expansions (9) used in Sec. 3 and Sec. 4 do not work and have to be modified. To this end, we look for modified small-amplitude expansions in the form

$$\mathbf{w} = \varepsilon \tilde{\mathbf{w}} + \varepsilon^2 \mathbf{w}^{(2)} + \varepsilon^3 \mathbf{w}^{(3)} + \cdots,$$
(25)

$$b = \tilde{b} + \varepsilon b^{(1)} + \varepsilon^2 b^{(2)} + \cdots$$
(26)

Substituting Eqs. (25)–(26) in (5) and using (24), in the ϵ^2 -order we obtain $\mathbf{w}^{(2)} = b^{(1)}\mathbf{v}$. The third order equation reads

$$b^{(2)}\tilde{\mathbf{w}} + \left(b^{(1)}\right)^2 \mathbf{v} = (H - \tilde{b})\mathbf{w}^{(3)} + F(\tilde{\mathbf{w}})\tilde{\mathbf{w}}.$$
(27)

Multiplying this equation from the left by $\tilde{\mathbf{w}}^*$, we obtain

$$\left(b^{(1)}\right)^2 = \frac{\langle \tilde{\mathbf{w}}^*, F(\tilde{\mathbf{w}})\tilde{\mathbf{w}} \rangle}{\langle \tilde{\mathbf{w}}^*, \mathbf{v} \rangle}.$$
(28)

Let us now assume that the nonlinear operator $F(\mathbf{w})$ is of the NL_{wPT} type [see (13)]. Then, due to (8), we conclude that the right hand side of (28) is real. Unlike the case of simple and semi-simple eigenvalues, however, for the condition (28) to have sense one has to require its r.h.s. to be nonnegative. This additional condition, which does not appear in the case of the ordinary points, is indeed restrictive, as we will illustrate in Sec. 5.2.3. Another important feature is that (28) indicates that families bifurcating from the eigenvalue \tilde{b} appear in *pairs* (corresponding to two opposite signs of $b^{(1)}$).

5.2. Example: Quadrimer with the nearest-neighbor interactions

5.2.1. "Phase diagram" Let us illustrate the above ideas on the example of a quadrimer with nearest-neighbor interactions. The linear part of (1) is now given by the Hamiltonian

$$H_{nn}(\gamma_1, \gamma_2) = \begin{pmatrix} i\gamma_1 & 1 & 0 & 0\\ 1 & i\gamma_2 & 1 & 0\\ 0 & 1 & -i\gamma_2 & 1\\ 0 & 0 & 1 & -i\gamma_1 \end{pmatrix},$$
(29)

where the subscript "nn" stays for "nearest-neibor". The linear operator $H_{nn}(\gamma_1, \gamma_2)$ is $\mathcal{P}_{11}\mathcal{T}$ -symmetric with $\mathcal{P}_{11} = \sigma_1 \otimes \sigma_1$ [see Eq. (15)], and the nonlinear part is given by the operator $F_K(\mathbf{w}) = \text{diag}(|w_1|^2, |w_2|^2, |w_3|^2, |w_4|^2)$ (where subscript K stands for Kerr nonlinearity).

Properties of the linear operator $H_{nn}(\gamma_1, \gamma_2)$ can be visualized conveniently by means of the "phase-diagram" [4] shown in Fig. 2, panel (PD). Depending on γ_1 and γ_2 , the phase diagram features three domains: (i) unbroken or *exact* \mathcal{PT} symmetry, when all the eigenvalues \tilde{b}_i , j = 1, 2, 3, 4, of $H_{nn}(\gamma_1, \gamma_2)$ are real; (ii) broken \mathcal{PT} symmetry with two real and two complex conjugated eigenvalues (notice that in the particular case $\gamma_1 = \gamma_2$ domain (ii) can not be encountered); (iii) broken \mathcal{PT} symmetry with all \tilde{b}_j complex. Varying parameters $\gamma_{1,2}$, one can "travel" across the phase diagram visiting domains with different phases. An interesting feature of the phase diagram shown in Fig. 2 (PD) is that in some cases increase of the total dissipation brings the system from the phase of broken \mathcal{PT} symmetry to the unbroken \mathcal{PT} symmetry. For example, fixing value of one of the coefficients as $\gamma_1 = 1.2$, than observes that for small γ_2 (say $\gamma_2 = 0$) the \mathcal{PT} symmetry is broken, but when γ_2 becomes sufficiently large, then the system enters the phase of the unbroken \mathcal{PT} symmetry. Further increase of γ_2 leads to another phase transition and the \mathcal{PT} symmetry becomes broken again. A similar scenario with two \mathcal{PT} transitions was recently reported in [41].

Boundaries separating different domains of the phase diagram correspond to the exceptional points. In particular, the boundaries contain exactly four triple points T_j , $j = 1, \ldots, 4$, where the three domains touch. The triple points correspond to values $\gamma_{1,2}$ for which $\tilde{b}_{1,\ldots,4} = 0$, and the canonical form of $H_{nn}(\gamma_1, \gamma_2)$ consists of the only 4×4 Jordan block. Depending on how $\gamma_{1,2}$ change in the vicinity of T_j , either the \mathcal{PT} -symmetric phase or one of the \mathcal{PT} symmetry broken phases arise. All the other points of the boundaries are the double points separating two different phases. They are characterized by the presence of one or two 2×2 Jordan blocks in the canonical form of $H_{nn}(\gamma_1, \gamma_2)$. The double points can be further sub-classified as belonging to boundaries separating either phases (i) and (ii), or phases (i) and (iii), or phases (ii) and (iii). Here however we do not intend to perform a complete classification and consider only the double points are likely to be more probable "candidates" to give birth to families of stable nonlinear modes [comparing to the double points which are not adjacent to the phase (i)].

5.2.2. Existence of nonlinear modes The nonlinear modes that obey $\mathcal{P}_{11}\mathcal{T}\mathbf{w} = \mathbf{w}$ have the following property: $w_1 = w_4^*$, $w_2 = w_3^*$. It is a simple exercise to ensure that the nonlinear part of the system $F_K(\mathbf{w})$ is of the $NL_{w\mathcal{PT}}(\mathcal{P}_{11})$ type, i.e. obeys property (13), see also Sec. 3 for discussion of the properties of $F_K(\mathbf{w})$. It was shown [4] that in this case the nonlinear modes can be found as roots of an eight-degree polynomial whose coefficients depend on b. The nonlinear modes constitute continuous families, which can be visualized as dependencies U vs. b, where $U = \frac{1}{4} \sum_{j=1}^{4} |w_j|^2$ can be associated with the norm of the solution (or with the total energy flow in the optical context).

5.2.3. Double points between T_1 and T_2 (between T_3 and T_4) Let us now consider bifurcations of nonlinear modes from the double points belonging to the boundary separating phases (i) and (iii). In the panel (PD) of Fig. 2 such points are situated on the boundary between the points T_1 and T_2 , e.g. point D_1 (or on the boundary between T_3 and T_4 which is considered analogously). The values of $\gamma_{1,2}$ corresponding to such points are given by the equation

$$(\gamma_1^2 - \gamma_2^2)^2 - 2(\gamma_1^2 + 4\gamma_1\gamma_2 + 3\gamma_2^2) + 5 = 0.$$
(30)

In this case the spectrum of the Hamiltonian $H_{nn}(\gamma_1, \gamma_2)$ consists of two opposite real double eigenvalues \tilde{b}_{\pm} , $\tilde{b}_{\pm} = -\tilde{b}_{-}$. Both \tilde{b}_{\pm} and \tilde{b}_{-} correspond to a 2×2 Jordan block. Let $\tilde{\mathbf{w}}_{\pm}$ and \mathbf{v}_{\pm} be the eigenvector and the generalized eigenvector corresponding to




Figure 2. Panel (PD): "Phase diagram" of the Hamiltonian $H_{nn}(\gamma_1, \gamma_2)$ defined by Eq. (29). The dark-grey diamond-shaped domain corresponds to unbroken \mathcal{PT} symmetry; in the light-gray domains there are two real and two complex eigenvalues; in the white domains all eigenvalues are complex. Other panels: families of nonlinear modes on the plane U vs. b for double points $D_{1,2,3}$, and for triple points T_1 and T_2 . Stable (unstable) modes are shown by solid blue (dotted red) lines.

the eigenvalue \tilde{b}_+ (respectively, $\tilde{\mathbf{w}}_-$ and \mathbf{v}_- correspond to \tilde{b}_-). Then we can write down two Jordan chains:

$$(H_{nn} - \tilde{b}_{\pm})\tilde{\mathbf{w}}_{\pm} = 0, \quad (H_{nn} - \tilde{b}_{\pm})\mathbf{v}_{\pm} = \tilde{\mathbf{w}}_{\pm}.$$
(31)

One can easily check that if one chooses $\mathcal{P}_{11}\mathcal{T}\tilde{\mathbf{w}}_{\pm} = \tilde{\mathbf{w}}_{\pm}$, then the eigenvectors $\tilde{\mathbf{w}}_{+}$ and $\tilde{\mathbf{w}}_{-}$ can be also chosen to be related by the following relation: $\tilde{\mathbf{w}}_{+} = G\mathcal{P}_{11}\tilde{\mathbf{w}}_{-}$, where G = diag(-i, i, -i, i). Therefore, the generalized eigenvectors are related through the relation $\mathbf{v}_{+} = \mathcal{P}_{11}G\mathbf{v}_{-}$ [this can be checked by direct substitution to Eqs. (31) and using the relations $H_{nn}\mathcal{P}_{11}G = G\mathcal{P}_{11}H_{nn}$, $G^2 = -I$, $(\mathcal{P}_{11}G)^2 = I$]. Then from Eq. (28) one obtains

$$\begin{pmatrix} b_{+}^{(1)} \end{pmatrix}^{2} = \frac{\langle \tilde{\mathbf{w}}_{+}^{*}, F_{K}(\tilde{\mathbf{w}}_{+})\tilde{\mathbf{w}}_{+} \rangle}{\langle \tilde{\mathbf{w}}_{+}^{*}, \mathbf{v}_{+} \rangle} = \frac{\langle -G\mathcal{P}_{11}\tilde{\mathbf{w}}_{-}^{*}, F_{K}(\tilde{\mathbf{w}}_{-})G\mathcal{P}_{11}\tilde{\mathbf{w}}_{-} \rangle}{\langle -G\mathcal{P}_{11}\tilde{\mathbf{w}}_{-}^{*}, \mathcal{P}_{11}G\mathbf{v}_{-} \rangle} = \frac{\langle \tilde{\mathbf{w}}_{-}^{*}, F_{K}(\tilde{\mathbf{w}}_{-})\tilde{\mathbf{w}}_{-} \rangle}{\langle \mathcal{P}_{11}GG\mathcal{P}_{11}\tilde{\mathbf{w}}_{-}^{*}, \mathbf{v}_{-} \rangle}$$
$$= -\frac{\langle \tilde{\mathbf{w}}_{-}^{*}, F_{K}(\tilde{\mathbf{w}}_{-})\tilde{\mathbf{w}}_{-} \rangle}{\langle \tilde{\mathbf{w}}_{-}^{*}, \mathbf{v}_{-} \rangle} = -\left(b_{-}^{(1)}\right)^{2}.$$

Here we have additionally used that $F_K(\tilde{\mathbf{w}}_+) = F_K(\tilde{\mathbf{w}}_-)$, $(G\mathcal{P}_{11})^2 = I$, $\mathcal{P}_{11}^2 = I$, and that the operators $\mathcal{P}_{11}G$ and $G\mathcal{P}_{11}$ are Hermitian. The obtained result indicates that if $b_+^{(1)} \neq 0$, then the bifurcation of nonlinear modes is possible only from one eigenstate, either from \tilde{b}_+ or from $\tilde{b}_- = -\tilde{b}_+$.

As an example, in Fig. 2 we report numerically obtained families of nonlinear modes at the double point D_1 where $\gamma_1 = \gamma_2 = \sqrt{5}/4$. Two families of unstable

nonlinear modes bifurcate from the positive eigenvalue and no nonlinear modes bifurcates from the negative one (hereafter the stability of nonlinear modes have been investigated by means of analysis of the spectrum of the linearized problem).

5.2.4. Double points between T_2 and T_3 (between T_1 and T_4). Such double points lie on the hyperbola $\gamma_2 = 1 - 1/\gamma_1$ (for the double points located between the triple points T_2 and T_3), or on $\gamma_2 = -1 - 1/\gamma_1$ (for the double points between T_1 and T_4). In this case spectrum of $H_{nn}(\gamma_1, \gamma_2)$ contains a double zero eigenvalue $\tilde{b}_0 = 0$ corresponding to a 2 × 2 Jordan block, and hence

$$\det H_{nn}(\gamma_1, \gamma_2) = 0. \tag{32}$$

Two other eigenvalues of $H_{nn}(\gamma_1, \gamma_2)$ are simple, real and have opposite values.

The peculiarity of these double points is that due to the particular structure of the linear eigenvector $\tilde{\mathbf{w}}$ and the nonlinearity $F_K(\tilde{\mathbf{w}})$, Eq. (28) yields that the coefficient $b^{(1)}$ is equal to zero. Then for any $b^{(2)}$ Eq. (27) has a nontrivial solution $\mathbf{w}^{(3)}$. This also implies that $\mathbf{w}^{(2)} = 0$. Therefore, for a more detail description of the families bifurcating from the double eigenvalue \tilde{b}_0 one has to proceed to next orders of the expansions (25)–(26). The modified expansions read

$$\mathbf{w} = \varepsilon \tilde{\mathbf{w}} + \varepsilon^3 \mathbf{w}^{(3)} + \varepsilon^4 \mathbf{w}^{(4)} + \varepsilon^5 \mathbf{w}^{(5)} \cdots,$$
(33)

$$b = \varepsilon^2 b^{(2)} + \varepsilon^3 b^{(3)} + \varepsilon^4 b^{(4)} \cdots,$$
(34)

which in the ε^4 -order gives $\mathbf{w}^{(4)} = b^{(3)}\mathbf{v}_1$, while in the ε^5 -order we obtain

$$b^{(4)}\tilde{\mathbf{w}} + b^{(2)}\mathbf{w}^3 = H_{nn}\tilde{\mathbf{w}}^{(5)} + \mathbf{f}_5, \tag{35}$$

where $\mathbf{f_5}$ is the ε^5 -order contribution of the nonlinear term $F_K(\mathbf{w})\mathbf{w}$:

$$\mathbf{f_5} = \tilde{\mathbf{w}} \circ \tilde{\mathbf{w}} \circ \left(\mathbf{w}^{(3)}\right)^* + 2\tilde{\mathbf{w}} \circ \tilde{\mathbf{w}}^* \circ \mathbf{w}^{(3)},\tag{36}$$

where we used "o" to designate element-wise multiplication of vectors. From Eq. (35) we obtain

$$b^{(2)}\langle \tilde{\mathbf{w}}^*, \mathbf{w}^{(3)} \rangle = \langle \tilde{\mathbf{w}}^*, \mathbf{f}_5 \rangle. \tag{37}$$

This equation together with Eq. (27) can be used to compute $b^{(2)}$. To this end, let us first solve Eq. (27) with $b^{(2)} = 0$, i.e.

$$H_{nn}\mathbf{w}^{(3)} = -F_K(\tilde{\mathbf{w}})\tilde{\mathbf{w}} \tag{38}$$

with respect to $\mathbf{w}^{(3)}$. In spite of the equality (32), Eq. (38) does have a \mathcal{PT} -invariant solution. Indeed, if \mathbf{u} is an arbitrary solution of Eq. (38), then applying \mathcal{PT} operator two both sides of (38) and using the fact that the nonlinearity $F_K(\mathbf{w})$ is weakly \mathcal{PT} symmetric [i.e. obeys (13)], we find that $\mathcal{PT}\mathbf{u}$ is also solution of the same equation. Thus a \mathcal{PT} -invariant solution of Eq. (38) is given by $\mathbf{u}_0 = \frac{1}{2}(\mathbf{u} + \mathcal{PT}\mathbf{u})$.

On the other hand, to construct a solution of Eq. (38) we let the first entry of $\mathbf{w}^{(3)}$ to be a free parameter and express all other entries of $\mathbf{w}^{(3)}$ in terms of the first one. Introducing $c = w_1^{(3)}$ (where c is the free parameter and $w_1^{(3)}$ is the first entry of $\mathbf{w}^{(3)}$), for the next entries we find $w_2^{(3)} = -i\gamma_1 c - |\tilde{w}_1|^2 \tilde{w}_1$, and $w_3^{(3)} = i\gamma_2 |\tilde{w}_1|^2 \tilde{w}_1 - \gamma_1 c - |\tilde{w}_2|^2 \tilde{w}_2$, where \tilde{w}_j are the entries of the linear eigenvector $\tilde{\mathbf{w}}$. (Here we have also used relation $\gamma_1 \gamma_2 + 1 = \gamma_1$ which is valid for the double points on the boundary between T_2 and T_3 ; for the double points situated on the boundary between T_1 and T_4 the resulting expressions will be slightly different but having the same structure.) Looking for a \mathcal{PT} -invariant solution (which is shown to exist) we Nonlinear lattices with PT-symmetric linear part

require $(w_3^{(3)})^* = w_2^{(3)}$, which results in a system of two linear equations with respect to $c_1 = \operatorname{Re} c$ and $c_2 = \operatorname{Im} c$:

$$\gamma_1(c_1 + c_2) = \operatorname{Re}\left(|\tilde{w}_1|^2 \tilde{w}_1 - |\tilde{w}_2|^2 \tilde{w}_2^* - i\gamma_2 |\tilde{w}_1|^2 \tilde{w}_1^* \right), \tag{39}$$

$$\gamma_1(c_1 + c_2) = \operatorname{Im} \left(|\tilde{w}_1|^2 \tilde{w}_1 - |\tilde{w}_2|^2 \tilde{w}_2^* - i\gamma_2 |\tilde{w}_1|^2 \tilde{w}_1^* \right).$$

$$(40)$$

While the determinant of latter system is zero, the system has to be compatible (otherwise, the \mathcal{PT} -invariant solution would not exist). We set $c_2 = 0$, and then the parameter c is fixed as follows:

$$c = \operatorname{Re} c = \frac{1}{\gamma_1} \operatorname{Re} \left(|\tilde{w}_1|^2 \tilde{w}_1 - |\tilde{w}_2|^2 (\tilde{w}_2)^* - i\gamma_2 |\tilde{w}_1|^2 \tilde{w}_1^* \right).$$
(41)

Therefore, the \mathcal{PT} -invariant solution for Eq. (38) is written down as

$$\mathbf{u}_{0} = \begin{pmatrix} c \\ -i\gamma_{1}c - |\tilde{w}_{1}|^{2}\tilde{w}_{1} \\ i\gamma_{1}c - |\tilde{w}_{1}|^{2}\tilde{w}_{1}^{*} \\ c \end{pmatrix}.$$
(42)

Notice that choice $c_2 = 0$ was not restrictive since the most general \mathcal{PT} -invariant solution of Eq. (38) can be found as $\mathbf{w}^{(3)} = \mathbf{u}_0 + d\tilde{\mathbf{w}}$, where the particular solution \mathbf{u}_0 is fixed by Eqs. (41)–(42), and where d is arbitrary real number, which can always be set equal to zero by means of rescaling of the small parameter ε .

Solution of Eq. (27) with arbitrary $b^{(2)}$ can be found as $\mathbf{w}^{(3)} = \mathbf{u}_0 + b^{(2)}\mathbf{v}$. Substituting the latter expression into Eq. (37), one obtains a quadratic equation with respect to $b^{(2)}$ (notice that all the coefficients of the latter equation are real and expressed in terms of $\tilde{\mathbf{w}}$, \mathbf{v} and $\gamma_{1,2}$).

Resorting at this stage to numerics, we consider two double points: $D_2 = (\frac{1}{2}, -1)$ and $D_3 = (\frac{3}{2}, \frac{1}{3})$. Finding for both of them the quadratic equation for $b^{(2)}$ and computing its roots, we observe that both for D_2 and D_3 the roots are real, nonzero, and distinct from each other. Moreover, for the point D_2 both roots are of the same sign, while for D_3 the roots have opposite signs. Therefore, both for D_2 and D_3 two families of nonlinear modes are expected to bifurcate from the double eigenvalue b = 0. This is confirmed by numerical results in Fig. 2 where we show families of nonlinear modes both for the points D_2 and D_3 . Bifurcation diagrams for D_2 and D_3 are similar locally near the double eigenvalue b = 0 in terms of existence: both for D_2 and D_3 there are two families bifurcating from b. Notice however that for the point D_2 both the families bifurcating from b bifurcate to the right (i.e. to the half-plane b > 0), while at the point D_3 one of the families bifurcates to the right and another one bifurcates to the left (i.e. to the half-plane b < 0). This behavior is in agreement with different signs of the roots of the quadratic equations for $b^{(2)}$. We also notice that stability of the bifurcating families is different: for the point D_2 one of the families is stable (in the vicinity of the bifurcation), while another one is unstable; on the other hand, for the point D_3 both the bifurcating families are stable in the sufficiently small vicinity of the bifurcation. Existence of stable nonlinear modes in spite of the presence of the exceptional point singularity is quite remarkable.

5.2.5. Triple points There exist exactly four triple points T_j , j = 1, 2, 3, 4. Each triple point corresponds to the situation when the Hamiltonian $H_{nn}(\gamma_1, \gamma_2)$ has a zero eigenvalue $\tilde{b}_0 = 0$ with multiplicity equal to 4. The canonical Jordan representation of $H_{nn}(\gamma_1, \gamma_2)$ in the triple points is given by a 4×4 Jordan block. Therefore, our analysis

in Eqs. (24)–(28) is not applicable in this case. However, for the sake of completeness of our studies, we report numerical results on the behavior of the nonlinear modes bifurcating from the triple points.

The points T_2 and T_3 correspond to values of γ_1 given as real roots of the equation $\gamma_1^4 - 2\gamma_1^2 - 2\gamma_1 + 1 = 0$, and $\gamma_2 = 1 - 1/\gamma_1$. For the triple points T_1 and T_4 one has the equation $\gamma_1^4 - 2\gamma_1^2 + 2\gamma_1 + 1 = 0$, and $\gamma_2 = -1 - 1/\gamma_1$. Notice that for each triple points T_j one has $\gamma_1^2 + \gamma_2^2 = 3$.

In Fig. 2 we show numerical results for points $T_1 \approx (-0.37, 1.69)$ and $T_2 \approx (1.68, 0.41)$ which feature different bifurcation diagrams. All the found modes are unstable.

6. Nonlinearities allowing for integrals of motion

6.1. General idea

It is known that the integrals of motion are of fundamental importance for the conservative systems, while they do not necessarily exist for the dissipative ones. Although the dissipative systems typically do not conserve the total energy, they can admit other conserved quantities and even be integrable [1]. For linear \mathcal{PT} -symmetric systems, some integrals of motion can be found in an explicit form [42], but in the nonlinear case the only system with known integrals of motion (to the best of the authors' knowledge) is the exactly integrable dimer [3] (see also [43] where some of the results stemming from the integrable dynamics of a dimer were generalized to the chain of coupled dimers).

So far, we considered how stationary nonlinear modes depend on the character of the linear eigenstate they bifurcate from and on the type of the nonlinearity $F(\mathbf{q})$. Let us now consider how the choice of nonlinearity $F(\mathbf{q})$ can affect dynamical properties of the system. More specifically, we address the existence of the conserved quantities of the nonlinear system (1). In this section, we find a condition which must be satisfied by the nonlinear operator $F(\mathbf{q})$ for the nonlinear system (1) to support at least one integral of motion. Using this result, we report several integrals of motion for a \mathcal{PT} symmetric quadrimer.

A motivation for our consideration is the known fact that a linear \mathcal{PT} system [which formally corresponds to Eq. (1) with $F(\mathbf{q}) = 0$] has an integral which is given as the "pseudo-power" $Q = \langle \mathcal{P}\mathbf{q}(t), \mathbf{q}(t) \rangle$ [42] [this fact can be also easily verified using Eqs. (2)]. However, the equality $\dot{Q} = 0$ generically does not hold for the nonlinear system (1) with $F(\mathbf{q}) \neq 0$. To establish conditions for the nonlinear system to admit an integral of motion, we look for a conserved quantity in the form $Q = \langle A\mathbf{q}(t), \mathbf{q}(t) \rangle$ where A is an arbitrary (so far) time-independent linear operator. Then (1) yields

$$i\dot{Q} = -\langle A[H + F(\mathbf{q})]\mathbf{q}, \mathbf{q} \rangle + \langle [H^{\dagger} + F^{\dagger}(\mathbf{q})]A\mathbf{q}, \mathbf{q} \rangle.$$
(43)

For Q to be a conserved quantity it is sufficient to require

$$AH = H^{\dagger}A \tag{44}$$

and

$$AF(\mathbf{q}) = F^{\dagger}(\mathbf{q})A \quad \text{for all } \mathbf{q}.$$

$$\tag{45}$$

A particularly interesting case corresponds to a situation when the properties (44)–(45) hold for $A = \mathcal{P}$, where \mathcal{P} is a parity operator. Then condition (44) is

equivalent to the pseudo-Hermiticity [30] of H, see also Eq. (2). Equation (45) now gives

$$F^{\dagger}(\mathbf{q}) = \mathcal{P}F(\mathbf{q})\mathcal{P} \quad \text{for all } \mathbf{q},\tag{46}$$

i.e. for all \mathbf{q} the nonlinear operator $F(\mathbf{q})$ must be pseudo-Hermitian with respect to the same \mathcal{P} operator as the linear operator H. The nonlinearities $F(\mathbf{q})$ obeying to the property (46) will be said to be of the pseudo-Hermitian type and the class of such nonlinearities will be denoted as $NL_{pH}(\mathcal{P})$ (with the respective parity operator \mathcal{P} indicated in the brackets).

The following proposition establishes a simple relation between pseudo-Hermitian and \mathcal{PT} -symmetric nonlinearities introduced in Sec. 3.

Proposition 3. If $F(\mathbf{q}) \in NL_{\mathcal{PT}}(\mathcal{P})$ and $F(\mathbf{q}) = F^T(\mathbf{q})$, then $F \in NL_{pH}(\mathcal{P})$.

6.2. Example: PT-symmetric quadrimer with integrals of motion

Turning now to particular examples, we consider dynamics of the system (1) described by the linear Hamiltonian $H_{bc}(\gamma_1, \gamma_2)$ defined by Eq. (19) and by the nonlinearity $F_{bc}(\mathbf{q})$ given by (20). As it has already been noticed in Sec. 4.3, the linear operator $H_{bc}(\gamma_1, \gamma_2)$ is $\mathcal{P}_{10}\mathcal{T}$ -symmetric [see Eq. (21) for the definition of \mathcal{P}_{10}], and the nonlinearity $F_{bc}(\mathbf{q})$ is $\mathcal{P}\mathcal{T}$ symmetric, i.e. $F_{bc}(\mathbf{q}) \in NL_{\mathcal{P}\mathcal{T}}(\mathcal{P}_{10})$. Due to Proposition 3, the latter fact together with the diagonal structure of $F_{bc}(\mathbf{q})$ implies that $F_{bc}(\mathbf{q}) \in NL_{pH}(\mathcal{P}_{10})$ (this can also be checked in a straightforward manner). Hence there exists at least one integral of motion given by

$$Q_1 = \langle \mathcal{P}_{10} \mathbf{q}, \mathbf{q} \rangle = 2 \operatorname{Re} \left(q_1^* q_3 + q_2^* q_4 \right).$$
(47)

Furthermore, the model (19)–(20) possesses another integral of motion. To find it, we notice that operator $H_{bc}(\gamma_1, \gamma_2)$ is also $\mathcal{P}_{20}\mathcal{T}$ -symmetric with respect to

$$\mathcal{P}_{20} = \sigma_2 \otimes \sigma_0 = \begin{pmatrix} 0 & 0 & -i & 0 \\ 0 & 0 & 0 & -i \\ i & 0 & 0 & 0 \\ 0 & i & 0 & 0 \end{pmatrix}.$$
(48)

Notice that while the operator \mathcal{P}_{20} is Hermitian, it can not be considered as a conventional parity operator because it does not commute with the operator \mathcal{T} which in our case is the complex conjugation. However, one can check that $F_{bc}^{\dagger}(\mathbf{w}) = F_{bc}(\mathbf{w}) = \mathcal{P}_{20}F_{bc}(\mathbf{w})\mathcal{P}_{20}$, i.e. $F_{bc}(\mathbf{q}) \in NL_{pH}(\mathcal{P}_{20})$. Thus the second integral of motion is readily found to be

$$Q_2 = \langle \mathcal{P}_{20} \mathbf{q}, \mathbf{q} \rangle = 2 \mathrm{Im} \, (q_1^* q_3 + q_2^* q_4). \tag{49}$$

Obviously, Q_1 and Q_2 can be combined in a single complex-valued integral

$$Q = q_1^* q_3 + q_2^* q_4. ag{50}$$

The integral (50) can be also obtained from a more general consideration. Indeed, considering the operator

$$A = \begin{pmatrix} 0 & 0 & a_1 & 0 \\ 0 & 0 & 0 & a_1 \\ a_2 & 0 & 0 & 0 \\ 0 & a_2 & 0 & 0 \end{pmatrix}$$
(51)

where $a_{1,2}$ are arbitrary, one can verify that the both conditions (44)–(45) hold. Then a conserved quantity is given as

$$\langle A\mathbf{q}, \mathbf{q} \rangle = a_1(q_1^*q_3 + q_2^*q_4) + a_2(q_1q_3^* + q_2q_4^*).$$
(52)

Setting $a_2 = 0$ one readily obtains integral (50).

It turns out, however, that in a particular case when the linear part is given by $H_{bc}(\gamma, -\gamma)$, and $\varkappa = 1$ in Eq. (20), the obtained quantity Q is not the only integral of motion. Indeed, let us consider the model (18a)-(18d) with $\varkappa = 1$:

$$i\dot{q}_1 = i\gamma q_1 + q_2 + (|q_1|^2 + |q_3|^2)q_1, \tag{53a}$$

$$i\dot{q}_2 = -i\gamma q_2 + q_1 + (|q_2|^2 + |q_4|^2)q_2, \tag{53b}$$

$$i\dot{q}_3 = -i\gamma q_3 + q_4 + (|q_3|^2 + |q_1|^2)q_3, \tag{53c}$$

$$i\dot{q}_4 = i\gamma q_4 + q_3 + (|q_4|^2 + |q_2|^2)q_4.$$
(53d)

To construct a new integral, we rewrite Eqs. (53a)-(53d) in terms of the Stokes components (this approach was used in [3] in the study of the nonlinear \mathcal{PT} -symmetric dimer):

$$\begin{split} S^0_1 &= |q_1|^2 + |q_2|^2, \qquad S^0_2 &= |q_3|^2 + |q_4|^2, \\ S^1_1 &= q^*_1 q_2 + q_1 q^*_2 \qquad S^1_2 &= q^*_4 q_3 + q_4 q^*_3, \\ S^2_1 &= i(q_1 q^*_2 - q^*_1 q_2), \qquad S^2_2 &= i(q_4 q^*_3 - q^*_4 q_3), \\ S^3_1 &= |q_1|^2 - |q_2|^2, \qquad S^3_2 &= |q_4|^2 - |q_3|^2. \end{split}$$

We also introduce the Stokes vectors $\mathbf{S}_j = (S_j^1, S_j^2, S_j^3)$ with j = 1, 2, and notice that

$$\mathbf{S}_j \cdot \mathbf{S}_j = (S_j^0)^2. \tag{54}$$

Next, we introduce the pseudo-electric field $\mathbf{E} = (0, 0, 2\gamma)$, pseudo-magnetic fields $\mathbf{B}_0 = (-2, 0, 0)$ and $\mathbf{B} = (0, 0, S_2^3 - S_1^3)$, and rewrite the system (53*a*)-(53*d*) in the form

$$\dot{S}_1^0 = \mathbf{E} \cdot \mathbf{S}_1, \quad \dot{S}_2^0 = \mathbf{E} \cdot \mathbf{S}_2, \tag{55}$$

$$\dot{\mathbf{S}}_1 = S_1^0 \mathbf{E} + \mathbf{S}_1 \times (\mathbf{B}_0 + \mathbf{B}), \quad \dot{\mathbf{S}}_2 = S_2^0 \mathbf{E} + \mathbf{S}_2 \times (\mathbf{B}_0 - \mathbf{B}).$$
(56)

Further, multiplying Eqs. (56) by **B**, we obtain (j = 1, 2)

$$\mathbf{B} \cdot \dot{\mathbf{S}}_{j} = S_{j}^{0} \mathbf{B} \cdot \mathbf{E} + \mathbf{S}_{j} \cdot (\mathbf{B}_{0} \times \mathbf{B}), \tag{57}$$

and multiplying Eqs. (56) by \mathbf{B}_0 we obtain

$$\mathbf{B}_0 \cdot \dot{\mathbf{S}}_1 = -\mathbf{S}_1 \cdot (\mathbf{B}_0 \times \mathbf{B}), \qquad \mathbf{B}_0 \cdot \dot{\mathbf{S}}_2 = \mathbf{S}_2 \cdot (\mathbf{B}_0 \times \mathbf{B}).$$
(58)

Combining (57) and (58) yields the relations

$$\mathbf{B} \cdot \dot{\mathbf{S}}_1 = S_1^0 \, \mathbf{B} \cdot \mathbf{E} - \mathbf{B}_0 \cdot \dot{\mathbf{S}}_1, \quad \mathbf{B} \cdot \dot{\mathbf{S}}_2 = S_2^0 \, \mathbf{B} \cdot \mathbf{E} + \mathbf{B}_0 \cdot \dot{\mathbf{S}}_2. \tag{59}$$

After subtracting one of Eqs. (59) from another and using the definitions of the vectors \mathbf{B} , \mathbf{B}_0 , and \mathbf{E} , we arrive at

$$\begin{split} 0 &= -\mathbf{B} \cdot (\dot{\mathbf{S}}_1 - \dot{\mathbf{S}}_2) + (S_1^0 - S_2^0) \,\mathbf{B} \cdot \mathbf{E} - \mathbf{B}_0 \cdot (\dot{\mathbf{S}}_1 + \dot{\mathbf{S}}_2) \\ &= (S_2^3 - S_1^3) \frac{d}{dt} (S_2^3 - S_1^3) - (S_1^0 - S_2^0) \frac{d}{dt} (S_1^0 - S_2^0) + 2 \frac{d}{dt} (S_1^1 + S_2^1) \\ &= \dot{J}, \end{split}$$

where integral of motion J is given by

$$J = \frac{1}{2}(S_2^3 - S_1^3)^2 - \frac{1}{2}(S_2^0 - S_1^0)^2 + 2(S_2^1 + S_1^1).$$
(60)

Using Eqs. (54) one can express J only through the components of the threedimensional Stokes vectors $\mathbf{S}_{1,2}$.

Notice that in the limit $q_3 = q_4 = 0$, i.e. $S_2^0 = 0$ and $\mathbf{S}_2 = 0$, the obtained integral is reduced to one of the known integrals for a \mathcal{PT} -symmetric dimer [3].

Finally, we notice that system (53a)-(53d) admits a solution in quadratures in the particular case when Q = 0. Indeed, this implies that $|q_1|^2|q_3|^2 = |q_2|^2|q_4|^2$, which is equivalent to $S_1^0 S_2^3 = S_1^3 S_2^0$. Combing the latter relation with Eqs. (55), we have $S_1^0 \dot{S}_2^0 = S_2^0 \dot{S}_1^0$, and hence $S_1^0 = CS_2^0$, where C is a constant. Subsequently $\mathbf{E} \cdot \mathbf{S}_1 = C\mathbf{E} \cdot \mathbf{S}_2$, i.e. $S_1^3 = CS_2^3$ and $\mathbf{B} = (1 - C)(0, 0, S_2^3)$. The latter formula means that the dynamical equation for \mathbf{S}_2 is singled out and acquires the form of the equation for the dimer. Solutions for such a dimer in an implicit form have been found in [3].

7. Conclusion

In the presented paper we have investigated some of nonlinear properties of finitedimensional systems respecting \mathcal{PT} symmetry. In contrast to the most part of the previous studies, our discussion has not been focused on a nonlinearity of any given form. Instead, we have emphasized how nonlinearities of different classes can affect stationary and dynamical properties of the system. First, we have considered a class of nonlinearities with weak \mathcal{PT} symmetry, the latter appearing to be necessary (and in many cases sufficient) for the existence of the families of stationary nonlinear modes. We have paid particular attention to analysis of the bifurcations of nonlinear modes from the multiple eigenstates of the underlying linear problem. In a situation when the underlying linear system has a semi-simple eigenstate, we have shown that the invariant subspace associated with the degenerate eigenvalue can always be spanned by a complete basis of \mathcal{PT} -invariant eigenvectors. The established fact has been used to construct formal expansions for the nonlinear modes bifurcating from the degenerate eigenstate. Next, we considered bifurcations of stationary modes from exceptional points, which typically occur at the phase transition between unbroken and broken \mathcal{PT} symmetries. We have shown that the generalized eigenvector associated with the multiple eigenvalue can be chosen \mathcal{PT} invariant. Then we have developed smallamplitude expansions for the bifurcations of nonlinear modes and demonstrated that the possibility of the bifurcations as well as stability of the modes depend on both the nonlinearity and the character of the coalescing eigenstates.

To complete the above picture, we notice that the nonlinear modes can also exist in the case when \mathcal{PT} symmetry of the underlying linear problem is broken, i.e. the spectrum of the underlying linear operator $H(\gamma)$ contains complex eigenvalues. If all the linear eigenvalues are complex, then the nonlinear modes obviously cannot bifurcate from the linear eigenstates, i.e. such nonlinear modes (if exist) do not have linear counterpart. Moreover, in the case of a finite-dimensional system (like a \mathcal{PT} symmetric oligomer) nonlinear modes can be stable even if the \mathcal{PT} symmetry of the linear problem is broken [4].

Next, we have turned to dynamical properties of nonlinear \mathcal{PT} -symmetric lattices and indicated another important class of nonlinearities, termed pseudo-Hermitian nonlinearities, which allow for the nonlinear system to admit at least one integral of motion. Using this idea, we have found several integrals for a \mathcal{PT} -symmetric nonlinear quadrimer and demonstrated that (at least in some cases) it admits a solution in quadratures.

Acknowledgements

Authors are grateful to P. G. Kevrekidis and U. Günther for stimulating discussion. The work was supported by FCT (Portugal) through the FCT grants PEst-OE/FIS/UI0618/2011, PTDC/FIS-OPT/1918/2012, and SFRH/BPD/64835/2009.

References

- Akhmediev N and Ankiewicz A 2005 Dissipative Solitons in the Complex Ginzburg–Landau and Swift-Hohenberg Equations Dissipative Solitons (Lect. Notes Phys. vol 661) ed N Akhmediev and A Ankiewicz (Springer, Berlin Heidelberg) p 1-17
- [2] Bender C M 2007 Rep. Prog. Phys. 70 947
- [3] Ramezani H, Kottos T, El-Ganainy R, and Christodoulides D 2010 Phys. Rev. A 82 043803
- [4] Zezyulin D A and Konotop V V 2012 Phys. Rev. Lett. **108** 213906
- [5] Li K, Zezyulin D A, Konotop V V, and Kevrekidis P G 2013 Phys. Rev. A 87 033812
- [6] Kevrekidis P G, Pelinovsky D E, and Tyugin D Y 2013 Nonlinear Stationary States in PTsymmetric Lattices *Preprint* arXiv:1303.3298 [nlin.PS]
- [7] Leykam D, Konotop V V, and Desyatnikov A S 2013 Opt. Lett. 38 371
- [8] Li K and Kevrekidis P G 2011 Phys. Rev. E 83 066608
- [9] Li K, Kevrekidis P G, Malomed B A, and Günther U 2012 J. Phys. A: Math. Theor. 45 444021
- [10] Suchkov S V, Malomed B A, Dmitriev S V, and Kivshar Yu S 2011 Phys. Rev. E 84 046609
- [11] Konotop V V, Pelinovsky D E, and Zezyulin D A 2012 EPL 100 56006
- [12] Musslimani Z H, Makris K G, El-Ganainy R, and Christodoulides D N 2008 Phys. Rev. Lett. 100 030402
- [13]~ Nixon S, Ge L, and Yang J 2012 Phys. Rev. A ${\bf 85}$ 023822
- [14] Moreira F C, Konotop V V, Malomed B A 2013 Phys. Rev. A 87 013832
- [15] Zezyulin D A and Konotop V V 2012 Phys. Rev. A 85 043840
- [16] Wang H and Wang J 2011 Opt. Express 19 4030
- [17] Moreira F C, Abdullaev F K, Konotop V V, and Yulin A V 2012 Phys. Rev. A 86 053815
- [18] Zhu X, Wang H, Zheng L-X, Li H, and He Y-J 2011 Opt. Lett. 36 2680
- [19] Abdullaev F K, Kartashov Y V, Konotop V V, and Zezyulin D A 2011 Phys. Rev. A 83 041805
- [20] Zezyulin D A, Kartashov Y V and Konotop V V 2011 EPL 96 64003
- [21] Miroshnichenko A E, Malomed B A, and Kivshar Yu S, 2011 Phys. Rev. A 84 012123
- [22] Duanmu M, Li K, Horne R L, Kevrekidis P G and Whitaker N 2013 Phil. Trans. R. Soc. A 371 20120171
- [23] He Y, Zhu X, Mihalache D, Liu J, and Chen Z 2012 Phys. Rev. A 85 013831
- [24] Driben R, and Malomed B A 2011 Opt. Lett. 36, 4223
- [25] Barashenkov I V, Suchkov S V, Sukhorukov A A, Dmitriev S V, and Kivshar Yu S 2012 Phys. Rev. A 86 053809
- [26] Barashenkov I V, Baker L and Alexeeva N V 2013 Phys. Rev. A 87 033819
- [27] Graefe E-M 2012 J. Phys. A: Math. Theor. 45 444015
- [28] Cartarius H, Wunner G 2012 Phys. Rev. A 86 013612
- [29] Kreibich M, Main J, Cartarius H, and Wunner G 2013 Phys. Rev. A 87 051601(R)
- [30] Mostafazadeh A 2002 J. Math. Phys. 43 205
- [31] Kato T 1980 Perturbation Theory for Linear Operators (Springer-Verlag, Berlin)
- [32] Bender C M, Meisinger P N, and Wang Q 2003 J. Phys. A: Math. Gen. 36 6791
- [33] Graefe E-M, and Jones H (2011) Phys. Rev. A 84 013818
- [34] Ramezani H, Kottos T, Kovanis V, and Christodoulides D (2012) Phys. Rev. A 85 013818
- [35] Bender C M, Brody D C, and Jones H F 2002 Phys. Rev. Lett. 89 270401
- [36] Heiss W D 2012 J. Phys. A 45 44016
- [37] MacKay R S and Aubry S 1994 Nonlinearity 7 1623
- [38] Nixon S, Zhu Y, and Yang J 2012 Opt. Lett. 37 4874
- [39] Nixon S and Yang J 2013 Opt. Lett. 38 1933
- [40] Moiseyev N 2011 Non-Hermitian Quantum Mechanics (Cambridge: Cambridge University Press)
- Bender C M and Gianfreda M 2013 Twofold Transition in *PT*-Symmetric Coupled Oscillators Preprint arXiv:1305.7107 [hep-th]
- [42] Bagchi B, Quesne C, and Znojil M 2001 Mod. Phys. Lett. A 16 2047
- [43] Kevrekidis P G, Pelinovsky D E, and Tyugin D Y 2013 Nonlinear Dynamics in PT-Symmetric Lattices Preprint arXiv:1307.2973 [nlin.SI]