

The differential $d_4(h_6^3)$ in the Adams
spectral sequence for spheres

By
Pomin Wu

Adviser
Dr. Wên-Hsiung Lin

A Doctorial Dissertation
Presented to
Department of Mathematics
at
National Tsing-Hua University

In Partial Fulfillment
of the Requirement for the Degree of
Doctoral of Philosophy

Sep, 2011

Acknowledgements

I would like to thank my advisor, Dr. Wên-Hsiung Lin, for his infinite support and patience. It may seem impossible for me to fully comprehend every technique there is to learn from my teacher. Yet it is among the non-technical things that I believe I have learned the most valuable lesson. The image of a mathematician which I have seen from Dr. Lin will always aspire me to pursuit a deeper understanding of the subject. Therefore I feel I am most fortunate to have studied under his direction. I would also like to thank Mr. Tai-Wei Chen for his help on some of the most complicated computations in this thesis.

I am also greatly in debt to many friends who in the last few years have supported me in one way or another. You are my connections to this world, and it is only by these connections that a pursuit in abstract knowledge acquires its full meaning. Thank you for being here with me.

I must also thank my parents and my sister, for all the support a family can give to one of its members. You are the closest thing to a definition of me in this world.

Contents

1	Introduction	1
2	Outline of the proof of Theorem 1.4	3
3	Calculations of some Ext groups	15
4	Proofs of Theorem 2.39 and Theorem 2.40	49
5	Proof of Proposition 1.6, Proposition 1.7, and Proposition 2.46	66
6	Modification of the proof without θ_6	76

1 Introduction

Let \mathcal{A} denote the mod 2 Steenrod algebra. Let X be a stable complex in the stable category ([1]). Let $H_*(X)$ and $H^*(X)$ denote respectively the mod 2 reduced homology and cohomology of X . The Ext groups $\text{Ext}_{\mathcal{A}}^{*,*}(H^*(X), \mathbb{Z}/2)$, which will be simply denoted by $\text{Ext}_{\mathcal{A}}^{*,*}(X)$, form the E_2 term of the mod 2 Adams spectral sequence (to be abbreviated as ASS) for computing the 2-primary stable homotopy groups $\pi_*^S(X)$ of X ([1, 2]). In the case $X = S^0$, the sphere spectrum in stable dimension zero, the Ext groups $\text{Ext}_{\mathcal{A}}^{*,*}(H^*(S^0), \mathbb{Z}/2) = \text{Ext}_{\mathcal{A}}^{*,*}(\mathbb{Z}/2, \mathbb{Z}/2)$ will be simply denoted by $\text{Ext}_{\mathcal{A}}^{*,*}$, and the 2-primary stable homotopy groups $\pi_*^S(S^0)$ by π_*^S .

For each $i \geq 0$, let $h_i \in \text{Ext}_{\mathcal{A}}^{1,2^i} = \text{Ext}_{\mathcal{A}}^{1,2^i}(\mathbb{Z}/2, \mathbb{Z}/2)$ be the class corresponding to the generator $Sq^{2^i} \in \mathcal{A}$. It is also known that $h_i^2 \neq 0$, $h_i^3 \neq 0$ for all $i \geq 0$ ([1]). The classes h_0 , h_1 , h_2 , and h_3 are known to detect homotopy elements $2\iota \in \pi_0^S$, $\eta \in \pi_1^S$, $\nu \in \pi_3^S$, and $\sigma \in \pi_7^S$, where η , ν , and σ are the three Hopf classes. So h_0^2 , h_1^2 , h_2^2 , and h_3^2 detect respectively the homotopy elements $4\iota = (2\iota)^2$, η^2 , ν^2 , and σ^2 , and h_0^3 , h_1^3 , h_2^3 , and h_3^3 detect respectively the homotopy elements $8\iota = (2\iota)^3$, η^3 , ν^3 , and σ^3 . (All of these homotopy elements are non-zero in π_*^S ([21, 22])) J. F. Adams also proves in [1] that for each $i > 3$, h_i does not detect homotopy elements in π_*^S by showing that

$$(1.1) \quad d_2(h_i) = h_{i-1}^2 h_0 \neq 0 \quad \text{for } i \geq 4 \text{ in the ASS for } \pi_*^S.$$

So it is not immediate whether h_i^2 detect homotopy elements for $i \geq 4$. Any homotopy element in $\pi_{2^{i+1}-2}^S$ for $i \geq 1$ which is detected by h_i^2 is denoted by θ_i . θ_i , if exists, is known as a Kervaire invariant element [7, 14]. Thus, $\eta^2 = \theta_1$, $\nu^2 = \theta_2$, and $\sigma^2 = \theta_3$. Mahowald and Tangora prove in [19] that h_4^2 does detect homotopy elements in π_{30}^S . That is, there exists a θ_4 . In fact they have shown that $\pi_{30}^S = \mathbb{Z}/2(\theta_4)$. So $2\theta_4 = 0$. Barratt, Jones, and Mahowald prove in [4] that h_5^2 also detects homotopy elements in π_{62}^S . Kochman and Mahowald have shown in [15] that there actually exists a θ_5 with $2\theta_5 = 0$. Recently, M. Hill, M. Hopkins, and D. Ravenel have shown that h_i^2 does not detect homotopy elements for $i \geq 7$. The status of h_6^2 is still open.

For the family $\{h_i^3\}_{i \geq 0}$, h_0^3 , h_1^3 , h_2^3 , and h_3^3 are known to detect homotopy elements as already mentioned above. Barratt, Mahowald, and Tangora have shown in [4] that h_4^3 detects homotopy elements in π_{45}^S . W. H. Lin has shown in [17] that h_7^3 does not detect homotopy elements. More precisely, Lin proves that $d_4(h_7^3)$ is essentially $h_0^3 g_5 \neq 0$ in the ASS for spheres. We refer to [17] for the details of this ‘‘essentiality.’’ Here we recall [17] that for each $i \geq 1$, there is a non-zero class $g_i \in \text{Ext}_{\mathcal{A}}^{4,2^{i+3}+2^{i+2}}$, and $\text{Ext}_{\mathcal{A}}^{4,2^{i+3}+2^{i+2}} = \mathbb{Z}/2(g_i)$. It is known ([24]) that $h_0^2 g_1 \neq 0$, $h_0^2 g_2 \neq 0$, and $h_0^3 g_1 = 0$, $h_0^3 g_2 = 0$. Lin has shown in [17] that $h_0^3 g_5 \neq 0$ as has just been mentioned above. If one can show $h_0^3 g_{i-2} \neq 0$ for $i \geq 8$, then Lin’s method in [17] actually also proves that $d_4(h_i^3)$ is essentially $h_0^3 g_{i-2}$ for all $i \geq 8$. He also remarks in [17] that his method does not include the cases $i = 5$ and 6. In [13] Kan has shown that $h_0^3 g_3 \neq 0$ and that $d_4(h_5^3) = h_0^3 g_3 \neq 0$ in the ASS for spheres.

It is the purpose of this paper to show the following.

Theorem 1.2. *In the mod 2 Adams spectral sequence for spheres, $d_4(h_6^3) = h_0^3 g_4 \neq 0$ in $\text{Ext}_{\mathcal{A}}^{7,195}$.*

This together with [4, 13, 17, 21, 22] mentioned above completely settle the differentials of the h_i^3 family.

The plan for this paper goes as follows. We will prove the following Ext group result in section 3.

(1.3) $h_0^3 g_4 \neq 0$ in $\text{Ext}_{\mathcal{A}}^{7,195}$.

Granting this, in section 2 we will outline the proof of the following result.

Theorem 1.4. $h_0^3 g_4$ is a boundary in the ASS for spheres.

In section 3 we will also prove the following Ext group results.

Proposition 1.5. (1) $\text{Ext}_{\mathcal{A}}^{s,s+189} = 0$ for $0 \leq s \leq 2$.

(2) $\text{Ext}_{\mathcal{A}}^{s,s+189} = \mathbb{Z}/2(h_0^{s-3} h_6^3)$ for $3 \leq s \leq 4$.

(3) $\text{Ext}_{\mathcal{A}}^{5,194} = \mathbb{Z}/2(h_0^2 h_6^3) \oplus \mathbb{Z}/2(h_1 g_4)$.

In section 4 we will compute the stable homotopy groups $\pi_{62}^S(P_{46}^{62})$ and $\pi_{62}^S(P_{47}^{62})$ (P_k^l are the stunted projective spaces) which we need to complete the proof of Theorem 1.4. In section 5 we will complete the proof of Theorem 1.4, and also prove the following results in the ASS for spheres.

Proposition 1.6. In the ASS for spheres, $d_r(h_6^3) = 0 \in \text{Ext}_{\mathcal{A}}^{3+r,191+r}$ for $2 \leq r \leq 3$.

Proposition 1.7. In the ASS for spheres, $d_2(h_1 g_4) \neq h_0^3 g_4$.

Proof of Theorem 1.2. By Theorem 1.4 $d_r(\alpha) = h_0^3 g_4$ for some $\alpha \in \text{Ext}_{\mathcal{A}}^{7-r,196-r}$, $2 \leq r \leq 7$. From 1.6, 1.7, and from the fact that h_0 is an infinite cycle in the ASS for spheres, it is easy to see that $d_2(h_0^2 h_6^3) = 0$, $d_2(h_1 g_4) \neq h_0^3 g_4$, $d_3(h_0 h_6^3) = 0$. So by Proposition 1.5, the only possibility is $d_4(h_6^3) = h_0^3 g_4$. This completes the proof of Theorem 1.2. \square

2 Outline of the proof of Theorem 1.4

We recall again that in the stable homotopy groups $\pi_*^S = \pi_*^S(S^0)$, there exist homotopy elements $\theta_4 \in \pi_{30}^S = \mathbb{Z}/2$ and $\theta_5 \in \pi_{62}^S$ detected respectively by h_4^2 and h_5^2 in the ASS for π_*^S , with $2\theta_4 = 0$, $2\theta_5 = 0$ ([4, 15, 19]). Throughout this paper we fix a θ_5 with $2\theta_5 = 0$. We also assume that there exists a $\theta_6 \in \pi_{126}^S$. (Recall that it is still not known whether θ_6 exists or not.) Our proof of the main theorem is more conveniently presented when θ_6 is assumed to be existent. The method of our proof can be modified to work without the knowledge of θ_6 , and this will be given in section 6 of the paper. Again, in what follows, we fix a θ_6 .

A key step in Kan's proof of the result $d_4(h_5^3) = h_0^3 g_3$ in his thesis [13] is to show that $\theta_4 \theta_5 \in \pi_{92}^S$ is non-zero and is detected by $h_0^2 g_3 \in \text{Ext}_{\mathcal{A}}^{6,98}$. Our proof of Theorem 1.4 is based on a similar result which is Theorem 2.2 below. To state it, we recall ([16]) that for each $i \geq 0$, there is a non-zero class $D_3(i) \in \text{Ext}_{\mathcal{A}}^{4,2^{i+6}+2^i}$. Consider the classes $h_0^2 g_4$ and $h_5^2 D_3(1)$ in $\text{Ext}_{\mathcal{A}}^{6,194}$. In section 3 we will show the following Ext group result.

(2.1) $h_0^2 g_4 \neq 0$, $h_5^2 D_3(1) \neq 0$, and they are linearly independent in $\text{Ext}_{\mathcal{A}}^{6,194}$.

The main work of this section is to prove the following theorem.

Theorem 2.2. $\theta_5 \theta_6$ in π_{188}^S is detected by $h_0^2 g_4 + \varepsilon h_5^2 D_3(1) \neq 0$ in $\text{Ext}_{\mathcal{A}}^{6,194}$, where $\varepsilon = 0$ or 1.

From this we will deduce Theorem 1.4 at the end of this section.

Remark 2.3. From Proposition 1.5 and Proposition 1.6, we see $h_0^2 g_4 + \varepsilon h_5^2 D_3(1)$ is not a boundary in the ASS for π_*^S .

Theorem 2.2 is a consequence of Theorem 2.17 later, which is a result about the stable homotopy group $\pi_{188}^S(P)$ parallel to the result about π_{188}^S in Theorem 2.2 above. Here P is the suspension spectrum of the infinite real projective space P^∞ . We need to recall the Kahn-Priddy theorem ([12]) in order to state Theorem 2.17, and also to see what the "parallel" is. In the next few paragraphs we do this recalling.

For any integer $n \geq 0$, the real projective space P^n is the space of all lines in \mathbb{R}^{n+1} through the origin. There is an inclusion $P^n \subset P^{n+1}$ induced by the natural inclusion $\mathbb{R}^{n+1} \subset \mathbb{R}^{n+2}$. Regard P^n as a subspace of P^{n+1} with this inclusion for each $n \geq 0$, and let $P^\infty = \bigcup_{n=0}^\infty P^n$. The suspension spectrum of the space P^n is also denoted by P^n , and the suspension spectrum of the space P^∞ is denoted by P as mentioned above.

For any integer $n \geq 0$ there is a space map $\Sigma^{n+1} P^n \xrightarrow{t_n} S^{n+1}$ which is called the n th transfer map. These transfer maps are compatible with respect to n , inducing a stable map $P \xrightarrow{t} S^0$, called the transfer map, from the spectrum P to the sphere spectrum S^0 . These are described in (2.5) through Definition 2.9 below.

Given a point $L \in P^n$, which is a line through the origin of \mathbb{R}^{n+1} , let L^\perp be the n -dimensional hyperplane through the origin of \mathbb{R}^{n+1} that is perpendicular to L , and let $\mathbb{R}^{n+1} \xrightarrow{r_L} \mathbb{R}^{n+1}$ be the map which is the reflection with respect to L^\perp . Then r_L is an element of $O(n+1)$, the group of orthogonal transformations

from \mathbb{R}^{n+1} to itself. Clearly the map $P^n \xrightarrow{r_n} O(n+1)$ defined by $r_n(L) = r_L$ is continuous for each n . Note that

$$(2.4) \quad r_n(L) \in O(n+1) - SO(n+1) \quad \forall L \in P^n,$$

where $SO(n+1) \subset O(n+1)$ is the subgroup of orthogonal transformations of determinant 1. These maps are compatible with respect to n in the sense that the following diagram, where the vertical maps are obvious inclusions, is commutative:

$$(2.5) \quad \begin{array}{ccc} P^n & \xrightarrow{r_n} & O(n+1) \\ \downarrow & & \downarrow \\ P^{n+1} & \xrightarrow{r_{n+1}} & O(n+2) \end{array}$$

S^{n+1} can be regarded as $\mathbb{R}^{n+1} \cup \{\infty\}$, the one-point compactification of \mathbb{R}^{n+1} . Then each element in $O(n+1)$ can be thought as a continuous function from $S^{n+1} = \mathbb{R}^{n+1} \cup \{\infty\}$ to itself, sending ∞ to ∞ . The $(n+1)$ -fold loop space $\Omega^{n+1}S^{n+1}$ is the function space of all the maps from S^{n+1} to itself preserving the base point ∞ . Thus we get a natural map $O(n+1) \xrightarrow{i_n} \Omega^{n+1}S^{n+1}$ for each n , and these maps are compatible with respect to n in the sense that the following diagram, where the vertical maps are obvious inclusions, is commutative:

$$(2.6) \quad \begin{array}{ccc} O(n) & \xrightarrow{i_{n-1}} & \Omega^n S^n \\ \downarrow & & \downarrow \\ O(n+1) & \xrightarrow{i_n} & \Omega^{n+1} S^{n+1} \end{array}$$

Note that

(2.7) $P^n \xrightarrow{i_n r_n} \Omega^{n+1} S^{n+1}$ maps P^n to the “-1” component of $\Omega^{n+1} S^{n+1}$ by (2.4), since each element of $O(n+1) - S(n+1)$ when considered as a self-map of S^{n+1} is a degree -1 map.

From (2.5) and (2.6) we have the following commutative diagram.

$$\begin{array}{ccccc} P^n & \xrightarrow{r_n} & O(n+1) & \xrightarrow{i_n} & \Omega^{n+1} S^{n+1} \\ \downarrow & & \downarrow & & \downarrow \\ P^{n+1} & \xrightarrow{r_{n+1}} & O(n+2) & \xrightarrow{i_{n+1}} & \Omega^{n+2} S^{n+2} \end{array}$$

Definition 2.8. The n th transfer map $\Sigma^{n+1} P^n \xrightarrow{t_n} S^{n+1}$ is the $(n+1)$ -fold adjoint map to $P^n \xrightarrow{i_n r_n} \Omega^{n+1} S^{n+1}$.

It is clear that t_n is also compatible with respect to n in the sense that the following diagram, where the left vertical map is the obvious inclusion, is commutative:

$$\begin{array}{ccc} \Sigma^{n+2} P^n & \xrightarrow{\Sigma t_n} & \Sigma S^{n+1} \\ \downarrow & & \parallel \\ \Sigma^{n+2} P^{n+1} & \xrightarrow{t_{n+1}} & S^{n+2} \end{array}$$

Definition 2.9. *The family of n th transfer maps $\{t_n\}$ induces a stable map $P \xrightarrow{t} S^0$ from the suspension spectrum P to the sphere spectrum S^0 . This map t is called the transfer map.*

Consider the induced map $\pi_*^S(P) \xrightarrow{t_\#} \pi_*^S$ of $P \xrightarrow{t} S^0$. Kahn-Priddy theorem ([12]) says that $t_\#$ is onto in the 2-primary components for $* > 0$. This (Kahn-Priddy theorem) is seen as follows. Take the $(n+1)$ loop map

$$(2.10) \quad \Omega^{n+1}\Sigma^{n+1}P^n \xrightarrow{\Omega^{n+1}t_n} \Omega^{n+1}S^{n+1}$$

of the n th transfer map $\Sigma^{n+1}P^n \xrightarrow{t_n} S^{n+1}$. Let $(\Omega^{n+1}S^{n+1})_{-1}$ be the “ -1 ” component of $\Omega^{n+1}S^{n+1}$ as mentioned in (2.7). Let $\Omega^\infty P = \lim_n \Omega^{n+1}\Sigma^{n+1}P^n$, $\Omega^\infty S^0 = \lim_n \Omega^{n+1}S^{n+1}$ and $(\Omega^\infty S^0)_{-1} = \lim_n (\Omega^{n+1}S^{n+1})_{-1}$. Letting $n \rightarrow \infty$ we get a space map

$$(2.11) \quad \Omega^\infty P \xrightarrow{\Omega^\infty t} \Omega^\infty S^0.$$

The stable homotopy groups $\pi_*^S(P)$ (resp. π_*^S) is one-to-one correspondent to the unstable homotopy groups $\pi_*(\Omega^\infty P)$ (resp. $\pi_*(\Omega^\infty S^0)$), and the group homomorphism $\pi_*^S(P) \xrightarrow{t_\#} \pi_*^S$ corresponds to the induced homomorphism $\pi_*(\Omega^\infty P) \xrightarrow{(\Omega^\infty t)_\#} \pi_*(\Omega^\infty S^0)$. Note that

(2.12) $\pi_*(\Omega^\infty S^0)$ is equal to $\pi_*((\Omega^\infty S^0)_{-1})$ for all $* > 0$.

By Sullivan’s theory of localization ([23]) there is a space ${}_2(\Omega^\infty S^0)_{-1}$ whose homotopy groups are 2-primary, and also a map $(\Omega^\infty S^0)_{-1} \xrightarrow{L} {}_2(\Omega^\infty S^0)_{-1}$ such that $\pi_*((\Omega^\infty S^0)_{-1}) \xrightarrow{L_\#} \pi_*({}_2(\Omega^\infty S^0)_{-1})$ is a 2-primary isomorphism for all $*$. Let $\Omega^\infty S^0 \xrightarrow{p} (\Omega^\infty S^0)_{-1}$ be the projection to the “ -1 ” component. Then Kahn-Priddy theorem says the following composite

$$\Omega^\infty P \xrightarrow{\Omega^\infty t} \Omega^\infty S^0 \xrightarrow{p} (\Omega^\infty S^0)_{-1} \xrightarrow{L} {}_2(\Omega^\infty S^0)_{-1}$$

is a retraction map. That is, there is a map ${}_2(\Omega^\infty S^0)_{-1} \xrightarrow{i} \Omega^\infty P$ such that the composite ${}_2(\Omega^\infty S^0)_{-1} \xrightarrow{i} \Omega^\infty P \xrightarrow{Lp\Omega^\infty t} (\Omega^\infty S^0)_{-1}$ is homotopic to the identity map. Therefore $\Omega^\infty P \xrightarrow{\Omega^\infty t} \Omega^\infty S^0$ induces a homomorphism $\pi_*^S(P) \xrightarrow{t_\#} \pi_*^S$ that is surjective on the 2-primary homotopy groups for $* > 0$. Summarizing, we have the following result.

(2.13) Given a 2-primary homotopy element $\alpha \in \pi_*^S$ with order $k \geq 1$ for $* > 0$. Then there exists an element $\beta \in \pi_*^S(P)$ also with order k , such that $t_\#(\beta) = \alpha$.

The transfer map $P \xrightarrow{t} S^0$ also induces an Ext group map

$$(2.14) \quad \text{Ext}_{\mathcal{A}}^{s,t}(P) \xrightarrow{t_*} \text{Ext}_{\mathcal{A}}^{s+1,t+1}.$$

This Ext group map (2.14) is a realization of the map $\pi_*^S(P) \xrightarrow{t_\#} \pi_*^S$. The algebraic Kahn-Priddy theorem by Lin [18] says that t_* is also onto for $t-s > 0$. In particular we have the following.

(2.15) For $n \geq 1$ there exists a class $\widehat{h}_n \in \text{Ext}_{\mathcal{A}}^{0,2^n-1}(P)$ such that $t_*(\widehat{h}_n) = h_n$. So $t_*(\widehat{h}_n h_n) = h_n^2$. In fact it is easy to see that $\text{Ext}_{\mathcal{A}}^{0,2^n-1}(P) = \mathbb{Z}/2(\widehat{h}_n)$, and $\text{Ext}_{\mathcal{A}}^{1,2^{n+1}-1}(P) = \mathbb{Z}/2(\widehat{h}_n h_n)$, for $n \geq 1$.

There is a result in $\pi_*^S(P)$ that is parallel to Theorem 2.2 by Kahn-Priddy theorem. To describe the result in $\pi_*^S(P)$ which is Theorem 2.17 below, first we note the following observation.

(2.16) Suppose there is an element $\theta_n \in \pi_{2^{n+1}-2}^S$ detected by h_n^2 in the ASS for π_*^S with $2\theta_n = 0$. Then by (2.13) and (2.15) there exists an element $\widehat{\theta}_n \in \pi_{2^{n+1}-2}^S(P)$ detected by $\widehat{h}_n h_n$ such that $t_{\#}(\theta_n) = \widehat{\theta}_n$ and $2\widehat{\theta}_n = 0$. By dimensional reason, $\widehat{\theta}_n \in \pi_{2^{n+1}-2}^S(P)$ can be pulled back to an element in $\pi_{2^{n+1}-2}^S(P^{2^{n+1}-2})$, which will also be denoted by $\widehat{\theta}_n$.

We already know there exist $\theta_4 \in \pi_{30}^S$ and $\theta_5 \in \pi_{62}^S$ with $2\theta_4 = 0$, $2\theta_5 = 0$, so by (2.16) there exist $\widehat{\theta}_4 \in \pi_{30}^S(P)$ and $\widehat{\theta}_5 \in \pi_{62}^S(P)$ with $2\widehat{\theta}_4 = 0$, $2\widehat{\theta}_5 = 0$. Also, these latter $\widehat{\theta}_4$ and $\widehat{\theta}_5$ have pullbacks $\widehat{\theta}_4 \in \pi_{30}^S(P^{30})$ and $\widehat{\theta}_5 \in \pi_{62}^S(P^{62})$ respectively. We will see in a moment ((2.18) below) that these pullbacks $\widehat{\theta}_4$, $\widehat{\theta}_5$ do not have $2\widehat{\theta}_4 = 0$, $2\widehat{\theta}_5 = 0$. Throughout this paper we fix a $\theta_4 \in \pi_{30}^S$, a $\theta_5 \in \pi_{62}^S$ with $2\theta_4 = 0$, $2\theta_5 = 0$, and a $\widehat{\theta}_4 \in \pi_{30}^S(P)$, a $\widehat{\theta}_5 \in \pi_{62}^S(P)$ with $2\widehat{\theta}_4 = 0$, $2\widehat{\theta}_5 = 0$. We also fix a pullback $\widehat{\theta}_4 \in \pi_{30}^S(P^{30})$, and a pullback $\widehat{\theta}_5 \in \pi_{62}^S(P^{62})$.

In [16] it is shown that for each $i \geq 1$ there is a non-zero class $\widehat{g}_i \in \text{Ext}_{\mathcal{A}}^{3,2^{i+3}+2^{i+2}-1}(P)$ with $t_*(\widehat{g}_i) = g_i \in \text{Ext}_{\mathcal{A}}^{4,2^{i+3}+2^{i+2}}$ where t_* is as in (2.14). In section 3 we will show that $\widehat{g}_4 \in \text{Ext}_{\mathcal{A}}^{3,191}(P)$ pulls back to a class in $\text{Ext}_{\mathcal{A}}^{3,191}(P^{62})$, which is also denoted by \widehat{g}_4 . There is a class $\widehat{h}_5 \in \text{Ext}_{\mathcal{A}}^{1,32}(P)$ by (2.15). It is clear that $\widehat{h}_5 \in \text{Ext}_{\mathcal{A}}^{1,32}(P)$ pulls back uniquely to a class in $\text{Ext}_{\mathcal{A}}^{1,32}(P^{62})$, which is also denoted by \widehat{h}_5 . We will see in a moment that $\widehat{g}_4 h_0^2, \widehat{h}_5 h_5 D_3(1) \in \text{Ext}_{\mathcal{A}}^{5,193}(P^{62})$ are linearly independent. The following is the result in $\pi_*^S(P)$ parallel to Theorem 2.2.

Theorem 2.17. *Let $\widehat{\theta}_5 \in \pi_{62}^S(P^{62})$ be as in (2.16). In $\pi_{188}^S(P^{62})$, $\widehat{\theta}_5 \theta_6 \neq 0$ and is detected in the ASS for π_*^S by a class*

$$\tau = \widehat{g}_4 h_0^2 + \varepsilon \widehat{h}_5 h_5 D_3(1) + \overline{\tau} \neq 0 \quad \text{in } \text{Ext}_{\mathcal{A}}^{5,193}(P^{62})$$

where $\varepsilon = 0$ or 1 . Here $\overline{\tau} \in \text{Ext}_{\mathcal{A}}^{5,193}(P^{62})$ is a class such that $t_*(\overline{\tau}) = 0$ in $\text{Ext}_{\mathcal{A}}^{6,194}$. Moreover, $t_*(\widehat{g}_4 h_0^2) = h_0^2 g_4$, and $t_*(\widehat{h}_5 h_5 D_3(1)) = h_5^2 D_3(1)$.

We remark that t_* in Theorem 2.17 is the restriction to P^{62} of the t_* in (2.14). And also note that from (2.1) and Theorem 2.17 we see that the classes $\widehat{g}_4 h_0^2$, $\widehat{h}_5 h_5 D_3(1)$, and $\overline{\tau}$ are linearly independent in $\text{Ext}_{\mathcal{A}}^{5,193}(P^{62})$ (if $\overline{\tau} \neq 0$), because $t_*(\widehat{g}_4 h_0^2) = h_0^2 g_4$, $t_*(\widehat{h}_5 h_5 D_3(1)) = h_5^2 D_3(1)$, and $t_*(\overline{\tau}) = 0$. It is easy to see then that Theorem 2.2 is a direct consequence of Theorem 2.17 and Remark 2.3.

To prove Theorem 2.17 we will need to recall a well-known result on the stable liftings of the stable map $S^{62} \xrightarrow{2\widehat{\theta}_5} P^{62}$, and to make an analysis on the ‘‘Adams’ filtration’’ of the homotopy element $\widehat{\theta}_5 \theta_6 \in \pi_{188}^S(P^{62})$ which is represented by the composite $S^{188} \xrightarrow{\theta_6} S^{62} \xrightarrow{\widehat{\theta}_5} P^{62}$. These are recalled in the next few paragraphs.

First note that the stable map $S^{62} \xrightarrow{2\widehat{\theta}_5} P^{62}$ is non-zero, because $\widehat{h}_5 h_0 h_5 \in \text{Ext}_{\mathcal{A}}^{2,64}(P^{62})$ is a non-zero infinite cycle detecting $2\widehat{\theta}_5$ in the ASS for $\pi_*^S(P^{62})$. In general, we have the following.

(2.18) If $S^{2^{i+1}-2} \xrightarrow{\widehat{\theta}_i} P^{2^{i+1}-2}$ is a lifting of a Kervaire invariant element $S^{2^{i+1}-2} \xrightarrow{\theta_i} S^0$ with $2\theta_i = 0$ ($i \geq 4$), then the stable map $S^{2^{i+1}-2} \xrightarrow{2\widehat{\theta}_i} P^{2^{i+1}-2}$ is non-zero as $2\widehat{\theta}_i \in \pi_{2^{i+1}-2}^S(P^{2^{i+1}-2})$ is detected in the ASS for $\pi_*^S(P^{2^{i+1}-2})$ by $\widehat{h}_i h_0 h_i$.

For $0 \leq l \leq m$, we say a stable map $S^m \xrightarrow{\varphi} P^m$ can be lifted to P^l if there is a stable map $S^m \xrightarrow{\psi} P^l$ such that the composite $S^m \xrightarrow{\psi} P^l \xrightarrow{i} P^m$ is φ as shown in the following diagram (2.19), where $P^l \xrightarrow{i} P^m$ is the inclusion map. ψ is said to be a lifting of φ to P^l .

$$(2.19) \quad \begin{array}{ccc} & & P^l \\ & \nearrow \psi & \downarrow i \\ S^m & \xrightarrow{\varphi} & P^m \end{array}$$

The result we are going to use is the following.

Proposition 2.20. *The stable map $S^{62} \xrightarrow{2\widehat{\theta}_5} P^{62}$ can be lifted to P^{51} and no further.*

The proof of Proposition 2.20 is based on the theory of vector fields on spheres. We recall this theory as follows.

For each $n \geq 1$, a non-zero vector field on S^{n-1} is a continuous map $S^{n-1} \xrightarrow{\sigma} S^{n-1}$ such that $\sigma(v) \perp v$ for all $v \in S^{n-1}$. A set of non-zero vector fields $\{\sigma_1, \dots, \sigma_r\}$ on S^{n-1} is said to be linearly independent if the set of vectors $\{\sigma_1(v), \dots, \sigma_r(v)\}$ is linearly independent for all $v \in S^{n-1}$. In the case where there is a linearly independent set $\{\sigma_1, \dots, \sigma_r\}$ of vector fields on S^{n-1} , we say S^{n-1} admits an r -field.

(2.21) Given an integer $n \geq 2$, write $n = 2^b(2a + 1)$, and let $b = c + 4d$, where $0 \leq c \leq 3$. Let $\rho(n) = 2^c + 8d$. It is classical that for each $n \geq 2$, S^{n-1} admits a $(\rho(n) - 1)$ -field. ([10])

J. F. Adams proves the following result in [3].

Theorem 2.22. *Let $\rho(n)$ be as defined in (2.21). S^{n-1} does not admit a $\rho(n)$ -field.*

For $n \geq 1$, the Whitehead square $[\iota_n, \iota_n] = \omega_n \in \pi_{2n-1}(S^n)$ is the homotopy element of the following composite of maps,

$$S^{2n-1} \xrightarrow{\sigma} S^n \vee S^n \xrightarrow{F} S^n,$$

where σ is the attaching map of the top cell of $S^n \times S^n = (S^n \vee S^n) \cup_{\sigma} e^{2n}$, and F is the folding map. We say ω_n can be desuspended r times or ω_n is an r -fold suspension (for $r \geq 0$), if there is a map $S^{2n-1-r} \xrightarrow{\widehat{\omega}_n} S^{n-r}$ such that $[\Sigma^r \widehat{\omega}_n] = \omega_n$. The problem of how far can ω_n be desuspended is connected to the vector field problem on the spheres in the following content as shown in [11].

Theorem 2.23. *Given an integer $r \geq 0$ with $r \leq n$. If S^n admits an r -field then ω_n is an r -fold suspension. Conversely if ω_n is an r -fold suspension, and if $n > 2r$, then S^n admits an r -field.*

For $n \geq 1$ with $n \leq 5$, let $m = 2^{n+1} - 2$, consider a space map $\Sigma^{m+1}S^m \xrightarrow{2\widehat{\theta}_n} \Sigma^{m+1}P^m$ (which is within the stable range) that represents $2\widehat{\theta}_n \in \pi_m^S(P^m)$. We refer to [5] for the following.

(2.24) For $r < \frac{m+2}{2}$, ω_{m+1} can be desuspended r times if and only if there is a lifting $\Sigma^{m+1}S^m \xrightarrow{\psi} \Sigma^{m+1}P^{m-r}$ of the stable map $\Sigma^{m+1}S^m \xrightarrow{2\widehat{\theta}_n} \Sigma^{m+1}P^m$.

Combining the results in Theorem 2.22, Theorem 2.23, and (2.24), we have the following

Corollary 2.25. *There is a (stable) lifting $S^{2^{n+1}-2} \xrightarrow{\psi} P^{2^{n+1}-2-k}$ of $S^{2^{n+1}-2} \xrightarrow{2\widehat{\theta}_n} P^{2^{n+1}-2}$ if and only if $k \leq \rho(2^{n+1}) - 1$.*

When $n = 5$, $\rho(64) = 12$, so Proposition 2.20 is a special case of Corollary 2.25.

Next we define ‘‘Adams’ filtration’’ $AF(\alpha)$ of a homotopy element $\alpha \in \pi_*^S(X)$ as follows.

Let K be the Eilenberg-MacLane spectrum $K(\mathbb{Z}/2)$, that is, $\pi_0^S(K) = \mathbb{Z}/2$ and $\pi_*^S(K) = 0$ for $* \neq 0$. Given a stable complex X , recall that the mod 2 Adams resolution of X consists of the followings: for each $s \geq 0$ there is a spectrum X_s , a map $X_{s+1} \xrightarrow{p_{s+1}} X_s$, a set of integers $I_s \subset \mathbb{N}$, a spectrum $K_s = \bigvee_{q \in I_s} \Sigma^q K$, and a map $X_s \xrightarrow{f_s} K_s$, such that $X_0 = X$, and for each $s \geq 0$, the map $X_s \xrightarrow{f_s} K_s$ induces epimorphism on the mod 2 cohomology, and there is a cofibration sequence

$$X_{s+1} \xrightarrow{p_{s+1}} X_s \xrightarrow{f_s} K_s$$

Given a map $Y \xrightarrow{\varphi} X$, we say $AF(\varphi)$, the Adams’ filtration of φ , is l if $l \geq 0$ and l is the largest integer such that there is a lifting $Y \xrightarrow{\varphi_l} X_l$ of φ to X_l in

the following tower (l may be ∞):

$$\begin{array}{ccccc}
& & \vdots & & \\
& & \downarrow & & \\
& & \Sigma^{-1}K_l \xrightarrow{j_{l+1}} X_{l+1} \xrightarrow{f_{l+1}} K_{l+1} & & \\
& & \downarrow p_{l+1} & & \\
& & \Sigma^{-1}K_{l-1} \xrightarrow{j_l} X_l \xrightarrow{f_l} K_l & & \\
& & \downarrow p_l & & \\
& & X_{l-1} \xrightarrow{f_{l-1}} K_{l-1} & & \\
& & \downarrow p_{l-1} & & \\
& & \vdots & & \\
& & \downarrow & & \\
& & \Sigma^{-1}K_1 \xrightarrow{j_2} X_2 \xrightarrow{f_2} K_2 & & \\
& & \downarrow p_2 & & \\
& & \Sigma^{-1}K_0 \xrightarrow{j_1} X_1 \xrightarrow{f_1} K_1 & & \\
& & \downarrow p_1 & & \\
& & X_0 \xrightarrow{f_0} K_0 & & \\
& \nearrow \varphi_l & & & \\
Y & \xrightarrow{\varphi} & X_0 & \xrightarrow{f_0} & K_0
\end{array}$$

Let $Z \xrightarrow{\psi} Y$ be another map, then it is not difficult to show that $AF(\varphi\psi) \geq AF(\varphi)$.

In order to prove Theorem 2.17, we have to prove several propositions first. For any $0 \leq k \leq n$ let $P_k^n = P^n/P^{k-1}$. Let $P^{62} \xrightarrow{q_k} P_k^{62}$ be the collapsing map and $P^k \xrightarrow{i_k} P^{62}$ be the inclusion map.

Proposition 2.26. *For any integer $k \geq 1$, the composite of stable maps $S^{62} \xrightarrow{2\iota} S^{62} \xrightarrow{\widehat{\theta}_5} P^{62} \xrightarrow{q_k} P_k^{62}$ is essential if and only if $k \leq 51$.*

Proof. The composite $S^{62} \xrightarrow{2\iota} S^{62} \xrightarrow{\widehat{\theta}_5} P^{62} \xrightarrow{q_k} P_k^{62}$ is zero if and only if there is a lifting of $S^{62} \xrightarrow{2\widehat{\theta}_5} P^{62}$ to P^{k-1} because $P_k^{62} = P^{62}/P^{k-1}$. By Proposition 2.20 such a lifting exists if and only if $k-1 \geq 51$. So the composite $S^{62} \xrightarrow{2\iota} S^{62} \xrightarrow{\widehat{\theta}_5} P^{62} \xrightarrow{q_k} P_k^{62}$ is zero if and only if $k \geq 52$, or equivalently, the composite $S^{62} \xrightarrow{2\iota} S^{62} \xrightarrow{\widehat{\theta}_5} P^{62} \xrightarrow{q_k} P_k^{62}$ is essential if and only if $k \leq 51$. This proves Proposition 2.26. \square

Corollary 2.27. *The composite of the stable maps $S^{62} \xrightarrow{\widehat{\theta}_5} P^{62} \xrightarrow{q_{47}} P_{47}^{62}$ is essential.*

Proof. By Proposition 2.26, $S^{62} \xrightarrow{q_{47}(2\widehat{\theta}_5)} P_{47}^{62}$ is essential, so $S^{62} \xrightarrow{q_{47}\widehat{\theta}_5} P_{47}^{62}$ must also be essential. \square

The homotopy element $q_{47}\widehat{\theta}_5 \in \pi_{62}^S(P_{47}^{62})$ will be described more clearly in Theorem 2.39 through Proposition 2.41 when we give a concrete result on the group $\pi_{62}^S(P_{47}^{62})$. For this purpose we are going to obtain a partial result on the Adams filtration $AF(q_{47}\widehat{\theta}_5)$ in Proposition 2.32. To this end we need to consider a (stable) complex \widetilde{P}^{62} and also a map $P^{62} \xrightarrow{\widetilde{q}} \widetilde{P}^{62}$. These are described in the next paragraph.

Recall the stable homotopy element $\widehat{\theta}_4 \in \pi_{30}^S(P)$ with $2\widehat{\theta}_4 = 0$ in (2.16). Consider a pullback $\widehat{\theta}_4 \in \pi_{30}^S(P^{31})$ of $\widehat{\theta}_4 \in \pi_{30}^S(P)$. From the cofibration sequence

$$\cdots \rightarrow \Sigma^{-1}P_{32} \rightarrow P^{31} \rightarrow P \rightarrow P_{32} \rightarrow \cdots,$$

we have $\pi_{30}^S(P^{31}) \cong \pi_{30}^S(P)$ since $\pi_{30}^S(P_{32}) = 0$ and $\pi_{31}^S(P_{32}) = 0$. So $2\widehat{\theta}_4 = 0 \in \pi_{30}^S(P^{31})$ also. Now consider the pullback of $\widehat{\theta}_4 \in \pi_{30}^S(P)$ to $\pi_{30}^S(P^{30})$ as defined in (2.16) which has $2\widehat{\theta}_4 \neq 0 \in \pi_{30}^S(P^{30})$ by (2.18). The composite

$$S^{30} \xrightarrow{2\iota} S^{30} \xrightarrow{\widehat{\theta}_4} P^{30}$$

is non-zero. Let $P^{30} \xrightarrow{i_{30}} P^{31}$ be the inclusion map, then the composite

$$S^{30} \xrightarrow{2\iota} S^{30} \xrightarrow{\widehat{\theta}_4} P^{30} \xrightarrow{i_{30}} P^{31}$$

is zero as already shown above. So $S^{30} \xrightarrow{i_{30}\widehat{\theta}_4} P^{31}$ has an extension \overline{h}'_5 :

$$S^{30} \cup_{2\iota} e^{31} \xrightarrow{\overline{h}'_5} P^{31}.$$

Let $P^{31} \xrightarrow{i_{31}} P^{62}$ be the inclusion map, and let \overline{h}_5 be the composite

$$(2.28) \quad \overline{h}_5 : S^{30} \cup_{2\iota} e^{31} \xrightarrow{\overline{h}'_5} P^{31} \xrightarrow{i_{31}} P^{62}.$$

Definition 2.29. Define \widetilde{P}^{62} to be the mapping cone $C_{\overline{h}_5} = P^{62} \cup_{\overline{h}_5} C(S^{30} \cup_{2\iota} e^{31})$, and define $P^{62} \xrightarrow{\widetilde{q}} \widetilde{P}^{62} = C_{\overline{h}_5}$ to be the inclusion map.

There are two properties of this complex \widetilde{P}^{62} , which are (2.30) and (2.31) below, that will be important to our proof later.

(2.30) \overline{h}_5 induces an isomorphism on the mod 2 homology groups $H_{31}(S^{30} \cup_{2\iota} e^{31}) \xrightarrow[\cong]{(\overline{h}_5)_*} H_{31}(P^{62})$.

Recall from (2.15) that $\text{Ext}_{\mathcal{A}}^{0,31}(P^{62}) \cong \text{Ext}_{\mathcal{A}}^{0,31}(P) = \mathbb{Z}/2(\widehat{h}_5)$. Then (2.30) implies

(2.31) $\text{Ext}_{\mathcal{A}}^{0,31}(P^{62}) \xrightarrow{\tilde{q}_*} \text{Ext}_{\mathcal{A}}^{0,31}(\tilde{P}^{62})$ has $\tilde{q}_*(\widehat{h}_5) = 0$.

(2.30) is shown as follows. We have the following diagram, where $S^{30} \xrightarrow{\bar{\sigma}} \Sigma^{-1}P^{31}$ is the attaching map of the top cell of $\Sigma^{-1}P^{32}$, and all other horizontal maps are cofibrations.

$$\begin{array}{ccccccc}
& & S^{30} & \xrightarrow{2\iota} & S^{30} & \longrightarrow & S^{30} \cup_{2\iota} e^{31} & \longrightarrow & S^{31} \\
& \nearrow^{m\iota} & \downarrow n\iota & \downarrow \widehat{\theta}_4 & \downarrow \widehat{h}'_5 & & \downarrow n\iota & & \downarrow n\iota \\
S^{30} & \xrightarrow{\bar{\sigma}} & \Sigma^{-1}P^{31} & \xrightarrow{\Sigma^{-1}q} & S^{30} & \xrightarrow{\sigma} & P^{30} & \xrightarrow{i_{30}} & P^{31} & \xrightarrow{q} & S^{31} \\
& \searrow_{2\iota} & & & & & & & \downarrow i_{31} & & \\
& & & & & & & & P^{62} & &
\end{array}$$

$S^{30} \xrightarrow{i_{30}\widehat{\theta}_4(2\iota)} P^{31}$ is zero implies that there is a pullback $S^{30} \xrightarrow{n\iota} S^{30}$ of $S^{30} \xrightarrow{2\widehat{\theta}_4} P^{30}$ for some $n \in \mathbb{Z}$, and $n\iota \neq 0$ because $S^{30} \xrightarrow{2\widehat{\theta}_4} P^{30}$ is non-zero. This pullback $n\iota$ is subjected to an indeterminacy $S^{30} \rightarrow \Sigma^{-1}P^{31} \xrightarrow{\Sigma^{-1}q} S^{30}$. Note that the composite $S^{30} \xrightarrow{\bar{\sigma}} \Sigma^{-1}P^{31} \xrightarrow{\Sigma^{-1}q} S^{30}$ is 2ι . So $S^{30} \xrightarrow{\Sigma^{-1}q\bar{\sigma}(m\iota)=2m\iota} S^{30}$ is an indeterminacy for any $m \in \mathbb{Z}$. Thus we can choose a pullback $S^{30} \xrightarrow{n\iota} S^{30}$ of $S^{30} \xrightarrow{2\widehat{\theta}_4} P^{30}$ with $n = 1$. By this choice, $H_{31}(S^{30} \cup_{2\iota} e^{31}; \mathbb{Z}) \xrightarrow{(\widehat{h}_5)_*} H_{31}(P^{62}; \mathbb{Z})$ is an isomorphism. This proves (2.30).

We can now prove a partial result on the Adams filtration of $S^{62} \xrightarrow{q_{47}\theta_5} P_{47}^{62}$.

Proposition 2.32. $AF(q_{47}\widehat{\theta}_5) \geq 3$ for the composite $S^{62} \xrightarrow{\theta_5} P^{62} \xrightarrow{q_{47}} P_{47}^{62}$.

To prove Proposition 2.32 we will use the following results (2.33) and (2.34). Let \tilde{P}^{46} be the 46-skeleton of \tilde{P}^{62} . From (2.28) we see that $\tilde{P}^{62}/\tilde{P}^{46} = P_{47}^{62}$. Let $\tilde{P}^{62} \xrightarrow{q'_{47}} \tilde{P}^{62}/\tilde{P}^{46} = P_{47}^{62}$ be the collapsing map. We have the following diagram consisting of a commutative square together with the map $S^{62} \xrightarrow{\widehat{\theta}_5} P^{62}$:

$$(2.33) \quad \begin{array}{ccc}
S^{62} & \xrightarrow{\widehat{\theta}_5} & P^{62} & \xrightarrow{q_{47}} & P_{47}^{62} \\
& & \downarrow \tilde{q} & & \parallel \\
& & \tilde{P}^{62} & \xrightarrow{q'_{47}} & \tilde{P}^{62}/\tilde{P}^{46}
\end{array}$$

In section 3 we will prove the following result

(2.34) $\text{Ext}_{\mathcal{A}}^{2,64}(\tilde{P}^{62}) = \mathbb{Z}/2(\overline{\tilde{e}_{31}h_5h_0}) \oplus \mathbb{Z}/2(\overline{\tilde{e}_{47}h_0h_4})$, and $(q'_{47})_*(\overline{\tilde{e}_{31}h_5h_0}) = (q'_{47})_*(\overline{\tilde{e}_{47}h_0h_4}) = 0$ in $\text{Ext}_{\mathcal{A}}^{2,64}(P_{47}^{62})$.

Proof of Proposition 2.32. From (2.33) we see that $q_{47}\widehat{\theta}_5 = q'_{47}\widehat{q}\widehat{\theta}_5$. $\widehat{\theta}_5$ is detected by \widehat{h}_5h_5 in the ASS for $\pi_*^S(P^{62})$. By (2.31), $\tilde{q}_*(\widehat{h}_5) = 0$. This implies $AF(\widehat{q}\widehat{\theta}_5) \geq 2$. If $AF(\widehat{q}\widehat{\theta}_5) = 2$, then $\widehat{q}\widehat{\theta}_5$ is detected by some class $\tau_1 \in \text{Ext}_{\mathcal{A}}^{2,64}(\tilde{P}^{62})$. By (2.34), $(q'_{47})_*(\tau_1) = 0$; so $AF(q'_{47}\widehat{q}\widehat{\theta}_5) \geq 3$. If $AF(\widehat{q}\widehat{\theta}_5) \geq 3$ then clearly $AF(q'_{47}\widehat{q}\widehat{\theta}_5) \geq 3$. In both cases we have $AF(q_{47}\widehat{\theta}_5) = AF(q'_{47}\widehat{q}\widehat{\theta}_5) \geq 3$. This proves Proposition 2.32. \square

We have to describe the map $S^{62} \xrightarrow{q_{47}\hat{\theta}_5} P_{47}^{62}$ more precisely than the result in (2.32) in order to prove Theorem 2.17. First we describe some specific homotopy elements $y, x, w \in \pi_{62}^S(P_{46}^{62})$, and $v \in \pi_{62}^S(P_{47}^{62})$ in (2.35) through (2.38) below. In (2.35), (2.36), and (2.38) the horizontal sequences are the obvious cofibration sequences.

(2.35) Define $y \in \pi_{62}^S(P_{46}^{62})$ as follows. Recall the Hopf classes $S^1 \xrightarrow{\eta} S^0$ and $S^7 \xrightarrow{\sigma} S^0$. Since $2\sigma^2 = 0$ and $\eta\sigma^2 = 0$, and since $P_{46}^{48} = (S^{46} \vee S^{47}) \cup_{\eta \vee 2\iota} e^{48}$, there is a coextension $\tilde{\sigma}^2$:

$$\begin{array}{ccccccc} \dots & \longrightarrow & S^{46} \vee S^{47} & \xrightarrow{i} & P_{46}^{48} & \xrightarrow{q} & S^{48} \xrightarrow{\eta \vee 2\iota} S^{47} \vee S^{48} \longrightarrow \dots \\ & & & & \swarrow \tilde{\sigma}^2 & & \uparrow \sigma^2 \\ & & & & & & S^{62} \end{array}$$

Let y be the composite $S^{62} \xrightarrow{\tilde{\sigma}^2} P_{46}^{48} \xrightarrow{i_{48}} P_{46}^{62}$, where i_{48} is the inclusion map. Then $y \in \pi_{62}^S(P_{46}^{62})$.

(2.36) Define $w \in \pi_{62}^S(P_{46}^{62})$ as follows. Consider the subcomplex $S^{47} \cup_{\sigma} e^{55} \subset P_{46}^{62}$. Since $\sigma(2\sigma) = 0$ there is a non-zero coextension $\overline{2\sigma}$:

$$\begin{array}{ccccccc} \dots & \longrightarrow & S^{47} & \xrightarrow{i'} & S^{47} \cup_{\sigma} e^{55} & \xrightarrow{q'} & S^{55} \xrightarrow{\sigma} S^{48} \longrightarrow \dots \\ & & & & \swarrow \overline{2\sigma} & & \uparrow 2\sigma \\ & & & & & & S^{62} \end{array}$$

Let w be the composite $S^{62} \xrightarrow{\overline{2\sigma}} S^{47} \cup_{\sigma} e^{55} \xrightarrow{j} P_{46}^{62}$, where j is the inclusion map. Then $w \in \pi_{62}^S(P_{46}^{62})$.

(2.37) Define $x = w + y \in \pi_{62}^S(P_{46}^{62})$.

(2.38) Define $v \in \pi_{62}^S(P_{47}^{62})$ as follows. The cells in dimensions 47, 51, and 53 in P_{47}^{62} form a subcomplex which is $S^{47} \cup_{\nu} e^{51} \cup_{\eta} e^{53} \subset P_{47}^{62}$. Recall ([25]) there is a stable map $S^8 \xrightarrow{\varepsilon} S^0$ which has the properties $\eta\varepsilon \neq 0$ and $\eta^2\varepsilon = 0$. It is not difficult to see that the Toda bracket $\langle \nu, \eta, \eta\varepsilon \rangle$ is zero. So there is a non-zero coextension $\tilde{\eta}\varepsilon$ of $\eta\varepsilon$:

$$\begin{array}{ccccccc} \dots & \longrightarrow & S^{47} \cup_{\nu} e^{51} & \xrightarrow{i''} & S^{47} \cup_{\nu} e^{51} \cup_{\eta} e^{53} & \xrightarrow{q''} & S^{53} \longrightarrow S^{48} \cup_{\nu} e^{52} \longrightarrow \dots \\ & & & & \swarrow \tilde{\eta}\varepsilon & & \uparrow \eta\varepsilon \\ & & & & & & S^{62} \end{array}$$

Let v be the composite $S^{62} \xrightarrow{\tilde{\eta}\varepsilon} S^{47} \cup_{\nu} e^{51} \cup_{\eta} e^{53} \xrightarrow{j'} P_{47}^{62}$, where j' is the inclusion map. Then v is an element in $\pi_{62}^S(P_{47}^{62})$.

We will show in section 4 that the elements $y, x, w \in \pi_{62}^S(P_{46}^{62})$ and $v \in \pi_{62}^S(P_{47}^{62})$ are non-zero. In fact, in section 4 we will the following Theorem 2.39 and Theorem 2.40.

Theorem 2.39. $\pi_{62}^S(P_{46}^{62}) = \mathbb{Z}/4(y) \oplus \mathbb{Z}/8(x)$. Moreover $AF(y) = 1$, $AF(w) = 1$, $AF(2y) = 2$, and $AF(x = y + w) = 2$.

Let $P_{46}^{62} \xrightarrow{q_1} P_{47}^{62}$ be the collapsing map, and consider the induced homomorphism $\pi_{62}^S(P_{46}^{62}) \xrightarrow{(q_1)_\#} \pi_{62}^S(P_{47}^{62})$. Let $\tilde{y} = (q_1)_\#(y)$, $\tilde{w} = (q_1)_\#(w)$, and $\tilde{x} = (q_1)_\#(x = y + w) = \tilde{y} + \tilde{w}$ in $\pi_{62}^S(P_{47}^{62})$.

Theorem 2.40. $\pi_{62}^S(P_{47}^{62}) = \mathbb{Z}/2(\tilde{y}) \oplus \mathbb{Z}/8(\tilde{x}) \oplus \mathbb{Z}/2(v - 2\tilde{x})$. Moreover $AF(\tilde{y}) = 1$, $AF(\tilde{x}) = 2$, and $AF(v) = 4$.

Now we can describe precisely the map $q_{47}\hat{\theta}_5$ considered in Corollary 2.27.

Proposition 2.41. $q_{47}\hat{\theta}_5 = 2\tilde{x} \in \pi_{62}^S(P_{47}^{62}) = \mathbb{Z}/2(\tilde{y}) \oplus \mathbb{Z}/8(\tilde{x}) \oplus \mathbb{Z}/2(v - 2\tilde{x})$.

Proof. Let $P_{46}^{62} \xrightarrow{q_{46}} P_{46}^{62}$ be the collapsing map. It is clear that $(q_{46})_*(\hat{h}_5) = 0$, and since $\hat{\theta}_5$ is detected by $\hat{h}_5 h_5$, it follows that $AF(q_{46}\hat{\theta}_5) \geq 2$. From Theorem 2.39 we thus see that $q_{46}\hat{\theta}_5 = ay + bx$ where $a = 0$ or 2 and $0 \leq b \leq 7$ (if $a = 1$ or 3 then $AF(q_{46}\hat{\theta}_5) = 1$). Since $2\tilde{y} = 0$ in $\pi_{62}^S(P_{47}^{62})$, we have

$$q_{47}\hat{\theta}_5 = q_1 q_{46}\hat{\theta}_5 = q_1(ay + bx) = a\tilde{y} + b\tilde{x} = b\tilde{x} \in \pi_{62}^S(P_{47}^{62})$$

If b is odd then $AF(q_{47}\hat{\theta}_5) = 2$ by Theorem 2.40. But Proposition 2.32 says $AF(q_{47}\hat{\theta}_5) \geq 3$ so this is a contradiction. Therefore $b = 0, 2$, or 4 . By Corollary 2.27, $q_{47}\hat{\theta}_5$ is essential so $b \neq 0$. If $b = 4$ then $2(q_{47}\hat{\theta}_5) = 2b\tilde{x} = 8\tilde{x} = 0$ in $\pi_{62}^S(P_{47}^{62})$, because the order of \tilde{x} is 8 . By Proposition 2.26 this is impossible. Therefore $b = 2$, and $q_{47}\hat{\theta}_5 = 2\tilde{x}$. This proves Proposition 2.41. \square

We are almost ready to prove the main result of this section which is Theorem 2.17. In section 4 we will prove the following.

(2.42) $2\tilde{x} \in \pi_{62}^S(P_{47}^{62})$ is detected by a non-zero class $\overline{e_{48}h_0h_3^2} \in \text{Ext}_{\mathcal{A}}^{3,65}(P_{47}^{62})$.

Note that $\overline{e_{48}h_0h_3^2h_6^2} \in \text{Ext}_{\mathcal{A}}^{5,193}(P_{47}^{62})$. In section 3 we will obtain the following Ext groups results (2.43) through (2.44).

(2.43) (1) $\text{Ext}_{\mathcal{A}}^{5,193}(P^{62}) = \mathbb{Z}/2(\hat{h}_5 h_5 D_3(1)) \oplus \mathbb{Z}/2(\hat{h}_4 h_0^2 c_4) \oplus \mathbb{Z}/2(\hat{h}_4 h_1 f_3) \oplus \mathbb{Z}/2(\gamma_{61} h_7) \oplus \mathbb{Z}/2(\hat{g}_4 h_0^2)$.

(2) $\text{Ext}_{\mathcal{A}}^{s,189+s}(P_{47}^{62}) = 0$ for $0 \leq s \leq 2$.

(3) $\text{Ext}_{\mathcal{A}}^{3,192}(P_{47}^{62}) = \mathbb{Z}/2(\overline{e_{62}h_1h_6^2})$.

(4) $\text{Ext}_{\mathcal{A}}^{3,192}(P^{62}) \xrightarrow{t_*} \text{Ext}_{\mathcal{A}}^{4,193}$ is given by $t_*(\overline{e_{62}h_1h_6^2}) = h_0 h_6^3 \neq 0$.

(2.44) The map $\text{Ext}_{\mathcal{A}}^{5,193}(P^{62}) \xrightarrow{t_*} \text{Ext}_{\mathcal{A}}^{6,194}$ as defined in (2.14) is given by the following. $t_*(\hat{g}_4 h_0^2) = h_0^2 g_4 \neq 0$, and $t_*(\hat{h}_5 h_5 D_3(1)) = h_5^2 D_3(1) \neq 0$, and $t_*(\hat{h}_4 h_0^2 c_4) = t_*(\hat{h}_4 h_1 f_3) = t_*(\gamma_{61} h_7) = 0$ in $\text{Ext}_{\mathcal{A}}^{6,194}$.

(2.45) The map $\text{Ext}_{\mathcal{A}}^{5,193}(P^{62}) \xrightarrow{(q_{47})_*} \text{Ext}_{\mathcal{A}}^{5,193}(P_{47}^{62})$ is given by the following. $(q_{47})_*(\hat{h}_5 h_5 D_3(1)) = (q_{47})_*(\hat{h}_4 h_0^2 c_4) = (q_{47})_*(\hat{h}_4 h_1 f_3) = (q_{47})_*(\gamma_{61} h_7) = 0$ and $(q_{47})_*(\hat{g}_4 h_0^2) = e_{48} h_0 h_3^2 h_6^2 \neq 0$ in $\text{Ext}_{\mathcal{A}}^{5,193}(P_{47}^{62})$.

Finally, in section 5 we will prove the following result from the Ext group information in section 3.

Proposition 2.46. For the composite $S^{188} \xrightarrow{\theta_6} S^{62} \xrightarrow{\widehat{\theta}_5} P^{62}$, $AF(\widehat{\theta}_5\theta_6) \geq 5$.

Now we prove Theorem 2.17 which is restated below.

Theorem 2.17. Let $\widehat{\theta}_5 \in \pi_{62}^S(P)$ be as in (2.16). In $\pi_{188}^S(P^{62})$, $\widehat{\theta}_5\theta_6 \neq 0$ and is detected in the ASS for π_*^S by a class

$$\tau = \widehat{g}_4 h_0^2 + \varepsilon \widehat{h}_5 h_5 D_3(1) + \overline{\tau} \neq 0 \quad \text{in } \text{Ext}_{\mathcal{A}}^{5,193}(P^{62})$$

where $\varepsilon = 0$ or 1 . Here $\overline{\tau} \in \text{Ext}_{\mathcal{A}}^{5,193}(P^{62})$ is a class such that $t_*(\overline{\tau}) = 0$ in $\text{Ext}_{\mathcal{A}}^{6,194}$. Moreover, $t_*(\widehat{g}_4 h_0^2) = h_0^2 g_4$, and $t_*(\widehat{h}_5 h_5 D_3(1)) = h_5^2 D_3(1)$.

Proof. By (2.41) and (2.42), $q_{47}\widehat{\theta}_5 \in \pi_{62}^S(P_{47}^{62})$ is detected by $\overline{e_{48}h_0h_3^2}$ in the ASS for $\pi_*^S(P_{47}^{62})$. By (2.45), $e_{48}h_0h_3^2h_6^2 = (q_{47})_*(\widehat{g}_4h_0^2) \neq 0$ in $\text{Ext}_{\mathcal{A}}^{5,193}(P_{47}^{62})$. $(q_{47})_*(\widehat{g}_4h_0^2)$ is not a boundary in the ASS for $\pi_*^S(P_{47}^{62})$ by (2.43) (2), (3), (4), by $t_*(\widehat{g}_4h_0^2) = h_0^2g_4 \neq 0$ in (2.44), and by the differential $d_2(h_0h_6^3) = h_0^2h_5^2h_6^2 = 0$ in the ASS for π_*^S . Therefore $q_{47}\widehat{\theta}_5\theta_6 \in \pi_{188}^S(P_{47}^{62})$ is non-zero and is detected by $(q_{47})_*(\widehat{g}_4h_0^2)$. It follows that $\widehat{\theta}_5\theta_6 \neq 0$ and $AF(\widehat{\theta}_5\theta_6) \leq AF(q_{47}\widehat{\theta}_5\theta_6) = 5$. By (2.46), $AF(\widehat{\theta}_5\theta_6) \geq 5$. So $AF(\widehat{\theta}_5\theta_6) = 5$. From this and (2.43) (1) we infer that $\widehat{\theta}_5\theta_6$ is detected by

$$\tau = \varepsilon_1 \widehat{g}_4 h_0^2 + \varepsilon_2 \widehat{h}_5 h_5 D_3(1) + \varepsilon_3 \widehat{h}_4 h_0^2 c_4 + \varepsilon_4 \widehat{h}_4 h_1 f_3 + \varepsilon_5 \gamma_{61} h_7$$

for some ε_i where $\varepsilon_i = 0$ or 1 for $1 \leq i \leq 5$ and $(\varepsilon_1, \varepsilon_2, \varepsilon_3, \varepsilon_4, \varepsilon_5) \neq (0, 0, 0, 0, 0)$. From (2.42) and (2.45) we see $\varepsilon_1 = 1$. So let $\overline{\tau} = \varepsilon_3 \widehat{h}_4 h_0^2 c_4 + \varepsilon_4 \widehat{h}_4 h_1 f_3 + \varepsilon_5 \gamma_{61} h_7$, and

$$\tau = \widehat{g}_4 h_0^2 + \varepsilon_2 \widehat{h}_5 h_5 D_3(1) + \overline{\tau} \in \text{Ext}_{\mathcal{A}}^{5,193}(P^{62}).$$

By (2.44), we see $t_*(\widehat{g}_4 h_0^2) = h_0^2 g_4$, $t_*(\widehat{h}_5 h_5 D_3(1)) = h_5^2 D_3(1)$, and $t_*(\overline{\tau}) = 0$ in $\text{Ext}_{\mathcal{A}}^{6,194}$. This proves Theorem 2.17. \square

From Remark 2.3 we see Theorem 2.17 implies the following theorem.

Theorem 2.2. $\theta_5\theta_6$ in π_{188}^S is detected by $h_0^2 g_4 + \varepsilon h_5^2 D_3(1) \neq 0$ in $\text{Ext}_{\mathcal{A}}^{6,194}$, where $\varepsilon = 0$ or 1 .

Finally we prove Theorem 1.4, which is restated below.

Theorem 1.4. $h_0^3 g_4$ is a boundary in the ASS for spheres.

Proof. Recall [16] that there is a relation $h_0 D_3(1) = 0$ in $\text{Ext}_{\mathcal{A}}^{5,131}$. Let $h_0^2 g_4 + \varepsilon h_5^2 D_3(1) \in \text{Ext}_{\mathcal{A}}^{6,194}$ be the class that detects $\theta_5\theta_6 \in \pi_{188}^S$ in Theorem 2.2, and we have

$$h_0^3 g_4 = h_0(h_0^2 g_4 + \varepsilon h_5^2 D_3(1)),$$

where $h_0^3 g_4 \neq 0$ in $\text{Ext}_{\mathcal{A}}^{7,195}$ by (1.3). Since h_0 detects 2ι , and since $h_0^2 g_4 + \varepsilon h_5^2 D_3(1)$ detects $\theta_5\theta_6$ by Theorem 2.2, $h_0^3 g_4 = h_0(h_0^2 g_4 + \varepsilon h_5^2 D_3(1))$ must be a boundary because $2\theta_5\theta_6 = 0$. This proves Theorem 1.4. \square

3 Calculations of some Ext groups

In this section we will describe and compute the Ext group results that we already have used in section 1, section 2, and will also be needed in section 5. There are many results, and we organize them as follows. In (3.1) and (3.2) we list respectively all the results about $\text{Ext}_{\mathcal{A}}^{*,*}$ and $\text{Ext}_{\mathcal{A}}^{*,*}(P)$ that we need, and in (3.3) are the results about $\text{Ext}_{\mathcal{A}}^{*,*}(P^{62})$. In (3.4) are the results about $\text{Ext}_{\mathcal{A}}^{*,*}(P_k^l)$ for some stunted projective spaces $P_k^l = P^l/P^{k-1}$. Recall the complex $\tilde{P}^{62} = P^{62} \cup_{h_5} C(S^{30} \cup_{2i} e^{31})$ in (2.29). The subcomplex $P^{16} \subset P^{62}$ is also a subcomplex of \tilde{P}^{62} . Let $\tilde{P}_{17}^{62} = \tilde{P}^{62}/P^{16}$. In (3.5) and (3.6) are the results about $\text{Ext}_{\mathcal{A}}^{*,*}(\tilde{P}^{62})$ and $\text{Ext}_{\mathcal{A}}^{*,*}(\tilde{P}_{17}^{62})$ respectively. These are all the Ext group results that we will compute in this section.

The cohomology classes that appear in (3.1) and (3.2), except $\beta_{187} \in \text{Ext}_{\mathcal{A}}^{5,192}(P)$ in (3.2) (8), are as defined in [8, 16], and these will be recalled in a moment: the classes $h_i, c_i, f_i, g_{i+1}, D_3(i), V_i$ for $i \geq 0$ in (3.1) and (3.2) will be recalled in (3.13), and the classes $\hat{h}_i, \hat{g}_{i+1}, \gamma_{61}(i), \widehat{D}_3(i)$ for $i \geq 0$ in (3.2) will be recalled in (3.21). The class β_{187} will be defined in (3.23). We will simply write D_3 for $D_3(0)$, γ_{61} for $\gamma_{61}(0)$, and \widehat{D}_3 for $\widehat{D}_3(0)$.

- (3.1)** (1) $\text{Ext}_{\mathcal{A}}^{4,192} = \mathbb{Z}/2(g_4)$.
(2) $\text{Ext}_{\mathcal{A}}^{5,193} = \mathbb{Z}/2(h_0 g_4)$.
(3) $\text{Ext}_{\mathcal{A}}^{s,s+189} = 0$ for $0 \leq s \leq 2$.
(4) $\text{Ext}_{\mathcal{A}}^{s,s+189} = \mathbb{Z}/2(h_0^{s-3} h_6^3)$, $3 \leq s \leq 4$.
(5) $\text{Ext}_{\mathcal{A}}^{5,194} = \mathbb{Z}/2(h_0^2 h_6^3) \oplus \mathbb{Z}/2(h_1 g_4)$.
(6) $\text{Ext}_{\mathcal{A}}^{s,s+190} = 0$ for $0 \leq s \leq 2$.
(7) $\text{Ext}_{\mathcal{A}}^{6,193} = \mathbb{Z}/2(h_5 V_0)$.
(8) $\text{Ext}_{\mathcal{A}}^{6,194} = \mathbb{Z}/2(h_0^2 g_4) \oplus \mathbb{Z}/2(h_5^2 D_3(1))$.
(9) $h_0^3 g_4, h_1 h_5 V_0$ are linearly independent in $\text{Ext}_{\mathcal{A}}^{7,195}$.

Recall the map $P \xrightarrow{t} S^0$ and the induced homomorphism $\text{Ext}_{\mathcal{A}}^{s,t}(P) \xrightarrow{t_*} \text{Ext}_{\mathcal{A}}^{s+1,t+1}$ mentioned in (2.14). (3.2) (j) (a) for $1 \leq j \leq 9$ below describe $\text{Ext}_{\mathcal{A}}^{s,t}(P)$ for some particular (s, t) . (3.2) (j) (b) for $j = 1, 5, 8, 9$, (3.2) (j) (c) for $2 \leq j \leq 9$ and (3.2) (8) (d) describe $t_*(x) \in \text{Ext}_{\mathcal{A}}^{s+1,t+1}$ for the classes x in (3.2) (j) (a) for $1 \leq j \leq 9$.

- (3.2)** (1) (a) $\text{Ext}_{\mathcal{A}}^{2,48}(P) = \mathbb{Z}/2(\hat{h}_4 h_0 h_5)$,
(b) $t_*(\hat{h}_4 h_0 h_5) = 0$ in $\text{Ext}_{\mathcal{A}}^{3,49}$.
(2) (a) $\text{Ext}_{\mathcal{A}}^{3,64}(P) = \mathbb{Z}/2(\widehat{D}_3)$,
(c) $t_*(\widehat{D}_3) = D_3 \neq 0$ in $\text{Ext}_{\mathcal{A}}^{4,65}$.
(3) (a) $\text{Ext}_{\mathcal{A}}^{2,64}(P) = \mathbb{Z}/2(\hat{h}_5 h_0 h_5)$,
(c) $t_*(\hat{h}_5 h_0 h_5) = h_0 h_5^2 \neq 0$ in $\text{Ext}_{\mathcal{A}}^{3,65}$.
(4) (a) $\text{Ext}_{\mathcal{A}}^{3,191}(P) = \mathbb{Z}/2(\hat{g}_4)$,
(c) $t_*(\hat{g}_4) = g_4 \neq 0$ in $\text{Ext}_{\mathcal{A}}^{4,192}$.
(5) (a) $\text{Ext}_{\mathcal{A}}^{4,192}(P) = \mathbb{Z}/2(\hat{h}_4 h_0 c_4) \oplus \mathbb{Z}/2(\widehat{D}_3 h_7) \oplus \mathbb{Z}/2(\hat{g}_4 h_0) \oplus \mathbb{Z}/2(\hat{h}_7 D_3)$,

- (b) $t_*(\widehat{h}_4 h_0 c_4) = 0$ in $\text{Ext}_{\mathcal{A}}^{5,193}$,
(c) $t_*(\widehat{D}_3 h_7) = t_*(\widehat{h}_7 D_3) = D_3 h_7$, $t_*(\widehat{g}_4 h_0) = h_0 g_4$, and $D_3 h_7, h_0 g_4$ are linearly independent in $\text{Ext}_{\mathcal{A}}^{5,193}$.
- (6) (a) $\text{Ext}_{\mathcal{A}}^{2,191}(P) = \mathbb{Z}/2(\widehat{h}_5 h_5 h_7)$,
(c) $t_*(\widehat{h}_5 h_5 h_7) = h_5^2 h_7 = h_6^3 \neq 0$ in $\text{Ext}_{\mathcal{A}}^{3,192}$.
- (7) (a) $\text{Ext}_{\mathcal{A}}^{3,192}(P) = \mathbb{Z}/2(\widehat{h}_5 h_0 h_5 h_7) \oplus \mathbb{Z}/2(\widehat{h}_6 h_0 h_6^2)$,
(c) $t_*(\widehat{h}_5 h_0 h_5 h_7) = t_*(\widehat{h}_6 h_0 h_6^2) = h_0 h_6^3 \neq 0$ in $\text{Ext}_{\mathcal{A}}^{4,193}$.
- (8) (a) $\text{Ext}_{\mathcal{A}}^{5,192}(P) = \mathbb{Z}/2(\beta_{187}) \oplus \mathbb{Z}/2(\widehat{h}_4 h_0 f_3) \oplus \mathbb{Z}/2(\widehat{h}_5 V_0)$,
(b) $t_*(\widehat{h}_4 h_0 f_3) = 0$ in $\text{Ext}_{\mathcal{A}}^{6,193}$,
(c) $t_*(\widehat{h}_5 V_0) = h_5 V_0 \neq 0$ in $\text{Ext}_{\mathcal{A}}^{6,193}$,
(d) $t_*(\beta_{187}) = h_5 V_0 \neq 0$ in $\text{Ext}_{\mathcal{A}}^{6,193}$.
- (9) (a) $\text{Ext}_{\mathcal{A}}^{5,193}(P) = \mathbb{Z}/2(\widehat{h}_5 h_5 D_3(1)) \oplus \mathbb{Z}/2(\widehat{h}_4 h_0^2 c_4) \oplus \mathbb{Z}/2(\widehat{h}_4 h_1 f_3) \oplus \mathbb{Z}/2(\gamma_{61} h_7) \oplus \mathbb{Z}/2(\widehat{g}_4 h_0^2)$,
(b) $t_*(\widehat{h}_4 h_0^2 c_4) = t_*(\widehat{h}_4 h_1 f_3) = t_*(\gamma_{61} h_7) = 0$ in $\text{Ext}_{\mathcal{A}}^{6,194}$,
(c) $t_*(\widehat{h}_5 h_5 D_3(1)) = h_5^2 D_3(1)$, $t_*(\widehat{g}_4 h_0^2) = h_0^2 g_4$, and $h_5^2 D_3(1), h_0^2 g_4$ are linearly independent in $\text{Ext}_{\mathcal{A}}^{6,194}$.

Let $P^{62} \xrightarrow{i_{62}} P$ be the inclusion map, and $\text{Ext}_{\mathcal{A}}^{s,t}(P^{62}) \xrightarrow{(i_{62})_*} \text{Ext}_{\mathcal{A}}^{s,t}(P)$, $s, t \geq 0$, be the induced homomorphisms. (3.3) (i) (j) for $1 \leq j \leq 9$ below describe the groups $\text{Ext}_{\mathcal{A}}^{s,t}(P^{62})$ for some $s, t \geq 0$. Each of the classes in $\text{Ext}_{\mathcal{A}}^{s,t}(P^{62})$ in (3.3) (i) (j) is a pullback through $(i_{62})_*$ of the class of the same name in $\text{Ext}_{\mathcal{A}}^{s,t}(P)$ given in (3.2) (j) (a) for $1 \leq j \leq 9$, except the following two classes: $\overline{e_{62} h_1 h_6^2} \in \text{Ext}_{\mathcal{A}}^{3,192}(P^{62})$ in (3.3) (i) (7) is a pullback of $\widehat{h}_6 h_0 h_6^2 \in \text{Ext}_{\mathcal{A}}^{3,192}(P)$ in (3.2) (7), and $\overline{e_{62} D_3(1)} \in \text{Ext}_{\mathcal{A}}^{4,192}(P^{62})$ in (3.3) (i) (5) is a pullback of $\widehat{h}_7 D_3 \in \text{Ext}_{\mathcal{A}}^{4,192}(P)$ in (3.2) (5). How to take these pullbacks will be explained in (3.46).

- (3.3)** (i) (1) $\text{Ext}_{\mathcal{A}}^{2,48}(P^{62}) = \mathbb{Z}/2(\widehat{h}_4 h_0 h_5)$,
(2) $\text{Ext}_{\mathcal{A}}^{3,64}(P^{62}) = \mathbb{Z}/2(\widehat{D}_3)$,
(3) $\text{Ext}_{\mathcal{A}}^{2,64}(P^{62}) = \mathbb{Z}/2(\widehat{h}_5 h_0 h_5)$,
(4) $\text{Ext}_{\mathcal{A}}^{3,191}(P^{62}) = \mathbb{Z}/2(\widehat{g}_4)$,
(5) $\text{Ext}_{\mathcal{A}}^{4,192}(P^{62}) = \mathbb{Z}/2(\widehat{h}_4 h_0 c_4) \oplus \mathbb{Z}/2(\widehat{D}_3 h_7) \oplus \mathbb{Z}/2(\widehat{g}_4 h_0) \oplus \mathbb{Z}/2(\overline{e_{62} D_3(1)})$,
(6) $\text{Ext}_{\mathcal{A}}^{2,191}(P^{62}) = \mathbb{Z}/2(\widehat{h}_5 h_5 h_7)$,
(7) $\text{Ext}_{\mathcal{A}}^{3,192}(P^{62}) = \mathbb{Z}/2(\widehat{h}_5 h_0 h_5 h_7) \oplus \mathbb{Z}/2(\overline{e_{62} h_1 h_6^2})$,
(8) $\text{Ext}_{\mathcal{A}}^{5,192}(P^{62}) = \mathbb{Z}/2(\beta_{187}) \oplus \mathbb{Z}/2(\widehat{h}_4 h_0 f_3) \oplus \mathbb{Z}/2(\widehat{h}_5 V_0)$,
(9) $\text{Ext}_{\mathcal{A}}^{5,193}(P^{62}) = \mathbb{Z}/2(\widehat{h}_5 h_5 D_3(1)) \oplus \mathbb{Z}/2(\widehat{h}_4 h_0^2 c_4) \oplus \mathbb{Z}/2(\widehat{h}_4 h_1 f_3) \oplus \mathbb{Z}/2(\gamma_{61} h_7) \oplus \mathbb{Z}/2(\widehat{g}_4 h_0^2)$.
- (ii) $\text{Ext}_{\mathcal{A}}^{s,t}(P^{62}) \xrightarrow[\cong]{(i_{62})_*} \text{Ext}_{\mathcal{A}}^{s,t}(P)$ for all the (s, t) in (3.3) (i).
- (iii) $\text{Ext}_{\mathcal{A}}^{s,t} \xrightarrow{t_*} \text{Ext}_{\mathcal{A}}^{s+1,t+1}$ for all (s, t) in (3.3) (i) are given by (3.2) (j) (b) for $j = 1, 5, 8, 9$ and (3.2) (j) (c) for $2 \leq j \leq 9$, except $t_*(\overline{e_{62} D_3(1)}) = D_3 h_7 \neq 0$ and $t_*(\overline{e_{62} h_1 h_6^2}) = h_0 h_6^3 \neq 0$.

The cohomology classes in the following (3.4) will be denoted by their \overline{E}_2 representatives in a spectral sequence $\{\overline{E}_r^{i,s,t}\}_{r \geq 1}$, whose group of infinite cycles is $\text{Ext}_{\mathcal{A}}^{s,t+s}(P_k^l = P^l/P^{k-1})$. This spectral sequence $\{\overline{E}_r^{i,s,t}\}_{r \geq 1}$ will be introduced in (3.45). The E_2 term of $\{\overline{E}_r^{i,s,t}\}_{r \geq 1}$ is generated over $\mathbb{Z}/2$ by elements of the form $e_i \alpha$, $\alpha \in \text{Ext}_{\mathcal{A}}^{s,t+s}$ (α could be $1 \in \text{Ext}_{\mathcal{A}}^{0,0} = \mathbb{Z}/2$) and e_i is a $\mathbb{Z}/2$ -basis element in $H_*(P_k^l)$. A permanent cycle represented by $e_i \alpha$ is denoted by $\overline{e_i \alpha}$, and is a non-zero class in $\text{Ext}_{\mathcal{A}}^{s,t+s+i}(P_k^l)$. The cohomology classes in (3.4) will be defined in (3.48).

- (3.4)** (1) $\text{Ext}_{\mathcal{A}}^{1,63}(P_{17}^{62}) = \mathbb{Z}/2(\overline{e_{31} h_5})$.
(2) $\text{Ext}_{\mathcal{A}}^{2,32}(P_{17}^{62}) = \mathbb{Z}/2(\overline{e_{23} h_0 h_3})$.
(3) $\text{Ext}_{\mathcal{A}}^{3,64}(P_{17}^{62}) = \mathbb{Z}/2(\overline{e_{23} h_0 h_3 h_5})$,
 $\text{Ext}_{\mathcal{A}}^{3,64}(P_{17}^{62}) \xrightarrow{(q_{17})_*} \text{Ext}_{\mathcal{A}}^{3,64}(P_{17}^{62})$ is given by $(q_{17})_*(\widehat{D}_3) = \overline{e_{23} h_0 h_3 h_5}$.
(4) $\text{Ext}_{\mathcal{A}}^{0,47}(P_{33}^{62}) = \mathbb{Z}/2(\overline{e_{47}})$.
(5) $\text{Ext}_{\mathcal{A}}^{2,48}(P_{33}^{62}) = \mathbb{Z}/2(\overline{e_{39} h_0 h_3})$.
(6) $\text{Ext}_{\mathcal{A}}^{1,31}(P^{16}) = \mathbb{Z}/2(\overline{e_{15} h_4})$.
(7) $\text{Ext}_{\mathcal{A}}^{2,16}(P^{16}) = \mathbb{Z}/2(\overline{e_7 h_0 h_3})$.
(8) $\text{Ext}_{\mathcal{A}}^{3,32}(P^{16}) = \mathbb{Z}/2(\overline{e_{15} h_0 h_3^2})$.
(9) $\text{Ext}_{\mathcal{A}}^{0,31}(P^{32}) = \mathbb{Z}/2(\overline{e_{31}})$.
(10) $\text{Ext}_{\mathcal{A}}^{1,47}(P^{32}) = \mathbb{Z}/2(\overline{e_{31} h_4})$.
(11) $\text{Ext}_{\mathcal{A}}^{2,32}(P^{32}) = \mathbb{Z}/2(\overline{e_{15} h_0 h_4})$.
(12) $\text{Ext}_{\mathcal{A}}^{3,48}(P^{32}) = \mathbb{Z}/2(\overline{e_{31} h_0 h_3^2}) \oplus \mathbb{Z}/2(\overline{e_{15} h_0 h_4^2})$.
(13) $\text{Ext}_{\mathcal{A}}^{s,189+s}(P_{47}^{62}) = 0$ for $0 \leq s \leq 2$.
(14) $\text{Ext}_{\mathcal{A}}^{2,64}(P_{47}^{62}) = \mathbb{Z}/2(\overline{e_{54} h_1 h_3})$.
(15) $\text{Ext}_{\mathcal{A}}^{3,65}(P_{47}^{62}) = \mathbb{Z}/2(\overline{e_{48} h_0 h_3^2})$.
(16) $\text{Ext}_{\mathcal{A}}^{3,192}(P_{47}^{62}) = \mathbb{Z}/2(\overline{e_{62} h_1 h_6^2})$,
 $\text{Ext}_{\mathcal{A}}^{3,192}(P_{47}^{62}) \xrightarrow{(q_{17})_*} \text{Ext}_{\mathcal{A}}^{3,192}(P_{47}^{62})$ is given by $(q_{17})_*(\overline{e_{62} h_1 h_6^2}) = \overline{e_{62} h_1 h_6^2}$.
(17) $\text{Ext}_{\mathcal{A}}^{5,193}(P_{47}^{62}) = \mathbb{Z}/2(\overline{e_{48} h_0 h_3^2 h_6^2})$,
 $\text{Ext}_{\mathcal{A}}^{5,193}(P_{47}^{62}) \xrightarrow{(q_{47})_*} \text{Ext}_{\mathcal{A}}^{5,193}(P_{47}^{62})$ is given by $(q_{47})_*(\widehat{h_5 h_5 D_3}(1)) = (q_{47})_*(\widehat{h_4 h_0^2 c_4}) = (q_{47})_*(\widehat{h_4 h_1 f_3}) = (q_{47})_*(\gamma_{61} h_7) = 0$, and $(q_{47})_*(\widehat{g_4 h_0^2}) = \overline{e_{48} h_0 h_3^2 h_6^2} \neq 0$.

Recall the complex \widetilde{P}^{62} , the map $P^{62} \xrightarrow{\widetilde{q}} \widetilde{P}^{62}$, and the induced homomorphism $\text{Ext}_{\mathcal{A}}^{*,*}(P^{62}) \xrightarrow{\widetilde{q}_*} \text{Ext}_{\mathcal{A}}^{*,*}(\widetilde{P}^{62})$ in (2.29). In (3.5) below the cohomology classes are all in the image of $\text{Ext}_{\mathcal{A}}^{*,*}(P^{62}) \xrightarrow{\widetilde{q}_*} \text{Ext}_{\mathcal{A}}^{*,*}(\widetilde{P}^{62})$ except those generated (using the obvious $\text{Ext}_{\mathcal{A}}^{*,*}$ -action) by the following four classes: $\overline{e_{31} h_5} \in \text{Ext}_{\mathcal{A}}^{1,63}(\widetilde{P}^{62})$, $\overline{e_{47}} \in \text{Ext}_{\mathcal{A}}^{0,47}(\widetilde{P}^{62})$, and $\overline{e_{39} h_3} \in \text{Ext}_{\mathcal{A}}^{2,48}(\widetilde{P}^{62})$, and $\overline{e_{31} h_4} \in \text{Ext}_{\mathcal{A}}^{1,31}(\widetilde{P}^{62})$. These four new classes will be defined in (3.52) later.

- (3.5)** (1) $\text{Ext}_{\mathcal{A}}^{0,47}(\widetilde{P}^{62}) = \mathbb{Z}/2(\overline{e_{47}})$.

- (2) $\text{Ext}_{\mathcal{A}}^{2,48}(\tilde{P}^{62}) = \mathbb{Z}/2(\overline{e_{39}h_3h_0}) \oplus \mathbb{Z}/2(\tilde{q}_*(\widehat{h_4h_0h_5}))$.
- (3) $\text{Ext}_{\mathcal{A}}^{3,64}(\tilde{P}^{62}) = \mathbb{Z}/2(\overline{\tilde{e}_{31}h_4^2h_0}) \oplus \mathbb{Z}/2(\tilde{q}_*(\widehat{D_3}))$,
 $\overline{e_{39}h_3h_0h_4} = \overline{\tilde{e}_{31}h_4^2h_0} + \tilde{q}_*(\widehat{D_3})$ in $\text{Ext}_{\mathcal{A}}^{3,64}(\tilde{P}^{62})$.
- (4) $\text{Ext}_{\mathcal{A}}^{1,63}(\tilde{P}^{62}) = \mathbb{Z}/2(\overline{\tilde{e}_{31}h_5}) \oplus \mathbb{Z}/2(\overline{e_{47}h_4})$.
- (5) $\text{Ext}_{\mathcal{A}}^{2,64}(\tilde{P}^{62}) = \mathbb{Z}/2(\overline{\tilde{e}_{31}h_5h_0}) \oplus \mathbb{Z}/2(\overline{e_{47}h_0h_4})$.
- (6) $\text{Ext}_{\mathcal{A}}^{4,192}(\tilde{P}^{62}) = \mathbb{Z}/2(\tilde{q}_*(\widehat{h_4h_0c_4})) \oplus \mathbb{Z}/2(\tilde{q}_*(\widehat{D_3}h_7)) \oplus \mathbb{Z}/2(\tilde{q}_*(\widehat{g_4h_0})) \oplus$
 $\mathbb{Z}/2(\tilde{q}_*(\overline{e_{62}D_3(1)})) \oplus \mathbb{Z}/2(\overline{\tilde{e}_{31}h_4^2h_0h_7})$.
- (7) We have the following results on $\text{Ext}_{\mathcal{A}}^{5,193}(\tilde{P}^{62})$ (see (3.3) (i) (9)):
- (a) $\tilde{q}_*(\widehat{h_5h_5D_3(1)}) = 0$ in $\text{Ext}_{\mathcal{A}}^{5,193}(\tilde{P}^{62})$,
- (b) $\tilde{q}_*(\widehat{h_4h_0^2c_4})$, $\tilde{q}_*(\widehat{h_4h_1f_3})$, $\tilde{q}_*(\gamma_{61}h_7)$, $\tilde{q}_*(\widehat{g_4h_0^2})$, $\overline{e_{31}h_4^2h_0^2h_7}$ are linearly independent in $\text{Ext}_{\mathcal{A}}^{5,193}(\tilde{P}^{62})$.
- (8) $\text{Ext}_{\mathcal{A}}^{0,189}(\tilde{P}^{62}) = 0$.
- (9) $\text{Ext}_{\mathcal{A}}^{1,190}(\tilde{P}^{62}) = 0$.
- (10) $\text{Ext}_{\mathcal{A}}^{2,191}(\tilde{P}^{62}) = \mathbb{Z}/2(\overline{\tilde{e}_{31}h_5h_7}) \oplus \mathbb{Z}/2(\overline{e_{47}h_4h_7})$.
- (11) $\text{Ext}_{\mathcal{A}}^{3,192}(\tilde{P}^{62}) = \mathbb{Z}/2(\overline{\tilde{e}_{31}h_5h_0h_7}) \oplus \mathbb{Z}/2(\overline{e_{47}h_4h_7h_0}) \oplus \mathbb{Z}/2(\tilde{q}_*(\overline{e_{62}h_1h_6^2}))$.

For each k with $1 \leq k \leq 62$ let \tilde{P}^k be the k -skeleton of \tilde{P}^{62} , and let $\tilde{P}_k^{62} = \tilde{P}^{62}/\tilde{P}^{k-1}$. Then $\tilde{P}^k = P^k$ and $\tilde{P}_k^{62} = \tilde{P}^{62}/P^k$ for $1 \leq k \leq 29$, and $\tilde{P}_k^{62} = P_k^{62}$ for $33 \leq k \leq 62$. Let $\tilde{P}^{62} \xrightarrow{q'_k} \tilde{P}_k^{62}$ be the pinching map. Then $P^{62} \xrightarrow{\tilde{q}} \tilde{P}^{62}$ induces a map $P_k^{62} \xrightarrow{\tilde{q}_k} \tilde{P}_k^{62}$ with the following commutative diagram.

$$\begin{array}{ccc} P^{62} & \xrightarrow{\tilde{q}} & \tilde{P}^{62} \\ \downarrow q_k & & \downarrow q'_k \\ P_k^{62} & \xrightarrow{\tilde{q}_k} & \tilde{P}_k^{62} \end{array}$$

In (3.6) (1), (2), (3) below we describe $\text{Ext}_{\mathcal{A}}^{s,t}(\tilde{P}_{17}^{62})$ for $(s, t) = (1, 63), (2, 32), (3, 64)$ respectively. For the classes in the image of $\text{Ext}_{\mathcal{A}}^{*,*}(P_{17}^{62}) \xrightarrow{(\tilde{q}_{17})_*} \text{Ext}_{\mathcal{A}}^{*,*}(\tilde{P}_{17}^{62})$ we denote them as such in (3.6) (1) through (3.6) (3). The classes generated by $\overline{\tilde{e}_{31}} \in \text{Ext}_{\mathcal{A}}^{0,31}(\tilde{P}_{17}^{62})$ in (3.6) (1) and (3.6) (3), and by $\overline{e_{47}} \in \text{Ext}_{\mathcal{A}}^{0,47}(\tilde{P}_{17}^{62})$ in (3.6) (1), are not in the image of $(\tilde{q}_{17})_*$. These classes will be defined in (3.60) later. We also describe the map $\text{Ext}_{\mathcal{A}}^{*,*}(\tilde{P}^{62}) \xrightarrow{(q'_{17})_*} \text{Ext}_{\mathcal{A}}^{*,*}(\tilde{P}_{17}^{62})$ at the dimensions specified in (3.6) (1) through (3.6) (3). For $k \geq 33$, $\tilde{P}_k^{62} = P_k^{62}$. In (3.6) (4), (5), (6) below we describe the map $\text{Ext}_{\mathcal{A}}^{*,*}(\tilde{P}^{62}) \xrightarrow{(q'_k)_*} \text{Ext}_{\mathcal{A}}^{*,*}(\tilde{P}_k^{62} = P_k^{62})$ for $(s, t, k) = (2, 48, 33), (0, 47, 33), (2, 64, 47)$ respectively. We refer to (3.4) and (3.5) for the classes in these descriptions.

- (3.6)** (1) $\text{Ext}_{\mathcal{A}}^{1,63}(\tilde{P}_{17}^{62}) = \mathbb{Z}/2(\overline{\tilde{e}_{31}h_5}) \oplus \mathbb{Z}/2(\overline{e_{47}h_4})$,
 $\text{Ext}_{\mathcal{A}}^{1,63}(\tilde{P}^{62}) \xrightarrow{(q'_{17})_*} \text{Ext}_{\mathcal{A}}^{1,63}(\tilde{P}_{17}^{62})$ is given by $(q'_{17})_*(\overline{e_{31}h_5}) = \overline{e_{31}h_5}$
and $(q'_{17})_*(\overline{e_{47}h_4}) = \overline{e_{47}h_4}$.
- (2) $\text{Ext}_{\mathcal{A}}^{2,32}(\tilde{P}_{17}^{62}) = \mathbb{Z}/2((\tilde{q}_{17})_*(\overline{e_{23}h_0h_3}))$.

- (3) $\text{Ext}_{\mathcal{A}}^{3,64}(\tilde{P}_{17}^{62}) = \mathbb{Z}/2(\overline{e_{31}h_0h_4^2}) \oplus \mathbb{Z}/2((\tilde{q}_{17})_*(\overline{e_{23}h_0h_3h_5}))$,
 $\text{Ext}_{\mathcal{A}}^{3,64}(\tilde{P}^{62}) \xrightarrow{(q'_{17})_*} \text{Ext}_{\mathcal{A}}^{3,64}(\tilde{P}_{17}^{62})$ is given by $(q'_{17})_*(\overline{e_{31}h_4^2h_0}) = \overline{e_{31}h_0h_4^2}$,
and
 $(q'_{17})_*(\tilde{q}_*(\widehat{D}_3)) = (\tilde{q}_{17})_*(\overline{e_{23}h_0h_3h_5})$.
- (4) $\text{Ext}_{\mathcal{A}}^{2,48}(\tilde{P}^{62}) \xrightarrow{(q'_{33})_*} \text{Ext}_{\mathcal{A}}^{2,48}(\tilde{P}_{33}^{62} = P_{33}^{62})$ is given by $(q'_{33})_*(\overline{e_{39}h_3h_0}) = \overline{e_{39}h_3h_0}$, and $(q'_{33})_*(\tilde{q}_*(\widehat{h}_4h_0h_5)) = 0$.
- (5) $\text{Ext}_{\mathcal{A}}^{0,47}(\tilde{P}^{62}) \xrightarrow{(q'_{33})_*} \text{Ext}_{\mathcal{A}}^{0,47}(\tilde{P}_{33}^{62} = P_{33}^{62})$ is given by $(q'_{33})_*(\overline{e_{47}}) = \overline{e_{47}}$.
- (6) $\text{Ext}_{\mathcal{A}}^{2,64}(\tilde{P}^{62}) \xrightarrow{(q'_{47})_*} \text{Ext}_{\mathcal{A}}^{2,64}(\tilde{P}_{47}^{62} = P_{47}^{62})$ is given by $(q'_{47})_*(\overline{e_{31}h_5h_5}) = \overline{e_{31}h_5h_5}$ and $(q'_{47})_*(\overline{e_{47}h_0h_4}) = 0$.

This completes the statements of all the Ext groups results in this section.

Remark 3.7. As will be seen later, the results in (3.2) (8) (c), (3.2) (9) (c) on $\text{Ext}_{\mathcal{A}}^{*,*}(P)$ imply the results in (3.1) (7), (8) respectively.

In the first half of this section we will describe and prove (3.1) and (3.2), and in the second half we will describe and prove (3.3) through (3.6).

(3.1) (1) through (3.1) (6) follow from some well-known results on $\text{Ext}_{\mathcal{A}}^{*,*}$ obtained in [2, 8, 16]. These known results are summarized in Theorem 3.14 below. In order to describe the cohomology classes in (3.1) and to state Theorem 3.14, we need to recall our main tool for calculating the Ext groups in (3.1), which is the mod 2 lambda algebra Λ ([6]). Recall that Λ is the bigraded differential algebra generated by $\lambda_i \in \Lambda^{1,i}$ for $i \geq 0$ with relation

$$(3.8) \quad \lambda_i \lambda_{2i+1+m} = \sum_{\nu \geq 0} \binom{m-1-\nu}{\nu} \lambda_{m+i-\nu} \lambda_{2i+1+\nu} \quad \text{for } m \geq 0.$$

The differential $\Lambda \xrightarrow{\delta} \Lambda$ is given by

$$(3.9) \quad \delta(\lambda_i) = \sum_{\nu \geq 0} \binom{i-1-\nu}{\nu+1} \lambda_{i-1-\nu} \lambda_{\nu} \quad \text{on the generators } \lambda_i.$$

Then

$$(3.10) \quad H^{s,t}(\Lambda) = H^{s,t}(\Lambda, \delta) = \text{Ext}_{\mathcal{A}}^{s,t+s}.$$

A monomial $\lambda_{i_1} \cdots \lambda_{i_s} \in \Lambda$ will simply be denoted by λ_I where $I = (i_1, \dots, i_s)$ is a sequence of non-negative integers.

(3.11) From (3.8) we see that the set of monomials $\lambda_I = \lambda_{i_1} \cdots \lambda_{i_s}$ with $2i_j \geq i_{j+1}$, $1 \leq j \leq s-1$, $s \geq 0$ ($\lambda_I = 1$ if $s = 0$), is a basis for Λ . This basis is called the admissible basis and the monomials in this basis are said to be admissible monomials.

There is an operation $\Lambda \xrightarrow{Sq^0} \Lambda$ given by

$$(3.12) \quad Sq^0(\lambda_{j_1} \cdots \lambda_{j_s}) = \lambda_{2j_1+1} \cdots \lambda_{2j_s+1},$$

where $\lambda_{j_1} \cdots \lambda_{j_s}$ is not necessarily admissible. This operation is compatible with the relations (3.8) and commutes with the differential (3.9), so it induces an Ext groups map

$$H^{s,t}(\Lambda) = \text{Ext}_{\mathcal{A}}^{s,t+s} \xrightarrow{Sq^0} H^{s,2t+s}(\Lambda) = \text{Ext}_{\mathcal{A}}^{s,2t+2s}.$$

Let $\text{Ext}_{\mathcal{A}}^{*,*} \xrightarrow{(Sq^0)^i} \text{Ext}_{\mathcal{A}}^{*,*}$ denote the composite $\underbrace{Sq^0 \cdots Sq^0}_i$ for $i \geq 1$, and let

$\text{Ext}_{\mathcal{A}}^{*,*} \xrightarrow{(Sq^0)^0} \text{Ext}_{\mathcal{A}}^{*,*}$ be the identity map.

For the following cohomology classes we refer to [8, 16].

- (3.13)**
- (1) $h_i = \{\lambda_{2^{i-1}} = (Sq^0)^i(\lambda_0)\} \in \text{Ext}_{\mathcal{A}}^{1,2^i}, i \geq 0.$
 - (2) $c_i = \{(Sq^0)^i(c_0^* = \lambda_2 \lambda_3^2)\} \in \text{Ext}_{\mathcal{A}}^{3,2^{i+3}+2^{i+1}+2^i}, i \geq 0.$
 - (3) $d_i = \{(Sq^0)^i(d_0^* = \lambda_6 \lambda_2 \lambda_3^2 + \lambda_4^2 \lambda_3^2 + \lambda_2 \lambda_4 \lambda_5 \lambda_3)\} \in \text{Ext}_{\mathcal{A}}^{4,2^{i+4}+2^{i+1}}, i \geq 0.$
 - (4) $e_i = \{(Sq^0)^i(\lambda_8 \lambda_3^3 + \lambda_4(\lambda_5^2 \lambda_3 + \lambda_7 \lambda_3^2) + \lambda_2(\lambda_9 \lambda_3^2 + \lambda_3^2 \lambda_9))\} \in \text{Ext}_{\mathcal{A}}^{4,2^{i+4}+2^{i+2}+2^i}, i \geq 0.$
 - (5) $f_i = \{(Sq^0)^i(\lambda_4 \lambda_0 \lambda_7^2 + \lambda_3(\lambda_9 \lambda_3^2 + \lambda_3 \lambda_5 \lambda_7) + \lambda_2 \lambda_4 \lambda_5 \lambda_7)\} \in \text{Ext}_{\mathcal{A}}^{4,2^{i+4}+2^{i+2}+2^{i+1}}, i \geq 0.$
 - (6) $g_{i+1} = \{(Sq^0)^i(\lambda_6 \lambda_0 \lambda_7^2 + \lambda_5(\lambda_9 \lambda_3^2 + \lambda_3^2 \lambda_9) + \lambda_3(\lambda_{11} \lambda_3^2 + \lambda_5 \lambda_9 \lambda_3))\} \in \text{Ext}_{\mathcal{A}}^{4,2^{i+4}+2^{i+3}}, i \geq 0.$
 - (7) $p_i = \{(Sq^0)^i(\lambda_0(\lambda_{19} \lambda_7^2 + \lambda_7^2 \lambda_{19}))\} \in \text{Ext}_{\mathcal{A}}^{4,2^{i+5}+2^{i+2}+2^i}, i \geq 0.$
 - (8) $D_3(i) = \{(Sq^0)^i(\lambda_0 \lambda_{23} \lambda_7 \lambda_{31})\} \in \text{Ext}_{\mathcal{A}}^{4,2^{i+6}+2^i}, i \geq 0.$
 - (9) $p'_i = \{(Sq^0)^i(\lambda_0(\lambda_{39} \lambda_{15}^2 + \lambda_{15}^2 \lambda_{39}))\} \in \text{Ext}_{\mathcal{A}}^{4,2^{i+6}+2^{i+3}+2^i}, i \geq 0.$
 - (10) $P^1 h_1 = \{\lambda_2 \lambda_0^3 \lambda_7 + \lambda_1(\lambda_2 \lambda_4 \lambda_1^2 + \lambda_1 \lambda_2 \lambda_4 \lambda_1 + \lambda_2 \lambda_1^2 \lambda_4)\} \in \text{Ext}_{\mathcal{A}}^{5,14}.$
 - (11) $P^1 h_2 = \{\lambda_4 \lambda_0^3 \lambda_7 + \lambda_3(\lambda_5 \lambda_1^3 + \lambda_2 \lambda_4 \lambda_1^2 + \lambda_1 \lambda_2 \lambda_4 \lambda_1 + \lambda_2 \lambda_1^2 \lambda_4) + \lambda_2^2 \lambda_0^2 \lambda_7 + \lambda_1^2 \lambda_2 \lambda_0 \lambda_7\} \in \text{Ext}_{\mathcal{A}}^{5,16}.$
 - (12) $n_i = \{(Sq^0)^i(\lambda_9 \lambda_3 \lambda_5 \lambda_7^2 + \lambda_6 \lambda_0 \lambda_3 \lambda_{15} \lambda_7 + \lambda_3 \lambda_5 \lambda_1 \lambda_{15} \lambda_7)\} \in \text{Ext}_{\mathcal{A}}^{5,2^{i+5}+2^{i+2}}, i \geq 0.$
 - (13) $x_i = \{(Sq^0)^i(\lambda_{14} \lambda_2 \lambda_3^2 \lambda_{15} + \lambda_{12} \lambda_4 \lambda_7^3 + \lambda_8^2 \lambda_7^3 + \lambda_6 \lambda_2 \lambda_3^2 \lambda_{23} + \lambda_4^2 \lambda_3^2 \lambda_{23} + \lambda_2 \lambda_4 \lambda_1 \lambda_{15}^2)\} \in \text{Ext}_{\mathcal{A}}^{5,2^{i+5}+2^{i+3}+2^{i+1}}, i \geq 0.$
 - (14) $D_1(i) = \{(Sq^0)^i(\lambda_4 \lambda_7 \lambda_{11} \lambda_{15}^2)\} \in \text{Ext}_{\mathcal{A}}^{5,2^{i+5}+2^{i+4}+2^{i+3}+2^i}, i \geq 0.$
 - (15) $H_1(i) = \{(Sq^0)^i(\lambda_{14} \lambda_7 \lambda_{11} \lambda_{15}^2 + \lambda_{10} \lambda_{11}^2 \lambda_{15}^2 + \lambda_8 \lambda_1 \lambda_7 \lambda_{31} \lambda_{15} + \lambda_6 \lambda_7 \lambda_3 \lambda_{31} \lambda_{15} + \lambda_4 \lambda_5 \lambda_7 \lambda_{31} \lambda_{15})\} \in \text{Ext}_{\mathcal{A}}^{5,2^{i+6}+2^{i+1}+2^i}, i \geq 0.$
 - (16) $Q_3(i) = \{(Sq^0)^i((\lambda_6 \lambda_0 \lambda_7^2 + \lambda_5(\lambda_9 \lambda_3^2 + \lambda_3^2 \lambda_9) + \lambda_3 \lambda_5 \lambda_9 \lambda_3 + \lambda_3 \lambda_{11} \lambda_3^2) \lambda_{47} + \lambda_5(\lambda_{21} \lambda_{11} \lambda_{15}^2 + \lambda_9 \lambda_{15} \lambda_7 \lambda_{31}) + \lambda_3 \lambda_{23} \lambda_{11} \lambda_{15}^2)\} \in \text{Ext}_{\mathcal{A}}^{5,2^{i+6}+2^{i+3}}, i \geq 0.$
 - (17) $K_i = \{(Sq^0)^i(\lambda_0^2 \lambda_{47} \lambda_{15} \lambda_{63} + (\lambda_9 \lambda_3^2 + \lambda_3^2 \lambda_9) \lambda_{47} \lambda_{63} + \lambda_{21} \lambda_{11} \lambda_{31}^3)\} \in \text{Ext}_{\mathcal{A}}^{5,2^{i+7}+2^{i+1}}, i \geq 0.$
 - (18) $J_i = \{(Sq^0)^i(\lambda_0(\lambda_{19} \lambda_7^2 + \lambda_7^2 \lambda_{19}) \lambda_{95} + \lambda_0 \lambda_{43} \lambda_{23} \lambda_{31}^2 + \lambda_0 \lambda_{19} \lambda_{31} \lambda_{15} \lambda_{63})\} \in \text{Ext}_{\mathcal{A}}^{5,2^{i+7}+2^{i+2}+2^i}, i \geq 0.$

$$\begin{aligned}
(19) \quad T_i &= \{(Sq^0)^i(\lambda_0^2(\lambda_{79}\lambda_{31}^2 + \lambda_{31}^2\lambda_{79}) + (\lambda_9\lambda_3^2 + \lambda_3^2\lambda_9)\lambda_{63}^2)\} \in \text{Ext}_{\mathcal{A}}^{5,2^{i+7}+2^{i+4}+2^{i+1}}, \\
&\quad i \geq 0. \\
(20) \quad V_i &= \{(Sq^0)^i(\lambda_0\lambda_{31}\lambda_{47}\lambda_{15}\lambda_{63})\} \in \text{Ext}_{\mathcal{A}}^{5,2^{i+7}+2^{i+5}+2^i}, \quad i \geq 0. \\
(21) \quad V'_i &= \{(Sq^0)^i(\lambda_0\lambda_{23}\lambda_7\lambda_{31}\lambda_{191} + \lambda_0\lambda_{47}\lambda_{15}\lambda_{127}\lambda_{63})\} \in \text{Ext}_{\mathcal{A}}^{5,2^{i+8}+2^i}, \\
&\quad i \geq 0. \\
(22) \quad U_i &= \{(Sq^0)^i(\lambda_0(\lambda_{39}\lambda_{15}^2 + \lambda_{15}^2\lambda_{39})\lambda_{191} + \lambda_0\lambda_{87}\lambda_{47}\lambda_{63}^2 + \lambda_0\lambda_{39}\lambda_{63}\lambda_{31}\lambda_{127})\} \in \\
&\quad \text{Ext}_{\mathcal{A}}^{5,2^{i+8}+2^{i+3}+2^i}, \quad i \geq 0.
\end{aligned}$$

Theorem 3.14 ([2, 8, 16]). (i) The algebra $\text{Ext}_{\mathcal{A}}^{s,*}$ for $s \leq 3$ is generated by $h_i \neq 0$ and $c_i \neq 0$ for $i \geq 0$, and subject only to the relations $h_i h_{i+1} = 0$, $h_i h_{i+2}^2 = 0$, and $h_{i+1}^3 = h_i^2 h_{i+2}$ for $i \geq 0$.

(ii) The subalgebra E' of the algebra $\text{Ext}_{\mathcal{A}}^{s,*}$ for $s \leq 4$ generated by h_i and c_i for $i \geq 0$ is subject only to the relations in (i) together with the relations $h_i^2 h_{i+3}^2 = 0$, and $h_j c_i = 0$ for $j = i - 1, i, i + 2$ and $i + 3$.

(iii) The set of classes $d_i, e_i, f_i, g_{i+1}, p_i, D_3(i)$ and p'_i for $i \geq 0$ is a $\mathbb{Z}/2$ -basis for the indecomposable elements in $\text{Ext}_{\mathcal{A}}^{4,*}$.

(iv) The subalgebra E'' of the algebra $\text{Ext}_{\mathcal{A}}^{s,*}$ for $s \leq 5$ generated by $h_1, c_i, d_i, e_i, f_i, g_{i+1}, p_i, D_3(i)$ and p'_i for $i \geq 0$ is subject only to the relations in (i) and (iii) together with the following relations (1) through (28), where $j \geq 0$.

$$\begin{aligned}
(1) \quad h_i^2 c_j &= 0 \text{ for } i = j + 1, j + 4, & (15) \quad h_{j+1} h_{j+4} c_j &= h_{j+3} e_j, \\
(2) \quad h_j h_{j+3} c_{j+2} &= 0, & (16) \quad h_j h_{j+4} c_{j+3} &= h_{j+5} p'_j, \\
(3) \quad h_j d_{j+1} &= 0, & (17) \quad h_{j+5}^2 c_j &= h_{j+1} p'_j, \\
(4) \quad h_i d_j &= 0 \text{ for } i = j + 3, j + 4, & (18) \quad h_j d_{j+2} &= h_{j+3} D_3(j), \\
(5) \quad h_j e_{j+1} &= 0, & (19) \quad h_{j+2} d_{j+1} &= h_{j+4} g_{j+1}, \\
(6) \quad h_{j+4} e_j &= 0, & (20) \quad h_{j+1} d_{j+1} &= h_j p_j, \\
(7) \quad h_i f_j &= 0 \text{ for } i = j + 1, j + 3, j + 4, & (21) \quad h_{j+2} d_j &= h_j e_j, \\
(8) \quad h_{j+3} g_{j+1} &= 0, & (22) \quad h_{j+1} e_j &= h_j f_j, \\
(9) \quad h_j p_{j+1} &= 0, & (23) \quad h_{j+2} e_{j+1} &= h_{j+1} f_{j+1} = \\
(10) \quad h_i p_j &= 0 \text{ for } i = j + 1, j + 2, j + & \quad h_j^2 c_{j+2}, \\
&\quad 4, j + 5, & (24) \quad h_{j+2} e_j &= h_j g_{j+1}, \\
(11) \quad h_j D_3(j + 1) &= 0, & (25) \quad h_j f_{j+2} &= h_{j+4} p'_j, \\
(12) \quad h_i D_3(j) &= 0 \text{ for } i = j, j + 5, j + & (26) \quad h_j f_{j+1} &= h_{j+3} p_j, \\
&\quad 6, & (27) \quad h_{j+2} f_j &= h_{j+1} g_{j+1}, \\
(13) \quad h_j p'_{j+1} &= 0, & (28) \quad h_{j+3} g_{j+2} &= h_{j+5} g_{j+1}. \\
(14) \quad h_i p'_j &= 0 \text{ for } i = j + 2, j + 3, j + & \\
&\quad 6, &
\end{aligned}$$

(v) The set of classes $P^1 h_1, P^1 h_2, n_i, x_i, D_1(i), H_1(i), Q_3(i), K_i, J_i, T_i, V_i, V'_i$ and U_i for $i \geq 0$ is a $\mathbb{Z}/2$ -basis for the indecomposable elements in $\text{Ext}_{\mathcal{A}}^{5,*}$.

It is not difficult to see that Theorem 3.14 covers (3.1) (1) through (3.1) (6).

(3.2) (1) through (3.2) (7) also follow from some “already known” results on $\text{Ext}_{\mathcal{A}}^{s,*}(P)$ for $s \leq 4$ in [8, 16]. These results will be summarized in Theorem 3.24 in a moment. To state these results, recall ([9]) that the lambda algebra Λ can also be used to calculate the Ext groups $\text{Ext}_{\mathcal{A}}^{*,*}(X)$ for any stable complex X , in particular for $X = P$. This is described as follows.

Given a stable complex X , the mod 2 Steenrod algebra \mathcal{A} acts on the mod 2 reduced homology group $H_*(X)$ from the right. Consider the bigraded differential module $H_*(X) \otimes \Lambda$ with bigrading $(H_*(X) \otimes \Lambda)^{s,t} = \sum_k H_k(X) \otimes \Lambda^{s,t-k}$. For $\alpha \in H_k(X)$, we write $\alpha \lambda_I$ to denote $\alpha \otimes \lambda_I \in H_*(X) \otimes \Lambda$, and write α to denote $\alpha 1 = \alpha \otimes 1 \in H_*(X) \otimes \Lambda$. Then we have the following.

(3.15) (1) $H_*(X) \otimes \Lambda$ is a differential Λ -module with differential $H_*(X) \otimes \Lambda \xrightarrow{\delta} H_*(X) \otimes \Lambda$ given by

$$\delta(\alpha \lambda_I) = \alpha \delta(\lambda_I) + \sum_{j \geq 0} \alpha S q^{j+1} \lambda_j \lambda_I.$$

(2) $H^{s,t}(H_*(X) \otimes \Lambda) = H^{s,t}(H_*(X) \otimes \Lambda, \delta) = \text{Ext}_{\mathcal{A}}^{s,t+s}(X) = \text{Ext}_{\mathcal{A}}^{s,t+s}(H_*(X), \mathbb{Z}/2)$.

Note that since $H_*(X) \otimes \Lambda$ is a differential Λ -module, $\text{Ext}_{\mathcal{A}}^{*,*}(X)$ is an $\text{Ext}_{\mathcal{A}}^{*,*}$ -module in the obvious way.

To calculate $\text{Ext}_{\mathcal{A}}^{*,*}(P)$, when $X = P$, let e_k denote the generator of $H_k(P) \cong \mathbb{Z}/2$ for $k \geq 1$. The mod 2 Steenrod algebra \mathcal{A} acts on $e_k \in H_k(P)$ from the right by

$$(3.16) \quad e_k S q^{\nu+1} = \binom{k-\nu-1}{\nu+1} e_{k-\nu-1}.$$

From (3.15) (1) and (3.16) we thus see that the differential $H_*(P) \otimes \Lambda \xrightarrow{\delta} H_*(P) \otimes \Lambda$ of the differential Λ -module $H_*(P) \otimes \Lambda = \bigoplus_{k \geq 1} e_k \Lambda$ is given by

$$(3.17) \quad \delta(e_k \lambda_I) = e_k \delta(\lambda_I) + \sum_{\nu \geq 0} \binom{k-1-\nu}{\nu+1} e_{k-\nu-1} \lambda_{\nu} \lambda_I.$$

And $H^{s,t}(H_*(P) \otimes \Lambda, \delta) = \text{Ext}_{\mathcal{A}}^{s,t+s}(P)$.

Analogous to the operation $\Lambda \xrightarrow{S q^0} \Lambda$ in (3.12), there is also an operation $H_*(P) \otimes \Lambda \xrightarrow{S q^0} H_*(P) \otimes \Lambda$ given by

$$(3.18) \quad S q^0(e_k \lambda_{j_1} \cdots \lambda_{j_s}) = e_{2k+1} \lambda_{2j_1+1} \cdots \lambda_{2j_s+1}.$$

This operation is compatible with the relations (3.8) and commutes with the differential (3.17), so it induces a map

$$H^{s,t}(H_*(P) \otimes \Lambda) = \text{Ext}_{\mathcal{A}}^{s,t+s}(P) \xrightarrow{S q^0} H^{s,2t+s+1}(H_*(P) \otimes \Lambda) = \text{Ext}_{\mathcal{A}}^{s,2t+2s+1}(P).$$

We also let $\text{Ext}_{\mathcal{A}}^{*,*}(P) \xrightarrow{(S q^0)^i} \text{Ext}_{\mathcal{A}}^{*,*}(P)$ denote the composite $\underbrace{S q^0 \cdots S q^0}_i$ for

$i \geq 1$, and let $\text{Ext}_{\mathcal{A}}^{*,*}(P) \xrightarrow{(S q^0)^0} \text{Ext}_{\mathcal{A}}^{*,*}(P)$ be the identity map.

Recall the transfer map $P \xrightarrow{t} S^0$ in (2.9). It is not difficult to show the following.

(3.19) $P \xrightarrow{t} S^0$ induces a differential Λ -map $H_*(P) \otimes \Lambda \xrightarrow{\bar{t}} \Lambda$ given by $\bar{t}(e_k \lambda_I) = \lambda_k \lambda_I$ for any $I = (i_1, \dots, i_s)$. It gives rise to an Ext groups map

$$\text{Ext}_{\mathcal{A}}^{s,t}(P) \xrightarrow{t_*} \text{Ext}_{\mathcal{A}}^{s+1,t+1}, \quad \text{for all } s \geq 0, t \geq 0.$$

From (3.12), (3.18) and (3.19) one easily sees the following.

(3.20) There is a commutative diagram

$$\begin{array}{ccc} \text{Ext}_{\mathcal{A}}^{s,t}(P) & \xrightarrow{t_*} & \text{Ext}_{\mathcal{A}}^{s+1,t+1} \\ \downarrow (Sq^0)^i & & \downarrow (Sq^0)^i \\ \text{Ext}_{\mathcal{A}}^{s,2^i t+2^i-1}(P) & \xrightarrow{t_*} & \text{Ext}_{\mathcal{A}}^{s+1,2^i(t+1)} \end{array}$$

for all $i \geq 1$.

A class $\alpha \in \text{Ext}_{\mathcal{A}}^{s,t}(P)$ is a decomposable element in the $\text{Ext}_{\mathcal{A}}^{*,*}$ -module $\text{Ext}_{\mathcal{A}}^{*,*}(P)$ if $\alpha = \sum_i a_i \alpha_i$ for some $a_i \in \text{Ext}_{\mathcal{A}}^{s_i, t_i}$ with $s_i > 0$ and $\alpha_i \in \text{Ext}_{\mathcal{A}}^{s_i, t_i}(P)$. Otherwise α is an indecomposable element. In (3.22) below we recall from [8, 16] all the indecomposable elements in $\text{Ext}_{\mathcal{A}}^{s,*}(P)$ for $s \leq 4$. Only the classes $\widehat{h}_i, \widehat{g}_{i+1}, \widehat{D}_3(i), \gamma_{61}(i)$ in (3.22) (i) (1), (3.22) (ii) (4), (3.22) (ii) (6), (3.22) (iii) (11) for $i \geq 0$ will be explicitly described in terms of cycle representations in $H_*(P) \otimes \Lambda$ as in (3.21) below. The cycle representations of other classes will not be recorded here because they will not be used in this paper.

- (3.21) (1) $\widehat{h}_i = (Sq^0)^{i-1}(\widehat{h}_1) = \{e_{2^i-1} = (Sq^0)^{i-1}(e_1)\} \in \text{Ext}_{\mathcal{A}}^{0,2^i-1}(P), i \geq 1$.
(2) $\widehat{g}_{i+1} = (Sq^0)^i(\widehat{g}_1) = \{(Sq^0)^i(e_6 \lambda_0 \lambda_7^2 + e_5(\lambda_9 \lambda_3^2 + \lambda_3^2 \lambda_9) + e_3 \lambda_{11} \lambda_3^2 + e_3 \lambda_5 \lambda_9 \lambda_3)\} \in \text{Ext}_{\mathcal{A}}^{3,2^{i+4}+2^{i+3}-1}(P), i \geq 0$.
(3) $\widehat{D}_3(i) = (Sq^0)^i(\widehat{D}_3) = \{(Sq^0)^i(e_{22} \lambda_1 \lambda_7 \lambda_{31} + e_{16} \lambda_{15}^3 + e_{14} \lambda_9 \lambda_7 \lambda_{31})\} \in \text{Ext}_{\mathcal{A}}^{3,2^{i+6}+2^i-1}(P), i \geq 0$.
(4) $\gamma_{61}(i) = (Sq^0)^i(\gamma_{61} = \{\gamma_{61}^*\}) \in \text{Ext}_{\mathcal{A}}^{4,2^{i+6}+2^{i+1}-1}(P), i \geq 0$, where

$$\begin{aligned} \gamma_{61}^* = & e_{29}(\lambda_{13} \lambda_5 \lambda_7^2 + \lambda_9^2 \lambda_7^2 + \lambda_5^2 \lambda_{15} \lambda_7) + e_{28} \lambda_0 (\lambda_{19} \lambda_7^2 + \lambda_7^2 \lambda_{19}) + e_{27} \lambda_9 \lambda_3 \lambda_{15} \lambda_7 + \\ & e_{26} \lambda_2 (\lambda_{19} \lambda_7^2 + \lambda_7^2 \lambda_{19}) + e_{26} \lambda_4 \lambda_1 \lambda_{15}^2 + e_{25} \lambda_5 \lambda_1 \lambda_{15}^2 + e_{23} \lambda_0^3 \lambda_{15} \lambda_{23} + \\ & e_{23} \lambda_{17} \lambda_7^3 + e_{23} \lambda_9 \lambda_{15} \lambda_7^2 + e_{23} \lambda_7 \lambda_9 \lambda_{15} \lambda_7 + e_{23} \lambda_3 \lambda_5 \lambda_{15}^2 + e_{22} \lambda_6 (\lambda_{19} \lambda_7^2 + \\ & \lambda_7^2 \lambda_{19}) + e_{22} \lambda_8 \lambda_1 \lambda_{15}^2 + e_{22} \lambda_2 \lambda_3^2 \lambda_{31} + e_{21} \lambda_7 (\lambda_{19} \lambda_7^2 + \lambda_7^2 \lambda_{19}) + e_{21} (\lambda_{19} \lambda_7^2 + \\ & \lambda_7^2 \lambda_{19}) \lambda_7 + e_{20} \lambda_0 \lambda_{11} \lambda_{15}^2 + e_{19} \lambda_0^2 \lambda_{11} \lambda_{31} + e_{19} \lambda_5 \lambda_{15}^2 \lambda_7 + e_{19} \lambda_7 \lambda_5 \lambda_{15}^2 + \\ & e_{19} \lambda_9 \lambda_7 \lambda_{11} \lambda_{15} + e_{18} \lambda_2 \lambda_{11} \lambda_{15}^2 + e_{15} \lambda_1 \lambda_{15} \lambda_{23} \lambda_7 + e_{15} \lambda_8 \lambda_0 \lambda_{15} \lambda_{23} + \\ & e_{15} \lambda_0 \lambda_8 \lambda_{15} \lambda_{23} + e_{15} \lambda_9 \lambda_{15}^2 \lambda_7 + e_{15} \lambda_{11} \lambda_{17} \lambda_3 \lambda_{15} + e_{15} \lambda_{11} \lambda_{13} \lambda_{15} \lambda_7 + \\ & e_{15} \lambda_{11} \lambda_9 \lambda_{11} \lambda_{15} + e_{14} \lambda_{10} \lambda_3^2 \lambda_{31} + e_{14} \lambda_8 \lambda_1 \lambda_7 \lambda_{31} + e_{14} \lambda_5 \lambda_0 \lambda_{11} \lambda_{31} + \\ & e_{14} \lambda_4 \lambda_5 \lambda_7 \lambda_{31} + e_{14} \lambda_3 \lambda_2 \lambda_{11} \lambda_{31} + e_{14} \lambda_6 \lambda_{11} \lambda_{15}^2 + e_{14} \lambda_{14} (\lambda_{19} \lambda_7^2 + \lambda_7^2 \lambda_{19}) + \\ & e_{14} \lambda_{16} \lambda_1 \lambda_{15}^2 + e_{14} \lambda_2 \lambda_{15}^3 + e_{13} \lambda_{15} (\lambda_{19} \lambda_7^2 + \lambda_7^2 \lambda_{19}) + e_{13} \lambda_3 \lambda_{27} \lambda_3 \lambda_{15} + \\ & e_{13} \lambda_3^2 \lambda_{35} \lambda_7 + e_{13} \lambda_{15} \lambda_{11} \lambda_{15} \lambda_7 + e_{13} \lambda_0 \lambda_6 \lambda_{11} \lambda_{31} + e_{12} \lambda_8 \lambda_{11} \lambda_{15}^2 + \\ & e_{11} \lambda_8 \lambda_0 \lambda_{11} \lambda_{31} + e_{11} \lambda_0 \lambda_8 \lambda_{11} \lambda_{31} + e_{11} \lambda_1^3 \lambda_{47} + e_{11} \lambda_{13} \lambda_{15}^2 \lambda_7 + e_{11} \lambda_{15} \lambda_5 \lambda_{15}^2 + \\ & e_{11} \lambda_{17} (\lambda_{19} \lambda_7^2 + \lambda_7^2 \lambda_{19}) + e_{11} \lambda_0^2 \lambda_{19} \lambda_{31} + e_{10} \lambda_{10} \lambda_{11} \lambda_{15}^2 + e_7 \lambda_6^2 \lambda_{11} \lambda_{31} + \\ & e_7 \lambda_4 \lambda_0 \lambda_{19} \lambda_{31} + e_7 \lambda_0 \lambda_4 \lambda_{19} \lambda_{31} + e_7 \lambda_4 \lambda_8 \lambda_{11} \lambda_{31} + e_7 \lambda_{11} \lambda_5 \lambda_7 \lambda_{31} + \\ & e_7 \lambda_7 \lambda_9 \lambda_7 \lambda_{31} + e_7 \lambda_{17} \lambda_3^2 \lambda_{31} + e_7 \lambda_9 \lambda_{11} \lambda_3 \lambda_{31} + e_7 \lambda_5 \lambda_1^2 \lambda_{47} + e_7 \lambda_{13} \lambda_{15} \lambda_{11} \lambda_{15} + \\ & e_7 \lambda_{11} \lambda_{13} \lambda_{15}^2 + e_7 \lambda_{17} \lambda_3 \lambda_{19} \lambda_{15} + e_7 \lambda_9 \lambda_{11} \lambda_{19} \lambda_{15} + e_6 \lambda_2 \lambda_3^2 \lambda_{47} + e_5 \lambda_1^2 \lambda_{23} \lambda_{31} + \\ & e_3 \lambda_0^2 \lambda_{11} \lambda_{47}. \end{aligned}$$

- (3.22) (i) (1) $\widehat{h}_i = (Sq^0)^{i-1}(\widehat{h}_1) \in \text{Ext}_{\mathcal{A}}^{0,2^i-1}(P), i \geq 1.$
(2) $\widehat{c}_i = (Sq^0)^i(\widehat{c}_0) \in \text{Ext}_{\mathcal{A}}^{2,2^{i+3}+2^{i+1}+2^i-1}(P), i \geq 0.$
- (ii) (1) $\widehat{d}_i = (Sq^0)^i(\widehat{d}_0) \in \text{Ext}_{\mathcal{A}}^{3,2^{i+4}+2^{i+1}-1}(P), i \geq 0.$
(2) $\widehat{e}_i = (Sq^0)^i(\widehat{e}_0) \in \text{Ext}_{\mathcal{A}}^{3,2^{i+4}+2^{i+2}+2^i-1}(P), i \geq 0.$
(3) $\widehat{f}_i = (Sq^0)^i(\widehat{f}_0) \in \text{Ext}_{\mathcal{A}}^{3,2^{i+4}+2^{i+2}+2^{i+1}-1}(P), i \geq 0.$
(4) $\widehat{g}_{i+1} = (Sq^0)^i(\widehat{g}_1) \in \text{Ext}_{\mathcal{A}}^{3,2^{i+4}+2^{i+3}-1}(P), i \geq 0.$
(5) $\widehat{p}_i = (Sq^0)^i(\widehat{p}_0) \in \text{Ext}_{\mathcal{A}}^{3,2^{i+5}+2^{i+2}+2^i-1}(P), i \geq 0.$
(6) $\widehat{D}_3(i) = (Sq^0)^i(\widehat{D}_3) \in \text{Ext}_{\mathcal{A}}^{3,2^{i+6}+2^i-1}(P), i \geq 0.$
(7) $\widehat{p}'_i = (Sq^0)^i(\widehat{p}'_0) \in \text{Ext}_{\mathcal{A}}^{3,2^{i+6}+2^{i+3}+2^i-1}(P), i \geq 0.$
(8) $\alpha_{16}(i) = (Sq^0)^i(\alpha_{16}) \in \text{Ext}_{\mathcal{A}}^{3,2^{i+4}+2^{i+2}-1}(P), i \geq 0.$
(9) $\alpha_{21}(i) = (Sq^0)^i(\alpha_{21}) \in \text{Ext}_{\mathcal{A}}^{3,2^{i+4}+2^{i+3}+2^i-1}(P), i \geq 0.$
(10) $\xi_{31}(i) = (Sq^0)^i(\xi_{31}) \in \text{Ext}_{\mathcal{A}}^{3,2^{i+5}+2^{i+1}+2^i-1}(P), i \geq 0.$
- (iii) (1) $\widehat{P^1 h_1} \in \text{Ext}_{\mathcal{A}}^{4,13}(P).$
(2) $\widehat{P^1 h_2} \in \text{Ext}_{\mathcal{A}}^{4,15}(P).$
(3) $\delta_{30}(i) = (Sq^0)^i(\delta_{30}) \in \text{Ext}_{\mathcal{A}}^{4,2^{i+5}+2^{i+1}+2^i-1}(P), i \geq 0.$
(4) $\widehat{n}_i = (Sq^0)^i(\widehat{n}_0) \in \text{Ext}_{\mathcal{A}}^{4,2^{i+5}+2^{i+2}-1}(P), i \geq 0.$
(5) $\gamma_{37}(i) = (Sq^0)^i(\gamma_{37}) \in \text{Ext}_{\mathcal{A}}^{4,2^{i+5}+2^{i+3}+2^{i+1}-1}(P), i \geq 0.$
(6) $\widehat{x}_i = (Sq^0)^i(\widehat{x}_0) \in \text{Ext}_{\mathcal{A}}^{4,2^{i+5}+2^{i+3}+2^{i+1}-1}(P), i \geq 0.$
(7) $\alpha_{40}(i) = (Sq^0)^i(\alpha_{40}) \in \text{Ext}_{\mathcal{A}}^{4,2^{i+5}+2^{i+3}+2^{i+2}+2^i-1}(P), i \geq 0.$
(8) $\phi_{45}(i) = (Sq^0)^i(\phi_{45}) \in \text{Ext}_{\mathcal{A}}^{4,2^{i+5}+2^{i+4}+2^{i+1}-1}(P), i \geq 0.$
(9) $\gamma_{50}(i) = (Sq^0)^i(\gamma_{50}) \in \text{Ext}_{\mathcal{A}}^{4,2^{i+5}+2^{i+4}+2^{i+2}+2^{i+1}+2^i-1}(P), i \geq 0.$
(10) $\widehat{D}_1(i) = (Sq^0)^i(\widehat{D}_1) \in \text{Ext}_{\mathcal{A}}^{4,2^{i+5}+2^{i+4}+2^{i+3}+2^i-1}(P), i \geq 0.$
(11) $\gamma_{61}(i) = (Sq^0)^i(\gamma_{61}) \in \text{Ext}_{\mathcal{A}}^{4,2^{i+6}+2^{i+1}-1}(P), i \geq 0.$
(12) $\widehat{H}_1(i) = (Sq^0)^i(\widehat{H}_1) \in \text{Ext}_{\mathcal{A}}^{4,2^{i+6}+2^{i+1}+2^i-1}(P), i \geq 0.$
(13) $\gamma_{63}(i) = (Sq^0)^i(\gamma_{63}) \in \text{Ext}_{\mathcal{A}}^{4,2^{i+6}+2^{i+1}+2^i-1}(P), i \geq 0.$
(14) $\phi_{65}(i) = (Sq^0)^i(\phi_{65}) \in \text{Ext}_{\mathcal{A}}^{4,2^{i+6}+2^{i+2}+2^{i+1}-1}(P), i \geq 0.$
(15) $\delta_{67}(i) = (Sq^0)^i(\delta_{67}) \in \text{Ext}_{\mathcal{A}}^{4,2^{i+6}+2^{i+3}-1}(P), i \geq 0.$
(16) $\widehat{Q}_3(i) = (Sq^0)^i(\widehat{Q}_3) \in \text{Ext}_{\mathcal{A}}^{4,2^{i+6}+2^{i+3}-1}(P), i \geq 0.$
(17) $\rho_{80}(i) = (Sq^0)^i(\rho_{80}) \in \text{Ext}_{\mathcal{A}}^{4,2^{i+6}+2^{i+4}+2^{i+2}+2^i-1}(P), i \geq 0.$
(18) $\widehat{K}_i = (Sq^0)^i(\widehat{K}_0) \in \text{Ext}_{\mathcal{A}}^{4,2^{i+7}+2^{i+1}-1}(P), i \geq 0.$
(19) $\gamma_{128}(i) = (Sq^0)^i(\gamma_{128}) \in \text{Ext}_{\mathcal{A}}^{4,2^{i+7}+2^{i+2}+2^i-1}(P), i \geq 0.$
(20) $\delta_{128}(i) = (Sq^0)^i(\delta_{128}) \in \text{Ext}_{\mathcal{A}}^{4,2^{i+7}+2^{i+2}+2^i-1}(P), i \geq 0.$
(21) $\widehat{J}_i = (Sq^0)^i(\widehat{J}_0) \in \text{Ext}_{\mathcal{A}}^{4,2^{i+7}+2^{i+2}+2^i-1}(P), i \geq 0.$
(22) $\gamma_{132}(i) = (Sq^0)^i(\gamma_{132}) \in \text{Ext}_{\mathcal{A}}^{4,2^{i+7}+2^{i+3}+2^i-1}(P), i \geq 0.$
(23) $\gamma_{140}(i) = (Sq^0)^i(\gamma_{140}) \in \text{Ext}_{\mathcal{A}}^{4,2^{i+7}+2^{i+4}+2^i-1}(P), i \geq 0.$

$$\begin{aligned}
(24) \quad \widehat{T}_i &= (Sq^0)^i(\widehat{T}_0) \in \text{Ext}_{\mathcal{A}}^{4,2^{i+7}+2^{i+4}+2^{i+1}-1}(P), i \geq 0. \\
(25) \quad \delta_{144}(i) &= (Sq^0)^i(\delta_{144}) \in \text{Ext}_{\mathcal{A}}^{4,2^{i+7}+2^{i+4}+2^{i+2}+2^i-1}(P), i \geq 0. \\
(26) \quad \gamma_{148}(i) &= (Sq^0)^i(\gamma_{148}) \in \text{Ext}_{\mathcal{A}}^{4,2^{i+7}+2^{i+4}+2^{i+3}+2^i-1}(P), i \geq 0. \\
(27) \quad \delta_{156}(i) &= (Sq^0)^i(\delta_{156}) \in \text{Ext}_{\mathcal{A}}^{4,2^{i+7}+2^{i+5}+2^i-1}(P), i \geq 0. \\
(28) \quad \widehat{V}_i &= (Sq^0)^i(\widehat{V}_0) \in \text{Ext}_{\mathcal{A}}^{4,2^{i+7}+2^{i+5}+2^i-1}(P), i \geq 0. \\
(29) \quad \gamma_{164}(i) &= (Sq^0)^i(\gamma_{164}) \in \text{Ext}_{\mathcal{A}}^{4,2^{i+7}+2^{i+5}+2^{i+3}+2^i-1}(P), i \geq 0. \\
(30) \quad \widehat{V}'_i &= (Sq^0)^i(\widehat{V}'_0) \in \text{Ext}_{\mathcal{A}}^{4,2^{i+8}+2^i-1}(P), i \geq 0. \\
(31) \quad \widehat{U}_i &= (Sq^0)^i(\widehat{U}_0) \in \text{Ext}_{\mathcal{A}}^{4,2^{i+8}+2^{i+3}+2^i-1}(P), i \geq 0.
\end{aligned}$$

We also describe the class $\beta_{187} \in \text{Ext}_{\mathcal{A}}^{5,192}(P)$ in (3.2) (8) as follows. β_{187} is an indecomposable element too.

$$(3.23) \quad \beta_{187} = \{e_8 \lambda_3^2 \lambda_{47} \lambda_{63}^2\} \in \text{Ext}_{\mathcal{A}}^{5,192}(P).$$

Thus all the classes in (3.2) can be read from (3.13), (3.21) and (3.23), noting that $\text{Ext}_{\mathcal{A}}^{s,*}(P)$ is an $\text{Ext}_{\mathcal{A}}^{*,*}$ -module.

Now we state the ‘‘already known’’ results on $\text{Ext}_{\mathcal{A}}^{s,*}$ for $s \leq 4$ from [8, 16] as follows.

Theorem 3.24 ([8, 16]). *(i) Modulo indecomposable elements in $\text{Ext}_{\mathcal{A}}^{3,*}(P)$, $\text{Ext}_{\mathcal{A}}^{s,*}(P)$ for $s \leq 3$ is generated by $\widehat{h}_i, \widehat{c}_i, h_i, c_i$ subject to the relations in Theorem 3.14 (i), and the following relations.*

$$\begin{aligned}
(1) \quad \widehat{h}_i h_{i-1} &= 0, & (6) \quad \widehat{h}_i c_j &= 0 \text{ for } j = i-2, i-3, \\
(2) \quad \widehat{h}_{i+2} h_i^2 &= \widehat{h}_{i+1} h_{i+1}^2, & (7) \quad \widehat{c}_i h_j &= 0 \text{ for } j = i-1, i, i+2, i+3, \\
(3) \quad \widehat{h}_{i+2} h_{i+2} h_i &= 0, & & \\
(4) \quad \widehat{h}_{i+3} h_{i+3} h_i^2 &= 0, & & \\
(5) \quad \widehat{h}_{i+2} h_{i+3}^2 h_i &= 0, & (8) \quad \widehat{c}_i h_{i+1} &= \widehat{h}_{i+1} c_i \neq 0.
\end{aligned}$$

(ii) *The elements in (3.22) (ii) is a $\mathbb{Z}/2$ -base for the submodule of indecomposable elements in the $\mathbb{Z}/2$ -module $\text{Ext}_{\mathcal{A}}^{3,*}(P)$.*

(iii) *The decomposable elements in $\text{Ext}_{\mathcal{A}}^{4,*}(P)$ is the $\mathbb{Z}/2$ -submodule generated by the following elements.*

$$\begin{aligned}
(1) \quad \widehat{h}_{4+n} h_0^4, \quad n &\geq 0, & (8) \quad \widehat{h}_n h_{j-1}^2 h_k^2, \quad 3 &\leq j+2 \leq n \leq k, \\
(2) \quad \widehat{h}_l h_j^3 h_k, \quad 1 &\leq l \leq j-1 < k-3, & \quad j &< k-3, \\
(3) \quad \widehat{h}_n h_{j-1}^3 h_k, \quad 3 &\leq j+2 \leq n \leq k, & (9) \quad \widehat{c}_{k-2} h_{j-1}^2, \quad 1 &\leq j < k-3, \\
\quad \quad \quad j &< k-2, & (10) \quad \widehat{h}_{k+1+n} h_{j-1}^2 h_{k-1}^2, \quad n &\geq 1, 1 \leq \\
(4) \quad \widehat{h}_{k+n} h_{j-1}^3 h_{k-1}, \quad n &\geq 1, 1 \leq j < & \quad j &< k-3, \\
\quad \quad \quad k-2, & & (11) \quad \widehat{c}_{j-2} h_k^2, \quad 2 &\leq j < k-3, \\
(5) \quad \widehat{c}_{j-2} h_k, \quad 2 &\leq j < k-2, & (12) \quad \widehat{h}_l h_j^2 h_k h_i, \quad 1 &\leq l \leq j < k-2 < \\
(6) \quad \widehat{c}_{j-3} h_{j+1} h_k, \quad 3 &\leq j < k-2, & \quad i-3, & \\
(7) \quad \widehat{h}_l h_j^2 h_k^2, \quad 1 &\leq l \leq j < k-3, & (13) \quad \widehat{h}_n h_{j-1}^2 h_k h_i, \quad 3 &\leq j+2 \leq n \leq \\
& & \quad k, j &< k-2 < i-3,
\end{aligned}$$

- (14) $\widehat{h}_n h_{j-1}^2 h_{k-1} h_i, k+1 \leq n \leq i,$
 $1 \leq j < k-2 < i-3,$
- (15) $\widehat{h}_{i+n+1} h_{j-1}^2 h_{k-1} h_{i-1}, n \geq 0,$
 $1 \leq j < k-2 < i-3,$
- (16) $\widehat{c}_{j-2} h_k h_i, 2 \leq j < k-2 < i-3,$
- (17) $\widehat{h}_l h_j h_k^2 h_i, 1 \leq l \leq j < k-2 <$
 $i-4,$
- (18) $\widehat{f}_j h_i, 0 \leq j < i-5,$
- (19) $\alpha_{16}(j) h_i, 0 \leq j < i-5,$
- (20) $\widehat{g}_{j+1} h_i, 0 \leq j < i-5,$
- (21) $\widehat{h}_n h_{j-1} h_k^2 h_i, 2 \leq j+1 \leq n \leq$
 $k, j < k-2 < i-4,$
- (22) $\widehat{c}_{k-2} h_{j-1} h_i, 1 \leq j < k-3 <$
 $i-5,$
- (23) $\widehat{p}'_{j-1} h_i, 1 \leq j < i-5,$
- (24) $\widehat{h}_n h_{j-1} h_{k-1}^2 h_i, k+2 \leq n \leq i,$
 $1 \leq j < k-2 < i-4,$
- (25) $\widehat{h}_{i+1+n} h_{j-1} h_{k-1}^2 h_{i-1}, n \geq 0,$
 $1 \leq j < k-2 < i-4,$
- (26) $\widehat{h}_l h_j h_k h_i^2, 1 \leq l \leq j < k-1 <$
 $i-3,$
- (27) $\widehat{h}_n h_{j-1} h_k h_i^2, 2 \leq j+1 \leq n \leq$
 $k, j < k-1 < i-3,$
- (28) $\widehat{f}_k h_{j-1}, 1 \leq j < k-1,$
- (29) $\alpha_{16}(k) h_{j-1}, 1 \leq j < k-1,$
- (30) $\widehat{g}_{k+1} h_{j-1}, 1 \leq j < k-1,$
- (31) $\widehat{h}_n h_{j-1} h_{k-1} h_i^2, k+1 \leq n \leq i,$
 $1 \leq j < k-1 < i-3,$
- (32) $\widehat{c}_{i-2} h_{j-1} h_{k-1}, 1 \leq j < k-1 <$
 $i-3,$
- (33) $\widehat{p}'_{k-1} h_{j-1}, 1 \leq j < k-1,$
- (34) $\widehat{h}_{i+1+n} h_{j-1} h_{k-1} h_{i-1}^2, n \geq 1,$
 $1 \leq j < k-1 < i-3,$
- (35) $\widehat{h}_l h_j h_k^3, 1 \leq l \leq j < k-3,$
- (36) $\widehat{f}_j h_{j+5}, j \geq 0,$
- (37) $\alpha_{16}(j) h_{j+5}, j \geq 0,$
- (38) $\widehat{g}_{j+1} h_{j+5}, j \geq 0,$
- (39) $\widehat{h}_n h_{j-1} h_k^3, 2 \leq j+1 \leq n \leq$
 $k-1, j < k-3,$
- (40) $\widehat{c}_{k-3} h_{j-1} h_{k+1}, 1 \leq j < k-3,$
- (41) $\widehat{e}_{k-2} h_{j-1}, 1 \leq j < k-3,$
- (42) $\widehat{h}_{k+2+n} h_{j-1} h_{k-1}^3, n \geq 0, 1 \leq$
 $j < k-3,$
- (43) $\widehat{h}_l h_j h_k h_i h_p, 1 \leq l \leq j < k-$
 $1 < i-2 < p-3,$
- (44) $\widehat{h}_n h_{j-1} h_k h_i h_p, 2 \leq j+1 \leq$
 $n \leq k, j < k-1 < i-2 < p-3,$
- (45) $\widehat{h}_n h_{j-1} h_{k-1} h_i h_p, k+1 \leq n \leq$
 $i, 1 \leq j < k-1 < i-2 < p-3,$
- (46) $\widehat{h}_n h_{j-1} h_{k-1} h_{i-1} h_p, i+1 \leq$
 $n \leq p, 1 \leq j < k-1 < i-2 <$
 $p-3,$
- (47) $\widehat{h}_{p+n} h_{j-1} h_{k-1} h_{i-1} h_{p-1}, n \geq$
 $1, 1 \leq j < k-1 < i-2 < p-3,$
- (48) $\widehat{D}_3(k-1) h_{j-1}, 1 \leq j < k-1,$
- (49) $\widehat{D}_3(j-1) h_p, 1 \leq j < p-5,$
- (50) $\widehat{d}_j h_j, j \geq 0,$
- (51) $\widehat{h}_l c_j h_{j+1}, 1 \leq l \leq j,$
- (52) $\xi_{31}(j-1) h_j, j \geq 1,$
- (53) $\widehat{h}_{j+3+n} h_j c_{j-1}, n \geq 0, j \geq 1,$
- (54) $\widehat{h}_l h_j c_k, 1 \leq l \leq j < k-1,$
- (55) $\widehat{h}_n h_{j-1} c_k, 2 \leq j+1 \leq n \leq$
 $k+1, j < k-1,$
- (56) $\widehat{h}_{k+3+n} h_{j-1} c_{k-1}, n \geq 0, 1 \leq$
 $j < k-1,$
- (57) $\xi_{31}(k-1) h_{j-1}, 1 \leq j < k-1,$
- (58) $\alpha_{21}(k-1) h_{j-1}, 1 \leq j < k-1,$
- (59) $\widehat{p}_{k-1} h_{j-1}, 1 \leq j < k-1,$
- (60) $\widehat{d}_k h_{j-1}, 1 \leq j < k-1,$
- (61) $\widehat{h}_l c_j h_k, 1 \leq l \leq j+1 < k-2,$
- (62) $\widehat{d}_j h_k, 0 \leq j < k-4,$
- (63) $\alpha_{21}(j-1) h_k, 1 \leq j < k-3,$
- (64) $\widehat{p}_{j-1} h_k, 1 \leq j < k-4,$
- (65) $\widehat{e}_{j+1} h_{j-1}, j \geq 1,$
- (66) $\widehat{h}_n c_{j-1} h_k, 4 \leq j+3 \leq n \leq k,$
 $j < k-3,$
- (67) $\widehat{h}_{k+1+n} c_{j-1} h_{k-1}, n \geq 0, 1 \leq$
 $j < k-4,$
- (68) $\xi_{31}(j-1) h_k, 1 \leq j < k-3,$
- (69) $\widehat{h}_l d_j, 1 \leq l \leq j+2,$
- (70) $\widehat{h}_{j+4+n} d_{j-1}, n \geq 0, j \geq 1,$

$$\begin{array}{ll}
(71) \ \xi_{31}(j-1)h_{j+1}, \ j \geq 1, & (81) \ \widehat{g}_{j+2}h_j, \ j \geq 0, \\
(72) \ \xi_{31}(j-1)h_{j+2}, \ j \geq 1, & (82) \ \widehat{h}_{j+1}p'_{j-1}, \ j \geq 1, \\
(73) \ \widehat{h}_l e_j, \ 1 \leq l \leq j+2, & (83) \ \widehat{h}_{j+5+n}p_{j-1}, \ n \geq 0, \ j \geq 1, \\
(74) \ \widehat{h}_{j+4+n}e_{j-1}, \ n \geq 0, \ j \geq 1, & (84) \ \widehat{h}_{j+3}D_3(j-1), \ j \geq 1, \\
(75) \ \widehat{h}_l f_j, \ 1 \leq l \leq j+2, & (85) \ \widehat{h}_l D_3(j), \ 1 \leq l \leq j+3, \\
(76) \ \widehat{h}_{j+4+n}f_{j-1}, \ n \geq 0, \ j \geq 1, & (86) \ \widehat{h}_{j+6+n}D_3(j-1), \ n \geq 0, \ j \geq 1, \\
(77) \ \widehat{p}'_{j-1}h_{j-1}, \ j \geq 1, & (87) \ \widehat{h}_l p'_j, \ 1 \leq l \leq j+1 \text{ and } j+3 \leq \\
(78) \ \widehat{h}_l g_{j+1}, \ 1 \leq l \leq j+2, & \quad \quad \quad l \leq j+4, \\
(79) \ \widehat{h}_{j+4+n}g_j, \ n \geq 1, \ j \geq 1, & (88) \ \widehat{g}_{j+3}h_j, \ j \geq 0, \\
(80) \ \widehat{h}_l p_j, \ 1 \leq l \leq j \text{ and } l = j+3, & (89) \ \widehat{h}_{j+6+n}p'_{j-1}, \ n \geq 0, \ j \geq 1.
\end{array}$$

(iv) The elements in (3.22) (iii) is a $\mathbb{Z}/2$ -base for the submodule of indecomposable elements in the $\mathbb{Z}/2$ -module $\text{Ext}_{\mathcal{A}}^{4,*}(P)$.

It is not difficult to see that Theorem 3.24 covers (3.2) (j) (a) for $1 \leq j \leq 7$.

Let $\text{Ext}_{\mathcal{A}}^{s,t}(P) \xrightarrow{t_*} \text{Ext}_{\mathcal{A}}^{s+1,t+1}$ be as in (3.19). From (3.13), (3.19), (3.20) and (3.21) it is easy to see the following (3.25) (1) through (3.25) (3). (3.25) (4) is proved in [8, 16].

(3.25) $\text{Ext}_{\mathcal{A}}^{s,t}(P) \xrightarrow{t_*} \text{Ext}_{\mathcal{A}}^{s+1,t+1}$ has the following properties:

$$\begin{array}{ll}
(1) \ t_*(\widehat{h}_i) = h_i, \ i \geq 0. & (3) \ t_*(\widehat{D}_3(i)) = D_3(i), \ i \geq 0. \\
(2) \ t_*(\widehat{g}_{i+1}) = g_{i+1}, \ i \geq 0. & (4) \ t_*(\gamma_{61}(i)) = 0, \ i \geq 0.
\end{array}$$

It is easy to see that the results in (3.2) (j) (b) for $j = 1, 5$, (3.2) (j) (c) for $2 \leq j \leq 7$, (3.2) (8) (b) and (3.2) (9) (b) follow from (3.14) and (3.25). The equations in (3.2) (8) (c) and (3.2) (9) (c) also follow straightforwardly from (3.25). The equation $t_*(\beta_{187}) = h_5 V_0$ in (3.2) (8) (d) is proved as follows. By (3.13), $V_0 = \{\lambda_0 \lambda_{31} \lambda_{47} \lambda_{15} \lambda_{63}\}$. So $h_5 V_0 = V_0 h_5 = \{\lambda_0 \lambda_{31} \lambda_{47} \lambda_{15} \lambda_{63} \lambda_{31}\}$. By (3.8), (3.19) and (3.23) we have $t_*(\beta_{187}) = \{\lambda_8 \lambda_3^2 \lambda_{47} \lambda_{63}^2 = \lambda_0 \lambda_{15} \lambda_7^2 \lambda_{63} \lambda_{95}\}$. Straightforward calculations show that $\delta(\lambda_0 \lambda_{23} \lambda_7 \lambda_{63} \lambda_{95}) = \lambda_0 \lambda_{15} \lambda_7^2 \lambda_{63} \lambda_{95} + \lambda_0 \lambda_{31} \lambda_{47} \lambda_{15} \lambda_{63} \lambda_{31}$. So $t_*(\beta_{187}) = h_5 V_0$. Nontriviality and linear independence as claimed in (3.2) (8) (c), (3.2) (8) (d), and (3.2) (9) (c) will be proved later.

We recall the algebraic Kahn-Priddy theorem of Lin mentioned in section 2 as follows.

Theorem 3.26 (algebraic Kahn-Priddy theorem). $\text{Ext}_{\mathcal{A}}^{s,t}(P) \xrightarrow{t_*} \text{Ext}_{\mathcal{A}}^{s+1,t+1}$ is onto for all t, s with $t - s > 0$.

From Theorem 3.26, it is easy to see that the Ext group results in (3.1) (7), (8) follow from (3.2) (8) and (3.2) (9) (cf. Remark 3.7). Thus to complete the proof of (3.1) and (3.2) it suffices to prove (3.1) (9), (3.2) (8) (a) and (3.2) (9) (a) (which we will simply say “the remaining (3.2) (a)”), and (3.2) (8) (c) and (3.2) (9) (c) (which we will simply say “the remaining (3.2) (c)”).

For this purpose we need to recall how Theorem 3.26 is proved. First of all define $\Lambda(n) \subset \Lambda$ as follows.

(3.27) For each $n \geq 0$, let $\Lambda(n)$ be the $\mathbb{Z}/2$ -submodule of Λ generated by the admissible monomials $\lambda_{i_1} \cdots \lambda_{i_s}$ with $i_1 \leq n$.

It is not difficult to see that each $\Lambda(n)$ is a subcomplex of Λ . So we have an increasing filtration $1 = \Lambda(0) \subset \Lambda(1) \subset \Lambda(2) \subset \cdots \subset \Lambda(n) \subset \Lambda(n+1) \subset \cdots$ of subcomplexes of Λ . The differential Λ -module $H_*(P) \otimes \Lambda$ also has an increasing filtration of subcomplexes $F(0) = 0 \subset F(1) \subset F(2) \subset \cdots \subset F(n) \subset F(n+1) \subset \cdots$ defined as follows.

(3.28) $F(n) = \sum_{1 \leq k \leq n} H_k(P) \otimes \Lambda$, for $n \geq 1$.

It is easy to see the following.

(3.29) $F(i)/F(i-1) \cong \Sigma^i \Lambda$.

In [18] Lin has shown the following.

(3.30) There is a chain map $\Lambda \xrightarrow{\phi} H_*(P) \otimes \Lambda$ that goes from $\Lambda^{s+1, t-s}$ to $(H_*(P) \otimes \Lambda)^{s, t-s}$, and so induces an Ext group map $\text{Ext}_{\mathcal{A}}^{s+1, t+1} \xrightarrow{\phi_*} \text{Ext}_{\mathcal{A}}^{s, t}(P)$ for $t-s \geq 0$. The chain map ϕ has the following properties.

- (1) $\phi(\Lambda(i)) \subset F(i)$ for all $i \geq 0$.
- (2) $\phi(\lambda_{i_1} \lambda_{i_2} \cdots \lambda_{i_s}) \equiv e_{i_1} \lambda_{i_2} \cdots \lambda_{i_s} \pmod{F(i_1 - 1)}$ for any admissible monomial $\lambda_{i_1} \cdots \lambda_{i_s}$ with $i_1 \geq 1$.
- (3) Let ψ be the composite $\Lambda \xrightarrow{\phi} H_*(P) \otimes \Lambda \xrightarrow{\bar{t}} \Lambda$ where \bar{t} is as in (3.19). Then for each admissible monomial $\lambda_{i_1} \lambda_{i_2} \cdots \lambda_{i_s} \in \Lambda$ with $i_1 \geq 1$ there is the relation

$$\psi(\lambda_{i_1} \lambda_{i_2} \cdots \lambda_{i_s}) \equiv \lambda_{i_1} \lambda_{i_2} \cdots \lambda_{i_s} \pmod{\Lambda(i_1 - 1)}.$$

It is easy to see that Theorem 3.26 follows from (3.20) and (3.30) (3). To complete our proofs for (3.1) and (3.2), we are going to use a spectral sequence $\{E_r^{i, s, t}\}_{r \geq 1}$ for computing $\text{Ext}_{\mathcal{A}}^{s, t}(P)$ described in (3.32) below to prove the remaining (3.2) (a). Then we will use these results, and the map ϕ_* in (3.30) to deduce (3.1) (9) and the remaining (3.2) (c).

The spectral sequence for computing $\text{Ext}_{\mathcal{A}}^{s, t+s}(P) = H^{s, t}(H_*(P) \otimes \Lambda)$ is the one defined by the filtration $\{F(i)\}_{i \geq 0}$ for $H_*(P) \otimes \Lambda$ in (3.28). From (3.29) we see that

$$(3.31) \quad H^{s, t}(F(i)/F(i-1)) \cong H^{s, t}(\Sigma^i \Lambda) \cong \Sigma^i \text{Ext}_{\mathcal{A}}^{s, t+s-i}.$$

So we have the following.

(3.32) The filtration $\{F(i)\}_{i \geq 0}$ gives rise to a spectral sequence $\{E_r^{i, s, t}\}_{r \geq 1}$ with $E_1^{i, s, t} = H^{s, t}(F(i)/F(i-1)) \cong \Sigma^i \text{Ext}_{\mathcal{A}}^{s, t+s-i}$, $i \geq 1$, and $\bigoplus_{i \geq 1} E_{\infty}^{i, s, t} \cong \text{Ext}_{\mathcal{A}}^{s, t+s}(P)$ as $\mathbb{Z}/2$ -modules. The differential d_r of this spectral sequence goes from $E_r^{i, s, t}$ to $E_r^{i-r, s+1, t-1}$.

We will simply write $E_r^{*, s, *} \xrightarrow{d_r} E_r^{*, s+1, *}$ to indicate that we are considering the differentials for a fixed s and for all i, t, r . From (3.32) we see that for a given $s > 1$, if one knows completely the Ext groups $\text{Ext}_{\mathcal{A}}^{s, *}$ and has computed

all the differentials $E_r^{*,s-1,*} \xrightarrow{d_r} E_r^{*,s,*}$ and $E_r^{*,s,*} \xrightarrow{d_r} E_r^{*,s+1,*}$, then one can determine the Ext groups $\text{Ext}_{\mathcal{A}}^{s,*}(P)$. This is done in [9], [16] and [8], where the differentials $E_r^{*,s,*} \xrightarrow{d_r} E_r^{*,s+1,*}$ with $0 \leq s \leq 2$, with $s = 3$ and with $s = 4$ are computed respectively. To deduce the remaining (3.2) (a) we have to compute the differentials $E_r^{i,s,t} \xrightarrow{d_r} E_r^{i-r,s+1,t-1}$ for $(s,t) = (4,188), (4,189)$ and for $(s,t) = (5,187), (5,188)$, all i,r . These will be described in (3.36) and (3.37) below. As will be seen later in (3.44), to deduce (3.1) (9) we also need to compute the above differential for $(s,t) = (4,190)$ and for $(s,t) = (5,189)$. This will be described in (3.38) below.

To state (3.36), (3.37) and (3.38) we use the following conventions. By (3.32), if α is a basis element in $\text{Ext}_{\mathcal{A}}^{s,*}$ then $e_i\alpha = e_i \otimes \alpha$ is a basis element in $E_1^{i,s,*}$. We will write $e_i\alpha \rightarrow e_{i-r}\beta$, where $e_{i-r}\beta$ is some basis element in $E_1^{i-r,s+1,*}$, to indicate that both $e_i\alpha$ and $e_{i-r}\beta$ survive to $E_r^{*,*,*}$ and $d_r(e_i\alpha) = e_{i-r}\beta$ in the spectral sequence. In (3.36), (3.37) and (3.38) below there are two families of non-trivial differentials: the differentials in (j) (a) for $j = 3.36, 3.37, 3.38$ (which will be referred to as the “(a)-differentials”) that go from $E_r^{*,4,*}$ to $E_r^{*,5,*}$, and the differentials in (j) (b) for $j = 3.36, 3.37, 3.38$ (which will be referred to as the “(b)-differentials”) that go from $E_r^{*,5,*}$ to $E_r^{*,6,*}$. Any basis element $e_i\alpha$ in $E_1^{*,5,*}$ that is not in the targets of the (a)-differentials nor in the sources of the (b)-differentials is a cycle that survives to $E_\infty^{*,5,*}$. These permanent cycles are listed in (j) (c) for $j = 3.36, 3.37, 3.38$ (which will be referred to as the “(c)-cycles”). For each (c)-cycle $e_i\alpha$, we will write $e_i\alpha \longleftrightarrow \beta$ to mean that $\beta \in \text{Ext}_{\mathcal{A}}^{*,*}(P)$ is the cohomology class represented by this cycle.

All of the (a)-differentials are covered by [8] ((4.1.1) through (4.1.289) in [8]). In (3.36), (3.37) and (3.38) below we simply list these differentials and omit the calculations. Computing the (b)-differentials is the main work of this section from which we are going to deduce the remaining (3.2) (a) and (3.2) (c). These calculations are lengthy and will not be given in this paper. Also, the targets of the (b)-differentials are of the form $e_i\beta$ where $\beta \in \text{Ext}_{\mathcal{A}}^{6,*}$. So to state the (b)-differentials we need to describe some indecomposable elements in $\text{Ext}_{\mathcal{A}}^{6,*}$ (elements which are not sums of products of elements in (3.13)). These elements are listed in (3.33) below.

- (3.33)** (1) $\overline{\lambda_{19}D_1} = \{\overline{\lambda_{19}D_1}^* = \lambda_{19}D_1^* + \lambda_{15}\lambda_1\lambda_2\lambda_3^2\lambda_{47} + (\lambda_{14}\lambda_9\lambda_7 + \lambda_{11}\lambda_{12}\lambda_7 + \lambda_7\lambda_{16}\lambda_7 + \lambda_7\lambda_7\lambda_{16})\lambda_{11}\lambda_{15}^2\}$.
(2) $\overline{\lambda_{39}D_1(1)} = Sq^0(\overline{\lambda_{19}D_1})$.
(3) $\overline{\lambda_{40}D_1(1)} = \{\lambda_2\lambda_3^2(\lambda_{79}\lambda_{31}^2 + \lambda_{31}^2\lambda_{79})\}$.
(4) $A = \{\lambda_9(\lambda_4\lambda_7\lambda_{11}\lambda_{15}^2) + \lambda_5\lambda_1\lambda_2\lambda_3^2\lambda_{47} + \lambda_3\lambda_0^3\lambda_{11}\lambda_{47}\}$.
(5) $A' = \{\lambda_{20}\lambda_0^2c_2^* + \lambda_{18}\lambda_2\lambda_0c_2^* + \lambda_{17}\lambda_1\lambda_2c_2^* + \lambda_{15}\lambda_4\lambda_7\lambda_5\lambda_{15}^2 + \lambda_{14}\lambda_6\lambda_0c_2^* + \lambda_{13}(\lambda_5\lambda_2 + \lambda_3\lambda_4)c_2^* + \lambda_{12}(\lambda_8\lambda_0 + \lambda_0\lambda_8)c_2^* + \lambda_{11}(\lambda_8\lambda_9(\lambda_{19}\lambda_7^2 + \lambda_7^2\lambda_{19}) + \lambda_3\lambda_6c_2^* + \lambda_0\lambda_7\lambda_{13}\lambda_{15}^2) + \lambda_{12}f_0^*\lambda_{31} + \lambda_{10}(\lambda_2\lambda_4\lambda_{15}^3 + \lambda_{10}\lambda_0c_2^* + \lambda_2\lambda_8c_2^*) + \lambda_9(\lambda_9\lambda_2c_2^* + \lambda_1\lambda_{10}c_2^*) + \lambda_7(\lambda_4\lambda_7\lambda_{13}\lambda_{15}^2 + \lambda_7\lambda_6c_2^* + \lambda_5\lambda_8c_2^*) + \lambda_6(\lambda_6\lambda_8c_2^* + \lambda_{10}\lambda_0\lambda_{15}^3 + \lambda_9^2\lambda_3^2 + \lambda_9\lambda_3^2\lambda_9 + \lambda_7\lambda_{11}\lambda_3^2\lambda_{31} + \lambda_6\lambda_4\lambda_{15}^3 + \lambda_7\lambda_5\lambda_9\lambda_3\lambda_{31})\}$.
(6) $D_2 = \{\lambda_0^4\lambda_{11}\lambda_{47}\}$.
(7) $t_0 = \{\lambda_5(\lambda_9\lambda_3\lambda_5\lambda_7^2 + \lambda_6\lambda_0\lambda_3\lambda_{15}\lambda_7 + \lambda_3\lambda_5\lambda_1\lambda_{15}\lambda_7) + \lambda_3(\lambda_{11}\lambda_3\lambda_5\lambda_7^2 + \lambda_5\lambda_1\lambda_{13}\lambda_7^2 + \lambda_3\lambda_5\lambda_1^2\lambda_{23})\}$.
(8) $q_0 = \{\lambda_2\lambda_0^3\lambda_{15}^2 + \lambda_1((\lambda_5\lambda_1^3 + \lambda_2\lambda_4\lambda_1^2 + \lambda_1\lambda_2\lambda_4\lambda_1 + \lambda_2\lambda_1^2\lambda_4)\lambda_{23} + \lambda_6\lambda_0\lambda_3\lambda_{15}\lambda_7 + \lambda_2\lambda_{10}\lambda_5\lambda_7^2 + \lambda_2\lambda_4\lambda_7\lambda_3\lambda_{15} + \lambda_9\lambda_3\lambda_5\lambda_7^2)\}$.

We also need to show the following.

(3.34) Let S be the set of all the classes β such that $e_i\beta$ appears in the targets of a (b)-differential in (3.36) through (3.38). Then S is linearly independent in $\text{Ext}_{\mathcal{A}}^{6,*}$.

(3.34) will be proved in (3.43) later.

Finally we make some remarks on the (c)-cycles. From (3.32), a permanent cycle in the spectral sequence represents a class in $\text{Ext}_{\mathcal{A}}^{*,*}(P)$. One needs some further calculations to describe the class represented by a permanent cycle in terms of the cohomology classes in (3.13) and (3.21). These calculations will not be given in this paper. Instead, we illustrate our calculations for two examples of these. The first example is $e_{14}h_1f_3$ in (3.36) (c) (2) which represents the cohomology class $\widehat{h}_4h_0f_3 \in \text{Ext}_{\mathcal{A}}^{5,192}(P)$. From (3.13), $f_3 = \{f_3^*\} \in \text{Ext}_{\mathcal{A}}^{4,176}$ where

$$f_3^* = (Sq^0)^3(f_0^*) = \lambda_{39}\lambda_7\lambda_{63}^2 + \lambda_{31}(\lambda_{79}\lambda_{31} + \lambda_{31}\lambda_{47}\lambda_{63}) + \lambda_{23}\lambda_{39}\lambda_{47}\lambda_{63}.$$

From (3.21) we have $\widehat{h}_4 = \{e_{15}\}$, so $\widehat{h}_4h_0 = \{e_{15}\lambda_0\} = \{e_{14}\lambda_1 + e_{12}\lambda_3 + e_8\lambda_7\}$, and $\widehat{h}_4h_0f_3 = \{(e_{14}\lambda_1 + e_{12}\lambda_3 + e_8\lambda_7)f_3^*\}$. It follows that the permanent cycle $e_{14}h_1f_3$ represents the class $\widehat{h}_4h_0f_3$.

The other example is the following.

(3.35) The cycle $e_{47}T_0$ represents the cohomology class $\widehat{g}_4h_0^2 \in \text{Ext}_{\mathcal{A}}^{5,193}(P)$ in the the spectral sequence of $\text{Ext}_{\mathcal{A}}^{*,*}(P)$.

The result (3.35) will be used to prove (3.4) (17). It is explained as follows. From (3.21) we have $\widehat{g}_4 = \{\widehat{g}_4^*\} \in \text{Ext}_{\mathcal{A}}^{3,191}(P)$ where

$$\widehat{g}_4^* = (Sq^0)^3(\widehat{g}_1^*) = e_{55}\lambda_7\lambda_{63}^2 + e_{47}(\lambda_{79}\lambda_{31}^2 + \lambda_{31}^2\lambda_{79}) + e_{31}\lambda_{47}\lambda_{79}\lambda_{31},$$

so $\widehat{g}_4h_0^2 = \{\widehat{g}_4^*\lambda_0^2\}$. From (3.13) we have $T_0 = \{T_0^*\} \in \text{Ext}_{\mathcal{A}}^{5,146}$ where

$$T_0^* = \lambda_0^2(\lambda_{79}\lambda_{31}^2 + \lambda_{31}^2\lambda_{79}) + (\lambda_9\lambda_3^2 + \lambda_3^2\lambda_9)\lambda_{63}^2.$$

The infinite cycle $e_{47}T_0$ represents $\widehat{g}_4h_0 \in \text{Ext}_{\mathcal{A}}^{5,193}(P)$ because $\delta(B_0) \equiv \widehat{g}_4^*\lambda_0^2 + e_{47}T_0^* \pmod{F(46)}$ where

$$\begin{aligned} B_0 = & e_{55}\lambda_7\lambda_{63}\lambda_{64}\lambda_0 + e_{55}\lambda_7\lambda_{63}\lambda_0\lambda_{64} + e_{55}\lambda_7\lambda_{64}\lambda_0\lambda_{63} + e_{55}\lambda_7\lambda_0\lambda_{64}\lambda_{63} + e_{55}\lambda_8\lambda_0\lambda_{63}^2 + \\ & e_{55}\lambda_0\lambda_8\lambda_{63}^2 + e_{56}\lambda_0\lambda_7\lambda_{63}^2 + e_{54}\lambda_2\lambda_7\lambda_{63}^2 + e_{57}\lambda_3^2\lambda_{63}^2 + e_{52}\lambda_4\lambda_7\lambda_{63}^2 + e_{51}\lambda_5\lambda_7\lambda_{63}^2 + \\ & e_{47}(\lambda_{79}\lambda_{31}\lambda_{32}\lambda_0 + \lambda_{79}\lambda_{31}\lambda_0\lambda_{32} + \lambda_{79}\lambda_{32}\lambda_0\lambda_{31} + \lambda_{79}\lambda_0\lambda_{32}\lambda_{31} + \lambda_{80}\lambda_0\lambda_{31}^2 + \lambda_{63}\lambda_{16}\lambda_{32}\lambda_{31} + \\ & \lambda_{64}\lambda_{16}\lambda_{31}^2 + \lambda_0\lambda_{80}\lambda_{31}^2 + \lambda_{31}\lambda_{47}\lambda_{64}\lambda_0 + \lambda_{31}\lambda_{47}\lambda_0\lambda_{64} + \lambda_{31}\lambda_{48}\lambda_0\lambda_{63} + \lambda_{31}\lambda_0\lambda_{48}\lambda_{63} + \\ & \lambda_0\lambda_{32}\lambda_{47}\lambda_{63} + \lambda_{32}\lambda_0\lambda_{47}\lambda_{63} + \lambda_9\lambda_7\lambda_{63}^2 + \lambda_7^2\lambda_{64}^2 + \lambda_7\lambda_8\lambda_{64}\lambda_{63} + \lambda_7\lambda_8\lambda_{63}\lambda_{64} + \lambda_8\lambda_7\lambda_{64}\lambda_{63} + \\ & \lambda_8^2\lambda_{63}^2 + \lambda_8\lambda_7\lambda_{63}\lambda_{64} + \lambda_{47}\lambda_{31}\lambda_{32}^2 + \lambda_{48}\lambda_{31}\lambda_{32}\lambda_{31} + \lambda_{48}\lambda_{31}^2\lambda_{32}^2 + \lambda_{47}\lambda_{32}^2\lambda_{31} + \lambda_{47}\lambda_{32}\lambda_{31}\lambda_{32} + \\ & \lambda_{31}^2\lambda_{16}\lambda_{64} + \lambda_{31}\lambda_{32}\lambda_{15}\lambda_{64} + \lambda_{32}\lambda_{31}\lambda_{15}\lambda_{64} + \lambda_{32}\lambda_{31}\lambda_{16}\lambda_{63} + \lambda_{32}^2\lambda_{15}\lambda_{63}). \end{aligned}$$

It follows that the infinite cycle $e_{47}T_0$ represents $\widehat{g}_4h_0^2$.

Now we describe the differentials $E_r^{i,4,188} \xrightarrow{d_r} E_r^{i-r,5,187} \xrightarrow{d_r} E_r^{i-2r,6,186}$, $E_r^{i,4,189} \xrightarrow{d_r} E_r^{i-r,5,188} \xrightarrow{d_r} E_r^{i-2r,6,187}$, and $E_r^{i,4,190} \xrightarrow{d_r} E_r^{i-r,5,189} \xrightarrow{d_r} E_r^{i-2r,6,188}$ in (3.36), (3.37), (3.38) below respectively (in the spectral sequence (3.32) for $\text{Ext}_{\mathcal{A}}^{*,*}(P)$).

(3.36) (a)

- | | |
|--|---|
| (1) $e_{12}h_2c_4 \rightarrow e_{11}h_0h_2c_4$ | (27) $e_{104}f_2 \rightarrow e_{103}h_0f_2$ |
| (2) $e_{16}f_3 \rightarrow e_{15}h_0f_3$ | (28) $e_{108}h_1^2h_4h_6 \rightarrow e_{106}h_1^3h_2h_4h_6$ |
| (3) $e_{16}h_4^3h_7 \rightarrow e_{15}h_0h_4^3h_7$ | (29) $e_{108}e_2 \rightarrow e_{107}h_0e_2$ |
| (4) $e_{20}c_2h_7 \rightarrow e_{19}h_0c_2h_7$ | (30) $e_{110}h_0^2h_4h_6 \rightarrow e_{109}h_0^3h_4h_6$ |
| (5) $e_{24}h_2^2h_5h_7 \rightarrow e_{20}h_2^3h_5h_7$ | (31) $e_{118}p_1 \rightarrow e_{111}h_4D_3$ |
| (6) $e_{24}e_3 \rightarrow e_{23}h_0e_3$ | (32) $e_{117}c_0h_6 \rightarrow e_{115}h_1c_0h_6$ |
| (7) $e_{28}h_1^2h_5h_7 \rightarrow e_{26}h_1^3h_5h_7$ | (33) $e_{118}h_0^2h_3h_6 \rightarrow e_{117}h_0^3h_3h_6$ |
| (8) $e_{30}h_0^2h_5h_7 \rightarrow e_{29}h_0^3h_5h_7$ | (34) $e_{120}d_2 \rightarrow e_{119}h_0d_2$ |
| (9) $e_{46}p'_1 \rightarrow e_{31}V_0$ | (35) $e_{124}h_1^2h_5 \rightarrow e_{122}h_1^3h_5^2$ |
| (10) $e_{42}c_1h_7 \rightarrow e_{38}h_2c_1h_7$ | (36) $e_{126}h_0^2h_5^2 \rightarrow e_{125}h_0^3h_5^2$ |
| (11) $e_{54}h_1h_3h_6^2 \rightarrow e_{40}h_3^3h_7$ | (37) $e_{138}c_1h_5 \rightarrow e_{134}h_2c_1h_5$ |
| (12) $e_{44}h_1^2h_4h_7 \rightarrow e_{42}h_1^3h_4h_7$ | (38) $e_{140}h_3c_2 \rightarrow e_{135}D_1$ |
| (13) $e_{44}p_2 \rightarrow e_{43}h_0p_2$ | (39) $e_{155}p_0 \rightarrow e_{142}h_1g_2$ |
| (14) $e_{46}h_0^2h_4h_7 \rightarrow e_{45}h_0^3h_4h_7$ | (40) $e_{144}h_0g_2 \rightarrow e_{143}h_0g_2$ |
| (15) $e_{48}d_3 \rightarrow e_{47}h_0d_3$ | (41) $e_{150}e_1 \rightarrow e_{146}h_2e_1$ |
| (16) $e_{53}c_0h_7 \rightarrow e_{51}h_1c_0h_7$ | (42) $e_{148}f_1 \rightarrow e_{147}h_0f_1$ |
| (17) $e_{56}h_2^2h_6^2 \rightarrow e_{52}h_2^3h_6^2$ | (43) $e_{149}c_0h_5 \rightarrow e_{147}h_1c_0h_5$ |
| (18) $e_{54}h_0^2h_3h_7 \rightarrow e_{53}h_0^3h_3h_7$ | (44) $e_{150}h_0^2h_3h_5 \rightarrow e_{149}h_0^3h_3h_5$ |
| (19) $e_{60}h_1^2h_6^2 \rightarrow e_{58}h_1^3h_6^2$ | (45) $e_{156}d_1 \rightarrow e_{154}h_1d_1$ |
| (20) $e_{62}h_0^2h_6^2 \rightarrow e_{61}h_0^3h_6^2$ | (46) $e_{158}h_0^2h_4^2 \rightarrow e_{157}h_0^3h_4^2$ |
| (21) $e_{102}h_1c_3 \rightarrow e_{78}D_1(1)$ | (47) $e_{165}c_0h_4 \rightarrow e_{163}h_1c_0h_4$ |
| (22) $e_{84}c_2h_6 \rightarrow e_{83}h_0c_2h_6$ | (48) $e_{171}e_0 \rightarrow e_{166}h_1g_1$ |
| (23) $e_{88}h_4c_3 \rightarrow e_{87}h_0h_4c_3$ | (49) $e_{168}g_1 \rightarrow e_{167}h_0g_1$ |
| (24) $e_{119}p'_0 \rightarrow e_{94}h_1g_3$ | (50) $e_{170}f_0 \rightarrow e_{169}h_0f_0$ |
| (25) $e_{96}g_3 \rightarrow e_{95}h_0g_3$ | (51) $e_{174}d_0 \rightarrow e_{173}h_0d_0$ |
| (26) $e_{106}c_1h_6 \rightarrow e_{102}h_2c_1h_6$ | (52) $e_{188}h_0^4 \rightarrow e_{187}h_0^5$ |
-
- | | |
|--|--|
| (b) (1) $e_{10}c_1h_5h_7 \rightarrow e_6h_2c_1h_5h_7$ | (17) $e_{32}h_4d_3 \rightarrow e_{31}h_0h_4d_3$ |
| (2) $e_{12}h_1^2c_4 \rightarrow e_{10}h_1^3c_4$ | (18) $e_{37}c_0h_4h_7 \rightarrow e_{35}h_1c_0h_4h_7$ |
| (3) $e_{12}h_5p_2 \rightarrow e_{11}h_0h_5p_2$ | (19) $e_{40}h_3d_3 \rightarrow e_{20}h_2x_2$ |
| (4) $e_{12}h_3c_2h_7 \rightarrow e_7D_1h_7$ | (20) $e_{40}g_1h_7 \rightarrow e_{39}h_0g_1h_7$ |
| (5) $e_{14}h_0^2c_4 \rightarrow e_{13}h_0^3c_4$ | (21) $e_{42}f_0h_7 \rightarrow e_{41}h_0f_0h_7$ |
| (6) $e_{16}h_5d_3 \rightarrow e_{15}h_0h_5d_3$ | (22) $e_{42}c_1h_6^2 \rightarrow e_{38}h_2c_1h_6^2$ |
| (7) $e_{16}g_2h_7 \rightarrow e_{15}h_0g_2h_7$ | (23) $e_{43}e_0h_7 \rightarrow e_{38}h_2f_0h_7$ |
| (8) $e_{20}f_1h_7 \rightarrow e_{19}h_0f_1h_7$ | (24) $e_{44}p'_1h_1 \rightarrow e_{42}h_0^2p_2$ |
| (9) $e_{21}c_0h_5h_7 \rightarrow e_{19}h_1c_0h_5h_7$ | (25) $e_{46}h_1d_3 \rightarrow e_{30}h_1h_4d_3$ |
| (10) $e_{22}h_1e_3 \rightarrow e_8h_3^2e_3$ | (26) $e_{46}d_0h_7 \rightarrow e_{45}h_0d_0h_7$ |
| (11) $e_{22}e_1h_7 \rightarrow e_{18}h_2e_1h_7$ | (27) $e_{46}T_0 \rightarrow e_{45}h_0T_0$ |
| (12) $e_{22}h_0^2h_3h_5h_7 \rightarrow e_{21}h_0^3h_3h_5h_7$ | (28) $e_{48}Q_3(1) \rightarrow e_{47}h_0Q_3(1)$ |
| (13) $e_{24}x_2 \rightarrow e_{23}h_0x_2$ | (29) $e_{48}n_2 \rightarrow e_{47}h_0n_2$ |
| (14) $e_{27}p_0h_7 \rightarrow e_{14}h_1g_2h_7$ | (30) $e_{53}c_0h_6^2 \rightarrow e_{51}h_1c_0h_6^2$ |
| (15) $e_{28}d_1h_7 \rightarrow e_{26}h_1d_1h_7$ | (31) $e_{54}h_0^2h_3h_6^2 \rightarrow e_{53}h_0^3h_3h_6^2$ |
| (16) $e_{30}h_0^2h_4^2h_7 \rightarrow e_{29}h_0^3h_4^2h_7$ | (32) $e_{54}h_3D_3(1) \rightarrow e_{46}h_3^2D_3(1)$ |

- (33) $e_{58}H_1(1) \rightarrow e_{54}h_2H_1(1)$ (58) $e_{117}c_0h_5^2 \rightarrow e_{115}h_1c_0h_5^2$
(34) $e_{58}h_2D_3(1) \rightarrow e_{51}h_3J_0$ (59) $e_{118}h_0p'_0 \rightarrow e_{117}h_0^2p'_0$
(35) $e_{59}J_0 \rightarrow e_{46}h_1Q_3(1)$ (60) $e_{120}n_1 \rightarrow e_{119}h_0n_1$
(36) $e_{60}h_0^4h_7 \rightarrow e_{59}h_0^5h_7$ (61) $e_{120}Q_3 \rightarrow e_{119}h_0Q_3$
(37) $e_{62}K_0 \rightarrow e_{61}h_0K_0$ (62) $e_{123}h_2D_3 \rightarrow e_{115}\overline{\lambda_{19}D_1}$
(38) $e_{80}g_2h_6 \rightarrow e_{79}h_0g_2h_6$ (63) $e_{124}h_0^4h_6 \rightarrow e_{123}h_0^5h_6$
(39) $e_{84}f_1h_6 \rightarrow e_{83}h_0f_1h_6$ (64) $e_{125}h_1D_3 \rightarrow e_{118}h_1Q_3$
(40) $e_{86}e_1h_6 \rightarrow e_{82}h_2e_1h_6$ (65) $e_{125}H_1 \rightarrow e_{123}h_1H_1$
(41) $e_{88}h_4f_2 \rightarrow e_{87}h_0h_4f_2$ (66) $e_{136}g_1h_5 \rightarrow e_{135}h_0g_1h_5$
(42) $e_{91}p_0h_6 \rightarrow e_{78}h_1g_2h_6$ (67) $e_{138}f_0h_5 \rightarrow e_{137}h_0f_0h_5$
(43) $e_{92}d_1h_6 \rightarrow e_{90}h_1d_1h_6$ (68) $e_{139}e_0h_5 \rightarrow e_{134}h_1g_1h_5$
(44) $e_{92}h_2g_3 \rightarrow e_{91}h_0h_2g_3$ (69) $e_{140}h_2g_2 \rightarrow e_{139}h_0h_2g_2$
(45) $e_{94}h_0^2h_5^3 \rightarrow e_{93}h_0^3h_5^3$ (70) $e_{142}d_0h_5 \rightarrow e_{141}h_0d_0h_5$
(46) $e_{100}h_2f_2 \rightarrow e_{95}h_4^2D_3$ (71) $e_{148}h_1e_1 \rightarrow e_{146}h_1^2e_1$
(47) $e_{101}c_0h_4h_6 \rightarrow e_{99}h_1c_0h_4h_6$ (72) $e_{150}x_0 \rightarrow e_{149}h_0x_0$
(48) $e_{102}h_0^2c_3 \rightarrow e_{101}h_0^3c_3$ (73) $e_{152}h_2d_1 \rightarrow e_{148}h_2^2d_1$
(49) $e_{102}h_1f_2 \rightarrow e_{86}h_0^2h_4c_3$ (74) $e_{156}n_0 \rightarrow e_{150}t_0$
(50) $e_{104}h_2e_2 \rightarrow e_{100}h_2^2e_2$ (75) $e_{156}h_0^4h_5 \rightarrow e_{155}h_0^5h_5$
(51) $e_{104}g_1h_6 \rightarrow e_{103}h_0g_1h_6$ (76) $e_{164}h_2g_1 \rightarrow e_{163}h_0h_2g_1$
(52) $e_{106}f_0h_6 \rightarrow e_{105}h_0f_0h_6$ (77) $e_{170}h_0e_0 \rightarrow e_{169}h_0^2e_0$
(53) $e_{107}e_0h_6 \rightarrow e_{102}h_1g_1h_6$ (78) $e_{172}h_0^4h_4 \rightarrow e_{171}h_0^5h_4$
(54) $e_{108}x_1 \rightarrow e_{107}h_0x_1$ (79) $e_{172}h_1d_0 \rightarrow e_{170}h_1^2d_0$
(55) $e_{110}d_0h_6 \rightarrow e_{109}h_0d_0h_6$ (80) $e_{176}P^1h_2 \rightarrow e_{175}h_0P^1h_2$
(56) $e_{112}h_3d_2 \rightarrow e_{111}h_0h_3d_2$ (81) $e_{178}P^1h_1 \rightarrow e_{172}h_0^2d_0$
(57) $e_{116}h_1p_1 \rightarrow e_{106}h_1x_1$

- (c) (1) $e_8h_2^2c_4 \longleftrightarrow \beta_{187}$ (3) $e_{30}D_3(1)h_5 \longleftrightarrow \widehat{h}_5V_0$
(2) $e_{14}h_1f_3 \longleftrightarrow \widehat{h}_4h_0f_3$

- (3.37) (a) (1) $e_{21}c_2h_7 \rightarrow e_{13}h_3c_2h_7$ (14) $e_{57}h_2^2h_6^2 \rightarrow e_{53}h_2^3h_6^2$
(2) $e_{16}h_0c_4 \rightarrow e_{15}h_0^2c_4$ (15) $e_{56}h_0h_3h_6^2 \rightarrow e_{55}h_0^2h_3h_6^2$
(3) $e_{17}f_3 \rightarrow e_{15}h_1f_3$ (16) $e_{60}h_0h_2h_6^2 \rightarrow e_{59}h_1^3h_6^2$
(4) $e_{25}h_2^2h_5h_7 \rightarrow e_{21}h_2^3h_5h_7$ (17) $e_{62}h_0^3h_7 \rightarrow e_{61}h_0^4h_7$
(5) $e_{24}h_0h_3h_5h_7 \rightarrow e_{23}h_0^2h_3h_5h_7$ (18) $e_{85}c_2h_6 \rightarrow e_{77}h_3c_2h_6$
(6) $e_{25}e_3 \rightarrow e_{23}h_1e_3$ (19) $e_{89}h_4c_3 \rightarrow e_{79}D_1(1)$
(7) $e_{28}h_0h_2h_5h_7 \rightarrow e_{27}h_1^3h_5h_7$ (20) $e_{109}h_1^2h_4h_6 \rightarrow e_{63}K_0$
(8) $e_{32}h_0h_4^2h_7 \rightarrow e_{31}h_0^2h_4^2h_7$ (21) $e_{96}h_0h_5^3 \rightarrow e_{95}h_0^2h_5^3$
(9) $e_{45}p_2 \rightarrow e_{31}D_3(1)h_5$ (22) $e_{97}g_3 \rightarrow e_{95}h_1g_3$
(10) $e_{48}h_0h_2^3h_7 \rightarrow e_{38}c_0h_4h_7$ (23) $e_{104}h_0c_3 \rightarrow e_{103}h_0^2c_3$
(11) $e_{44}h_0h_2h_4h_7 \rightarrow e_{43}h_1^3h_4h_7$ (24) $e_{105}f_2 \rightarrow e_{103}h_1f_2$
(12) $e_{49}d_3 \rightarrow e_{47}h_1d_3$ (25) $e_{107}c_1h_6 \rightarrow e_{103}h_2c_1h_6$
(13) $e_{54}c_0h_7 \rightarrow e_{47}d_0h_7$ (26) $e_{108}h_0h_2h_4h_6 \rightarrow e_{107}h_0^2h_2h_4h_6$ \rightarrow

- (27) $e_{109}e_2 \rightarrow e_{101}h_3e_2$ (39) $e_{149}f_1 \rightarrow e_{141}h_3f_1$
(28) $e_{109}h_1^2h_4h_6 \rightarrow e_{63}K_0$ (40) $e_{150}c_0h_5 \rightarrow e_{143}d_0h_5$
(29) $e_{118}c_0h_6 \rightarrow e_{111}d_0h_6$ (41) $e_{151}e_1 \rightarrow e_{139}f_0h_5$
(30) $e_{119}p_1 \rightarrow e_{93}h_2g_3$ (42) $e_{156}p_0 \rightarrow e_{155}h_0p_0$
(31) $e_{120}p'_0 \rightarrow e_{119}h_0p'_0$ (43) $e_{157}d_1 \rightarrow e_{126}H_1$
(32) $e_{121}d_2 \rightarrow e_{117}h_2d_2$ (44) $e_{158}h_0^3h_5 \rightarrow e_{157}h_0^4h_5$
(33) $e_{124}h_0h_2h_5^2 \rightarrow e_{123}h_0^2h_2h_5^2$ (45) $e_{169}g_1 \rightarrow e_{167}h_1g_1$
(34) $e_{126}h_0^3h_6 \rightarrow e_{125}h_0^4h_6$ (46) $e_{171}f_0 \rightarrow e_{165}h_2g_1$
(35) $e_{128}D_3 \rightarrow e_{126}h_1D_3$ (47) $e_{172}e_0 \rightarrow e_{171}h_0e_0$
(36) $e_{139}c_1h_5 \rightarrow e_{135}h_2c_1h_5$ (48) $e_{174}h_0^3h_4 \rightarrow e_{173}h_0^4h_4$
(37) $e_{145}g_2 \rightarrow e_{143}h_1g_2$ (49) $e_{180}h_1c_0 \rightarrow e_{177}P^1h_2$
(38) $e_{148}h_0c_2 \rightarrow e_{147}h_0^2c_2$ (50) $e_{182}h_0^3h_3 \rightarrow e_{179}P^1h_1$
- (b) (1) $e_{11}c_1h_5h_7 \rightarrow e_7h_2c_1h_5h_7$ (31) $e_{59}h_2D_3(1) \rightarrow e_{45}h_2Q_3(1)$
(2) $e_{12}h_0h_2c_4 \rightarrow e_{11}h_1^3c_4$ (32) $e_{60}J_0 \rightarrow e_{59}h_0J_0$
(3) $e_{16}h_0f_3 \rightarrow e_{15}h_0^2f_3$ (33) $e_{62}h_0^3h_6^2 \rightarrow e_{61}h_0^4h_6^2$
(4) $e_{17}h_3e_3 \rightarrow e_9h_3^2e_3$ (34) $e_{81}g_2h_6 \rightarrow e_{79}h_1g_2h_6$
(5) $e_{17}g_2h_7 \rightarrow e_{15}h_1g_2h_7$ (35) $e_{84}h_0c_2h_6 \rightarrow e_{83}h_0^2c_2h_6$
(6) $e_{20}h_0c_2h_7 \rightarrow e_{19}h_0^2c_2h_7$ (36) $e_{85}f_1h_6 \rightarrow e_{77}h_3f_1h_6$
(7) $e_{21}f_1h_7 \rightarrow e_{13}h_3f_1h_7$ (37) $e_{88}h_0h_4c_3 \rightarrow e_{87}h_0^2h_4c_3$
(8) $e_{22}e_0h_5h_7 \rightarrow e_{15}d_0h_5h_7$ (38) $e_{92}p_0h_6 \rightarrow e_{91}h_0p_0h_6$
(9) $e_{23}e_1h_7 \rightarrow e_{11}f_0h_5h_7$ (39) $e_{93}d_1h_6 \rightarrow e_{62}H_1h_6$
(10) $e_{24}h_0e_3 \rightarrow e_{23}h_0^2e_3$ (40) $e_{96}h_0g_3 \rightarrow e_{95}h_0^2g_3$
(11) $e_{25}x_2 \rightarrow e_{23}h_1x_2$ (41) $e_{104}h_0f_2 \rightarrow e_{103}h_0^2f_2$
(12) $e_{28}p_0h_7 \rightarrow e_{27}h_0p_0h_7$ (42) $e_{105}h_2e_2 \rightarrow e_{101}h_2^2e_2$
(13) $e_{30}h_0^3h_5h_7 \rightarrow e_{29}h_0^4h_5h_7$ (43) $e_{105}g_1h_6 \rightarrow e_{103}h_1g_1h_6$
(14) $e_{32}V_0 \rightarrow e_{31}h_0V_0$ (44) $e_{107}f_0h_6 \rightarrow e_{101}h_2g_1h_6$
(15) $e_{33}g_3h_6 \rightarrow e_{31}h_1g_3h_6$ (45) $e_{108}e_0h_6 \rightarrow e_{107}h_0e_0h_6$
(16) $e_{41}h_3d_3 \rightarrow e_{21}h_2x_2$ (46) $e_{108}h_0e_2 \rightarrow e_{107}h_0^2e_2$
(17) $e_{41}g_1h_7 \rightarrow e_{39}h_1g_1h_7$ (47) $e_{109}x_1 \rightarrow e_{93}h_1^2g_3$
(18) $e_{43}f_0h_7 \rightarrow e_{37}h_2g_1h_7$ (48) $e_{110}h_0^3h_4h_6 \rightarrow e_{109}h_0^4h_4h_6$
(19) $e_{44}h_0p_2 \rightarrow e_{43}h_0^2p_2$ (49) $e_{112}h_4D_3 \rightarrow e_{110}h_1h_4D_3$
(20) $e_{44}e_0h_7 \rightarrow e_{43}h_0e_0h_7$ (50) $e_{113}h_3d_2 \rightarrow e_{105}d_1h_4^2$
(21) $e_{45}h_1p'_1 \rightarrow e_{30}h_1V_0$ (51) $e_{116}c_0h_1h_6 \rightarrow e_{113}P^1h_2h_6$
(22) $e_{46}h_0^3h_4h_7 \rightarrow e_{45}h_0^4h_4h_7$ (52) $e_{118}h_0^3h_3h_6 \rightarrow e_{115}P^1h_1h_6$
(23) $e_{48}h_0d_3 \rightarrow e_{47}h_0^2d_3$ (53) $e_{121}Q_3 \rightarrow e_{119}h_1Q_3$
(24) $e_{49}Q_3(1) \rightarrow e_{47}h_1Q_3(1)$ (54) $e_{121}n_1 \rightarrow e_{109}t_1$
(25) $e_{49}n_2 \rightarrow e_{47}h_1n_2$ (55) $e_{124}h_2D_3 \rightarrow e_{117}h_2Q_3$
(26) $e_{52}h_1c_0h_7 \rightarrow e_{49}P^1h_2h_7$ (56) $e_{126}h_0^3h_5^2 \rightarrow e_{125}h_0^4h_5^2$
(27) $e_{54}h_0^3h_3h_7 \rightarrow e_{51}P^1h_1h_7$ (57) $e_{136}D_1 \rightarrow e_{129}D_2$
(28) $e_{54}c_0h_6^2 \rightarrow e_{47}d_0h_6^2$ (58) $e_{137}g_1h_5 \rightarrow e_{135}h_1g_1h_5$
(29) $e_{55}h_3D_3(1) \rightarrow e_{39}\overline{\lambda_{39}}D_1(1)$ (59) $e_{140}e_0h_5 \rightarrow e_{139}h_0e_0h_5$
(30) $e_{59}H_1(1) \rightarrow e_{55}h_2H_1(1)$ (60) $e_{144}h_0g_2 \rightarrow e_{143}h_0^2g_2$
(61) $e_{148}h_0f_1 \rightarrow e_{147}h_0^2f_1$
(62) $e_{148}h_1c_0h_5 \rightarrow e_{145}P^1h_2h_5$

$$\begin{aligned}
(63) \quad & e_{149}h_3d_1 \rightarrow e_{140}h_0h_2g_2 & (69) \quad & e_{168}h_0g_1 \rightarrow e_{167}h_0^2g_1 \\
(64) \quad & e_{150}h_0^3h_3h_5 \rightarrow e_{147}P^1h_1h_5 & (70) \quad & e_{170}h_1e_0 \rightarrow e_{164}h_2^2e_0 \\
(65) \quad & e_{151}x_0 \rightarrow e_{138}h_1e_0h_5 & (71) \quad & e_{173}h_1d_0 \rightarrow e_{171}h_1^2d_0 \\
(66) \quad & e_{153}h_2d_1 > e_{149}h_2^2d_1 & (72) \quad & e_{174}h_0d_0 \rightarrow e_{173}h_0^2d_0 \\
(67) \quad & e_{158}h_0^3h_4^2 \rightarrow e_{155}q_0 & (73) \quad & e_{188}h_0^5 \rightarrow e_{187}h_0^6 \\
(68) \quad & e_{164}h_1c_0h_4 \rightarrow e_{161}h_2^2g_1
\end{aligned}$$

$$\begin{aligned}
(c) \quad (1) \quad & e_9h_2^2c_4 \longleftrightarrow \widehat{h}_5h_5D_3(1) & (3) \quad & e_{13}h_2f_3 \longleftrightarrow \widehat{h}_4h_1f_3 \\
& & (4) \quad & e_{29}d_1h_7 \longleftrightarrow \gamma_{61}h_7 \\
(2) \quad & e_{13}h_1^2c_4 \longleftrightarrow \widehat{h}_4h_0^2c_4 & (5) \quad & e_{47}T_0 \longleftrightarrow \widehat{g}_4h_0^2
\end{aligned}$$

$$\begin{aligned}
(3.38) \quad (a) \quad (1) \quad & e_2g_4 \rightarrow e_1h_0g_4 & (28) \quad & e_{106}f_2 \rightarrow e_{105}h_0f_2 \\
(2) \quad & e_{14}h_2c_4 \rightarrow e_{13}h_0h_2c_4 & (29) \quad & e_{108}c_1h_6 \rightarrow e_{94}d_1h_6 \\
(3) \quad & e_{16}h_1c_4 \rightarrow e_{14}h_1^2c_4 & (30) \quad & e_{110}e_2 \rightarrow e_{109}h_0e_2 \\
(4) \quad & e_{18}f_3 \rightarrow e_{17}h_0f_3 & (31) \quad & e_{112}h_0^2h_4h_6 \rightarrow e_{111}h_0^3h_4h_6 \\
(5) \quad & e_{18}h_4^3h_7 \rightarrow e_{17}h_0h_4^3h_7 & (32) \quad & e_{120}h_0^2h_3h_6 \rightarrow e_{119}h_0^3h_3h_6 \\
(6) \quad & e_{22}c_2h_7 \rightarrow e_{21}h_0c_2h_7 & (33) \quad & e_{120}p_1 \rightarrow e_{118}h_1p_1 \\
(7) \quad & e_{24}h_1h_3h_5h_7 \rightarrow e_{22}h_2^3h_5h_7 & (34) \quad & e_{121}p'_0 \rightarrow e_{119}p'_0h_1 \\
(8) \quad & e_{26}h_2^2h_5h_7 \rightarrow e_{23}c_0h_5h_7 & (35) \quad & e_{122}d_2 \rightarrow e_{121}h_0d_2 \\
(9) \quad & e_{26}e_3 \rightarrow e_{25}h_0e_3 & (36) \quad & e_{128}h_0^2h_5^2 \rightarrow e_{127}h_0^3h_5^2 \\
(10) \quad & e_{32}h_1h_4^2h_7 \rightarrow e_{29}p_0h_7 & (37) \quad & e_{129}D_3 \rightarrow e_{127}h_1D_3 \\
(11) \quad & e_{32}h_0^2h_5h_7 \rightarrow e_{31}h_0^3h_5h_7 & (38) \quad & e_{140}c_1h_5 \rightarrow e_{125}h_2D_3 \\
(12) \quad & e_{44}c_1h_7 \rightarrow e_{30}d_1h_7 & (39) \quad & e_{142}h_3c_2 \rightarrow e_{127}H_1 \\
(13) \quad & e_{46}p_2 \rightarrow e_{45}h_0p_2 & (40) \quad & e_{146}g_2 \rightarrow e_{145}h_0g_2 \\
(14) \quad & e_{48}h_0^2h_4h_7 \rightarrow e_{47}h_0^3h_4h_7 & (41) \quad & e_{150}f_1 \rightarrow e_{149}h_0f_1 \\
(15) \quad & e_{48}p'_1 \rightarrow e_{46}h_1p'_1 & (42) \quad & e_{152}e_1 \rightarrow e_{150}h_1e_1 \\
(16) \quad & e_{50}d_3 \rightarrow e_{49}h_0d_3 & (43) \quad & e_{152}h_0^2h_3h_5 \rightarrow e_{151}h_0^3h_3h_5 \\
(17) \quad & e_{56}h_0^2h_3h_7 \rightarrow e_{55}h_0^3h_3h_7 & (44) \quad & e_{156}h_0^2h_2h_5 \rightarrow e_{124}h_1^3h_5^2 \\
(18) \quad & e_{56}h_1h_3h_6^2 \rightarrow e_{54}h_1^2h_3h_6^2 & (45) \quad & e_{160}h_0^2h_4^2 \rightarrow e_{159}h_0^3h_4^2 \\
(19) \quad & e_{58}h_2^2h_6^2 \rightarrow e_{55}c_0h_6^2 & (46) \quad & e_{168}h_2c_1 \rightarrow e_{158}n_0 \\
(20) \quad & e_{64}h_0^2h_6^2 \rightarrow e_{63}h_0^3h_6^2 & (47) \quad & e_{170}g_1 \rightarrow e_{169}h_0g_1 \\
(21) \quad & e_{64}D_3(1) \rightarrow e_{61}J_0 & (48) \quad & e_{172}h_1^2h_4 \rightarrow e_{141}e_0h_5 \\
(22) \quad & e_{86}c_2h_6 \rightarrow e_{85}h_0c_2h_6 & (49) \quad & e_{172}f_0 \rightarrow e_{171}h_0f_0 \\
(23) \quad & e_{90}h_4c_3 \rightarrow e_{89}h_0h_4c_3 & (50) \quad & e_{173}e_0 \rightarrow e_{165}h_3e_0 \\
(24) \quad & e_{96}h_1h_5^3 \rightarrow e_{93}p_0h_6 & (51) \quad & e_{176}d_0 \rightarrow e_{175}h_0d_0 \\
(25) \quad & e_{98}g_3 \rightarrow e_{97}h_0g_3 & (52) \quad & e_{181}h_1c_0 \rightarrow e_{174}h_1d_0 \\
(26) \quad & e_{104}h_1c_3 \rightarrow e_{102}h_1^2c_3 & (53) \quad & e_{190}h_0^4 \rightarrow e_{189}h_0^5 \\
(27) \quad & e_{106}h_3^3h_6 \rightarrow e_{103}c_0h_4h_6
\end{aligned}$$

$$\begin{aligned}
(b) \quad (1) \quad & e_{10}h_2c_4 \rightarrow e_7c_0c_4 & (4) \quad & e_{16}h_1f_3 \rightarrow e_{14}h_1^2f_3 \\
(2) \quad & e_{14}h_5p_2 \rightarrow e_{13}h_0h_5p_2 & (5) \quad & e_{18}g_2h_7 \rightarrow e_{17}h_0g_2h_7 \\
(3) \quad & e_{16}h_0^2c_4 \rightarrow e_{15}h_0^3c_4 & (6) \quad & e_{22}f_1h_7 \rightarrow e_{21}h_0f_1h_7
\end{aligned}$$

$$\begin{array}{ll}
(7) e_{24}e_1h_7 \rightarrow e_{22}h_1e_1h_7 & (40) e_{104}h_1f_2 \rightarrow e_{102}h_1^2f_2 \\
(8) e_{24}h_0^2h_3h_5h_7 \rightarrow e_{23}h_0^3h_3h_5h_7 & (41) e_{106}h_2e_2 \rightarrow e_{102}h_2^2e_2 \\
(9) e_{24}h_1e_3 \rightarrow e_{22}h_1^2e_3 & (42) e_{106}g_1h_6 \rightarrow e_{105}h_0g_1h_6 \\
(10) e_{32}h_5D_3(1) \rightarrow e_{29}h_5J_0 & (43) e_{108}f_0h_6 \rightarrow e_{107}h_0f_0h_6 \\
(11) e_{32}h_0^2h_4^2h_7 \rightarrow e_{31}h_0^3h_4^2h_7 & (44) e_{109}e_0h_6 \rightarrow e_{101}h_3e_0h_6 \\
(12) e_{33}V_0 \rightarrow e_{31}h_1V_0 & (45) e_{110}x_1 \rightarrow e_{109}h_0x_1 \\
(13) e_{34}g_3h_6 \rightarrow e_{33}h_0g_3h_6 & (46) e_{112}d_0h_6 \rightarrow e_{111}h_0d_0h_6 \\
(14) e_{40}h_2c_1h_7 \rightarrow e_{30}n_0h_7 & (47) e_{113}h_4D_3 \rightarrow e_{111}h_1h_4D_3 \\
(15) e_{42}h_2p_2 \rightarrow e_{39}\lambda_{40}D_1(1) & (48) e_{114}h_3d_2 \rightarrow e_{113}h_0h_3d_2 \\
(16) e_{42}g_1h_7 \rightarrow e_{41}h_0g_1h_7 & (49) e_{117}h_1c_0h_6 \rightarrow e_{110}h_1d_0h_6 \\
(17) e_{44}f_0h_7 \rightarrow e_{43}h_0f_0h_7 & (50) e_{120}h_0p'_0 \rightarrow e_{119}h_0^2p'_0 \\
(18) e_{44}h_1^3h_4h_7 \rightarrow e_{31}e_0h_5h_7 & (51) e_{122}n_1 \rightarrow e_{121}h_0n_1 \\
(19) e_{45}e_0h_7 \rightarrow e_{38}h_3e_0h_7 & (52) e_{122}Q_3 \rightarrow e_{121}h_0Q_3 \\
(20) e_{48}h_1d_3 \rightarrow e_{45}h_4J_0 & (53) e_{126}h_0^4h_6 \rightarrow e_{125}h_0^5h_6 \\
(21) e_{48}d_0h_7 \rightarrow e_{47}h_0d_0h_7 & (54) e_{136}h_2c_1h_5 \rightarrow e_{126}h_5n_0 \\
(22) e_{48}T_0 \rightarrow e_{47}h_0T_0 & (55) e_{137}D_1 \rightarrow e_{127}A \\
(23) e_{50}Q_3(1) \rightarrow e_{49}h_0Q_3(1) & (56) e_{138}g_1h_5 \rightarrow e_{137}h_0g_1h_5 \\
(24) e_{50}n_2 \rightarrow e_{49}h_0n_2 & (57) e_{140}f_0h_5 \rightarrow e_{139}h_0f_0h_5 \\
(25) e_{53}h_1c_0h_7 \rightarrow e_{46}h_1d_0h_7 & (58) e_{142}h_2g_2 \rightarrow e_{141}h_0h_2g_2 \\
(26) e_{56}h_0^2h_3h_6^2 \rightarrow e_{55}h_0^3h_3h_6^2 & (59) e_{144}h_1g_2 \rightarrow e_{136}h_1g_1h_5 \\
(27) e_{56}h_3D_3(1) \rightarrow e_{53}h_3J_0 & (60) e_{144}d_0h_5 \rightarrow e_{143}h_0d_0h_5 \\
(28) e_{62}h_0^4h_7 \rightarrow e_{61}h_0^5h_7 & (61) e_{148}h_0^2c_2 \rightarrow e_{127}A' \\
(29) e_{64}K_0 \rightarrow e_{63}h_0K_0 & (62) e_{149}h_1c_0h_5 \rightarrow e_{142}h_1d_0h_5 \\
(30) e_{78}c_2h_3h_6 \rightarrow e_{63}H_1h_6 & (63) e_{152}x_0 \rightarrow e_{151}h_0x_0 \\
(31) e_{82}g_2h_6 \rightarrow e_{81}h_0g_2h_6 & (64) e_{154}h_2d_1 \rightarrow e_{150}h_2^2d_1 \\
(32) e_{86}f_1h_6 \rightarrow e_{85}h_0f_1h_6 & (65) e_{156}h_1d_1 \rightarrow e_{125}h_1H_1 \\
(33) e_{88}e_1h_6 \rightarrow e_{86}h_1e_1h_6 & (66) e_{158}h_0^4h_5 \rightarrow e_{157}h_0^5h_5 \\
(34) e_{90}h_4f_2 \rightarrow e_{89}h_0h_4f_2 & (67) e_{166}h_2g_1 \rightarrow e_{165}h_0h_2g_1 \\
(35) e_{94}h_2g_3 \rightarrow e_{93}h_0h_2g_3 & (68) e_{168}h_1g_1 \rightarrow e_{162}h_2^2g_1 \\
(36) e_{96}h_1g_3 \rightarrow e_{94}h_1^2g_3 & (69) e_{172}h_0e_0 \rightarrow e_{171}h_0^2e_0 \\
(37) e_{96}h_0^2h_5^3 \rightarrow e_{95}h_0^3h_5^3 & (70) e_{174}h_0^4h_4 \rightarrow e_{173}h_0^5h_4 \\
(38) e_{104}h_0^2c_3 \rightarrow e_{103}h_0^3c_3 & (71) e_{178}P^1h_2 \rightarrow e_{177}h_0P^1h_2 \\
(39) e_{104}h_2c_1h_6 \rightarrow e_{94}h_6n_0 & (72) e_{180}P^1h_1 \rightarrow e_{178}h_1P^1h_1
\end{array}$$

$$\begin{array}{ll}
(c) (1) e_1D_3h_7 \longleftrightarrow \widehat{h}_1D_3h_7 & (5) e_{60}H_1(1) \longleftrightarrow \widehat{h}_7H_1 \\
(2) e_{12}c_1h_5h_7 \longleftrightarrow \xi_{31}h_5h_7 & (6) e_{60}h_2D_3(1) \longleftrightarrow \widehat{h}_7h_1D_3 \\
(3) e_{14}h_3c_2h_7 \longleftrightarrow \widehat{H}_1h_7 & (7) e_{60}h_1^3h_6^2 \longleftrightarrow \widehat{h}_6h_0^3h_6^2 \\
(4) e_{28}h_1^3h_5h_7 \longleftrightarrow \widehat{h}_5h_0^3h_5h_7 &
\end{array}$$

The remaining (3.2) (a) ((3.2) (8) (a), (3.2) (9) (a)) follow from the results in (3.36) (c) and (3.37) (c) above. This proves the remaining (3.2) (a).

To prove the remaining (3.2) (c) ((3.2) (8) (c), (3.2) (9) (c)), (3.34), and (3.1) (9) we recall in (3.39) and (3.40) below from [8, 16] an important notion about the spectral sequence $\{E_r^{i,s,t}\}$ in (3.32).

(3.39) Given a (possibly zero) class $\alpha \in \text{Ext}_{\mathcal{A}}^{s,t}$ with $t - s > 0$ and $s \geq 1$. Let x be a cycle in $\Lambda^{s,t-s}$ representing the class α with $x \neq 0$. Since $t - s > 0$ we have $x \equiv \lambda_j \lambda_I \pmod{\Lambda(j-1)}$ for some $j \geq 1$, where $\lambda_I \in \Lambda^{s-1,t-s-j}$ is a nonzero chain such that for any admissible monomial $\lambda_{j_1} \cdots \lambda_{j_{s-1}}$ appearing in the admissible expansion of λ_I , $\lambda_j \lambda_{j_1} \cdots \lambda_{j_{s-1}}$ is admissible.

Since x is a cycle and $\Lambda(j-1)$ is a subcomplex of Λ , λ_I in (3.39) is actually a cycle. Recall the map $\text{Ext}_{\mathcal{A}}^{s+1,t+1} \xrightarrow{\phi_*} \text{Ext}_{\mathcal{A}}^{s,t}(P)$ in (3.30). It is easy to see from (3.30) (2) that

$$\phi(x) \equiv e_j \lambda_I \pmod{F(j-1)}.$$

Since λ_I is a cycle, $e_j \{\lambda_I\}$ is an element in $E_1^{*,s-1,*}$ in the spectral sequence considered in (3.32). And since $\phi(x)$ is a cycle, $e_j \{\lambda_I\}$ is an infinite cycle in the spectral sequence which may not be a nontrivial one. To show that $\alpha \neq 0$ it suffices to show that $\phi_*(\alpha) \neq 0$. It is not difficult to see that there are two cases for showing this.

(3.40) (1) If $e_j \{\lambda_I\}$ is non-zero in the spectral sequence, then α is a non-zero class. In this case we will use $\overline{\phi}(\alpha)$ to denote $e_j \{\lambda_I\}$, and use the correspondence

$$\alpha = \{x\} = \{\overline{\lambda_j \lambda_I}\} \rightarrow e_j \{\lambda_I\} = \overline{\phi}(\alpha),$$

to indicate that the class $\phi_*(\alpha)$ is represented by the infinite cycle $e_j \{\lambda_I\} = \overline{\phi}(\alpha)$ (in the spectral sequence in (3.32) for $\text{Ext}_{\mathcal{A}}^{**}(P)$).

(2) If $e_j \{\lambda_I\}$ is a boundary in the spectral sequence, but $\phi(x)$ is homologous to a cycle z such that $z \equiv e_k \lambda_J \pmod{F(k-1)}$ for some k with $1 \leq k < j$ and some $\lambda_J \in \Lambda^{s-1,*}$, and such that $e_k \{\lambda_J\}$ is non-zero in the spectral sequence, then α is also a non-zero class. In this case we will use $\tilde{\phi}(\alpha)$ to denote $e_k \{\lambda_J\}$, and use the correspondence

$$\alpha = \{x\} = \{\overline{\lambda_j \lambda_I}\} \rightarrow e_k \{\lambda_J\} = \tilde{\phi}(\alpha).$$

to indicate that the class $\phi_*(\alpha)$ is represented by the infinite cycle $e_k \{\lambda_J\} = \tilde{\phi}(\alpha)$.

(3.41) and (3.42) below give the correspondences (defined in (3.40)) for the elements in the remaining (3.2) (c).

$$\mathbf{(3.41)} \quad (1) \quad h_5 V_0 \rightarrow e_{30} D_3(1) h_5 = \overline{\phi}(h_5 V_0)$$

$$\mathbf{(3.42)} \quad (1) \quad h_0^2 g_4 \rightarrow e_{47} T_0 = \overline{\phi}(h_0^2 g_4)$$

$$(2) \quad h_5^2 D_3(1) \rightarrow e_9 h_2^2 c_4 = \tilde{\phi}(h_5^2 D_3(1))$$

(3.41) shows that $\phi_*(h_5 V_0) \neq 0$, and (3.42) shows that $\phi_*(h_0^2 g_4), \phi_*(h_5^2 D_3(1))$ are linearly independent. This proves the remaining (3.2) (c).

To prove (3.34), we list in (3.43) below each $\beta \in \text{Ext}_{\mathcal{A}}^{6,*}$ such that $e_i \beta$ appears in the targets of the (b)-differentials in (3.36) and (3.37). The calculations for these correspondences are lengthy and will not be given in this paper.

- (3.43) (1) $A \rightarrow e_9 D_1 = \overline{\phi}(A)$
(2) $A' \rightarrow e_{20} h_0^2 c_2 = \overline{\phi}(A')$
(3) $\overline{\lambda_{19} D_1} \rightarrow e_{19} D_1 = \overline{\phi}(\overline{\lambda_{19} D_1})$
(4) $\overline{\lambda_{39} D_1(1)} \rightarrow e_{39} D_1(1) = \overline{\phi}(\overline{\lambda_{39} D_1(1)})$
(5) $\overline{\lambda_{40} D_1(1)} \rightarrow e_{40} D_1(1) = \overline{\phi}(\overline{\lambda_{40} D_1(1)})$
(6) $D_2 \rightarrow e_{28} h_0^3 h_4^2 = \overline{\phi}(D_2)$
(7) $q_0 \rightarrow e_2 h_0^2 h_4^2 = \tilde{\phi}(q)$
(8) $t_0 \rightarrow e_5 n_0 = \tilde{\phi}(t_0)$
(9) $h_0^3 h_3 h_5 h_7 \rightarrow e_{84} h_1^3 h_4 h_6 = \overline{\phi}(h_0^3 h_3 h_5 h_7)$
(10) $h_0^3 h_3 h_6^2 \rightarrow e_{53} h_1^3 h_4 h_6 = \overline{\phi}(h_0^3 h_3 h_6^2)$
(11) $h_0^3 h_5^3 \rightarrow e_{28} h_1^3 h_5^2 = \overline{\phi}(h_0^3 h_4^2 h_6)$
(12) $h_0^3 h_4^2 h_7 \rightarrow e_{92} h_1^3 h_5^2 = \overline{\phi}(h_0^3 h_4^2 h_7)$
(13) $h_0^4 h_4 h_6 \rightarrow e_{40} h_0^3 h_3 h_5 = \tilde{\phi}(h_0^4 h_4 h_6)$
(14) $h_0^4 h_4 h_7 \rightarrow e_{104} h_0^3 h_3 h_5 = \tilde{\phi}(h_0^4 h_4 h_7)$
(15) $h_0^4 h_5^2 \rightarrow e_{24} h_0^3 h_3 h_5 = \tilde{\phi}(h_0^4 h_5^2)$
(16) $h_0^4 h_5 h_7 \rightarrow e_{88} h_0^3 h_3 h_6 = \tilde{\phi}(h_0^4 h_5 h_7)$
(17) $h_0^4 h_6^2 \rightarrow e_{56} h_0^3 h_3 h_6 = \tilde{\phi}(h_0^4 h_6^2)$
(18) $h_0^5 h_4 \rightarrow e_6 P^1 h_1 = \tilde{\phi}(h_0^5 h_4)$
(19) $h_0^5 h_5 \rightarrow e_{22} P^1 h_1 = \tilde{\phi}(h_0^5 h_5)$
(20) $h_0^5 h_6 \rightarrow e_{54} P^1 h_1 = \tilde{\phi}(h_0^5 h_6)$
(21) $h_0^5 h_7 \rightarrow e_{118} P^1 h_1 = \tilde{\phi}(h_0^5 h_7)$
(22) $h_0^2 c_2 h_6 \rightarrow e_{17} h_1^2 c_3 = \overline{\phi}(h_0^2 c_2 h_6)$
(23) $h_0^2 c_2 h_7 \rightarrow e_{81} h_1^2 c_3 = \overline{\phi}(h_0^2 c_2 h_7)$
(24) $h_0^2 h_4 c_3 \rightarrow e_{13} h_1^2 c_3 = \overline{\phi}(h_0^2 h_4 c_3)$
(25) $h_0^3 c_3 \rightarrow e_{20} h_1^3 h_5^2 = \overline{\phi}(h_0^3 c_3)$
(26) $h_0^3 c_4 \rightarrow e_{44} h_1^3 h_6^2 = \overline{\phi}(h_0^3 c_4)$
(27) $h_1^3 c_4 \rightarrow e_{41} h_2^3 h_6^2 = \overline{\phi}(h_1^3 c_4)$
(28) $h_1 c_0 h_5^2 \rightarrow e_{18} h_2 c_1 h_5 = \overline{\phi}(h_1 c_0 h_5^2)$
(29) $h_1 c_0 h_6^2 \rightarrow e_{50} h_2 c_1 h_6 = \overline{\phi}(h_1 c_0 h_6^2)$
(30) $h_2 c_1 h_6^2 \rightarrow e_{37} h_3 c_2 h_6 = \overline{\phi}(h_2 c_1 h_6^2)$
(31) $h_1 c_0 h_4 h_6 \rightarrow e_{34} h_2 c_1 h_5 = \overline{\phi}(h_1 c_0 h_4 h_6)$
(32) $h_1 c_0 h_4 h_7 \rightarrow e_{98} h_2 c_1 h_5 = \overline{\phi}(h_1 c_0 h_4 h_7)$
(33) $h_1 c_0 h_5 h_7 \rightarrow e_{82} h_2 c_1 h_6 = \overline{\phi}(h_1 c_0 h_5 h_7)$
(34) $h_2 c_1 h_5 h_7 \rightarrow e_{69} h_3 c_2 h_6 = \overline{\phi}(h_2 c_1 h_5 h_7)$
(35) $c_0 c_4 \rightarrow e_{36} c_1 h_6^2 = \overline{\phi}(c_0 c_4)$
(36) $h_0^2 d_3 = h_4 K_0 + D_1(1) h_5 \rightarrow e_{15} K_0 = \tilde{\phi}(e_{15} K_0)$
(37) $h_0^2 d_0 = h_2 P^1 h_2 \rightarrow e_3 P^1 h_2 = \overline{\phi}(h_2 P^1 h_2)$

- (38) $h_1^2 d_0 \rightarrow e_1 h_1 d_0 = \tilde{\phi}(h_1^2 d_0)$
(39) $h_2^2 d_1 \rightarrow e_3 h_2 d_1 = \tilde{\phi}(h_2^2 d_1)$
(40) $d_0 h_6^2 \rightarrow e_{45} d_1 h_6 = \bar{\phi}(d_0 h_6^2)$
(41) $h_0 d_0 h_5 \rightarrow e_{12} h_1 d_1 = \bar{\phi}(h_0 d_0 h_5)$
(42) $h_0 d_0 h_6 \rightarrow e_{44} h_1 d_1 = \bar{\phi}(h_0 d_0 h_6)$
(43) $h_0 d_0 h_7 \rightarrow e_{108} h_1 d_1 = \bar{\phi}(h_0 d_0 h_7)$
(44) $h_1 d_1 h_6 = h_0 p_0 h_6 \rightarrow e_{25} h_2 d_2 = \bar{\phi}(h_1 d_1 h_6)$
(45) $h_1 d_1 h_7 = h_0 p_0 h_7 \rightarrow e_{89} h_2 d_2 = \bar{\phi}(h_1 d_1 h_7)$
(46) $h_1 d_0 h_5 \rightarrow e_{11} h_2 d_1 = \bar{\phi}(h_1 d_0 h_5)$
(47) $h_1 d_0 h_6 \rightarrow e_{43} h_2 d_1 = \bar{\phi}(h_1 d_0 h_6)$
(48) $h_1 d_0 h_7 \rightarrow e_{107} h_2 d_1 = \bar{\phi}(h_1 d_0 h_7)$
(49) $d_0 h_5 h_7 \rightarrow e_{77} d_1 h_6 = \bar{\phi}(d_0 h_5 h_7)$
(50) $h_0 h_3 d_2 \rightarrow e_4 h_2 d_2 = \tilde{\phi}(h_0 h_3 d_2)$
(51) $h_1 h_4 d_3 \rightarrow e_9 h_3 d_3 = \tilde{\phi}(h_1 h_4 d_3)$
(52) $h_0 h_4 d_3 \rightarrow e_{14} h_1 d_3 = \tilde{\phi}(h_0 h_4 d_3)$
(53) $h_0 h_5 d_3 \rightarrow e_{30} h_1 d_3 = \bar{\phi}(h_0 h_5 d_3)$
(54) $h_0^2 e_0 = h_0 h_2 d_0 \rightarrow e_2 h_1 d_0 = \bar{\phi}(h_0 h_2 d_0)$
(55) $h_1^2 e_1 = h_1 h_3 d_1 = h_0^2 f_1 = h_0 h_3 p_0 \rightarrow e_3 x_0 = \tilde{\phi}(h_1 h_3 d_1)$
(56) $h_2^2 e_2 = h_2 h_4 d_2 = h_1^2 f_2 = h_1 h_4 p_1 \rightarrow e_7 x_1 = \tilde{\phi}(h_2 h_4 d_2)$
(57) $h_3^2 e_3 = h_3 h_5 d_3 = h_2^2 f_3 = h_2 h_5 p_2 \rightarrow e_{15} x_2 = \tilde{\phi}(h_3 h_5 d_3)$
(58) $h_0 h_3 g_1 = h_2^2 e_0 \rightarrow e_2 h_1 g_1 = \bar{\phi}(h_2^2 e_0)$
(59) $h_0^2 e_2 = h_1 x_1 \rightarrow e_{27} h_2 c_1 h_5 = \bar{\phi}(h_0^2 e_2)$
(60) $h_0^2 e_3 \rightarrow e_{61} f_1 h_6 = \tilde{\phi}(h_0^2 e_3)$
(61) $h_1^2 e_3 = h_2 x_2 \rightarrow e_{55} h_3 c_2 h_6 = \bar{\phi}(h_1^2 e_3)$
(62) $h_0 e_0 h_5 = h_2 d_0 h_5 \rightarrow e_9 h_3 d_1 = \bar{\phi}(h_0 e_0 h_5)$
(63) $h_0 e_0 h_6 = h_2 d_0 h_6 \rightarrow e_{41} h_3 d_1 = \bar{\phi}(h_0 e_0 h_6)$
(64) $h_0 e_0 h_7 = h_2 d_0 h_7 \rightarrow e_{105} h_3 d_1 = \bar{\phi}(h_0 e_0 h_7)$
(65) $h_1 e_1 h_6 = h_3 d_1 h_6 \rightarrow e_{19} h_4 d_2 = \bar{\phi}(h_1 e_1 h_6)$
(66) $h_1 e_1 h_7 = h_3 d_1 h_7 \rightarrow e_{85} h_4 d_2 = \bar{\phi}(h_1 e_1 h_7)$
(67) $h_2 e_1 h_6 = h_1 f_1 h_6 \rightarrow e_{17} h_3 e_2 = \bar{\phi}(h_2 e_1 h_6)$
(68) $h_2 e_1 h_7 = h_1 f_1 h_7 \rightarrow e_{81} h_3 e_2 = \bar{\phi}(h_2 e_1 h_7)$
(69) $h_3 e_0 h_6 \rightarrow e_{34} c_1 h_2 h_5 = \bar{\phi}(h_3 e_0 h_6)$
(70) $h_3 e_0 h_7 \rightarrow e_{98} c_1 h_2 h_5 = \bar{\phi}(h_3 e_0 h_7)$
(71) $e_0 h_5 h_7 \rightarrow e_{74} e_1 h_6 = \bar{\phi}(e_0 h_5 h_7)$
(72) $h_0^2 f_2 = h_0 h_4 p'_0 \rightarrow e_{14} h_1 p'_0 = \bar{\phi}(h_0^2 f_2)$
(73) $h_0^2 f_3 = h_5 T_0 \rightarrow e_{31} T_0 = \bar{\phi}(h_5 T_0)$
(74) $h_1^2 f_3 = h_1 h_5 p'_1 \rightarrow e_{29} h_2 p'_1 = \bar{\phi}(h_1^2 f_3)$

- (75) $h_0f_0h_5 = h_1e_0h_5 \rightarrow e_8h_2e_1 = \overline{\phi}(h_0f_0h_5)$
(76) $h_0f_0h_6 = h_1e_0h_6 \rightarrow e_{40}h_2e_1 = \overline{\phi}(h_0f_0h_6)$
(77) $h_0f_0h_7 = h_1e_0h_7 \rightarrow e_{104}h_2e_1 = \overline{\phi}(h_0f_0h_7)$
(78) $h_0h_4f_2 \rightarrow e_{14}h_1f_2 = \overline{\phi}(h_0h_4f_2)$
(79) $h_1h_4f_2 \rightarrow e_{13}h_2f_2 = \overline{\phi}(h_1h_4f_2)$
(80) $h_0f_1h_6 = h_3p_0h_6 \rightarrow e_{18}p_1h_4 = \overline{\phi}(h_0f_1h_6)$
(81) $h_0f_1h_7 = h_3p_0h_7 \rightarrow e_{82}p_1h_4 = \overline{\phi}(h_0f_1h_7)$
(82) $f_0h_5h_7 \rightarrow e_{73}f_1h_6 = \overline{\phi}(f_0h_5h_7)$
(83) $h_2f_0h_7 = h_1g_1h_7 \rightarrow e_{101}h_3f_1 = \overline{\phi}(h_2f_0h_7)$
(84) $h_3f_1h_6 = h_2g_2h_6 = h_2g_3h_4 = e_1h_3^2 \rightarrow e_{11}h_4f_2 = \overline{\phi}(h_3f_1h_6)$
(85) $h_3f_1h_7 = Sq^0(h_2f_0h_6) \rightarrow e_{75}h_4f_2 = \overline{\phi}(h_3f_1h_7)$
(86) $h_0^2g_1 = h_2^2d_0 = h_0h_2e_0 \rightarrow e_2h_1e_0 = \tilde{\phi}(h_2^2d_0)$
(87) $h_0^2g_2 = h_3x_0 \rightarrow e_7x_0 = \overline{\phi}(h_3x_0)$
(88) $h_0^2g_3 \rightarrow e_{22}h_1p'_0 = \tilde{\phi}(h_0^2g_3)$
(89) $h_1^2g_3 = Sq^0(h_0^2g_2) \rightarrow e_{15}x_1 = \overline{\phi}(h_4x_1)$
(90) $h_4P^1h_2 = h_2^2g_1 \rightarrow e_3h_2g_1 = \overline{\phi}(h_2^2g_1)$
(91) $h_0g_1h_5 \rightarrow e_6h_1g_2 = \overline{\phi}(h_0g_1h_5)$
(92) $h_1h_4g_3 = h_1g_2h_6 = e_1h_3h_6 \rightarrow e_{13}h_2g_3 = \overline{\phi}(h_1h_4g_3)$
(93) $h_1g_2h_7 = h_3e_1h_7 \rightarrow e_{77}h_4e_2 = \overline{\phi}(h_1g_2h_7)$
(94) $h_0g_1h_6 = h_2e_0h_6 \rightarrow e_{38}e_1h_3 = \overline{\phi}(h_0g_1h_6)$
(95) $h_0g_1h_7 = h_2e_0h_7 \rightarrow e_{102}e_1h_3 = \overline{\phi}(h_0g_1h_7)$
(96) $h_0g_2h_6 \rightarrow e_{14}h_1g_3 = \overline{\phi}(h_0g_2h_6)$
(97) $h_0g_2h_7 \rightarrow e_{78}h_1g_3 = \overline{\phi}(h_0g_2h_7)$
(98) $h_1g_3h_6 = h_1h_4d_3 \rightarrow e_9h_3d_3 = \tilde{\phi}(h_1h_4d_3)$
(99) $h_0g_3h_6 = h_0h_4d_3 \rightarrow e_{14}h_1d_3 = \tilde{\phi}(h_0h_4d_3)$
(100) $h_0h_2g_2 = h_0h_3f_1 \rightarrow e_6h_1f_1 = \tilde{\phi}(h_0h_2g_2)$
(101) $h_1g_1h_5 \rightarrow e_5h_2g_2 = \overline{\phi}(h_1g_1h_5)$
(102) $h_1g_1h_6 = h_2f_0h_6 \rightarrow e_{37}h_3f_1 = \overline{\phi}(h_1g_1h_6)$
(103) $h_1g_1h_7 = h_2f_0h_7 \rightarrow e_{101}h_3f_1 = \overline{\phi}(h_1g_1h_7)$
(104) $h_2g_1h_6 \rightarrow e_{35}h_3g_2 = \overline{\phi}(h_2g_1h_6)$
(105) $h_2g_1h_7 \rightarrow e_{99}h_3g_2 = \overline{\phi}(h_2g_1h_7)$
(106) $h_0h_2g_1 \rightarrow e_2h_1g_2 = \overline{\phi}(h_0h_2g_1)$
(107) $h_0h_2g_3 = h_0h_4e_2 \rightarrow e_{12}h_2e_2 = \overline{\phi}(h_0h_2g_3)$
(108) $h_0^2p_2 \rightarrow e_{49}h_4e_2 = \tilde{\phi}(h_0^2p_2)$
(109) $h_0h_5p_2 \rightarrow e_{28}h_2p_2 = \tilde{\phi}(h_0h_5p_2)$
(110) $h_4^2D_3 \rightarrow e_4h_1^2c_3 = \tilde{\phi}(h_4^2D_3)$
(111) $h_3^2D_3(1) \rightarrow e_7h_3D_3(1) = \tilde{\phi}(h_3^2D_3(1))$

- (112) $h_1h_4D_3 \rightarrow e_1h_4D_3 = \tilde{\phi}(h_1h_4D_3)$
(113) $h_0^2p'_0 \rightarrow e_{24}h_3e_1 = \tilde{\phi}(h_0^2p'_0)$
(114) $h_0J_0 \rightarrow e_{57}h_1p_1 = \tilde{\phi}(h_0J_0)$
(115) $h_3J_0 \rightarrow e_{50}h_4p_0 = \tilde{\phi}(h_3J_0)$
(116) $h_4J_0 \rightarrow e_{15}J_0 = \tilde{\phi}(h_4J_0)$
(117) $h_5J_0 \rightarrow e_{30}H_1(1) = \tilde{\phi}(h_5J_0)$
(118) $h_0T_0 \rightarrow e_{76}h_1^3h_5^2 = \tilde{\phi}(h_0T_0)$
(119) $h_0K_0 \rightarrow e_{44}h_0^2h_2h_4h_6 = \tilde{\phi}(h_0K_0)$
(120) $h_0n_1 = h_4D_1 = h_2^2D_3 \rightarrow e_3h_3D_3 = \tilde{\phi}(h_4D_1)$
(121) $h_0n_2 \rightarrow e_{30}D_1(1) = \tilde{\phi}(h_0n_2)$
(122) $h_1n_2 = h_5D_1(1) = h_3^2D_3(1) \rightarrow e_7h_4D_3(1) = \tilde{\phi}(h_5D_1(1))$
(123) $h_0Q_3 \rightarrow e_{22}h_1g_2 = \tilde{\phi}(h_0Q_3)$
(124) $h_0Q_3(1) \rightarrow e_{46}h_1g_3 = \tilde{\phi}(h_0Q_3(1))$
(125) $h_1Q_3 \rightarrow e_{21}h_2g_2 = \tilde{\phi}(h_1Q_3)$
(126) $h_1Q_3(1) \rightarrow e_{45}h_2g_2 = \tilde{\phi}(h_1Q_3(1))$
(127) $h_2Q_3(1) \rightarrow e_{43}h_3g_3 = \tilde{\phi}(h_2Q_3(1))$
(128) $h_0x_0 \rightarrow e_{13}h_1c_0h_4 = \tilde{\phi}(h_0x_0)$
(129) $h_1x_1 \rightarrow e_{27}h_2c_1h_5 = \tilde{\phi}(h_1x_1)$
(130) $h_2x_2 \rightarrow e_{55}h_3c_2h_6 = \tilde{\phi}(h_2x_2)$
(131) $h_0x_1 = h_2h_4D_3 \rightarrow e_3h_4D_3 = \tilde{\phi}(h_2h_4D_3)$
(132) $h_1x_2 = h_3h_5D_3(1) \rightarrow e_7h_5D_3(1) = \tilde{\phi}(h_3h_5D_3(1))$
(133) $h_1V_0 \rightarrow e_{28}h_2D_3(1) = \tilde{\phi}(h_1V_0)$
(134) $h_0V_0 = h_5K_0 \rightarrow e_{15}T_0 = \tilde{\phi}(h_5K_0)$
(135) $h_1H_1 \rightarrow e_{11}D_1 = \tilde{\phi}(h_1H_1)$
(136) $h_2H_1(1) \rightarrow e_{23}D_1(1) = \tilde{\phi}(h_2H_1(1))$

It is not difficult to see that (3.43) implies (3.34).

To prove (3.1) (9), we give in (3.44) below the correspondences (defined in (3.40)) for $h_0^3g_4, h_1h_5V_0 \in \text{Ext}_{\mathcal{A}}^{7,195}$.

- (3.44)** (1) $h_0^3g_4 \rightarrow e_{45}e_0h_6^2 = \tilde{\phi}(h_0^3g_4)$
(2) $h_1h_5V_0 \rightarrow e_{29}h_2V_0 = \tilde{\phi}(h_1h_5V_0)$

A check against the differentials in (3.38) (b) shows that $e_{45}e_0h_6^2, e_{29}h_2V_0$ are not boundaries in the spectral sequence. It follows that $e_{45}e_0h_6^2, e_{29}h_2V_0$ are infinite cycles which represent non-trivial classes in $\text{Ext}_{\mathcal{A}}^{6,194}(P)$, and these classes are linearly independent. This proves (3.1) (9).

This completes the proofs of (3.1) and (3.2).

The remainder of this section is devoted to proving (3.3), (3.4), (3.5), and (3.6) (simply “(3.3) through (3.6)”).

First we prove (3.3). For $0 \leq l \leq m$ we can calculate $\text{Ext}_{\mathcal{A}}^{s,t}(P_l^m)$ using the spectral sequence in (3.32) with the following modifications. The mod 2 reduced homology group $H_*(P_l^m)$ has the same right \mathcal{A} -module structure as that of $H_*(P)$ given in (3.16) except that $e_k = 0$ for $k < l$ and for $k > m$. Therefore we have the following.

(3.45) Let $\overline{F}(0) = 0 \subset \overline{F}(l) \subset \overline{F}(l+1) \cdots \subset \overline{F}(m) = H_*(P_l^m)$ be the increasing filtration of subcomplexes given by $\overline{F}(n) = \Sigma_{l \leq k \leq n} H_k(P_l^m) \otimes \Lambda$, $l \leq n \leq m$, then $\{\overline{F}(i)\}_{l \leq i \leq m}$ defines a spectral sequence $\{\overline{E}_r^{i,s,t}\}_{r \geq 1}$ with $\overline{E}_1^{i,s,t} \cong \Sigma^i \text{Ext}_{\mathcal{A}}^{s,t+s-i}$ and $\bigoplus_{i \geq 1} \overline{E}_\infty^{i,s,t} \cong \text{Ext}_{\mathcal{A}}^{s,t+s}(P_l^m)$, which is identical to the one in (3.32) except that $\overline{E}_r^{i,s,t} = 0$ for $i < l$ and for $i > m$.

Computing the spectral sequence $\{\overline{E}_r^{i,s,t}\}_{r \geq 1}$ is equivalent to computing $\{E_r^{i,s,t}\}_{r \geq 1}$ (put $e_k = 0$ for $k < l$ and for $k > m$ for $\overline{E}_r^{i,s,t}$.) Thus the results (3.3) (i) (8), (9) follow from the spectral sequence calculations in (3.36) and (3.37). It is also easy to see the following.

(3.46) Let $P^m \xrightarrow{i_m} P$ be the inclusion map and $\text{Ext}_{\mathcal{A}}^{s,t}(P^m) \xrightarrow{(i_m)_*} \text{Ext}_{\mathcal{A}}^{s,t}(P)$ be the induced homomorphism. If $\alpha \in \text{Ext}_{\mathcal{A}}^{s,t}(P)$ is a non-zero class represented in the spectral sequence $\{E_r^{i,s,t}\}_{r \geq 0}$ by an infinite cycle $e_k \beta$ with $k \leq m$, then α can be pulled back to a non-zero class $\overline{\alpha} \in \text{Ext}_{\mathcal{A}}^{s,t}(P^m)$ represented in the spectral sequence $\{\overline{E}_r^{i,s,t}\}_{r \geq 0}$ also by the infinite cycle $e_k \beta$.

From (3.21) and (3.46), it is not difficult to see that each class in (3.2), except $\widehat{h}_6 h_0 h_6^2 \in \text{Ext}_{\mathcal{A}}^{3,192}(P)$ and $\widehat{h}_7 D_3 \in \text{Ext}_{\mathcal{A}}^{4,192}(P)$, can be pulled back to $\text{Ext}_{\mathcal{A}}^{*,*}(P^{62})$. The classes $\widehat{h}_6 h_0 h_6^2, \widehat{h}_7 D_3$ can also be pulled back to $\text{Ext}_{\mathcal{A}}^{*,*}(P^{62})$ because of the following cycle representations for these classes.

$$(3.47) \quad (1) \quad e_{62} h_1 h_6^2 \longleftrightarrow \widehat{h}_6 h_0 h_6^2 \\ (2) \quad e_{62} D_3(1) \longleftrightarrow \widehat{h}_7 D_3$$

Thus all the classes in (3.2) for $\text{Ext}_{\mathcal{A}}^{*,*}(P)$ can be pulled back to $\text{Ext}_{\mathcal{A}}^{*,*}(P^{62})$ via $\text{Ext}_{\mathcal{A}}^{s,t}(P^{62}) \xrightarrow{(i_{62})_*} \text{Ext}_{\mathcal{A}}^{s,t}(P)$. And the pullbacks of $\widehat{h}_6 h_0 h_6^2 \in \text{Ext}_{\mathcal{A}}^{3,192}(P)$ and $\widehat{h}_7 D_3 \in \text{Ext}_{\mathcal{A}}^{4,192}(P)$ are denoted by $\overline{e_{62} h_1 h_6^2} \in \text{Ext}_{\mathcal{A}}^{3,192}(P^{62})$ and $\overline{e_{62} D_3(1)} \in \text{Ext}_{\mathcal{A}}^{3,192}(P^{62})$ respectively.

To prove (3.3) (i) (j) for $1 \leq j \leq 9$ (simply (3.3) (i)) and (3.3) (ii), consider the cofibration sequence $\Sigma(P_{m+1} = P/P^m) \rightarrow P^m \xrightarrow{i_m} P$ and the induced long exact sequence of Ext groups.

$$\cdots \rightarrow \text{Ext}_{\mathcal{A}}^{s-1,t}(P_{m+1}) \xrightarrow{\delta} \text{Ext}_{\mathcal{A}}^{s,t}(P^m) \xrightarrow{(i_m)_*} \text{Ext}_{\mathcal{A}}^{s,t}(P) \rightarrow \text{Ext}_{\mathcal{A}}^{s,t}(P_{m+1}) \rightarrow \cdots$$

Since each of the classes in $\text{Ext}_{\mathcal{A}}^{s,t}(P^{62})$ in (3.3) is a pullback from $\text{Ext}_{\mathcal{A}}^{s,t}(P)$ via $(i_m)_*$, we need to show that for $m = 62$ and for each of the (s, t) in (3.3) (i) (1) through (9), the map $\text{Ext}_{\mathcal{A}}^{s-1,t}(P_{63}) \xrightarrow{\delta} \text{Ext}_{\mathcal{A}}^{s,t}(P^{62})$ is zero. This is equivalent to saying that in the spectral sequence $\{E_r^{i,s,t-s}\}_{r \geq 1}$ for $\text{Ext}_{\mathcal{A}}^{s,t}(P)$, each of the differentials $E_r^{i,s-1,t-s} \xrightarrow{d_r} E_r^{i-r,s,t-s}$ that starts with the basis element $e_i \alpha$ with $i > 62$ must either be a zero differential or hit a target $e_{i-r} \beta$ with $i-r > 62$. A check against the list of differentials in [16] shows that this is indeed the case. This proves (3.3) (i) and (3.3) (ii). (3.3) (iii) follows easily from (3.3) (i) and (3.3) (ii). This proves (3.3).

The result (3.4) (1) through (16) also follows from the list of the differentials in [16] once we describe cycle representations for the classes in (3.4) as follows.

- (3.48) (1) $\overline{e_{23}} = \{e_{23}\} \in \text{Ext}_{\mathcal{A}}^{0,23}(P_{17}^{62})$.
(2) $\overline{e_{31}} = \{e_{31}\} \in \text{Ext}_{\mathcal{A}}^{0,31}(P_{17}^{62}) = \text{Ext}_{\mathcal{A}}^{0,31}(P^{32})$.
(3) $\overline{e_{39}} = \{e_{39}\} \in \text{Ext}_{\mathcal{A}}^{0,39}(P_{33}^{62})$.
(4) $\overline{e_{47}} = \{e_{47}\} \in \text{Ext}_{\mathcal{A}}^{0,47}(P_{33}^{62})$.
(5) $\overline{e_{15}} = \{e_{15}\} \in \text{Ext}_{\mathcal{A}}^{0,15}(P^{16}) = \text{Ext}_{\mathcal{A}}^{0,15}(P^{32})$.
(6) $\overline{e_7} = \{e_7\} \in \text{Ext}_{\mathcal{A}}^{0,7}(P^{16})$.
(7) $\overline{e_{62}h_1} = \{e_{62}\lambda_1 + e_{60}\lambda_3 + e_{56}\lambda_7 + e_{48}\lambda_{15} + e_{32}\lambda_{31}\} \in \text{Ext}_{\mathcal{A}}^{1,64}(P_{47}^{62})$.
(8) $\overline{e_{48}h_0h_3^2} = \{e_{48}\lambda_0\lambda_7^2 + e_{47}(\lambda_9\lambda_3^2 + \lambda_3^2\lambda_9)\} \in \text{Ext}_{\mathcal{A}}^{3,65}(P_{47}^{62})$.
(9) $\overline{e_{54}h_1h_3} = \{e_{54}\lambda_1\lambda_7 + e_{48}\lambda_7^2\} \in \text{Ext}_{\mathcal{A}}^{2,64}(P_{47}^{62})$.

To prove

$$(3.4) (17) \quad (q_{47})_*(\widehat{g}_4h_0^2) = \overline{e_{48}h_0h_3^2h_6^2} \neq 0 \text{ in } \text{Ext}_{\mathcal{A}}^{5,193}(P_{47}^{62}),$$

we recall from (3.35) that $\widehat{g}_4h_0^2 \in \text{Ext}_{\mathcal{A}}^{5,193}(P)$ is represented by $e_{47}T_0$ in the spectral sequence for $\text{Ext}_{\mathcal{A}}^{*,*}(P)$. Also recall from (3.13) (19) that

$$T_0 = \{T_0^* = \lambda_0^2(\lambda_{79}\lambda_{31}^2 + \lambda_{31}^2\lambda_{79}) + (\lambda_9\lambda_3^2 + \lambda_3^2\lambda_9)\lambda_{63}^2\} \in \text{Ext}_{\mathcal{A}}^{5,146}.$$

It follows that $(q_{47})_*(\widehat{g}_4h_0^2) = \{e_{47}T_0^*\} \in \text{Ext}_{\mathcal{A}}^{5,193}$. By (3.48) (11), $\overline{e_{48}h_0h_3^2} = \{e_{48}\lambda_0\lambda_7^2 + e_{47}(\lambda_9\lambda_3^2 + \lambda_3^2\lambda_9)\}$. So $e_{48}h_0h_3^2h_6^2 = (q_{47})_*(\widehat{g}_4h_0^2)$ because $\delta(e_{48}\lambda_0(\lambda_{79}\lambda_{31}^2 + \lambda_{31}^2\lambda_{79})) = e_{48}\lambda_0\lambda_7^2\lambda_{63}^2 + e_{47}(\lambda_9\lambda_3^2 + \lambda_3^2\lambda_9)\lambda_{63}^2 + e_{47}T_0^*$ in $H_*(P_{47}^{62}) \otimes \Lambda$.

This completes the proof of (3.4).

Next we prove (3.5). To describe cohomology classes in $\text{Ext}_{\mathcal{A}}^{s,t}(X)$ for $X = \widetilde{P}^{62}$ we need to study $H_*(X = \widetilde{P}^{62})$. Recall from (2.29) that \widetilde{P}^{62} is the mapping cone $C_{\overline{h}_5}$ of a map $S^{30} \cup_{2\iota} e^{31} \xrightarrow{\overline{h}_5} P^{62}$ which is an extension of the map $S^{30} \xrightarrow{\widehat{\theta}_4} P^{62}$, and that $P^{62} \xrightarrow{\widetilde{q}} \widetilde{P}^{62} = P^{62} \cup_{\overline{h}_5} C(S^{30} \cup_{2\iota} e^{31})$ is the inclusion map. Consider the following long exact sequence.

$$\begin{aligned} \cdots \rightarrow H_{31}(S^{30} \cup_{2\iota} e^{31}) \xrightarrow{(\overline{h}_5)_*} H_{31}(P^{62}) \xrightarrow{\widetilde{q}_*} H_{31}(\widetilde{P}^{62}) \\ \xrightarrow{\Delta} H_{30}(S^{30} \cup_{2\iota} e^{31}) \xrightarrow{(\overline{h}_5)_*} H_{30}(P^{62}) \xrightarrow{\widetilde{q}_*} H_{30}(\widetilde{P}^{62}) \rightarrow \cdots \end{aligned}$$

By (2.30), $H_{31}(S^{30} \cup_{2\iota} e^{31}) \xrightarrow{(\overline{h}_5)_*} H_{31}(P^{62})$ is an isomorphism. $H_{30}(S^{30} \cup_{2\iota} e^{31}) \xrightarrow{(\overline{h}_5)_*} H_{30}(P^{62})$ is zero because $S^{30} \xrightarrow{\widehat{\theta}_4} P$ is detected by \widehat{h}_4h_4 in the ASS of $\pi_*^S(P)$. So from the above long exact sequence we see the following.

- (3.49) (1) $\mathbb{Z}/2 \cong H_k(P^{62}) \xrightarrow{\widetilde{q}_*} H_k(\widetilde{P}^{62}) \cong \mathbb{Z}/2$ for $k \neq 31$.
(2) $\mathbb{Z}/2 \cong H_{31}(P^{62}) \xrightarrow{\widetilde{q}_*=0} H_{31}(\widetilde{P}^{62}) \cong \mathbb{Z}/2$.

To describe the right \mathcal{A} -module $H_*(\widetilde{P}^{62})$, denote by \widetilde{e}_{31} the generator of $H_{31}(\widetilde{P}^{62}) \cong \mathbb{Z}/2$, and denote by e_k the generator of $H_k(\widetilde{P}^{62}) \cong \mathbb{Z}/2$ for $1 \leq k \leq 62$ with $k \neq 31$. Thus

$$(3.50) \quad H_k(\widetilde{P}^{62}) = \begin{cases} \mathbb{Z}/2(e_k) & \text{for } 1 \leq k \leq 62 \text{ and } k \neq 31. \\ \mathbb{Z}/2(\widetilde{e}_{31}) & \text{for } k = 31 \\ 0 & \text{otherwise} \end{cases}$$

All the maps in (3.54), except $H_*(P^{62}) \xrightarrow{\tilde{q}_*} H_*(\tilde{P}^{62})$, are algebraic homomorphisms, meaning that they are not induced by topological maps. It follows that (3.54) induces the following long exact sequences of Ext groups (we denote by $\text{Ext}_{\mathcal{A}}^{s,t}(M)$ the Ext group $\text{Ext}_{\mathcal{A}}^{s,t}(M, \mathbb{Z}/2)$):

$$(3.55) \quad \begin{array}{ccccccc} & & & & & \text{Ext}_{\mathcal{A}}^{s-1,t-31} & \\ & & & & & \swarrow \Delta_2 & \\ \dots & \longrightarrow & \text{Ext}_{\mathcal{A}}^{s,t-31} & \xrightarrow{(i_1)_*} & \text{Ext}_{\mathcal{A}}^{s,t}(P^{62}) & \xrightarrow{(q_1)_*} & \text{Ext}_{\mathcal{A}}^{s,t}(M) & \xrightarrow{\Delta_1} & \text{Ext}_{\mathcal{A}}^{s+1,t-31} & \longrightarrow & \dots \\ & & & & \downarrow \tilde{q}_* & \swarrow (i_2)_* & & & & & \\ & & & & \text{Ext}_{\mathcal{A}}^{s,t}(\tilde{P}^{62}) & & & & & & \\ & & & & & \swarrow (q_2)_* & & & & & \\ & & & & & \text{Ext}_{\mathcal{A}}^{s,t-31} & & & & & \end{array}$$

And from this we see:

(3.56) To calculate $\text{Ext}_{\mathcal{A}}^{s,t}(\tilde{P}^{62})$ from the exact sequences in (3.55) we need the following information: (i) $\text{Ext}_{\mathcal{A}}^{s,t-31}$, (ii) $\text{Ext}_{\mathcal{A}}^{s,t}(P^{62})$, (iii) $\text{Ext}_{\mathcal{A}}^{s+1,t-31}$, and (iv) $\text{Ext}_{\mathcal{A}}^{s-1,t-31}$. We also need to know the maps $\text{Ext}_{\mathcal{A}}^{s,t}(M) \xrightarrow{\Delta_1} \text{Ext}_{\mathcal{A}}^{s+1,t-31}$ and $\text{Ext}_{\mathcal{A}}^{s-1,t-31} \xrightarrow{\Delta_2} \text{Ext}_{\mathcal{A}}^{s,t}(M)$.

Now we prove the Ext group results in (3.5). The equation $\overline{e_{39}h_3h_0h_4} = \overline{e_{31}h_4h_0} + \tilde{q}_*(\overline{D_3})$ in (3.5) (3) will be proved later. To see

$$(3.5) \quad (8) \quad \text{Ext}_{\mathcal{A}}^{0,189}(\tilde{P}^{62}) = 0,$$

$$(3.5) \quad (9) \quad \text{Ext}_{\mathcal{A}}^{1,190}(\tilde{P}^{62}) = 0,$$

we note from Theorem 3.14 and Theorem 3.24 that the Ext groups in (3.56) (i) through (iv) are all zero for $(s, t) = (0, 189), (1, 190)$. Therefore $\text{Ext}_{\mathcal{A}}^{0,189}(\tilde{P}^{62}) = 0$ and $\text{Ext}_{\mathcal{A}}^{1,190}(\tilde{P}^{62}) = 0$. This proves (3.5) (8) and (3.5) (9). It is not difficult to see

$$(3.5) \quad (1) \quad \text{Ext}_{\mathcal{A}}^{0,47}(\tilde{P}^{62}) = \mathbb{Z}/2(\overline{e_{47}}),$$

$$(3.5) \quad (4) \quad \text{Ext}_{\mathcal{A}}^{1,63}(\tilde{P}^{62}) = \mathbb{Z}/2(\overline{e_{31}h_5}) \oplus \mathbb{Z}/2(\overline{e_{47}h_4}),$$

from the differential $H_*(\tilde{P}^{62}) \otimes \Lambda \xrightarrow{\delta} H_*(\tilde{P}^{62}) \otimes \Lambda$ as given by (3.51). To see (3.5) (2),(3),(5),(6),(10),(11), we will use the knowledge of (3.56) (i), (iii), and (iv) as given by (3.13), and that of (3.56) (ii) as given by (3.3). In (3.57) below we collect this information in (3.57) (j) (i), (ii), (iii), (iv) for each $j = 2, 3, 5, 6, 10, 11$. These will imply (3.57) (j) (v) and (3.57) (j) (vi), and note that (3.57) (j) (vi) is precisely (3.5) (j), $j = 2, 3, 5, 6, 10, 11$.

$$(3.57) \quad (2) \quad \begin{array}{l} \text{(i)} \quad \text{Ext}_{\mathcal{A}}^{2,17} = \mathbb{Z}/2(h_0h_4), \\ \text{(ii)} \quad \text{Ext}_{\mathcal{A}}^{2,48}(P^{62}) = \mathbb{Z}/2(\widehat{h_4}h_0h_5), \end{array}$$

- (iii) $\text{Ext}_{\mathcal{A}}^{3,17} = \mathbb{Z}/2(h_0h_3^2)$,
 - (iv) $\text{Ext}_{\mathcal{A}}^{1,17} = 0$,
 - (v) $\text{Ext}_{\mathcal{A}}^{2,48}(M) = \mathbb{Z}/2((q_1)_*(\widehat{h}_4h_0h_5))$,
 - (vi) $\text{Ext}_{\mathcal{A}}^{2,48}(\widetilde{P}^{62}) = \mathbb{Z}/2(\overline{e_{39}h_3h_0}) \oplus \mathbb{Z}/2(q_*(\widehat{h}_4h_0h_5))$.
- (3)
- (i) $\text{Ext}_{\mathcal{A}}^{3,33} = \mathbb{Z}/2(h_0h_4^2)$,
 - (ii) $\text{Ext}_{\mathcal{A}}^{3,64}(P^{62}) = \mathbb{Z}/2(\widehat{D}_3)$,
 - (iii) $\text{Ext}_{\mathcal{A}}^{4,33} = 0$,
 - (iv) $\text{Ext}_{\mathcal{A}}^{2,33} = \mathbb{Z}/2(h_0h_5)$,
 - (v) $\text{Ext}_{\mathcal{A}}^{3,64}(M) = \mathbb{Z}/2((q_1)_*(\widehat{D}_3))$,
 - (vi) $\text{Ext}_{\mathcal{A}}^{3,64}(\widetilde{P}^{62}) = \mathbb{Z}/2(\overline{\widetilde{e}_{31}h_4^2h_0}) \oplus \mathbb{Z}/2(\widetilde{q}_*(\widehat{D}_3))$.
- (5)
- (i) $\text{Ext}_{\mathcal{A}}^{2,33} = \mathbb{Z}/2(h_0h_5)$,
 - (ii) $\text{Ext}_{\mathcal{A}}^{2,64}(P^{62}) = \mathbb{Z}/2(\widehat{h}_5h_0h_5)$,
 - (iii) $\text{Ext}_{\mathcal{A}}^{3,33} = \mathbb{Z}/2(h_0h_4^2)$,
 - (iv) $\text{Ext}_{\mathcal{A}}^{1,33} = 0$,
 - (v) $\text{Ext}_{\mathcal{A}}^{2,64}(M) = \mathbb{Z}/2(\overline{e_{47}h_0h_4})$,
 - (vi) $\text{Ext}_{\mathcal{A}}^{2,64}(\widetilde{P}^{62}) = \mathbb{Z}/2(\overline{\widetilde{e}_{31}h_5h_0}) \oplus \mathbb{Z}/2(\overline{e_{47}h_0h_4})$.
- (6)
- (i) $\text{Ext}_{\mathcal{A}}^{4,161} = \mathbb{Z}/2(h_0h_4^2h_7)$,
 - (ii) $\text{Ext}_{\mathcal{A}}^{4,192}(P^{62}) = \mathbb{Z}/2(\widehat{h}_4h_0c_4) \oplus \mathbb{Z}/2(\widehat{D}_3h_7) \oplus \mathbb{Z}/2(\widehat{g}_4h_0) \oplus \mathbb{Z}/2(\overline{e_{62}D_3(1)})$,
 - (iii) $\text{Ext}_{\mathcal{A}}^{5,161} = \mathbb{Z}/2(V_0)$,
 - (iv) $\text{Ext}_{\mathcal{A}}^{3,161} = h_0h_5h_7$,
 - (v) $\text{Ext}_{\mathcal{A}}^{4,192}(M) = \mathbb{Z}/2((q_1)_*(\widehat{h}_4h_0c_4)) \oplus \mathbb{Z}/2((q_1)_*(\widehat{D}_3h_7)) \oplus \mathbb{Z}/2((q_1)_*(\widehat{g}_4h_0)) \oplus \mathbb{Z}/2((q_1)_*(\overline{e_{62}D_3(1)}))$,
 - (vi) $\text{Ext}_{\mathcal{A}}^{4,192}(\widetilde{P}^{62}) = \mathbb{Z}/2(\widetilde{q}_*(\widehat{h}_4h_0c_4)) \oplus \mathbb{Z}/2(\widetilde{q}_*(\widehat{D}_3h_7)) \oplus \mathbb{Z}/2(\widetilde{q}_*(\widehat{g}_4h_0)) \oplus \mathbb{Z}/2(\widetilde{q}_*(\overline{e_{62}D_3(1)})) \oplus \mathbb{Z}/2(\overline{\widetilde{e}_{31}h_4^2h_0h_7})$.
- (10)
- (i) $\text{Ext}_{\mathcal{A}}^{2,160} = \mathbb{Z}/2(h_5h_7)$,
 - (ii) $\text{Ext}_{\mathcal{A}}^{2,191}(P^{62}) = \mathbb{Z}/2(\widehat{h}_5h_5h_7)$,
 - (iii) $\text{Ext}_{\mathcal{A}}^{3,160} = \mathbb{Z}/2(h_4^2h_7)$,
 - (iv) $\text{Ext}_{\mathcal{A}}^{1,160} = 0$,
 - (v) $\text{Ext}_{\mathcal{A}}^{2,191}(M) = \mathbb{Z}/2(\overline{e_{47}h_4h_7})$,
 - (vi) $\text{Ext}_{\mathcal{A}}^{2,191}(\widetilde{P}^{62}) = \mathbb{Z}/2(\overline{\widetilde{e}_{31}h_5h_7}) \oplus \mathbb{Z}/2(\overline{e_{47}h_4h_7})$.
- (11)
- (i) $\text{Ext}_{\mathcal{A}}^{3,161} = \mathbb{Z}/2(h_0h_5h_7)$,
 - (ii) $\text{Ext}_{\mathcal{A}}^{3,192}(P^{62}) = \mathbb{Z}/2(\widehat{h}_5h_0h_5h_7) \oplus \mathbb{Z}/2(\overline{e_{62}h_1h_6^2})$,
 - (iii) $\text{Ext}_{\mathcal{A}}^{4,161} = \mathbb{Z}/2(h_0h_4^2h_7)$,
 - (iv) $\text{Ext}_{\mathcal{A}}^{2,161} = 0$,
 - (v) $\text{Ext}_{\mathcal{A}}^{3,192}(M) = \mathbb{Z}/2((q_1)_*(\overline{e_{62}h_1h_6^2})) \oplus \mathbb{Z}/2(\overline{e_{47}h_4h_0h_7})$,
 - (vi) $\text{Ext}_{\mathcal{A}}^{3,192}(\widetilde{P}^{62}) = \mathbb{Z}/2(\overline{\widetilde{e}_{31}h_5h_0h_7}) \oplus \mathbb{Z}/2(\overline{e_{47}h_4h_0h_7}) \oplus \mathbb{Z}/2(\widetilde{q}_*(\overline{e_{62}h_1h_6^2}))$.

We explain how we obtain (3.57) (2) (v) and (3.57) (2) (vi) from (3.57) (2) (i), (ii), (iii), (iv) as an illustration of such deductions. From the definition of $\mathbb{Z}/2(e_{31}) \xrightarrow{i_1} H_*(P^{62})$ we have $(i_1)_*(h_0h_4) = \widehat{h}_5h_0h_4 = 0$ in $\text{Ext}_{\mathcal{A}}^{2,48}(P^{62})$ and

$(i_1)_*(h_0h_3^2) = \widehat{h}_5h_0h_3^2 = \widehat{h}_4h_4^2h_0 \neq 0$ in $\text{Ext}_{\mathcal{A}}^{3,48}(P^{62})$. From this we deduce (3.57) (2) (v). Since $\Delta_2(h_0h_4) = \widehat{h}_4h_4^2h_0 = \widehat{h}_5h_3^2h_0 = 0$ in $\text{Ext}_{\mathcal{A}}^{3,48}(M)$, $h_0h_4 \in \text{Ext}_{\mathcal{A}}^{2,17}$ is pulled back to a class in $\text{Ext}_{\mathcal{A}}^{2,48}(\widetilde{P}^{62})$ via $(q_2)_*$. From (3.52) we see this class is $\overline{e_{39}h_3h_0}$. Thus $\text{Ext}_{\mathcal{A}}^{2,48}(\widetilde{P}^{62}) = \mathbb{Z}/2(\overline{e_{39}h_3h_0}) \oplus \mathbb{Z}/2(q_*(\widehat{h}_4h_0h_5))$ as claimed in (3.57) (2) (vi). (3.57) (j) (vi) for $j = 3, 5, 6, 10, 11$ are similarly deduced. This proves (3.5) (2), (3), (5), (6), (10), (11). To see

$$(3.5) (7) (a) \quad \widetilde{q}_*(\widehat{h}_5h_5D_3(1)) = 0 \text{ in } \text{Ext}_{\mathcal{A}}^{5,193}(\widetilde{P}^{62}),$$

$$(3.5) (7) (b) \quad \widetilde{q}_*(\widehat{h}_4h_0^2c_4), \widetilde{q}_*(\widehat{h}_4h_1f_3), \widetilde{q}_*(\gamma_{61}h_7), \widetilde{q}_*(\widehat{g}_4h_0^2), \overline{\widetilde{e}_{31}h_4^2h_0^2h_7} \text{ are linearly independent in } \text{Ext}_{\mathcal{A}}^{5,193}(\widetilde{P}^{62}),$$

we also use the diagram (3.55) with following the knowledge of (3.56) (i) and (iv) for $(s, t) = (5, 193)$ as given by (3.13), and that of (3.56) (iv) for $(s, t) = (5, 193)$ as given by (3.3) (i) (9).

$$(3.58) \quad (i) \quad \text{Ext}_{\mathcal{A}}^{5,162} = \mathbb{Z}/2(h_0^2h_4^2h_7) \oplus \mathbb{Z}/2(h_5D_3(1)).$$

$$(ii) \quad \text{Ext}_{\mathcal{A}}^{5,193}(P^{62}) = \mathbb{Z}/2(\widehat{h}_5h_5D_3(1)) \oplus \mathbb{Z}/2(\widehat{h}_4h_0^2c_4) \oplus \mathbb{Z}/2(\widehat{h}_4h_1f_3) \oplus \mathbb{Z}/2(\gamma_{61}h_7) \oplus \mathbb{Z}/2(\widehat{g}_4h_0^2).$$

$$(iv) \quad \text{Ext}_{\mathcal{A}}^{4,162} = \mathbb{Z}/2(h_1h_4^2h_7) \oplus \mathbb{Z}/2(h_0^2h_5h_7).$$

The map $\text{Ext}_{\mathcal{A}}^{5,162} \xrightarrow{(i_1)_*} \text{Ext}_{\mathcal{A}}^{5,193}(P^{62})$ is given by $(i_1)_*(h_5D_3(1)) = \widehat{h}_5h_5D_3(1) \in \text{Ext}_{\mathcal{A}}^{5,193}(P^{62})$, and $(i_1)_*(h_0^2h_4^2h_7) = \widehat{h}_5h_0^2h_4^2h_7 = 0$ in $\text{Ext}_{\mathcal{A}}^{5,193}(P^{62})$. So we see $(q_1)_*(\widehat{h}_5h_5D_3(1)) = 0$ and therefore $\widetilde{q}_*(\widehat{h}_5h_5D_3(1)) = (i_2)_*(q_1)_*(\widehat{h}_5h_5D_3(1)) = 0$. This proves (3.5) (7) (a). We also see that $(q_1)_*(\widehat{h}_4h_0^2c_4), (q_1)_*(\widehat{h}_4h_1f_3), (q_1)_*(\gamma_{61}h_7), (q_1)_*(\widehat{g}_4h_0^2)$ are linearly independent in $\text{Ext}_{\mathcal{A}}^{5,193}(M)$. The map $\text{Ext}_{\mathcal{A}}^{4,162} \xrightarrow{\Delta_2} \text{Ext}_{\mathcal{A}}^{5,193}(M)$ is given by $\Delta_2(h_1h_4^2h_7) = \widehat{h}_4h_4h_1h_4^2h_7 = \widehat{h}_4h_1h_3^2h_5h_7 = 0$ and $\Delta_2(h_0^2h_5h_7) = \widehat{h}_4h_4h_0^2h_5h_7 = 0$ in $\text{Ext}_{\mathcal{A}}^{5,193}(M)$, so the map $\text{Ext}_{\mathcal{A}}^{4,162} \xrightarrow{(i_2)_*} \text{Ext}_{\mathcal{A}}^{5,193}(\widetilde{P}^{62})$ is a monomorphism. From (3.58) (i) we also have $h_0^2h_4^2h_7 \in \text{Ext}_{\mathcal{A}}^{5,162}$ with $\Delta_2(h_0^2h_4^2h_7) = \widehat{h}_4h_4h_0^2h_4^2h_7 = \widehat{h}_4h_0^2h_3^2h_5h_7 = 0$ in $\text{Ext}_{\mathcal{A}}^{6,193}(M)$, so $h_0^2h_4^2h_7 \in \text{Ext}_{\mathcal{A}}^{5,162}$ has a pullback $\overline{\widetilde{e}_{31}h_4^2h_0^2h_7} \in \text{Ext}_{\mathcal{A}}^{5,193}(\widetilde{P}^{62})$. From these it is not difficult to see that $\widetilde{q}_*(\widehat{h}_4h_0^2c_4), \widetilde{q}_*(\widehat{h}_4h_1f_3), \widetilde{q}_*(\gamma_{61}h_7), \widetilde{q}_*(\widehat{g}_4h_0^2), \overline{\widetilde{e}_{31}h_4^2h_0^2h_7}$ are linearly independent in $\text{Ext}_{\mathcal{A}}^{5,193}(\widetilde{P}^{62})$. This proves (3.5) (7) (b).

The equation

$$(3.5) (3) \quad \overline{e_{39}h_3h_0h_4} = \overline{\widetilde{e}_{31}h_4^2h_0} + \widetilde{q}_*(\widehat{D}_3) \text{ in } \text{Ext}_{\mathcal{A}}^{3,64}(\widetilde{P}^{62}).$$

is proved by the following (3.59).

$$(3.59) \quad (1) \quad \text{From (3.21) (3), } \widehat{D}_3 = \{e_{22}\lambda_1\lambda_7\lambda_{31} + e_{16}\lambda_{15}^3 + e_{14}\lambda_9\lambda_7\lambda_{31}\} \in \text{Ext}_{\mathcal{A}}^{3,64}(P), \text{ and } \{e_{22}\lambda_1\lambda_7\lambda_{31} + e_{16}\lambda_{15}^3 + e_{14}\lambda_9\lambda_7\lambda_{31}\} = \{e_{23}\lambda_7\lambda_{31}\lambda_0 + e_{15}(\lambda_{15}^2\lambda_{16} + \lambda_{15}\lambda_{16}\lambda_{15} + \lambda_{16}\lambda_{15}^2) + e_{16}\lambda_{15}^3\}. \text{ So } \widetilde{q}_*(\widehat{D}_3) = \{e_{23}\lambda_7\lambda_{31}\lambda_0 + e_{15}(\lambda_{15}^2\lambda_{16} + \lambda_{15}\lambda_{16}\lambda_{15} + \lambda_{16}\lambda_{15}^2) + e_{16}\lambda_{15}^3\}.$$

$$(2) \quad \text{It is not difficult to see that } \overline{\widetilde{e}_{31}h_4^2h_0} = \{\widetilde{e}_{31}\lambda_{15}^2\lambda_0 + e_{15}(\lambda_{15}^2\lambda_{16} + \lambda_{15}\lambda_{16}\lambda_{15} + \lambda_{16}\lambda_{15}^2) + e_{16}\lambda_{15}^3\}.$$

$$(3) \quad \text{It is not difficult to see that } \overline{e_{39}h_3} = \{e_{39}\lambda_7 + e_{23}\lambda_{23} + \widetilde{e}_{31}\lambda_{15}\}. \text{ So } \overline{e_{39}h_3h_0h_4} = \overline{e_{39}h_3h_4h_0} = \{(e_{39}\lambda_7 + e_{23}\lambda_{23} + \widetilde{e}_{31}\lambda_{15})\lambda_{15}\lambda_0\} = \{e_{23}\lambda_7\lambda_{31}\lambda_0 + \widetilde{e}_{31}\lambda_{15}^2\lambda_0\}.$$

(4) Then from (1), (2), and (3) we have $\widetilde{q}_*(\widehat{D}_3) + \overline{\widetilde{e}_{31}h_4^2h_0} = \{e_{23}\lambda_7\lambda_{31}\lambda_0 + \widetilde{e}_{31}\lambda_{15}^2\lambda_0\} = e_{39}h_3h_0h_4$.

This completes the proof of (3.5).

We proceed to prove (3.6) (1), (2), (3). In (3.60) below we define some cohomology classes in $\text{Ext}_{\mathcal{A}}^{*,*}(\widetilde{P}_{17}^{62})$.

- (3.60)** (1) $\overline{e_{31}} = \{\widetilde{e}_{31}\} \in \text{Ext}_{\mathcal{A}}^{0,31}(\widetilde{P}_{17}^{62})$.
(2) $\overline{e_{47}} = \{e_{47}\} \in \text{Ext}_{\mathcal{A}}^{0,47}(\widetilde{P}_{17}^{62})$.

$P_{17}^{62} = P^{62}/P^{16} \xrightarrow{\widetilde{q}_{17}} \widetilde{P}_{17}^{62} = \widetilde{P}^{62}/P^{16}$ is induced by $P^{62} \xrightarrow{\widetilde{q}} \widetilde{P}^{62}$, so we can modify diagram (3.55) for P^{62} to P_{17}^{62} as follows,

$$(3.61) \quad \begin{array}{ccccccc} & & & & & \text{Ext}_{\mathcal{A}}^{s-1,t-31} & \\ & & & & & \swarrow \Delta'_2 & \\ & & & & & \text{Ext}_{\mathcal{A}}^{s+1,t-31} & \\ & & & & & \swarrow \Delta'_1 & \\ \dots & \longrightarrow & \text{Ext}_{\mathcal{A}}^{s,t-31} & \xrightarrow{(i_1)_*} & \text{Ext}_{\mathcal{A}}^{s,t}(P_{17}^{62}) & \xrightarrow{(q_1)_*} & \text{Ext}_{\mathcal{A}}^{s,t}(M_{17}) & \xrightarrow{\Delta'_1} & \text{Ext}_{\mathcal{A}}^{s+1,t-31} & \longrightarrow & \dots \\ & & & & \downarrow (\widetilde{q}_{17})_* & \swarrow (i_2)_* & & & & & \\ & & & & \text{Ext}_{\mathcal{A}}^{s,t}(\widetilde{P}_{17}^{62}) & & & & & & \\ & & & & \swarrow (q_2)_* & & & & & & \\ & & & & \text{Ext}_{\mathcal{A}}^{s,t-31} & & & & & & \end{array}$$

where $M_{17} = \frac{H_*(P_{17}^{62})}{\mathbb{Z}/2\langle e_{31} \rangle}$. To calculate (3.6) (1), (2), (3) we use the same method as in the proof of (3.5). The information in (3.62) (j) (i), (ii), (iii), (iv) for $1 \leq j \leq 3$ below are collected from (3.14) and (3.4) (1), (2), (3). From these we derive (3.62) (j) (v) and (3.62) (j) (vi), and note that (3.62) (j) (vi) are precisely (3.6) (j) for $1 \leq j \leq 3$.

- (3.62)** (1) (i) $\text{Ext}_{\mathcal{A}}^{1,32} = \mathbb{Z}/2\langle h_5 \rangle$,
(ii) $\text{Ext}_{\mathcal{A}}^{1,63}(P_{17}^{62}) = \mathbb{Z}/2\langle \overline{e_{31}}h_5 \rangle$,
(iii) $\text{Ext}_{\mathcal{A}}^{2,32} = \mathbb{Z}/2\langle h_4^2 \rangle$,
(iv) $\text{Ext}_{\mathcal{A}}^{0,32} = 0$,
(v) $\text{Ext}_{\mathcal{A}}^{1,63}(M_{17}) = \mathbb{Z}/2\langle \overline{e_{47}}h_4 \rangle$,
(vi) $\text{Ext}_{\mathcal{A}}^{1,63}(\widetilde{P}_{17}^{62}) = \mathbb{Z}/2\langle \overline{e_{31}}h_5 \rangle \oplus \mathbb{Z}/2\langle \overline{e_{47}}h_4 \rangle$.
(2) (i) $\text{Ext}_{\mathcal{A}}^{2,1} = 0$,
(ii) $\text{Ext}_{\mathcal{A}}^{2,32}(P_{17}^{62}) = \mathbb{Z}/2\langle \overline{e_{23}}h_0h_3 \rangle$,
(iii) $\text{Ext}_{\mathcal{A}}^{3,1} = 0$,
(iv) $\text{Ext}_{\mathcal{A}}^{1,1} = \mathbb{Z}/2\langle h_0 \rangle$,
(v) $\text{Ext}_{\mathcal{A}}^{2,32}(M_{17}) = \mathbb{Z}/2\langle (q_1)_*(\overline{e_{23}}h_0h_3) \rangle$,
(vi) $\text{Ext}_{\mathcal{A}}^{2,32}(\widetilde{P}_{17}^{62}) = \mathbb{Z}/2\langle (\widetilde{q}_{17})_*(\overline{e_{23}}h_0h_3) \rangle$.
(3) (i) $\text{Ext}_{\mathcal{A}}^{3,33} = \mathbb{Z}/2\langle h_0h_4^2 \rangle$,

- (ii) $\text{Ext}_{\mathcal{A}}^{3,64}(P_{17}^{62}) = \mathbb{Z}/2(\overline{e_{23}h_0h_3h_5})$,
- (iii) $\text{Ext}_{\mathcal{A}}^{4,33} = 0$,
- (iv) $\text{Ext}_{\mathcal{A}}^{2,33} = \mathbb{Z}/2(h_0h_5)$,
- (v) $\text{Ext}_{\mathcal{A}}^{3,64}(M_{17}) = \mathbb{Z}/2((q_1)_*(\overline{e_{23}h_0h_3h_5}))$,
- (vi) $\text{Ext}_{\mathcal{A}}^{3,64}(\tilde{P}_{17}^{62}) = \mathbb{Z}/2(\overline{\tilde{e}_{31}h_0h_4^2}) \oplus \mathbb{Z}/2((\tilde{q}_{17})_*(\overline{e_{23}h_0h_3h_5}))$.

This proves the Ext group results in (3.6) (1), (2), (3). It is not difficult to see that the map $\text{Ext}_{\mathcal{A}}^{1,63}(\tilde{P}^{62}) \xrightarrow{(q'_{17})_*} \text{Ext}_{\mathcal{A}}^{1,63}(\tilde{P}_{17}^{62})$ is as given in (3.6) (1), and the map $\text{Ext}_{\mathcal{A}}^{3,64}(\tilde{P}^{62}) \xrightarrow{(q'_{17})_*} \text{Ext}_{\mathcal{A}}^{3,64}(\tilde{P}_{17}^{62})$ is as given in (3.6) (3). Note that $\overline{\tilde{e}_{31}h_4} = \{\tilde{e}_{31}\lambda_{15}\} \in \text{Ext}_{\mathcal{A}}^{1,63}(\tilde{P}_{17}^{62})$ is a decomposable element because there is a class $\overline{\tilde{e}_{31}} = \{\tilde{e}_{31}\} \in \text{Ext}_{\mathcal{A}}^{0,31}(\tilde{P}_{17}^{62})$ (but \tilde{e}_{31} is not a cycle in $H_*(\tilde{P}^{62}) \otimes \Lambda$). This proves (3.6) (1), (2), (3). Finally, it is not difficult to see

$$(3.6) (4) \quad \text{Ext}_{\mathcal{A}}^{2,48}(\tilde{P}^{62}) \xrightarrow{(q'_{33})_*} \text{Ext}_{\mathcal{A}}^{2,48}(\tilde{P}_{33}^{62} = P_{33}^{62}) \text{ is given by } (q'_{33})_*(\overline{e_{39}h_3h_0}) = \overline{e_{39}h_3h_0}, (q'_{33})_*(\tilde{q}_*(\tilde{h}_4h_0h_5)) = 0,$$

$$(3.6) (5) \quad \text{Ext}_{\mathcal{A}}^{0,47}(\tilde{P}^{62}) \xrightarrow{(q'_{33})_*} \text{Ext}_{\mathcal{A}}^{0,47}(\tilde{P}_{33}^{62} = P_{33}^{62}) \text{ is given by } (q'_{33})_*(\overline{e_{47}}) = \overline{e_{47}},$$

$$(3.6) (6) \quad \text{Ext}_{\mathcal{A}}^{2,64}(\tilde{P}^{62}) \xrightarrow{(q'_{47})_*} \text{Ext}_{\mathcal{A}}^{2,64}(\tilde{P}_{47}^{62} = P_{47}^{62}) \text{ is given by } (q'_{47})_*(\overline{\tilde{e}_{31}h_5h_5}) = (q'_{47})_*(\overline{e_{47}h_0h_4}) = 0,$$

from the cycle representations of these classes.

This completes the proof of (3.6).

4 Proofs of Theorem 2.39 and Theorem 2.40

In this section we prove Theorem 2.39 and Theorem 2.40 which are about the stable homotopy groups $\pi_{62}^S(P_{46}^{62})$ and $\pi_{62}^S(P_{47}^{62})$. The precise statements of these theorems will be recalled later separately when we come to prove them. Our tool for proving Theorem 2.39 and Theorem 2.40 is the Adams spectral sequence for $\pi_*^S(P_l^m)$ with $1 \leq l < m$. In the first half of this section we will compute

(4.1) The Ext groups $\text{Ext}_{\mathcal{A}}^{s,t}(P_{46}^{62})$ and $\text{Ext}_{\mathcal{A}}^{s,t}(P_{47}^{62})$ for $s \geq 0$, $61 \leq t - s \leq 63$.

In the second half we will then prove Theorem 2.39 and Theorem 2.40.

To compute (4.1) we will use a long exact sequence in Ext groups as in (4.2) below. Let $Y \xrightarrow{f} X \xrightarrow{g} Z$ be a cofibration sequence, and suppose $0 \rightarrow H_*(Y) \xrightarrow{f_*} H_*(X) \xrightarrow{g_*} H_*(Z) \rightarrow 0$ is an exact sequence in mod 2 homology. Then we have a long exact sequence of Ext groups for all s, t .

$$(4.2) \quad \begin{array}{c} \cdots \rightarrow \text{Ext}_{\mathcal{A}}^{s-1,t}(Z) \xrightarrow{\delta} \text{Ext}_{\mathcal{A}}^{s,t}(Y) \xrightarrow{f_*} \text{Ext}_{\mathcal{A}}^{s,t}(X) \\ \xrightarrow{g_*} \text{Ext}_{\mathcal{A}}^{s,t}(Z) \xrightarrow{\delta} \text{Ext}_{\mathcal{A}}^{s+1,t}(Y) \rightarrow \cdots \end{array}$$

Then to compute $\text{Ext}_{\mathcal{A}}^{s,t}(X)$ for certain s, t it suffices to know $\ker(\text{Ext}_{\mathcal{A}}^{s,t}(Z) \xrightarrow{\delta} \text{Ext}_{\mathcal{A}}^{s+1,t}(Y))$ and $\text{coker}(\text{Ext}_{\mathcal{A}}^{s-1,t}(Z) \xrightarrow{\delta} \text{Ext}_{\mathcal{A}}^{s,t}(Y))$.

The computations of (4.1) will be accomplished in the following steps.

- (4.3) (1) Compute $\text{Ext}_{\mathcal{A}}^{s,t}(M_2 = S^0 \cup_{2^l} e^1)$ for $s \geq 0$, $0 \leq t - s \leq 2$ and $13 \leq t - s \leq 17$ with the cofibration sequence $S^0 \rightarrow M_2 = S^0 \cup_{2^l} e^1 \rightarrow S^1$.
- (2) Compute $\text{Ext}_{\mathcal{A}}^{s,t}(P_{15})$ for $s \geq 0$, $15 \leq t - s \leq 17$ with $\text{Ext}_{\mathcal{A}}^{s,t}(M_2 = S^0 \cup_{2^l} e^1)$ for $s \geq 0$, $0 \leq t - s \leq 2$ in (1).
- (3) Compute $\text{Ext}_{\mathcal{A}}^{s,t}(P^{14})$ for $s \geq 0$, $13 \leq t - s \leq 16$ with the cofibration sequence $P^{14} \rightarrow P \rightarrow P_{15}$.
- (4) Compute $\text{Ext}_{\mathcal{A}}^{s,t}(P_{47}^{62})$ for $s \geq 0$, $61 \leq t - s \leq 63$ with the cofibration sequence $P_{47}^{48} \simeq \Sigma^{47} M_2 \rightarrow P_{47}^{62} \rightarrow P_{49}^{62}$.
- (5) Compute $\text{Ext}_{\mathcal{A}}^{s,t}(P_{46}^{48})$ for $s \geq 0$, $60 \leq t - s \leq 63$ with the cofibration sequence $S^{46} \rightarrow P_{46}^{48} \rightarrow P_{47}^{48} \simeq \Sigma^{47} M_2$.
- (6) Compute $\text{Ext}_{\mathcal{A}}^{s,t}(P_{46}^{62})$ for $s \geq 0$, $61 \leq t - s \leq 63$ with the cofibration sequence $P_{46}^{48} \rightarrow P_{46}^{62} \rightarrow P_{49}^{62}$.

We begin with (4.3) (1). Since $\text{Ext}_{\mathcal{A}}^{s,t}(S^l) \cong \text{Ext}_{\mathcal{A}}^{s,t-l}$ for all s, t, l , from (4.2) we have the following long exact sequence.

$$(4.4) \quad \begin{array}{c} \cdots \rightarrow \text{Ext}_{\mathcal{A}}^{s-1,t-1} \xrightarrow{(2^l)_*} \text{Ext}_{\mathcal{A}}^{s,t} \rightarrow \text{Ext}_{\mathcal{A}}^{s,t}(M_2 = S^0 \cup_{2^l} e^1) \\ \rightarrow \text{Ext}_{\mathcal{A}}^{s,t-1} \xrightarrow{(2^l)_*} \text{Ext}_{\mathcal{A}}^{s+1,t} \rightarrow \cdots \end{array}$$

Here the group homomorphisms $\text{Ext}_{\mathcal{A}}^{s-1,t-1} \xrightarrow{(2^l)_*} \text{Ext}_{\mathcal{A}}^{s,t}$ and $\text{Ext}_{\mathcal{A}}^{s,t-1} \xrightarrow{(2^l)_*} \text{Ext}_{\mathcal{A}}^{s+1,t}$ are described as follows.




Figure 1: $\text{Ext}_{\mathcal{A}}^{s,t}$, $s \geq 0$, $13 \leq t - s \leq 17$

(4.5) Since $h_0 \in \text{Ext}_{\mathcal{A}}^{1,1}$ corresponds to $Sq^1 \in \mathcal{A}$ and Sq^1 detects the stable map $S^0 \xrightarrow{2\iota} S^0$, it follows that the induced homomorphism $\text{Ext}_{\mathcal{A}}^{s,t} \xrightarrow{(2\iota)_*} \text{Ext}_{\mathcal{A}}^{s+1,t+1}$ is given by

$$(2\iota)_*(x) = h_0 x, \quad \text{for all } x \in \text{Ext}_{\mathcal{A}}^{s,t} \text{ and for all } s, t.$$

We recall from [24] the Ext groups $\text{Ext}_{\mathcal{A}}^{s,t}$ for $s \geq 0$, $0 \leq t - s \leq 17$ as in Figure 1. Here, and in all figures of this section, the horizontal axis will be the $(t - s)$ -axis and the vertical axis will be the s -axis. From (4.4), (4.5) and Figure 1 it is not difficult to deduce the structure of $\text{Ext}_{\mathcal{A}}^{s,t}(M_2)$ for $s \geq 0$, $0 \leq t - s \leq 17$. The results are given in Figure 2. We need to explain the notation for the classes in Figure 2, and also the h_i -extensions, $i = 0, 1$ among some of these classes. We recall from (3.45) that there is a spectral sequence $\{\overline{E}_r^{i,s,t}\}_{r \geq 1}$ with $\overline{E}_1^{i,s,t} \cong \Sigma^i \text{Ext}_{\mathcal{A}}^{s,t+s-i}$ for all i, s, t and $\bigoplus_{i \geq 1} \overline{E}_\infty^{i,s,t} \cong \text{Ext}_{\mathcal{A}}^{s,t+s}(P_l^m)$ for all s, t . A cohomology class represented by $e_i \alpha \in \overline{E}_1^{i,s,t}$ for $l \leq i \leq m$ and $\alpha \in \text{Ext}_{\mathcal{A}}^{s,t+s-i}$ in this spectral sequence will be denoted by $\overline{e}_i \alpha \in \text{Ext}_{\mathcal{A}}^{s,t+s}(P_l^m)$. In particular, since $M_2 = S^0 \cup_{2\iota} e^1 \simeq \Sigma^{-l}(P_l^{l+1} = S^l \cup_{2\iota} e^{l+1})$ (or equivalently, $P_l^{l+1} = \Sigma^l M_2$) for $l = 2k + 1 > 0$, in Figure 2 we display $\text{Ext}_{\mathcal{A}}^{s,t}(P_l^{l+1})$ for $s \geq 0$, $l \leq t - s \leq l + 5$ and $l + 13 \leq t - s \leq l + 17$ for all $l = 2k + 1$, from which one can read $\text{Ext}_{\mathcal{A}}^{s,t}(M_2)$ via $\text{Ext}_{\mathcal{A}}^{s,t}(M_2) \cong \text{Ext}_{\mathcal{A}}^{s,t+l}(P_l^{l+1})$. And the cohomology classes in Figure 2 are denoted by the same notations of the classes in $\text{Ext}_{\mathcal{A}}^{s,t}(P_l^m)$ just mentioned. The cases that we will use are $\text{Ext}_{\mathcal{A}}^{s,t}(P_{15}^{16})$ (for $l = 15$) and $\text{Ext}_{\mathcal{A}}^{s,t}(P_{47}^{48})$ (for $l = 47$).

We also note some of the group extensions in $\text{Ext}_{\mathcal{A}}^{*,*}(P_l^{l+1})$. In Figure 2 there are the following relations, which are denoted by dashed lines.




Figure 2: $\text{Ext}_{\mathcal{A}}^{s,t}(P_l^{l+1})$, $l = 2k+1$, $s \geq 0$, $l \leq t-s \leq l+5$ and $l+13 \leq t-s \leq l+17$

- (4.6) (1) $\overline{e_{l+1} h_1 h_0} = \overline{e_l h_1^2}$ in $\text{Ext}_{\mathcal{A}}^{2,l+4}(P_l^{l+1})$.
(2) $\overline{e_{l+1} h_0^2 d_0 h_1} = \overline{e_l P^1 c_0}$ in $\text{Ext}_{\mathcal{A}}^{7,l+23}(P_l^{l+1})$.
(3) $\overline{e_{l+1} h_0^7 h_4 h_1} = \overline{e_l P^2 h_1}$ in $\text{Ext}_{\mathcal{A}}^{9,l+26}(P_l^{l+1})$.
(4) $\overline{e_{l+1} P^1 c_0 h_0} = \overline{e_l h_1 P^1 c_0}$ in $\text{Ext}_{\mathcal{A}}^{8,l+25}(P_l^{l+1})$.

Note that the relations $\overline{e_{l+1} h_1 h_0 h_4} = \overline{e_l h_1^2 h_4}$ in $\text{Ext}_{\mathcal{A}}^{3,l+20}(P_l^{l+1})$, and $\overline{e_{l+1} h_1 h_0 d_0} = \overline{e_l h_1^2 d_0}$ in $\text{Ext}_{\mathcal{A}}^{6,l+22}(P_l^{l+1})$ in Figure 2 also follow from (4.6) (1). We will only prove the relation (4.6) (1) below because the other two relations in (4.6) will not be used in this paper. For $l = 2k + 1 > 0$ it is not difficult to see the chain representations of the following classes in $\text{Ext}_{\mathcal{A}}^{*,*}(P_l^{l+1})$.

- (4.7) (1) $\overline{e_l} = \{e_l\} \in \text{Ext}_{\mathcal{A}}^{0,l}(P_l^{l+1})$.
(2) $\overline{e_{l+1} h_1} = \{e_{l+1} \lambda_1\} \in \text{Ext}_{\mathcal{A}}^{1,l+3}(P_l^{l+1})$.

The relation (4.6) (1) follows from (4.7) (1), (2), and from $\delta(e_{l+1} \lambda_2) = e_{l+1} \lambda_1 \lambda_0 + e_l \lambda_1^2$ in $H_*(P_l^{l+1}) \otimes \Lambda$.

This completes the computations for (4.3) (1).

To compute $\text{Ext}_{\mathcal{A}}^{s,t}(P_{15})$ for $s \geq 0$, $15 \leq t-s \leq 17$ in (4.3) (2), note that by dimensional reasons, $\text{Ext}_{\mathcal{A}}^{s,t}(P_{15}) \cong \text{Ext}_{\mathcal{A}}^{s,t}(P_{15}^{18})$ for $s \geq 0$, $15 \leq t-s \leq 17$. Since $P_{15}^{18} = (S^{15} \cup_{2\iota} e^{16}) \cup_{\overline{\eta}} C(S^{16} \cup_{2\iota} e^{17})$, where $S^{16} \cup_{2\iota} e^{17} \xrightarrow{\overline{\eta}} S^{15}$ is an extension of the map $S^{16} \xrightarrow{\eta} S^{15}$, there is a cofibration sequence $\Sigma^{15} M_2 \cong S^{15} \cup_{2\iota} e^{16} \rightarrow P_{15}^{18} \rightarrow \Sigma^{17} M_2 \cong S^{17} \cup_{2\iota} e^{18}$ such that $0 \rightarrow H_*(\Sigma^{15} M_2) \rightarrow H_*(P_{15}^{18}) \rightarrow H_*(\Sigma^{17} M_2) \rightarrow 0$ is exact. Thus by (4.2) we have the following long

exact sequence.

$$(4.8) \quad \begin{aligned} \cdots \rightarrow \text{Ext}_{\mathcal{A}}^{s-1, t-17}(M_2) \xrightarrow{\delta} \text{Ext}_{\mathcal{A}}^{s, t-15}(M_2) \rightarrow \text{Ext}_{\mathcal{A}}^{s, t}(P_{15}^{18}) \\ \rightarrow \text{Ext}_{\mathcal{A}}^{s, t-17}(M_2) \xrightarrow{\delta} \text{Ext}_{\mathcal{A}}^{s+1, t-15}(M_2) \rightarrow \cdots \end{aligned}$$

From (4.8), Figure 2 (put $l = 15$ and $l = 17$) and the differentials $\delta(\overline{e_{17}}) = \overline{e_{15}}h_1$, $\delta(\overline{e_{18}h_1}) = 0$ in the spectral sequence for $\text{Ext}_{\mathcal{A}}^{*,*}(P_{15}^{18})$, we deduce the structure of $\text{Ext}_{\mathcal{A}}^{s,t}(P_{15}^{18})$ for $s \geq 0$, $15 \leq t - s \leq 17$. The results are given in Figure 3. This completes the computations for (4.3) (2).

We proceed to compute $\text{Ext}_{\mathcal{A}}^{s,t}(P^{14})$ for $s \geq 0$, $13 \leq t - s \leq 16$ in (4.3) (3). From the cofibration sequence $P^{14} \xrightarrow{i_{14}} P \xrightarrow{q_{15}} P_{15}$, and from (4.2) we have the following long exact sequence.

$$(4.9) \quad \begin{aligned} \cdots \rightarrow \text{Ext}_{\mathcal{A}}^{s-1, t}(P) \xrightarrow{(q_{15})^*} \text{Ext}_{\mathcal{A}}^{s-1, t}(P_{15}) \xrightarrow{\delta} \text{Ext}_{\mathcal{A}}^{s, t}(P^{14}) \\ \rightarrow \text{Ext}_{\mathcal{A}}^{s, t}(P) \xrightarrow{(q_{15})^*} \text{Ext}_{\mathcal{A}}^{s, t}(P_{15}) \rightarrow \cdots \end{aligned}$$

We recall from [26] the Ext groups $\text{Ext}_{\mathcal{A}}^{s,t}(P)$ for $s \geq 0$, $13 \leq t - s \leq 16$ as in Figure 4. From (4.9), Figure 3 and Figure 4 it is not difficult to deduce $\text{Ext}_{\mathcal{A}}^{s,t}(P^{14})$ for $s \geq 0$, $13 \leq t - s \leq 16$, noting that the map $\text{Ext}_{\mathcal{A}}^{s,t}(P) \xrightarrow{(q_{15})^*} \text{Ext}_{\mathcal{A}}^{s,t}(P_{15})$ is given by $(q_{15})_*(\widehat{h}_4) = \overline{e_{15}} \in \text{Ext}_{\mathcal{A}}^{0,15}(P_{15})$, and $\overline{e_{16}h_1} \in \text{Ext}_{\mathcal{A}}^{1,18}(P_{15})$ is not in the image of $(q_{15})_*$. These results are given in Figure 5. This completes the computations of (4.3) (3).

To compute $\text{Ext}_{\mathcal{A}}^{s,t}(P_{47}^{62})$ for $s \geq 0$, $61 \leq t - s \leq 63$ for (4.3) (4), we use the cofibration sequence $P_{47}^{48} \rightarrow P_{47}^{62} \rightarrow P_{49}^{62}$. Since $H^*(P_{49}^{62}) \cong \Sigma^{48}H^*(P^{14})$ as left \mathcal{A} -modules, $\text{Ext}_{\mathcal{A}}^{s,t}(P_{49}^{62}) \cong \text{Ext}_{\mathcal{A}}^{s, t-48}(P^{14})$ for all s, t . So by (4.2) we have the following long exact sequence.

$$(4.10) \quad \begin{aligned} \cdots \rightarrow \text{Ext}_{\mathcal{A}}^{s-1, t-48}(P^{14}) \xrightarrow{\delta} \text{Ext}_{\mathcal{A}}^{s, t}(P_{47}^{48}) \rightarrow \text{Ext}_{\mathcal{A}}^{s, t}(P_{47}^{62}) \\ \rightarrow \text{Ext}_{\mathcal{A}}^{s, t-48}(P^{14}) \xrightarrow{\delta} \text{Ext}_{\mathcal{A}}^{s+1, t}(P_{47}^{48}) \rightarrow \cdots \end{aligned}$$

From (4.10), Figure 2 (put $l = 47$) and Figure 5, it is not difficult to deduce $\text{Ext}_{\mathcal{A}}^{s,t}(P_{47}^{62})$ for $s \geq 0$, $61 \leq t - s \leq 63$. The results are given in Figure 6. The group extensions in Figure 6 are given in the following (4.11).

- (4.11) (1) $\overline{e_{55}h_0h_3h_0} = \overline{e_{48}h_0h_3^2}$ by the following (a), (b), and (c).
- (a) $\overline{e_{55}h_0h_3} = \{\overline{e_{55}h_0h_3}^*\}$ where $\overline{e_{55}h_0h_3}^* \equiv e_{55}\lambda_0\lambda_7 + e_{48}\lambda_7^2 \pmod{F(47)}$,
 - (b) $\overline{e_{48}h_0h_3^2} = \{\overline{e_{48}h_0h_3^2}^*\}$ where $\overline{e_{48}h_0h_3^2}^* \equiv e_{48}\lambda_7^2\lambda_0 \pmod{F(47)}$,
 - (c) $\overline{e_{55}h_0h_3}^*\lambda_0 + \overline{e_{48}h_0h_3^2}^* \equiv \delta(e_{55}\lambda_0\lambda_8 + e_{56}\lambda_0\lambda_7 + e_{54}\lambda_2\lambda_7 + e_{57}\lambda_3^2 + e_{52}\lambda_4\lambda_7 + e_{51}\lambda_5\lambda_7) \pmod{F(47)}$.
- (2) The extension from $\overline{e_{48}h_0h_3^2}$ to $\overline{e_{51}P^1h_2}$ in Figure 6 is nontrivial and will be explained in the proof of Theorem 2.39 later.
- (3) $\overline{e_{53}h_1c_0h_0} = \overline{e_{51}P^1h_2}$ by the following.
- (a) $P^1h_2 = \langle h_1, h_1c_0, h_0 \rangle \in \text{Ext}_{\mathcal{A}}^{5,16}$.
- (4) $\overline{e_{51}P^1h_2h_0} = \overline{e_{48}h_0^2d_0}$ because the following (a), (b), (c), and (d).




Figure 3: $\text{Ext}_A^{s,t}(P_{15}^{18})$, $s \geq 0$, $15 \leq t - s \leq 17$




Figure 4: $\text{Ext}_A^{s,t}(P)$, $s \geq 0$, $13 \leq t - s \leq 16$




Figure 5: $\text{Ext}_A^{s,t}(P^{14})$, $s \geq 0$, $13 \leq t - s \leq 16$




Figure 6: $\text{Ext}_A^{s,t}(P_{47}^{62})$, $s \geq 0$, $61 \leq t - s \leq 63$

- (a) Recall the following
(3.13) (3) $d_0 = \{d_0^* = \lambda_6 \lambda_2 \lambda_3^2 + \lambda_4^2 \lambda_3^2 + \lambda_2 \lambda_4 \lambda_5 \lambda_3\} \in \text{Ext}_{\mathcal{A}}^{4,18}$
(b) $\overline{e_{51} P^1 h_2} = \{\overline{e_{51} P^1 h_2^*}\}$ where $\overline{e_{51} P^1 h_2^*} \equiv e_{51} \lambda_0^4 \lambda_{11} + e_{48} \lambda_0 d_0^*$
 $\text{mod } F(47)$,
(c) $\overline{e_{48} h_0^2 d_0} = \{\overline{e_{48} h_0^2 d_0^*}\}$ where $\overline{e_{48} h_0^2 d_0^*} \equiv e_{48} \lambda_0 d_0^* \lambda_0 \text{ mod } F(47)$,
(d) $\overline{e_{51} P^1 h_2^*} \lambda_0 + \overline{e_{48} h_0^2 d_0^*} \equiv \delta(e_{51} \lambda_0^4 \lambda_{12} + e_{52} \lambda_0^4 \lambda_{11} + e_{51} \lambda_0^2 \lambda_6 \lambda_3^2 + e_{50} (\lambda_2 \lambda_0^3 \lambda_{11} + \lambda_4 \lambda_1 \lambda_2 \lambda_3^2) + e_{49} (\lambda_1 \lambda_2 \lambda_0^2 \lambda_{11} + \lambda_2 \lambda_4 \lambda_2 \lambda_3^2 + \lambda_5 \lambda_1 \lambda_2 \lambda_3^2)) \text{ mod } F(47)$.

To compute $\text{Ext}_{\mathcal{A}}^{s,t}(P_{46}^{48})$ for $s \geq 0$, $60 \leq t - s \leq 63$ for (4.3) (5), consider the cofibration sequence $S^{46} \rightarrow P_{46}^{48} = S^{46} \cup_{\overline{\eta}} C(S^{46}) \cup_{2\iota} e^{47} \rightarrow S^{47} \cup_{2\iota} e^{48}$. By (4.2) this induces the following long exact sequence of Ext groups.

$$(4.12) \quad \begin{aligned} \cdots \rightarrow \text{Ext}_{\mathcal{A}}^{s-1,t}(S^{47} \cup_{2\iota} e^{48}) \xrightarrow{\delta} \text{Ext}_{\mathcal{A}}^{s,t}(S^{46}) \rightarrow \text{Ext}_{\mathcal{A}}^{s,t}(P_{46}^{48}) \\ \rightarrow \text{Ext}_{\mathcal{A}}^{s,t}(S^{47} \cup_{2\iota} e^{48}) \xrightarrow{\delta} \text{Ext}_{\mathcal{A}}^{s+1,t}(S^{46}) \rightarrow \cdots \end{aligned}$$

From (4.12), Figure 1 and Figure 2 (put $l = 47$) it is not difficult to deduce the structure of $\text{Ext}_{\mathcal{A}}^{s,t}(P_{46}^{48})$ for $s \geq 0$, $61 \leq t - s \leq 63$. These results are given in Figure 7. The only group extension in Figure 7 that will be used in this paper is explained in the following.

(4.13) $\overline{e_{47} h_4 h_0} = \overline{e_{47} h_0 h_4} = \overline{e_{46} h_1 h_4} = \overline{e_{46} h_4 h_1}$ because $e_{47} \lambda_0 + e_{46} \lambda_1 \equiv \delta(e_{48}) \text{ mod } F(45)$.

This completes the computation for (4.3) (5).

From Figure 5 (put $l = 47$) and Figure 7 it is not difficult to deduce $\text{Ext}_{\mathcal{A}}^{s,t-s}(P_{46}^{62})$ for $s \geq 0$, $61 \leq t - s \leq 63$. These results are given in Figure 8. The group extensions given in Figure 8 are obtained by the same computations in (4.11) (1) through (4), except the followings.

- (4.14)** (1) $\overline{e_{47} h_4 h_0} = \overline{e_{46} h_4 h_1}$ follows from (4.13).
(2) $\overline{e_{48} h_0^2 d_0 h_0} = \overline{e_{46} P^1 c_0}$ because the following.
(a) $P^1 c_0 = \langle h_1, h_0^2 d_0, h_0 \rangle$.

This completes the computations for (4.3) (6).

In order to prove Theorem 2.39 and 2.40, we first prove two differentials and also show an infinite cycle in the ASS for $\pi_*^S(P_{46}^{62})$ and also in the ASS for $\pi_*^S(P_{47}^{62})$. The ranges of these two spectral sequences of our discussion are shown in Figure 6 and Figure 8. The two differentials are stated in Proposition 4.16. The infinite cycle is given in Proposition 4.17. We also need the following Ext group information: $\text{Ext}_{\mathcal{A}}^{s,t}(P_{46})$ for $s \geq 0$ and $61 \leq t - s \leq 63$, and $\text{Ext}_{\mathcal{A}}^{s,t}(P_{46}^{49})$ for $0 \leq s \leq 4$ and $62 \leq t - s \leq 64$. These results are given in Figure 9 and Figure 10. The computations of these groups follow the same procedure as the computation of, for example, $\text{Ext}_{\mathcal{A}}^{s,t}(M_2)$ in Figure 2, and will not be repeated here.

To prove Proposition 4.16 we need Lemma 4.15. We recall the stable map $S^{62} \xrightarrow{\hat{\theta}_5} P^{62}$ in (2.16). For $1 \leq k \leq 62$ let $P^{62} \xrightarrow{q_k} P_k^{62}$ be the collapsing map and let $P^k \xrightarrow{i_k} P^{62}$ be the inclusion map. Let $P_{46}^{62} \xrightarrow{q'} P_{47}^{62}$ and $P_{46}^{62} \xrightarrow{q''} P_{49}^{62}$ also be the collapsing maps. Recall the following (2.26) and (2.32) in section 2.




Figure 7: $\text{Ext}_A^{s,t}(P_{46}^{48})$, $s \geq 0$, $61 \leq t - s \leq 63$




Figure 8: $\text{Ext}_A^{s,t}(P_{46}^{62})$, $s \geq 0$, $61 \leq t - s \leq 63$




Figure 9: $\text{Ext}_{\mathcal{A}}^{s,t}(P_{46})$, $s \geq 0$, $61 \leq t - s \leq 63$




Figure 10: $\text{Ext}_{\mathcal{A}}^{s,t}(P_{46}^{49})$, $0 \leq s \leq 4$, $62 \leq t - s \leq 64$

Proposition 2.26. *The composite $S^{62} \xrightarrow{2\iota} S^{62} \xrightarrow{\widehat{\theta}_5} P^{62} \xrightarrow{q_k} P_k^{62}$ is essential if and only if $1 \leq k \leq 51$.*

Proposition 2.32. *$AF(q_{47}\widehat{\theta}_5) \geq 3$ for the composite $S^{62} \xrightarrow{\theta_5} P^{62} \xrightarrow{q_{47}} P_{47}^{62}$.*

Lemma 4.15. *The stable homotopy element $2q_{46}\widehat{\theta}_5 \in \pi_{62}^S(P_{46}^{62})$ is detected by $\overline{e_{51}P^1h_2} \in \text{Ext}_{\mathcal{A}}^{5,67}(P_{46}^{62})$ in the ASS for $\pi_*^S(P_{46}^{62})$.*

Proof. Refer to Figure 8 for the ASS for $\pi_*^S(P_{46}^{62})$ in the range of our discussion. $\widehat{\theta}_5 \in \pi_{62}^S(P^{62})$ is detected by $\widehat{h}_5h_5 \in \text{Ext}_{\mathcal{A}}^{1,63}(P^{62})$, so $AF(\widehat{\theta}_5) = 1$ in the ASS for $\pi_*^S(P^{62})$. It is clear that $(q_{46})_*(\widehat{h}_5) = 0$. So $AF(q_{46}\widehat{\theta}_5) \geq 2$, and this implies that $AF(2q_{46}\widehat{\theta}_5) \geq 3$ for the ASS for $\pi_*^S(P_{46}^{62})$. From Figure 8 we see that $\overline{2q_{46}\widehat{\theta}_5}$ must be detected by one of the following four Ext group classes: $\overline{e_{48}h_0h_3^2} \in \text{Ext}_{\mathcal{A}}^{3,65}(P_{46}^{62})$, $\overline{e_{51}P^1h_2} \in \text{Ext}_{\mathcal{A}}^{5,67}(P_{46}^{62})$, $\overline{e_{48}h_0^2d_0} \in \text{Ext}_{\mathcal{A}}^{6,68}(P_{46}^{62})$, or $\overline{e_{46}P^1c_0} \in \text{Ext}_{\mathcal{A}}^{7,69}(P_{46}^{62})$. If $AF(2q_{46}\widehat{\theta}_5) \geq 6$, that is, if $2q_{46}\widehat{\theta}_5$ is detected by $\overline{e_{48}h_0^2d_0}$ or $\overline{e_{46}P^1c_0}$, then

$$(*) \quad AF(2q_{49}\widehat{\theta}_5 = q''(2q_{46}\widehat{\theta}_5)) \geq 6 \quad \text{in the ASS for } \pi_*^S(P_{49}^{62}).$$

Since $\text{Ext}_{\mathcal{A}}^{s,62+s}(P_{49}^{62}) \cong \text{Ext}_{\mathcal{A}}^{s,14+s}(P^{14})$, from Figure 5 we see that $\text{Ext}_{\mathcal{A}}^{s,62+s}(P_{49}^{62}) = 0$ for $s \geq 6$. This together with (*) imply $2q_{49}\widehat{\theta}_5 = 0$ in $\pi_{62}^S(P_{49}^{62})$. But this contradicts Proposition 2.26 (for $k = 49$). Therefore $AF(2q_{46}\widehat{\theta}_5) \leq 5$ in the ASS for $\pi_*^S(P_{46}^{62})$. It follows that $\overline{S^{62} \xrightarrow{2q_{46}\widehat{\theta}_5} P_{46}^{62}}$, which is non-zero by Proposition 2.20, must be detected by $\alpha = \overline{e_{48}h_0h_3^2}$ or $\overline{e_{51}P^1h_2}$ in Figure 8. From Figure 6 we see that $(q')_*(\alpha) \neq 0$ in $\text{Ext}_{\mathcal{A}}^{*,*}(P_{47}^{62})$, which is also denoted by $\overline{e_{48}h_0h_3^2}$ and $\overline{e_{51}P^1h_2}$ respectively, where $P_{46}^{62} \xrightarrow{q'} P_{47}^{62}$ is the collapsing map. Comparing Figure 8 and Figure 6 we see that $\text{Ext}_{\mathcal{A}}^{s,63+s}(P_{46}^{62}) \cong \text{Ext}_{\mathcal{A}}^{s,63+s}(P_{47}^{62})$ for $0 \leq s \leq 3$, so if α is not a boundary in the ASS for $\pi_*^S(P_{46}^{62})$, then $(q')_*(\alpha)$ is not a boundary in the ASS for $\pi_*^S(P_{47}^{62})$ either. Therefore $2q_{47}\widehat{\theta}_5 = q'(2q_{46}\widehat{\theta}_5)$ is detected by $(q')_*(\alpha)$ ($= \overline{e_{48}h_0h_3^2}$ or $\overline{e_{51}P^1h_2}$) in the ASS for $\pi_*^S(P_{47}^{62})$. By Proposition 2.32, $AF(q_{47}\widehat{\theta}_5) \geq 3$. So $AF(2q_{47}\widehat{\theta}_5) \geq 4$. Therefore $2q_{47}\widehat{\theta}_5$ must be detected by $\overline{e_{51}P^1h_2} \in \text{Ext}_{\mathcal{A}}^{5,67}(P_{47}^{62})$. So $2q_{46}\widehat{\theta}_5$ has to be detected by $\overline{e_{51}P^1h_2} \in \text{Ext}_{\mathcal{A}}^{5,67}(P_{46}^{62})$. This completes the proof of Lemma 4.15. \square

Proposition 4.16. *$d_5(\overline{e_{63}h_0}) = \overline{e_{48}h_0^2d_0}$ and $d_5(\overline{e_{63}h_0^2}) = \overline{e_{46}P^1c_0}$ in the ASS for $\pi_*^S(P_{46}^{62})$ and the ASS for $\pi_*^S(P_{47}^{62})$.*

Proof. By Lemma 4.15, $2q_{46}\widehat{\theta}_5 = q_{46}(2\widehat{\theta}_5)$ is detected by $\overline{e_{51}P^1h_2}$ in the ASS for $\pi_*^S(P_{46}^{62})$. Consider the inclusion map $P_{46}^{62} \xrightarrow{i_{62}} P_{46}$. Recall that $2\widehat{\theta}_5 = 0$ in $\pi_*^S(P)$, so that $i_{62}q_{46}(2\widehat{\theta}_5) = 0$ in $\pi_*^S(P_{46})$. From Figure 9 we see $(i_{62})_*(\overline{e_{51}P^1h_2}) \neq 0$ in $\text{Ext}_{\mathcal{A}}^{5,67}(P_{46})$, which is also denoted by $\overline{e_{51}P^1h_2}$. So $\overline{e_{51}P^1h_2}$ must be a boundary in the ASS for $\pi_*^S(P_{46})$. Comparing Figure 8 and Figure 9, we see $\text{Ext}_{\mathcal{A}}^{s,63+s}(P_{46}) \cong \text{Ext}_{\mathcal{A}}^{s,63+s}(P_{46}^{62})$ for $1 \leq s \leq 3$ and $\text{Ext}_{\mathcal{A}}^{0,63}(P_{46}) = \mathbb{Z}/2(\overline{e_{63}})$. Thus $d_5(\overline{e_{63}}) = \overline{e_{51}P^1h_2}$ in the ASS for $\pi_*^S(P_{46})$. From Figure 8, $\overline{e_{51}P^1h_2}h_0 = \overline{e_{48}h_0^2d_0}$ and $\overline{e_{51}P^1h_2}h_0^2 = \overline{e_{48}h_0^2d_0}h_0 = \overline{e_{46}P^1c_0}$, so it follows that $d_5(\overline{e_{63}h_0}) = \overline{e_{48}h_0^2d_0}$ and $d_5(\overline{e_{63}h_0^2}) = \overline{e_{46}P^1c_0}$ in the ASS for $\pi_*^S(P_{46})$. Comparing Figure 6, Figure 8, and Figure 9 we see that these two differentials also hold in the ASS for $\pi_*^S(P_{46}^{62})$ and the ASS for $\pi_*^S(P_{47}^{62})$. This proves Proposition 4.16. \square

We proceed to show an infinite cycle in the ASS for $\pi_*^S(P_{46}^{62})$ and the ASS for $\pi_*^S(P_{47}^{62})$.

Proposition 4.17. $\overline{e_{49}h_3^2}$ is an infinite cycle in the ASS for $\pi_*^S(P_{46}^{62})$ and the ASS for $\pi_*^S(P_{47}^{62})$.

Proof. Consider the complex $P_{46}^{49} = (S^{46} \vee S^{47}) \cup_{(\eta, 2\nu)} e^{48} \cup_{\eta} e^{49}$. We have the following cofibration sequence.

$$\begin{array}{ccccccc} \cdots & \longrightarrow & P_{46}^{49} = (S^{46} \vee S^{47}) \cup_{(\eta, 2\nu)} e^{48} \cup_{\eta} e^{49} & \longrightarrow & S^{49} & \longrightarrow & (S^{47} \vee S^{48}) \cup_{(\eta, 2\nu)} e^{49} \longrightarrow \cdots \\ & & & & \uparrow \sigma^2 & & \\ & & & & S^{63} & & \end{array}$$

$\swarrow \widetilde{\sigma}^2$

Since $\eta\sigma^2 = 0$ there is a coextension $S^{63} \xrightarrow{\widetilde{\sigma}^2} P_{46}^{49}$ to $S^{63} \xrightarrow{\sigma^2} S^{49}$. Since $AF(\sigma^2) = 2$ in the ASS for π_*^S , we have $AF(\widetilde{\sigma}^2) \leq 2$ in the ASS for $\pi_*^S(P_{46}^{49})$. From Figure 10 we see $S^{63} \xrightarrow{\widetilde{\sigma}^2} P_{46}^{49}$ is detected by $\overline{e_{49}h_3^2}$ in the ASS for $\pi_*^S(P_{46}^{49})$. So $\overline{e_{49}h_3^2} \in \text{Ext}_{\mathcal{A}}^{2,65}(P_{46}^{49})$ is an infinite cycle and in the ASS for $\pi_*^S(P_{46}^{49})$. It follows that $\overline{e_{49}h_3^2}$ is also an infinite cycle in the ASS for $\pi_*^S(P_{46}^{62})$ and also in the ASS for $\pi_*^S(P_{47}^{62})$. This completes the proof of Proposition 4.17. \square

We proceed to prove Theorem 2.39 and Theorem 2.40.

To describe the stable homotopy groups $\pi_{62}^S(P_{46}^{62})$ and $\pi_{62}^S(P_{47}^{62})$ in Theorem 2.39 and Theorem 2.40, we need to recall the stable homotopy elements $y, w, x \in \pi_{62}^S(P_{46}^{62})$ and $v \in \pi_{62}^S(P_{47}^{62})$ defined in (2.35), (2.36), (2.37), and (2.38) respectively. In Propositions 4.18, 4.19, 4.20, and 4.21 below we recall their definitions and show that they are essential. Recall the Hopf classes $S^1 \xrightarrow{\eta} S^0$, $S^3 \xrightarrow{\nu} S^0$ and $S^7 \xrightarrow{\sigma} S^0$. To show that the stable homotopy elements y, w, x and v are essential we need the information of the following Ext group structures: $\text{Ext}_{\mathcal{A}}^{s,t}(S^{47} \cup_{\sigma} e^{55})$ for $s \geq 0$ and $61 \leq t - s \leq 63$, $\text{Ext}_{\mathcal{A}}^{s,t}(S^{47} \cup_{\nu} e^{55} \cup_{\eta} e^{53})$ for $s \geq 0$ and $61 \leq t - s \leq 63$, and $\text{Ext}_{\mathcal{A}}^{s,t}(S^{47} \cup_{2\nu} e^{48} \cup_{\nu} e^{51} \cup_{\eta} e^{53})$ for $0 \leq s \leq 4$ and $61 \leq t - s \leq 63$. These are given Figure 11 and Figure 12. The computations of these groups are analogous to that of $\text{Ext}_{\mathcal{A}}^{s,t}(M_2)$ in Figure 2, and will not be repeated here.

Proposition 4.18. Let $P_{46}^{48} \xrightarrow{i_{48}} P_{46}^{62}$ be the inclusion map. Let $S^{62} \xrightarrow{\widetilde{\sigma}^2} P_{46}^{48}$ be some coextension of $S^{62} \xrightarrow{\sigma^2} S^{48}$ as shown in the following diagram.

$$\begin{array}{ccccccc} \cdots & \longrightarrow & P_{46}^{48} & \xrightarrow{q_{48}} & S^{48} & \xrightarrow{\eta \vee 2\nu} & S^{47} \vee S^{48} \longrightarrow \cdots \\ & & & & \uparrow \sigma^2 & & \\ & & & & S^{62} & & \end{array}$$

$\swarrow \widetilde{\sigma}^2$

Let y be the composite $S^{62} \xrightarrow{\widetilde{\sigma}^2} P_{46}^{48} \xrightarrow{i_{48}} P_{46}^{62}$. Then y is detected by $\overline{e_{47}h_4} \in \text{Ext}_{\mathcal{A}}^{1,63}(P_{46}^{62})$ in the ASS for $\pi_*^S(P_{46}^{62})$.

Proof. $S^{62} \xrightarrow{\sigma^2} S^{48}$ is detected by $h_3^2 \in \text{Ext}_{\mathcal{A}}^{2,16}$ in the ASS for π_*^S , so $AF(\sigma^2) = 2$. Thus $AF(\widetilde{\sigma}^2) \leq 2$ in the ASS for $\pi_*^S(P_{46}^{48})$. From Figure 7, the nonzero stable




Figure 11: $\text{Ext}_{\mathcal{A}}^{s,t}(S^{47} \cup_{\sigma} e^{55})$, $0 \leq s \leq 4$, $61 \leq t - s \leq 63$




Figure 12: $\text{Ext}_{\mathcal{A}}^{s,t}(W_1 = S^{47} \cup_{\nu} e^{51} \cup_{\eta} e^{53})$, $0 \leq s \leq 4$, $61 \leq t - s \leq 63$




Figure 13: $\text{Ext}_{\mathcal{A}}^{s,t}(W_2 = S^{47} \cup_{2\iota} e^{48} \cup_{\nu} e^{51} \cup_{\eta} e^{53})$, $0 \leq s \leq 4$, $61 \leq t - s \leq 63$

homotopy class $\widetilde{\sigma}^2$ must be detected by $\overline{e_{47}h_4} \in \text{Ext}_{\mathcal{A}}^{1,63}(P_{46}^{48})$ or $\overline{e_{47}h_0h_4} \in \text{Ext}_{\mathcal{A}}^{2,64}(P_{46}^{48})$. Suppose $\widetilde{\sigma}^2$ is detected by $\overline{e_{47}h_0h_4}$ in the ASS for $\pi_*^S(P_{46}^{48})$, then $AF(\widetilde{\sigma}^2) = 2$. It is not difficult to see that $(q_{48})_*(\overline{e_{47}h_0h_4}) = 0$ in $\text{Ext}_{\mathcal{A}}^{2,64}$, so it follows that $AF(\sigma^2 = q_{48}(\widetilde{\sigma}^2)) \geq 3$ in the ASS for π_*^S , but we have seen that $AF(\sigma^2) = 2$ so this is a contradiction. Therefore $\widetilde{\sigma}^2$ is detected by $\overline{e_{47}h_4}$ in the ASS for $\pi_*^S(P_{46}^{48})$. From Figure 8, $\overline{e_{47}h_4} \neq 0$ in $\text{Ext}_{\mathcal{A}}^{1,63}(P_{46}^{62})$ is not a boundary in the ASS for $\pi_*^S(P_{46}^{62})$, so $y = i\sigma^2 \in \pi_{62}^S(P_{46}^{62})$ is essential and is detected by $\overline{e_{47}h_4} \in \text{Ext}_{\mathcal{A}}^{1,63}(P_{46}^{62})$ in the ASS for $\pi_*^S(P_{46}^{62})$. This proves Proposition 4.18. \square

Proposition 4.19. *Let $S^{47} \cup_{\sigma} e^{55} \xrightarrow{j} P_{46}^{62}$ be the inclusion map. Let $S^{62} \xrightarrow{2\sigma} S^{47} \cup_{\sigma} e^{55}$ be some coextension of $S^{62} \xrightarrow{2\sigma} S^{55}$ as shown in the following diagram.*

$$\begin{array}{ccccccc} \dots & \longrightarrow & S^{47} \cup_{\sigma} e^{55} & \xrightarrow{q_{55}} & S^{55} & \xrightarrow{\sigma} & S^{48} \longrightarrow \dots \\ & & & & \uparrow 2\sigma & & \\ & & & & S^{62} & & \\ & & \swarrow \overline{2\sigma} & & & & \end{array}$$

Let w be the composite $S^{62} \xrightarrow{2\sigma} S^{47} \cup_{\sigma} e^{55} \xrightarrow{j} P_{46}^{62}$. Then w is detected by $\overline{e_{47}h_4} \in \text{Ext}_{\mathcal{A}}^{1,63}(P_{46}^{62})$ in the ASS for $\pi_*^S(P_{46}^{62})$.

Proof. $S^{62} \xrightarrow{2\sigma} S^{55}$ is detected by h_0h_3 in the ASS for π_*^S , so $AF(2\sigma) = 2$. Thus $AF(\overline{2\sigma}) \leq 2$ in the ASS for $\pi_*^S(S^{47} \cup_{\sigma} e^{55})$. From Figure 11, the nonzero stable homotopy class $\overline{2\sigma}$ must be detected by $\overline{e_{47}h_4} \in \text{Ext}_{\mathcal{A}}^{1,63}(S^{47} \cup_{\sigma} e^{55})$ or $\overline{e_{47}h_0h_4} \in \text{Ext}_{\mathcal{A}}^{2,64}(S^{47} \cup_{\sigma} e^{55})$. Suppose $\overline{2\sigma}$ is detected by $\overline{e_{47}h_0h_4}$ in the ASS for $\pi_*^S(S^{47} \cup_{\sigma} e^{55})$, then $AF(\overline{2\sigma}) = 2$. It is not difficult to see that $(q_{55})_*(\overline{e_{47}h_0h_4}) = 0$ in $\text{Ext}_{\mathcal{A}}^{2,64}$, so it follows that $AF(2\sigma = q_{55}\overline{2\sigma}) \geq 3$, but we already have shown that $AF(2\sigma) = 2$ so this is a contradiction. Therefore $\overline{2\sigma}$ is detected by $\overline{e_{47}h_4}$ in the ASS for $\pi_*^S(S^{47} \cup_{\sigma} e^{55})$. From Figure 8, $\overline{e_{47}h_4} \neq 0$ in $\text{Ext}_{\mathcal{A}}^{1,63}(P_{46}^{62})$ is not a boundary in the ASS for $\pi_*^S(P_{46}^{62})$, so $w = j\overline{2\sigma} \in \pi_{62}^S(P_{46}^{62})$ is essential and is detected by $\overline{e_{47}h_4} \in \text{Ext}_{\mathcal{A}}^{1,63}(P_{46}^{62})$ in the ASS for $\pi_*^S(P_{46}^{62})$. This proves Proposition 4.19. \square

Proposition 4.20. $x = y + w \in \pi_{62}^S(P_{46}^{62})$ is detected by $\overline{e_{55}h_0h_3} + \varepsilon\overline{e_{47}h_0h_4} \in \text{Ext}_{\mathcal{A}}^{2,64}(P_{46}^{62})$, $\varepsilon = 0$ or 1.

Proof. Let $P_{46}^{62} \xrightarrow{q_{49}} P_{49}^{62}$ be the quotient map and consider the composite $S^{62} \xrightarrow{x=y+w} P_{46}^{62} \xrightarrow{q_{49}} P_{49}^{62}$. From the definition of y it is not difficult to see that $q_{49}y = 0$ in $\pi_{62}^S(P_{49}^{62})$, as q_{49} pinches P_{46}^{48} . It is also not difficult to see that S^{55} is a subcomplex of P_{49}^{62} . Let $S^{55} \xrightarrow{i_{55}} P_{49}^{62}$ be the inclusion map. Then from the definition of x and w , $q_{49}x = q_{49}(y + w) = q_{49}w$ is the composite $S^{62} \xrightarrow{2\sigma} S^{55} \xrightarrow{i_{55}} P_{49}^{62}$. $S^{62} \xrightarrow{2\sigma} S^{55}$ is detected by h_0h_3 in the ASS for π_*^S . Recall that, since $H_*(P_{49}^{62}) \cong H_{*-48}(P^{14})$ as left- \mathcal{A} modules, $\text{Ext}_{\mathcal{A}}^{s,t}(P_{49}^{62}) \cong \text{Ext}_{\mathcal{A}}^{s,t-48}(P^{14})$. $\overline{e_{55}h_0h_3} \in \text{Ext}_{\mathcal{A}}^{2,64}(P_{49}^{62})$ corresponds to $\widehat{h_3h_0h_3} \in \text{Ext}_{\mathcal{A}}^{2,16}(P^{14})$ under this isomorphism. So from Figure 5, we deduce that $\overline{e_{55}h_0h_3} \in \text{Ext}_{\mathcal{A}}^{2,64}(P_{49}^{62})$ is not a boundary in the ASS for $\pi_*^S(P_{49}^{62})$ and $\text{Ext}_{\mathcal{A}}^{2,64}(P_{49}^{62}) = \mathbb{Z}/2(\overline{e_{55}h_0h_3})$. So $q_{49}x = i_{55}(2\sigma) \in \pi_{62}^S(P_{49}^{62})$ is detected by $\overline{e_{55}h_0h_3} \in \text{Ext}_{\mathcal{A}}^{2,64}(P_{49}^{62})$, and $AF(q_{49}x = i_{55}(2\sigma)) = 2$ in the ASS for $\pi_*^S(P_{49}^{62})$. It

- (3) In the ASS for $\pi_*^S(P_{46}^{62})$, $d_5(\overline{e_{63}h_0}) = \overline{e_{48}h_0^2d_0}$, and $d_5(\overline{e_{63}h_0h_0}) = \overline{e_{46}P^1c_0}$ by Proposition 4.16, so the classes $\overline{e_{48}h_0^2d_0}, \overline{e_{46}P^1c_0}$ do not represent homotopy classes. It follows that the order of x is at most 8.
- (4) To show that $2x \neq 0$ in $\pi_*^S(P_{46}^{62})$, recall from (4.11) (1) that there is an extension $\overline{e_{55}h_0h_3h_0} = \overline{e_{48}h_0h_3^2}$ and that $\overline{e_{47}h_0^2h_4} = 0$ in $\text{Ext}_A^{3,65}(P_{46}^{62})$. Since x is detected by $\overline{e_{55}h_0h_3} + \varepsilon\overline{e_{47}h_0h_4}$, $(\overline{e_{55}h_0h_3} + \varepsilon\overline{e_{47}h_0h_4})h_0 = \overline{e_{55}h_0h_3h_0} = \overline{e_{48}h_0h_3^2}$ is an infinite cycle in the ASS for $\pi_*^S(P_{46}^{62})$, and from Figure 8 and (3) we see $\overline{e_{48}h_0h_3^2}$ is not a boundary. So $2x$ is detected by $\overline{e_{48}h_0h_3^2} = \overline{e_{55}h_0h_3h_0}$ in the ASS for $\pi_*^S(P_{46}^{62})$.
- (5) To show that $4x \neq 0$, note that $4y = 0$ by (2), so $4x = 4(w + y) = 4w$. Consider the following diagram, where $S^{62} \xrightarrow{2\sigma} S^{47} \cup_\sigma e^{55}$ is the coextension as in Proposition 4.19, the map $S^{47} \cup_\sigma e^{55} \xrightarrow{i'} P_{46}^{55}$ is the inclusion of the subcomplex $S^{47} \cup_\sigma e^{55}$, and for each $51 \leq l \leq 55$, $S^l \xrightarrow{f_l} P_{46}^l \xrightarrow{i_l} P_{46}^{l+1} \xrightarrow{q_{l+1}} S^{l+1}$ is a cofibration sequence.

$$\begin{array}{ccccccc}
& & & S^{51} & \xrightarrow{\sigma_{51}} & P_{46}^{51} & \xrightarrow{q_{51}} & S^{51} \\
& & & & & \downarrow i_{51} & & \\
& & & & & \vdots & & \\
& & & S^{54} & \xrightarrow{\sigma_{54}} & P_{46}^{54} & \xrightarrow{q_{54}} & S^{54} \\
& & & & & \downarrow i_{54} & & \\
& & & S^{55} & \xrightarrow{\sigma_{55}} & P_{46}^{55} & \xrightarrow{q_{55}} & S^{55} \\
& & & & & \downarrow i_{55} & & \\
& & & S^{56} & \xrightarrow{\sigma_{56}} & P_{46}^{56} & \xrightarrow{q_{56}} & S^{56} \\
& & & & & \downarrow i_{56} & & \\
& & & & & \vdots & & \\
& & & & & \downarrow & & \\
S^{62} & \xrightarrow{2\iota} & S^{62} & \xrightarrow{2\iota} & S^{62} & \xrightarrow{w} & P_{46}^{62}
\end{array}$$

$\widetilde{4w}$ (curved arrow from S^{62} to S^{51})
 \overline{w}' (curved arrow from S^{62} to P_{46}^{55})
 σ (curved arrow from S^{62} to P_{46}^{56})
 $\overline{w} = (i'(2\sigma))$ (curved arrow from S^{62} to P_{46}^{55})

$S^{62} \xrightarrow{\overline{w} = i'(2\sigma)} P_{46}^{55}$ is a lifting of $S^{62} \xrightarrow{w} P_{46}^{62}$. Note that from the cell structure of P_{46}^{56} , $q_{55}\sigma_{55} = 2\iota$. So by choosing another lifting $S^{62} \xrightarrow{\overline{w}' = \overline{w} + \sigma_{55}\sigma} P_{46}^{55}$ of $S^{62} \xrightarrow{w} P_{46}^{62}$, we have $q_{55}\overline{w}' = 0$, and $S^{62} \xrightarrow{\overline{w}'} P_{46}^{55}$ can be lifted to $S^{62} \xrightarrow{\overline{w}''} P_{46}^{54}$ with $q_{54}\overline{w}'' = \eta\sigma$. From this and following similar arguments as in (1), it is not difficult to see that $S^{62} \xrightarrow{4\iota} S^{62} \xrightarrow{w} P_{46}^{62}$ has a lifting $S^{62} \xrightarrow{4\widetilde{w}} P_{46}^{51}$ with $q_{51}(4\widetilde{w}) = \langle \eta, \eta^2\sigma, 2\iota \rangle$. The Toda bracket $\langle \eta, \eta^2\sigma, 2\iota \rangle$ is

detected by $P^1 h_2 \in \text{Ext}_{\mathcal{A}}^{5,16}$ in the ASS for π_*^S . By Proposition 4.17 and (3), $\overline{e_{51} P^1 h_2} \in \text{Ext}_{\mathcal{A}}^{5,67}(P_{46}^{62})$ is not a boundary in the ASS for $\pi_*^S(P_{46}^{62})$. It follows that $4x$ is detected by $\overline{e_{51} P^1 h_2}$ in the ASS for $\pi_*^S(P_{46}^{62})$. This together with (3) proves that the order of x is 8.

This completes the proof of Theorem 2.39. \square

In order to prove Theorem 2.40 we need to recall the stable homotopy element $v \in \pi_{62}^S(P_{47}^{62})$ in (2.38). In Proposition 4.21 below we recall its definition and show that it is essential.

Proposition 4.21. $W_1 = S^{47} \cup_{\nu} e^{51} \cup_{\eta} e^{53}$ is a subcomplex of P_{47}^{62} . Let $W_1 \xrightarrow{j'} P_{47}^{62}$ be the inclusion map. Let $S^{62} \xrightarrow{\tilde{\eta}\varepsilon} W_1$ be some coextension of $S^{62} \xrightarrow{\eta\varepsilon} S^{53}$ as in the following diagram.

$$\begin{array}{ccccccc} \dots & \longrightarrow & W_1 = S^{47} \cup_{\nu} e^{51} \cup_{\eta} e^{53} & \xrightarrow{q_4} & S^{53} & \longrightarrow & S^{48} \cup_{\nu} e^{52} \longrightarrow \dots \\ & & & & \uparrow \eta\varepsilon & & \\ & & & & S^{62} & & \\ & & \nwarrow \tilde{\eta}\varepsilon & & & & \end{array}$$

Let v be the composite $S^{62} \xrightarrow{\tilde{\eta}\varepsilon} W_1 \xrightarrow{j'} P_{47}^{62}$. Then v is detected by $\overline{e_{53} h_1 c_0} \in \text{Ext}_{\mathcal{A}}^{4,66}(P_{47}^{62})$ in the ASS for $\pi_*^S(P_{47}^{62})$.

Proof. $S^{62} \xrightarrow{\eta\varepsilon} S^{53}$ is detected by $h_1 c_0 \in \text{Ext}_{\mathcal{A}}^{4,13}$ in the ASS for π_*^S , so $AF(\tilde{\eta}\varepsilon) \leq 4$ in the ASS for $\pi_*^S(W_1)$. From Figure 12 we see that $\text{Ext}_{\mathcal{A}}^{1,63}(W_1) = 0$, $\text{Ext}_{\mathcal{A}}^{2,64}(W_1) = \mathbb{Z}/2(\overline{e_{47} h_4})$, and $\text{Ext}_{\mathcal{A}}^{3,65}(W_1) = \mathbb{Z}/2(\overline{e_{47} h_0 h_4})$. It is well-known that in the ASS for π_*^S , $d_2(h_4) = h_0 h_3^2$ and $d_3(h_0 h_4) = h_0 d_0$, so in the ASS for $\pi_*^S(S^{47} \cup_{\nu} e^{51} \cup_{\eta} e^{53})$, $d_2(\overline{e_{47} h_4}) = \overline{e_{47} h_0 h_3^2} \neq 0$ and $d_3(\overline{e_{47} h_0 h_4}) = \overline{e_{47} h_0 d_0} \neq 0$. It follows that $AF(\tilde{\eta}\varepsilon) \geq 3$.

Consider the subcomplex $W_2 = S^{47} \cup_{2\nu} e^{48} \cup_{\nu} e^{51} \cup_{\eta} e^{53} \subset P_{47}^{62}$. Let $W_1 \xrightarrow{i_5} W_2 \xrightarrow{j_1} P_{47}^{62}$ be inclusion maps and let $W_2 \xrightarrow{q_5} S^{53}$ be the collapsing map. Then since $AF(\tilde{\eta}\varepsilon) \geq 3$ in the ASS for $\pi_*^S(W_1)$ and since $AF(q_5(i_5 \tilde{\eta}\varepsilon)) = \eta\varepsilon = 4$, we see $3 \leq AF(i_5 \tilde{\eta}\varepsilon) \leq 4$ in the ASS for $\pi_*^S(W_2)$. If $AF(i_5 \tilde{\eta}\varepsilon) = 3$ in the ASS for $\pi_*^S(W_2)$ then from Figure 13 we see that $i_5 \tilde{\eta}\varepsilon$ must be detected by $\overline{e_{53} h_1^2 h_3} \in \text{Ext}_{\mathcal{A}}^{3,65}(W_2)$. But it is not difficult to see that $(q_5)_*(\overline{e_{53} h_1^2 h_3}) = h_1^2 h_3 \neq 0$ in $\text{Ext}_{\mathcal{A}}^{3,12}$, so it follows that $AF(q_5(i_5 \tilde{\eta}\varepsilon)) = q_5 \tilde{\eta}\varepsilon = \eta\varepsilon = 3$, and this is a contradiction because $AF(\eta\varepsilon) = 4$. Therefore $AF(i_5 \tilde{\eta}\varepsilon) = 4$. From Figure 13 we see that $i_5 \tilde{\eta}\varepsilon$ must be detected by $\overline{e_{53} h_1 c_0} \in \text{Ext}_{\mathcal{A}}^{3,65}(W_2)$. From Figure 6, Proposition 4.16, and Proposition 4.17, $\overline{e_{53} h_1 c_0} \neq 0$ is not a boundary in the ASS for $\pi_*^S(P_{47}^{62})$, so $v = j_1(i_5 \tilde{\eta}\varepsilon)$ is detected by $\overline{e_{53} h_1 c_0} \in \text{Ext}_{\mathcal{A}}^{4,66}(P_{47}^{62})$. This proves Proposition 4.21. \square

Now we prove Theorem 2.40, which is restated as follows.

Theorem 2.40. Let $S^{62} \xrightarrow{\tilde{y}} P_{47}^{62}$ be the composite $S^{62} \xrightarrow{y} P_{46}^{62} \xrightarrow{q_1} P_{47}^{62}$, $S^{62} \xrightarrow{\tilde{w}} P_{47}^{62}$ be the composite $S^{62} \xrightarrow{w} P_{46}^{62} \xrightarrow{q_1} P_{47}^{62}$, and $S^{62} \xrightarrow{\tilde{x}=\tilde{y}+\tilde{w}} P_{47}^{62}$. Then

$$\pi_{62}^S(P_{47}^{62}) = \mathbb{Z}/2(\tilde{y}) \oplus \mathbb{Z}/8(\tilde{x}) \oplus \mathbb{Z}/2(v - 2\tilde{x}).$$

And $AF(\tilde{y}) = 1$, $AF(\tilde{x}) = 2$, and $AF(v) = 4$.

Proof of Theorem 2.40. (1) By Proposition 4.21, $AF(v) = 4$ in the ASS for $\pi_*^S(P_{47}^{62})$. From Proposition 4.18 we see that $\overline{e_{47}h_4} \in \text{Ext}_{\mathcal{A}}^{1,63}(P_{46}^{62})$ detects the stable homotopy class $y \in \pi_{62}^S(P_{46}^{62})$. From Figure 6, $\overline{e_{47}h_4}$ is not a boundary in the ASS for $\pi_*^S(P_{47}^{62})$, so $\overline{e_{47}h_4}$ detects $\tilde{y} = q_1y$, and $AF(\tilde{y}) = 1$.

(2) $2\tilde{y} = 0$ because we have shown in the proof of Theorem 2.39 that $S^{62} \xrightarrow{2\iota} S^{62} \xrightarrow{y} P_{48}^{62}$ can be lifted to $S^{46} \subset P_{46}^{48}$.

(3) We already have shown in the proof of Theorem 2.39 that $\overline{e_{55}h_0h_3} \in \text{Ext}_{\mathcal{A}}^{2,64}(P_{46}^{62})$, $\overline{e_{48}h_0h_3^2} \in \text{Ext}_{\mathcal{A}}^{3,65}(P_{46}^{62})$, $\overline{e_{51}P^1h_2} \in \text{Ext}_{\mathcal{A}}^{5,67}(P_{46}^{62})$ detect the stable homotopy classes $x, 2x, 4x \in \pi_{62}^S(P_{46}^{62})$ respectively. From Figure 6, Proposition 4.16, and Proposition 4.17, the classes $\overline{e_{55}h_0h_3}, \overline{e_{48}h_0h_3^2}, \overline{e_{51}P^1h_2}$ are not boundaries in the ASS for $\pi_*^S(P_{47}^{62})$, so $\overline{e_{55}h_0h_3}$ detects $\tilde{x} = q_1x$ and $AF(\tilde{x}) = 2$, $\overline{e_{48}h_0h_3^2}$ detects $2\tilde{x} = q_1(2x)$, and $\overline{e_{51}P^1h_2}$ detects $4\tilde{x} = q_1(4x)$.

(4) We have shown in Proposition 4.21 that $\overline{e_{53}h_1c_0} \in \text{Ext}_{\mathcal{A}}^{4,66}(P_{47}^{62})$ detects $v \in \pi_{62}^S(P_{47}^{62})$. In (4.11) (3) we have seen $\overline{e_{53}h_1c_0h_0} = \overline{e_{51}P^1h_2}$. We have seen that $\overline{e_{51}P^1h_2}$ is not a boundary in the ASS for $\pi_*^S(P_{47}^{62})$. Therefore $2v$ is detected by $\overline{e_{55}P^1h_2} \in \text{Ext}_{\mathcal{A}}^{5,67}(P_{47}^{62})$.

(5) We have $d_5(e_{63}h_0) = \overline{e_{48}h_0^2d_0}$ by Proposition 4.16. So from Figure 6, $2v = 4\tilde{x}$ and $4v = 8\tilde{x} = 0$.

From (3), (4), and (5) it follows that $AF(v - 2\tilde{x}) = 3$, and that $2(v - 2\tilde{x}) = 2v - 4\tilde{x} = 0$. So the order of $v - 2\tilde{x}$ is 2. From $AF(\tilde{y}) = 1$, $AF(\tilde{x}) = 2$, $AF(2\tilde{x}) = 3$, $AF(4\tilde{x}) = 5$, and $AF(v - 2\tilde{x}) = 3$, it is not difficult to see that $\tilde{y}, \tilde{x}, v - 2\tilde{x}$ are linearly independent. This completes the proof of Theorem 2.40. \square

5 Proof of Proposition 1.6, Proposition 1.7, and Proposition 2.46

In this section we prove the technical results Proposition 1.6, Proposition 1.7, and Proposition 2.46. These are the remaining theorems in section 1 and section 2 that need to be proved to complete the proof of Theorem 1.2.

Proposition 1.6. *In the ASS for spheres, $d_r(h_6^3) = 0 \in \text{Ext}_{\mathcal{A}}^{3+r, 191+r}$ for $2 \leq r \leq 3$.*

Proposition 1.7. *In the ASS for spheres, $d_2(h_1g_4) \neq h_0^3g_4$.*

Proposition 2.46. *The composite $S^{188} \xrightarrow{\theta_6} S^{62} \xrightarrow{\hat{\theta}_5} P$ has $AF = 5$ in the Adams spectral sequence for $P = P^\infty$ ($AF = \text{Adams filtration}$).*

The order of our proving these propositions is 2.46, 1.6, and 1.7. Before we start proving Proposition 2.46, we first recall the following differential in the ASS for π_*^S mentioned in (1.1).

$$(5.1) \quad d_2(h_i) = h_{i-1}^2 h_0 \neq 0 \quad \text{in the ASS for } \pi_*^S.$$

We remark that it is not difficult to see the following differential (5.2) in the ASS for $\pi_*^S(P)$ from (5.1), (3.19), (3.25), and the structure of $\text{Ext}_{\mathcal{A}}^{s,t}(P)$ for $0 \leq s \leq 3$ as given in Theorem 3.24.

$$(5.2) \quad d_2(\hat{h}_i) = \hat{h}_{i-1} h_0 h_{i-1} \neq 0 \quad \text{in the ASS for } \pi_*^S(P).$$

To prove Proposition 2.46 we begin by proving the following result in Lemma 5.3. Recall the complex \tilde{P}^{62} and the map $P^{62} \xrightarrow{\tilde{q}} \tilde{P}^{62}$ in (2.29) which has the property that $\tilde{q}_*(\hat{h}_5 = \overline{e_{31}}) = 0$ for $\hat{h}_5 \in \text{Ext}_{\mathcal{A}}^{0,31}(P^{62}) = \mathbb{Z}/2$.

Lemma 5.3. *The composite $S^{188} \xrightarrow{\theta_6} S^{62} \xrightarrow{\hat{\theta}_5} P^{62} \xrightarrow{\tilde{q}} \tilde{P}^{62}$ has $AF \geq 5$ in the ASS for $\pi_*^S(\tilde{P}^{62})$.*

Proof. Recall (2.16) that $\hat{\theta}_5$ is detected by $\hat{h}_5 h_5$, so $AF(\hat{\theta}_5) = 1$. Since $\tilde{q}_*(\hat{h}_5) = 0$, $AF(\tilde{q}\hat{\theta}_5) \geq 2$. Consider the following two cases.

- (1) If $AF(\tilde{q}\hat{\theta}_5) = 2$, then $\tilde{q}\hat{\theta}_5 \in \pi_{62}^S(\tilde{P}^{62})$ is detected by a non-zero class $\alpha \in \text{Ext}_{\mathcal{A}}^{2,64}(\tilde{P}^{62}) = \mathbb{Z}/2(\overline{e_{31}h_3h_0}) \oplus \mathbb{Z}/2(\overline{e_{47}h_0h_4})$ ((3.5) (2)). So

$$\alpha = \varepsilon_1 \overline{e_{31}h_3h_0} + \varepsilon_2 \overline{e_{47}h_0h_4}, \quad \varepsilon_1, \varepsilon_2 = 0 \text{ or } 1 \text{ and } (\varepsilon_1, \varepsilon_2) \neq (0, 0).$$

It is not difficult to see that $\overline{e_{31}h_3} = \{\tilde{e}_{31}\lambda_{31}\} \in \text{Ext}_{\mathcal{A}}^{1,63}(\tilde{P}^{62})$ and $\overline{e_{47}} = \{e_{47}\} \in \text{Ext}_{\mathcal{A}}^{0,47}(\tilde{P}^{62})$. Thus

$$\begin{aligned} (\overline{e_{31}h_3h_0})h_6^2 &= \overline{e_{31}h_3}h_6^2h_0 = \{\tilde{e}_{31}\lambda_{31}\lambda_{63}^2\lambda_0\} = 0, \\ (\overline{e_{47}h_0h_4})h_6^2 &= \overline{e_{47}h_0}h_4h_6^2 = 0 \quad (\text{since } h_4h_6^2 = 0). \end{aligned}$$

So $\alpha h_6^2 = 0$. Since θ_6 is detected by h_6^2 in the ASS for π_*^S , we see $AF(\tilde{q}\hat{\theta}_5\theta_6) \geq 5$.

(2) If $AF(\widehat{q}\theta_5) \geq 3$ then clearly $AF(\widehat{q}\theta_5\theta_6) \geq 3 + 2 = 5$.

This proves the Lemma 5.3. \square

The scheme to prove Proposition 2.46 is as follows. Since $\widehat{\theta}_5, \theta_6$ are detected by $\widehat{h}_5 h_5 \in \text{Ext}_{\mathcal{A}}^{1,63}(P^{62})$ and $h_6^2 \in \text{Ext}_{\mathcal{A}}^{2,128}$ in the ASS for $\pi_*^S(P^{62})$ and for π_*^S respectively, and since $h_5 h_6 = 0$ in $\text{Ext}_{\mathcal{A}}^{*,*}$, it follows that $AF(\widehat{\theta}_5\theta_6) \geq 4$. If $AF(\widehat{\theta}_5\theta_6) = 4$ then $\widehat{\theta}_5\theta_6$ must be detected by a non-zero class $\tau \in \text{Ext}_{\mathcal{A}}^{4,192}(P^{62})$. Consider the induced homomorphism $\text{Ext}_{\mathcal{A}}^{*,*}(P^{62}) \xrightarrow{\widehat{q}_*} \text{Ext}_{\mathcal{A}}^{*,*}(\widetilde{P}^{62})$. By Lemma 5.3, $AF(\widehat{q}\theta_5\theta_6) = 5$. Thus either $q_*(\tau) = 0$ in $\text{Ext}_{\mathcal{A}}^{4,192}(\widetilde{P}^{62})$ or $q_*(\tau)$ is a boundary in the ASS for $\pi_*^S(\widetilde{P}^{62})$. To show that neither of these two cases is possible we need the following Lemma 5.4. To state it, we recall the following.

$$(3.5) \quad (4) \quad \text{Ext}_{\mathcal{A}}^{2,191}(\widetilde{P}^{62}) = \mathbb{Z}/2(\overline{e_{31}h_5h_7}) \oplus \mathbb{Z}/2(\overline{e_{47}h_4h_7}).$$

$$(3.5) \quad (6) \quad \text{Ext}_{\mathcal{A}}^{4,192}(\widetilde{P}^{62}) = \mathbb{Z}/2(\widetilde{q}_*(\widehat{h}_4h_0c_4)) \oplus \mathbb{Z}/2(\widetilde{q}_*(\widehat{D}_3h_7)) \oplus \mathbb{Z}/2(\widetilde{q}_*(\widehat{g}_4h_0)) \oplus \mathbb{Z}/2(\widetilde{q}_*(\overline{e_{62}D_3(1)})) \oplus \mathbb{Z}/2(\overline{e_{31}h_4^2h_0h_7}).$$

Lemma 5.4. *In the ASS for $\pi_*^S(\widetilde{P}^{62})$,*

$$(1) \quad d_2(\overline{e_{31}h_5h_7}) = \widetilde{q}_*(\widehat{D}_3h_7) + \overline{e_{31}h_0h_4^2h_7},$$

$$(2) \quad d_2(\overline{e_{47}h_4h_7}) = \widetilde{q}_*(\widehat{D}_3h_7) + \overline{e_{31}h_0h_4^2h_7}.$$

The proof of this Lemma 5.4 is rather lengthy. We will postpone this proof to the end of this section. Now we prove Proposition 2.46.

Proof of Proposition 2.46. We want to show the composite $S^{188} \xrightarrow{\theta_6} S^{62} \xrightarrow{\widehat{\theta}_5} P^{62}$ has $AF \geq 5$. We already know that $AF(\widehat{\theta}_5\theta_6) \geq 4$ (since $\widehat{h}_5h_5h_6^2 = 0$). Recall

$$(3.3) \quad (i) \quad (5) \quad \text{Ext}_{\mathcal{A}}^{4,192}(P^{62}) = \mathbb{Z}/2(\widehat{h}_4h_0c_4) \oplus \mathbb{Z}/2(\widehat{D}_3h_7) \oplus \mathbb{Z}/2(\widehat{g}_4h_0) \oplus \mathbb{Z}/2(\overline{e_{62}D_3(1)}).$$

Suppose $AF(\widehat{\theta}_5\theta_6) = 4$. Then

$$(5.5) \quad \widehat{\theta}_5\theta_6 \text{ is detected by } \tau = \varepsilon_1\widehat{h}_4h_0c_4 + \varepsilon_2\widehat{D}_3h_7 + \varepsilon_3\widehat{g}_4h_0 + \varepsilon_4\widehat{h}_7D_3, \text{ where } \varepsilon_i = 0 \text{ or } 1 \text{ for } 1 \leq i \leq 4, \text{ and } (\varepsilon_1, \varepsilon_2, \varepsilon_3, \varepsilon_4) \neq (0, 0, 0, 0).$$

We are going to show that this would lead to a contradiction. Recall

$$(3.5) \quad (6) \quad \text{Ext}_{\mathcal{A}}^{4,192}(\widetilde{P}^{62}) = \mathbb{Z}/2(\widetilde{q}_*(\widehat{h}_4h_0c_4)) \oplus \mathbb{Z}/2(\widetilde{q}_*(\widehat{D}_3h_7)) \oplus \mathbb{Z}/2(\widetilde{q}_*(\widehat{g}_4h_0)) \oplus \mathbb{Z}/2(\widetilde{q}_*(\widehat{h}_7D_3)) \oplus \mathbb{Z}/2(\overline{e_{31}h_4^2h_0h_7})$$

Let $P^{62} \xrightarrow{\widehat{q}} \widetilde{P}^{62}$ be as in Lemma 5.3. Then from (5.5) we have:

$$(5.6) \quad \widetilde{q}_*(\tau) = \varepsilon_1\widetilde{q}_*(\widehat{h}_4h_0c_4) + \varepsilon_2\widetilde{q}_*(\widehat{D}_3h_7) + \varepsilon_3\widetilde{q}_*(\widehat{g}_4h_0) + \varepsilon_4\widetilde{q}_*(\widehat{h}_7D_3), \text{ where } \varepsilon_i = 0 \text{ or } 1 \text{ for } 1 \leq i \leq 4, \text{ and } (\varepsilon_1, \varepsilon_2, \varepsilon_3, \varepsilon_4) \neq (0, 0, 0, 0). \widetilde{q}_*(\tau) \text{ is non-zero in } \text{Ext}_{\mathcal{A}}^{4,192}(\widetilde{P}^{62}) \text{ by (3.5) (6) and the fact that } (\varepsilon_1, \varepsilon_2, \varepsilon_3, \varepsilon_4) \neq (0, 0, 0, 0).$$

By Lemma 5.3, $\widetilde{q}_*(\tau)$ has to be a boundary (if $\widetilde{q}_*(\tau)$ is not a boundary then $AF(\widehat{q}\theta_5\theta_6) = 4$). So $d_r(\beta) = \widetilde{q}_*(\tau)$ for some $\beta \in \text{Ext}_{\mathcal{A}}^{r-4,185+r}(\widetilde{P}^{62})$ with $2 \leq r \leq 4$. Recall

$$(3.5) \quad (8) \quad \text{Ext}_{\mathcal{A}}^{0,189}(\widetilde{P}^{62}) = 0.$$

$$(3.5) (9) \quad \text{Ext}_{\mathcal{A}}^{1,190}(\tilde{P}^{62}) = 0.$$

$$(3.5) (10) \quad \text{Ext}_{\mathcal{A}}^{2,191}(\tilde{P}^{62}) = \mathbb{Z}/2(\overline{\tilde{e}_{31}h_5h_7}) \oplus \mathbb{Z}/2(\overline{\tilde{e}_{47}h_4h_7}).$$

Thus the only possibility is $d_2(\delta_1\overline{\tilde{e}_{31}h_5h_7} + \delta_2\overline{\tilde{e}_{47}h_4h_7}) = \tilde{q}_*(\tau)$ where $(\delta_1, \delta_2) \neq (0, 0)$. By Lemma 5.4,

$$\begin{aligned} d_2(\delta_1\overline{\tilde{e}_{31}h_5h_7} + \delta_2\overline{\tilde{e}_{47}h_4h_7}) &= \delta_1(\tilde{q}_*(\widehat{D}_3h_7) + \overline{\tilde{e}_{31}h_0h_4^2h_7}) + \delta_2(\tilde{q}_*(\widehat{D}_3h_7) + \overline{\tilde{e}_{31}h_0h_4^2h_7}) \\ &= 0 \quad \text{or} \quad \tilde{q}_*(\widehat{D}_3h_7) + \overline{\tilde{e}_{31}h_0h_4^2h_7}. \end{aligned}$$

By (3.5) (6) and (5.6) above, we see $\tilde{q}_*(\tau) = \varepsilon_1\tilde{q}_*(\widehat{h}_4h_0c_4) + \varepsilon_2\tilde{q}_*(\widehat{D}_3h_7) + \varepsilon_3\tilde{q}_*(\widehat{g}_4h_0) + \varepsilon_4\tilde{q}_*(\widehat{h}_7D_3) \neq \tilde{q}_*(\widehat{D}_3h_7) + \overline{\tilde{e}_{31}h_0h_4^2h_7}$. This contradiction proves that $AF(\widehat{\theta}_5\theta_6) \geq 5$. This concludes the proof of Proposition 2.46. \square

We proceed to prove Proposition 1.6. We want to prove for this proposition that in the ASS for spheres, $d_r(h_6^3) = 0$ for $r = 2$ and 3 . That $d_2(h_6^3) = 0$ is quite easy as $d_2(h_6^3 = h_6h_6^2) = h_0h_5^2h_6^2 = 0$. We have $d_3(h_6^3) \in E_3^{6,194}$ which is a subquotient of $E_2^{6,194} = \text{Ext}_{\mathcal{A}}^{6,194}$. Recall

$$(3.1) (8) \quad \text{Ext}_{\mathcal{A}}^{6,194} = \mathbb{Z}/2(h_0^2g_4) \oplus \mathbb{Z}/2(h_5^2D_3(1)).$$

So $d_3(h_6^3) = \varepsilon_1h_0^2g_4 + \varepsilon_2h_5^2D_3(1)$ with $\varepsilon_i = 0$ or 1 , $1 \leq i \leq 2$. We have to show $\varepsilon_1 = \varepsilon_2 = 0$. For this purpose we first prove the following case that says that $\varepsilon_1 = 0$ and $\varepsilon_2 = 1$ can not happen.

Lemma 5.7. *In the ASS for π_*^S , $d_3(h_6^3) \neq h_5^2D_3(1)$.*

Proof. Recall that the composite of stable maps $S^{63} \xrightarrow{\theta_5} S^1 \xrightarrow{2\iota} S^1$ is zero. So for the following cofibration

$$(5.8) \quad S^0 \xrightarrow{i_1} S^0 \cup_{2\iota} e^1 \xrightarrow{q_1} S^1,$$

there is a coextension

$$(5.9) \quad S^{63} \xrightarrow{\bar{\theta}_5} S^0 \cup_{2\iota} S^1.$$

We have $h_6^3 \in \text{Ext}_{\mathcal{A}}^{3,192}$, and $(i_1)_*(h_6^3) \in \text{Ext}_{\mathcal{A}}^{3,192}(S^0 \cup_{2\iota} e^1)$ for the $S^0 \xrightarrow{i_1} S^0 \cup_{2\iota} e^1$ in (5.8). We claim the following.

- (5.10) (1) $(i_1)_*(h_6^3) \neq 0$ in $\text{Ext}_{\mathcal{A}}^{3,192}(S^0 \cup_{2\iota} e^1)$.
- (2) The composite $S^{189} \xrightarrow{\theta_6} S^{63} \xrightarrow{\bar{\theta}_5} S^0 \cup_{2\iota} e^1$ is detected by $(i_1)_*(h_6^3)$ in the ASS for $\pi_*^S(S^0 \cup_{2\iota} e^1)$.
- (3) $\text{Ext}_{\mathcal{A}}^{4,193}(S^0 \cup_{2\iota} e^1) = 0$.

Assuming (5.10), we show $d_3(h_6^3) \neq h_5^2D_3(1)$ in the ASS for π_*^S as follows. If $d_3(h_6^3) = h_5^2D_3(1)$ in the ASS for π_*^S , then since $(i_1)_*(h_6^3) \neq 0$ is an infinite cycle in the ASS for $\pi_*^S(S^0 \cup_{2\iota} e^1)$ (by (5.10)), we must have either of the following two cases.

- (5.11) (1) $(i_1)_*(h_5^2 D_3(1)) = 0$ in $\text{Ext}_{\mathcal{A}}^{6,194}(S^0 \cup_{2\iota} e^1)$, or
(2) $(i_1)_*(h_5^2 D_3(1)) \neq 0$ in $\text{Ext}_{\mathcal{A}}^{6,194}(S^0 \cup_{2\iota} e^1)$ and $d_2(\alpha) = (i_1)_*(h_5^2 D_3(1))$
for some $\alpha \in \text{Ext}_{\mathcal{A}}^{4,193}(S^0 \cup_{2\iota} e^1)$ in the ASS for $\pi_*^S(S^0 \cup_{2\iota} e^1)$.

By (5.10) (3), (5.11) (2) is impossible. To show that (5.11) (1) is impossible, recall from (4.4) that since (5.8) induces a short exact sequence $0 \rightarrow H_*(S^0) \xrightarrow{(i_1)_*} H_*(S^0 \cup_{2\iota} e^1) \xrightarrow{(q_1)_*} H_*(S^1) \rightarrow 0$, there is a long exact sequence

$$(5.12) \quad \cdots \rightarrow \text{Ext}_{\mathcal{A}}^{s-1, t-1} \xrightarrow{(2\iota)_*} \text{Ext}_{\mathcal{A}}^{s, t} \xrightarrow{(i_1)_*} \text{Ext}_{\mathcal{A}}^{s, t}(S^0 \cup_{2\iota} e^1) \xrightarrow{(q_1)_*} \text{Ext}_{\mathcal{A}}^{s, t-1} \xrightarrow{(2\iota)_*} \text{Ext}_{\mathcal{A}}^{s+1, t} \rightarrow \cdots$$

So $(i_1)_*(h_5^2 D_3(1)) = 0$ in $\text{Ext}_{\mathcal{A}}^{6,194}(S^0 \cup_{2\iota} e^1)$ for $(s, t) = (6, 194)$ implies that there is $\beta \in \text{Ext}_{\mathcal{A}}^{5,193}$ such that $(2\iota)_*(\beta) = h_0\beta = h_5^2 D_3(1)$ in $\text{Ext}_{\mathcal{A}}^{6,194}$. From (3.1) (2) and (8), there is no such $\beta \in \text{Ext}_{\mathcal{A}}^{5,193}$. So (5.11) (1) is impossible. This shows that $d_3(h_6^3) \neq h_5^2 D_3(1)$.

To complete the proof of Lemma 5.7 it remains to show (5.10). (5.10) (1), (3) can also be proved by using the long exact sequence in (5.12). To show $(i_1)_*(h_6^3) \neq 0$ as claimed in (5.10) (1), note that $h_6^3 \in \text{Ext}_{\mathcal{A}}^{3,192}$ for $(s, t) = (3, 192)$, and by (3.1) (3), $\text{Ext}_{\mathcal{A}}^{2,191} = 0$. So, by (5.12), $(i_1)_*(h_6^3) \neq 0$ in $\text{Ext}_{\mathcal{A}}^{3,192}(S^0 \cup_{2\iota} e^1)$. This proves (5.10) (1). To show $\text{Ext}_{\mathcal{A}}^{4,193}(S^0 \cup_{2\iota} e^1) = 0$ for $(s, t) = (4, 193)$ as claimed in (5.10) (3), recall the following.

$$(3.1) (1) \quad \text{Ext}_{\mathcal{A}}^{4,192} = \mathbb{Z}/2(g_4).$$

$$(3.1) (2) \quad \text{Ext}_{\mathcal{A}}^{5,193} = \mathbb{Z}/2(h_0 g_4).$$

$$(3.1) (4) \quad \text{Ext}_{\mathcal{A}}^{s, s+189} = \mathbb{Z}/2(h_0^{s-3} h_6^3), \quad 3 \leq s \leq 4.$$

So $\text{Ext}_{\mathcal{A}}^{4,192} = \mathbb{Z}/2(g_4)$ with $(2\iota)_*(g_4) = h_0 g_4 \neq 0$ in $\text{Ext}_{\mathcal{A}}^{5,193}$, and $\text{Ext}_{\mathcal{A}}^{4,193} = \mathbb{Z}/2(h_0 h_6^3)$ with $(2\iota)_*(h_6^3) = h_0 h_6^3$. From (5.12) we thus see $\text{Ext}_{\mathcal{A}}^{4,193}(S^0 \cup_{2\iota} e^1) = 0$. This proves (5.10) (3). Finally we show that $S^{188} \xrightarrow{\theta_6} S^{62} \xrightarrow{\bar{\theta}_5} S^0 \cup_{2\iota} e^1$ is detected by $(i_1)_*(h_6^3)$ as claimed in (5.10) (2). $S^{62} \xrightarrow{\bar{\theta}_5} S^0 \cup_{2\iota} e^1$ is detected by $(i_1)_*(h_6)$ by [1]. Then since $(i_1)_*(h_6)h_6^2 = (i_1)_*(h_6^3) \neq 0$ as in (5.10) and since $\text{Ext}_{\mathcal{A}}^{s, 190+s} = 0$ for $0 \leq s \leq 1$ ((3.1) (6)) we see that $(i_1)_*(h_6^3)$ is not a boundary in the spectral sequence and that $\bar{\theta}_5 \theta_6$ is detected by $(i_1)_*(h_6^3)$ in the ASS for $\pi_*(S^0 \cup_{2\iota} e^1)$. This proves Lemma 5.7. \square

Now we can prove Proposition 1.6.

Proof of Proposition 1.6. Recall the transfer map $P \xrightarrow{t} S^0$ and the induced homomorphism $\text{Ext}_{\mathcal{A}}^{s, t}(P) \xrightarrow{t_*} \text{Ext}_{\mathcal{A}}^{s+1, t+1}$ in (2.14). There is a class $\widehat{h}_5 h_5 h_7 \in \text{Ext}_{\mathcal{A}}^{2,191}(P^{62})$ such that $t_*(\widehat{h}_5 h_5 h_7) = h_5^2 h_7 = h_6^3$. Since $d_2(h_6^3) = 0$ in the ASS for π_*^S , and $d_2(\widehat{h}_5 h_5 h_7) = \widehat{h}_5 h_4^2 h_0 h_7 = 0$ in the ASS for $\pi_*^S(P)$, we can consider the following diagram.

$$\begin{array}{ccc} \text{Ext}_{\mathcal{A}}^{2,191}(P^{62}) & \xrightarrow{t_*} & \text{Ext}_{\mathcal{A}}^{3,192} \\ \downarrow d_3 & & \downarrow d_3 \\ \text{Ext}_{\mathcal{A}}^{5,193}(P^{62}) & \xrightarrow{t_*} & \text{Ext}_{\mathcal{A}}^{6,194} \end{array}$$

Suppose $d_3(h_6^3) = \alpha \neq 0$ in $\text{Ext}_{\mathcal{A}}^{6,194}$ in the ASS for π_*^S . We show this would lead to a contradiction. $d_3(h_6^3) = \alpha$ in the ASS for π_*^S implies $d_3(\widehat{h}_5 h_5 h_7) = \alpha' \neq 0$ in $\text{Ext}_{\mathcal{A}}^{5,193}(P^{62})$ with $t_*(\alpha') = \alpha$ in the ASS for $\pi_*^S(P^{62})$. We recall the following.

$$(3.3) \text{ (i) (9)} \quad \text{Ext}_{\mathcal{A}}^{5,193}(P^{62}) = \mathbb{Z}/2(\widehat{h}_5 h_5 D_3(1)) \oplus \mathbb{Z}/2(\widehat{h}_4 h_0^2 c_4) \oplus \mathbb{Z}/2(\widehat{h}_4 h_1 f_3) \oplus \mathbb{Z}/2(\gamma_{61} h_7) \oplus \mathbb{Z}/2(\widehat{g}_4 h_0^2).$$

Therefore

$$(5.13) \quad d_3(\widehat{h}_5 h_5 h_7) = \alpha' = \varepsilon_1 \widehat{h}_5 h_5 D_3(1) + \varepsilon_2 \widehat{h}_4 h_0^2 c_4 + \varepsilon_3 \widehat{h}_4 h_1 f_3 + \varepsilon_4 \gamma_{61} h_7 + \varepsilon_5 \widehat{g}_4 h_0^2, \quad \varepsilon_i = 0 \text{ or } 1, 1 \leq i \leq 5, \\ \text{with } (\varepsilon_1, \varepsilon_2, \varepsilon_3, \varepsilon_4, \varepsilon_5) \neq (0, 0, 0, 0, 0).$$

Recall the complex \widetilde{P}^{62} and the map $P^{62} \xrightarrow{\widetilde{q}} \widetilde{P}^{62}$. Consider the following diagram.

$$\begin{array}{ccc} \text{Ext}_{\mathcal{A}}^{2,191}(P^{62}) & \xrightarrow{\widetilde{q}_*} & \text{Ext}_{\mathcal{A}}^{2,191}(\widetilde{P}^{62}) \\ \downarrow d_3 & & \downarrow d_3 \\ \text{Ext}_{\mathcal{A}}^{5,193}(P^{62}) & \xrightarrow{\widetilde{q}_*} & \text{Ext}_{\mathcal{A}}^{5,193}(\widetilde{P}^{62}) \end{array}$$

We recall the following.

$$(3.5) \text{ (7) (a)} \quad \widetilde{q}_*(\widehat{h}_5 h_5 D_3(1)) = 0 \text{ in } \text{Ext}_{\mathcal{A}}^{5,193}(\widetilde{P}^{62}).$$

$$(3.5) \text{ (7) (b)} \quad \widetilde{q}_*(\widehat{h}_4 h_0^2 c_4), \widetilde{q}_*(\widehat{h}_4 h_1 f_3), \widetilde{q}_*(\gamma_{61} h_7), \widetilde{q}_*(\widehat{g}_4 h_0^2), \overline{e_{31} h_4^2 h_0^2 h_7} \text{ are linearly independent in } \text{Ext}_{\mathcal{A}}^{5,193}(\widetilde{P}^{62}).$$

Since $\widetilde{q}_*(\widehat{h}_5 h_5 h_7) = 0$, we have $\widetilde{q}_*(\alpha') = \widetilde{q}_*(d_3(\widehat{h}_5 h_5 h_7)) = d_3(\widetilde{q}_*(\widehat{h}_5 h_5 h_7)) = 0$ in the ASS for $\pi_*^S(\widetilde{P}^{62})$. There are two cases for this to be possible.

- (1) If $\widetilde{q}_*(\alpha') = 0$ in $\text{Ext}_{\mathcal{A}}^{5,193}(\widetilde{P}^{62})$, from (3.3) (i) (9), (5.13) and (3.5) (7) (a), (b) we see that $\alpha' = \widehat{h}_5 h_5 D_3(1)$, that is, $\varepsilon_1 = 1$, $\varepsilon_i = 0$ for $2 \leq i \leq 5$. Then $d_3(h_6^3) = t_*(\alpha' = \widehat{h}_5 h_5 D_3(1)) = h_5^2 D_3(1)$ in the ASS for π_*^S . By Lemma 5.7 this is impossible.

- (2) If $\widetilde{q}_*(\alpha') \neq 0$ in $\text{Ext}_{\mathcal{A}}^{5,193}(\widetilde{P}^{62})$, then $\widetilde{q}_*(\alpha')$ must be a boundary in the ASS for $\pi_*^S(\widetilde{P}^{62})$. Let $\alpha'' = \widetilde{q}_*(\alpha')$, then there is $\beta \in \text{Ext}_{\mathcal{A}}^{3,192}(\widetilde{P}^{62})$ with $d_2(\beta) = \alpha''$. By

$$(3.5) \text{ (11)} \quad \text{Ext}_{\mathcal{A}}^{3,192}(\widetilde{P}^{62}) = \mathbb{Z}/2(\overline{e_{31} h_5 h_0 h_7}) \oplus \mathbb{Z}/2(\overline{e_{47} h_4 h_7 h_0}) \oplus \mathbb{Z}/2(\widetilde{q}_*(\overline{e_{62} h_1 h_6^2}))$$

we have $d_2(\widetilde{q}_*(\overline{e_{62} h_1 h_6^2})) = 0$ ($\overline{e_{62} h_1 h_6^2} \in \text{Ext}_{\mathcal{A}}^{3,192}(P^{62})$ is a pullback of $\widehat{h}_6 h_0 h_6^2 \in \text{Ext}_{\mathcal{A}}^{3,192}(P)$, and $d_2(\widehat{h}_6) = \widehat{h}_5 h_0 h_5$ and $h_5 h_6 = 0$ in the ASS for $\pi_*^S(P)$). And by Lemma 5.4,

$$d_2(\overline{e_{31} h_5 h_7}) = \widetilde{q}_*(\widehat{D}_3 h_7) + \overline{e_{31} h_0 h_4^2 h_7}, \\ d_2(\overline{e_{47} h_4 h_7}) = \widetilde{q}_*(\widehat{D}_3 h_7) + \overline{e_{31} h_0 h_4^2 h_7}.$$

Therefore

$$\begin{aligned} d_2(\overline{\tilde{e}_{31}h_5}h_7h_0) &= \tilde{q}_*(\widehat{D}_3h_7)h_0 + \overline{\tilde{e}_{31}h_0h_4^2}h_0h_7 = \overline{\tilde{e}_{31}h_0h_4^2}h_0h_7, \\ d_2(\overline{\tilde{e}_{47}h_4}h_7h_0) &= \tilde{q}_*(\widehat{D}_3h_7)h_0 + \overline{\tilde{e}_{31}h_0h_4^2}h_0h_7 = \overline{\tilde{e}_{31}h_0h_4^2}h_0h_7. \end{aligned}$$

It follows that the only non-trivial boundary in the ASS for $\pi_*^S(\tilde{P}^{62})$ in $\text{Ext}_{\mathcal{A}}^{5,193}(\tilde{P}^{62})$ is $\overline{\tilde{e}_{31}h_0h_4^2}h_0h_7$. Since $\tilde{q}_*(\alpha') = \varepsilon_2\tilde{q}_*(\widehat{h}_4h_0^2c_4) + \varepsilon_3\tilde{q}_*(\widehat{h}_4h_1f_3) + \varepsilon_4\tilde{q}_*(\gamma_{61}h_7) + \varepsilon_5\tilde{q}_*(\widehat{g}_4h_0^2) \neq \overline{\tilde{e}_{31}h_0h_4^2}h_0h_7$ for any $\alpha' \in \text{Ext}_{\mathcal{A}}^{5,193}(P^{62})$ (by (3.5) (7) (b)), it follows that $\tilde{q}_*(\alpha')$ is not a boundary in the ASS for $\pi_*^S(\tilde{P}^{62})$.

This completes the proof of Proposition 1.6. \square

Finally we prove Proposition 1.7.

Proof of Proposition 1.7. In the ASS for π_*^S we have $d_2(h_1) = 0$, so $d_2(h_1g_4) = h_1d_2(g_4) \in \text{Ext}_{\mathcal{A}}^{7,195}$. There are two cases to consider.

- (1) If $d_2(g_4) = 0$ in $\text{Ext}_{\mathcal{A}}^{6,193}$ then $d_2(h_1g_4) = 0$ and the proof is done.
- (2) If $d_2(g_4) \neq 0$ in $\text{Ext}_{\mathcal{A}}^{6,193}$, then since $\text{Ext}_{\mathcal{A}}^{6,193} = \mathbb{Z}/2(h_5V_0)$ (by (3.1) (7)) we see $d_2(g_4) = h_5V_0$. And from (3.1) (9), $h_1h_5V_0 \neq h_0^3g_4$. So $d_2(h_1g_4) = h_1d_2(g_4) = h_1h_5V_0 \neq h_0^3g_4$ in $\text{Ext}_{\mathcal{A}}^{7,195}$.

This proves Proposition 1.7. \square

The remainder of this section is devoted to proving Lemma 5.4. First we prove

$$(5.4) \quad (1) \quad \text{In the ASS for } \pi_*^S(\tilde{P}^{62}), \quad d_2(\overline{\tilde{e}_{31}h_5}h_7) = \tilde{q}_*(\widehat{D}_3h_7) + \overline{\tilde{e}_{31}h_0h_4^2}h_7.$$

Since $d_2(h_7) = h_0h_6^2$ in the ASS for π_*^S (by (5.1)), we have the following differential in the ASS for $\pi_*^S(\tilde{P}^{62})$.

$$(5.14) \quad \begin{aligned} d_2(\overline{\tilde{e}_{31}h_5}h_7) &= d_2(\overline{\tilde{e}_{31}h_5})h_7 + \overline{\tilde{e}_{31}h_5}d_2(h_7) \\ &= d_2(\overline{\tilde{e}_{31}h_5})h_7 + \overline{\tilde{e}_{31}h_5}h_0h_6^2 \\ &= d_2(\overline{\tilde{e}_{31}h_5})h_7. \end{aligned}$$

Here $\overline{\tilde{e}_{31}h_5}h_0h_6^2 = 0$ in $\text{Ext}_{\mathcal{A}}^{4,192}(\tilde{P}^{62})$ because, from (3.52) (1) we have $\overline{\tilde{e}_{31}h_5} = \{\tilde{e}_{31}\lambda_{31}\} \in \text{Ext}_{\mathcal{A}}^{1,63}(\tilde{P}^{62})$, and we see that $\overline{\tilde{e}_{31}h_5}h_0h_6^2 = \overline{\tilde{e}_{31}h_5}h_6h_0h_6 = \{\tilde{e}_{31}\lambda_{31}\lambda_{63}\lambda_0\lambda_{63}\} = 0$. Thus to compute $d_2(\overline{\tilde{e}_{31}h_5}h_7)$ we need to compute $d_2(\overline{\tilde{e}_{31}h_5})$ in the ASS for $\pi_*^S(\tilde{P}^{62})$. Consider the following cofibration.

$$(5.15) \quad \tilde{P}^{62} \xrightarrow{q'_{17}} \tilde{P}_{17}^{62} \xrightarrow{\Delta} \Sigma P^{16}.$$

Recall

$$(3.5) \quad (4) \quad \text{Ext}_{\mathcal{A}}^{1,63}(\tilde{P}^{62}) = \mathbb{Z}/2(\overline{\tilde{e}_{31}h_5}) \oplus \mathbb{Z}/2(\overline{\tilde{e}_{47}h_4}).$$

$$(3.6) \quad (1) \quad \text{Ext}_{\mathcal{A}}^{1,63}(\tilde{P}_{17}^{62}) = \mathbb{Z}/2(\overline{\tilde{e}_{31}h_5}) \oplus \mathbb{Z}/2(\overline{\tilde{e}_{47}h_4}).$$

$\text{Ext}_{\mathcal{A}}^{1,63}(\tilde{P}^{62}) \xrightarrow{(q'_{17})_*} \text{Ext}_{\mathcal{A}}^{1,63}(\tilde{P}_{17}^{62})$ is given by $(q'_{17})_*(\overline{\tilde{e}_{31}h_5}) = \overline{\tilde{e}_{31}h_5}$ and $(q'_{17})_*(\overline{\tilde{e}_{47}h_4}) = \overline{\tilde{e}_{47}h_4}$.

$$(3.5) \quad (3) \quad \text{Ext}_{\mathcal{A}}^{3,64}(\tilde{P}^{62}) = \mathbb{Z}/2(\overline{\tilde{e}_{31}h_4^2h_0}) \oplus \mathbb{Z}/2(\tilde{q}_*(\widehat{D}_3))$$

$$(3.6) \quad (3) \quad \text{Ext}_{\mathcal{A}}^{3,64}(\tilde{P}_{17}^{62}) = \mathbb{Z}/2(\overline{\tilde{e}_{31}h_0h_4^2}) \oplus \mathbb{Z}/2((\tilde{q}_{17})_*(\overline{\tilde{e}_{23}h_0h_3h_5})).$$

$$\text{Ext}_{\mathcal{A}}^{3,64}(\tilde{P}^{62}) \xrightarrow{(q'_{17})_*} \text{Ext}_{\mathcal{A}}^{3,64}(\tilde{P}_{17}^{62}) \text{ is given by } (q'_{17})_*(\overline{\tilde{e}_{31}h_4^2h_0}) = \overline{\tilde{e}_{31}h_0h_4^2},$$

$$\text{and } (q'_{17})_*(\tilde{q}_*(\widehat{D}_3)) = (\tilde{q}_{17})_*(\overline{\tilde{e}_{23}h_0h_3h_5}).$$

Note that $\text{Ext}_{\mathcal{A}}^{1,63}(\tilde{P}^{62}) \xrightarrow{(q'_{17})_*} \text{Ext}_{\mathcal{A}}^{1,63}(\tilde{P}_{17}^{62})$ and $\text{Ext}_{\mathcal{A}}^{3,64}(\tilde{P}^{62}) \xrightarrow{(q'_{17})_*} \text{Ext}_{\mathcal{A}}^{3,64}(\tilde{P}_{17}^{62})$ are both isomorphisms. By naturality of the differentials in the Adams spectral sequence, we have the following equation in the ASS for $\pi_*^S(\tilde{P}_{17}^{62})$.

$$(5.16) \quad (q'_{17})_*(d_2(\overline{\tilde{e}_{31}h_5})) = d_2((q'_{17})_*(\overline{\tilde{e}_{31}h_5}))$$

$$= d_2(\overline{\tilde{e}_{31}h_5}) \quad (\text{by (3.6) (1)})$$

$$= d_2(\overline{\tilde{e}_{31}})h_5 + \overline{\tilde{e}_{31}}d_2(h_5)$$

$$= d_2(\overline{\tilde{e}_{31}})h_5 + \overline{\tilde{e}_{31}}h_0h_4^2 \quad (\text{by (3.6) (3)})$$

Thus to compute $(q'_{17})_*(d_2(\overline{\tilde{e}_{31}h_5}))$ we need to compute $d_2(\overline{\tilde{e}_{31}})$ in the ASS for $\pi_*^S(\tilde{P}_{17}^{62})$. Recall

$$(3.6) \quad (2) \quad \text{Ext}_{\mathcal{A}}^{2,32}(\tilde{P}_{17}^{62}) = \mathbb{Z}/2((\tilde{q}_{17})_*(\overline{\tilde{e}_{23}h_0h_3})).$$

We claim

$$(5.17) \quad d_2(\overline{\tilde{e}_{31}}) \neq 0 \text{ in the ASS for } \pi_*^S(\tilde{P}_{17}^{62}).$$

Assuming (5.17), from (3.6) (2) we must have $d_2(\overline{\tilde{e}_{31}}) = (\tilde{q}_{17})_*(\overline{\tilde{e}_{23}h_0h_3})$. Then we have

$$(5.18) \quad (q'_{17})_*(d_2(\overline{\tilde{e}_{31}h_5})) = d_2(\overline{\tilde{e}_{31}})h_5 + \overline{\tilde{e}_{31}}h_0h_4^2 \quad (\text{by (5.16)})$$

$$= (\tilde{q}_{17})_*(\overline{\tilde{e}_{23}h_0h_3})h_5 + \overline{\tilde{e}_{31}}h_0h_4^2 \quad (\text{by } d_2(\overline{\tilde{e}_{31}}) = (\tilde{q}_{17})_*(\overline{\tilde{e}_{23}h_0h_3}))$$

$$= (\tilde{q}_{17})_*(\overline{\tilde{e}_{23}h_0h_3h_5}) + \overline{\tilde{e}_{31}}h_0h_4^2$$

$$= (q'_{17})_*(\tilde{q}_*(\widehat{D}_3) + \overline{\tilde{e}_{31}h_4^2h_0}) \quad (\text{by (3.6) (3)})$$

And since $\text{Ext}_{\mathcal{A}}^{3,64}(\tilde{P}^{62}) \xrightarrow{(q'_{17})_*} \text{Ext}_{\mathcal{A}}^{3,64}(\tilde{P}_{17}^{62})$ is an isomorphism, we have

$$(5.19) \quad d_2(\overline{\tilde{e}_{31}h_5}) = \tilde{q}_*(\widehat{D}_3) + \overline{\tilde{e}_{31}h_4^2h_0}.$$

We then have the following (5.20).

$$(5.20) \quad d_2(\overline{\tilde{e}_{31}h_5h_7}) = d_2(\overline{\tilde{e}_{31}h_5})h_7 \quad (\text{by (5.14)})$$

$$= (\tilde{q}_*(\widehat{D}_3) + \overline{\tilde{e}_{31}h_4^2h_0})h_7 \quad (\text{by (5.19)})$$

$$= \tilde{q}_*(\widehat{D}_3h_7) + \overline{\tilde{e}_{31}h_4^2h_0h_7}.$$

This proves (5.4) (1) modulo (5.17). The proof of (5.17) is given as follows.

We use the $\tilde{P}_{17}^{62} \xrightarrow{\Delta} \Sigma P^{16}$ map in (5.15). Recall from (3.51) that in the bigraded differential module $H_*(\tilde{P}^{62}) \otimes \Lambda$ we have the differential $\delta(\tilde{e}_{31}) = e_{15}\lambda_{15} \in H_*(\tilde{P}^{62}) \otimes \Lambda$ (put $\lambda_I = 1$ in (3.51)). We have $\text{Ext}_{\mathcal{A}}^{1,32}(\Sigma P^{16}) \cong \text{Ext}_{\mathcal{A}}^{1,31}(P^{16})$, and recall

$$(3.4) (6) \quad \text{Ext}_{\mathcal{A}}^{1,31}(P^{16}) = \mathbb{Z}/2(\overline{e_{15}}h_4).$$

So we have

$$(5.21) \quad \Delta_*(\overline{e_{31}} = \{\tilde{e}_{31}\}) = \{\delta(\tilde{e}_{31})\} = \{e_{15}\lambda_{15}\} = \overline{e_{15}}h_4 \in \text{Ext}_{\mathcal{A}}^{1,31}(P^{16}) \cong \text{Ext}_{\mathcal{A}}^{1,32}(\Sigma P^{16}).$$

Recall

$$(3.4) (7) \quad \text{Ext}_{\mathcal{A}}^{2,16}(P^{16}) = \mathbb{Z}/2(\overline{e_7}h_0h_3).$$

$$(3.4) (8) \quad \text{Ext}_{\mathcal{A}}^{3,32}(P^{16}) = \mathbb{Z}/2(\overline{e_{15}}h_0h_3^2).$$

Since $\overline{e_{15}} \in \text{Ext}_{\mathcal{A}}^{0,15}(P^{16})$ and $\overline{e_7} \in \text{Ext}_{\mathcal{A}}^{0,7}(P^{16})$ are pullbacks of $\widehat{h}_4 \in \text{Ext}_{\mathcal{A}}^{0,15}(P)$ and $\widehat{h}_3 \in \text{Ext}_{\mathcal{A}}^{0,7}(P)$, by naturality of the differentials in the Adams spectral sequence, we have the following equation in the ASS for $\pi_*^S(P^{16})$.

$$(5.22) \quad \begin{aligned} \Delta_*(d_2(\overline{e_{31}})) &= d_2(\Delta_*(\overline{e_{31}})) \\ &= d_2(\overline{e_{15}}h_4) && \text{(by (5.21))} \\ &= d_2(\overline{e_{15}})h_4 + \overline{e_{15}}d_2(h_4) \\ &= \overline{e_7}h_0h_3h_4 + \overline{e_{15}}h_0h_3^2 && \text{(by (3.4) (7) and (5.1))} \\ &= \overline{e_{15}}h_0h_3^2 \neq 0 && \text{(by (3.4) (8)).} \end{aligned}$$

It follows that we must have $d_2(\overline{e_{31}}) \neq 0$ in the ASS for $\pi_*^S(\widetilde{P}_{17}^{62})$. This proves (5.17).

This completes the proof of Lemma 5.4 (1).

Next we prove

$$(5.4) (2) \quad \text{In the ASS for } \pi_*^S(\widetilde{P}^{62}), d_2(\overline{e_{47}}h_4h_7) = \tilde{q}_*(\widehat{D}_3h_7) + \overline{e_{31}}h_0h_4^2h_7.$$

Since $d_2(h_4) = h_0h_3^2$ and $d_2(h_7) = h_0h_6^2$ (by (5.1)), we have the following differential in the ASS for $\pi_*^S(\widetilde{P}^{62})$.

$$(5.23) \quad \begin{aligned} d_2(\overline{e_{47}}h_4h_7) &= d_2(\overline{e_{47}})h_4h_7 + \overline{e_{47}}d_2(h_4)h_7 + \overline{e_{47}}h_4d_2(h_7) \\ &= d_2(\overline{e_{47}})h_4h_7 + \overline{e_{47}}h_0h_3^2h_7 + \overline{e_{47}}h_4h_0h_6^2 \\ &= d_2(\overline{e_{47}})h_4h_7. \end{aligned}$$

Here $\overline{e_{47}}h_4h_0h_6^2 = \overline{e_{47}}h_0h_4h_6^2 = 0$ in $\text{Ext}_{\mathcal{A}}^{4,192}(\widetilde{P}^{62})$ because $h_4h_6^2 = 0$, and $\overline{e_{47}}h_0h_3^2h_7 = 0$ in $\text{Ext}_{\mathcal{A}}^{4,192}(\widetilde{P}^{62})$ because $\overline{e_{47}}h_0h_3^2h_7 = \overline{e_{47}}h_3^2h_0h_7 = \{e_{47}\lambda_7^2\lambda_0\lambda_{127}\}$ and $\delta(e_{55}\lambda_7\lambda_0\lambda_{127}) = e_{47}\lambda_7^2\lambda_0\lambda_{127}$ in $H_*(\widetilde{P}^{62}) \otimes \Lambda$. So to compute $d_2(\overline{e_{47}}h_4h_7)$ we need to compute $d_2(\overline{e_{47}})$ in the ASS for $\pi_*^S(\widetilde{P}^{62})$. Consider the following diagram where the horizontal maps are cofibrations.

$$(5.24) \quad \begin{array}{ccccccc} P^{32} & \xrightarrow{i} & P^{62} & \xrightarrow{q_{33}} & P_{33}^{62} & \xrightarrow{\Delta} & \Sigma P^{32} \\ \downarrow & & \downarrow \tilde{q} & & \parallel & & \\ \tilde{P}^{32} & \xrightarrow{\tilde{i}} & \tilde{P}^{62} & \xrightarrow{q'_{33}} & P_{33}^{62} & & \end{array}$$

We also recall the following.

$$(3.4) (4) \quad \text{Ext}_{\mathcal{A}}^{0,47}(P_{33}^{62}) = \mathbb{Z}/2(\overline{e_{47}}).$$

$$(3.4) (5) \quad \text{Ext}_{\mathcal{A}}^{2,48}(P_{33}^{62}) = \mathbb{Z}/2(\overline{e_{39}h_0h_3}).$$

$$(3.5) (1) \quad \text{Ext}_{\mathcal{A}}^{0,47}(\tilde{P}^{62}) = \mathbb{Z}/2(\overline{e_{47}}).$$

$$(3.5) (2) \quad \text{Ext}_{\mathcal{A}}^{2,48}(\tilde{P}^{62}) = \mathbb{Z}/2(\overline{e_{39}h_3h_0}) \oplus \mathbb{Z}/2(\tilde{q}_*(\widehat{h_4h_0h_5})).$$

$$(3.5) (3) \quad \overline{e_{39}h_3h_0h_4} = \tilde{q}_*(\widehat{D_3}) + \overline{\tilde{e}_{31}h_0h_4^2} \text{ in } \text{Ext}_{\mathcal{A}}^{3,64}(\tilde{P}^{62}).$$

$$(3.6) (4) \quad \text{Ext}_{\mathcal{A}}^{2,48}(\tilde{P}^{62}) \xrightarrow{(q'_{33})_*} \text{Ext}_{\mathcal{A}}^{2,48}(\tilde{P}_{33}^{62} = P_{33}^{62}) \text{ is given by } (q'_{33})_*(\overline{e_{39}h_3h_0}) = \overline{e_{39}h_3h_0}, (q'_{33})_*(\tilde{q}_*(\widehat{h_4h_0h_5})) = 0.$$

$$(3.6) (5) \quad \text{Ext}_{\mathcal{A}}^{0,47}(\tilde{P}^{62}) \xrightarrow{(q'_{33})_*} \text{Ext}_{\mathcal{A}}^{0,47}(\tilde{P}_{33}^{62} = P_{33}^{62}) \text{ is given by } (q'_{33})_*(\overline{e_{47}}) = \overline{e_{47}}.$$

By naturality of the differentials in the Adams spectral sequence, we have the following equations in the ASS for $\pi_*^S(P_{33}^{62})$.

$$(5.25) \quad \begin{aligned} (q'_{33})_*(d_2(\overline{e_{47}})) &= d_2((q'_{33})_*(\overline{e_{47}})) \\ &= d_2(\overline{e_{47}}) \end{aligned} \quad (\text{by (3.6) (5)})$$

We claim

(5.26) In the ASS for $\pi_*^S(P_{33}^{62})$, $d_2(\overline{e_{47}}) \neq 0$.

Assuming (5.26), by (3.4) (5) we must have $d_2(\overline{e_{47}}) = \overline{e_{39}h_0h_3}$ in the ASS for $\pi_*^S(P_{33}^{62})$. From this and (5.25) we have $(q'_{33})_*(d_2(\overline{e_{47}})) = \overline{e_{39}h_0h_3}$, and together with (3.6) (4) we see $d_2(\overline{e_{47}}) = \overline{e_{39}h_3h_0} + \varepsilon\tilde{q}_*(\widehat{h_4h_0h_5})$ in the ASS for $\pi_*^S(\tilde{P}^{62})$, with $\varepsilon = 0$ or 1 . So we have the following equation in the ASS for $\pi_*^S(\tilde{P}^{62})$.

$$(5.27) \quad \begin{aligned} d_2(\overline{e_{47}}h_4h_7) &= d_2(\overline{e_{47}})h_4h_7 && (\text{by (5.23)}) \\ &= (\overline{e_{39}h_3h_0} + \varepsilon\tilde{q}_*(\widehat{h_4h_0h_5}))h_4h_7 && (\text{by } d_2(\overline{e_{47}}) = \overline{e_{39}h_3h_0} + \varepsilon\tilde{q}_*(\widehat{h_4h_0h_5})) \\ &= \overline{e_{39}h_3h_0h_4h_7} \\ &= (\tilde{q}_*(\widehat{D_3}) + \overline{\tilde{e}_{31}h_0h_4^2})h_7 && (\text{by (3.5) (3)}) \\ &= \tilde{q}_*(\widehat{D_3}h_7) + \overline{\tilde{e}_{31}h_0h_4^2}h_7 \end{aligned}$$

This proves (5.4) (2) modulo (5.26). The proof of (5.26) is given as follows.

We use the map $P_{33}^{62} \xrightarrow{\Delta} \Sigma P^{32}$ in (5.24). Recall from (3.17) that $\delta(e_{47}) = e_{31}\lambda_{15}$ in $H_*(P) \otimes \Lambda$. We have $\text{Ext}_{\mathcal{A}}^{1,46}(\Sigma P^{32}) \cong \text{Ext}_{\mathcal{A}}^{1,47}(P^{32})$. Recall

$$(3.4) (10) \quad \text{Ext}_{\mathcal{A}}^{1,47}(P^{32}) = \mathbb{Z}/2(\overline{e_{31}h_4})$$

So we have

$$(5.28) \quad \Delta_*(\overline{e_{47}} = \{e_{47}\}) = \{\delta(e_{47})\} = \{e_{31}\lambda_{15}\} = \overline{e_{31}h_4} \in \text{Ext}_{\mathcal{A}}^{1,47}(P^{32}) \cong \text{Ext}_{\mathcal{A}}^{1,46}(\Sigma P^{32}).$$

Recall

$$(3.4) (11) \quad \text{Ext}_{\mathcal{A}}^{2,32}(P^{32}) = \mathbb{Z}/2(\overline{e_{15}}h_0h_4).$$

$$(3.4) (12) \quad \text{Ext}_{\mathcal{A}}^{3,48}(P^{32}) = \mathbb{Z}/2(\overline{e_{31}}h_0h_3^2) \oplus \mathbb{Z}/2(\overline{e_{15}}h_0h_4^2).$$

Since $\overline{e_{31}} \in \text{Ext}_{\mathcal{A}}^{0,31}(P^{32})$ is a pullback of $\widehat{h}_5 \in \text{Ext}_{\mathcal{A}}^{0,31}(P)$, by naturality of the differentials in the Adams spectral sequence, we have the following equations in the ASS for $\pi_*^S(P^{32})$.

$$\begin{aligned}
(5.29) \quad \Delta_*(d_2(\overline{e_{47}})) &= d_2(\Delta_*(\overline{e_{47}})) \\
&= d_2(\overline{e_{31}}h_4) \\
&= d_2(\overline{e_{31}})h_4 + \overline{e_{31}}d_2(h_4) \\
&= \overline{e_{15}}h_0h_4^2 + \overline{e_{31}}h_0h_3^2 && \text{(by (3.4) (11) and (5.1))} \\
&\neq 0 && \text{(by (3.4) (12))}
\end{aligned}$$

It follows that $d_2(\overline{e_{47}}) \neq 0$ in the ASS for $\pi_*^S(P_{33}^{62})$. This proves (5.26).

This completes the proof of Lemma 5.4 (2).

6 Modification of the proof without θ_6

In this section we discuss how to modify our proof of Theorem 1.2 without using the knowledge of θ_6 . Recall that the Kervaire invariant element $\theta_6 \in \pi_{126}^S$ is a stable homotopy element detected by $h_6^2 \in \text{Ext}_{\mathcal{A}}^{2,128}$ in the ASS for π_*^S . In section 1 through section 4 we have assumed the existence of a $\theta_6 \in \pi_{126}^S$ when proving the following theorems.

(6.1) Theorem 1.4, Theorem 2.2, Theorem 2.17, Proposition 2.46, Lemma 5.3, and Lemma 5.7.

Let $X = S^{124} \cup_{\eta} e^{126}$. We will show later in Proposition 6.8 that there is a stable map $X \xrightarrow{\bar{\theta}_6} S^0$ that is detected by h_6^2 on the top cell e^{126} . Then we will show that the methods that we proved the results in (6.1) assuming the existence of a θ_6 still works when $S^{126} \xrightarrow{\theta_6} S^0$ is replaced by $X \xrightarrow{\bar{\theta}_6} S^0$.

To define the stable map $X \xrightarrow{\bar{\theta}_6} S^0$ we need to use the quadratic construction $D_2(Y)$ of a stable complex Y with base point ([5, 20]). These are recalled as follows. Given a stable complex with base point $(Y, *)$, for each $n \geq 0$ denote the points in $S^n \times (Y \wedge Y)$ by $(\lambda, x \wedge y)$, for $\lambda \in S^n$ and $x, y \in Y$. Define an involution $S^n \times (Y \wedge Y) \xrightarrow{T} S^n \times (Y \wedge Y)$ by

$$T(\lambda, x \wedge y) = (-\lambda, y \wedge x), \quad \text{where } -\lambda \in S^n \text{ is the antipodal point of } \lambda.$$

Then the group $\{1, T\}$ defines a \mathbb{Z}_2 -action on $S^n \times (Y \wedge Y)$. Let $\overline{D}_2^n(Y) = S^n \times_{\mathbb{Z}_2} (Y \wedge Y) = S^n \times (Y \wedge Y) / \{1, T\}$. Let $D_2^n(Y) = \frac{\overline{D}_2^n(Y)}{S^n \times_{\mathbb{Z}_2} \{*\}}$ with base point $* = S^n \times_{\mathbb{Z}_2} \{*\}$. Denote the points in $D_2^n(Y)$ by $[\lambda, x \wedge y]$, where $\lambda \in S^n$ and $x, y \in Y$. It is not difficult to see that $D_2^n(Y) \subset D_2^{n+1}(Y)$ for each $n \geq 0$.

(6.2) The quadratic construction of Y is the complex $D_2(Y) = \cup_{n \geq 0} D_2^n(Y)$.

(6.3) Given a based stable map $Y \xrightarrow{f} X$ where Y, X are stable complexes with base points, $D_2(Y) \xrightarrow{D_2(f)} D_2(X)$ is the base point preserving map defined by

$$D_2(f)([\lambda, x \wedge y]) = [\lambda, f(x) \wedge f(y)], \quad \text{where } \lambda \in S^n, x, y \in Y.$$

Given a stable map $Y \xrightarrow{f} S^0$, consider the induced map $D_2(Y) \xrightarrow{D_2(f)} D_2(S^0)$. It is not difficult to see that $D_2(S^0) = P \cup \{*\}$ where P is the infinite real projective space. Let $P \cup \{*\} \xrightarrow{g} S^0 = \{*, 1\}$ be the map defined by $g(*) = *$, $g(P) = 1$.

(6.4) Let $D_2(Y) \xrightarrow{\gamma(f)} S^0$ be the following composite.

$$\gamma(f) : D_2(Y) \xrightarrow{D_2(f)} D_2(S^0) \xrightarrow{g} S^0.$$

To describe the stable map $X \xrightarrow{\bar{\theta}_6} S^0$, we need to recall from [5] the following Theorem 6.6 and Corollary 6.7. To state these results, suppose there is a Kervaire invariant element $\theta_k \in \pi_n^S$ with $2\theta_k = 0$, where $k \geq 1$ and $n = 2^{k+1} - 2$. Let

Theorem 2.17. *Let $\widehat{\theta}_5 \in \pi_{62}^S(P^{62})$ be as in (2.16). In $\pi_{188}^S(P^{62})$, $\widehat{\theta}_5\theta_6 \neq 0$ and is detected in the ASS for π_*^S by a class*

$$\tau = \widehat{g}_4 h_0^2 + \varepsilon \widehat{h}_5 h_5 D_3(1) + \overline{\tau} \neq 0 \quad \text{in } \text{Ext}_{\mathcal{A}}^{5,193}(P^{62})$$

where $\varepsilon = 0$ or 1 . Here $\overline{\tau} \in \text{Ext}_{\mathcal{A}}^{5,193}(P^{62})$ is a class such that $t_*(\overline{\tau}) = 0$ in $\text{Ext}_{\mathcal{A}}^{6,194}$. Moreover, $t_*(\widehat{g}_4 h_0^2) = h_0^2 g_4$, and $t_*(\widehat{h}_5 h_5 D_3(1)) = h_5^2 D_3(1)$.

Proposition 2.46. *For the composite $S^{188} \xrightarrow{\theta_6} S^{62} \xrightarrow{\widehat{\theta}_5} P^{62}$, $AF(\widehat{\theta}_5\theta_6) \geq 5$.*

Lemma 5.3. *The composite $S^{188} \xrightarrow{\theta_6} S^{62} \xrightarrow{\widehat{\theta}_5} P^{62} \xrightarrow{\widetilde{q}} \widetilde{P}^{62}$ has $AF \geq 5$ in the ASS for $\pi_*^S(\widetilde{P}^{62})$.*

Denote $\Sigma^{62}(X = S^{124} \cup_{\eta} e^{126}) \cong S^{186} \cup_{\eta} e^{188}$ still by X , and $X = S^{186} \cup_{\eta} e^{188} \xrightarrow{\Sigma^{62}\overline{\theta}_6} S^{62}$ still by θ_6 . In each of Theorem 2.2, Theorem 2.17, Proposition 2.46, and Lemma 5.3, replace S^{188} with X , and $S^{188} \xrightarrow{\theta_6} S^{62}$ with $X \xrightarrow{\overline{\theta}_6} S^{62}$. To see that the arguments in the original proofs still work after this modification, consider the cofibration $S^{186} \xrightarrow{i_1} X \xrightarrow{q_1} S^{188}$. From (6.5) we see that the composite $S^{186} \xrightarrow{i_1} X \xrightarrow{\overline{\theta}_6} S^{62}$ is $S^{186} \xrightarrow{\theta_5^2} S^{62}$. Since θ_5 is detected by h_5^2 in the ASS for π_*^S and since $h_5^4 = 0$ in $\text{Ext}_{\mathcal{A}}^{*,*}$, we see $AF(\theta_5^2 = \overline{\theta}_6 i_1) \geq 5$. If $AF(\theta_5^2) = 5$ then θ_5^2 is detected by a cohomology class $\alpha \in \text{Ext}_{\mathcal{A}}^{5,129}$. From (3.13) we deduce

$$(6.9) \quad \text{Ext}_{\mathcal{A}}^{5,129} = 0.$$

Therefore we have

$$(6.10) \quad AF(\theta_5^2 = \overline{\theta}_6 i_1) \geq 6.$$

It follows that the composite $S^{186} \xrightarrow{i_1} X \xrightarrow{\overline{\theta}_6} S^{62} \xrightarrow{\widehat{\theta}_5} P^{62}$ has $AF \geq 7$. Then it is not difficult to see that all the arguments for proving Theorem 2.2, Theorem 2.17, Proposition 2.46, and Lemma 5.3 carry through if we replace $S^{188} \xrightarrow{\theta_6} S^{62}$ by $X \xrightarrow{\overline{\theta}_6} S^{62}$.

We proceed to describe the modification for the following

Lemma 5.7. *In the ASS for π_*^S , $d_3(h_6^3) \neq h_5^2 D_3(1)$.*

In the proof of Lemma 5.7 we have assumed a Kervaire invariant element $S^{189} \xrightarrow{\theta_6} S^{63}$ in the following content.

$$(5.10) \quad (2) \quad \text{The composite } S^{189} \xrightarrow{\theta_6} S^{63} \xrightarrow{\overline{\theta}_5} S^0 \cup_{2\iota} e^1 \text{ is detected by } (i_1)_*(h_6^3) \text{ in the ASS for } \pi_*^S(S^0 \cup_{2\iota} e^1).$$

We replace S^{189} and $S^{189} \xrightarrow{\theta_6} S^{63}$ by $\Sigma X = S^{187} \cup_{\eta} e^{189}$ and $\Sigma X \xrightarrow{\Sigma\overline{\theta}_6} S^{63}$ respectively in the proof of (5.10) (2). From (6.10) it is not difficult to see that the modified proof of (5.10) carries through. This completes the modification for Lemma 5.7.

Finally we describe the modification for Theorem 1.4.

Theorem 1.4. *$h_0^3 g_4$ is a boundary in the ASS for spheres.*

We have assumed a Kervaire invariant element θ_6 in the proof of Theorem 1.4 when using Theorem 2.2 and the property $2\theta_5\theta_6 = 0$. We replace Theorem 2.2 with the modified version, and replace each $S^{188} \xrightarrow{\theta_6} S^{62}$ by $X \xrightarrow{\bar{\theta}_6} S^{62}$. We also have $2\theta_5\bar{\theta}_6 = 0$ because $2\theta_5 = 0$. Then the modified proof for Theorem 1.4 carries through.

This completes the modifications to the proof of Theorem 1.2 with $S^{126} \xrightarrow{\theta_6} S^0$ replaced by $X \xrightarrow{\bar{\theta}_6} S^0$.

References

- [1] J. F. Adams, *On the non-existence of elements of Hopf invariant one*, Ann. of Math. (2) **72** (1960), 20–104.
- [2] ———, *On the structure and applications of the Steenrod algebra*, Comment. Math. Helv. **32** (1958), 180–214.
- [3] ———, *Vector fields on spheres*, Ann. of Math. (2) **75** (1962), 603–632.
- [4] M. G. Barratt, M. E. Mahowald, and M. C. Tangora, *Some differentials in the Adams spectral sequence. II*, Topology **9** (1970), 309–316.
- [5] M. G. Barratt, J. D. S. Jones, and M. E. Mahowald, *The Kervaire invariant problem*, Proceedings of the Northwestern Homotopy Theory Conference (Evanston, Ill., 1982), 1983, pp. 9–22.
- [6] A. K. Bousfield, E. B. Curtis, D. M. Kan, D. G. Quillen, D. L. Rector, and J. W. Schlesinger, *The mod $-p$ lower central series and the Adams spectral sequence*, Topology **5** (1966), 331–342.
- [7] William Browder, *The Kervaire invariant of framed manifolds and its generalization*, Ann. of Math. (2) **90** (1969), 157–186.
- [8] Tai-Wei Chen, *Determination of $\text{Ext}_{\mathbb{A}}^{5,*}(\mathbb{Z}/2, \mathbb{Z}/2)$* , Topology and its Applications **158** (2011), no. 5, 660–689.
- [9] Ralph L. Cohen, Wên-Hsiung Lin, and M. E. Mahowald, *The Adams spectral sequence of the real projective spaces*, Pacific J. Math. **134** (1988), no. 1, 27–55.
- [10] Dale Husemoller, *Fibre bundles*, Third, Graduate Texts in Mathematics, vol. 20, Springer-Verlag, New York, 1994.
- [11] I. M. James, *Whitehead products and vector-fields on spheres*, Proc. Cambridge Philos. Soc. **53** (1957), 817–820.
- [12] Daniel S. Kahn and Stewart B. Priddy, *Applications of the transfer to stable homotopy theory*, Bulletin of the American Mathematical Society **78** (1972), 981–987.
- [13] Chung-Wei Kan, *A differential $d_4(h_5^3)$ in the Adams spectral sequence for spheres*, Ph.D. Thesis, National Tsing Hua University, 2001.
- [14] Michel A. Kervaire, *A manifold which does not admit any differentiable structure*, Commentarii Mathematici Helvetici **34** (1960), 257–270.
- [15] Stanley O. Kochman and Mark Mahowald, *On the computation of stable stems*, The Čech centennial (Boston, MA, 1993), 1995, pp. 299–316.
- [16] Wên-Hsiung Lin, *$\text{Ext}_{\mathbb{A}}^{4,*}(\mathbb{Z}/2, \mathbb{Z}/2)$ and $\text{Ext}_{\mathbb{A}}^{5,*}(\mathbb{Z}/2, \mathbb{Z}/2)$* , Topology Appl. **155** (2008), no. 5, 459–496.
- [17] ———, *A differential in the Adams spectral sequence for spheres*, Stable and unstable homotopy (Toronto, ON, 1996), 1998, pp. 205–239.
- [18] ———, *Algebraic Kahn-Priddy theorem*, Pacific J. Math. **96** (1981), no. 2, 435–455.
- [19] Mark Mahowald and Martin Tangora, *Some differentials in the Adams spectral sequence*, Topology **6** (1967), 349–369.
- [20] R. James Milgram, *Unstable homotopy from the stable point of view*, Lecture Notes in Mathematics, vol. 368, Springer-Verlag, Berlin, 1974.

- [21] Mamoru Mimura and Hiroshi Toda, *The $(n+20)$ -th homotopy groups of n -spheres*, Journal of Mathematics of Kyoto University **3** (1963), 37–58.
- [22] Mamoru Mimura, *On the generalized Hopf homomorphism and the higher composition. II. $\pi_n^i(S^n)$ for $i = 21$ and 22* , Journal of Mathematics of Kyoto University **4** (1965), 301–326.
- [23] Dennis Sullivan, *Genetics of homotopy theory and the Adams conjecture*, Ann. of Math. (2) **100** (1974), 1–79.
- [24] Martin C. Tangora, *On the cohomology of the Steenrod algebra*, Math. Z. **116** (1970), 18–64.
- [25] Hiroshi Toda, *Composition methods in homotopy groups of spheres*, Annals of Mathematics Studies, No. 49, Princeton University Press, Princeton, N.J., 1962.
- [26] George W. Whitehead, *Recent advances in homotopy theory*, American Mathematical Society, Providence, R.I., 1970. Conference Board of the Mathematical Sciences Regional Conference Series in Mathematics, No. 5.