SU(2)-CYCLIC SURGERIES ON KNOTS

JIANFENG LIN

ABSTRACT. A surgery on a knot in S^3 is called SU(2)-cyclic if it gives a manifold whose fundamental group has no non-cyclic SU(2) representations. Using holonomy perturbations on the Chern-Simons functional, we prove that two SU(2)-cyclic surgery coefficients $\frac{p_1}{q_1}$ and $\frac{p_2}{q_2}$ should satisfy $|p_1q_2-p_2q_1| \leq |p_1|+|p_2|$. This is an analog of Culler-Gordon-Luecke-Shalen's cyclic surgery theorem.

1. Introduction

Definition 1.1. A closed orientable 3 manifold M, M is called SU(2)-cyclic (or SO(3)-cyclic) if there exists no homomorphism $\phi: \pi_1(M) \to SU(2)$ (or SO(3)) with non-cyclic image.

Suppose $K \subset S^3$ is a knot. For $r \in \mathbb{Q}$, we denote the manifold obtained by doing r-surgery on K by K(r).

Definition 1.2. A surgery on K with coefficient r is called SU(2)-cyclic (or SO(3)-cyclic) if K(r) is SU(2)-cyclic (or SO(3)-cyclic).

We have the following exact sequence:

$$0 \to \mathbb{Z}_2 \to SU(2) \to SO(3) \to 0$$

It's easy to see that an SO(3)-cyclic surgery is always an SU(2)-cyclic surgery. Using some basic obstruction theory, we get:

Lemma 1.3. If $r = \frac{p}{q}$ is an SU(2)-cyclic surgery with p odd, then r is an SO(3)-cyclic surgery.

In [5], Kronheimer and Mrowka proved the following theorem:

Theorem 1.4 (Kronheimer, Mrowka 2003 [5]). Any r-surgery on a nontrivial knot with surgery coefficient $|r| \le 2$ is not SU(2)-cyclic.

In particular, this theorem gave a proof for the Property-P Conjecture:

Corollary 1.5 (Kronheimer, Mrowka 2003 [5]). A nontrivial surgery on a nontrivial knot does not give simply connected 3-manifold.

Obviously, lens spaces are all SU(2)-cyclic and SO(3)-cyclic. Thus all cyclic surgeries (the surgeries which give lens spaces) are SO(3)-cyclic. Therefore, we have:

Example 1.6. The $pq - \frac{1}{r}$ $(r \in \mathbb{Z})$ surgeries on the (p,q)-torus knot are cyclic and hence SO(3)-cyclic.

Dunfield [4] gives the following example:

Example 1.7. The $18, \frac{37}{2}, 19$ surgeries on the (-2,3,7)-pretzel knot are SO(3)-cyclic. The 18, 19 surgeries give lens spaces, while $K(\frac{37}{2})$ is a graph manifold obtained by gluing the left-handed trefoil knot complement and the right-handed trefoil complement.

Another related theorem is Culler-Gordon-Luecke-Shalen's cyclic surgery theorem (we only state the case for knot surgery):

Theorem 1.8 (Culler-Gordon-Luecke-Shalen [8]). Suppose that K is not a torus knot and r, s are both cyclic surgeries, then $\triangle(r, s) \le 1$.

Here for two rational numbers $r = \frac{p_1}{q_1}$ and $s = \frac{p_2}{q_2}$, the distance $\triangle(r, s)$ is defined to be $|p_1q_2 - p_2q_1|$.

Since $\frac{1}{0}$ -surgery is always cyclic, this theorem implies that when K is not a torus knot, r-surgery can be cyclic only if $r \in \mathbb{Z}$. Moreover, there are at most two such integers, and if there are two then they must be successive.

Although Example 1.7 shows that Theorem 1.8 is not true for SU(2)-cyclic or SO(3)-cyclic surgeries, we have the following analogous result, which is the main theorem of this paper.

Theorem 1.9. Consider a nontrivial knot $K \subset S^3$ and two surgeries with coefficients $r_1 = p_1/q_2$ and $r_2 = p_2/q_2$. We have the following:

- If r_1, r_2 are both SU(2)-cyclic, then $\triangle(r_1, r_2) \le |p_1| + |p_2|$.
- If r_1, r_2 are both SO(3)-cyclic, then $2\triangle(r_1, r_2) \leq |p_1| + |p_2|$.

Combining this theorem with Lemma 1.3, we get the following corollaries.

Corollary 1.10. Suppose r_1, r_2 are both SU(2)-cyclic. If p_1 is odd, then $2\triangle(r_1, r_2) \le 2|p_1| + |p_2|$. If p_1, p_2 are both odd, then $2\triangle(r_1, r_2) \le |p_1| + |p_2|$.

Corollary 1.11. If r_1, r_2 on K are both SO(3)-cyclic surgeries, then $r_1r_2 > 0$. If r_1, r_2 on K are both SU(2)-cyclic surgeries, then $r_1r_2 > 0$ unless r_1 and r_2 are both even integers.

Corollary 1.12. For a nontrivial surgery on a nontrivial amphichiral knot K with coefficient r, we have the following:

- It can never be SO(3)-cyclic.
- If it is SU(2)-cyclic, then r is an even integer and some $\frac{r}{2}$ -th root of unity is a root of Δ_K (the Alexander polynomial of K).

Remark 1.13. Actually, we haven't found any examples of SU(2)-cyclic surgeries on an amphichiral knot. It would be interesting to know whether there exists such a surgery.

We know that $\Delta_K(1) = \pm 1$ for any knot K while $\Phi_p(1) = p$ for any prime number p (Φ_p is the p-th cyclotomic polynomial). Therefor the Alexander polynomial Δ_K never has the p-th root of unity as its root. We get:

Example 1.14. If p is prime, then the 2p-surgery on an non-trivial amphichiral knot is not SU(2)-cyclic.

Remark 1.15. In [5], Kronheimer and Mrowka asked whether there exists SU(2)-cyclic surgery with coefficient 3 or 4. We see that there exist no such surgeries for nontrivial amphichiral knots.

Using the criterion in Corollary 1.12, we checked the amphichiral knots with crossing number ≤ 10 and get:

Example 1.16. All the nontrivial amphichiral knots with crossing number ≤ 10 except perhaps 8_{18} and 10_{99} in Rolfsen's knot table admit no SU(2)-cyclic surgeries. For 8_{18} and 10_{99} , we have no examples of SU(2)-cyclic surgeries.

Corollary 1.17. Given a nontrivial knot K and an integer q, there exist at most finitely many $p \in \mathbb{Z}$ such that (p,q) = 1 and the $\frac{p}{q}$ -surgery on K is SO(3)-cyclic. For the SU(2) case, the only possible exception is when q = 1 and infinitely many even p.

In particular, any nontrivial knot admits only finitely many integer SO(3)-cyclic surgeries and only finite many odd SU(2)-cyclic surgeries.

The paper is organized as follows: in section 2, we review some preliminaries and basic constructions related to holonomy perturbations. In section 3, we prove the main theorem and the corollaries.

Acknowledgement The author wishes to thank Nathan Dunfield, Cameron Gordon, Yi Ni and Yi Liu for valuable discussions and comments. The author is especially grateful to Ciprian Manolescu for inspiring conversations and helpful suggestions in writing this paper.

2. Preliminaries

In this section, we review the basic facts about holonomy perturbations. Most details can be found in [5] and [9]. The constructions are very similar to [13], but for completeness, we review them again here.

Consider the closed manifold K(0). We have $b_1(K(0)) = 1$. Let E be the rank 2 unitary bundle over K(0) with $c_1(E)$ the Poincaré dual of the meridian m of K. Let \mathfrak{g}_E be the bundle whose sections are traceless, skew-hermitian endomorphisms of E. Let \mathcal{A} be the affine space of SO(3) connections of \mathfrak{g}_E . Let \mathcal{G} be the group of gauge transformations on E with determinant 1. Notice that \mathcal{G} is slightly smaller than the SO(3)-gauge transformation group of \mathfrak{g}_E .

Fix a reference connection A_0 on \mathfrak{g}_E . Then for any connection A on \mathfrak{g}_E , $A - A_0$ can be identified with $\omega \in \Omega^1(\mathfrak{g}_E)$. We have the Chern-Simons functional:

$$CS: \mathcal{A} \to \mathbb{R}$$

$$CS(A) = \frac{1}{4} \int_{X_0} Tr(2\omega \wedge F_{A_0} + \omega \wedge d\omega + \frac{2}{3}\omega \wedge \omega \wedge \omega)$$

Here F_{A_0} is the curvature of A_0 .

The critical points of the Chern-Simons functional are the flat connections.

Floer introduced the holonomy perturbations as follows. Take a function $\phi: SU(2) \to \mathbb{R}$ which is invariant under conjugation. Then it is uniquely determined by the even, 2π -periodic function:

$$f(x) := \phi \begin{pmatrix} e^{ix} & 0 \\ 0 & e^{-ix} \end{pmatrix} \tag{1}$$

Let D be a compact 2-manifold with boundary. Consider an embedding $D \times S^1$ in K(0) such that \mathfrak{g}_E is trivial over it. Fix a trivialization of \mathfrak{g}_E over $D \times S^1$ and take a 2-form μ which is supported in the interior of D with integral 1. Using the trivialization, we can lift A to a connection \overline{A} on the trivialized SU(2)-bundle \widetilde{P} over $D \times S^1$. We consider the functional:

$$\Phi: \mathcal{A} \to \mathbb{R}$$

$$\Phi(A) := \int_{p \in D} \phi(\operatorname{Hol}_{\{p\} \times S^1}(\bar{A})) \mu(p)$$
(2)

Here $\operatorname{Hol}_{\{p\}\times S^1}$ is the holonomy along $\{p\}\times S^1.$

We decompose K(0) into three parts: $(S^3 - N(K)) \underset{\{0\} \times l \times m}{\cup} ([0,1] \times l \times m) \underset{\{1\} \times l \times m}{\cup} (D^2 \times l \times m)$

m). We have meridians and longitudes on both side of the thicken torus. Denote them by m_0 , l_0 , m_1 , l_1 respectively.

We should be careful that m_0 is the meridian of the knot complement but m_1 the longitude of the attached solid torus. Also l_0 is the longitude of the knot complement but the l_1 is the meridian of the attached solid torus.

For our purpose, we will do two types of perturbations:

- Set $D \cong D^2$ and $i_1(D^2 \times S^1) = (D^2 \times m) \subset K(0)$. That means we use the holonomy along m to do the perturbation. We denote this perturbation by Φ_1
- Set $D \cong m \times [0,1]$ (D is an annulus) and $i_2(D \times S^1) = (m \times [0,1]) \times l \subset K(0)$. That means we embed a thickened torus and use the holonomy along l to do the perturbation. We denote this perturbation by Φ_2 .

We choose a trivialization of \mathfrak{g}_E over $(D^2 \times m) \cup (m \times [0,1] \times l)$ and use it to lift the connection A to a SU(2)-connection \overline{A} on \widetilde{P} . Now use Formula (2) and consider the perturbed Chern-Simons functional $\widehat{CS} = CS + \Phi_1 + \Phi_2 : \mathcal{A} \to \mathbb{R}$.

The following theorem was first proved in [7]:

Theorem 2.1 (Kronheimer, Mrowka [7]). If K is a nontrivial knot, then for any holonomy perturbation, the perturbed Chern-Simons functional \widehat{CS} over K(0) always has at least one critical point.

Remark 2.2. The proof of this theorem is highly nontrivial. It combines Gabai's result about taut foliation in [2], Eliashberg-Thurston's theorem about symplectic filling in [15] and [16], Taubes's result about the Seiberg-Witten invariants of the symplectic four-manifold in [11], Feehan and Leness's work about Witten's conjecture in [10] and Kronheimer-Mrowka's work about the refinement of Eliashberg-Thurston's theorem in [7].

The critical points can be completed determined:

Lemma 2.3. If $A \in \mathcal{A}$ is a critical point of \widehat{CS} , then:

- A is flat on $S^3 N(K) \subset K(0)$.
- We can choose suitable trivialization of the SU(2)-bundle \widetilde{P} such that: $Hol_{m_0}(\overline{A}) = \begin{pmatrix} e^{i\theta_0} & 0 \\ 0 & e^{-i\theta_0} \end{pmatrix}, Hol_{m_1}(\overline{A}) = \begin{pmatrix} e^{i\theta_1} & 0 \\ 0 & e^{-i\theta_1} \end{pmatrix}, Hol_{l_0}(\overline{A}) = \begin{pmatrix} e^{i\eta_0} & 0 \\ 0 & e^{-i\eta_0} \end{pmatrix}$ and $Hol_{l_1}(\overline{A}) = \begin{pmatrix} e^{i\eta_1} & 0 \\ 0 & e^{-i\eta_1} \end{pmatrix}$.
- $\eta_0 = \eta_1 = -f_2'(\theta_1) + 2\mathbb{Z}\pi \text{ and } \theta_0 \theta_1 = -f_1'(\eta_0) + 2\mathbb{Z}\pi.$

Remark 2.4. Recall that we chose $\phi_i : SU(2) \to \mathbb{R}$ to define the perturbation Φ_i (i = 1, 2), which gives us $f_i : \mathbb{R} \to \mathbb{R}$ by formula (1).

Proof of Lemma 2.3. By Lemma 4 in [9] and Lemma 2.2 in [13], A is flat on $S^3 - N(K) \subset K(0)$ and near $(m \times l \times \{0\}) \cup (m \times l \times \{1\})$. Moreover, we can choose a suitable trivialization of \widetilde{P} such that $Hol_{m_0}(\overline{A}) = \begin{pmatrix} e^{i\theta_0} & 0 \\ 0 & e^{-i\theta_0} \end{pmatrix}$, $Hol_{m_1}(\overline{A}) = \begin{pmatrix} e^{i\theta_1} & 0 \\ 0 & e^{-i\theta_1} \end{pmatrix}$, $Hol_{l_0}(\overline{A}) = \begin{pmatrix} e^{i\eta_1} & 0 \\ 0 & e^{-i\eta_1} \end{pmatrix}$ and $\theta_0 - \theta_1 = -f'_1(\eta_0) + 2\mathbb{Z}\pi$. Also, we can choose another trivialization of \widetilde{P} such that $Hol_{m_1}(\overline{A}) = \begin{pmatrix} e^{i\theta'_1} & 0 \\ 0 & e^{-i\theta'_1} \end{pmatrix}$, $Hol_{l_1}(\overline{A}) = \begin{pmatrix} e^{i\eta'_1} & 0 \\ 0 & e^{-i\eta'_1} \end{pmatrix}$ and $\eta'_1 = -f'_2(\theta'_1) + 2\mathbb{Z}\pi$. Since different trivializations give the same holonomy modulo conjugation. We have $(\theta'_1, \eta'_1) = \pm (\theta_1, \eta_1)$. Since f'_2 is an odd function, we have $\eta_1 = -f'_2(\theta_1) + 2\mathbb{Z}\pi$.

Now suppose A is a critical point. Since \mathfrak{g}_E is trivial over $\pi_1(S^3 - N(K))$, we fix a trivialization of $\mathfrak{g}_E|_{S^3-N(K)}$. Using this trivialization, we lift the connection A to a SU(2)-connection \widetilde{A} over $S^3 - N(K)$.

Remark 2.5. This trivialization of $\mathfrak{g}_E|_{\pi_1(S^3-N(K))}$ does not agree with the trivialization over $(D^2 \times m) \cup (m \times [0,1] \times l)$ (the trivialization which we chose to define the holonomy perturbations) on the torus boundary. They differ by a map $f: \partial(S^3-N(K)) \to SO(3)$ such that

$$f_*(m) = 1 \in \pi_1(SO(3)), f_*(l) = -1 \in \pi_1(SO(3))$$
 (3)

By taking the holonomy of \widetilde{A} , we get a representation $\rho: \pi_1(S^3 - N(K)) \to SU(2)$.

Definition 2.6. We define a subset R_K of $(\mathbb{R}/2\pi\mathbb{Z}) \bigoplus (\mathbb{R}/2\pi\mathbb{Z})$ as follows:

$$\{(\theta,\eta)|\exists \rho: \pi_1(S^3-N(K)) \to SU(2) \text{ s.t. } \rho(m) = \left(\begin{smallmatrix} e^{i\theta} & 0 \\ 0 & e^{-i\theta} \end{smallmatrix}\right); \rho(l) = \left(\begin{smallmatrix} e^{i\eta} & 0 \\ 0 & e^{-i\eta} \end{smallmatrix}\right)\}$$

We can also describe R_K as:

$$\{(\theta,\eta)|\exists \text{ flat connection } \widetilde{A} \text{ over } S^3-N(K) \text{ s.t. } \operatorname{Hol}_m(\widetilde{A}) = \left(\begin{smallmatrix} e^{i\theta} & 0 \\ 0 & e^{-i\theta} \end{smallmatrix}\right); \operatorname{Hol}_l(\widetilde{A}) = \left(\begin{smallmatrix} e^{i\eta} & 0 \\ 0 & e^{-i\eta} \end{smallmatrix}\right)\}$$

Notation 2.7. Let $S \subset (\mathbb{R}/2\pi\mathbb{Z}) \bigoplus (\mathbb{R}/2\pi\mathbb{Z})$ be a subset. If h is a function with period 2π , we denote the set $\{(\theta, \eta + h(\theta)) | (\theta, \eta) \in S\}$ by S + (*, h) and the set $\{(\theta + h(\eta), \eta) | (\theta, \eta) \in S\}$ by S + (h, *). We also denote the set $\{(\theta + a, \eta + b) | (\theta, \eta) \in S\}$ by S + (a, b) for constant a, b.

The following lemma is proved in [5]. We change the statement a little. For completeness, we give the proof here.

Lemma 2.8. R_K has the following properties:

- 1) Any point in R_K off the line $\{\eta = 2\pi \mathbb{Z}\}$ gives some non-cyclic representation.
- 2) R_K is a closed subset of $(\mathbb{R}/2\pi\mathbb{Z}) \bigoplus (\mathbb{R}/2\pi\mathbb{Z})$.
- $3)R_K = R_K + (\pi, 0).$
- $4)R_K \cap \{\theta = k\pi\} = (k\pi, 2k'\pi), (k, k' \in \mathbb{Z}).$
- 5) $\exists \epsilon > 0$ such that $\forall k \in \mathbb{Z}, R_K \cap \{\theta \in [k\pi \epsilon, k\pi + \epsilon]\} \cap \{\eta \neq 2\mathbb{Z}\pi\} = \emptyset$.

Proof. 1) Any point in R_K gives a representation $\rho: \pi_1(S^3 - N(K)) \to SU(2)$. If ρ is cyclic, then ρ factors through $H_1(S^3 - N(K))$, which implies that $\rho(l) = 1 \in SU(2)$.

- 2) R_K is closed because $\pi_1(S^3 K)$ is finitely generated and SU(2) is compact. 3) We have a map $\rho_0: \pi_1(S^3 K) \to H_1(S^3 K) \to \mathbb{Z}_2 \subset SU(2)$ such that $\rho_0(m) =$ $-1 \in SU(2)$ and $\rho_1(l) = 1 \in SU(2)$. For any homomorphism $\rho : \pi_1(S^3 - N(K)) \to SU(2)$, we can multiply it by ρ_0 to get another representation ρ' such that $\rho'(l) = \rho(l)$ and $\rho'(m) = -\rho(m)$. By definition of R_K , we get $R_K = R_K + (\pi, 0)$.
- 4) Suppose ρ is given by a point with $\theta = 0$, then $\rho(m) = 1 \in SU(2)$ and ρ factors through $\pi_1(S^3)$, which is a trivial. We get $\rho(l) = 1$ and $\eta = 2k'\pi$. For the case $\theta = \pi$, we use 3).
- 5) Look at a small neighborhood U of $(0,0) \in R_K$ in $(\mathbb{R}/2\pi\mathbb{Z}) \bigoplus (\mathbb{R}/2\pi\mathbb{Z})$. The point (0,0) is given by the restriction of the trivial representation ρ_1 . The deformations of ρ_1 are governed by $H^1(\pi_1(S^3-K),\mathbb{R}^3)\cong\mathbb{R}^3$. But every vector in this \mathbb{R}^3 can be realized by the some reducible representation. We see that in a small neighborhood of ρ_1 , all the representations are reducible. Thus $U \cap R_K \cap \{\eta \neq 2\mathbb{Z}\pi\} = \emptyset$ if U is small enough. Use 4) and the compactness of R_K , we prove 5) for the case k=0. Then we use 3) to prove the case k = 1.

Lemma 2.9. If A is a critical point of the perturbed Chern-Simons functional, then $(\theta_0, \eta_0) \in R_K + (0, -\pi)$. $(\theta_0 \text{ and } \eta_0 \text{ are defined in Lemma 2.3})$

Proof. On the torus $m \times l \times \{0\}$, we use two different trivializations to lift A to two SU(2)connections \overline{A} and A. Because of formula (3), we see that $(\operatorname{Hol}_{m_0}(A), \operatorname{Hol}_{l_0}(A)) \in SU(2) \times I$ SU(2) is conjugate with $(\operatorname{Hol}_{m_0}(\overline{A}), -\operatorname{Hol}_{l_0}(\overline{A}))$. So we can change the trivialization of \mathfrak{g}_E over $S^3 - N(K)$ so that $(\operatorname{Hol}_{m_0}(\overline{A}), -\operatorname{Hol}_{l_0}(\overline{A})) = (\operatorname{Hol}_{m_0}(\widetilde{A}), \operatorname{Hol}_{l_0}(\overline{A}))$. Then we use the second description of R_K .

Combining this lemma and Lemma 2.3, we get:

Lemma 2.10. If A is a critical point of the perturbed Chern-Simons functional, then $(\theta_1, \eta_1) \in (R_K + (0, -\pi) + (f_1', *)) \cap \{\eta = -f_2'(\theta)\}.$

3. Proof of the Theorem and Corollaries

3.1. **Proof of the main theorem.** Now suppose $K \subset S^3$ is a nontrivial knot. Denote the set $R_K \cap \{\eta \neq 2\mathbb{Z}\pi\}$ by R_K^* . For $r = \frac{p}{q}$, we define the subsets S(r) and $\widehat{S}(r)$ to be:

$$S(r) := \{ (\theta, \eta) | (p\theta + q\eta) \in 2\mathbb{Z}\pi \text{ or } (p\theta + p\pi + q\eta) \in 2\mathbb{Z}\pi \}$$
$$\widehat{S}(r) := \{ (\theta, \eta) | (p\theta + q\eta) \in \mathbb{Z}\pi \}$$

Remark 3.1. When p is odd, we have $\widehat{S}(r) = S(r)$.

Lemma 3.2. If r is an SU(2)-cyclic surgery, then $R_K^* \cap S(r) = \emptyset$. If r is an SO(3)-cyclic surgery, then $R_K^* \cap \widehat{S}(r) = \emptyset$.

Proof. If $(\theta, \eta) \in R_K^*$ satisfies $p\theta + q\eta \in 2\mathbb{Z}\pi$, then it gives a representation $\rho : \pi_1(S^3 - N(K)) \to SU(2)$ such that $\rho(pm + ql) = 1 \in SU(2)$. Thus ρ factors through $\pi_1(K(r))$. By (1) of Lemma 2.8, ρ is non-cyclic. We get the contradiction since r is a SU(2)-cyclic surgery. We see that $R_K^* \cap \{(\theta, \eta) | (p\theta + q\eta) \in 2\mathbb{Z}\pi\} = \emptyset$. By (3) of Lemma2.8, we have $R_K^* + (\pi, 0) = R_K^*$. Thus we also have $R_K^* \cap \{(\theta, \eta) | (p\theta + p\pi + q\eta) \in 2\mathbb{Z}\pi\} = \emptyset$. We proved the first assertion. The second assertion can be proved similarly.

Since we are considering the subsets of $(\mathbb{R}/2\pi\mathbb{Z}) \bigoplus (\mathbb{R}/2\pi\mathbb{Z})$, it will be convenient to fix a region $W = \{(\theta, \eta) | \theta \in (-\infty, \infty), \eta \in [0, 2\pi]\} \subset \mathbb{R}^2$. We define W^* to be $\{(\theta, \eta) | \theta \in (-\infty, \infty), \eta \in (0, 2\pi)\}$. We can work in W and W^* and then project to $(\mathbb{R}/2\pi\mathbb{Z}) \bigoplus (\mathbb{R}/2\pi\mathbb{Z})$.

For two different numbers $r_1 = \frac{p_1}{q_1}, r_2 = \frac{p_2}{q_2}$. We define another two numbers:

$$d_1(r_1, r_2) = \begin{cases} \frac{2\pi |p_1|}{\Delta(r_1, r_2)} & \text{if } p_2 \text{ is even} \\ \frac{\pi |p_1|}{\Delta(r_1, r_2)} & \text{if } p_2 \text{ is odd} \end{cases}; d_2(r_1, r_2) = \begin{cases} \frac{2\pi |p_2|}{\Delta(r_1, r_2)} & \text{if } p_1 \text{ is even} \\ \frac{\pi |p_2|}{\Delta(r_1, r_2)} & \text{if } p_1 \text{ is odd} \end{cases}$$

The intersection $S(r_i) \cap W^*$ are just some line segments of slope $-r_i$ and $S(r_1) \cap S(r_2) \cap W^*$ consists of isolated points. We say two intersection points in $S(r_1) \cap S(r_2) \cap W^*$ are adjacent in $S(r_i)$ (i = 1, 2) if they lie in the same component of $S(r_i) \cap W^*$ and there is no intersection point between them. We define two intersection points in $\widehat{S}(r_1) \cap \widehat{S}(r_2) \cap W^*$ to be adjacent in $\widehat{S}(r_i)$ in a similar way.

The following lemma is easy to prove:

Lemma 3.3. (1) If two intersection points $(\theta, \eta), (\theta', \eta') \in S(r_1) \cap S(r_2) \cap W^*$ are adjacent in $S(r_i)$, then $|\eta - \eta'| = d_i(r_1, r_2)$ (i = 1, 2).

- (2) If two intersection points $(\theta, \eta), (\theta', \eta') \in \widehat{S}(r_1) \cap \widehat{S}(r_2) \cap W^*$ are adjacent in $\widehat{S}(r_i)$, then $|\eta \eta'| = \frac{\pi |p_i|}{\Delta(r_1, r_2)}$ (i = 1, 2).
- (3) For $(\theta, \eta) \in S(r_1) \cap S(r_2) \cap W^*$, if $\eta > d_i(r_1, r_2)$, then we can find $(\theta', \eta') \in S(r_1) \cap S(r_2) \cap W^*$ such that they are adjacent in $S(r_i)$ and $\eta' < \eta$. If $\eta < 2\pi d_i(r_1, r_2)$, then we can find $(\theta', \eta') \in S(r_1) \cap S(r_2) \cap W^*$ such that they are adjacent in $S(r_i)$ and $\eta' > \eta$.
- (4) For $(\theta, \eta) \in \widehat{S}(r_1) \cap \widehat{S}(r_2) \cap W^*$, if $\eta > \frac{\pi |p_i|}{\Delta(r_1, r_2)}$, then we can find $(\theta', \eta') \in \widehat{S}(r_1) \cap \widehat{S}(r_2) \cap W^*$ such that they are adjacent in $\widehat{S}(r_i)$ and $\eta' < \eta$. If $\eta < 2\pi \frac{\pi |p_i|}{\Delta(r_1, r_2)}$, then we can find $(\theta', \eta') \in \widehat{S}(r_1) \cap \widehat{S}(r_2) \cap W^*$ such that they are adjacent in $\widehat{S}(r_i)$ and $\eta' > \eta$.

Now we can start the proof of our main theorem:

Proof of Theorem 1.9. Let r_1, r_2 be two SU(2)-cyclic surgeries. Since the theorem is trivial when $r_1 = r_2$, we always assume that $r_1 \neq r_2$. By Theorem 1.4, we have $|r_i| > 2$. Moreover, when r_1 or r_2 equals $\frac{1}{0}$, the identities in the theorem and corollaries can be easily deduced

from Theorem 1.4. Thus we can assume $p_i \neq 0$ and $q_i \neq 0$. Suppose $d_1(r_1, r_2) + d_2(r_1, r_2) < 2\pi$. By Lemma 3.2 and (4) of Lemma 2.8, we have $R_K^* \cap (S(r_1) \bigcup S(r_2) \bigcup \{\theta = k\pi\}) = \emptyset$. We will construct a broken line $L : [-1, 1] \to W$ such that $Im(L) \subset S(r_1) \bigcup S(r_2) \bigcup \{\theta = k\pi\}$. There are two cases:

(1) Suppose $r_1 < -2 < 2 < r_2$. Let $L(0) = (0, \pi)$. Then as t increases, L first goes up along $\theta = 0$ to $(0, 2\pi)$. Since $(0, 2\pi) \in S(r_2)$, L can go down along $S(r_2)$ to the lowest intersection point $(\theta_1, \eta_1) \in S(r_1) \cap S(r_2) \cap W^*$ on this line segment of $S(r_2)$. By (3) of Lemma 3.3, we have $\eta_1 \leq d_2(r_1, r_2)$. By our assumption, we have $\eta_1 < 2\pi - d_1(r_2, r_2)$.

Again by Lemma 3.3, (θ_1, η_1) is not the highest intersection point in the component of $S(r_1) \cap W^*$ which contains it. Thus L can go along $S(r_1)$ to the highest intersection point. Notice that this point is still in W^* . After that, L again goes along $S(r_2)$ to the lowest intersection point. Repeat this procedure until L hits the line $\theta = \pi$. Then L goes along $\theta = \pi$ to the point (π, π) . We have defined L(t) for $t \in [0, 1]$. Reflecting along $(0, \pi)$, we can define L(t) for $t \in [-1, 0]$.




FIGURE 1. L (left) and \widehat{L} (right) when $r_1 = -3, r_2 = 4$

(2) Suppose r_1, r_2 are of the same sign. We do the case $2 < r_1 < r_2$ and the other case is similar. Set $L(0) = (0, \pi)$ and let L goes along $\theta = 0$ to $(0, 2\pi)$. Then L moves down alone $S(r_1)$ to the lowest intersection point in W^* . After that L moves along $S(r_2)$ to the highest intersection point. The difference from case (1) is that we repeat this procedure until L intersects the line segment $l \subset S(r_1)$ which passes through $(\pi, 0)$. It is easy to see that this happens before L hits $\theta = \pi$. Then L goes along l to $(\pi, 0)$ and then goes along $\theta = \pi$ to (π, π) . By reflecting along the point $(0, \pi)$, we define L(t) for all $t \in [-1, 1]$.

We denote the image of L by $Im(L) \subset W$. In both cases, we have $Im(L) \subset S(r_1) \cup S(r_2) \cup \{\theta = k\pi\}$. Thus $R_K^* \cap Im(L) = \emptyset$. The image of L intersects the line $\eta = 0$ and $\eta = 2\pi$ at $(0,0), (0,2\pi)$ in case (1) and at $(0,0), (0,2\pi), (\pi,0), (-\pi,2\pi)$ in case (2). We need to do small modification around these points. Take the point $(0,2\pi)$ for example. We choose a small neighborhood U of $(0,2\pi)$ and remove $Im(L) \cap U$. Then we replace it




FIGURE 2. L (left) and \widehat{L} (right) when $r_1 = \frac{7}{3}, r_2 = 5$

with a short horizontal line segment $\eta = 2\pi - \varepsilon$. By 5) of Lemma 2.8, after doing this modification, we still get a map $\widehat{L}: [-1,1] \to W^*$ such that $Im(\widehat{L}) \cap R_K = \emptyset$. Moreover, $Im(\widehat{L})$ is symmetric under the reflection about $(0,\pi)$. Suppose $\widehat{L}(t) = (\theta(t),\eta(t))$. By the compactness of R_K , there exists a small neighborhood N of $Im(\widehat{L})$ such that $N \cap R_K = \emptyset$. In case (1), " \widehat{L} goes forward", which means that $\theta(t) \geq \theta(t')$ if $t \geq t'$. Since $(0,\pi), (\pm \pi,\pi) \in Im(\widehat{L})$ and $Im(\widehat{L})$ is symmetric under the reflection of $(0,\pi)$, there exists a smooth odd function g_2 with period 2π such that the graph $\{\eta = g_2(\theta)\}$ is contained in $N + (0, -\pi)$. Thus $R_K + (0, -\pi)$ does not intersect the graph of g_2 . We can choose an even, 2π -periodic function f_2 such that $f'_2 = -g_2$. Set $f_1 \equiv 0$ and use Lemma 2.10. We see that \widehat{CS} has no critical point, which contradicts Theorem 2.1.

In case (2), \widehat{L} does not always go forward and our argument needs to be modified. Take the case $2 < r_1 < r_2$ for example (see Figure 3). By the construction of \widehat{L} , there exists a small $\epsilon > 0$ such that $Im(\widehat{L})$ is contained in the region $\epsilon < \eta < 2\pi - \epsilon$. Choose a number $r_0 \in (r_1, r_2)$. There exist an odd, periodic- 2π function $\theta = g_1(\eta)$ such that $g_1(\eta) = \frac{\eta}{r_0}, \forall \eta \in [\epsilon, 2\pi - \epsilon]$. The image of \widehat{L} only consists of the following 4 types of segments:

- i) horizontal line that goes forward,
- ii) going down line of slope $-r_1$,
- iii) going up line of slope $-r_2$,
- iv) going up line of slope $+\infty$.

Therefore, it is not difficult to see that $Im(\widehat{L}) + (0, -\pi) + (g_1, *)$ is a broken line that goes forward (it's just a shearing of $Im(\widehat{L}) + (0, -\pi)$). Thus we can find an odd, 2π -periodic function g_2 such that the graph $\{\eta = g_2(\theta)\}$ is contained in $N + (0, -\pi) + (g_1, *)$. We see that $R_K + (0, -\pi) + (g_1, *)$ does not intersect the graph $\{\eta = g_2(\theta)\}$. We can find even,




FIGURE 3. The iamge of $\widehat{L} + (0, -\pi)$ (left) and $\widehat{L} + (0, -\pi) + (g_1, *)$ (right) when $r_1 = \frac{7}{3}, r_2 = 5$

 2π -periodic function f_1, f_2 such that $f'_1 = g_1, f'_2 = -g_2$. Using Lemma 2.10 and Theorem 2.1, we get the contradiction again.

The SO(3)-cyclic case can be proved similarly by considering $\widehat{S}(r_i)$ instead of $S(r_i)$. \square

Remark 3.4. Actually, we have proved that if r_1, r_2 are both SU(2)-cyclic, then $d_1(r_1, r_2) + d_2(r_1, r_2) \ge 2\pi$. When p_i is odd, this gives the conclusions of Corollary 1.10.

Corollary 1.10, Corollary 1.11 and Corollary 1.17 are easy to prove using the main theorem.

3.2. Relation with the Alexander polynomial. In this subsection, we will give some relations between the SU(2)-cyclic surgeries and the Alexander polynomial and prove Corollary 1.12.

Suppose $d_1(r_1, r_2) + d_2(r_1, r_2) = 2\pi$ (for example $r_1 = -r_2 = 2k$) and r_1, r_2 are both SU(2)-cyclic. Let's try to repeat the argument as before. We do the case $r_2 < 0 < r_1$ and the other cases are similar. Consider $S(r_i) \subset (\mathbb{R}/2\pi\mathbb{Z}) \bigoplus (\mathbb{R}/2\pi\mathbb{Z})$ (i = 1, 2), then $R_K^* \cap S(r_i) = \emptyset$. We now construct $L : [-1, 1] \to W$. Set $L(0) = (0, \pi)$ and L goes upwards along $\theta = 0$ to $(0, 2\pi)$. Then L goes down along $S(r_2)$ the the lowest intersection point $(\theta_1, \eta_1) \in S(r_1) \cap S(r_2) \cap W$. After that, L goes up along $S(r_1)$ to the highest intersection point $(\theta_2, \eta_2) \in S(r_1) \cap S(r_2) \cap W$. As shown in Figure 4, it is possible that the lowest intersection point in $(\theta_2, \eta_2) \in S(r_1) \cap S(r_2) \cap W^*$ is also the highest one. Thus we can only work in W instead of W^* . This is different from the case when $d_1(r_1, r_2) + d_2(r_1, r_2) < 2\pi$. We repeat this procedure and get $L : [-1,1] \to W$. Then we need to modify L to \widehat{L} whose image is contained in W^* . The trouble appears: L may contain some points like $(\theta_0, 0)$ or $(\theta_0, 2\pi)$ with $\theta_0 \neq 0$ or $\pm \pi$. In general, we don't have the result like 5) of Lemma 2.8 which allows us to modify L near these points without intersecting R_K .




FIGURE 4. When $r_1 = 4, r_2 = -4$, we can modify L near $(\theta_0, 0)$ in the left picture but we can't modify L in the right picture.

We just do the $(\theta_0,0)$ case and the $(\theta_0,2\pi)$ case is similar. Suppose that we can choose a small neighborhood U of $(\theta_0,0)$ such that $R_K^* \cap U = \emptyset$. We just replace $Im(L) \cap U$ by some short, horizontal line $l \subset U \cap W^*$. If we can do this for every point in $Im(L) \cap (W \setminus W^*)$, we can construct \widehat{L} and get the contradiction as before. If we can't do this for some point $(\theta_0,0) \in S(r_1)$, then there exist a sequence $(\theta_n,\eta_n) \in R_K^*$ converging to $(\theta_0,0)$ as $n \to \infty$. Each (θ_n,η_n) gives an irreducible representation $\rho_n:\pi_1(S^3-N(K))\to SU(2)$. It is easy to see that these representations are also irreducible as $SL(2,\mathbb{C})$ representations. By the compactness of SU(2) representation variety, ρ_n converge to some ρ_0 after taking a subsequence. We will get $(\theta_0,0) \in S(r_1)$ if we restrict ρ_0 to the boundary. Recall that we have a representation $\pi_1(S^3-K)\to \pm 1\to SU(2)$ such that m is mapped to -1. After multiplying ρ_0 by this representation if necessary, we get a representation of ρ'_0 $\pi_1(S^3-N(K))$ such that $\rho'_0(p_1m+q_1l)=1$. Since r_1 is an SU(2)-cyclic surgery, this representation must be cyclic. In particular, this implies that ρ_0 is cyclic. Thus we get a sequence of irreducible $SL(2,\mathbb{C})$ representations of $\pi_1(S^3-N(K))$ converging to a reducible $SL(2,\mathbb{C})$ representation ρ_0 with $\rho_0(m)=\begin{pmatrix} e^{i\theta_0} & 0 \\ 0 & e^{-i\theta_0} \end{pmatrix}$.

We apply the following proposition in [12]:

Proposition 3.5 ([12]). Let M be the complement of a knot K in a homology 3-sphere. Suppose that ρ is a reducible representation of $\pi_1(M)$ such that the character of ρ lies on a component of $\chi(M)$ which contains the character of an irreducible representation. Then $\rho(m)$ has an eigenvalue whose square is a root of Δ_K (the Alexander polynomial of K).

Using this theorem, we see that $e^{2i\theta_0}$ is a root of Δ_K . Since $(\theta_0, 0) \in S(r_1)$, we see that Δ_K has a root which is a p_1 -th root of unity for odd p_1 and $\frac{p_1}{2}$ -th root of unity for even p_1 .

By considering the intersection point $(\theta_0, 2\pi) \in Im(L) \cap \{\eta = 2\pi\}$, we can get the same conclusion for p_2 . In particular, we get the following:

Proposition 3.6. Suppose that $r_1 = \frac{p_1}{q_1}$, $r_2 = \frac{p_2}{q_2}$ are two SU(2)-cyclic surgeries with $d_1(r_1, r_2) + d_2(r_1, r_2) = 2\pi$ and p_1, p_2 even, then the Alexander polynomial of K has a root which is either a $\frac{p_1}{2}$ -th or a $\frac{p_2}{2}$ -th root of unity.

Notice that if K is amphichiral, then the r-surgery is SU(2)-cyclic implies that the -r-surgery is also SU(2)-cyclic. By Corollary 1.11, we get r is an even integer and $d_1(r,-r)+d_2(r,-r)=2\pi$. Therefore, Corollary 1.12 is a straightforward consequence of the proposition above.

References

- [1] A.Floer, *Instanton homology and Dehn surgery*. The Floer memorial volume, Progr. Math., vol.133, Birkhauser, Basel, 1995,pp. 195-256 no.3,77-97.
- [2] D.Gabai, Foliation and the topology of 3-manifolds. J. Differential Geom.26(1987), no.3, 479-536.
- [3] Y.Ni, Link Floer homology detects the Thurston norm. Geom. Topol. 13, No.5, 2991-3019 (2009).
- [4] N.M.Dunfield, Private communication.
- [5] P.B.Kronheimer, T.S.Mrowka, Dehn surgery, the fundamental group and SU(2). Math. Res. Lett. 11 (2004), no. 5-6, 741C754.
- [6] P.B.Kronheimer, T.S.Mrowka, Embedded surfaces and the structure of Donaldson's polynomial invariants, J. Differential Geom. 41(1995), no.3,573-734.
- [7] P.B.Kronheimer, T.S.Mrowka, Witten's conjeture and Property P, Geom. Topol. 8(2004) 295-310.
- [8] M Culler, CMcA Gordon, J Luecke, PB Shalen, Dehn surgery on knots, Ann. of Math. 125 (1987) 237C300
- [9] P.J.Braam and S.K.Donaldson, Floer's work on instanton homology, knots and surgery. The Floer memorial volume, Progr. Math., vol.133, Birkhauser, Basel, 1995,pp. 195-256
- [10] P.M.N.Feehan and T.G.Leness, A general SO(3)-monopole corbordism formula relating Donaldson and Seiberg-Witten invariants. Preprint, arXiv:math.DG/0203047,2003
- [11] C.H.Taubes, The Seiberg-Witten invariants and symplectic forms, Math. Res. Lett.1(1994),no.6,809-822.
- [12] D.Cooper, M.Culler, H.Gillet, D.Long and P.Shalen, Plane curves associated to character varieties of 3-manifolds. Invent. Math. 118(1994) 47-84.
- [13] J. Lin, The A-polynomial and holonomy perturbations. Preprint, arXiv:1304.7232.
- [14] C McA Gordon, J Luecke, Knots are determined by their complements, J.Amer. Math. Soc. 2 (1989) 371C415
- [15] Y. Eliashberg, Few remarks about symplectic filling. Geom. Topol. 8 (2004), 277C293.
- [16] Y. Eliashberg and W.Thurston, Confoliations. University Lecture Series, no. 13, American Mathematical Society, (1998).

Jianfeng Lin

Department of Mathematics, University of California Los Angeles, Los Angeles, US juliuslin@math.ucla.edu