

# The Evolution of Beliefs over Signed Social Networks

Guodong Shi, Alexandre Proutiere,  
Mikael Johansson, John S. Baras, and Karl Henrik Johansson

## Abstract

We study the evolution of opinions (or beliefs) over a social network modeled as a signed graph. The sign attached to an edge in this graph characterizes whether the corresponding individuals or end nodes are friends (positive links) or enemies (negative links). Pairs of nodes are randomly selected to interact over time, and when two nodes interact, each of them updates its opinion based on the opinion of the other node and the sign of the corresponding link. This model generalizes DeGroot model to account for negative links: when two enemies interact, their opinions go in opposite directions. We provide conditions for convergence and divergence in expectation, in mean-square, and in almost sure sense, and exhibit phase transition phenomena for these notions of convergence depending on the parameters of the opinion update model and on the structure of the underlying graph. We establish a *no-survivor* theorem, stating that the difference in opinions of any two nodes diverges whenever opinions in the network diverge as a whole. We also prove a *live-or-die* lemma, indicating that almost surely, the opinions either converge to an agreement or diverge. Finally, we extend our analysis to cases where opinions have hard lower and upper limits. In these cases, we study when and how opinions may become asymptotically clustered to the belief boundaries, and highlight the crucial influence of (strong or weak) structural balance of the underlying network on this clustering phenomenon.

**Keywords:** opinion dynamics, signed graph, social networks, opinion clustering

## 1 Introduction

### 1.1 Motivation

We all form opinions about economical, political, and social events that take place in society. These opinions can be binary (e.g., whether one supports a candidate in an election or not) or continuous (to what degree one expects a prosperous future economy). Our opinions are revised when we interact with each other over various social networks. Characterizing the evolution of opinions, and understanding the dynamic and asymptotic behavior of the social belief, are fundamental challenges in the theoretical study of social networks.

Building a good model on how individuals interact and influence each other is essential for studying opinion dynamics. In interaction models, it is natural that a trusted friend should have a different influence on the opinion formation than a dubious stranger. The observation that sentiment influences opinions can be traced back to the 1940's when Heider (1946) introduced the theory of signed social networks, where each interaction link in the social network is associated with a sign (positive or negative) indicating whether two individuals are friends or enemies. Efforts to understand the properties of signed social networks have led to the development of structural balance theory, with seminal contributions by Cartwright and Harary (1956) and Davis (1963, 1967). A fundamental insight from these studies, formalized in Harary's theorem Harary (1953), is that local structural properties imply hard global constraints on the social network formation.

In this paper, we attempt to model the evolution of opinions in signed social networks when local hostile or antagonistic relations influence the global social belief. The relative strengths and structures of positive and negative relations are shown to have an essential effect on opinion convergence. In some cases, tight conditions for convergence and divergence can be established.

## 1.2 Related Work

The concept of signed social networks was introduced by Heider (1946). His objective was to formally distinguish between friendly (positive) and hostile (negative) relationships. The notion of structural balance was introduced to understand local interactions, and formalize intricate local scenarios (e.g., *two of my friends are enemies*). A number of classical results on social balance was established by Harary (1953), Cartwright and Harary (1956), Davis (1963, 1967), who derived critical conditions on the global structure of the social network which ensure structural balance. Social balance theory has since become an important topic in the study of social networks. On one hand, efforts are made to characterize and compute the degree of balance for real-world large social networks, e.g. Facchetti et al. (2011). On the other hand, dynamical models are proposed for the signs of social links with the aim of describing stable equilibria or establishing asymptotic convergence for the sign patterns, e.g., Galam (1996) (where a signed structure was introduced as a revised Ising model of political coalitions, where two competing world coalitions were shown to have one unique stable formation), Macy et al. (2003) (who verified convergence to structural balances numerically for a Hopfield model), and Marvel et al. (2011) (where a continuous-time dynamical model for the link signs was proposed under which convergence to structural balance was proven).

Opinion dynamics is another long-standing topic in the study of social networks, see Jackson (2008) and Easley and Kleinberg (2010) for recent textbooks. Following the survey Acemoglu and Ozdaglar

(2011), we classify opinion evolution models into Bayesian and non-Bayesian updating rules. Their main difference lies in whether each node has access to and acts according to a global model or not. We refer to Banerjee (1992), Bikhchandani et al. (1992) and, more recent work Acemoglu et al. (2011) for Bayesian opinion dynamics. In non-Bayesian models, nodes follow simple local updating strategies. DeGroot’s model (DeGroot (1974)) is a classical non-Bayesian model of opinion dynamics, where each node updates its belief as a convex combination of its neighbors’ beliefs, e.g., DeMarzo et al. (2003), Golub and Jackson (2010), Jadbabaie et al. (2012). Note that DeGroot’s model relates to averaging consensus algorithms, e.g., Tsitsiklis (1984), Xiao and Boyd (2004), Boyd et al. (2006), Tahbaz-Salehi and Jadbabaie (2008), Fagnani and Zampieri (2008), Touri and Nedić (2011), Matei et al. (2013). Non-consensus asymptotic behaviors, e.g., clustering, disagreement, and polarization, have been investigated for linear or nonlinear variations of DeGroot-type update rules, Krause (1997), Blondel et al. (2009, 2010), Dandekar et al. (2013), Shi et al. (2013), Li et al. (2013). Various models from statistical physics have also been applied to study social opinion dynamics, please refer to Castellano et al. (2009) for a survey.

The influence of misbehaving nodes in social networks have been studied only to some extent. For instance, in Acemoglu et al. (2010), a model of the spread of misinformation in large societies was discussed. There, some individuals are *forceful*, meaning that they influence the beliefs of some of the other individuals they meet, but do not change their own opinions. In Acemoglu et al. (2013), the authors studied the propagation of opinion disagreement under DeGroot’s model, when some nodes stick to their initial beliefs during the entire evolution. This idea was extended to binary opinion dynamics under the voter model in Yildiz et al. (2013). In Altafini (2012, 2013), the author proposed a linear model for belief dynamics over signed graphs. In Altafini (2013), it was shown that a bipartite agreement, i.e., clustering of opinions, is reached as long as the signed social graph is strongly balanced in the sense of the classical structural balance theory (Cartwright and Harary (1956)), which presents an important link between opinion dynamics and structure balance. However, in the model studied in Altafini (2012, 2013), all beliefs converge to a common value, equal to zero, if the graph is not strongly balanced. This behavior seems to be difficult to interpret and justify from real-world observations. A game-theoretical approach for studying the interplay between good and bad players in collaborative networks was introduced in Theodorakopoulos and Baras (2008).

### 1.3 Contribution

We propose and analyze a new model for belief dynamics over signed social networks. Nodes randomly execute pairwise interactions to update their beliefs. In case of a positive link (representing that the two

interacting nodes are friends), the update follows DeGroot’s update rule which drives the two beliefs closer to each other. On the contrary, in case of a negative link (i.e., when the two nodes are enemies), the update increases the difference between the two beliefs. Thus, two opposite types of opinion updates are defined, and the beliefs are driven not only by random node interactions but also by the type of relationship of the interacting nodes. Under this simple attraction–repulsion model for opinions on signed social networks, we establish a number of fundamental results on belief convergence and divergence, and study the impact of the parameters of the update rules and of the network structure on the belief dynamics.

Using classical spectral methods, we derive conditions for mean and mean-square convergence and divergence of beliefs. We establish phase transition phenomena for these notions of convergence, and study how the thresholds depend on the parameters of the opinion update model and on the structure of the underlying graph. We derive phase transition conditions for almost sure convergence and divergence of beliefs. The proofs are based on what we call the *Triangle lemma*, which characterizes the evolution of the beliefs held by three different nodes. We utilize probabilistic tools such as the Borel-Cantelli lemma, the Martingale convergence theorems, the strong law of large numbers, and sample-path arguments.

We establish two counter-intuitive results about the way beliefs evolve: (i) a *no-survivor* theorem which states that the difference between opinions of any two nodes tends to infinity almost surely (along a subsequence of instants) whenever the difference between the maximum and the minimum beliefs in the network tends to infinity (along a subsequence of instants); (ii) a *live-or-die* lemma which demonstrates that almost surely, the opinions either converge to an agreement or diverge. We also show that networks whose positive component includes an hypercube are (essentially, the only) robust networks in the sense that almost sure convergence of beliefs holds irrespective of the number of negative links, their positions in the network, and the strength of the negative update.

The considered model is extended to cases where updates may be asymmetric (in the sense that when two nodes interact, only one of them updates its belief), and where beliefs have hard lower and upper constraints. The latter boundedness constraint adds slight nonlinearity to the belief evolution. It turns out in this case that the classical social network structural balance theory plays a fundamental role in determining the asymptotic formation of opinions:

- If the social network is structurally balanced (strongly balanced, or complete and weakly balanced), i.e., the network can be divided into subgroups with positive links inside each subgroup and negative links among different subgroups, then almost surely, the beliefs within the same subgroup will be clustered to one of the belief boundaries, when the strength of the negative updates is sufficiently large.

- In the absence of structural balance, and if the positive graph of the social network is connected, then almost surely, the belief of each node oscillates between the lower and upper bounds and touches the two belief boundaries an infinite number of times.

For balanced social networks, the boundary clustering results are established based on the almost sure happening of suitable *separation* events, i.e., the node beliefs for a subgroup become group polarized (either larger or smaller than the remaining nodes' beliefs). From this argument such events tend to happen more easily in the presence of small subgroups. As a result, small subgroups contribute to faster clustering of the social beliefs, which is consistent with the study of minority influence in social psychology Nemeth (1986), Clark and Maass (1990) suggesting that consistent minorities can substantially influence opinions. For unbalanced social networks, the established opinion oscillation contributes to a new type of belief formation which complements polarization, disagreement, and consensus Dandekar et al. (2013).

## 1.4 Paper Organization

In Section 2, we present the signed social network model, specify the dynamics along positive and negative links, and define the problem of interest. Section 3 focuses on the mean and mean-square convergence and divergence analysis, and Section 4 considers convergence and divergence in the almost sure sense. In Section 5, we study a model with upper and lower belief bounds and asymmetric updates. It is shown how structural balance determines the clustering of opinions. Finally concluding remarks are given in Section 6.

## Notation and Terminology

An undirected graph is denoted by  $G = (V, E)$ . Here  $V = \{1, \dots, n\}$  is a finite set of vertices (nodes). Each element in  $E$  is an unordered pair of two distinct nodes in  $V$ , called an edge. The edge between nodes  $i, j \in V$  is denoted by  $\{i, j\}$ . Let  $V_* \subseteq V$  be a subset of nodes. The *induced* graph of  $V_*$  on  $G$ , denoted  $G_{V_*}$ , is the graph  $(V_*, E_{V_*})$  with  $\{u, v\} \in E_{V_*}$ ,  $u, v \in V_*$  if and only if  $\{u, v\} \in E$ . A path in  $G$  with length  $k$  is a sequence of distinct nodes,  $v_1 v_2 \dots v_{k+1}$ , such that  $\{v_m, v_{m+1}\} \in E$ ,  $m = 1, \dots, k$ . The length of a shortest path between two nodes  $i$  and  $j$  is called the distance between the nodes, denoted  $d(i, j)$ . The greatest length of all shortest paths is called the diameter of the graph, denoted  $\text{diam}(G)$ . The *degree* matrix of  $G$ , denoted  $D(G)$ , is the diagonal matrix  $\text{diag}(d_1, \dots, d_n)$  with  $d_i$  denoting the number of nodes sharing an edge with  $i$ ,  $i \in V$ . The *adjacency* matrix  $A(G)$  is the symmetric  $n \times n$  matrix such that  $[A(G)]_{ij} = 1$  if  $\{i, j\} \in E$  and  $[A(G)]_{ij} = 0$  otherwise. The matrix  $L(G) := D(G) - A(G)$  is called the *Laplacian* of  $G$ . Two graphs containing the same number of vertices are called *isomorphic* if they are identical subject to a permutation of vertex labels.

All vectors are column vectors and denoted by lower case letters. Matrices are denoted with upper case letters. Given a matrix  $M$ ,  $M'$  denotes its transpose and  $M^k$  denotes the  $k$ -th power of  $M$  when it is a square matrix. The  $ij$ -entry of a matrix  $M$  is denoted  $[M]_{ij}$ . Given a matrix  $M \in \mathbb{R}^{mn}$ , the vectorization of  $M$ , denoted by  $\mathbf{vec}(M)$ , is the  $mn \times 1$  column vector  $([M]_{11}, \dots, [M]_{m1}, \dots, [M]_{1n}, \dots, [M]_{mn})'$ . We have  $\mathbf{vec}(ABC) = (C' \otimes A)\mathbf{vec}(B)$  for all real matrices  $A, B, C$  with  $ABC$  well defined. A square matrix  $M$  is called a stochastic matrix if all of its entries are non-negative and the sum of each row of  $M$  equals one. A stochastic matrix  $M$  is doubly stochastic if  $M'$  is also a stochastic matrix. With the universal set prescribed, the complement of a given set  $S$  is denoted  $S^c$ . The orthogonal complement of a subspace  $S$  in a vector space is denoted  $S^\perp$ . Depending on the argument,  $|\cdot|$  stands for the absolute value of a real number, the Euclidean norm of a vector, and the cardinality of a set. Similarly with argument well defined,  $\sigma(\cdot)$  represents the  $\sigma$ -algebra of a random variable (vector), or the spectrum of a matrix. The smallest integer no smaller than a given real number  $a$  is denoted  $\lceil a \rceil$ . We use  $\mathbb{P}(\cdot)$  to denote the probability,  $\mathbb{E}\{\cdot\}$  the expectation,  $\mathbb{V}\{\cdot\}$  the variance of their arguments, respectively.

## 2 Opinion Dynamics over Signed Social Networks

In this section, we present our model of interaction between nodes in a signed social network, and describe the resulting dynamics of the beliefs held by each node.

### 2.1 Signed Social Network and Peer Interactions

We consider a social network with  $n \geq 3$  members, each labeled by a unique integer in  $\{1, 2, \dots, n\}$ . The network is represented by an undirected graph  $G = (\mathbf{V}, \mathbf{E})$  whose node set  $\mathbf{V} = \{1, 2, \dots, n\}$  corresponds to the members and whose edge set  $\mathbf{E}$  describes potential interactions between the members. Each edge in  $\mathbf{E}$  is assigned a unique label, either  $+$  or  $-$ . In classical social network theory, a  $+$  label indicates a friend relation, while a  $-$  label indicates an enemy relation (Heider (1946), Cartwright and Harary (1956)). The graph  $G$  equipped with a sign on each edge is then called a *signed graph*. Let  $\mathbf{E}_{\text{pst}}$  and  $\mathbf{E}_{\text{neg}}$  be the collection of the positive and negative edges, respectively; clearly,  $\mathbf{E}_{\text{pst}} \cap \mathbf{E}_{\text{neg}} = \emptyset$  and  $\mathbf{E}_{\text{pst}} \cup \mathbf{E}_{\text{neg}} = \mathbf{E}$ . We call  $G_{\text{pst}} = (\mathbf{V}, \mathbf{E}_{\text{pst}})$  and  $G_{\text{neg}} = (\mathbf{V}, \mathbf{E}_{\text{neg}})$  the positive and the negative graph, respectively; see Figure 1 for an illustration. Without loss of generality, we adopt the following assumption throughout the paper.

**Assumption 1** *The underlying graph  $G$  is connected, and the negative graph  $G_{\text{neg}}$  is nonempty.*

Actual interactions follow the model introduced in Boyd et al. (2006): each node initiates interactions at the instants of a rate-one Poisson process, and at each of these instants, picks a node at random to




Figure 1: A signed social network.

interact with. Under this model, at a given time, at most one node initiates an interaction. This allows us to order interaction events in time and to focus on modeling the node pair selection at interaction times. The node selection process is characterized by an  $n \times n$  stochastic matrix  $P = [p_{ij}]$  complying with the graph  $\mathbf{G}$ , in the sense that  $p_{ij} > 0$  always implies  $\{i, j\} \in \mathbf{E}$  for  $i \neq j \in \mathbf{V}$ . The  $p_{ij}$  represents the probability that node  $i$  initiates an interaction with node  $j$ . The node pair selection is then performed as follows.

**Definition 1** *At each interaction event  $k \geq 0$ , (i) a node  $i \in \mathbf{V}$  is drawn uniformly at random, i.e., with probability  $1/n$ ; (ii) node  $i$  picks node  $j$  with probability  $p_{ij}$ . In this case, we say that the unordered node pair  $\{i, j\}$  is selected.*

The node pair selection process is assumed to be i.i.d., i.e., the nodes that initiate an interaction and the selected node pairs are identically distributed and independent over  $k \geq 0$ . Formally, the node selection process can be analyzed using the following probability spaces. Let  $(\mathbf{E}, \mathcal{S}, \mu)$  be the probability space, where  $\mathcal{S}$  is the discrete  $\sigma$ -algebra on  $\mathbf{E}$ , and  $\mu$  is the probability measure defined by  $\mu(\{i, j\}) = \frac{p_{ij} + p_{ji}}{n}$  for all  $\{i, j\} \in \mathbf{E}$ . The node selection process can then be seen as a random event in the product probability space  $(\Omega, \mathcal{F}, \mathbb{P})$ , where  $\Omega = \mathbf{E}^{\mathbb{N}} = \{\omega = (\omega_0, \omega_1, \dots) : \forall k, \omega_k \in \mathbf{E}\}$ ,  $\mathcal{F} = \mathcal{S}^{\mathbb{N}}$ , and  $\mathbb{P}$  is the product probability measure (uniquely) defined by: for any finite subset  $K \subset \mathbb{N}$ ,  $\mathbb{P}((\omega_k)_{k \in K}) = \prod_{k \in K} \mu(\omega_k)$  for any  $(\omega_k)_{k \in K} \in \mathbf{E}^{|K|}$ . For any  $k \in \mathbb{N}$ , we define the coordinate mapping  $G_k : \Omega \rightarrow \mathbf{E}$  by  $G_k(\omega) = \omega_k$ , for all  $\omega \in \Omega$  (note that  $\mathbb{P}(G_k = \omega_k) = \mu(\omega_k)$ ), and we refer to  $(G_k, k = 0, 1, \dots)$  as the *node pair selection process*. We further refer to  $\mathcal{F}_k = \sigma(G_0, \dots, G_k)$  as the  $\sigma$ -algebra capturing the  $(k + 1)$  first interactions of the selection process.

## 2.2 Positive and Negative Dynamics

Each node maintains a scalar real-valued *opinion*, or *belief*, which it updates whenever it interacts with other nodes. We let  $x(k) \in \mathbb{R}^n$  denote the vector of the beliefs held by nodes at the interaction event  $k$ .

The belief update depends on the relationship between the interacting nodes. Suppose that node pair  $\{i, j\}$  is selected at time  $k$ . The nodes that are not selected keep their beliefs unchanged, whereas the beliefs held by nodes  $i$  and  $j$  are updated as follows:

- (*Positive Update*) If  $\{i, j\} \in \mathbf{E}_{\text{pst}}$ , either node  $m \in \{i, j\}$  updates its belief as

$$x_m(k+1) = x_m(k) + \alpha(x_{-m}(k) - x_m(k)) = (1 - \alpha)x_m(k) + \alpha x_{-m}(k), \quad (1)$$

where  $-m \in \{i, j\} \setminus \{m\}$  and  $0 \leq \alpha \leq 1$ .

- (*Negative Update*) If  $\{i, j\} \in \mathbf{E}_{\text{neg}}$ , either node  $m \in \{i, j\}$  updates its belief as

$$x_m(k+1) = x_m(k) - \beta(x_{-m}(k) - x_m(k)) = (1 + \beta)x_m(k) - \beta x_{-m}(k), \quad (2)$$

where  $\beta \geq 0$ .

The positive update is consistent with the classical DeGroot model (DeGroot (1974)), where each node iteratively updates its belief as a convex combination of the previous beliefs of itself and of the neighbor with which it interacts. This update naturally reflects *trustful* or *cooperative* relationships. It is sometimes referred to as naïve learning in social networks, under which *wisdom* can be held by the crowds (Golub and Jackson (2010)). The positive update tends to drive node beliefs closer to each other and can be thought of as the attraction of the beliefs.

The dynamics on the negative edges, on the other hand, is not yet universally agreed upon in the literature. Considerable efforts have been made to characterize these *mistrustful* or *antagonistic* relationships, which has led to a number of insightful models, e.g., Acemoglu et al. (2010, 2013), Altafini (2012, 2013). Our negative update rule enforces belief differences between interacting nodes, and is the *opposite* of the attraction of beliefs represented by the positive update.

## 2.3 Model Rationale

### 2.3.1 Relation to Non-Bayesian Rules

Our underlying signed graph is a *prescribed* world with fixed trust or mistrustful relations where nodes do not switch their relations. Two nodes holding the same opinion can be enemies, and vice versa. This



contrasts Krause’s model, where trustful relations are state-dependent and nodes only interact with nodes which hold similar opinions, i.e., whose beliefs are within a given distance.

In our model, the signed graph classifies the social interactions into two categories, positive and negative, each with its own type of dynamics. Studies of stubborn agents in social network Acemoglu et al. (2013), Yildiz et al. (2013) also classify nodes into two categories, but stubborn agents do not account for the opinion of its neighbors. Our model is more similar to the one introduced by Altafini in Altafini (2013), where the author proposed a different update rule for two nodes sharing a negative link. The model in Altafini (2013) is written in continuous time (beliefs evolve along some ODE), but its corresponding discrete-time update across a negative link  $\{i, j\} \in \mathbf{E}_{\text{neg}}$  is:

$$x_m(k+1) = x_m(k) - \beta(x_{-m}(k) + x_m(k)) = (1 - \beta)x_m(k) - \beta x_{-m}(k), \quad m \in \{i, j\}, \quad (3)$$

where  $\beta \in (0, 1)$  represents the negative strength. This update rule admits the following interpretations:

- Node  $i$  attempts to *trick* her negative neighbors  $j$ , by flipping the sign of her true belief (i.e.,  $x_i(k)$  to  $-x_i(k)$ ) before revealing it to  $j$ ;
- Node  $i$  recognizes  $j$  as her negative neighbor and upon observing  $j$ ’s true belief,  $x_j(k)$ , she tries to get closer to the opposite view of  $j$  since  $x_i(k+1)$  is a convex combination of  $x_i(k)$  and  $-x_j(k)$ .

In both of the two interpretations of the Altafini model, the belief origin must be of some particular significance in the nodes’ belief space. This is not the case for our model, where the positive/negative dynamics describe *choices* intended to *keeping close to friends* and *keeping distance from enemies*. When nodes  $i$  and  $j$  perform a negative update in our model, if  $x_i(k) > x_j(k)$  then  $x_i(k+1) > x_i(k)$  and if  $x_i(k) < x_j(k)$  then  $x_i(k+1) < x_i(k)$ . That is, in either case, the node’s updated opinion is in a direction away from the opinion of the interacting node (i.e., nodes make an effort to “keep distance from the enemies” and do not assign any special meaning to the belief origin).

**Remark 1** *The Altafini model Altafini (2013) and the current work are intended for building theories to opinion dynamics over signed social networks. Indeed nontrivial efforts have been made to model the dynamics of signed social networks themselves Galam (1996), Macy et al. (2003), Marvel et al. (2011). It is intriguing to ask how opinions and social networks shape each other in the presence of trustful/mistrustful relations, where fundamental difficulty arises in how to properly model such couplings as well as the challenges brought by the couplings.*

### 2.3.2 Relation to Bayesian Rules

Bayesian opinion dynamics assume that there is a global *model* of the world and individuals aim to realize asymptotic learning of the underlying world Banerjee (1992), Bikhchandani et al. (1992), Acemoglu et al. (2011). It has been shown that DeGroot update can also serve as a naive learning approach as long as the network somehow contains no dictators Golub and Jackson (2010).

We argue here our model corresponds to the situation where nodes naively follow the code of keeping distance with enemies and keeping close to friends, rather than having interest in some underlying world model. Our definition of the negative dynamics becomes quite natural if one views the DeGroot type of update as the approach of keeping close to friends. This simple yet informative model leads to a number of nontrivial belief formations in terms of convergence or divergence for unconstrained evolution, consensus, clustering, or oscillation under boundedness constraint.

We note that it is an interesting open challenge to find a proper model for Bayesian learning over signed social networks, since nodes must learn in the presence of *negative* interactions, on the one hand, and may try to prevent their enemies from asymptotic learning, on the other.

## 3 Mean and Mean-square Convergence/Divergence

Let  $x(k) = (x_1(k) \dots x_n(k))'$ ,  $k = 0, 1, \dots$  be the (random) vector of beliefs at time  $k$  resulting from the node interactions. The initial beliefs  $x(0)$ , also denoted as  $x^0$ , is assumed to be deterministic. In this section, we investigate the mean and mean-square evolution of the beliefs for the considered signed social network. We introduce the following definition.

**Definition 2** (i) *Belief convergence is achieved in expectation if  $\lim_{k \rightarrow \infty} \mathbb{E}\{x_i(k) - x_j(k)\} = 0$  for all  $i$  and  $j$ ; in mean square if  $\lim_{k \rightarrow \infty} \mathbb{E}\{(x_i(k) - x_j(k))^2\} = 0$  for all  $i$  and  $j$ .*

(ii) *Belief divergence is achieved in expectation if  $\limsup_{k \rightarrow \infty} \max_{i,j} |\mathbb{E}\{x_i(k) - x_j(k)\}| = \infty$ ; in mean square if  $\limsup_{k \rightarrow \infty} \max_{i,j} \mathbb{E}\{(x_i(k) - x_j(k))^2\} = \infty$ .*

The belief dynamics as described above can be written as:

$$x(k+1) = W(k)x(k), \quad (4)$$

where  $W(k)$ ,  $k = 0, 1, \dots$  are i.i.d. random matrices satisfying

$$\begin{aligned} \mathbb{P}\left(W(k) = W_{ij}^+ := I - \alpha(e_i - e_j)(e_i - e_j)'\right) &= \frac{p_{ij} + p_{ji}}{n}, \quad \{i, j\} \in E_{\text{pst}}, \\ \mathbb{P}\left(W(k) = W_{ij}^- := I + \beta(e_i - e_j)(e_i - e_j)'\right) &= \frac{p_{ij} + p_{ji}}{n}, \quad \{i, j\} \in E_{\text{neg}}, \end{aligned} \quad (5)$$

and  $e_m = (0 \dots 0 \ 1 \ 0 \dots 0)'$  is the  $n$ -dimensional unit vector whose  $m$ -th component is 1. In this section, we use spectral properties of the linear system (4) to study convergence and divergence in mean and mean-square. Our results can be seen as extensions of existing convergence results on deterministic consensus algorithms, e.g., Xiao and Boyd (2004).

### 3.1 Convergence in Mean

We first provide conditions for convergence and divergence in mean. We then exploit these conditions to establish the existence of a phase transition for convergence when the negative update parameter  $\beta$  increases. These results are illustrated at the end of this subsection. For technical reasons we adopt the following assumption in this subsection.

**Assumption 2** *There holds either (i)  $p_{ii} \geq 1/2$  for all  $i \in \mathcal{V}$ , or (ii)  $P = [p_{ij}]$  is doubly stochastic with  $n \geq 4$ .*

Generalization to the case when Assumption 2 does not hold is essentially straightforward but under a bit more careful treatment.

#### 3.1.1 Convergence/Divergence Conditions

Denote  $P^\dagger = (P + P')/n$ . We write  $P^\dagger = P_{\text{pst}}^\dagger + P_{\text{neg}}^\dagger$ , where  $P_{\text{pst}}^\dagger$  and  $P_{\text{neg}}^\dagger$  correspond to the positive and negative graphs, respectively. Specifically,  $[P_{\text{pst}}^\dagger]_{ij} = [P^\dagger]_{ij}$  if  $\{i, j\} \in \mathbf{E}_{\text{pst}}$  and  $[P_{\text{pst}}^\dagger]_{ij} = 0$  otherwise, while  $[P_{\text{neg}}^\dagger]_{ij} = [P^\dagger]_{ij}$  if  $\{i, j\} \in \mathbf{E}_{\text{neg}}$  and  $[P_{\text{neg}}^\dagger]_{ij} = 0$  otherwise. We further introduce the degree matrix  $D_{\text{pst}}^\dagger = \text{diag}(d_1^+ \dots d_n^+)$  of the positive graph, where  $d_i^+ = \sum_{j=1, j \neq i}^n [P_{\text{pst}}^\dagger]_{ij}$ . Similarly, the degree matrix of the negative graph is defined as  $D_{\text{neg}}^\dagger = \text{diag}(d_1^- \dots d_n^-)$  with  $d_i^- = \sum_{j=1, j \neq i}^n [P_{\text{neg}}^\dagger]_{ij}$ . Then  $L_{\text{pst}}^\dagger = D_{\text{pst}}^\dagger - P_{\text{pst}}^\dagger$  and  $L_{\text{neg}}^\dagger = D_{\text{neg}}^\dagger - P_{\text{neg}}^\dagger$  represent the (weighted) Laplacian matrices of the positive graph  $\mathbf{G}_{\text{pst}}$  and negative graph  $\mathbf{G}_{\text{neg}}$ , respectively. It can be easily deduced from (5) that

$$\mathbb{E}\{W(k)\} = I - \alpha L_{\text{pst}}^\dagger + \beta L_{\text{neg}}^\dagger. \quad (6)$$

Clearly,  $\mathbf{1}'\mathbb{E}\{W(k)\} = \mathbb{E}\{W(k)\}\mathbf{1} = \mathbf{1}$  where  $\mathbf{1} = (1 \dots 1)'$  denotes the  $n \times 1$  vector of all ones, but  $\mathbb{E}\{W(k)\}$  is not necessarily a stochastic matrix since it may contain negative entries.

Introduce  $y_i(k) = x_i(k) - \sum_{s=1}^n x_s(k)/n$  and let  $y(k) = (y_1(k) \dots y_n(k))'$ . Define  $U := \mathbf{1}\mathbf{1}'/n$  and note that  $y(k) = (I - U)x(k)$ ; furthermore,  $(I - U)W(k) = W(k)(I - U) = W(k) - U$  for all possible realizations of  $W(k)$ . Hence, the evolution of  $\mathbb{E}\{y(k)\}$  is linear:

$$\mathbb{E}\{y(k+1)\} = \mathbb{E}\{(I - U)W(k)x(k)\} = \mathbb{E}\{(I - U)W(k)(I - U)x(k)\} = (\mathbb{E}\{W(k)\} - U)\mathbb{E}\{y(k)\}.$$

The following elementary inequalities

$$|\mathbb{E}\{x_i(k) - x_j(k)\}| \leq |\mathbb{E}\{y_i(k)\}| + |\mathbb{E}\{y_j(k)\}|, \quad |\mathbb{E}\{y_i(k)\}| \leq \frac{1}{n} \mathbb{E} \sum_{s=1}^n |x_i(k) - x_s(k)| \quad (7)$$

imply that belief convergence in expectation is equivalent to  $\lim_{k \rightarrow \infty} |\mathbb{E}\{y(k)\}| = 0$ , and belief divergence is equivalent to  $\limsup_{k \rightarrow \infty} |\mathbb{E}\{y(k)\}| = \infty$ . Belief convergence or divergence is hence determined by the spectral radius of  $\mathbb{E}\{W(k)\} - U$ .

With Assumption 2, there always holds that

$$d_i^+ = \sum_{j=1, j \neq i}^n [P_{\text{pst}}^\dagger]_{ij} \leq \sum_{j=1, j \neq i}^n (p_{ij} + p_{ji})/n \leq 1/2.$$

As a result, Geršhgorin's Circle Theorem (see, e.g., Theorem 6.1.1 in Horn and Johnson (1985)) guarantees that each eigenvalue of  $I - \alpha L_{\text{pst}}^\dagger$  is nonnegative. It then follows that each eigenvalue of  $I - \alpha L_{\text{pst}}^\dagger - U$  is nonnegative since  $L_{\text{pst}}^\dagger U = U L_{\text{pst}}^\dagger = 0$  and the two matrices  $I - \alpha L_{\text{pst}}^\dagger$  and  $U$  share the same eigenvector  $\mathbf{1}$  for eigenvalue one. Moreover, it is well known in algebraic graph theory that  $L_{\text{pst}}^\dagger$  and  $L_{\text{neg}}^\dagger$  are positive semi-definite matrices. As a result, Weyl's inequality (see Theorem 4.3.1 in Horn and Johnson (1985)) further ensures that each eigenvalue of  $\mathbb{E}\{W(k)\} - U$  is also nonnegative. To summarize, we have shown that:

**Proposition 1** *Let Assumption 2 hold. Belief convergence is achieved in expectation for all initial values if  $\lambda_{\max}(I - \alpha L_{\text{pst}}^\dagger + \beta L_{\text{neg}}^\dagger - U) < 1$ ; belief divergence is achieved in expectation for almost all initial values if  $\lambda_{\max}(I - \alpha L_{\text{pst}}^\dagger + \beta L_{\text{neg}}^\dagger - U) > 1$ .*

In the above proposition and what follows,  $\lambda_{\max}(M)$  denotes the largest eigenvalue of the real symmetric matrix  $M$ , and by ‘‘almost all initial conditions,’’ we mean that the property holds for any initial condition  $y(0)$  except if  $y(0)$  is perfectly orthogonal to the eigenspace of  $\mathbb{E}\{W(k)\} - U$  corresponding to its maximal eigenvalue  $\lambda_{\max}(I - \alpha L_{\text{pst}}^\dagger + \beta L_{\text{neg}}^\dagger - U)$ . Hence the set of initial conditions where the property does not hold has zero Lebesgue measure.

The Courant-Fischer Theorem (see Theorem 4.2.11 in Horn and Johnson (1985)) implies

$$\begin{aligned} \lambda_{\max}(I - \alpha L_{\text{pst}}^\dagger + \beta L_{\text{neg}}^\dagger - U) &= \sup_{|z|=1} z' (I - \alpha L_{\text{pst}}^\dagger + \beta L_{\text{neg}}^\dagger - U) z \\ &= 1 + \sup_{|z|=1} \left[ -\alpha \sum_{\{i,j\} \in \mathbf{E}_{\text{pst}}} [P^\dagger]_{ij} (z_i - z_j)^2 + \beta \sum_{\{i,j\} \in \mathbf{E}_{\text{neg}}} [P^\dagger]_{ij} (z_i - z_j)^2 - \frac{1}{n} \left( \sum_{i=1}^n z_i \right)^2 \right]. \end{aligned} \quad (8)$$

We see from (8) that the influence of  $\mathbf{G}_{\text{pst}}$  and  $\mathbf{G}_{\text{neg}}$  on the belief convergence/divergence in mean are *separated*: links in  $\mathbf{E}_{\text{pst}}$  contribute to belief convergence, while links in  $\mathbf{E}_{\text{neg}}$  contribute to belief divergence. As will be shown later on, this separation property no longer holds for mean-square convergence, and there may be a non-trivial correlation between the influence of  $\mathbf{E}_{\text{pst}}$  and that of  $\mathbf{E}_{\text{neg}}$ .

### 3.1.2 Phase Transition

Next we study the impact of update parameters  $\alpha$  and  $\beta$  on the convergence in expectation. Define  $f(\alpha, \beta) := \lambda_{\max}(I - \alpha L_{\text{pst}}^\dagger + \beta L_{\text{neg}}^\dagger - U)$ . The function  $f$  has the following properties under Assumption 2:

- (i) (*Convexity*) Since both  $L_{\text{pst}}^\dagger$  and  $L_{\text{neg}}^\dagger$  are symmetric,  $f(\alpha, \beta)$  is the spectral norm of  $I - \alpha L_{\text{pst}}^\dagger + \beta L_{\text{neg}}^\dagger - U$ . As every matrix norm is convex, we have

$$f(\gamma(\alpha_1, \beta_1) + (1 - \gamma)(\alpha_2, \beta_2)) \leq \gamma f(\alpha_1, \beta_1) + (1 - \gamma)f(\alpha_2, \beta_2) \quad (9)$$

for all  $\gamma \in [0, 1]$  and  $\alpha_1, \alpha_2, \beta_1, \beta_2 \in \mathbb{R}$ . This implies that  $f(\alpha, \beta)$  is convex in  $(\alpha, \beta)$ .

- (ii) (*Monotonicity*) From (8),  $f(\alpha, \beta)$  is non-increasing in  $\alpha$  for fixed  $\beta$ , and non-decreasing in  $\beta$  for fixed  $\alpha$ . As a result, setting  $\alpha = 1$  provides the *fastest* convergence whenever belief convergence in expectation is achieved (for a given fixed  $\beta$ ). Note that when  $\alpha = 1$ , when two nodes interact, they simply switch their beliefs.

When  $G_{\text{pst}}$  is connected, the second smallest eigenvalue of  $L_{\text{pst}}^\dagger$ , denoted by  $\lambda_2(L_{\text{pst}}^\dagger)$ , is positive. We can readily see that  $f(\alpha, 0) = 1 - \alpha \lambda_2(L_{\text{pst}}^\dagger) < 1$ . From (8), we also have  $f(\alpha, \beta) \rightarrow \infty$  as  $\beta \rightarrow \infty$  provided that  $G_{\text{neg}}$  is nonempty. Combining these observations with the monotonicity of  $f$ , we conclude that:

**Proposition 2** *Assume that  $G_{\text{pst}}$  is connected and let Assumption 2 hold. Then for any fixed  $\alpha \in (0, 1]$ , there exists a threshold value  $\beta_\star > 0$  (that depends on  $\alpha$ ) such that*

- (i) *Belief convergence in expectation is achieved for all initial values if  $0 \leq \beta < \beta_\star$ ;*  
 (ii) *Belief divergence in expectation is achieved for almost all initial values if  $\beta > \beta_\star$ .*

We remark that belief divergence can only happen for almost all initial values since if the initial beliefs of all the nodes are identical, they do not evolve over time.

### 3.1.3 Examples

An interesting question is to determine how the phase transition threshold  $\beta_\star$  scales with the network size. Answering this question seems challenging. However there are networks for which we can characterize  $\beta_\star$  exactly. Next we derive explicit expressions for  $\beta_\star$  when  $G$  is a complete graph or a ring graph. These two topologies represent the most dense and almost the most sparse structures for a connected network.

**Example 1 (Complete Graph)** Let  $G = K_n$ , where  $K_n$  is the complete graph with  $n$  nodes, and consider the node pair selection matrix  $P = (\mathbf{1}\mathbf{1}' - I)/(n - 1)$ . Let  $L(K_n) = nI - \mathbf{1}\mathbf{1}'$  be the Laplacian of  $K_n$ . Then  $L(K_n)$  has eigenvalue 0 with multiplicity 1 and eigenvalue  $n$  with multiplicity  $n - 1$ . Define  $L(G_{\text{neg}})$  as the standard Laplacian of  $G_{\text{neg}}$ . Observe that

$$\begin{aligned} I - \alpha L_{\text{pst}}^\dagger + \beta L_{\text{neg}}^\dagger - U &= I - \alpha(L_{\text{pst}}^\dagger + L_{\text{neg}}^\dagger) + (\alpha + \beta)L_{\text{neg}}^\dagger - U \\ &= I - \frac{2\alpha}{n(n-1)}L(K_n) + \frac{2(\alpha + \beta)}{n(n-1)}L(G_{\text{neg}}) - U. \end{aligned} \quad (10)$$

Also note that  $L(G_{\text{neg}})L(K_n) = L(K_n)L(G_{\text{neg}}) = nL(G_{\text{neg}})$ . From these observations, we can then readily conclude that:

$$\beta_\star = \frac{n\alpha}{\lambda_{\max}(L(G_{\text{neg}}))} - \alpha. \quad (11)$$

**Example 2 (Erdős-Rényi Negative Graph over Complete Graph)** Let  $G = K_n$  with

$$P = (\mathbf{1}\mathbf{1}' - I)/(n - 1).$$

Let  $G_{\text{neg}}$  be the Erdős-Rényi random graph (Erdős and Rényi (1960)) where for any  $i, j \in V$ ,  $\{i, j\} \in E_{\text{neg}}$  with probability  $p$  (independently of other links). Note that since  $G_{\text{neg}}$  is a random subgraph, the function  $f(\alpha, \beta)$  becomes a random variable, and we denote by  $\mathbf{P}$  the probability measure related to the randomness of the graph in Erdős-Rényi's model. Spectral theory for random graphs suggests that (Ding and Jiang (2010))

$$\frac{\lambda_{\max}(L(G_{\text{neg}}))}{pn} \rightarrow 1, \text{ as } n \rightarrow \infty \quad (12)$$

in probability. Now for fixed  $p$ , we deduce from (11) and (12) that the threshold  $\beta_\star$  converges, as  $n$  grows large, to  $\alpha/p$  in probability. Now let us fix the update parameters  $\alpha$  and  $\beta$ , and investigate the impact of the probability  $p$  on the convergence in mean.

- If  $p < \frac{\alpha}{\alpha + \beta}$ , we show that  $\mathbf{P}[f(\alpha, \beta) < 1] \rightarrow 1$ , when  $n \rightarrow \infty$ , i.e., when the network is large, we likely achieve convergence in mean. Let  $\epsilon < \frac{\alpha}{(\alpha + \beta)p} - 1$ . It follows from (12) that

$$\begin{aligned} \mathbf{P}(f(\alpha, \beta) < 1) &= \mathbf{P}\left(1 - \frac{2\alpha}{n(n-1)}n + \frac{2(\alpha + \beta)}{n(n-1)}\lambda_{\max}(L(G_{\text{neg}})) < 1\right) \\ &= \mathbf{P}\left((\alpha + \beta)\lambda_{\max}(L(G_{\text{neg}})) < \alpha n\right) \\ &= \mathbf{P}\left(\frac{\lambda_{\max}(L(G_{\text{neg}}))}{pn} < \frac{\alpha}{(\alpha + \beta)p}\right) \\ &\geq \mathbf{P}\left(\left|\frac{\lambda_{\max}(L(G_{\text{neg}}))}{pn} - 1\right| < \epsilon\right) \rightarrow 1, \text{ as } n \rightarrow \infty. \end{aligned} \quad (13)$$

- If  $p > \frac{\alpha}{\alpha + \beta}$ , we similarly establish that  $\mathbf{P}(f(\alpha, \beta) > 1) \rightarrow 1$ , when  $n \rightarrow \infty$ , i.e., when the network is large, we observe divergence in mean with high probability.

Hence we have a sharp phase transition between convergence and divergence in mean when the proportion of negative links  $p$  increases and goes above the threshold  $p_\star = \alpha/(\alpha + \beta)$ .

**Example 3 (Ring Graph)** Denote  $R_n$  as the ring graph with  $n$  nodes. Let  $A(R_n)$  and  $L(R_n)$  be the adjacency and Laplacian matrices of  $R_n$ , respectively. Let the underlying graph  $G = R_n$  with only one negative link (if one has more than two negative links, it is easy to see that divergence in expectation is achieved irrespective of  $\beta > 0$ ). Take  $P = A(R_n)/2$ . We know that  $L(R_n)$  has eigenvalues  $2 - 2\cos(2\pi k/n)$ ,  $0 \leq k \leq n/2$ . Applying Weyl's inequality we obtain  $f(\alpha, \beta) \geq 1 + (\beta - \alpha)/n$ . We conclude that  $\beta_\star < \alpha$ , irrespective of  $n$ .

### 3.2 Mean-square Convergence

We now turn our attention to the analysis of the mean-square convergence and divergence. Define:

$$\begin{aligned} \mathbb{E}\{|y(k)|^2\} &= \mathbb{E}\{x(k)'(I - U)x(k)\} \\ &= x(0)'\mathbb{E}\{W(0) \dots W(k-1)(I - U)W(k-1) \dots W(0)\}x(0). \end{aligned} \quad (14)$$

Again based on inequalities (7), we see that belief convergence in mean square is equivalent to

$$\lim_{k \rightarrow \infty} \mathbb{E}\{|y(k)|^2\} = 0,$$

and belief divergence to  $\limsup_{k \rightarrow \infty} \mathbb{E}\{|y(k)|^2\} = \infty$ . Define:

$$\Phi(k) = \begin{cases} \mathbb{E}\{W(0) \dots W(k-1)(I - U)W(k-1) \dots W(0)\}, & k \geq 1, \\ I - U, & k = 0. \end{cases} \quad (15)$$

Then,  $\Phi(k)$  evolves as a linear dynamical system (Fagnani and Zampieri (2008))

$$\begin{aligned} \Phi(k) &= \mathbb{E}\{W(0) \dots W(k-1)(I - U)W(k-1) \dots W(0)\} \\ &= \mathbb{E}\{W(0)(I - U)W(1) \dots W(k-1)(I - U)W(k-1) \dots W(1)(I - U)W(0)\} \\ &= \mathbb{E}\{(W(k) - U)\Phi(k-1)(W(k) - U)\}, \end{aligned} \quad (16)$$

where in the second equality we have used the facts that  $(I - U)^2 = I - U$  and  $(I - U)W(k) = W(k)(I - U) = W(k) - U$  for all possible realizations of  $W(k)$ , and the third equality is due to that  $W(k)$  and  $W(0)$  are i.i.d. We can rewrite (16) using an equivalent vector form:

$$\mathbf{vec}(\Phi(k)) = \Theta \mathbf{vec}(\Phi(k-1)), \quad (17)$$

where  $\Theta$  is the matrix in  $\mathbb{R}^{n^2 \times n^2}$  given by

$$\begin{aligned} \Theta &= \mathbb{E}\{(W(0) - U) \otimes (W(0) - U)\} \\ &= \sum_{\{i,j\} \in \mathbf{G}_{\text{pst}}} [P^\dagger]_{ij} \left( (W_{ij}^+ - U) \otimes (W_{ij}^+ - U) \right) + \sum_{\{i,j\} \in \mathbf{G}_{\text{neg}}} [P^\dagger]_{ij} \left( (W_{ij}^- - U) \otimes (W_{ij}^- - U) \right). \end{aligned}$$

Let  $S_\lambda$  be the eigenspace corresponding to an eigenvalue  $\lambda$  of  $\Theta$ . Define

$$\lambda_\star := \max\{\lambda \in \sigma(\Theta) : \mathbf{vec}(I - U) \notin S_\lambda^\perp\},$$

which denotes the spectral radius of  $\Theta$  restricted to the smallest invariant subspace containing  $\mathbf{vec}(I - U)$ , i.e.,  $S := \text{span}\{\Theta^k \mathbf{vec}(I - U), k = 0, 1, \dots\}$ . Then mean-square belief convergence/divergence is fully determined by  $\lambda_\star$ : convergence in mean square for all initial conditions is achieved if  $\lambda_\star < 1$ , and divergence for almost all initial conditions is achieved if  $\lambda_\star > 1$ .

Observing that  $\lambda \leq 1$  for every  $\lambda \in \sigma(W_{ij}^+)$  and  $\lambda \geq 1$  for every  $\lambda \in \sigma(W_{ij}^-)$ , we can also conclude that each link in  $\mathbf{E}_{\text{pst}}$  contributes positively to  $\lambda_{\max}(\Theta)$  and each link in  $\mathbf{E}_{\text{neg}}$  contributes negatively to  $\lambda_{\max}(\Theta)$ . However, unlike in the case of the analysis of convergence in expectation, although  $\lambda_\star$  defines a precise threshold for the phase-transition between mean-square convergence and divergence, it is difficult to determine the influence  $\mathbf{E}_{\text{pst}}$  and  $\mathbf{E}_{\text{neg}}$  have on  $\lambda_\star$ . The reason is that they are coupled in a nontrivial manner for the invariant subspace  $S$ . Nevertheless, we are still able to propose the following conditions for mean-square belief convergence and divergence:

**Proposition 3** *Belief convergence is achieved for all initial values in mean square if*

$$\lambda_{\max}(I - 2\alpha(1 - \alpha)L_{\text{pst}}^\dagger + 2\beta(1 + \beta)L_{\text{neg}}^\dagger - U) < 1;$$

*belief divergence is achieved in mean square for almost all initial values if*

$$\lambda_{\max}(I - \alpha L_{\text{pst}}^\dagger + \beta L_{\text{neg}}^\dagger - U) > 1$$

or

$$\lambda_{\min}(I - 2\alpha(1 - \alpha)L_{\text{pst}}^\dagger + 2\beta(1 + \beta)L_{\text{neg}}^\dagger - U) > 1.$$

The condition  $\lambda_{\max}(I - \alpha L_{\text{pst}}^\dagger + \beta L_{\text{neg}}^\dagger - U)$  is sufficient for mean square divergence, in view of Proposition 1 and the fact that  $\mathcal{L}_1$  divergence implies  $\mathcal{L}_p$  divergence for all  $p \geq 1$ . The other conditions are essentially consistent with the upper and lower bounds of  $\lambda_\star$  established in Proposition 4.4 of Fagnani and Zampieri (2008). Proposition 3 is a consequence of Lemma 3 (see Appendix), as explained in Remark 5.



## 4 Almost Sure Convergence/Divergence

In this section, we explore the almost sure convergence of beliefs in signed social networks. We introduce the following definition.

**Definition 3** *Belief convergence is achieved almost surely (a.s.) if  $\mathbb{P}(\lim_{k \rightarrow \infty} |x_i(k) - x_j(k)| = 0) = 1$  for all  $i$  and  $j$ ; Belief divergence is achieved almost surely if  $\mathbb{P}(\limsup_{k \rightarrow \infty} \max_{i,j} |x_i(k) - x_j(k)| = \infty) = 1$ .*

Basic probability theory tells us that mean-square belief convergence implies belief convergence in expectation (mean convergence), and similarly belief divergence in expectation implies belief divergence in mean square. However, in general there is no direct connection between almost sure convergence/divergence and mean or mean-square convergence/divergence. Finally observe that, a priori, it is not clear that either a.s. convergence or a.s. divergence should be achieved.

While the analysis of the convergence of beliefs in mean and square-mean mainly relied on spectral arguments, we need more involved probabilistic methods (e.g., sample-path arguments, martingale convergence theorems) to study almost sure convergence or divergence. We first establish two insightful properties of the belief evolutions: (i) the *no-survivor* property stating that in case of almost sure divergence, the difference between the beliefs of any two nodes in the network tends to infinity (along a subsequence of instants); (ii) the *live-or-die* property which essentially states that the maximum difference between the beliefs of any two nodes either grows to infinity, or vanishes to zero. We then show a zero-one law and a phase transition of almost sure convergence/divergence. Finally, we investigate the robustness of networks against negative links. More specifically, we show that when the graph  $G_{\text{pst}}$  of positive links contains an hypercube, and when the positive updates are truly averaging, i.e.,  $\alpha = 1/2$ , then almost sure belief convergence is reached in finite time, irrespective of the number of negative links, their positions in the network, and the negative update parameter  $\beta$ . We believe that these are the only networks enjoying this strong robustness property.

### 4.1 The No-Survivor Theorem

The following theorem establishes that in the case of almost sure divergence, there is no pair of nodes that can *survive* this divergence: for any two nodes, the difference in their beliefs grow arbitrarily large.

**Theorem 1** (*No-Survivor*) *Fix the initial condition and assume almost sure belief divergence. Then  $\mathbb{P}(\limsup_{k \rightarrow \infty} |x_i(k) - x_j(k)| = \infty) = 1$  for all  $i \neq j \in \mathcal{V}$ .*

Observe that the above result only holds for the almost sure divergence. We may easily build simple network examples where we have belief divergence in expectation (or mean square), but where some node pairs survive, in the sense that the difference in their beliefs vanishes (or at least bounded). The no-survivor theorem indicates that to check almost sure divergence, we may just observe the evolution of beliefs held at two arbitrary nodes in the network.

## 4.2 The Live-or-Die lemma and Zero-One Laws

Next we further classify the ways beliefs can evolve. Specifically, we study the following events: for any initial beliefs  $x^0$ ,

$$\begin{aligned}\mathcal{C}_{x^0} &\doteq \left\{ \limsup_{k \rightarrow \infty} \max_{i,j} |x_i(k) - x_j(k)| = 0 \right\}, & \mathcal{D}_{x^0} &\doteq \left\{ \limsup_{k \rightarrow \infty} \max_{i,j} |x_i(k) - x_j(k)| = \infty \right\}, \\ \mathcal{C}_{x^0}^* &\doteq \left\{ \liminf_{k \rightarrow \infty} \max_{i,j} |x_i(k) - x_j(k)| = 0 \right\}, & \mathcal{D}_{x^0}^* &\doteq \left\{ \liminf_{k \rightarrow \infty} \max_{i,j} |x_i(k) - x_j(k)| = \infty \right\},\end{aligned}$$

and

$$\begin{aligned}\mathcal{C} &\doteq \left\{ \limsup_{k \rightarrow \infty} \max_{i,j} |x_i(k) - x_j(k)| = 0 \text{ for all } x^0 \in \mathbb{R}^n \right\}, \\ \mathcal{D} &\doteq \left\{ \exists (\text{deterministic}) x^0 \in \mathbb{R}^n, \text{ s.t. } \limsup_{k \rightarrow \infty} \max_{i,j} |x_i(k) - x_j(k)| = \infty \right\}.\end{aligned}$$

We establish that the maximum difference between the beliefs of any two nodes either goes to  $\infty$ , or to zero. This result is referred to as *live-or-die* lemma:

**Lemma 1** (*Live-or-Die*) *Let  $\alpha \in (0, 1)$  and  $\beta > 0$ . Suppose  $\mathbf{G}_{\text{pst}}$  is connected. Then (i)  $\mathbb{P}(\mathcal{C}_{x^0}) + \mathbb{P}(\mathcal{D}_{x^0}) = 1$ ; (ii)  $\mathbb{P}(\mathcal{C}_{x^0}^*) + \mathbb{P}(\mathcal{D}_{x^0}^*) = 1$ .*

*As a consequence, almost surely, one of the following events happens:*

$$\begin{aligned}&\left\{ \lim_{k \rightarrow \infty} \max_{i,j} |x_i(k) - x_j(k)| = 0 \right\}; \\ &\left\{ \lim_{k \rightarrow \infty} \max_{i,j} |x_i(k) - x_j(k)| = \infty \right\}; \\ &\left\{ \liminf_{k \rightarrow \infty} \max_{i,j} |x_i(k) - x_j(k)| = 0; \limsup_{k \rightarrow \infty} \max_{i,j} |x_i(k) - x_j(k)| = \infty \right\}.\end{aligned}$$

The Live-or-Die lemma deals with events where the initial beliefs have been fixed. We may prove stronger results on the probabilities of events that hold for *any* initial condition, e.g.,  $\mathcal{C}$ , or for at least one initial condition, e.g.,  $\mathcal{D}$ :

**Theorem 2** (*Zero-One Law*) *Let  $\alpha \in [0, 1]$  and  $\beta > 0$ . Both  $\mathcal{C}$  and  $\mathcal{D}$  are trivial events (i.e., each of them occurs with probability equal to either 1 or 0) and  $\mathbb{P}(\mathcal{C}) + \mathbb{P}(\mathcal{D}) = 1$ .*

To prove this result, we show that  $\mathcal{C}$  is a tail event, and hence trivial in view of Kolmogorov's zero-one law (the same kind of arguments has been used by Tahbaz-Salehi and Jadbabaie (2008)). From the Live-or-Die lemma, we then simply deduce that  $\mathcal{D}$  is also a trivial event. Note that  $\mathcal{C}_{x^0}$  and  $\mathcal{D}_{x^0}$  may not be trivial events. In fact, we can build examples where  $\mathbb{P}(\mathcal{C}_{x^0}) = 1/2$  and  $\mathbb{P}(\mathcal{D}_{x^0}) = 1/2$ .

### 4.3 Phase Transition

As for the convergence in expectation, for fixed positive update parameter  $\alpha$ , we are able to establish the existence of thresholds for the value  $\beta$  of the negative update parameter, which characterizes the almost sure belief convergence and divergence.

**Theorem 3** (*Phase Transition*) *Suppose  $\mathbf{G}_{\text{pst}}$  is connected. Fix  $\alpha \in (0, 1)$  with  $\alpha \neq 1/2$ . Then*

- (i) *there exists  $\beta^\sharp(\alpha) > 0$  such that  $\mathbb{P}(\mathcal{C}) = 1$  if  $0 \leq \beta < \beta^\sharp$ ;*
- (ii) *there exists  $\beta^\sharp(\alpha) > 0$  such that  $\mathbb{P}(\liminf_{k \rightarrow \infty} \max_{i,j} |x_i(k) - x_j(k)| = \infty) = 1$  for almost all initial values if  $\beta > \beta^\sharp$ .*

It should be observed that the divergence condition in (ii) is stronger than our notion of almost sure belief divergence (the maximum belief difference between two nodes diverge almost surely to  $\infty$ ). Also note that  $\beta^\sharp \leq \beta^\sharp$ , and we were not able to show that the gap between these two thresholds vanishes (as in the case of belief convergence in expectation or mean-square).

### 4.4 Robustness to Negative Links: the Hypercube

We have seen in Theorem 3 that when  $\alpha \neq 1/2$ , one single negative link is capable of driving the network beliefs to almost sure divergence as long as  $\beta$  is sufficiently large. The following result shows that the evolution of the beliefs can be robust against negative links. This is the case when nodes can reach an agreement in finite time. In what follows, we provide conditions on  $\alpha$  and the structure of the graph under which finite time belief convergence is reached.

**Proposition 4** *Suppose there exist an integer  $T \geq 1$  and a finite sequence of node pairs  $\{i_s, j_s\} \in \mathbf{G}_{\text{pst}}, s = 1, 2, \dots, T$  such that  $W_{i_T j_T}^+ \cdots W_{i_1 j_1}^+ = U$ . Then  $\mathbb{P}(\mathcal{C}) = 1$  for all  $\beta \geq 0$ .*

Proposition 4 is a direct consequence of the Borel-Cantelli Lemma. If there is a finite sequence of node pairs  $\{i_s, j_s\} \in \mathbf{G}_{\text{pst}}, s = 1, 2, \dots, T$  such that  $W_{i_T j_T}^+ \cdots W_{i_1 j_1}^+ = U$ , then

$$\mathbb{P}\left(W(k+T) \cdots W(k+1) = U\right) \geq \left(\frac{p^*}{n}\right)^T,$$




Figure 2: The hypercubes  $H^1$ ,  $H^2$ , and  $H^3$ .

for all  $k \geq 0$ , where  $p_* = \min\{p_{ij} + p_{ji} : \{i, j\} \in E\}$ . Noting that  $UW(k) = W(k)U = U$  for all possible realizations of  $W(k)$ , the Borel-Cantelli Lemma guarantees that

$$\mathbb{P}\left(\lim_{k \rightarrow \infty} W(k) \cdots W(0) = U\right) = 1$$

for all  $\beta \geq 0$ , or equivalently,  $\mathbb{P}(\mathcal{C}) = 1$  for all  $\beta \geq 0$ . This proves Proposition 4.

The existence of such finite sequence of node pairs under which the beliefs of the nodes in the network reach a common value in finite time is crucial (we believe that this condition is actually necessary) to ensure that the influence of  $G_{\text{neg}}$  vanishes. It seems challenging to know whether this is at all possible. As it turns out, the structure of the positive graph plays a fundamental role. To see that, we first provide some definitions.

**Definition 4** Let  $G_1 = (V_1, E_1)$  and  $G_2 = (V_2, E_2)$  be a pair of graphs. The Cartesian product of  $G_1$  and  $G_2$ , denoted by  $G_1 \square G_2$ , is defined by

- (i) the vertex set of  $G_1 \square G_2$  is  $V_1 \times V_2$ , where  $V_1 \times V_2$  is the Cartesian product of  $V_1$  and  $V_2$ ;
- (ii) for any two vertices  $(v_1, v_2), (u_1, u_2) \in V_1 \times V_2$ , there is an edge between them in  $G_1 \square G_2$  if and only if either  $v_1 = u_1$  and  $\{v_2, u_2\} \in E_2$ , or  $v_2 = u_2$  and  $\{v_1, u_1\} \in E_1$ .

Let  $K_2$  be the complete graph with two nodes. The  $m$ -dimensional Hypercube  $H^m$  is then defined as

$$H^m = \underbrace{K_2 \square K_2 \cdots \square K_2}_{m \text{ times}}.$$

An illustration of hypercubes is in Figure 2.

The following result provides sufficient conditions to achieve finite-time convergence.

**Proposition 5** If  $\alpha = 1/2$ ,  $n = 2^m$  for some integer  $m > 0$ , and  $G_{\text{pst}}$  has a subgraph isomorphic with an  $m$ -dimensional hypercube, then there exists a sequence of  $(n \log_2 n)/2$  node pairs  $\{i_s, j_s\} \in G_{\text{pst}}$ ,  $s = 1, \dots, (n \log_2 n)/2$  such that  $W_{i_{(n \log_2 n)/2} j_{(n \log_2 n)/2}}^+ \cdots W_{i_1 j_1}^+ = U$ .

Next we derive necessary conditions for finite time convergence. Let us first recall the following definition.

**Definition 5** Let  $G = (V, E)$  be a graph. A matching of  $G$  is a set of pairwise non-adjacent edges in the sense that no two edges share a common vertex. A perfect matching of  $G$  is a matching which matches all vertices.

**Proposition 6** If there exist an integer  $T \geq 1$  and a sequence of node pairs  $\{i_s, j_s\} \in G_{\text{pst}}, s = 1, 2, \dots, T$  such that  $W_{i_T j_T}^+ \cdots W_{i_1 j_1}^+ = U$ , then  $\alpha = 1/2$ ,  $n = 2^m$ , and  $G_{\text{pst}}$  has a perfect matching.

In fact, in the proof of Proposition 6, we show that if  $W_{i_T j_T}^+ \cdots W_{i_1 j_1}^+ = U$ , then a subset of

$$\{\{i_1, j_1\}, \dots, \{i_T, j_T\}\}$$

forms a perfect matching of  $G_{\text{pst}}$ .

We have seen that the belief dynamics and convergence can be robust against negative links, but this robustness comes at the expense of strong conditions on the number of the nodes and the structure of the positive graph.

## 5 Belief Clustering and Structural Balance

So far we have studied the belief dynamics when the node interactions are symmetric, and the values of beliefs are unconstrained. The results illustrate that often either convergence or divergence can be predicted for the social-network beliefs. Although this symmetric and unconstrained belief update rule is plausible for ideal social network models, in reality these assumptions might not hold, that is: when  $\{i, j\}$  is selected, it might happen that only one of the two nodes in  $i$  and  $j$  updates its belief; there might be a hard constraint on beliefs:  $x_i(k) \in [-A, A]$  for all  $i$  and  $k$ , and for some  $A > 0$ .

In this section, we consider the following model for the updates of the beliefs. Define:

$$\mathcal{P}_A(z) = \begin{cases} -A, & \text{if } z < -A; \\ z, & \text{if } z \in [-A, A]; \\ A, & \text{if } z > A. \end{cases} \quad (18)$$

Let  $a, b, c > 0$  be three positive real numbers such that  $a + b + c = 1$ , and define the function  $\theta : E \rightarrow \mathbb{R}$  so that  $\theta(\{i, j\}) = \alpha$  if  $\{i, j\} \in E_{\text{pst}}$  and  $\theta(\{i, j\}) = -\beta$  if  $\{i, j\} \in E_{\text{neg}}$ . Assume that node  $i$  interacts with node  $j$  at time  $k$ . Nodes  $i$  and  $j$  update their beliefs as:

[Asymmetric and Constrained Belief Evolution]

$$\begin{aligned}
 x_i(k+1) &= \mathcal{P}_A((1-\theta)x_i(k) + \theta x_j(k)) \text{ and } x_j(k+1) = x_j(k), & \text{with probability } a; \\
 x_j(k+1) &= \mathcal{P}_A((1-\theta)x_j(k) + \theta x_i(k)) \text{ and } x_i(k+1) = x_i(k), & \text{with probability } b; \\
 x_m(k+1) &= \mathcal{P}_A((1-\theta)x_m(k) + \theta x_{-m}(k)), \quad m \in \{i, j\}, & \text{with probability } c.
 \end{aligned} \tag{19}$$

Enforcing the belief within the interval  $[-A, A]$  can be viewed as a social member's decision based on her fundamental model of the world. With asymmetric and constrained belief evolution, the dynamics become essentially nonlinear, which brings new challenges in the analysis. We continue to use  $\mathbb{P}$  to denote the overall probability measure capturing the randomness of the updates in the asymmetric constrained model.

## 5.1 Balanced Graphs and Clustering

We introduce the notion of *balance* for signed graphs, for which we refer to Wasserman and Faust (1994) for a comprehensive discussion.

**Definition 6** *Let  $G = (V, E)$  be a signed graph. Then*

(i)  *$G$  is weakly balanced if there is an integer  $k \geq 2$  and a partition of  $V = V_1 \cup V_2 \cdots \cup V_k$ , where  $V_1, \dots, V_k$  are nonempty and mutually disjoint, such that any edge between different  $V_i$ 's is negative, and any edge within each  $V_i$  is positive.*

(ii)  *$G$  is strongly balanced if it is weakly balanced with  $k = 2$ .*

Harary's balance theorem states that a signed graph  $G$  is strongly balanced if and only if there is no cycle with an odd number of negative edges in  $G$  (Cartwright and Harary (1956)), while  $G$  is weakly balanced if and only if no cycle has exactly one negative edge in  $G$  (Davis (1967)).

It turned out that, with certain balance of the underlying graph, clustering arises for the social-network beliefs. We make the following definition.

**Definition 7** (i) *Let  $G$  be strongly balanced subject to partition  $V = V_1 \cup V_2$ . Then almost sure boundary belief clustering for the initial value  $x^0$  is achieved if there are two random variables  $B_1^\dagger(x^0)$  and  $B_2^\dagger(x^0)$ , both taking values in  $\{-A, A\}$ , such that:*

$$\mathbb{P}\left(\lim_{k \rightarrow \infty} x_i(k) = B_1^\dagger(x^0), i \in V_1; \lim_{k \rightarrow \infty} x_i(k) = B_2^\dagger(x^0), i \in V_2\right) = 1. \tag{20}$$

(ii) *Let  $G$  be weakly balanced subject to partition  $V = V_1 \cup V_2 \cdots \cup V_m$  for some  $m \geq 2$ . Then almost sure boundary belief clustering for the initial value  $x^0$  is achieved if there are there are  $m$  random variables,*

$B_1^\sharp(x^0), \dots, B_m^\sharp(x^0)$ , each of which taking values in  $\{-A, A\}$ , such that:

$$\mathbb{P}\left(\lim_{k \rightarrow \infty} x_i(k) = B_j^\sharp(x^0), i \in V_j, j = 1, \dots, m\right) = 1. \quad (21)$$

In the case of strongly balanced graphs, we can show that beliefs are asymptotically clustered when  $\beta$  is large enough, as stated in the following theorem.

**Theorem 4** *Assume that  $\mathbf{G}$  is strongly balanced under partition  $\mathbf{V} = \mathbf{V}_1 \cup \mathbf{V}_2$ , and that  $\mathbf{G}_{\mathbf{V}_1}$  and  $\mathbf{G}_{\mathbf{V}_2}$  are connected. For any  $\alpha \in (0, 1) \setminus \{1/2\}$ , when  $\beta$  is sufficiently large, for almost all initial values  $x^0$ , almost sure boundary belief clustering is achieved under the update rule (19).*

In fact, there holds  $B_1^\dagger(x^0) + B_2^\dagger(x^0) = 0$  almost surely in the above boundary belief clustering for strongly balanced social networks. Theorem 4 states that, for strongly balanced social networks, beliefs are eventually polarized to the two opinion boundaries.

The analysis of belief dynamics in weakly balanced graphs is more involved, and we restrict our attention to complete graphs. In social networks, this case means that everyone knows everyone else – which constitutes a suitable model for certain social groups of small sizes (a classroom, a sport team, or the UN, see Easley and Kleinberg (2010)). As stated in the following theorem, for weakly balanced complete graphs, beliefs are again clustered.

**Theorem 5** *Assume that  $\mathbf{G}$  is a complete and weakly balanced graph under the partition  $\mathbf{V} = \mathbf{V}_1 \cup \mathbf{V}_2 \dots \cup \mathbf{V}_m$  with  $m \geq 2$ . Further assume that  $\mathbf{G}_{\mathbf{V}_j}, j = 1, \dots, m$  are connected. For any  $\alpha \in (0, 1) \setminus \{1/2\}$ , when  $\beta$  is sufficiently large, almost sure boundary belief clustering is achieved for almost all initial values under (19).*

**Remark 2** *Under the model (3), it can be shown (cf., Altafini (2013), Shi et al. (2015))*

(i) *if  $\mathbf{G}$  is strongly balanced and  $\beta \in (0, 1)$ , then there are two values  $z_1(x^0)$  and  $z_2(x^0)$  such that*

$$\mathbb{P}\left(\lim_{k \rightarrow \infty} x_i(k) = z_1(x^0), i \in \mathbf{V}_1, \lim_{k \rightarrow \infty} x_i(k) = z_2(x^0), i \in \mathbf{V}_2\right) = 1. \quad (22)$$

(ii) *if  $\mathbf{G}$  is not strongly balanced (i.e., even if it is weakly balanced) and  $\beta \in (0, 1)$ , then*

$$\mathbb{P}\left(\lim_{k \rightarrow \infty} x_i(k) = 0, i \in \mathbf{V}\right) = 1, \quad (23)$$

*where the impact of the initial beliefs are entirely erased from the asymptotic limit.*

Our Theorem 4 appears to be similar to (22), but the clustering in Theorem 4 is due to fundamentally different reasons: besides the strong balance of the social network, it is the nonlinearity in the constrained

update ( $\mathcal{P}_A(\cdot)$ ), and the sufficiently large  $\beta$  that makes the boundary clustering arise in Theorem 4. In contrast, (22) is resulted from the crucial condition that  $\beta \in (0, 1)$ . Under the Altafini model (3), even when  $\beta$  is sufficiently large, it is easy to see that the boundary clustering in Theorem 5 can never happen for weakly balanced graphs.

The distribution of the clustering limits established in Theorems 4 and 5 relies on the initial value. In this way, the initial beliefs make an impact on the final belief limit, which is either  $A$  or  $-A$ . The boundary clustering is due to the hard boundaries of the beliefs as well as the negative updates (ironically the larger the better), whose mechanism is fundamentally different with the opinion clustering phenomena resulted from missing of connectivity in Krause types of models Krause (1997), Blondel et al. (2009, 2010), Li et al. (2013), or nonlinear bias in the opinion evolution Dandekar et al. (2013).

**Remark 3** *Note that in the considered asymmetric and constrained belief evolution, we take symmetric belief boundaries  $[-A, A]$  just for simplifying the discussion. Theorems 4 and 5 continue to hold if the belief boundaries are chosen to be  $[A, B]$  for arbitrary  $-\infty < A < B < \infty$ <sup>1</sup>. Letting  $A = 0, B = 1$ , our boundary clustering results in Theorems 4 and 5 are then formally consistent with the belief polarization result, Theorem 3, in Dandekar et al. (2013). It is worth mentioning that Theorem 3 in Dandekar et al. (2013) relies on a type of strong balance (the two-island assumption) and that the initial beliefs should be separated, while Theorems 4 and 5 hold for almost arbitrary initial values.*

The proof of Theorems 4 and 5 is obtained by establishing the almost sure happening of suitable *separation* events, i.e., the node beliefs for a subnetwork become group polarized (either larger or smaller than the remaining nodes' beliefs). From the analysis it is clear that such events tend to happen more easily for small subnetworks in the partition of (strongly or weakly) balanced social networks. On the other hand, boundary belief clustering follows quickly after the separation event, even in the presence of large subgroups. For a large subgroup, the boundary clustering to a consensus for its members is more a consequence of the “push” by the already separated small subgroups, rather than the trustful interactions therein. This means, relatively small subgroups contribute to faster occurrence of the clustering of the entire social network beliefs. Therefore, these results are in strong consistency with the research of minority influence in social psychology Nemeth (1986), Clark and Maass (1990), which suggests that consistent minorities can substantially modify people's private attitudes and opinions.

---

<sup>1</sup>This further confirms that, in our model, the origin of the belief space has no special meaning at all, in contrast to the model of Altafini (2013).



## 5.2 When Balance is Missing

Since the boundary constraint only restricts the negative update, similar to Theorem 3, for sufficiently small  $\beta$ , almost sure state consensus can be guaranteed when the positive graph  $G_{\text{pst}}$  is connected.

In absence of any balance property for the underlying graph, belief clustering may not happen. However, we can establish that when the positive graph is connected, then clustering cannot be achieved when  $\beta$  is large enough. In fact, the belief of a given node touches the two boundaries  $-A$  and  $A$  an infinite number of times. Note that if the positive graph is connected, then the graph cannot be balanced.

**Theorem 6** *Assume that the positive graph  $G_{\text{pst}}$  is connected. For any  $\alpha \in (0, 1) \setminus \{1/2\}$ , when  $\beta$  is sufficiently large, for almost all initial beliefs, under (19), we have: for all  $i \in V$ ,*

$$\mathbb{P}\left(\liminf_{k \rightarrow \infty} x_i(k) = -A, \limsup_{k \rightarrow \infty} x_i(k) = A\right) = 1. \quad (24)$$

Theorem 6 suggests a new class of collective formation for the social beliefs beyond consensus, disagreement, or clustering studied in the literature.

**Remark 4** *The condition that  $\beta$  being sufficiently large in Theorems 4 and 5 is just a technical assumption ensuring almost sure boundary clustering. Practically one can often encounter such clustering even for a small  $\beta$ , as illustrated in the coming numerical examples. On the other hand, Theorem 6 relies crucially on a large  $\beta$ , while a small  $\beta$  leads to belief consensus even in the presence of the negative edges.*

## 5.3 Numerical Examples

We now provide a few numerical examples to illustrate the results established in this section. We take  $A = 1$  so that the node beliefs are restricted to the interval  $[-1, 1]$ . We take  $\alpha = 1/3$  for the positive dynamics and  $a = b = c = 1/3$  for the random asymmetric updates. The pair selection process is given by that when a node  $i$  is drawn, it will choose one of its neighbors with equal probability  $1/\text{deg}(i)$ , where  $\text{deg}(i)$  is the degree of node  $i$  in the underlying graph  $G$ .

First of all we select two social graphs, one is strongly balanced and the other is weakly balanced, as shown in Figure 3. We take  $\beta = 0.2$  and randomly select the nodes' initial values. It is observed that the boundary clustering phenomena established in Theorems 4 and 5 practically show up in every run of the random belief updates. We plot one of their typical sample paths in Figure 4, respectively, for the strongly balanced and weakly balanced graphs in Figure 3. In fact one can see that the clustering is achieved in around 300 steps.




Figure 3: Strongly balanced (left) and weakly balanced (right) social graphs. The negative links are shadowed. Nodes within the same subgraph in the balance partition are marked with the same color.




Figure 4: The evolution of beliefs for strongly balanced (left) and weakly balanced (right) graphs. The beliefs of nodes within the same subgraph in the balance partition are marked with the same color.




Figure 5: A social network which is neither strongly nor weakly balanced. The negative links are dashed.




Figure 6: The social network beliefs tend to a consensus with  $\beta = 0.2$ .

Next, we select a social graph which is neither strongly nor weakly balanced, as in Figure 5. In Figure 6, we plot one of the typical sample paths of the random belief evolution with  $\beta = 0.2$ , where clearly belief consensus is achieved. In Figure 7, we plot one of the typical sample paths of the random belief evolution for a selected node with  $\beta = 7$ , where the node belief alternatively touches the two boundaries  $-1$  and  $1$  in the plotted 5000 steps.

These numerical results are consistent with the results in Theorems 4, 5, and 6.

## 6 Conclusions

The evolution of opinions over signed social networks was studied. Each link marking interpersonal interaction in the network was associated with a sign indicating friend or enemy relations. The dynamics of opinions were defined along positive and negative links, respectively. We have presented a comprehensive analysis to the belief convergence and divergence under various modes: in expectation, in mean-square, and almost surely. Phase transitions were established with sharp thresholds for the mean and mean-




Figure 7: The belief oscillation for a particular node with  $\beta = 7$ .

square convergence. In the almost sure sense, some surprising results were presented. When opinions have hard lower and upper bounds with asymmetric updates, the classical structure balance properties were shown to play a key role in the belief clustering. We believe that these results have largely extended our understanding to how trustful and antagonistic relations shape social opinions.

Some interesting directions for future research include the following topics. Intuitively there is some natural coupling between the structure dynamics and the opinion evolution for signed networks. How this coupling determines the formation of the social structure is an interesting question bridging the studies on the dynamics of signed graphs (e.g., Marvel et al. (2011)) and the opinion dynamics on signed social networks (e.g., Altafini (2012, 2013)). It will also be interesting to ask what might be a proper model, and what the role of structure balance is, for Bayesian opinion evolution on signed social networks (e.g., Bikhchandani et al. (1992)).

## Appendix: Proofs of Statements

### A. The Triangle Lemma

We establish a key technical lemma on the relative beliefs of three nodes in the network in the presence of at least one link among the three nodes. Denote  $J_{ab}(k) := |x_a(k) - x_b(k)|$  for  $a, b \in \mathcal{V}$  and  $k \geq 0$ .

**Lemma 2** *Let  $i_0, i_1, i_2$  be three different nodes in  $\mathcal{V}$ . Suppose  $\{i_0, i_1\} \in \mathcal{E}$ . There exist a positive number  $\delta > 0$  and an integer  $Z > 0$ , such that*

(i) *there is a sequence of  $Z$  successive node pairs leading to  $J_{i_1 i_2}(Z) \geq \delta J_{i_0 i_1}(0)$ ;*

(ii) *there is a sequence of  $Z$  successive node pairs leading to  $J_{i_1 i_2}(Z) \geq \delta J_{i_0 i_2}(0)$ .*

Here  $\delta$  and  $Z$  are absolute constants in the sense that they do not depend on  $i_0, i_1, i_2$ , nor on the values held at these nodes.

*Proof.* We assume  $n \geq 5$ . Generality is not lost by making this assumption because for  $n = 3$  and  $n = 4$ , some (tedious but straightforward) analysis on each possible  $G$  leads to the desired conclusion.

(i). There are two cases:  $\{i_0, i_1\} \in E_{\text{pst}}$ , or  $\{i_0, i_1\} \in E_{\text{neg}}$ . We prove the desired conclusion for each of the two cases. Without loss of generality, we assume that  $x_{i_0}(0) < x_{i_1}(0)$ .

- Let  $\{i_0, i_1\} \in E_{\text{pst}}$ . If  $x_{i_2}(0) \in [\frac{3}{4}x_{i_0}(0) + \frac{1}{4}x_{i_1}(0), \frac{1}{4}x_{i_0}(0) + \frac{3}{4}x_{i_1}(0)]$ , we have  $J_{i_1i_2}(0) \geq \frac{1}{4}J_{i_0i_1}(0)$ . Thus, the desired conclusion holds for  $\delta = \frac{1}{4}$ , arbitrary  $Z > 0$ , and any node pair sequence over  $0, 1, \dots, Z - 1$  for which  $i_0, i_1, i_2$  are never selected.

On the other hand suppose  $x_{i_2}(0) \notin [\frac{3}{4}x_{i_0}(0) + \frac{1}{4}x_{i_1}(0), \frac{1}{4}x_{i_0}(0) + \frac{3}{4}x_{i_1}(0)]$ . Take

$$d_* = \begin{cases} \lceil \log_{|1-2\alpha|} \frac{1}{4} \rceil & \text{if } \alpha \neq \frac{1}{2}, \\ 1, & \text{if } \alpha = \frac{1}{2}. \end{cases} \quad (25)$$

If  $\{i_0, i_1\}$  is selected for  $0, 1, \dots, d_* - 1$ , we obtain  $J_{i_0i_1}(d_*) \leq \frac{1}{4}J_{i_0i_1}(0)$  which leads to

$$x_{i_1}(d_*) \in \left[ \frac{5}{8}x_{i_0}(0) + \frac{2}{8}x_{i_1}(0), \frac{3}{8}x_{i_0}(0) + \frac{5}{8}x_{i_1}(0) \right]; \quad x_{i_2}(d_*) = x_{i_2}(0).$$

This gives us  $J_{i_1i_2}(d_*) \geq \frac{1}{8}J_{i_0i_1}(0)$ .

- Let  $\{i_0, i_1\} \in E_{\text{neg}}$ . If  $x_{i_2}(0) \notin [\frac{1}{2}x_{i_0}(0) + \frac{1}{2}x_{i_1}(0), -\frac{1}{2}x_{i_0}(0) + \frac{3}{2}x_{i_1}(0)]$ , we have  $J_{i_1i_2}(0) \geq \frac{1}{2}J_{i_0i_1}(0)$ . The conclusion holds for  $\delta = \frac{1}{2}$ , arbitrary  $Z > 0$ , and any node pair sequence over  $0, 1, \dots, Z - 1$  for which  $i_0, i_1, i_2$  are never selected.

On the other hand let  $x_{i_2}(0) \in [\frac{1}{2}x_{i_0}(0) + \frac{1}{2}x_{i_1}(0), -\frac{1}{2}x_{i_0}(0) + \frac{3}{2}x_{i_1}(0)]$ . Take  $d^* = \lceil \log_{|1+2\beta|} 4 \rceil$ . Let  $\{i_0, i_1\}$  be selected for  $0, 1, \dots, d^* - 1$ . In this case,  $x_{i_0}(s)$  and  $x_{i_1}(s)$  are symmetric with respect to their center  $\frac{1}{2}x_{i_0}(0) + \frac{1}{2}x_{i_1}(0)$  for all  $s = 0, \dots, d^*$ , and  $J_{i_0i_1}(d^*) \geq 4J_{i_0i_1}(0)$ . Thus we have  $x_{i_2}(d^*) = x_{i_2}(0)$ , and

$$\begin{aligned} x_{i_1}(d^*) &\geq \frac{1}{2}x_{i_0}(0) + \frac{1}{2}x_{i_1}(0) + 2(x_{i_1}(0) - x_{i_0}(0)) \\ &= -\frac{3}{2}x_{i_0}(0) + \frac{5}{2}x_{i_1}(0). \end{aligned} \quad (26)$$

We can therefore conclude that  $J_{i_1i_2}(d^*) \geq J_{i_0i_1}(0)$ .

In summary, the desired conclusion holds for  $\delta = \frac{1}{8}$  and

$$Z = \begin{cases} \max\{\lceil \log_{|1+2\beta|} 4 \rceil, \lceil \log_{|1-2\alpha|} \frac{1}{4} \rceil\} & \text{if } \alpha \neq \frac{1}{2} \\ \lceil \log_{|1+2\beta|} 4 \rceil, & \text{if } \alpha = \frac{1}{2}. \end{cases} \quad (27)$$

(ii). We distinguish the cases  $\{i_0, i_1\} \in \mathbf{E}_{\text{pst}}$  and  $\{i_0, i_1\} \in \mathbf{E}_{\text{neg}}$ . Without loss of generality, we assume that  $x_{i_0}(0) < x_{i_2}(0)$ .

- Let  $\{i_0, i_1\} \in \mathbf{E}_{\text{pst}}$ . If  $x_{i_1}(0) \notin [\frac{1}{2}x_{i_0}(0) + \frac{1}{2}x_{i_2}(0), -\frac{1}{2}x_{i_0}(0) + \frac{3}{2}x_{i_2}(0)]$ , we have  $J_{i_1 i_2}(0) \geq \frac{1}{2}J_{i_0 i_2}(0)$ . The conclusion holds for  $\delta = \frac{1}{2}$ , arbitrary  $Z > 0$ , and any node pair sequence  $0, 1, \dots, Z - 1$  for which  $i_0, i_1, i_2$  are never selected.

Now let  $x_{i_1}(0) \in [\frac{1}{2}x_{i_0}(0) + \frac{1}{2}x_{i_2}(0), -\frac{1}{2}x_{i_0}(0) + \frac{3}{2}x_{i_2}(0)]$ . We write  $x_{i_1}(0) = (1 - \varsigma)x_{i_0}(0) + \varsigma x_{i_2}(0)$  with  $\varsigma \in [\frac{1}{2}, \frac{3}{2}]$ . Let  $\{i_0, i_1\}$  be the node pair selected for  $0, 1, \dots, d_* - 1$  with  $d_*$  defined by (25). Note that according to the structure of the update rule,  $x_{i_0}(s)$  and  $x_{i_1}(s)$  will be symmetric with respect to their center  $(1 - \frac{\varsigma}{2})x_{i_0}(0) + \frac{\varsigma}{2}x_{i_2}(0)$  for all  $s = 0, \dots, d_*$ , and  $J_{i_0 i_1}(d_*) \leq \frac{1}{4}J_{i_0 i_1}(0)$ . This gives us  $x_{i_2}(d_*) = x_{i_2}(0)$  and

$$\begin{aligned} x_{i_1}(d_*) &\in \left[ \left(1 - \frac{\varsigma}{2}\right)x_{i_0}(0) + \frac{\varsigma}{2}x_{i_2}(0) - \frac{1}{8}(x_{i_1}(0) - x_{i_0}(0)), \right. \\ &\quad \left. \left(1 - \frac{\varsigma}{2}\right)x_{i_0}(0) + \frac{\varsigma}{2}x_{i_2}(0) + \frac{1}{8}(x_{i_1}(0) - x_{i_0}(0)) \right] \\ &= \left[ \left(1 - \frac{3\varsigma}{8}\right)x_{i_0}(0) + \frac{3\varsigma}{8}x_{i_2}(0), \left(1 - \frac{5\varsigma}{8}\right)x_{i_0}(0) + \frac{5\varsigma}{8}x_{i_2}(0) \right], \end{aligned} \quad (28)$$

which implies

$$J_{i_1 i_2}(d_*) \geq \left(1 - \frac{5\varsigma}{8}\right)J_{i_0 i_2}(0) \geq \frac{1}{16}J_{i_0 i_2}(0). \quad (29)$$

- Let  $\{i_0, i_1\} \in \mathbf{E}_{\text{neg}}$ . If  $x_{i_1}(0) \notin [\frac{1}{2}x_{i_0}(0) + \frac{1}{2}x_{i_2}(0), -\frac{1}{2}x_{i_0}(0) + \frac{3}{2}x_{i_2}(0)]$ , the conclusion holds for the same reason as in the case where  $\{i_0, i_1\} \in \mathbf{E}_{\text{pst}}$ .

Now let  $x_{i_1}(0) \in [\frac{1}{2}x_{i_0}(0) + \frac{1}{2}x_{i_2}(0), -\frac{1}{2}x_{i_0}(0) + \frac{3}{2}x_{i_2}(0)]$ . We continue to use the notation  $x_{i_1}(0) = (1 - \varsigma)x_{i_0}(0) + \varsigma x_{i_2}(0)$  with  $\varsigma \in [\frac{1}{2}, \frac{3}{2}]$ . Let  $\{i_0, i_1\}$  be the node pair selected for  $0, 1, \dots, d^* - 1$  where  $d^* = \lceil \log_{1+2\beta} 4 \rceil$ . In this case,  $x_{i_0}(s)$  and  $x_{i_1}(s)$  are still symmetric with respect to their center  $(1 - \frac{\varsigma}{2})x_{i_0}(0) + \frac{\varsigma}{2}x_{i_2}(0)$  for all  $s = 0, 1, \dots, d^*$ , and  $J_{i_0 i_1}(d^*) \geq 4J_{i_0 i_1}(0)$ . This gives us  $x_{i_2}(d^*) = x_{i_2}(0)$  and

$$\begin{aligned} x_{i_1}(d^*) &\geq \left(1 - \frac{\varsigma}{2}\right)x_{i_0}(0) + \frac{\varsigma}{2}x_{i_2}(0) + 2(x_{i_1}(0) - x_{i_0}(0)) \\ &= \left(1 - \frac{5\varsigma}{2}\right)x_{i_0}(0) + \frac{5\varsigma}{2}x_{i_2}(0) \end{aligned} \quad (30)$$

which implies

$$J_{i_1 i_2}(d^*) \geq \left(\frac{5\varsigma}{2} - 1\right)J_{i_0 i_2}(0) \geq \frac{1}{4}J_{i_0 i_2}(0). \quad (31)$$

In summary, the desired conclusion holds for  $\delta = \frac{1}{16}$  with  $Z$  defined in (27).

## B. Proof of Theorem 1

Introduce

$$X_{\min}(k) = \min_{i \in \mathbf{V}} x_i(k); \quad X_{\max}(k) = \max_{i \in \mathbf{V}} x_i(k).$$

We define  $\mathfrak{X}(k) = X_{\max}(k) - X_{\min}(k)$ . Suppose belief divergence is achieved almost surely. Take a constant  $N_0$  such that  $N_0 > \mathfrak{X}(0)$ . Then almost surely,

$$K_1 := \inf_k \{\mathfrak{X}(k) \geq N_0\}$$

is a finite number. Then  $K_1$  is a *stopping time* for the node pair selection process  $G_k, k = 0, 1, 2, \dots$  since

$$\{K_1 = k\} \in \sigma(G_0, \dots, G_{k-1})$$

for all  $k = 1, 2, \dots$  due to the fact that  $\mathfrak{X}(k)$  is, indeed, a function of  $G_0, \dots, G_{k-1}$ . Strong Markov Property leads to:  $G_{K_1}, G_{K_1+1}, \dots$  are independent of  $\mathcal{F}_{K_1-1}$ , and they are i.i.d. with the same distribution as  $G_0$  (e.g., Theorem 4.1.3 in Durrett (2010)).

Now take two different (deterministic) nodes  $i_0$  and  $j_0$ . Since  $\mathfrak{X}(K_1) \geq N_0$ , there must be two different (random) nodes  $i_*$  and  $j_*$  satisfying  $x_{i_*}(K_1) < x_{j_*}(K_1)$  with  $J_{i_*j_*}(K_1) \geq N_0$ . We make the following claim.

*Claim.* There exist a positive number  $\delta_0 > 0$  and an integer  $Z_0 > 0$  ( $\delta_0$  and  $Z_0$  are deterministic constants) such that we can always select a sequence of node pairs for time steps  $K_1, K_1 + 1, K_1 + Z_0 - 1$  which guarantees  $J_{i_0j_0}(K_1 + Z_0) \geq \delta_0 N_0$ .

First of all note that  $i_*$  and  $j_*$  are independent with  $G_{K_1}, G_{K_1+1}, \dots$ , since  $i_*, j_* \in \mathcal{F}_{K_1-1}$ . Therefore, we can treat  $i_*$  and  $j_*$  as deterministic and prove the claim for all choices of such  $i_*$  and  $j_*$  (because we can always carry out the analysis conditioned on different events  $\{i_* = i, j_* = j\}, i, j \in \mathbf{V}$ ). We proceed the proof recursively taking advantage of the Triangle Lemma.

Suppose  $\{i_0, j_0\} = \{i_*, j_*\}$ , the claim holds trivially. Now suppose  $i_0 \notin \{i_*, j_*\}$ . Either  $J_{i_0i_*}(K_1) \geq \frac{N_0}{2}$  or  $J_{i_0j_*}(K_1) \geq \frac{N_0}{2}$  must hold. Without loss of generality we assume  $J_{i_0i_*}(K_1) \geq \frac{N_0}{2}$ . Since  $\mathbf{G}$  is connected, there is a path  $i_0i_1 \dots i_\tau j_0$  in  $\mathbf{G}$  with  $\tau \leq n - 2$ .

Based on Lemma 2, there exist  $\delta > 0$  and integer  $Z > 0$  such that a selection of node pair sequence for  $K_1, K_1 + 1, \dots, K_1 + Z - 1$  leads to

$$J_{i_0i_1}(K_1 + Z) \geq \delta J_{i_0i_*}(K_1) \geq \frac{\delta N_0}{2}$$

since  $\{i_0, i_1\} \in \mathbf{E}$ . Applying recursively the Triangle Lemma based on the fact that  $\{i_1, i_2\}, \dots, \{i_\tau, j_0\} \in \mathbf{E}$ , we see that a selection of node pair sequence for  $K_1, K_1 + 1, \dots, K_1 + (\tau + 1)Z - 1$  will give us

$$J_{i_0j_0}(K_1 + (\tau + 1)Z) \geq \delta^{\tau+1} J_{i_0i_*}(K_1) \geq \frac{\delta^{\tau+1} N_0}{2}.$$

Since  $\tau \leq n - 2$ , the claim always holds for  $\delta_0 = \frac{\delta^{n-1}}{2}$  and  $Z_0 = (n - 1)Z$ , independently of  $i_*$  and  $j_*$ .

Therefore, denoting  $p_* = \min\{p_{ij} + p_{ji} : \{i, j\} \in E\}$ , the claim we just proved yields that

$$\mathbb{P}\left(J_{i_0 j_0}(K_1 + (n - 1)Z) \geq \frac{\delta^{n-1}N_0}{2}\right) \geq \left(\frac{p_*}{n}\right)^{(n-1)Z}. \quad (32)$$

We proceed the analysis by recursively defining

$$K_{m+1} := \inf\{k \geq K_m + Z_0 : \mathfrak{X}(k) \geq N_0\}, \quad m = 1, 2, \dots$$

Given that belief divergence is achieved,  $K_m$  is finite for all  $m \geq 1$  almost surely. Thus,

$$\mathbb{P}\left(J_{i_0 j_0}(K_m + Z_0) \geq \frac{\delta^{n-1}N_0}{2}\right) \geq \left(\frac{p_*}{n}\right)^{Z_0}, \quad (33)$$

for all  $m = 1, 2, \dots$ . Moreover, the node pair sequence

$$G_{K_1}, \dots, G_{K_1+Z_0-1}; \dots; G_{K_m}, \dots, G_{K_m+Z_0-1}; \dots$$

are independent and have the same distribution as  $G_0$  (This is due to that  $\mathcal{F}_{K_1} \subseteq \mathcal{F}_{K_1+1} \subseteq \dots \subseteq \mathcal{F}_{K_1+Z_0-1} \subseteq \mathcal{F}_{K_2} \dots$  (cf. Theorem 4.1.4 in Durrett (2010))).

Therefore, we can finally invoke the second Borel-Cantelli Lemma (cf. Theorem 2.3.6 in Durrett (2010)) to conclude that almost surely, there exists an infinite subsequence  $K_{m_s}, s = 1, 2, \dots$ , satisfying

$$J_{i_0 j_0}(K_{m_s} + Z_0) \geq \frac{\delta^{n-1}N_0}{2}, \quad s = 1, 2, \dots, \quad (34)$$

conditioned on that belief divergence is achieved. Since  $\delta$  is a constant and  $N_0$  is arbitrarily chosen, (34) is equivalent to  $\mathbb{P}(\limsup_{k \rightarrow \infty} |x_{i_0}(k) - x_{j_0}(k)| = \infty) = 1$ , which completes the proof.

### C. Proof of Lemma 1

(i). It suffices to show that  $\mathbb{P}(\limsup_{k \rightarrow \infty} \mathfrak{X}(k) \in [a_*, b_*]) = 0$  for all  $0 < a_* < b_*$ . We prove the statement by contradiction. Suppose  $\mathbb{P}(\limsup_{k \rightarrow \infty} \mathfrak{X}(k) \in [a_*, b_*]) = p > 0$  for some  $0 < a_* < b_*$ .

Take  $0 < \varepsilon < 1$  and define  $a = a_*(1 - \varepsilon), b = b_*(1 + \varepsilon)$ . We introduce

$$T_1 := \inf_k \{\mathfrak{X}(k) \in [a, b]\}.$$

Then  $T_1$  is finite with probability at least  $p$ .  $T_1$  is a stopping time.  $G_{T_1}, G_{T_1+1}, \dots$  are independent of  $\mathcal{F}_{T_1-1}$ , and they are i.i.d. with the same distribution as  $G_0$ .

Now since  $G_{\text{neg}}$  is nonempty, we take a link  $\{i_*, j_*\} \in E_{\text{neg}}$ . Repeating the same analysis as the proof of Theorem 1, the following statement holds true conditioned on that  $T_1$  is finite: there exist a positive number  $\delta_0 > 0$  and an integer  $Z_0 > 0$  ( $\delta_0$  and  $Z_0$  are deterministic constants) such that we can always select a sequence of node pairs for time steps  $T_1, T_1 + 1, T_1 + Z_0 - 1$  which guarantees  $J_{i_*, j_*}(T_1 + Z_0) \geq \delta_0 a$ .



Here  $\delta_0$  and  $Z_0$  follow from the same definition in the proof of Theorem 1. Take

$$m_0 = \left\lceil \log_{2\beta+1} \frac{2b}{\delta_0 a} \right\rceil$$

and let  $\{i_*, j_*\}$  be selected for  $T_1 + Z_0, \dots, T_1 + Z_0 + m_0 - 1$ . Then noting that  $\{i_*, j_*\} \in \mathbf{E}_{\text{neg}}$ , the choice of  $m_0$  and the fact that  $J_{i_*, j_*}(s+1) = (2\beta+1)J_{i_*, j_*}(s)$ ,  $s = T_1 + Z_0, \dots, T_1 + Z_0 + m_0 - 1$  lead to

$$\mathfrak{X}(T_1 + Z_0 + m_0) \geq J_{i_*, j_*}(T_1 + Z_0 + m_0) \geq (2\beta+1)^{m_0} \delta_0 a \geq 2b \geq 2b_*.$$

We have proved that

$$\mathbb{P}\left(\mathfrak{X}(T_1 + Z_0 + m_0) \geq 2b_* \mid T_1 < \infty\right) \geq \left(\frac{p_*}{n}\right)^{Z_0 + m_0}. \quad (35)$$

Similarly, we proceed the analysis by recursively defining

$$T_{m+1} := \inf \{k \geq T_m + Z_0 + m_0 : \mathfrak{X}(k) \in [a, b]\}, \quad m = 1, 2, \dots$$

Given  $\mathbb{P}(\limsup_{k \rightarrow \infty} \mathfrak{X}(k) \in [a_*, b_*]) = p$ ,  $T_m$  is finite for all  $m \geq 1$  with probability at least  $p$ . Thus, there holds

$$\mathbb{P}\left(\mathfrak{X}(T_m + Z_0 + m_0) \geq 2b_* \mid T_m < \infty\right) \geq \left(\frac{p_*}{n}\right)^{Z_0 + m_0}, \quad m = 1, 2, \dots \quad (36)$$

The independence of

$$G_{T_1}, \dots, G_{T_1 + Z_0 + m_0 - 1}; \dots; G_{T_m}, \dots, G_{T_m + Z_0 + m_0 - 1}; \dots$$

once again allows us to invoke the Borel-Cantelli Lemma to conclude that almost surely, there exists an infinite subsequence  $T_{m_s}, s = 1, 2, \dots$ , satisfying

$$\mathfrak{X}(T_{m_s} + Z_0 + m_0) \geq 2b_*, \quad s = 1, 2, \dots, \quad (37)$$

given that  $T_m, m = 1, 2, \dots$ , are finite. In other words, we have obtained that

$$\mathbb{P}\left(\limsup_{k \rightarrow \infty} \mathfrak{X}(k) \geq 2b_* \mid \limsup_{k \rightarrow \infty} \mathfrak{X}(k) \in [a_*, b_*]\right) = 1, \quad (38)$$

which is impossible and the first part of the theorem has been proved.

(ii). It suffices to show that  $\mathbb{P}(\liminf_{k \rightarrow \infty} \mathfrak{X}(k) \in [a_*, b_*]) = 0$  for all  $0 < a_* < b_*$ . The proof is again by contradiction. Assume that  $\mathbb{P}(\liminf_{k \rightarrow \infty} \mathfrak{X}(k) \in [a_*, b_*]) = q > 0$ . Let  $a, b$ , and  $T_1 := \inf_k \{\mathfrak{X}(k) \in [a, b]\}$  as defined earlier.  $T_1$  is finite with probability at least  $q$ .

Let  $\ell_0 \in \mathbf{V}$  satisfying  $x_{\ell_0}(T_1) = X_{\min}(T_1)$ . There is a path from  $\ell_0$  to every other node in the network since  $\mathbf{G}_{\text{pst}}$  is connected. We introduce

$$\mathbf{V}_t^\dagger := \{j : d(\ell_0, j) = t \text{ in } \mathbf{G}_{\text{pst}}\}, \quad t = 0, \dots, \text{diam}(\mathbf{G}_{\text{pst}})$$

as a partition of  $V$ . We relabel the nodes in  $V \setminus \{\ell_0\}$  in the following manner.

$$\begin{aligned} \ell_s &\in V_1^\dagger, s = 1, \dots, |V_1^\dagger|; \\ \ell_s &\in V_2^\dagger, s = |V_1^\dagger| + 1, \dots, |V_1^\dagger| + |V_2^\dagger|; \\ &\dots\dots \\ \ell_s &\in V_{\text{diam}(\mathbf{G}_{\text{pst}})}^\dagger, s = \sum_{t=1}^{\text{diam}(\mathbf{G}_{\text{pst}})-1} |V_t^\dagger|, \dots, n-1. \end{aligned}$$

Then the definition of  $V_t^\dagger$  and the connectivity of  $\mathbf{G}_{\text{pst}}$  allow us to select a sequence of node pairs in the form of

$$G_{T_1+s} = \{\ell_\rho, \ell_{s+1}\}, \quad \{\ell_\rho, \ell_{s+1}\} \in \mathbf{E}_{\text{pst}} \text{ with } \rho \leq s,$$

for  $s = 0, \dots, n-2$ . Next we give an estimation for  $\mathfrak{X}$  under the selected sequence of node pairs.

- Since  $\{\ell_0, \ell_1\}$  is selected at time  $T_1$ , we have

$$\begin{aligned} x_{\ell_0}(T_1+1) &= (1-\alpha)x_{\ell_0}(T_1) + \alpha x_{\ell_1}(T_1) \leq (1-\alpha)X_{\min}(T_1) + \alpha X_{\max}(T_1); \\ x_{\ell_1}(T_1+1) &= (1-\alpha)x_{\ell_1}(T_1) + \alpha x_{\ell_0}(T_1) \leq (1-\alpha)X_{\max}(T_1) + \alpha X_{\min}(T_1). \end{aligned} \quad (39)$$

This leads to  $x_{\ell_s}(T_1+1) \leq (1-\alpha_*)X_{\min}(T_1) + \alpha_*X_{\max}(T_1)$ ,  $s = 0, 1$ , where  $\alpha_* = \max\{\alpha, 1-\alpha\}$ .

- Note that  $X_{\max}(T_1+1) = X_{\max}(T_1)$ , and that either  $\{\ell_0, \ell_2\}$  or  $\{\ell_1, \ell_2\}$  is selected at time  $T_1+1$ . We deduce:

$$\begin{aligned} x_{\ell_s}(T_1+2) &\leq (1-\alpha)[(1-\alpha_*)X_{\min}(T_1) + \alpha_*X_{\max}(T_1)] + \alpha X_{\max}(T_1) \\ &\leq (1-\alpha_*)^2 X_{\min}(T_1) + (1-(1-\alpha_*)^2)X_{\max}(T_1), \quad s = 0, 1; \\ x_{\ell_2}(T_1+2) &\leq \alpha[(1-\alpha_*)X_{\min}(T_1) + \alpha_*X_{\max}(T_1)] + (1-\alpha)X_{\max}(T_1) \\ &\leq (1-\alpha_*)^2 X_{\min}(T_1) + (1-(1-\alpha_*)^2)X_{\max}(T_1), \end{aligned} \quad (40)$$

Thus we obtain  $x_{\ell_s}(T_1+2) \leq (1-\alpha_*)^2 X_{\min}(T_1) + (1-(1-\alpha_*)^2)X_{\max}(T_1)$ ,  $s = 0, 1, 2$ .

- We carry on the analysis recursively, and finally get:

$$x_{\ell_s}(T_1+n-1) \leq (1-\alpha_*)^{n-1} X_{\min}(T_1) + (1-(1-\alpha_*)^{n-1})X_{\max}(T_1),$$

for  $s = 0, 1, 2, \dots, n-1$ . Equivalently:

$$X_{\max}(T_1+n-1) \leq (1-\alpha_*)^{n-1} X_{\min}(T_1) + (1-(1-\alpha_*)^{n-1})X_{\max}(T_1). \quad (41)$$

We conclude that:

$$\begin{aligned}
 \mathfrak{X}(T_1 + n - 1) &= X_{\max}(T_1 + n - 1) - X_{\min}(T_1 + n - 1) \\
 &= X_{\max}(T_1 + n - 1) - X_{\min}(T_1) \\
 &\leq r_0 \mathfrak{X}(T_1),
 \end{aligned} \tag{42}$$

where  $r_0 = 1 - (1 - \alpha_*)^{n-1}$  is a constant in  $(0, 1)$ .

With the above analysis taking

$$L_0 = \left\lceil \log_{r_0} \frac{a}{2b} \right\rceil,$$

and selecting the given pair sequence periodically for  $L_0$  rounds, we obtain

$$\mathfrak{X}(T_1 + (n - 1)L_0) \leq r_0^{L_0} \mathfrak{X}(T_1) \leq \frac{a}{2b} \cdot b = \frac{a}{2} < \frac{a_*}{2}. \tag{43}$$

In light of (43) and the selection of the node pair sequence, we have obtained that

$$\mathbb{P}\left(\mathfrak{X}(T_1 + (n - 1)L_0) \leq \frac{a_*}{2}\right) \geq \left(\frac{p_*}{n}\right)^{(n-1)L_0} \tag{44}$$

given that  $T_1$  is finite. We repeat the above argument for  $T_{m+1}$ ,  $m = 2, 3, \dots$ . Borel-Cantelli Lemma then implies

$$\mathbb{P}\left(\liminf_{k \rightarrow \infty} \mathfrak{X}(k) \leq \frac{a_*}{2} \mid \liminf_{k \rightarrow \infty} \bar{\mathfrak{X}}(k) \in [a_*, b_*]\right) = 1, \tag{45}$$

which is impossible and completes the proof.

#### D. Proof of Theorem 2

Let  $\omega \notin \mathcal{C}$ . Then there exists an initial value  $x^0 \in \mathbb{R}^n$  from which

$$\limsup_{k \rightarrow \infty} \mathfrak{X}(k)(\omega) > 0. \tag{46}$$

According to Lemma 1, (46) implies that

$$\mathbb{P}\left(\limsup_{k \rightarrow \infty} \mathfrak{X}(k) = \infty \mid \mathcal{C}^c\right) = \mathbb{P}\left(\mathcal{D} \mid \mathcal{C}^c\right) = 1, \tag{47}$$

which implies  $\mathbb{P}(\mathcal{C}) + \mathbb{P}(\mathcal{D}) = 1$ .

With  $\mathbb{P}(\mathcal{C}) + \mathbb{P}(\mathcal{D}) = 1$ ,  $\mathcal{D}$  is a trivial event as long as  $\mathcal{C}$  is a trivial event. Therefore, for completing the proof we just need to verify that  $\mathcal{C}$  is a trivial event.

We first show that  $\mathcal{C} = \{\lim_{k \rightarrow \infty} W_k \dots W_0 = U\}$ . In fact, if  $\limsup_{k \rightarrow \infty} \max_{i,j} |x_i(k) - x_j(k)| = 0$  under  $x^0 \in \mathbb{R}^n$ , then we have  $\lim_{k \rightarrow \infty} x(k) = \frac{1}{n} \mathbf{1} \mathbf{1}' x^0$  because the sum of the beliefs is preserved.

Therefore, we can restrict the analysis to  $x^0 = e_i$ ,  $i = 1, \dots, n$  and one can readily see that  $\mathcal{C} = \{\lim_{k \rightarrow \infty} W_k \dots W_0 = U\}$ .

Next, we apply the argument, which was originally introduced in Tahbaz-Salehi and Jadbabaie (2008) for establishing the weak ergodicity of product of random stochastic matrices with positive diagonal terms, to conclude that  $\mathcal{C}$  is a trivial event. A more general treatment to zero-one laws of random averaging algorithms can be found in Touri and Nedić (2011). Define a sequence of event  $\mathcal{C}_s = \{\lim_{k \rightarrow \infty} W_k \dots W_s = U\}$  for  $s = 1, 2, \dots$ . We see that

- $\mathbb{P}(\mathcal{C}_s) = \mathbb{P}(\mathcal{C})$  for all  $s = 1, 2, \dots$  since  $W_k, k = 0, 1, \dots$ , are i.i.d.
- $\mathcal{C}_{s+1} \subseteq \mathcal{C}_s$  for all  $s = 1, 2, \dots$  since  $\lim_{k \rightarrow \infty} W_k \dots W_{s+1} = U$  implies  $\lim_{k \rightarrow \infty} W_k \dots W_s = U$  due to the fact that  $UW_s \equiv U$ .

Therefore, we have  $\bigcap_{s=1}^{\infty} \mathcal{C}_s$  is a tail event within the tail  $\sigma$ -field  $\bigcap_{s=1}^{\infty} \sigma(G_s, G_{s+1}, \dots)$ . By Kolmogorov's zero-one law,  $\bigcap_{s=1}^{\infty} \mathcal{C}_s$  is a trivial event. Hence  $\mathbb{P}(\mathcal{C}) = \lim_{s \rightarrow \infty} \mathbb{P}(\mathcal{C}_s) = \mathbb{P}(\bigcap_{s=1}^{\infty} \mathcal{C}_s)$  is a trivial event, and the desired conclusion follows.

## E. Proof of Theorem 3

Theorem 3 is a direct consequence of the following lemmas.

**Lemma 3** *Suppose  $G_{\text{pst}}$  is connected. Then for every fixed  $\alpha \in (0, 1)$ , we have  $\mathbb{P}(\mathcal{C}) = 1$  for all  $0 \leq \beta < \beta^\ddagger$  with*

$$\beta^\ddagger := \sup \left\{ \beta : \beta(1 + \beta) < \frac{\lambda_2(L_{\text{pst}}^\dagger)}{\lambda_{\max}(L_{\text{neg}}^\dagger)} \alpha(1 - \alpha) \right\}.$$

*Proof.* Let  $x_{\text{ave}} = \sum_{i \in \mathcal{V}} x_i(0)/n$  be the average of the initial beliefs. We introduce  $V(k) = \sum_{i=1}^n |x_i(k) - x_{\text{ave}}|^2 = \|(I - U)x(k)\|^2$ . The evolution of  $V(k)$  follows from

$$\begin{aligned} \mathbb{E}\{V(k+1) | x(k)\} &= \mathbb{E}\{x(k+1)'(I - U)^2 x(k+1) | x(k)\} \\ &\stackrel{a)}{=} \mathbb{E}\{x(k)'W(k)(I - U)W(k)x(k) | x(k)\} \\ &\stackrel{b)}{=} \mathbb{E}\{x(k)'(I - U)[W(k)(I - U)W(k)](I - U)x(k) | x(k)\} \\ &\stackrel{c)}{\leq} \lambda_{\max}(\mathbb{E}\{W(k)(I - U)W(k)\}) \|(I - U)x(k)\|^2 \\ &\stackrel{d)}{=} \lambda_{\max}(\mathbb{E}\{W^2(k)\} - U)V(k), \end{aligned} \tag{48}$$

where *a)* is based on the facts that  $W(k)$  is symmetric and the simple fact  $(I - U)^2 = I - U$ , *b)* holds because  $(I - U)W(k) = W(k)(I - U)$  always holds and again  $(I - U)^2 = I - U$ , *c)* follows from Rayleigh-Ritz theorem (cf. Theorem 4.2.2 in Horn and Johnson (1985)) and the fact that  $W(k)$  is independent of  $x(k)$ , *d)* is based on simple algebra and  $W(k)U = UW(k) = U$ .

We now compute  $\mathbb{E}(W^2(k))$ . Note that

$$\begin{aligned} (I - \alpha(e_i - e_j)(e_i - e_j)')^2 &= I - 2\alpha(1 - \alpha)(e_i - e_j)(e_i - e_j)'; \\ (I + \beta(e_i - e_j)(e_i - e_j)')^2 &= I + 2\beta(1 + \beta)(e_i - e_j)(e_i - e_j)'. \end{aligned} \quad (49)$$

This observation combined with (5) leads to

$$\begin{aligned} \mathbb{P}\left(W^2(k) = I - 2\alpha(1 - \alpha)(e_i - e_j)(e_i - e_j)'\right) &= \frac{p_{ij} + p_{ji}}{n}, \quad \{i, j\} \in \mathbf{E}_{\text{pst}}; \\ \mathbb{P}\left(W^2(k) = I + 2\beta(1 + \beta)(e_i - e_j)(e_i - e_j)'\right) &= \frac{p_{ij} + p_{ji}}{n}, \quad \{i, j\} \in \mathbf{E}_{\text{neg}}. \end{aligned}$$

As a result, we have

$$\mathbb{E}\{W^2(k)\} = I - 2\alpha(1 - \alpha)L_{\text{pst}}^\dagger + 2\beta(1 + \beta)L_{\text{neg}}^\dagger. \quad (50)$$

Consequently, we have  $0 < \gamma := \lambda_{\max}(\mathbb{E}(W^2(k)) - U) < 1$  for all  $\beta$  satisfying

$$\beta(1 + \beta) < \frac{\lambda_2(L_{\text{pst}}^\dagger)}{\lambda_{\max}(L_{\text{neg}}^\dagger)}\alpha(1 - \alpha). \quad (51)$$

Since  $g(\beta) = \beta(1 + \beta)$  is nondecreasing, we conclude from (48) that

$$\mathbb{E}\{V(k+1)|x(k)\} < \gamma V(k) \quad (52)$$

with  $0 < \gamma < 1$  for all  $0 \leq \beta < \beta^\ddagger$ . This means that  $V(k)$  is a supermartingale as long as  $0 \leq \gamma \leq 1$  Durrett (2010), and  $V(k)$  converges to a limit almost surely by the martingale convergence theorem (Theorem 5.2.9, Durrett (2010)). Next we show that this limit is zero almost surely if  $0 \leq \gamma < 1$ . Let  $\epsilon > 0$  and  $0 \leq \gamma < 1$ . We have:

$$\begin{aligned} \mathbb{P}\left(V(k) > \epsilon \text{ infinitely often}\right) &\stackrel{a)}{=} \mathbb{P}\left(\sum_{k=0}^{\infty} \mathbb{P}(V(k+1) > \epsilon | x(k)) = \infty\right) \\ &\stackrel{b)}{\leq} \mathbb{P}\left(\frac{1}{\epsilon} \sum_{k=0}^{\infty} \mathbb{E}\{V(k+1)|x(k)\} = \infty\right) \\ &\stackrel{c)}{\leq} \mathbb{P}\left(\frac{\gamma}{\epsilon} \sum_{k=1}^{\infty} V(k) = \infty\right), \end{aligned} \quad (53)$$

where  $a)$  is straightforward application of the Second Borel-Cantelli Lemma (Theorem 5.3.2. in Durrett (2010)),  $b)$  is from the Markov's inequality, and  $c)$  holds directly from (52). Observing that

$$\sum_{k=1}^{\infty} \mathbb{E}\{V(k)\} \leq \sum_{k=1}^{\infty} \gamma^k V(0) \leq \frac{\gamma}{1 - \gamma} V(0) < \infty, \quad (54)$$

we obtain  $\mathbb{P}\left(\frac{\gamma}{\epsilon} \sum_{k=1}^{\infty} V(k) = \infty\right) = 0$ . Therefore, we have proved that  $\mathbb{P}(V(k) > \epsilon \text{ infinitely often}) = 0$ , or equivalently,  $\mathbb{P}(\lim_{k \rightarrow \infty} V(k) = 0) = 1$ .

Finally, observe that:

$$V(k) = \sum_{i=1}^n |x_i(k) - x_{\text{ave}}|^2 \geq |x_{\rho_1}(k) - x_{\text{ave}}|^2 + |x_{\rho_2}(k) - x_{\text{ave}}|^2 \geq \frac{1}{2} |x_{\rho_1}(k) - x_{\rho_2}(k)|^2 = \frac{1}{2} \mathfrak{X}^2(k),$$

where  $\rho_1$  and  $\rho_2$  are chosen such that  $x_{\rho_1}(k) = X_{\min}(k)$ ,  $x_{\rho_2}(k) = X_{\max}(k)$ . Hence  $\mathbb{P}(\lim_{k \rightarrow \infty} V(k) = 0) = 1$  implies  $\mathbb{P}(\lim_{k \rightarrow \infty} \mathfrak{X}(k) = 0) = 1$ . This completes the proof.  $\square$

**Remark 5** *We have shown that:*

$$\mathbb{E}\{V(k+1)\} \leq \lambda_{\max}(\mathbb{E}\{W^2(k)\} - U)\mathbb{E}\{V(k)\} \quad (55)$$

from (48). A symmetric analysis leads to:

$$\mathbb{E}\{V(k+1)\} \geq \lambda_{\min}(\mathbb{E}\{W^2(k)\} - U)\mathbb{E}\{V(k)\}. \quad (56)$$

Proposition 3 readily follows from these inequalities.

**Lemma 4** *Suppose  $\alpha \in [0, 1]$  with  $\alpha \neq 1/2$ . There exists a constant  $\beta^\sharp > 0$  such that*

$$\mathbb{P}(\liminf_{k \rightarrow \infty} \max_{i,j} |x_i(k) - x_j(k)| = \infty) = 1$$

for almost all initial beliefs when  $\beta > \beta^\sharp$ .

*Proof.* Suppose  $\mathfrak{X}(0) > 0$ . We have:

$$J_{ij}(k+1) = \begin{cases} |2\alpha - 1|J_{ij}(k), & \text{if } G_k = \{i, j\} \in \mathbf{E}_{\text{pst}} \\ |2\beta + 1|J_{ij}(k), & \text{if } G_k = \{i, j\} \in \mathbf{E}_{\text{neg}}. \end{cases} \quad (57)$$

Thus,  $\mathfrak{X}(k) > 0$  almost surely for all  $k$  as long as  $\mathfrak{X}(0) > 0$ . As a result, the following sequence of random variables is well defined:

$$\zeta_k = \frac{\mathfrak{X}(k+1)}{\mathfrak{X}(k)}, \quad k = 0, 1, \dots \quad (58)$$

The proof is based on the analysis of  $\zeta_k$ . We proceed in three steps.

Step 1. In this step, we establish some natural upper and lower bounds for  $\zeta_k$ . First of all, from (57), it is easy to see that:

$$\mathbb{P}\left(\zeta_k = \frac{\mathfrak{X}(k+1)}{\mathfrak{X}(k)} \geq |2\alpha - 1|\right) = 1 \quad (59)$$

and  $\mathbb{P}(\zeta_k < 1) \leq \mathbb{P}(\text{one link in } \mathbf{E}_{\text{pst}} \text{ is selected})$ .

On the other hand let  $\{i_0, j_0\} \in \mathbf{G}_{\text{neg}}$ . Suppose  $i_\star$  and  $j_\star$  are two nodes satisfying  $J_{i_\star j_\star} = \mathfrak{X}(0)$ . Repeating the analysis in the proof of Theorem 1 by recursively applying the Triangle Lemma, we conclude that there is a sequence of node pairs for time steps  $0, 1, \dots, (n-1)Z - 1$  which guarantees

$$J_{i_0 j_0}((n-1)Z) \geq \frac{\delta^{n-1}}{2} \mathfrak{X}(0) \quad (60)$$

where  $\delta = 1/16$  and  $Z = \max\{\lceil \log_{1+2\beta} 4 \rceil, \lceil \log_{|1-2\alpha|} \frac{1}{4} \rceil\}$  are defined in the Triangle Lemma. For the remaining of the proof we assume that  $\beta$  is sufficiently large so that  $\lceil \log_{1+2\beta} 4 \rceil \leq \lceil \log_{|1-2\alpha|} \frac{1}{4} \rceil$ , which means that we can select  $Z = \lceil \log_{|1-2\alpha|} \frac{1}{4} \rceil$  independently of  $\beta$ .

Now take an integer  $H_0 \geq 1$ . Continuing the previous node pair sequence, let  $\{i_0, j_0\}$  be selected at time steps  $(n-1)Z, \dots, (n-1)Z + H_0 - 1$ . It then follows from (57) and (60) that

$$\mathfrak{X}((n-1)Z + H_0) \geq J_{i_0 j_0}((n-1)Z + H_0) \geq \frac{(2\beta + 1)^{H_0} \delta^{n-1}}{2} \mathfrak{X}(0). \quad (61)$$

Denote  $Z_{H_0} = (n-1)Z + H_0$ . This node sequence for  $0, 1, \dots, Z_{H_0}$ , which leads to (61), is denoted  $\mathbf{S}_{i_0 j_0}([0, Z_{H_0}])$ .

Step 2. We now define a random variable  $Q_{Z_{H_0}}(0)$ , associated with the node pair selection process in steps  $0, \dots, Z_{H_0} - 1$ , by

$$Q_{Z_{H_0}}(0) = \begin{cases} |2\alpha - 1|^{Z_{H_0}}, & \text{if at least one link in } \mathbf{E}_{\text{pst}} \text{ is selected in steps } 0, 1, \dots, Z_{H_0} - 1; \\ \frac{(2\beta+1)^{H_0} \delta^{n-1}}{2}, & \text{if node sequence } \mathbf{S}_{i_0 j_0}([0, Z_{H_0}]) \text{ is selected in steps } 0, 1, \dots, Z_{H_0} - 1; \\ 1, & \text{otherwise.} \end{cases} \quad (62)$$

In view of (59) and (61), we have:

$$\mathbb{P}\left(\prod_{k=0}^{Z_{H_0}-1} \zeta_k = \frac{\mathfrak{X}(Z_{H_0})}{\mathfrak{X}(0)} \geq Q_{Z_{H_0}}(0)\right) = 1. \quad (63)$$

From direct calculation based on the definition of  $Q_{Z_{H_0}}(0)$ , we conclude that

$$\begin{aligned} \mathbb{E}\left\{\log Q_{Z_{H_0}}(0)\right\} &\geq \left(\frac{p^*}{n}\right)^{Z_{H_0}} \log \frac{(2\beta + 1)^{H_0} \delta^{n-1}}{2} + \left(1 - \left(1 - \frac{p^*}{n}\right)^{E_0 Z_{H_0}}\right) \log |2\alpha - 1|^{Z_{H_0}} \\ &:= C_{H_0} \end{aligned} \quad (64)$$

where  $p^* = \max\{p_{ij} + p_{ji} : \{i, j\} \in \mathbf{E}\}$  and  $E_0 = |\mathbf{E}_{\text{pst}}|$  denotes the number of positive links. Since  $Z$  does not depend on  $\beta$ , we see from (64) that for any fixed  $H_0$ , there is a constant  $\beta^\diamond(H_0) > 0$  with  $\lceil \log_{1+2\beta^\diamond} 4 \rceil \leq \lceil \log_{|1-2\alpha|} \frac{1}{4} \rceil$  guaranteeing that

$$\beta > \beta^\diamond(H_0) \Rightarrow C_{H_0} > 0.$$

Step 3. Recursively applying the analysis in the previous steps, node pair sequences  $S_{i_0 j_0}([sZ_{H_0}, (s+1)Z_{H_0}])$  can be found for  $s = 1, 2, \dots$ , and  $Q_{Z_{H_0}}(s), s = 1, 2, \dots$  can be defined associated with the node pair selection process (following the same definition of  $Q_{Z_{H_0}}(0)$ ). Since the node pair selection process is independent of time and node states,  $Q_{Z_{H_0}}(s), s = 0, 1, 2, \dots$ , are independent random variables (not necessarily i.i.d since  $S_{i_0 j_0}([sZ_{H_0}, (s+1)Z_{H_0}])$  may correspond to different pair sequences for different  $s$ .) The lower bound established in (64) holds for all  $s$ , i.e.,

$$\mathbb{E}\{\log Q_{Z_{H_0}}(s)\} \geq C_{H_0}, s = 0, 1, \dots \quad (65)$$

Moreover, we can prove as (63) was established that:

$$\mathbb{P}\left(\prod_{k=0}^{tZ_{H_0}-1} \zeta_k = \frac{\mathfrak{X}(tZ_{H_0})}{\mathfrak{X}(0)} \geq \prod_{s=0}^{t-1} Q_{Z_{H_0}}(s), t = 0, 1, 2, \dots\right) = 1. \quad (66)$$

It is straightforward to see that  $\mathbb{V}\{\log Q_{Z_{H_0}}(s)\}, s = 0, 1, \dots$  is bounded uniformly in  $s$ . Kolmogorov's strong law of large numbers (for a sequence of mutually independent random variables under Kolmogorov criterion, see Feller (1968)) implies that:

$$\mathbb{P}\left(\lim_{t \rightarrow \infty} \frac{1}{t} \sum_{s=0}^t \left(\log Q_{Z_{H_0}}(s) - \mathbb{E}\{\log Q_{Z_{H_0}}(s)\}\right) = 0\right) = 1. \quad (67)$$

Using (65), (67) further implies that:

$$\mathbb{P}\left(\liminf_{t \rightarrow \infty} \frac{1}{t} \sum_{s=0}^t \log Q_{Z_{H_0}}(s) \geq C_{H_0}\right) = 1. \quad (68)$$

The final part of the proof is based on (66). With the definition of  $\zeta_k$ , (66) yields:

$$\mathbb{P}\left(\log \mathfrak{X}((t+1)Z_{H_0}) - \log \mathfrak{X}(0) = \sum_{k=0}^{(t+1)Z_{H_0}-1} \log \zeta_k \geq \sum_{s=0}^t \log Q_{Z_{H_0}}(s), t = 0, 1, 2, \dots\right) = 1,$$

which together with (68) gives us:

$$\mathbb{P}\left(\liminf_{t \rightarrow \infty} \mathfrak{X}((t+1)Z_{H_0}) = \infty\right) = 1. \quad (69)$$

We can further conclude that:

$$\mathbb{P}\left(\liminf_{k \rightarrow \infty} \mathfrak{X}(k) = \infty\right) = 1 \quad (70)$$

since  $\mathbb{P}(\mathfrak{X}(k) \geq |2\alpha - 1|^{Z_{H_0}} \mathfrak{X}(\lceil \frac{k}{Z_{H_0}} \rceil Z_{H_0})) = 1$  in view of (59).

Therefore, for any integer  $H_0 \geq 1$ , we have proved that belief divergence is achieved for all initial condition satisfying  $\mathfrak{X}(0) > 0$  if  $\beta > \beta^\diamond(H_0)$ . Define

$$\beta^\sharp := \inf_{H_0 \geq 1} \beta^\diamond(H_0).$$

With this choice of  $\beta^\sharp$ , the desired conclusion holds.  $\square$



## F. Proof of Proposition 5

Note that there exist  $\{i_s, j_s\} \in \mathbf{G}_{\text{pst}}$ ,  $s = 1, 2, \dots, T$  with  $T \geq 1$  such that

$$\mathbf{W}_{i_T j_T}^+ \cdots \mathbf{W}_{i_1 j_1}^+ = \mathbf{U} \quad (71)$$

if and only if for any  $y(0) = y^0 = (y_1^0 \dots y_n^0)'$ , the dynamical system

$$y(k) = \mathbf{W}_{i_k j_k}^+ y(k-1), \quad k = 1, \dots, T \quad (72)$$

drives  $y(k) = (y_1(k), \dots, y_n(k))'$  to  $y(T) = \text{ave}(y(0))\mathbf{1}$  where  $\text{ave}(y(0)) = \sum_{i=1}^n y_i^0/n$ . Thus we may study the matrix equality (71) through individual node dynamics, which we leverage in the proof.

The claim follows from an induction argument. Assume that the desired sequence of node pairs with length  $T_k = k2^{k-1}$  exists for  $m = k$ . Assume that  $\mathbf{G}_{\text{pst}}$  has a subgraph isomorphic to an  $m+1$  dimensional hypercube. Without loss of generality we assume  $\mathbf{V}$  has been rewritten as  $\{0, 1\}^{k+1}$  following the definition of hypercube.

Now define

$$\mathbf{V}_0^\dagger := \{i_1 \times \cdots \times i_{k+1} \in \mathbf{V} : i_{k+1} = 0\}; \quad \mathbf{V}_1^\dagger := \{i_1 \times \cdots \times i_{k+1} \in \mathbf{V} : i_{k+1} = 1\}.$$

It is easy to see that each of the subgraphs  $\mathbf{G}_{\mathbf{V}_0^\dagger}$  and  $\mathbf{G}_{\mathbf{V}_1^\dagger}$  contains a positive subgraph isomorphic with an  $m$ -dimensional hypercube. Therefore, for any initial value of  $y(0)$ , the nodes in each set  $\mathbf{G}_{\mathbf{V}_s^\dagger}$ ,  $s = 0, 1$  can reach the same value, say  $C^0(y(0))$  and  $C^1(y(0))$ , respectively. Then we select the following  $2^k$  edges for updates from  $\mathbf{G}$ :

$$\{i_1 \times \cdots \times i_k \times 0, i_1 \times \cdots \times i_k \times 1\} : i_s \in \{0, 1\}, s = 1, \dots, k.$$

After these updates, all nodes reach the same value  $(C^0(y(0)) + C^1(y(0)))/2$  which has to be  $\text{ave}(y(0))$  since the sum of the node beliefs is constant during this process. Thus, the desired sequence of node pairs exists also for  $m = k + 1$ , with a length

$$T_{k+1} = 2T_k + 2^k = 2k2^{k-1} + 2^k = (k+1)2^k.$$

This proves the desired conclusion.

## G. Proof of Proposition 6

The requirement of  $\alpha = 1/2$  is obvious since otherwise  $\mathbf{W}_{ij}^+$  is nonsingular for all  $\{i, j\} \in \mathbf{E}_{\text{pst}}$ , while  $\text{rank} \mathbf{U} = 1$ . The necessity of  $m = 2^k$  for some  $k \geq 0$  was proved in Shi et al. (2014) through an elementary

number theory argument by constructing a particular initial value for which finite-time convergence can never be possible by pairwise averaging.

It remains to show that  $\mathbf{G}_{\text{pst}}$  has a perfect matching. Now suppose Eq. (71) holds. Without loss of generality we assume that Eq. (71) is *irreducible* in the sense that the equality will no longer hold if any (one or more) matrices are removed from that sequence. The idea of the proof is to analyze the dynamical system (72) backwards from the final step. In this way we will recover a perfect matching from  $\{\{i_1, j_1\}, \dots, \{i_T, j_T\}\}$ . We divide the remaining of the proof into three steps.

Step 1. We first establish some property associated with  $\{i_T, j_T\}$ . After the last step in (72), two nodes  $i_T$  and  $j_T$  reach the same value,  $\text{ave}(y^0)$ , along with all the other nodes. We can consequently write

$$y_{i_T}(T-1) = \text{ave}(y^0) + h_T(y^0), \quad y_{j_T}(T-1) = \text{ave}(y^0) - h_T(y^0),$$

where  $h_T(\cdot)$  is a real-valued function marking the error between  $y_{i_T}(T-1)$ ,  $y_{j_T}(T-1)$  and the true average  $\text{ave}(y^0)$ .

Indeed, the set  $\{y^0 : h_T(y^0) = 0\}$  is explicitly given by

$$\left\{ y^0 : (0 \dots \underbrace{1}_{i_T\text{'th}} \dots \underbrace{-1}_{j_T\text{'th}} \dots 0) W_{i_{T-1}j_{T-1}}^+ \dots W_{i_1j_1}^+ y^0 = 0 \right\},$$

which is a linear subspace with dimension  $n-1$  (recall that the equation  $W_{i_Tj_T}^+ \dots W_{i_1j_1}^+ = U$  is irreducible). Thus there must be  $h_T(y^0) \neq 0$  for some initial value  $y^0$ .

Step 2. If there are only two nodes in the network, we are done. Otherwise  $\{i_{T-1}, j_{T-1}\} \neq \{i_T, j_T\}$ . We make the following claim.

*Claim.*  $i_{T-1}, j_{T-1} \notin \{i_T, j_T\}$ .

Suppose without loss of generality that  $i_{T-1} = i_T$ . Then

$$y_{j_{T-1}}(T) = y_{j_{T-1}}(T-1) = y_{i_{T-1}}(T-1) = y_{i_T}(T-1) = \text{ave}(y^0) + h_T(y^0).$$

While on the other hand  $y_{j_{T-1}}(T) = \text{ave}(y^0)$  for all  $y^0$ . The claim holds observing that as we just established,  $\{h_T(y^0) \neq 0\}$  is a nonempty set.

We then write:

$$y_{i_{T-1}}(T-2) = \text{ave}(y^0) + h_{T-1}(y^0), \quad y_{j_{T-1}}(T-2) = \text{ave}(y^0) - h_{T-1}(y^0)$$

where  $h_{T-1}(\cdot)$  is again a real-valued function and  $h_{T-1}(y^0) \neq 0$  for some initial value  $y^0$  (applying the same argument as for  $h_T(y^0) \neq 0$ ). Note that

$$\{y^0 : h_T(y^0) \neq 0\} \cap \{y^0 : h_{T-1}(y^0) \neq 0\} = \left( \{y^0 : h_T(y^0) = 0\} \cup \{y^0 : h_{T-1}(y^0) = 0\} \right)^c$$

is nonempty because it is the complement of the union of two linear subspaces of dimension  $n - 1$  in  $\mathbb{R}^n$ .

Step 3. Again, if there are only four nodes in the network, we are done. Otherwise, we can define:

$$T_\star := \max \left\{ \tau : \{i_\tau, j_\tau\} \not\subseteq \{i_{T-1}, j_{T-1}, i_T, j_T\} \right\} \quad (73)$$

We emphasize that  $T_\star$  must exist since Eq. (71) holds. As before, we have  $i_{T_\star}, j_{T_\star} \notin \{i_{T-1}, j_{T-1}, i_T, j_T\}$  and  $h_{T_\star}(y^0)$  can be found with  $\{h_{T_\star}(y^0) = 0\}$  being another  $(n - 1)$ -dimensional subspace such that

$$y_{i_{T_\star}}(T_\star - 1) = \text{ave}(y^0) + h_{T_\star}(y^0), \quad y_{j_{T_\star}}(T_\star - 1) = \text{ave}(y^0) - h_{T_\star}(y^0).$$

We thus conclude that this argument can be proceeded recursively until we have found a perfect matching of  $\mathbf{G}_{\text{pst}}$  in  $\{\{i_1, j_1\}, \dots, \{i_T, j_T\}\}$ . We have now completed the proof.

## H. Proof of Theorem 4

We first state and prove intermediate lemmas that will be useful for the proofs of Theorems 4, 5, and 6.

**Lemma 5** *Assume that  $\alpha \in (0, 1)$ . Let  $i_1 \dots i_k$  be a path in the positive graph, i.e.,  $\{i_s, i_{s+1}\} \in \mathbf{G}_{\text{pst}}, s = 1, \dots, k - 1$ . Take a node  $i_\star \in \{i_1, \dots, i_k\}$ . Then for any  $\varepsilon > 0$ , there always exists an integer  $Z_\star(\varepsilon) \geq 1$ , such that we can select a sequence of node pairs from  $\{i_s, i_{s+1}\}, s = 1, \dots, k - 1$  under asymmetric updates which guarantees*

$$J_{i_\star i_s}(Z_\star) \leq 2A\varepsilon, \quad s \in \{1, \dots, k\}$$

for all initial condition  $x_{i_s}(0), s = 1, \dots, k$ .

*Proof.* The proof is easy and an appropriate sequence of node pairs can be built just observing that  $J_{i_\star i_s} \leq 2A$  for all  $s \in \{1, \dots, k\}$ .  $\square$

**Lemma 6** *Fix  $\alpha \in (0, 1)$  with  $\alpha \neq 1/2$ . Under belief dynamics (19), there exist an integer  $Z_0 \geq 1$  and a constant  $\vartheta_0 > 0$  such that*

$$\mathbb{P}(\exists \{i_\star, j_\star\} \in \mathbf{G}_{\text{neg}} \text{ s.t. } J_{i_\star j_\star}(Z_0) \geq \frac{1}{2n} \mathfrak{X}(0)) \geq \vartheta_0. \quad (74)$$

*Proof.* We can always uniquely divide  $\mathbf{V}$  into  $m_0 \geq 1$  mutually disjoint sets  $V_1, \dots, V_{m_0}$  such that  $\mathbf{G}_{\text{pst}}(V_k), k = 1, \dots, m_0$  are connected graphs, where  $\mathbf{G}_{\text{pst}}(V_k)$  is the induced graph of  $\mathbf{G}_{\text{pst}}$  by node set  $V_k$ . The idea is to treat each  $\mathbf{G}_{\text{pst}}(V_k)$  as a *super node* (an illustration of this partition is shown in Figure 8). Since  $\mathbf{G}$  is connected and  $\mathbf{G}_{\text{neg}}$  is nonempty, these super nodes form a connected graph whose edges are negative.




Figure 8: There is a unique partition of  $G$  into subgraphs following the connected components of  $G_{\text{pst}}$ . Viewing each subgraph as a super node, the graph is connected, and has only negative edges.

One can readily show that there exist two distinct nodes  $\eta_1, \eta_2 \in V$  with  $\eta_i \in V_{\nu_i}, i = 1, 2$  ( $V_{\nu_1}$  and  $V_{\nu_2}$  can be the same, of course) such that there is at least one *negative* edge between  $V_{\nu_1}$  and  $V_{\nu_2}$  and such that:

$$J_{\eta_1 \eta_2}(0) \geq \frac{1}{m_0} \mathfrak{X}(0). \quad (75)$$

Now select  $v_1 \in V_{\nu_1}$  and  $v_2 \in V_{\nu_2}$  such that  $\{v_1, v_2\} \in E_{\text{neg}}$ . In view of Lemma 5 and observing that asymmetric updates happen with a strictly positive probability, we can always find  $\vartheta_0 > 0$  and  $Z_0 \geq 1$  (both functions of  $(\alpha, n, a, b, c)$ ) such that:

$$\mathbb{P}\left(x_{\nu_i}(Z_0) = x_{\nu_i}(0), J_{v_i \nu_i}(Z_0) \leq \frac{1}{4n} \mathfrak{X}(0), i = 1, 2\right) \geq \vartheta_0, \quad (76)$$

(because  $G_{\text{pst}}(V_{\nu_i}), i = 1, 2$  are connected graphs). (74) follows from (75) and (76) since  $m_0 \leq n$ .  $\square$

**Lemma 7** Fix  $\alpha \in (0, 1)$  with  $\alpha \neq 1/2$ . Under belief dynamics (19), there exists  $\beta^\diamond(\alpha) > 0$  such that  $\mathbb{P}(\limsup_{k \rightarrow \infty} \mathfrak{X}(k) = 2A) = 1$  for almost all initial beliefs if  $\beta > \beta^\diamond$ .

*Proof.* In view of Lemma 6, we have:

$$\mathbb{P}\left(\mathfrak{X}(Z_0 + t) \geq \min\left\{\frac{(\beta + 1)^t}{2n} \mathfrak{X}(0), 2A\right\}\right) \geq \left(\frac{c p^*}{n}\right)^t \vartheta_0, \quad t = 0, 1, \dots \quad (77)$$

We can conclude that:

$$\mathbb{P}(\limsup_{k \rightarrow \infty} \mathfrak{X}(k) = 2A) + \mathbb{P}(\limsup_{k \rightarrow \infty} \mathfrak{X}(k) = 0) = 1 \quad (78)$$

as long as  $\beta > 0$  using the same argument as that used in the proof of statement (i) in Lemma 1.

With (77), we have:

$$\mathbb{P}\left(\mathfrak{X}(Z_0 + 1) \geq \frac{\beta + 1}{2n} \mathfrak{X}(0)\right) \geq \frac{cP^*}{n} \vartheta_0 \quad (79)$$

conditioned on  $\mathfrak{X}(0) \leq 4An/(1 + \beta)$ . Moreover, (59) still holds for belief dynamics (19). Therefore, we can invoke exactly the same argument as that used in the proof of Lemma 4 to conclude that there exists  $\beta^\circ(\alpha) > 0$  such that

$$\mathbb{P}\left(\limsup_{k \rightarrow \infty} \mathfrak{X}(k) \geq 4An/(1 + \beta)\right) = 1 \quad (80)$$

for all  $\beta > \beta^\circ(\alpha)$ . Combining (78) and (80), we get the desired result.  $\square$

**Lemma 8** *Assume that the graph is strongly balanced under partition  $\mathbf{V} = \mathbf{V}_1 \cup \mathbf{V}_2$ , and that  $\mathbf{G}_{\mathbf{V}_1}$  and  $\mathbf{G}_{\mathbf{V}_2}$  are connected. Let  $\alpha \in (0, 1) \setminus \{1/2\}$ . Fix the initial beliefs  $x^0$ . Then under belief dynamics (19), there are two random variables,  $B_1^\dagger(x^0), B_2^\dagger(x^0)$  both taking value in  $\{-A, A\}$ , such that*

$$\mathbb{P}\left(\lim_{k \rightarrow \infty} x_i(k) = B_1^\dagger, i \in \mathbf{V}_1; \lim_{k \rightarrow \infty} x_i(k) = B_2^\dagger, i \in \mathbf{V}_2 \mid \mathcal{E}_{\text{sep}}(\epsilon)\right) = 1 \quad (81)$$

for all  $\epsilon > 0$ , where by definition,  $\mathcal{E}_{\text{sep}}(\epsilon)$  is the  $\epsilon$ -separation event:

$$\mathcal{E}_{\text{sep}}(\epsilon) := \left\{ \limsup_{k \rightarrow \infty} \max_{i \in \mathbf{V}_1, j \in \mathbf{V}_2} |x_i(k) - x_j(k)| \geq \epsilon \right\}.$$

*Proof.* Suppose  $x_{i_1}(0) - x_{i_2}(0) \geq \epsilon > 0$  for  $i_1 \in \mathbf{V}_1$  and  $i_2 \in \mathbf{V}_2$ . By assumption,  $\mathbf{G}_{\mathbf{V}_1}$  and  $\mathbf{G}_{\mathbf{V}_2}$  are connected. Thus, from Lemma 5, there exist an integer  $Z_1 \geq 1$  and a constant  $\bar{p}$  (both depending on  $\epsilon, n, \alpha, a, b$ ) such that

$$\min_{i \in \mathbf{V}_1} x_i(Z_1) - \max_{i \in \mathbf{V}_2} x_i(Z_1) \geq \frac{\epsilon}{2} \quad (82)$$

happens with probability at least  $\bar{p}$ . Intuitively Eq. (82) characterizes the event where the beliefs in the two sets  $\mathbf{V}_1$  and  $\mathbf{V}_2$  are *completely separated*. Since all edges between the two sets are negative, conditioned on event (82), it is then straightforward to see that almost surely we have  $\lim_{k \rightarrow \infty} x_i(k) = A, i \in \mathbf{V}_1$  and  $\lim_{k \rightarrow \infty} x_i(k) = -A, i \in \mathbf{V}_2$ .

Given  $\mathcal{E}_{\text{sep}}(\epsilon)$ ,  $\{\exists i_1 \in \mathbf{V}_1, i_2 \in \mathbf{V}_2 \text{ s.t. } x_{i_1}(k) - x_{i_2}(k) \geq \epsilon \text{ for infinitely many } k\}$  is an almost sure event. Based on our previous discussion and by a simple stopping time argument, the Borel-Cantelli Lemma implies that the complete separation event happens almost surely given  $\mathcal{E}_{\text{sep}}(\epsilon)$ . This completes the proof.  $\square$

**Lemma 9** *Assume that the graph is strongly balanced under partition  $\mathbf{V} = \mathbf{V}_1 \cup \mathbf{V}_2$ , and that  $\mathbf{G}_{\mathbf{V}_1}$  and  $\mathbf{G}_{\mathbf{V}_2}$  are connected. Suppose  $\alpha \in (0, 1) \setminus \{1/2\}$ . Then under dynamics (19), there exists  $\beta$  sufficiently large such that  $\mathbb{P}(\mathcal{E}_{\text{sep}}(A/2)) = 1$  for almost all initial beliefs.*

*Proof.* Let us first focus on a fixed time instant  $k$ . Suppose  $x_i(k) - x_j(k) \geq A$  for some  $i, j \in \mathbf{V}$ . If  $i$  and  $j$  belong to different sets  $\mathbf{V}_1$  and  $\mathbf{V}_2$ , we already have  $\max_{i \in \mathbf{V}_1, j \in \mathbf{V}_2} |x_i(k) - x_j(k)| \geq A$ . Otherwise, say  $i, j \in \mathbf{V}_1$ . There must be another node  $l \in \mathbf{V}_2$ . We have  $\max_{i \in \mathbf{V}_1, j \in \mathbf{V}_2} |x_i(k) - x_j(k)| \geq A/2$  since either  $|x_i(k) - x_l(k)| \geq A/2$  or  $|x_j(k) - x_l(k)| \geq A/2$  must hold. Therefore, we conclude that

$$\mathfrak{X}(k) \geq A \implies \max_{i \in \mathbf{V}_1, j \in \mathbf{V}_2} |x_i(k) - x_j(k)| \geq A/2. \quad (83)$$

Then the desired conclusion follows directly from Lemma 7.  $\square$

Theorem 4 is a direct consequence of Lemmas 8 and 9.

## I. Proof of Theorem 5

The proof is similar to that of Theorem 4. We just provide the main arguments.

First by Lemma 7 we have  $\mathbb{P}(\limsup_{k \rightarrow \infty} \mathfrak{X}(k) = 2A) = 1$  for almost all initial values with sufficiently large  $\beta$ . Then as for (83), we have

$$\mathfrak{X}(k) \geq A \implies \max_{i \in \mathbf{V}_s, j \in \mathbf{V}_t, s \neq t \in \{1, \dots, m\}} |x_i(k) - x_j(k)| \geq \frac{A}{m}, \quad (84)$$

where  $m \geq 2$  comes from the definition of weak balance. Therefore, introducing

$$\mathcal{E}_{\text{sep}}^*(\epsilon) := \left\{ \limsup_{k \rightarrow \infty} \max_{i \in \mathbf{V}_s, j \in \mathbf{V}_t, s \neq t \in \{1, \dots, m\}} |x_i(k) - x_j(k)| \geq \epsilon \right\},$$

we can show that  $\mathbb{P}(\mathcal{E}_{\text{sep}}^*(A/m)) = 1$  for almost all initial beliefs, for sufficiently large  $\beta$ .

Next, suppose there exist a constant  $\eta > 0$  and two node sets  $\mathbf{V}_{i_1}$  and  $\mathbf{V}_{i_2}$  with  $i_1, i_2 \in \{1, \dots, m\}$  such that the complete separation event

$$\min_{i \in \mathbf{V}_{i_1}} x_i(k) - \max_{i \in \mathbf{V}_{i_2}} x_i(k) \geq \eta \quad (85)$$

happens. Recall that the underlying graph is complete. Then if  $(\beta + 1)\eta \geq 2A$ , we can always select  $Z_* := |\mathbf{V}_{i_1}| + |\mathbf{V}_{i_2}|$  negative edges between nodes in the sets  $\mathbf{V}_{i_1}$  and  $\mathbf{V}_{i_2}$ , so that after the corresponding updates:

$$x_i(k + Z_*) = A, i \in \mathbf{V}_{i_1}, \quad x_i(k + Z_*) = -A, i \in \mathbf{V}_{i_2}. \quad (86)$$

One can easily see that we can continue to build the (finite) sequence of edges for updates, such that nodes in  $\mathbf{V}_k$  will hold the same belief in  $\{-A, A\}$ , for all  $k = 1, \dots, m$ . After this sequence of updates, the beliefs held at the various nodes remain unchanged (two nodes with the same belief cannot influence each other, even in presence of a negative link; and two nodes with different beliefs are necessarily enemies). To summarize, conditioned on the complete separation event (85), we can select a sequence of node pairs under which belief clustering is reached, and this clustering state is an *absorbing* state.

Finally, the Borel-Cantelli Lemma and  $\mathbb{P}(\mathcal{E}_{\text{sep}}^*(A/m)) = 1$  guarantee that almost surely the complete separation event (85) happens an infinite number of times if  $\eta = A/2m$  in view of Lemma 5. The end of the proof is then done as in that of Theorem 4.

## J. Proof of Theorem 6

Again the result is obtained by combining Lemmas 5 and 7 with Borel-Cantelli lemma.

## Acknowledgement

This work has been supported in part by the Knut and Alice Wallenberg Foundation, the Swedish Research Council, and KTH SRA TNG. The authors gratefully thank Dr. Shuangshuang Fu for her generous help in preparing the numerical examples.

## References

- Acemoglu, D., Ozdaglar A., and ParandehGheibi A. (2010) Spread of (Mis)information in social networks. *Games and Economic Behavior* **70**: 194-227.
- Acemoglu, D., Dahleh, M.A., Lobel, I., and Ozdaglar, A. (2011) Bayesian learning in social networks. *Review of Economic Studies* **78**: 1201-1236.
- Acemoglu, D., Como, G., Fagnani F., and Ozdaglar A. (2013) Opinion fluctuations and disagreement in social networks. *Mathematics of Operations Research*, **38**: 1-27.
- Acemoglu, D. and Ozdaglar, A. (2011) Opinion dynamics and learning in social networks. *Dynamic Games and Applications*, **1**: 3-49.
- Altafini, C. (2012) Dynamics of opinion forming in structurally balanced social networks. *PLoS ONE*, **7(6)**: e38135.
- Altafini, C. (2013) Consensus problems on networks with antagonistic interactions. *IEEE Trans. Automatic Control*, **58**:935-946.
- Banerjee, A. (1992) A simple model of herd behavior. *The Quarterly Journal of Economics*, **107**: 797-817.
- Bikhchandani, S., Hirshleifer, D., and Welch, I. (1992) A theory of fads, fashion, custom, and cultural change as information cascades. *The Journal of Political Economy*, **100**: 992-1026.
- Boyd, S., Ghosh, A., Prabhakar B., and Shah, D. (2006) Randomized gossip algorithms. *IEEE Trans. on Inf. Theory*, **52**: 2508-2530.
- Blondel, V. D., Hendrickx, J. M., and Tsitsiklis, J. N. (2009) On Krause's multi-agent consensus model with state-dependent connectivity. *IEEE Trans. on Autom. Control*, **54**: 2586-2597.

- Blondel, V. D., Hendrickx, J. M., and Tsitsiklis, J. N. (2010) Continuous-time average-preserving opinion dynamics with opinion-dependent communications. *SIAM J. Control and Optimization*, **48(8)**: 5214-5240.
- Castellano, C., Fortunato, S., and Loreto, V. (2009) Statistical physics of social dynamics. *Rev. Modern Phys.*, **81(2)**:591-646.
- Cartwright, D., and Harary, F. (1956) Structural balance: A generalization of Heider's theory. *Psychol Rev.*, **63**:277-293.
- Clark, R. D., and Maass, A. (1990) The effects of majority size on minority influence. *European Journal of Social Psychology* **20(2)**: 99-117.
- Dandekara, P., Goel, A., and Lee D. T. (2013) Biased assimilation, homophily, and the dynamics of polarization. *Proceedings of the National Academy of Sciences*, **110(15)**: 5791-5796.
- Davis, J. A. (1963) Structural balance, mechanical solidarity, and interpersonal relations. *American Journal of Sociology*, **68**:444-462.
- Davis, J. A. (1967) Clustering and structural balance in graphs. *Human Relations*, **20**: 181-187.
- DeGroot, M. H. (1974) Reaching a consensus. *Journal of the American Statistical Association*, **69**: 118-121.
- DeMarzo, Vayanos, and Zwiebel (2003) Persuasion bias, social influence, and unidimensional opinions. *Quarterly Journal of Economics*, **118**:909-968.
- Ding, X. and Jiang, T. (2010) Spectral distributions of adjacency and Laplacian matrices of random graphs. *The Annals of Applied Probability*, **20**: 2086-2117.
- Durrett, R. (2010) *Probability Theory: Theory and Examples*. 4th ed. Cambridge University Press: New York.
- Easley, D. and Kleinberg, J. (2010) *Networks, Crowds, and Markets: Reasoning About a Highly Connected World*. Cambridge University Press.
- Erdős, P. and Rényi, A. (1960) On the evolution of random graphs. *Publications of the Mathematical Institute of the Hungarian Academy of Sciences*, 17-61.
- Facchetti, G., Iacono, G., and Altafini, C. (2011) Computing global structural balance in large-scale signed social networks. *Proceedings of the National Academy of Sciences*, **108(52)**:20953-8.
- Fagnani, F. and Zampieri, S. (2008) Randomized consensus algorithms over large scale networks. *IEEE J. Selected Areas in Communications*, **26**: 634-649.
- Feller, W. (1968) *An Introduction to Probability Theory and Its Applications*. 3rd ed. New York: Wiley.
- Galam, S. (1996) Fragmentation versus stability in bimodal coalitions. *Physica A*, **230**: 174-188.
- Golub, B. and Jackson, M. O. (2010) Naive learning in social networks and the wisdom of crowds. *American Economic Journal: Microeconomics*, **2(1)**: 112-149.
- Theodorakopoulos, G. and Baras, J. S. (2008) Game theoretic modeling of malicious users in collaborative networks. *IEEE J. Selected Areas in Communications*, **26**: 1317-1327.
- Heider, F. (1946) Attitudes and cognitive organization. *J Psychol*, **21**:107-112.



- Harary, F. (1953) On the notion of balance of a signed graph. *Michigan Math. Journal*, **2(2)**:143-146.
- Horn, R. A. and Johnson, C. R. (1985) *Matrix Analysis*. Cambridge University Press.
- Jackson, M. O. (2008) *Social and Economic Networks*. 1st ed. Princeton University Press.
- Jadbabaie, A., Molavi, P., Sandroni, A., and Tahbaz-Salehi, A. (2012) Non-Bayesian social learning. *Games and Economic Behavior*, **76**: 210-225.
- Krause, U. (1997) Soziale dynamiken mit vielen interagierenden Akteuren. Eine problemskizze, in *Proc. Modellierung Simul. von Dynamiken mit vielen interagierenden Akteuren*: 37-51.
- Li, L., Scaglione, A., Swami, A., and Zhao Q. (2013) Consensus, polarization and clustering of opinions in social networks. *IEEE J. Selected Areas in Communications*, **31**: 1072-1083.
- Macy, M. W., Kitts, J. A., Flache, A., and Benard, S. (2003) Polarization in dynamic networks: A Hopfield model of emergent structure. *Dynamic Social Network Modeling and Analysis*, 162-173.
- Mäs, M., Flache, A., and Helbing, D. (2010) Individualization as driving force of clustering phenomena in humans. *PLOS Computational Biology*, **6(10)**: e1000959.
- Marvel, S. A., Kleinberg, J., Kleinberg, R. D., and Strogatz, S. H. (2011) Continuous-time model of structural balance. *Proceedings of the National Academy of Sciences*, **108(5)**: 1751-1752.
- Matei, I., Baras, J. S., and Somarakis C. (2013) Convergence results for the linear consensus problem under Markovian random graphs. *SIAM J. on Control and Optimization*, **51(2)**: 1574-1591.
- Nemeth, C. J. (1986) Differential contributions of majority and minority influence. *Psychological Review*, **93(1)**: 23.
- Shi, G., Li, B., Johansson, M., and Johansson, K. H. (2014) When do gossip algorithms converge in finite time? in *The 21st International Symposium on Mathematical Theory of Networks and Systems (MTNS)*, Groningen, The Netherlands.
- Shi, G., Johansson, M., and Johansson, K. H. (2013) How agreement and disagreement evolve over random dynamic networks. *IEEE J. Selected Areas in Communications*, **31**: 1061-1071.
- Shi, G., Proutiere, A., Johansson, M., Baras, J. S., and Johansson, K. H. (2015) Emergent behaviors over signed random dynamical networks: state-flipping model. *IEEE Trans. on Control of Network Systems*, **2**: 142-153.
- Tahbaz-Salehi, A. and Jadbabaie, A. (2008) A necessary and sufficient condition for consensus over random networks. *IEEE Trans. on Autom. Control*, **53**: 791-795.
- Touri, B. and Nedić, A. (2011) On ergodicity, infinite flow and consensus in random models. *IEEE Trans. on Automatic Control*, **56**: 1593-1605.
- Tsitsiklis, J. N. (1984) *Problems in decentralized decision making and computation*. Ph.D. thesis, Dept. of Electrical Engineering and Computer Science, Massachusetts Institute of Technology, Boston, MA.
- Wasserman, S. and Faust, K. (1994) *Social Network Analysis: Methods and Applications, Structural Analysis in the Social Sciences*. Cambridge Univ Press, New York.
- Xiao, L., and Boyd, S. (2004) Fast linear iterations for distributed averaging. *Systems & Control Letters*, **53**: 65-78.

Yildiz, E., Acemoglu, D., Ozdaglar, A., Saberi, A., and Scaglione, A. (2013) Binary opinion dynamics with stubborn agents. *ACM Trans. Econ. Comp.*, **1(4)**: 19.

GUODONG SHI

Research School of Engineering, College of Engineering and Computer Science,

The Australian National University, Canberra ACT 0200, Australia

Email: `guodong.shi@anu.edu.au`

ALEXANDRE PROUTIERE, MIKAEL JOHANSSON, AND KARL H. JOHANSSON

ACCESS Linnaeus Centre, School of Electrical Engineering, KTH Royal Institute of Technology,

Stockholm 100 44, Sweden

Email: `alepro@kth.se`, `mikaelj@kth.se`, `kallej@kth.se`

JOHN S. BARAS

Institute for Systems Research, University of Maryland,

College Park, MD 20742, USA

Email: `baras@umd.edu`