

ANDRÉ-QUILLEN HOMOLOGY AND COMPLETE INTERSECTION FLAT DIMENSION

TIRDAD SHARIF

ABSTRACT. We study the relation between the vanishing of André-Quillen homology and complete intersection flat dimension and we extend some of the existing results in the literature.

1. INTRODUCTION

Convention. In this paper, rings are commutative, noetherian, and local with identity. In the entire paper, (R, \mathfrak{m}, k) , (S, \mathfrak{n}, ℓ) , and (T, \mathfrak{r}, v) are local rings where S and T are complete. When it is convenient, $\varphi: R \rightarrow S$ and $\sigma: S \rightarrow T$ will denote local ring homomorphisms.

Over forty years ago, André [1, 2] and Quillen [22, 23] introduced a homology theory for commutative algebras that is known as André-Quillen (or cotangent) homology. The n -th André-Quillen homology of the R -algebra S with coefficients in an S -module N , denoted $D_n(S | R, N)$, is the n -th homology module of $\mathcal{L}_\varphi \otimes_S N$, where \mathcal{L}_φ is the cotangent complex of φ , uniquely defined in the derived category of the category of S -modules. The vanishing of André-Quillen homology characterizes important classes of rings and ring homomorphisms; see [1, 3, 6, 9, 10, 12, 22, 23].

To study the relation between the vanishing of André-Quillen homology and the structure of a local ring homomorphism φ , Avramov and Iyengar [10] introduced the notion of André-Quillen dimension of the R -algebra S , denoted $\text{AQ-dim}_R S$, that is

$$\text{AQ-dim}_R S := \sup\{n \in \mathbb{N} \mid D_n(S | R, -) \neq 0\};$$

in particular, $\text{AQ-dim}_R S = -\infty$ if and only if $D_n(S | R, -) = 0$ for all integers n .

In this paper, we investigate the relation between the vanishing of André-Quillen homology and complete intersection flat dimension. As the first application of this study, we prove the following theorem which is one of our main results in this paper; see 3.2 for the proof.

Theorem A. *Assume that σ is surjective, $\sigma\varphi$ is complete intersection at \mathfrak{r} , and $\text{AQ-dim}_S T < \infty$. Then $\text{CI-dim}_S T < \infty$.*

It is worth mentioning that the surjectivity condition in this theorem can be removed if one can give an affirmative answer to a question posed by Foxby; see Remark 3.6.

2010 *Mathematics Subject Classification.* Primary: 13D03, 13D05; Secondary: 13B10, 14B25.

Key words and phrases. Algebra retract, André-Quillen dimension, André-Quillen homology, Betti numbers, complete intersection dimension, complexity, semidualizing module, trivial extension.

Tirdad Sharif was supported by a grant from IPM (No. 83130311).

The second main result of this paper is the following theorem, which extends a result of Soto [27, Proposition 12] and of Avramov, Henriques, and Şega [9, (2.5)(3)]; see 3.5 for the proof. Here, $H_1(K.(\varphi))$, denotes the first Koszul homology of φ ; see Definition 2.8

Theorem B. *Assume that φ is essentially of finite type and $\text{CI-fd}_R S < \infty$. Then $\text{AQ-dim}_R S \leq 2$ if and only if $H_1(K.(\varphi))$ is a free S -module.*

2. SOME BACKGROUND MATERIAL

This section lists important foundational material for use in this paper. We use the notations $\mu_R(M)$ and $\text{edim}(R)$ for the minimal number of generators of a finitely generated R -module M and the embedding dimension of the ring R , respectively.

The objects defined in the next definition are from the work of Avramov, Foxby, and B. Herzog [7].

2.1. A *regular factorization* of φ is a diagram of local ring homomorphisms $R \xrightarrow{\hat{\varphi}} R' \xrightarrow{\varphi'} S$ such that $\varphi = \varphi' \hat{\varphi}$, the map $\hat{\varphi}$ is flat, the closed fibre $R'/\mathfrak{m}R'$ is regular, and φ' is surjective. A *Cohen factorization* of φ is a regular factorization $R \xrightarrow{\hat{\varphi}} R' \xrightarrow{\varphi'} S$ of φ such that R' is complete. Since we assumed that S is complete, existence of a Cohen factorization of φ is guaranteed by [7, Theorem (1.1)].

2.2. The local ring homomorphism φ is called *complete intersection at \mathfrak{n}* if in a given Cohen factorization $R \rightarrow R' \xrightarrow{\varphi'} S$ of φ the ideal $\text{Ker } \varphi'$ is generated by an R' -regular sequence. The complete intersection property for φ is independent of the choice of Cohen factorization by [6, Remark (3.3)]. Also φ is called *locally complete intersection* if for each $\mathfrak{q} \in \text{Spec}(S)$, the local homomorphism $\varphi_{\mathfrak{q}} : R_{\mathfrak{p}} \rightarrow S_{\mathfrak{q}}$, in which $\mathfrak{p} = \mathfrak{q} \cap R$, is complete intersection at $\mathfrak{q}S_{\mathfrak{q}}$.

We proceed with introducing the complete intersection flat dimension as defined by Sather-Wagstaff [25].

2.3. Fix an arbitrary R -module M . Define

$$\text{CI-fd}_R M := \inf \left\{ \text{fd}_Q(R' \otimes_R M) - \text{pd}_Q(R') \mid \begin{array}{l} R \rightarrow R' \leftarrow Q \text{ is a} \\ \text{quasi-deformation} \end{array} \right\}.$$

Recall that a diagram $R \rightarrow R' \leftarrow Q$ of local ring homomorphisms is a *quasi-deformation of R* if $R \rightarrow R'$ is flat and $R' \leftarrow Q$ is surjective with kernel generated by a Q -regular sequence. It is clear that if M is finitely generated then $\text{CI-dim}_R M = \text{CI-fd}_R M$.

2.4. Here, we recall some notions from [10, 20].

Let $\overline{\varphi}: k \rightarrow \ell$ be the induced map between the residue fields. Then φ is called *almost small* if the kernel of the homomorphism

$$\text{Tor}_*^{\varphi}(\overline{\varphi}, \ell): \text{Tor}_*^R(k, \ell) \rightarrow \text{Tor}_*^S(\ell, \ell)$$

of graded algebras is generated by elements of degree 1. Assuming $k = \ell$, the homomorphism φ is called *large* if $\text{Tor}_*^{\varphi}(\overline{\varphi}, k)$ is surjective.

The next definition originates with work of Gulliksen and Levin [15]; see [5, §6] for details. It involves a different type of algebra extensions introduced by Tate [28]; see [5, 15] for details.

2.5. Let A be a DG R -algebra, and assume that φ is surjective. Let $A\langle Y \rangle$ denote a DG R -algebra obtained from A by adjunctions of sets of exterior variables of positive odd degrees and of divided power variables of positive even degrees. Then a factorization $R \rightarrow R\langle Y \rangle \xrightarrow{\cong} S$ of φ is called an *acyclic closure* of φ if $\partial(Y_1)$ minimally generates $\text{Ker } \varphi$, and $\{\text{cls}(\partial(y)) \mid y \in Y_{n+1}\}$ is a minimal generating set of $H_n(R\langle Y_{\leq n} \rangle)$ for all $n \geq 1$.

Fact 2.6. If $J = (x) = (y)$ is an ideal of R , such that x and y are two minimal set of generators for J , then the Koszul complexes associated to (x) and (y) are isomorphic, see [14], and hence homology isomorphism. Thus we define the Koszul complex, $K^{.R}(J)$, to be the Koszul complex associated to an arbitrary minimal set of generators for ideal J .

Fact 2.7. Let $R \xrightarrow{\varphi_j} R_j \xrightarrow{\varphi_j} S$, with $J_j = \text{Ker } \varphi_j$, for $j = 1, 2$, be two arbitrary Cohen factorization of φ . From [7, (1.2)] it follows that they have a common deformation $R \rightarrow R' \xrightarrow{\varphi'} S$, with $J' = \text{Ker } \varphi'$ such that $R_j = R'/I'_j$ and $J_j = J'/I'_j$, for $j = 1, 2$, where I'_j is an ideal generated by an R' -regular sequence that extends to a minimal set of generators for J' . Let \underline{x}_j be a minimal set of generators for I'_j , which can be extended to a minimal set of generators $\underline{x}_j, \underline{y}_j$ for J' . Then the residues \underline{y}'_j of \underline{y}_j in R_j form a minimal set of generators for J_j , by Nakayama's lemma. Now from the above fact we have

$$K^{.R'}(J') \cong K^{.R'}(\underline{x}_j) \otimes_{R'} K^{.R'}(\underline{y}_j) \simeq R_j \otimes_{R'} K^{.R'}(\underline{y}_j) \cong K^{.R_j}(\underline{y}'_j) \cong K^{.R_j}(J_j).$$

Thus $K^{.R'}(J') \simeq K^{.R_j}(J_j)$, as R' -complexes, for $j = 1, 2$. Since $J' \text{H}(K^{.R'}(J')) = 0$, we get $\text{H}(K^{.R'}(J')) \cong \text{H}(K^{.R_j}(J_j))$, as S -modules, for $j = 1, 2$.

By the above fact, we can give the following definition.

Definition 2.8. Let $R \rightarrow R' \xrightarrow{\varphi'} S$ be a Cohen factorizations of φ , with $J' = \text{Ker } \varphi'$. We define the *n th Koszul homology* of φ , denoted by $\text{H}_n(K.(\varphi))$, to be $\text{H}_n(K^{.R'}(J'))$.

Indeed, we can consider $\text{H}(K.(\varphi))$, as an invariant of the local homomorphism φ .

Lemma 2.9. *Let $R \rightarrow R' \xrightarrow{\varphi'} S$ be a Cohen factorization of φ . If E is the Koszul complex associated to an arbitrary set of generators \underline{v} for $J' := \text{Ker } \varphi'$. Then $\text{H}_1(E)$ is a free S -module if and only if $\text{H}_1(K.(\varphi))$ is a free S -module.*

Proof. Let $\underline{v} = x_1, \dots, x_p, x_{p+1}, \dots, x_n$, in which $\underline{w} = x_1, \dots, x_p$ is a minimal set of generators for J' . Then from [14, (1.6.21)] and Fact 2.6, we get

$$\text{H}_1(E) \cong \text{H}_1(K^{.R'}(\underline{w})) \oplus S^{\oplus(n-p)} \cong \text{H}_1(K^{.R'}(J')) \oplus S^{\oplus(n-p)}.$$

Now from the above fact, assertion holds. \square

2.10. Let M be a finitely generated R -module. The *complexity of M over R* , denoted by $\text{cx}_R M$, is the number

$$\text{cx}_R M := \inf\{d \in \mathbb{N} \mid \text{there exists } \gamma \in \mathbb{R} \text{ such that } \beta_n^R(M) \leq \gamma n^{d-1} \text{ for all } n \gg 0\}$$

where $\beta_n^R(M) := \text{rank}_k(\text{Tor}_n^R(M, k))$ is the n -th *Betti number* of M .

3. MAIN RESULTS

This section contains the proofs of Theorems A and B. We begin by a construction used by Iyengar and Sather-Wagstaff [19], essentially given by Avramov, Foxby, and B. Herzog [7]. We use this construction in the proof of Theorem A and in Remark 3.6.

Construction 3.1. Let $R \xrightarrow{\dot{\varphi}} R' \xrightarrow{\varphi'} S$ and $R' \xrightarrow{\dot{\rho}} R'' \xrightarrow{\rho'} T$ be Cohen factorizations of φ and $\sigma\varphi'$, respectively. Then ρ' factors through the tensor product $S' = R'' \otimes_{R'} S$ and gives the following commutative diagram of local ring homomorphisms

$$\begin{array}{ccccc}
 & & R'' & & \\
 & \overset{\dot{\rho}\dot{\varphi}}{\curvearrowright} & \nearrow \dot{\rho} & \searrow \varphi'' & \overset{\rho'=\sigma'\varphi''}{\curvearrowright} \\
 & R' & & S' & \\
 \nearrow \dot{\varphi} & & \searrow \varphi' & \nearrow \dot{\sigma} & \searrow \sigma' \\
 R & \xrightarrow{\varphi} & S & \xrightarrow{\sigma} & T
 \end{array}$$

in which $\dot{\sigma}$ and φ'' are the natural maps to the tensor product and the diagrams $S \rightarrow S' \rightarrow T$, $R \rightarrow R'' \rightarrow T$, and $R' \rightarrow R'' \rightarrow S'$ are Cohen factorizations.

Here is the proof of Theorem A.

3.2 (Proof of Theorem A). We use the notation from Construction 3.1. Since σ is surjective, we get the following diagram

$$R' \xrightarrow{\varphi'} S \xrightarrow{\sigma} T \quad (3.2.1)$$

of surjective local ring homomorphisms such that $R \xrightarrow{\dot{\varphi}} R' \xrightarrow{\sigma\varphi'} T$ is a Cohen factorization. Since $\sigma\varphi$ is complete intersection at \mathfrak{t} , from [6, (1.7), (1.8)] we get that $\sigma\varphi'$ is also complete intersection at \mathfrak{t} . Hence, by [10, Corollary (4.9)] we conclude that φ' is an almost small ring homomorphism. Since $D_n(T | S, v) = 0$ for all $n \gg 0$, from the Jacobi-Zariski exact sequence arising from diagram (3.2.1), we find that $D_n(S | R', v) = 0$ for all $n \gg 0$. Since φ' is almost small, it follows from [10, (6.4)] that φ' is complete intersection at \mathfrak{t} . Thus, $\text{Ker } \varphi'$ is generated by an R' -regular sequence, and the diagram $S \xrightarrow{\text{id}} S \xleftarrow{\varphi'} R'$ of local ring homomorphisms, is a quasi-deformation. Hence, we conclude that $\text{pd}_{R'}(T \otimes_S S) = \text{pd}_{R'} T < \infty$. Therefore, $\text{CI-dim}_S T < \infty$. \square

The following lemma is an extension of [24, Lemma 1(2)]. This lemma is well-known; see [4]. For convenience of the reader we give a proof based on the Cohen factorization.

Lemma 3.3. *If $\text{fd}_R S < \infty$, then the homomorphism $D_2(S | R, \ell) \xrightarrow{\theta} D_2(\ell | R, \ell)$ is trivial.*

Proof. Let $R \xrightarrow{\varphi} R' \xrightarrow{\varphi'} S$ be a Cohen factorization of φ , and consider the following commutative diagram of local ring homomorphisms

$$\begin{array}{ccccc} R & \xrightarrow{\varphi} & R' & \xrightarrow{\varphi'} & S \\ \varphi' \downarrow & & \downarrow \varphi & & \downarrow \pi \\ S & \xrightarrow{\text{id}} & S & \xrightarrow{\pi} & \ell \\ \pi \downarrow & & \downarrow \pi & & \\ \ell & \xrightarrow{\text{id}} & \ell & & \end{array}$$

where π is the natural surjection. This diagram induces a commutative diagram

$$\begin{array}{ccc} D_2(S \mid R, \ell) & \xrightarrow{\theta} & D_2(\ell \mid R, \ell) \\ \downarrow & & \downarrow \eta \\ D_2(S \mid R', \ell) & \xrightarrow{\alpha} & D_2(\ell \mid R', \ell) \end{array} \quad (3.3.1)$$

of ℓ -vector spaces.

Setting $\overline{R'} := R'/\mathfrak{m}R'$, from the Jacobi-Zariski exact sequence of $k \rightarrow \overline{R'} \rightarrow \ell$ we get the following exact sequence:

$$D_{n+1}(\ell \mid \overline{R'}, \ell) \rightarrow D_n(\overline{R'} \mid k, \ell) \rightarrow D_n(\ell \mid k, \ell) \quad (3.3.2)$$

From [6, (1.6)(2)-(3)] we get $D_n(\ell \mid k, \ell) = 0$ for all $n \geq 2$, and $D_{n+1}(\ell \mid \overline{R'}, \ell) = 0$ for all $n \geq 2$. Hence, by (3.3.2) we conclude that $D_n(\overline{R'} \mid k, \ell) = 0$ for all $n \geq 2$. On the other hand, by flat base change we have $D_n(\overline{R'} \mid k, \ell) \cong D_n(R' \mid R, \ell)$ for all integers n ; see [2, 4.54]. Therefore, $D_n(R' \mid R, \ell) = 0$ for all $n \geq 2$. Now, the Jacobi-Zariski exact sequence arising from $R \rightarrow R' \rightarrow \ell$ implies that η (from diagram (3.3.1)) is injective.

From [7, (3.2) Lemma] and our assumption, we have $\text{pd}_{R'} S < \infty$. Hence by [24, Lemma 1(2)] we get that α (from diagram (3.3.1)) is trivial. Therefore, commutativity of diagram (3.3.1) and the fact that η is injective imply that θ is trivial. \square

Lemma 3.4. *If $\text{AQ-dim}_R S \leq 2$, then $H_1(K(\varphi))$ is a free S -module.*

Proof. Let $R \xrightarrow{\varphi} R' \xrightarrow{\varphi'} S$ be a Cohen factorization of φ , and let $J' := \text{Ker } \varphi'$. By assumption we get $D_n(S \mid R, \ell) = 0$ for all $n \geq 3$. Hence, $D_n(S \mid R', \ell) = 0$ for all $n \geq 3$ by [6, Lemma (1.7)]. This implies that $D_n(S \mid R', -) = 0$ for all $n \geq 3$. Now [13, Corollary 3] implies that $H_1(E')$, in which E' is the Koszul complex associated to an arbitrary set of generators for J' , is a free S -module. The assertion now follows from Lemma 2.9. \square

Here is the proof of Theorem B.

3.5 (Proof of Theorem B). Since φ is essentially of finite type, it admits a regular factorization $R \rightarrow R[X]_{\mathfrak{M}} \rightarrow S$, in which $R[X] := R[x_1, x_2, \dots, x_n]$, for some $n \geq 1$, and \mathfrak{M} is a prime ideal of $R[X]$ lying over \mathfrak{m} . Since $\text{CI-fd}_R S < \infty$, there is a quasi-deformation $R \rightarrow R'' \xleftarrow{\eta} Q$ of R such that $\text{fd}_Q(S \otimes_R R'') < \infty$. Set $R' := R[X]_{\mathfrak{M}}$, $P := R' \otimes_R R''$, $U := S \otimes_R R''$, and consider the diagram $Q \xrightarrow{\alpha} P \xrightarrow{\beta} U$ of natural

ring homomorphisms, in which β is surjective. Let $\mathfrak{u} \in \text{Spec}(U)$, and set $\mathfrak{p} := \beta^{-1}(\mathfrak{u})$ and $\mathfrak{q} := \alpha^{-1}(\mathfrak{p})$. Then we have the diagram

$$Q_{\mathfrak{q}} \xrightarrow{\alpha_{\mathfrak{q}}} P_{\mathfrak{p}} \xrightarrow{\beta_{\mathfrak{p}}} U_{\mathfrak{u}} \quad (3.5.1)$$

of local ring homomorphisms that gives us the commutative diagram

$$\begin{array}{ccccc} Q_{\mathfrak{q}} & \xrightarrow{\alpha_{\mathfrak{q}}} & P_{\mathfrak{p}} & \xrightarrow{\beta_{\mathfrak{p}}} & U_{\mathfrak{u}} \\ \beta_{\mathfrak{p}} \alpha_{\mathfrak{q}} \downarrow & & \downarrow \beta_{\mathfrak{p}} & & \downarrow \\ U_{\mathfrak{u}} & \xrightarrow{\text{id}} & U_{\mathfrak{u}} & \longrightarrow & k(\mathfrak{u}) \\ \downarrow & & \downarrow & & \\ k(\mathfrak{u}) & \xrightarrow{\text{id}} & k(\mathfrak{u}) & & \end{array}$$

of local ring homomorphisms. This diagram induces the following commutative diagram of $k(\mathfrak{u})$ -vector spaces

$$\begin{array}{ccccc} D_3(k(\mathfrak{u}) | U_{\mathfrak{u}}, k(\mathfrak{u})) & \xrightarrow{f} & D_2(U_{\mathfrak{u}} | P_{\mathfrak{p}}, k(\mathfrak{u})) & \longrightarrow & D_2(k(\mathfrak{u}) | P_{\mathfrak{p}}, k(\mathfrak{u})) \\ \parallel & & \uparrow h & & \\ D_3(k(\mathfrak{u}) | U_{\mathfrak{u}}, k(\mathfrak{u})) & \xrightarrow{g} & D_2(U_{\mathfrak{u}} | Q_{\mathfrak{q}}, k(\mathfrak{u})) & \xrightarrow{\theta} & D_2(k(\mathfrak{u}) | Q_{\mathfrak{q}}, k(\mathfrak{u})) \\ & & \uparrow \lambda & & \\ & & D_2(P_{\mathfrak{p}} | Q_{\mathfrak{q}}, k(\mathfrak{u})) & & \end{array} \quad (3.5.2)$$

Since $\text{fd}_{Q_{\mathfrak{q}}} U_{\mathfrak{u}} < \infty$, it follows from Lemma 3.3 that θ is trivial and hence g is surjective, and since $H_1(K(\varphi))$ is a free S -module, we can see that $H_1(E)$ is a free $U_{\mathfrak{u}}$ -module, where E is the Koszul complex associated to the set of generators for the ideal $\text{Ker } \beta_{\mathfrak{p}}$ of $P_{\mathfrak{p}}$ arising from a set of generators for $\text{Ker } \beta$. Now from [14, (1.6.21)] it follows that $H_1(K^{P_{\mathfrak{p}}}(\text{Ker}(\beta_{\mathfrak{p}})))$ is a free $U_{\mathfrak{u}}$ -module. Thus, from [?, Theorem 1] we conclude that f is trivial. Since $D_n(R[X] | R, -) = 0$ for all $n \geq 1$, we get $D_n(R' | R, -) = 0$ for all $n \geq 1$. The faithful flatness of R'' over R also yields $D_n(P | R'', -) = 0$ for all $n \geq 1$. Hence, setting $\mathfrak{p}'' := \mathfrak{p} \cap R''$, we get $D_n(P_{\mathfrak{p}} | R''_{\mathfrak{p}''}, k(\mathfrak{p})) = 0$ for all $n \geq 1$.

Consider the diagram $Q_{\mathfrak{q}} \xrightarrow{\eta_{\mathfrak{q}}} R''_{\mathfrak{p}''} \rightarrow P_{\mathfrak{p}}$ of local ring homomorphisms. Since $\text{Ker } \eta_{\mathfrak{q}}$ is generated by a $Q_{\mathfrak{q}}$ -regular sequence, from the Jacobi-Zariski exact sequence we get $D_n(P_{\mathfrak{p}} | Q_{\mathfrak{q}}, k(\mathfrak{p})) = 0$ for all $n \geq 2$. Thus, h in diagram (3.5.2) is injective. Commutativity of diagram (3.5.2) implies that $D_2(U_{\mathfrak{u}} | Q_{\mathfrak{q}}, k(\mathfrak{u})) = 0$. Hence, by [6, Proposition (1.8)] we get $D_n(U_{\mathfrak{u}} | Q_{\mathfrak{q}}, k(\mathfrak{u})) = 0$ for all $n \geq 2$. Now, from the Jacobi-Zariski exact sequence of diagram (3.5.1) we conclude that $D_n(U_{\mathfrak{u}} | P_{\mathfrak{p}}, k(\mathfrak{u})) = 0$ for all $n \geq 3$. Since \mathfrak{u} was an arbitrary prime ideal of U , we conclude that $D_n(U | P, -) = 0$ for all $n \geq 3$; see [2]. Now, faithful flatness of R'' over R implies that $D_n(S | R', -) = 0$ for all $n \geq 3$. Thus, from [6, Lemma (1.7)] we have $D_n(S | R, \ell) = 0$ for all $n \geq 3$. Since φ is essentially of finite type, from [18, Lemma 8.7] we conclude that $\text{AQ-dim}_R S \leq 2$.

The converse follows immediately from Lemma 3.4. \square

Foxby asked the following question in his personal communications.

Foxby's Question. Let $R \xrightarrow{\varphi} R' \xrightarrow{\varphi'} S$ be a Cohen factorization of φ , and let M be a finitely generated S -module. Do the following inequalities hold?

$$\text{CI-fd}_R M \leq \text{CI-dim}_{R'} M \leq \text{CI-fd}_R N + \text{edim}(R'/\mathfrak{m}R')$$

In the next remark, we show that an affirmative answer to this question helps us to generalize Theorem A. In fact, validity of the left inequality of Foxby's Question enables us to remove the hypothesis “ σ is surjective” in Theorem A; see (3.6.1). Also, we show that an affirmative answer to the right inequality in Foxby's Question provides a much shorter proof for Theorem B, without the assumption of finiteness of φ ; see (3.6.2).

Remark 3.6. (3.6.1) Using the notation from Construction 3.1, there exists a diagram $R'' \xrightarrow{\varphi''} S' \xrightarrow{\sigma'} T$ of surjective local ring homomorphisms. Since $\sigma\varphi$ is complete intersection at \mathfrak{r} , from [6, (1.7), (1.8)] we get that $\sigma'\varphi''$ is also complete intersection at \mathfrak{r} . Now a similar argument as in the proof of Theorem A, yields $\text{CI-dim}_{S'} T < \infty$. Then the left inequality of Foxby's Question implies that $\text{CI-fd}_S T < \infty$, as desired.

(3.6.2) Let $R \xrightarrow{\varphi} R' \xrightarrow{\varphi'} S$ be a Cohen factorization of φ . From the right inequality in Foxby's Question, we get $\text{CI-dim}_{R'} S < \infty$. Since $H_1(K.(\varphi))$ is a free S -module, from Lemma 2.9 and [27, Proposition 12] we conclude that $D_n(S | R', \ell) = 0$ for all $n \geq 3$. Now, the assertion follows from [6, Lemma (1.7)] and [18, Lemma 8.7].

The next corollary of Soto [27, Proposition 12] and of Avramov, Henriques, and Sega [9, (2.5)(3)] follows from Theorem B and Lemma 2.9.

Corollary 3.7. *Assume that φ is surjective, and let $H_1(E)$ be the first homology of the Koszul complex E associated to an arbitrary set of generators for $J := \text{Ker } \varphi$. If $\text{CI-dim}_R S < \infty$ and $H_1(E)$ is a free S -module, then $\text{AQ-dim}_R S \leq 2$.*

Using Theorem A, Lemma 3.4 and Corollary 3.7 we have the next result due to Soto [27, Proposition 23 (i) \iff (ii)]. Recall that S is called an *algebra retract* of R if there exists a local ring homomorphism $\psi: S \rightarrow R$ such that $\varphi\psi = \text{id}_S$.

Corollary 3.8. *Let S be an algebra retract of R and let $H_1(E)$ be the first homology of the Koszul complex E associated to an arbitrary set of generators for $J := \text{Ker } \varphi$. Then $\text{AQ-dim}_R S \leq 2$ if and only if $\text{CI-dim}_R S < \infty$ and $H_1(E)$ is a free S -module.*

Fact 3.9. Let S be an algebra retract of R and $\text{cx}_R S < \infty$. If $\text{char}(k) = 0$ then S contains the rational numbers. On the other hand $\varphi: R \rightarrow S$ is a large homomorphism. Let $R\langle U \rangle$ be the acyclic closure of φ , which is a minimal free resolution for S as an R -module by [11, Corollary (2.7)]. Since $\text{cx}_R S < \infty$, it follows from [27, Lemma 16] that $D_{2n}(S | R, -) = 0$ for all $n \gg 0$. Now from [10, Theorem (7.5)(iv) \implies (ii)] we have $\text{AQ-dim}_R S \leq 2$. Thus by Corollary 3.8, $\text{CI-dim}_R S < \infty$.

The above fact answers Soto's Problem 24 in [27] when $\text{char}(k) = 0$. As an application of this fact we also have the following result.

Corollary 3.10. *Let $A := S \otimes_R S$, and assume that φ is a local flat homomorphism essentially of finite type. Let $\omega: A \rightarrow S$ be the natural surjection and $\mathfrak{q} = \omega^{-1}(\mathfrak{n})$. If $\text{char}(\ell) = 0$ and $\text{cx}_{A_{\mathfrak{q}}} S < \infty$, then φ is locally complete intersection.*

Proof. Consider the algebra retract $S \rightarrow A_q \rightarrow S$. Fact 3.9 implies that $\text{AQ-dim}_{A_q} S \leq 2$. Using Jacobi-Zariski exact sequence arising from $A \rightarrow A_q \rightarrow S$ and the localization property, we have $\text{AQ-dim}_A S \leq 2$. Now Jacobi-Zariski exact sequence arising from the natural diagram $S \rightarrow A \rightarrow S$, and the flat base change imply that $D_n(S | R, -) = 0$, for $n \geq 2$. Thus φ is locally complete intersection by [6, Theorem (1.2)]. \square

Remark 3.11. At this time, we do not know if the assumption $\text{char}(k) = 0$ is necessary in Fact 3.9. However, as we will see in Proposition 3.13, when we work with R as an algebra retract of its trivial extension by an R -module M with $\text{Ann}_R(M) = 0$, we are able to remove this assumption. This can be done using a result of Herzog.

Recall that the *trivial extension of R by an R -module M* , denoted $R \times M$, is the underlying R -module $R \oplus M$ equipped with a ring structure given by the multiplication rule $(r, m) \cdot (r', m') = (rr', rm' + r'm)$. There are ring homomorphisms $\rho: R \rightarrow R \times M$ with $\rho(r) = (r, 0)$ and $\pi: R \times M \rightarrow R$ with $\pi(r, m) = r$. Note that the composition $\pi\rho$ is the identity on R , that is, R is an algebra retract of $R \times M$.

Lemma 3.12. *Let M be a finitely generated R -module. Then for each non-negative integer i we have $(\mu_R(M))^i \leq \beta_i^{R \times M}(R)$.*

Proof. By [17, Corollary 1] we have $P_R^{R \times M}(t) = 1 + tP_R^{R \times M}(t)P_M^R(t)$. Hence, by induction on i , there exists a sequence $\{\gamma_i\}$ of natural numbers such that $\beta_i^{R \times M}(R) = \gamma_i + (\mu_R(M))^i$. Therefore, $(\mu_R(M))^i \leq \beta_i^{R \times M}(R)$. \square

The following result gives a characterization of local rings admitting non-trivial semidualizing modules. Recall that a finitely generated R -module C is called *semidualizing* if $R \cong \text{Hom}_R(C, C)$ and $\text{Ext}_R^i(C, C) = 0$ for all $i > 0$. For instance, every finitely generated projective R -module of rank 1 is semidualizing. Also, every dualizing module is semidualizing with finite injective dimension.

Proposition 3.13. *Let M be a finitely generated R -module with $\text{Ann}_R(M) = 0$. Then the following conditions are equivalent:*

- (i) $M \cong R$.
- (ii) $\text{AQ-dim}_{R \times M} R = 2$.
- (iii) $\text{CI-dim}_{R \times M} R < \infty$.
- (iv) $\text{cx}_{R \times M} R < \infty$.

If one of the above conditions holds, then $\text{CI-dim}_{R \times M} R = 0$ and $\text{cx}_{R \times M} R = 1$.

Proof. (i) \implies (ii) Assume that $M \cong R$. It is straightforward to see that $\text{Ker } \pi = (x)$, where x is an exact zero divisor in the sense of [16], that is, $\text{Ann}_{R \times M}(x) \cong (R \times M)/x(R \times M)$. Hence, $\text{AQ-dim}_{R \times M} R = 2$ by [9, 1.7, 3.1].

(ii) \implies (iii) follows from Theorem A.

(iii) \implies (iv) follows from [8, Theorem (5.3)].

(iv) \implies (i) Assume that $\text{cx}_{R \times M} R < \infty$. Hence, the sequence of the Betti numbers of R over $R \times M$ has polynomial growth. Since, by Lemma 3.12, for each non-negative integer i we have $(\mu_R(M))^i \leq \beta_i^{R \times M}(R)$, we conclude that $\mu_R(M) = 1$, which implies that $M \cong R$.

Note that when $\text{CI-dim}_{R \times M} R < \infty$, then $\text{CI-dim}_{R \times M} R = 0$ by [26, Lemma 3.14]. Note also that when $M \cong R$, $\text{Ker } \pi$ is generated by an exact zero divisor x . Therefore,

$$\cdots \xrightarrow{x} R \times R \xrightarrow{x} R \times R \xrightarrow{\pi} R \rightarrow 0$$

is a free resolution of R over $R \times R$, that is, $\text{cx}_{R \times R} R = 1$. \square

The following is an application of Proposition 3.13 in a concrete case.

Example 3.14. Let $S = R[\underline{x}]/J$, where $\underline{x} = x_1, \dots, x_n$ for some integer $n \geq 1$, and $J = \langle x_i^{n_i} x_j^{n_j} \mid n_i + n_j = 2, \text{ and } 1 \leq i \leq j \leq n \rangle$. Then one of the integers $\text{cx}_S R$, $\text{CI-dim}_S R$, or $\text{AQ-dim}_S R$ is finite if and only if $n = 1$. In this case we have $\text{cx}_S R = 1$, $\text{CI-dim}_S R = 0$, and $\text{AQ-dim}_S R = 2$.

ACKNOWLEDGMENT

The author is deeply grateful to Javier Majadas Soto for his comments on Fact 2.7, and he is also grateful to Saeed Nasseh for his comments on this paper.

REFERENCES

1. M. André, *Méthodes simpliciales en algèbre homologique et algèbre commutative*, Lecture Notes in Mathematics, **32**, Springer-Verlag, Berlin-New York 1967 iii+122 pp.
2. ———, *Homologie des algèbres commutatives*, Springer-Verlag, Berlin, 1974, Die Grundlehren der mathematischen Wissenschaften, Band 206.
3. ———, *Localisation de la lissité formelle*, Manuscripta Math. **13** (1974), 297–307.
4. L. L. Avramov, *Descente des déviations par homomorphismes locaux et génération des idéaux de dimension projective finie*, (French), C. R. Acad. Sci. Paris Sér. I Math. **295** (1982), 665–668.
5. ———, *Infinite free resolutions*, Six lectures on commutative algebra (Bellaterra, 1996), Progr. Math., vol. 166, Birkhäuser, Basel, 1998, pp. 1–118.
6. ———, *Locally complete intersection homomorphisms and a conjecture of Quillen on the vanishing of cotangent homology*, Ann. of Math. **150** (1999), 455–487.
7. L. L. Avramov, H.-B. Foxby, and B. Herzog, *Structure of local homomorphisms*, J. Algebra **164** (1994), 124–145.
8. ———, *Complete intersection dimension*, Inst. Hautes Études Sci. Publ. Math. (1997), no. 86, 67–114 (1998).
9. L. L. Avramov, I. B. Henriques, and L. M. Şega, *Quasi-complete intersection homomorphisms*, Pure and Applied Mathematics Quarterly **9** (2013), no. 4, 1–31.
10. L. L. Avramov and S. B. Iyengar, *André-Quillen homology of algebra retracts*, Ann. Sci. École Norm. Sup. (4) **36** (2003), 431–462.
11. ———, *Finite generation of Hochschild homology algebras*, Invent. Math. **140** (2000), 143–170.
12. ———, *Homological criteria for regular homomorphisms and for locally complete intersection homomorphisms*, Algebra, arithmetic and geometry, Part I, II (Mumbai, 2000), Tata Inst. Fund. Res. Stud. Math. 16 Tata Inst. Fund. Res., Bombay, 2002, 97–122.
13. A. Blanco, J. Majadas, and A. G. Rodicio, *On the acyclicity of the Tate complex*, J. Pure Appl. Algebra **131** (1998), no. 2, 125–132.
14. W. Bruns and J. Herzog, *Cohen-Macaulay rings*, revised ed., Studies in Advanced Mathematics, vol. 39, University Press, Cambridge, 1998.
15. T. H. Gulliksen and G. Levin, *Homology of local rings*, Queen’s Paper in Pure and Applied Mathematics **20**, Kingston, Ont. 1969 x+192 pp.
16. I. B. Henriques and L. M. Şega, *Free resolutions over short Gorenstein local rings*, Math. Z. **267** (2011), 645–663.
17. J. Herzog, *Algebra retracts and Poincaré-series*, Manuscripta Math., **21** (1977), 307–314.
18. S. B. Iyengar, *André-Quillen homology of commutative algebras*, Interactions between homotopy theory and algebra, Contemp. Math. **436**, Amer. Math. Soc., Providence, RI, 2007.
19. S. B. Iyengar and S. Sather-Wagstaff, *G-dimension over local homomorphisms. Applications to the Frobenius endomorphism*, Illinois J. Math. **48** (2004), no. 1, 241–272.
20. G. Levin, *Large homomorphisms of local rings*, Math. Scand. **46** (1980), 209–215.
21. J. Majadas, A. G. Rodicio, *Smoothness, Regularity and Complete Intersection*, London Mathematical Society Lecture Note Series, **373**, Cambridge University Press, Cambridge, 2010.

22. D. Quillen, *On the (co-) homology of commutative rings*, Applications of Categorical Algebra (Proc. Sympos. Pure Math., Vol. XVII, New York, 1968), Amer. Math. Soc., Providence, **RI**, 1970, 65–87.
23. D. Quillen, *Homology of commutative rings*, mimeographed notes, MIT 1968.
24. A. G. Rodicio, *On a result of Avramov*, Manuscripta Math. **62** (1988), 181–185.
25. ———, *Complete intersection dimensions and Foxby classes*, J. Pure Appl. Algebra **212** (2008), no. 12, 2594–2611.
26. S. Sather-Wagstaff, T. Sharif, and D. White, *Comparison of relative cohomology theories with respect to semidualizing modules*, Math. Z. **264** (2010), 571–600.
27. J. J. M. Soto, *Finite complete intersection dimension and vanishing of André-Quillen homology*, J. Pure Appl. Algebra **146** (2000), 197–207.
28. J. Tate, *Homology of Noetherian rings and local rings*, Illinois J. Math. **1** (1957), 14–27.

SCHOOL OF MATHEMATICS, INSTITUTE FOR RESEARCH IN FUNDAMENTAL SCIENCES (IPM), P. O. BOX 19395-5746, TEHRAN, IRAN.

E-mail address: `sharif@ipm.ir`