

New Proofs for Several Properties of Capacities

Guangyan Jia

Qilu Securities Institute for Financial Studies, Shandong University
250199 Jinan, People's Republic of China

E-mail: jiagy@sdu.edu.cn

Na Zhang*

College of Management and Economics, Tianjin University
China Center for Social Computing, Tianjin University
300072 Tianjin, People's Republic of China

E-mail: zna1225@163.com

Abstract In this note, we find a new way to prove several properties of 2-alternating capacities.

Keywords: Alternating capacity · Probability measure · Monotone capacity · Minimal member

AMS 2010 subject classifications: primary 28A12

1 Introduction

Let Ω denote the basic set and \mathcal{B} the σ -algebra on Ω . A set function $c : \mathcal{B} \rightarrow [0, 1]$ is called a capacity if it satisfies:

(C1). $c(\Omega) = 1, c(\emptyset) = 0$;

(C2)(monotonicity). $c(A) \leq c(B)$ for any $A \subseteq B, A, B \in \mathcal{B}$.

A capacity c is called 2-alternating, if $c(A \cup B) + c(A \cap B) \leq c(A) + c(B)$. It is called a probability measure if $c(A \cup B) + c(A \cap B) = c(A) + c(B)$. We usually denote a probability measure by P .

*corresponding author

For any expectation E , we can define a capacity c by $c(A) = E[I_A]$, $\forall A \in \mathcal{B}$; on the other hand, for any capacity c , we can define expectation through Choquet integral, i.e., $E[X] = \int Xdc$. Choquet integral was first introduced by Choquet in 1953. The readers can refer to [1] or [2] for more details. In [2], Denneberg proved the following result.

Lemma 1.1 ([2, Chapter 6]) *If the integral with respect to a capacity c is subadditive, i.e.,*

$$\int (X + Y)dc \leq \int Xdc + \int Ydc,$$

then c is 2-alternating. Conversely, let c be a 2-alternating capacity, then for any \mathcal{B} -measurable square integrable functions X, Y ,

$$\int (X + Y)dc \leq \int Xdc + \int Ydc.$$

In order to prove the above result, Denneberg proved the following result.

Lemma 1.2 ([2, Lemma 6]) *Suppose that A_1, A_2, \dots, A_n is a partition of Ω , \mathcal{B} is a σ -algebra generated by A_1, A_2, \dots, A_n and $c : \mathcal{B} \rightarrow [0, 1]$ is a capacity. For any permutation π of $(1, \dots, n)$, we define*

$$S_i^\pi := \bigcup_{j=1}^i A_{\pi_j}, \quad i = 1, \dots, n, \quad S_0^\pi := \emptyset.$$

We define a probability measure P^π on \mathcal{B} by

$$P^\pi(A_{\pi_i}) := \mu(S_i^\pi) - \mu(S_{i-1}^\pi), \quad i = 1, \dots, n.$$

Suppose X is a \mathcal{B} -measurable real valued function X defined on Ω . If μ is 2-alternating, then

$$\int Xd\mu \geq \int XdP^\pi.$$

If $X(A_{\pi_1}) \geq X(A_{\pi_2}) \geq \dots \geq X(A_{\pi_n})$, the above equality holds.

Since Choquet integral is positive homogeneous, any Choquet expectation generated by a 2-alternating capacity is sublinear expectation. Jia [3] defined a partial order " \leq " on the set of expectations as follows:

for any two expectations E_1 and E_2 , $E_1 \leq E_2$ if for any \mathcal{B} -measurable square integrable random variable X , $E_1[X] \leq E_2[X]$,

and proved the following results.

Lemma 1.3 ([3, Theorem 2.7 and Theorem 3.1]) *E is a minimal member of the set of all the sublinear expectations if and only if E is a linear expectation. Suppose E_1 is a subadditive expectation, E_2 is a superadditive expectation and $E_1 \geq E_2$, then there exists a linear expectation E_0 such that $E_1 \geq E_0 \geq E_2$.*

Therefore we may wonder if the minimal members of all the 2-alternating capacities are exactly the probability measures? In fact, the answer is positive, since the following result holds:

Suppose \mathcal{Q} is a set, \prec is a semiorder defined on \mathcal{Q} and \mathcal{U} denotes the set of all the minimal members in \mathcal{Q} . Thus for any set \mathcal{Z} satisfying $\mathcal{U} \subset \mathcal{Z} \subset \mathcal{Q}$, the set of all the minimal members of \mathcal{Z} is still \mathcal{U} .

In this note, we'll give another proof of the above results by means of capacity only. The method of constructing a probability measure step by step from a 2-alternating capacity is also given.

2 Main results

First we list the following definitions which will be used below. A capacity defined on (Ω, \mathcal{B}) is said to be:

- *2-monotone* if $c(A \cup B) + c(A \cap B) \geq c(A) + c(B)$;
- *n-alternating* if $c(\bigcap_{i=1}^n A_i) \leq \sum_{\emptyset \neq I \subset \{1, \dots, n\}} (-1)^{|I|+1} v(\bigcup_{i \in I} A_i), \forall A_1, \dots, A_n \in \mathcal{B}$;
- *n-monotone* if $c(\bigcup_{i=1}^n A_i) \geq \sum_{\emptyset \neq I \subset \{1, \dots, n\}} (-1)^{|I|+1} v(\bigcap_{i \in I} A_i), \forall A_1, \dots, A_n \in \mathcal{B}$;
- *∞ -alternating* if c is n -alternating, for all n ;
- *∞ -monotone* if c is n -monotone, for all n .

Furthermore, we have the following notations.

- \mathcal{A}_n denotes the set of n -alternating capacities, for any $n \geq 2$;
- \mathcal{M}_n denotes the set of n -monotone capacities, for any $n \geq 2$;
- \mathcal{P} denotes the set of probability measures;
- \mathcal{A}_∞ denotes the set of ∞ -alternating capacities;
- \mathcal{M}_∞ the set of ∞ -monotone capacities.

It is known that $\mathcal{P} \subseteq \mathcal{A}_\infty \subseteq \mathcal{A}_{n+1} \subseteq \mathcal{A}_n$, $\mathcal{P} \subseteq \mathcal{M}_\infty \subseteq \mathcal{M}_{n+1} \subseteq \mathcal{M}_n$ and $\mathcal{A}_n \cap \mathcal{M}_m = \mathcal{P}$ for any $n \geq 2$ and $m \geq 2$.

Now let us define the partial order " \leq ": for any two capacities c_1 and c_2 , $c_1 \leq c_2$ means that $c_1(A) \leq c_2(A)$, for all $A \in \mathcal{B}$. If $c_1 \leq c_2$, we can also denote by $c_2 \geq c_1$. If $c_1 \leq c_2$ and $c_1 \geq c_2$, we have $c_1 = c_2$.

The following lemma holds.

Lemma 2.1 *Let $\mathcal{T} \subset \mathcal{A}_2$ be a nonempty set and totally ordered (for each pair $c_1, c_2 \in \mathcal{T}$, one has either $c_1 \leq c_2$ or $c_2 \leq c_1$). Then the set function*

$$\nu(A) \triangleq \inf_{c \in \mathcal{T}} c(A), \quad A \in \mathcal{B},$$

is a 2-alternating capacity, that is $\nu \in \mathcal{A}_2$.

Proof. It is obvious that $\nu(\Omega) = 1, \nu(\emptyset) = 0$ and ν is monotone. We now prove that it is 2-alternating.

$$\begin{aligned} \nu(A \cap B) &= \inf_{c \in \mathcal{T}} c(A \cap B) \\ &\leq \inf_{c \in \mathcal{T}} \{c(A) + c(B) - c(A \cup B)\} \\ &\leq \inf_{c \in \mathcal{T}} \{c(A) + c(B)\} - \nu(A \cup B). \end{aligned}$$

Since \mathcal{T} is totally ordered, for $c_1, c_2 \in \mathcal{T}$, we suppose, without loss of generality, that $c_1 \leq c_2$, so $c_1(A) + c_2(B) \geq c_1(A) + c_1(B)$. Therefore,

$$\begin{aligned} \nu(A \cap B) &\leq \inf_{c_1, c_2 \in \mathcal{T}} \{c_1(A) + c_2(B)\} - \nu(A \cup B) \\ &= \inf_{c \in \mathcal{T}} \{c(A)\} + \inf_{c \in \mathcal{T}} \{c(B)\} - \nu(A \cup B) \\ &= \nu(A) + \nu(B) - \nu(A \cup B). \end{aligned}$$

Thus the result follows. \square

Theorem 2.1 *Any $P \in \mathcal{P}$ is a minimal member of \mathcal{A}_2 . Conversely, if c is a minimal member of \mathcal{A}_2 , then $c \in \mathcal{P}$.*

Proof. Suppose $c \in \mathcal{A}_2, c \leq P$. Then we have

$$\forall A \in \mathcal{B}, 1 - c(A^c) \leq c(A) \leq P(A) = 1 - P(A^c).$$

Since $c(A^c) \leq P(A^c)$, we have $c(A) = P(A)$, which means that there is no 2-alternating capacity c satisfying $c \leq P$, i.e., P is a minimal member of \mathcal{A}_2 .

If c is a minimal member of \mathcal{A}_2 , for a fixed $A \in \mathcal{B}$, we define

$$c^A(B) := c(A \cup B) + c(A \cap B) - c(A), \forall B \in \mathcal{B}.$$

Obviously, $c^A \leq c$, $c^A(\Omega) = c(\Omega) + c(A) - c(A) = 1$, $c^A(\emptyset) = c(A) + 0 - c(A) = 0$. The monotonicity of c^A can be easily deduced by the monotonicity of c . For any $B \in \mathcal{B}, F \in \mathcal{B}$,

$$\begin{aligned} c^A(B \cup F) + c^A(B \cap F) &= c(A \cup (B \cup F)) + c(A \cap (B \cup F)) - c(A) \\ &\quad + c(A \cup (B \cap F)) + c(A \cap (B \cap F)) - c(A) \\ &= c((A \cup B) \cup (A \cup F)) + c((A \cap B) \cup (A \cap F)) \\ &\quad + c((A \cup B) \cap (A \cup F)) + c((A \cap B) \cap (A \cap F)) - 2c(A) \\ &\leq c(A \cup B) + c(A \cup F) + c(A \cap B) + c(A \cap F) - 2c(A) \\ &= c^A(B) + c^A(F), \end{aligned}$$

i.e., $c^A \in \mathcal{A}_2$. Note that c is the minimal member of \mathcal{A}_2 , thus $c^A = c$, which means that, for any $B \in \mathcal{B}$, we have $c(A \cup B) + c(A \cap B) = c(A) + c(B)$. Since A can be any set in \mathcal{B} , c is a probability measure. \square

Remark 2.1 1. *By similar proof we can deduce that any minimal member of \mathcal{A}_n ($n \geq 2$) (resp. \mathcal{A}_∞) can only be probability measure and any probability measure is its minimal member.*

2. *The maximal member of \mathcal{M}_n ($n \geq 2$) (resp. \mathcal{M}_∞) can only be probability measure and any probability measure is its maximal member.*

Definition 2.1 *For a capacity c , we define the invariant subfield \mathcal{B}^c of c as follows:*

$$\mathcal{B}^c \triangleq \{A \in \mathcal{B} : \forall B \in \mathcal{B}, c(A \cup B) + c(A \cap B) = c(A) + c(B)\}$$

It is obvious that \mathcal{B}^c is nonempty, since $\Omega \in \mathcal{B}^c$ and $\emptyset \in \mathcal{B}^c$. A capacity c is a probability measure if and only if $\mathcal{B}^c = \mathcal{B}$. Note that if c is a 2-alternating capacity, then for all $A \in \mathcal{B}$, such that $c(A) = 0$, we have $A \in \mathcal{B}^c$. If c is a 2-monotone capacity, then for all $A \in \mathcal{B}$, such that $c(A) = 1$, we have $A \in \mathcal{B}^c$.

$\forall c \in \mathcal{A}_2$, it has been proved that $c^F \in \mathcal{A}_2$. Thus we can define the following mapping.

Definition 2.2 *For all $\mathcal{F} \in \mathcal{B}$, we define mapping $\Pi^{\mathcal{F}} : \mathcal{A}^2 \rightarrow \mathcal{A}^2$ as follows:*

$$\Pi^{\mathcal{F}} c = c^{\mathcal{F}}.$$

Proposition 2.1 *The following properties about invariant subfield and the above mapping hold.*

(i). $\forall c \in \mathcal{A}_2, \Pi^{\mathcal{F}}(c) \leq c$;

(ii). $\forall A \in \mathcal{B}$, if $A \subset \mathcal{F}$ or $\mathcal{F} \subset A$, one has $c^{\mathcal{F}}(A) = c(A)$;

(iii). $\forall A \in \mathcal{B}^c, c^{\mathcal{F}}(A) = c(A)$;

(iv). $\mathcal{F} \in \mathcal{B}^{c^{\mathcal{F}}}$;

(v). $\mathcal{B}^c \subset \mathcal{B}^{c^{\mathcal{F}}}$;

(vi). If $\mathcal{F} \in \mathcal{B}^c, c^{\mathcal{F}} = c$.

Proof. (i) and (vi) are obvious.

(ii). Without lost of generality, suppose that $A \subset \mathcal{F}$, thus $c^{\mathcal{F}}(A) = c(A \cup \mathcal{F}) + c(A \cap \mathcal{F}) - c(\mathcal{F}) = c(A) + c(\mathcal{F}) - c(\mathcal{F}) = c(A)$.

(iii). For all $A \in \mathcal{B}^c, c^{\mathcal{F}}(A) = c(\mathcal{F} \cup A) + c(\mathcal{F} \cap A) - c(\mathcal{F}) = c(A)$.

(iv). According to (ii), for all $A \in \mathcal{B}$,

$$\begin{aligned} & c^F(F \cup A) + c^F(F \cap A) - c^F(F) - c^F(A) \\ &= c(F \cup A) + c(A \cap F) - c(F) - (c(A \cup F) + c(A \cap F) - c(F)) = 0. \end{aligned}$$

(v). Suppose $A \in \mathcal{B}^c$, $B \in \mathcal{B}$,

$$\begin{aligned} & c^F(A \cup B) + c^F(A \cap B) - c^F(A) - c^F(B) \\ &= c(F \cup (A \cup B)) + c(F \cap (A \cup B)) - c(F) + c(F \cup (A \cap B)) \\ & \quad + c(F \cap (A \cap B)) - c(F) - c(A) - c(F \cup B) - c(F \cap B) + c(F) \\ &= [c(A) + c(F \cup B) - c(A \cap (F \cup B))] + c(F \cap (A \cup B)) - c(F) \\ & \quad + c(F \cup (A \cap B)) + [c(A) + c(F \cap B) - c(A \cup (F \cap B))] \\ & \quad - c(F) - c(A) - c(F \cup B) - c(F \cap B) + c(F) \\ &= c(A) - c(A \cap (F \cup B)) + c(F \cap (A \cup B)) - c(F) + c(F \cup (A \cap B)) \\ & \quad - c(A \cup (F \cap B)) \\ &= [c(F \cup (A \cap B)) - c(A \cap (F \cup B))] \\ & \quad + [c(F \cap (A \cup B)) - c(A \cup (F \cap B))] + c(A) - c(F) \\ &= [c(F \cup (A \cap B)) - c((F \cup (A \cap B)) \cap A)] \\ & \quad + [c(F \cap (A \cup B)) - c((F \cap (A \cup B)) \cup A)] + c(A) - c(F) \\ &= [c((F \cup (A \cap B)) \cup A) - c(A)] + [c((F \cap (A \cup B)) \cap A) - c(A)] \\ & \quad + c(A) - c(F) \\ &= [c(F \cup A) - c(A)] + [c(F \cap A) - c(A)] + c(A) - c(F) = 0. \end{aligned}$$

□

With the help of this mapping, we can prove Lemma 1.2, i.e., the following theorem, by way of capacity.

Theorem 2.2 *Consider (Ω, \mathcal{B}) . Suppose that \mathcal{B} is finite, c is a 2-alternating capacity defined on \mathcal{B} . Take $F_1, \dots, F_n \in \mathcal{B}$ such that $F_1 \subset F_2 \subset \dots \subset F_n$. Thus there exists a probability measure P , such that $P(F_i) = c(F_i)$, for all $i = 1, \dots, n$ and $P \leq c$.*

Proof. First, we design a cyclic program as follows.

Set $\mu = c$.

Step I: Check F_i , $i = 1, \dots, n$. If all the sets F_i belong to the invariant subfield of μ , go straight to Step III. Otherwise, suppose that F_i does not belong to the subfield of μ . By Proposition 2.1, the following result holds:

$$\mu^{F_i}(F_j) = \mu(F_j), \forall j = 1, \dots, n,$$

i.e., μ^{F_i} and μ are equal on F_j , $j = 1, \dots, n$;

$$F_i \in \mathcal{B}^{\mu^{F_i}}, \quad \mathcal{B}^\mu \subset \mathcal{B}^{\mu^{F_i}},$$

i.e., from μ to μ^{F_i} , the invariant subfield is enlarged and F_i is also included.

Step II: Update μ by c^{F_i} . The invariant subfield of μ is enlarged by Step I. Repeating the procedures in Step I.

Step III: We get the final μ , which satisfies $\mu \in \mathcal{A}_2$, $\mu \leq c$, and for all $i = 1, \dots, n$, $\mu(F_i) \equiv c(F_i)$, $F_i \in \mathcal{B}^\mu$, $\mathcal{B} \subset \mathcal{B}^\mu$.

Next, we consider μ , and design another cyclic program.

Step 1: Check \mathcal{B}^μ and \mathcal{B} . If they are the same, go straight to Step 3. Otherwise, suppose $A \in \mathcal{B}/\mathcal{B}^{cF}$. Consider the transformation of μ induced by A . By Proposition 2.1, we have $\mu^A(F_i) \equiv \mu(F_i)$, $i = 1, \dots, n$.

Step 2: Update μ by μ^A . The invariant subfield of μ is enlarged. Repeat the procedures in Step 1.

Step 3: μ satisfies the following conditions: for all $i = 1, \dots, n$, $\mu(F_i) = c(F_i)$, $\mu \leq c$. Furthermore, $\mathcal{B}^\mu = \mathcal{B}$, thus μ is just the probability measure satisfying the conditions needed. The proof is complete. \square

Theorem 2.3 Consider space (Ω, \mathcal{B}) . Suppose that \mathcal{B} is finite, μ is a 2-alternating capacity defined on \mathcal{B} , ν is a 2-monotone capacity defined on \mathcal{B} . If $\mu \geq \nu$, there exists a probability measure P such that $\mu \geq P \geq \nu$.

Proof. Since \mathcal{B} is finite, we can take $A \in \mathcal{B}/\mathcal{B}^\mu$ such that

$$\mu(A) - \nu(A) = \min_{B \in \mathcal{B}/\mathcal{B}^\mu} \{\mu(B) - \nu(B)\}.$$

Make transformation Π^A on μ , thus

$$\mu^A(B) = \mu(A \cup B) + \mu(A \cap B) - \mu(A) \geq \nu(A \cup B) + \nu(A \cap B) - \nu(A) \geq \nu(B),$$

i.e., $\mu \geq \mu^A \geq \nu$. By Proposition 2.1,

$$\mathcal{B}^\mu \subset \mathcal{B}^{\mu^A},$$

and $\mathcal{B}^\mu \neq \mathcal{B}^{\mu^A}$.

For μ^A , repeat the above procedure, until we get a capacity P , such that $\mathcal{B}^P = \mathcal{B}$. P satisfying that $\mu \geq P \geq \nu$. The proof is complete. \square

Remark 2.2 According the above theorem, we may get different probability measures if we make transformation by different sets or in a different order.

References

- [1] Choquet, G., Theory of capacities, *Annales de l'Institut Fourier*, 5 (1953-1954), 131-295.
- [2] Dieter Denneberg, *Non-additive measure and integral*, Kluwer Academic Publishers, Boston, 1994, 184 pp.
- [3] Guangyan Jia, The minimal sublinear expectations and their related properties, *Science in China Ser. A: Mathematics*. 39(2009), 79-87.