DISTRIBUTION OF THE ROOTS OF THE EQUATIONS Z(t)=0, Z'(t)=0 IN THE THEORY OF THE RIEMANN ZETA-FUNCTION

JAN MOSER

ABSTRACT. Let the symbols $\{\gamma\}$, $\{t_0\}$; $t_0 \neq \gamma$ denote the sequences of the roots of the equations

$$Z(t) = 0$$
, and $Z'(t) = 0$,

respectively, and

$$m(t_0) = \min\{\gamma'' - t_0, t_0 - \gamma'\}, \quad Q(t_0) = \max\{\gamma'' - t_0, t_0 - \gamma'\}, \quad \gamma' < t_0 < \gamma'',$$

where γ', γ'' are the neighboring zeroes. We have proved the following in this paper: on the Riemann hypothesis we have

$$\frac{Q(t_0)}{m(t_0)} < t_0 \ln^2 t_0 \ln_2 t_0 \ln_3 t_0, \quad t_0 \to \infty.$$

This paper is the English version of the paper of ref. [5].

1. Formulation of Theorem

We will denote by the symbols

$$\{\gamma\}, \{t_0\}$$

the sequences of the roots of the equations

$$Z(t) = 0, \quad Z'(t) = 0, \quad t_0 \neq \gamma,$$

where (see [6], pp. 79, 329)

(1.1)
$$Z(t) = e^{i\vartheta(t)} \zeta\left(\frac{1}{2} + it\right),$$

$$\vartheta(t) = -\frac{t}{2} \ln \pi + \operatorname{Im} \ln \Gamma\left(\frac{1}{4} + \frac{1}{2}it\right) =$$

$$= \frac{t}{2} \ln \frac{t}{2\pi} - \frac{t}{2} - \frac{\pi}{8} + \mathcal{O}\left(\frac{1}{t}\right).$$

Remark. On the Riemann hypothesis the points of the sequences $\{\gamma\}, \{t_0\}$ are separated each from other, i. e.

$$\gamma' < t_0 < \gamma''$$

where γ', γ'' are neighboring points of the sequence $\{\gamma\}$ (see [4], Corollary 3).

The sequence $\{t_0\}$ itself oscillates in a complicated manner around the sequence $\{\gamma\}$, (comp., for example, the graph of the function Z(t) in the neighborhood of the first Lehmer pair of zeroes in [1], p. 296). Let further

(1.2)
$$Q(t_0) = \max\{\gamma'' - t_0, t_0 - \gamma'\},\ m(t_0) = \min\{\gamma'' - t_0, t_0 - \gamma'\}.$$

Then the quotient

$$\frac{Q(t_0)}{m(t_0)}$$

measures the asymmetry of the point t_0 with respect to the points γ', γ'' .

In this paper we prove the following statement.

Theorem. On the Riemann hypothesis we have

(1.3)
$$\frac{Q(t_0)}{m(t_0)} < t_0 \ln^2 t_0 \ln_2 t_0 \ln_3 t_0, \quad t_0 \to \infty,$$

where

$$\ln_2 t_0 = \ln \ln t_0, \dots$$

The estimate (1.3) follows from our formula

(1.4)
$$\frac{\pi}{4} = \sum_{\gamma} \frac{t_0}{\gamma^2 - t_0^2} + \mathcal{O}\left(\frac{1}{t_0}\right)$$

that is a consequence of the Riemann hypothesis (see [3], p. 117, comp. [4]). The formula (1.4) is a conjugate formula with respect to the Riemann formula

$$c + 2 - \ln 4\pi = \sum_{\gamma} \frac{1}{\frac{1}{4} + \gamma^2},$$

where c stands for the Euler's constant. Let us remind, finally, that on the Riemann hypothesis also the Littlewood estimate

(1.5)
$$\gamma'' - \gamma' < \frac{A}{\ln \ln \gamma'}, \quad \gamma' \to \infty$$

holds true.

2. Lemmas

Lemma 1.

(2.1)
$$\sum_{t_0 + T < \gamma} \frac{1}{\gamma^2 - t_0^2} < A \frac{\ln T}{T},$$

where

$$(2.2) T = T(t_0) = t_0 \ln t_0 \ln_2 t_0.$$

Proof. Let N(t) denote the number of zeroes

$$\beta + i\gamma$$

of the function

$$\zeta(s)$$
, $s = \sigma + it$, $\beta \in (0,1)$, $\gamma \in (0,t]$.

Since (see [6], p. 178)

$$N(t+1) - N(t) < A \ln t,$$

then

$$\begin{split} & \sum_{t_0+T \leq \gamma} \frac{1}{\gamma^2 - t_0^2} = \sum_{n=1}^{\infty} \sum_{\{t_0+T+n-1 < \gamma \leq t_0+T+n\}} \frac{1}{\gamma^2 - t_0^2} < \\ & < A \sum_{n=1}^{\infty} \frac{\ln(n+T-1+t_0)}{(n+T-1)(n+T-1-2t_0)} < \\ & < A \frac{\ln(T+t_0)}{T(T+2t_0)} + A \int_{1}^{\infty} \frac{\ln(x+T-1+t_0)}{(x+T-1)(x+T-1+2t_0)} \mathrm{d}x. \end{split}$$

Following the identity

$$\int_{-\infty}^{\infty} \tau e^{-\tau} d\tau = e^{-a}(a+1)$$

we obtain

$$\int_{1}^{\infty} \frac{\ln(x+T-1+t_0)}{(x+T-1)(x+T-1+2t_0)} dx < \int_{1}^{\infty} \frac{\ln(x+T-1+t_0)}{(x+T-1)(x+T-1+t_0)} dx =$$

$$= \int_{\ln(T+t_0)}^{\infty} \frac{\tau}{e^{\tau}-t_0} d\tau = \int_{\ln(T+t_0)}^{\infty} \frac{e^{-\tau}\tau}{1-t_0e^{-\tau}} d\tau < \frac{T+t_0}{T} \int_{\ln(T+t_0)}^{\infty} e^{-\tau}\tau d\tau =$$

$$= \frac{1}{T} \left\{ \ln(T+t_0) + 1 \right\} < A \frac{\ln T}{T},$$

i. e. we have finished the proof.

Consequently, we have from (1.4) by (2.1), (2.2) the following

Lemma 2. On the Riemann hypothesis we have

(2.3)
$$\frac{\pi}{8t_0} = \sum_{0 < \gamma < t_0 + T} \frac{1}{\gamma^2 - t_0^2} + \mathcal{O}\left(\frac{1}{t_0 \ln_2 t_0}\right), \quad t_0 \to \infty,$$
$$T = T(t_0) = t_0 \ln t_0 \ln_2 t_0.$$

3. Proof of the Theorem

If there is any \tilde{t}_0 such that

(3.1)
$$K(\tilde{t}_0) = \frac{Q(\tilde{t}_0)}{m(\tilde{t}_0)} \ge \tilde{t}_0 \ln^2 \tilde{t}_0 \ln_2 \tilde{t}_0 \ln_3 \tilde{t}_0$$

then we have:

(A). If

(3.2)
$$\tilde{t}_0 - \tilde{\gamma}' = K(\tilde{t}_0)(\tilde{\gamma}'' - \tilde{t}_0)$$

then by (2.3) there is some A > 0 such that

(3.3)
$$\frac{A}{\tilde{t}_0} > \sum_{0 < \gamma < \tilde{t}_0 + \tilde{T}} \frac{1}{\gamma^2 - \tilde{t}_0^2}, \quad \tilde{T} = T(\tilde{t}_0).$$

Now $\tilde{\gamma}' < \tilde{t}_0 < \tilde{\gamma}''$ and the symbol $n(\gamma)$ denotes the order of the zero $\frac{1}{2} + i\gamma$. We have

$$(3.4) \sum_{0<\gamma\leq\tilde{t}_{0}+\tilde{T}}\frac{1}{\gamma^{2}-\tilde{t}_{0}^{2}} = \sum_{0<\gamma\leq\tilde{\gamma}'}\frac{1}{\gamma^{2}-\tilde{t}_{0}^{2}} + \frac{n(\tilde{\gamma}'')}{(\tilde{\gamma}'')^{2}-\tilde{t}_{0}^{2}} + \sum_{\tilde{\gamma}''<\gamma\leq\tilde{t}_{0}+\tilde{T}}\frac{1}{\gamma^{2}-\tilde{t}_{0}^{2}} > \\ > \sum_{0<\gamma<\tilde{\gamma}'}\frac{1}{\gamma^{2}-\tilde{t}_{0}^{2}} + \frac{1}{(\tilde{\gamma}'')^{2}-\tilde{t}_{0}^{2}}.$$

Since (see [6], p. 181)

$$(3.5) N(t) < At \ln t,$$

then

$$\sum_{0<\gamma<\tilde{\gamma}'} \frac{1}{\gamma^{2} - \tilde{t}_{0}^{2}} = \sum_{0<\gamma<\tilde{t}_{0}} \frac{1}{\gamma^{2} - \tilde{t}_{0}^{2}} = -\sum_{0<\gamma<\tilde{t}_{0}} \frac{1}{(\tilde{t}_{0} - \gamma)(\gamma + \tilde{t}_{0})} >$$

$$> -\frac{1}{\tilde{t}_{0} - \tilde{\gamma}'} \sum_{0<\gamma<\tilde{t}_{0}} \frac{1}{\gamma + \tilde{t}_{0}} > -\frac{1}{\tilde{t}_{0}(\tilde{t}_{0} - \tilde{\gamma}')} \sum_{0<\gamma<\tilde{t}_{0}} 1 >$$

$$> -\frac{1}{\tilde{t}_{0}(\tilde{t}_{0} - \tilde{\gamma}')} A \tilde{t}_{0} \ln \tilde{t}_{0} = -A \frac{\ln \tilde{t}_{0}}{\tilde{t}_{0} - \tilde{\gamma}'},$$

and by (3.2)

$$(3.7) \frac{1}{(\tilde{\gamma}'')^2 - \tilde{t}_0^2} = \frac{1}{(\tilde{\gamma}'' - \tilde{t}_0)(\tilde{\gamma}'' + \tilde{t}_0)} > \frac{1}{2\tilde{\gamma}''(\tilde{\gamma}'' - \tilde{t}_0)} > A \frac{K(\tilde{t}_0)}{\tilde{t}_0(\tilde{t}_0 - \tilde{\gamma}')}.$$

Consequently, from (3.4) by (1.5), (3.1), (3.6), (3.7) we have

(3.8)
$$\sum_{0 < \gamma \le \tilde{t}_0 + \tilde{T}} \frac{1}{\gamma^2 - \tilde{t}_0^2} > \frac{A}{\tilde{t}_0 - \tilde{\gamma}'} \left\{ \frac{K(\tilde{t}_0)}{\tilde{t}_0} - \ln \tilde{t}_0 \right\} >$$
$$> \frac{A}{\tilde{\gamma}'' - \tilde{\gamma}'} \ln^2 \tilde{t}_0 \ln_2 \tilde{t}_0 \ln_3 \tilde{t}_0 > A \ln^2 \tilde{t}_0 \ln_2^2 \tilde{t}_0 \ln_3 \tilde{t}_0,$$

and finally (see (3.3), (3.8))

$$\frac{1}{\tilde{t}_0} > A \ln^2 \tilde{t}_0 \ln_2^2 \tilde{t}_0 \ln_3 \tilde{t}_0,$$

i. e. we obtain the contradiction.

(B). If

(3.9)
$$\tilde{\gamma}'' - \tilde{t}_0 = K(\tilde{t}_0)(\tilde{t}_0 - \tilde{\gamma}')$$

then by (2.3) there is some A > 0 such that

$$\frac{A}{\tilde{t}_0} < \sum_{0 < \gamma < \tilde{t}_0 + \tilde{T}} \frac{1}{\gamma^2 - \tilde{t}_0^2}.$$

In this case we have the following lower estimate for the corresponding sums: by (2.2) and (3.5) one obtains

(3.11)
$$\sum_{0 < \gamma \leq \tilde{t}_0 + \tilde{T}} 1 < A\tilde{T} \ln \tilde{T} < A\tilde{t}_0 \ln^2 \tilde{t}_0 \ln_2 \tilde{t}_0,$$

next, by (3.9)

$$\sum_{0 < \gamma \le \tilde{\gamma}'} \frac{1}{\gamma^2 - \tilde{t}_0^2} < \frac{1}{(\tilde{\gamma}')^2 - \tilde{t}_0^2} = -\frac{K(\tilde{t}_0)}{(\tilde{\gamma}' + \tilde{t}_0)(\tilde{\gamma}'' - \tilde{t}_0)} < -A \frac{K(\tilde{t}_0)}{\tilde{t}_0(\tilde{\gamma}' - \tilde{t}_0)},$$

and finally (see (3.11))

$$\sum_{\tilde{\gamma}'' < \gamma \leq \tilde{t}_0 + \tilde{T}} \frac{1}{\gamma^2 - \tilde{t}_0^2} < \frac{1}{(\tilde{\gamma}'')^2 - \tilde{t}_0^2} \sum_{\tilde{\gamma}'' < \gamma \leq \tilde{t}_0 + \tilde{T}} 1 < A \frac{\ln^2 \tilde{t}_0 \ln_2 \tilde{t}_0}{\tilde{\gamma}'' - \tilde{t}_0}.$$

Consequently,

(3.12)
$$\sum_{0 < \gamma \le \tilde{t}_0 + \tilde{T}} \frac{1}{\gamma^2 - \tilde{t}_0^2} = \left\{ \sum_{0 < \gamma \le \tilde{\gamma}''} + \sum_{\tilde{\gamma}'' \le \gamma \le \tilde{t}_0 + \tilde{T}} \right\} \frac{1}{\gamma^2 - \tilde{t}_0^2} < \frac{A}{\tilde{\gamma}'' - \tilde{t}_0} \left\{ -\frac{K(\tilde{t}_0)}{\tilde{t}_0} + \ln^2 \tilde{t}_0 \ln_2 \tilde{t}_0 \right\}.$$

Hence, from (3.10) by (3.1), (3.12) we have that

$$\frac{1}{\tilde{t}_0} < \frac{A}{\tilde{\gamma}'' - \tilde{t}_0} \{ -\ln_3 \tilde{t}_0 + 1 \} \ln^2 \tilde{t}_0 \ln_2 \tilde{t}_0 < 0,$$

i. e. we have obtained the contradiction.

I would like to thank Michal Demetrian for helping me with the electronic version of this work.

REFERENCES

- [1] D.H. Lehmer, 'On the roots of the Riemann zeta-function', Acta. Math., 95 (1956), 291-298.
- [2] J.E. Littlewood, 'Two notes on the Riemann zeta function', Proc. Cambr. Phil. Soc., 22 (1924), 234-242.
- [3] J. Moser, 'Some consequences from the Riemann hypothesis', Acta F.R.N. Univ. Comeniana. Math., 29 (1974), 111-118, (in Russian).
- [4] J. Moser, 'Some properties of the Riemann zeta-function on the critical line', Acta Arith. 26 (1974), 33-39, (in Russian); arXiv: 0710.0943
- [5] J. Moser, 'Distribution of the roots of the equations Z(t) = 0, Z'(t) = 0 in the theory of the Riemann zeta-function', Acta Math. Univ. Comeniana, 52-53 (1988), 7-14.
- [6] E.C. Titchmarsh, 'The theory of the Riemann zeta-function', Clarendon Press, Oxford, 1951.

Department of Mathematical Analysis and Numerical Mathematics, Comenius University, Mlynska Dolina M105, 842 48 Bratislava, SLOVAKIA

E-mail address: jan.mozer@fmph.uniba.sk