The spread of the spectrum of a nonnegative matrix with a zero diagonal element \mathbb{R}

Roman Drnovšek

Department of Mathematics, Faculty of Mathematics and Physics, University of Ljubljana, Jadranska 19, SI-1000 Ljubljana, Slovenia

Abstract

Let $A = [a_{ij}]_{i,j=1}^n$ be a nonnegative matrix with $a_{11} = 0$. We prove some lower bounds for the spread $s(A)$ of A that is defined as the maximum distance between any two eigenvalues of A. If A has only two distinct eigenvalues, then $s(A) \geq \frac{n}{2(n-1)} r(A)$, where $r(A)$ is the spectral radius of A. Moreover, this lower bound is the best possible.

Keywords: nonnegative matrices, spectrum, spread *2010 MSC:* 15B48, 15A42

1. Introduction

Let A be a complex $n \times n$ matrix with the spectrum $\{\lambda_1, \lambda_2, \ldots, \lambda_n\}.$ The spectral radius and the trace of A are denoted by $r(A)$ and $tr(A)$, respectively. The spread $s(A)$ of A is the maximum distance between any two eigenvalues, that is, $s(A) = \max_{i,j} |\lambda_i - \lambda_j|$. This quantity was introduced by Mirsky [\[4](#page-9-0)], and it has been studied by several authors; see e.g. [\[3](#page-9-1)] and the references therein. Note that $s(\lambda A) = |\lambda|s(A)$ for every complex number λ and that the spread of a nilpotent matrix is zero. Thus, when studying the

[✩]The paper will appear in Linear Algebra and its Applications.

Email address: roman.drnovsek@fmf.uni-lj.si (Roman Drnovšek)

spread of a matrix A, there is no loss of generality in assuming that $r(A) = 1$.

Let \mathcal{C}_n (with $n \geq 2$) be the collection of all nonnegative $n \times n$ matrices $A = [a_{ij}]_{i,j=1}^n$ such that $a_{11} = 0$ and $r(A) = 1$. It is not difficult to prove (see e.g. Proposition [2.1\)](#page-1-0) that the spread of a matrix $A \in \mathcal{C}_n$ cannot be zero, that is, the number 1 cannot be the only point in the spectrum of A . This motivates searching for lower bounds for the spread of A. If A has only two distinct eigenvalues, we prove that $s(A) \geq \frac{n}{2(n-1)}$, and we provide a matrix for which this lower bound is achieved. Such a matrix is necessarily irreducible, that is, there exists no permutation matrix P such that

$$
P^T A P = \begin{bmatrix} A_{11} & A_{12} \\ 0 & A_{22} \end{bmatrix},
$$

where A_{11} and A_{22} are square matrices.

2. Results

We start with an easy observation.

Proposition 2.1. Let A be a nonnegative $n \times n$ matrix with the spectral *radius* $r(A) = 1$ *. If* A *has* k *zero* diagonal elements, then

$$
s(A) \geq \frac{k}{n} .
$$

In particular, if $A \in \mathcal{C}_n$ *then*

$$
s(A) \geq \frac{1}{n} .
$$

PROOF. Since A is a nonnegative matrix, the spectral radius $r(A) = 1$ is its Perron eigenvalue. We denote it by λ_1 , while the rest eigenvalues of A are denoted by $\lambda_2, \lambda_3, \ldots, \lambda_n$. For every $i = 1, 2, \ldots, n$ we have

$$
Re (1 - \lambda_i) \le |1 - \lambda_i| = |\lambda_1 - \lambda_i| \le s(A),
$$

and so $1 - s(A) \leq \text{Re }\lambda_i$. It follows that

$$
n(1 - s(A)) \le \sum_{i=1}^{n} \text{Re } \lambda_i = \sum_{i=1}^{n} \lambda_i = \text{tr}(A).
$$

However, $tr(A) = \sum_{i=1}^{n} a_{ii} \leq n - k$, as A has k zero diagonal elements and $a_{ii} \le r(A) = 1$ for all i. We thus obtain that $n(1 - s(A)) \le n - k$, and so $n s(A) \geq k$ as asserted.

Applying the known inequalities of Johnson, Loewy and London we will prove a better result for matrices in \mathcal{C}_n . Let A be a nonnegative $n \times n$ matrix and let $s_k := \text{tr}(A^k)$ for $k \in \mathbb{N}$. The JLL-inequalities (discovered independently by Loewy and London [\[2](#page-9-2)], and Johnson [\[1](#page-9-3)]) state that

$$
s_k^m \le n^{m-1} s_{km}
$$

for all positive integers k and m . A slight modification of their proof gives the following inequalities.

Proposition 2.2. Let A be a nonnegative $n \times n$ matrix with k zero diagonal *elements. Then*

$$
s_1^m \le (n-k)^{m-1} s_m
$$

for all $m \in \mathbb{N}$ *. In particular, if* $A \in \mathcal{C}_n$ *then*

$$
s_1^m \le (n-1)^{m-1} s_m
$$

for all $m \in \mathbb{N}$ *.*

PROOF. Since \overline{A} is a nonnegative matrix, we have

$$
s_m = \text{tr}(A^m) \ge \sum_{i=1}^n a_{ii}^m = \sum_{i \in J} a_{ii}^m,
$$

where $J = \{i \in \{1, 2, ..., n\} : a_{ii} > 0\}$. On the other hand, Hölder's inequality gives

$$
s_1^m = \left(\sum_{i \in J} a_{ii}\right)^m \le (n-k)^{m-1} \sum_{i \in J} a_{ii}^m,
$$

and so we conclude that $s_1^m \le (n-k)^{m-1}$ s_m .

Using Proposition [2.2](#page-2-0) we prove the following lower estimates for the spread of a matrix in \mathcal{C}_n .

Theorem 2.3. *If* $A \in \mathcal{C}_n$ *then*

$$
s(A) > \frac{2}{4 + \sqrt{2(n+3)}}
$$

for $n \geq 6$ *,*

$$
s(A) \ge \frac{5}{8 + \sqrt{74}}
$$

for $n = 5$, and

$$
s(A) \ge \frac{1}{3}
$$

for $n = 4$ *.*

PROOF. Since $s(A) > 0$ by Proposition [2.1](#page-1-0) and since the result is true if $s(A) \geq 1$, we may assume that $s := s(A) \in (0,1)$, and consequently the eigenvalues of A have positive real parts. Let $\lambda_1 = r(A) = 1, \lambda_2, \lambda_3, \ldots, \lambda_n$ be the spectrum of A. By Proposition [2.2,](#page-2-0) we have

$$
\left(\sum_{i=1}^{n} \lambda_i\right)^2 = s_1^2 \le (n-1)s_2 = (n-1)\sum_{i=1}^{n} \lambda_i^2.
$$

This inequality can be rewritten in the form

$$
\sum_{i=1}^{n} \lambda_i^2 \le \sum_{i=1}^{n-1} \sum_{j=i+1}^{n} (\lambda_i - \lambda_j)^2.
$$
 (1)

The right-hand side of [\(1\)](#page-3-0) is clearly at most $n(n-1)s^2/2$. To obtain a lower bound for the left-hand side of [\(1\)](#page-3-0), we choose any eigenvalue λ of A. Since $\lambda + \overline{\lambda} = 2 \operatorname{Re} \lambda \ge 2(1 - s) > 0$, we have

$$
\lambda^{2} + \overline{\lambda}^{2} = (\lambda + \overline{\lambda})^{2} - 2|\lambda|^{2} \ge (2(1 - s))^{2} - 2 = 4s^{2} - 8s + 2,
$$

and so we obtain the following lower bound for the left-hand side of [\(1\)](#page-3-0):

$$
\sum_{i=1}^{n} \lambda_i^2 = 1 + \sum_{i=2}^{n} \lambda_i^2 \ge 1 + \frac{n-1}{2} (4s^2 - 8s + 2).
$$

Therefore, the inequality [\(1\)](#page-3-0) gives the inequality

$$
\frac{n(n-1)}{2}s^2 \ge 1 + \frac{n-1}{2}(4s^2 - 8s + 2),
$$

which leads to the inequality

$$
(n-1)(n-4)s2 + 8(n-1)s - 2n \ge 0.
$$
 (2)

For $n=4$ we obtain that $s\geq \frac{1}{3}$ $\frac{1}{3}$, while for $n = 5$ we have

$$
2s^2 + 16s - 5 \ge 0,
$$

implying that

$$
s \ge \frac{-8 + \sqrt{74}}{2} = \frac{5}{8 + \sqrt{74}}.
$$

If $n\geq 6$ we rewrite the inequality [\(2\)](#page-4-0) to the form

$$
(n2 - 5n)s2 + 8ns - 2n \ge -4s2 + 8s = 4s(2 - s) > 0,
$$

and so

$$
(n-5)s^2 + 8s - 2 > 0.
$$

It follows that

$$
s > \frac{-4 + \sqrt{2(n+3)}}{n-5} = \frac{2}{4 + \sqrt{2(n+3)}}.
$$

This completes the proof.

For $n \in \{2,3\}$ we can obtain sharp lower bounds for the spread of a matrix in \mathcal{C}_n .

Proposition 2.4. *If* $A \in \mathcal{C}_2$ *then* $s(A) \geq 1$ *; if* $A \in \mathcal{C}_3$ *then* $s(A) \geq \frac{3}{4}$ 4 *. Both bounds are exact.*

PROOF. Let 1 and λ be the eigenvalues of $A \in \mathcal{C}_2$. By Proposition [2.2,](#page-2-0) we have

$$
(1 + \lambda)^2 = s_1^2 \le s_2 = 1 + \lambda^2,
$$

and so $\lambda \leq 0$ proving that $s(A) \geq 1$. The diagonal matrix diag $(0,1) \in C_2$ shows that this lower bound is exact.

In the case $n = 3$ we first suppose that a matrix $A \in \mathcal{C}_3$ has real eigenvalues 1, λ and μ . We may assume that $0 \leq \lambda \leq \mu \leq 1$. Then the inequality [\(1\)](#page-3-0) gives the inequality

$$
1 + \lambda^2 + \mu^2 \le (1 - \lambda)^2 + (1 - \mu)^2 + (\lambda - \mu)^2,
$$

and so

$$
2\lambda^2 \le 2\lambda\mu \le (1-\lambda)^2 + (1-\mu)^2 - 1 \le 2(1-\lambda)^2 - 1 = 2\lambda^2 - 4\lambda + 1.
$$

It follows that $\lambda \leq \frac{1}{4}$ $\frac{1}{4}$, so that $s(A) \geq \frac{3}{4}$ $\frac{3}{4}$.

Assume now that a matrix $A \in C_3$ has eigenvalues 1, $\lambda = a + ib$ and $\overline{\lambda} = a - ib$, where $a \in \mathbb{R}$ and $b > 0$. By Proposition [2.2,](#page-2-0) we have

$$
(1+2a)^2 = s_1^2 \le 2s_2 = 2(1+\lambda^2+\overline{\lambda}^2) = 2+4a^2-4b^2 \le 2+4a^2,
$$

and so $a \leq \frac{1}{4}$ $\frac{1}{4}$. This implies that $s(A) \geq \frac{3}{4}$ $\frac{3}{4}$ as asserted.

The exactness of this lower bound is proved by the matrix

$$
A = \frac{1}{4} \begin{bmatrix} 0 & 2 & 0 \\ 0 & 3 & 1 \\ 2 & 0 & 3 \end{bmatrix} \in \mathcal{C}_3
$$

the spectrum of which is $\{1, \frac{1}{4}\}$ $\frac{1}{4}$, $\frac{1}{4}$ 4 }.

For $n \geq 4$ it looks difficult to obtain exact lower bounds for the spread of matrices in \mathcal{C}_n . We thus restrict our attention to a special subset of \mathcal{C}_n . Proposition [2.1](#page-1-0) trivially implies that every matrix in \mathcal{C}_n has at least two distinct eigenvalues, that is, 1 is not the only point in its spectrum. Let \mathcal{D}_n (with $n \geq 2$) be the collection of all matrices in \mathcal{C}_n having exactly two distinct eigenvalues. We now prove sharp lower bounds for the spread of matrices in \mathcal{D}_n .

Theorem 2.5. *If* $A \in \mathcal{D}_n$ *then*

$$
s(A) \ge \frac{n}{2(n-1)}
$$

Moreover, this bound is the best possible, i.e., there is a (necessarily irreducible) matrix $A \in \mathcal{D}_n$ such that $s(A) = \frac{n}{2(n-1)}$.

PROOF. Assume first that a matrix $A \in \mathcal{D}_n$ is irreducible. Then 1 is a simple eigenvalue of A by the Perron-Frobenius theorem. Therefore, A also has an eigenvalue $\lambda \in (-1, 1)$ of multiplicity $n - 1$. In this case the inequality [\(1\)](#page-3-0) reads as follows:

$$
1 + (n - 1)\lambda^2 \le (n - 1)(1 - \lambda)^2.
$$

Simplifying it, we obtain

$$
\lambda \le \frac{n-2}{2(n-1)}.
$$

This implies that

$$
s(A) = 1 - \lambda \ge \frac{n}{2(n-1)}.
$$

Assume now that a matrix $A \in \mathcal{D}_n$ is reducible. Then, up to similarity with a permutation matrix, we may assume that

$$
A = \begin{bmatrix} A_{11} & A_{12} & A_{13} & \dots & A_{1m} \\ 0 & A_{22} & A_{23} & \dots & A_{2m} \\ 0 & 0 & A_{33} & \dots & A_{3m} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & A_{mm} \end{bmatrix}
$$

where each of A_{11} , A_{22} , ..., A_{mm} is either an irreducible (square) matrix or a 1×1 block. Let A_{kk} be one of these diagonal blocks that has a zero diagonal element. Without loss of generality we may assume that $s(A) < 1$, so that 0 is not in the spectrum of A implying that all 1×1 diagonal blocks are non-zero. Therefore, if A_{kk} is an $r \times r$ matrix, then $r \geq 2$, and so

$$
s(A) \ge s(A_{kk}) \ge \frac{r}{2(r-1)} > \frac{n}{2(n-1)}.
$$

This completes the proof of the first assertion of the theorem.

To show that the lower bound can be achieved, we define the matrix $A = [a_{i,j}]_{i,j=1}^n$ with nonzero elements: $a_{i,i+1} = n - i$ for $i = 1, 2, \ldots, n - 1$, $a_{i,i} = n$ for $i = 2, 3, \ldots, n$, and $a_{i,j} = 2$ if $i - j$ is an even positive integer. We also introduce the upper triangular matrix $U = [u_{i,j}]_{i,j=1}^n$ with nonzero elements: $u_{i,i+1} = n-i$ for $i = 1, 2, ..., n-1$, $u_{1,1} = 2(n-1)$ and $u_{i,i} = n-2$ for $i = 2, 3, \ldots, n$. For example, if $n = 5$ then

$$
A = \begin{bmatrix} 0 & 4 & 0 & 0 & 0 \\ 0 & 5 & 3 & 0 & 0 \\ 2 & 0 & 5 & 2 & 0 \\ 0 & 2 & 0 & 5 & 1 \\ 2 & 0 & 2 & 0 & 5 \end{bmatrix} \text{ and } U = \begin{bmatrix} 8 & 4 & 0 & 0 & 0 \\ 0 & 3 & 3 & 0 & 0 \\ 0 & 0 & 3 & 2 & 0 \\ 0 & 0 & 0 & 3 & 1 \\ 0 & 0 & 0 & 0 & 3 \end{bmatrix}
$$

.

The proof is complete if we show that A and U are similar matrices, because then we have $r(A) = 2(n-1)$, $s(A) = n$, and $\frac{1}{2(n-1)}A \in \mathcal{D}_n$. Define two nilpotent matrices

$$
N = \begin{bmatrix} 0 & 0 & 0 & 0 & \dots & 0 & 0 \\ 1 & 0 & 0 & 0 & \dots & 0 & 0 \\ 0 & 1 & 0 & 0 & \dots & 0 & 0 \\ 0 & 0 & 1 & 0 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & \dots & 0 & 0 \\ 0 & 0 & 0 & 0 & \dots & 1 & 0 \end{bmatrix}
$$

and

$$
M = \begin{bmatrix} 0 & n-1 & 0 & 0 & \dots & 0 & 0 \\ 0 & 0 & n-2 & 0 & \dots & 0 & 0 \\ 0 & 0 & 0 & n-3 & \dots & 0 & 0 \\ 0 & 0 & 0 & 0 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & \dots & 0 & 1 \\ 0 & 0 & 0 & 0 & \dots & 0 & 0 \end{bmatrix}.
$$

Introduce also the matrix

$$
S = (I + N)(I - N)^{-1} = (I + N)(I + N + N^2 + N^3 + \dots + N^{n-1}) =
$$

= $I + 2N + 2N^2 + 2N^3 + 2N^4 + \dots + 2N^{n-1} =$

$$
\begin{bmatrix} 1 & 0 & 0 & \dots & 0 & 0 \\ 2 & 1 & 0 & 0 & \dots & 0 & 0 \\ 2 & 2 & 1 & 0 & \dots & 0 & 0 \\ 2 & 2 & 2 & 1 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 2 & 2 & 2 & 2 & \dots & 1 & 0 \\ 2 & 2 & 2 & 2 & \dots & 2 & 1 \end{bmatrix}
$$

Let e_1, \ldots, e_n be the standard basis vectors, and let $e = e_1 + \ldots + e_n =$ $(1, 1, \ldots, 1)^T$. Observe that

$$
A = M + nI - ne_1e_1^T + 2(N^2 + N^4 + N^6 + ...) =
$$

= $M + (n - 2)I - ne_1e_1^T + 2(I - N^2)^{-1}$

and

$$
U = M + (n - 2)I + n e_1 e_1^T.
$$

Note also that $[N, M] := NM - MN = I - ne_1e_1^T$. By induction one can verify that $[N^k, M] = kN^{k-1} - ne_k e_1^T$ for $k = 1, 2, ..., n$. Then the commutator of S and M is

$$
[S,M] = 2\sum_{k=1}^{n} [N^k, M] = 2\sum_{k=1}^{n} kN^{k-1} - 2n\sum_{k=1}^{n} e_k e_1^T = 2(I-N)^{-2} - 2nee_1^T.
$$

Now we have

$$
SU - AS = [S, M] + n(Se_1)e_1^T + ne_1e_1^TS - 2(I - N^2)^{-1}S =
$$

= 2(I - N)⁻² - 2nee₁^T + n(2e - e₁)e₁^T + ne₁e₁^T - 2(I - N²)⁻¹(I + N)(I - N)⁻¹ =
= 2(I - N)⁻² - 2(I - N)⁻² = 0.

This proves that the matrices A and U are similar. \square

Acknowledgments.

The author was supported in part by the Slovenian Research Agency. He would like to thank Thomas Laffey and Helena Smigoc for pointing out that Proposition [2.2](#page-2-0) holds.

References

- [1] C.R. Johnson, Row stochastic matrices similar to doubly stochastic matrices, Linear and Multilinear Algebra 10 (1981), 113–130.
- [2] R. Loewy, D. London, A note on an inverse eigenvalue problem for nonnegative matrices, Linear and Multilinear Algebra 6 (1978/79), 83– 90.
- [3] J. K. Merikoski, R. Kumar, Characterizations and lower bounds for the spread of a normal matrix, Linear Algebra Appl. 364 (2003), 13–31.
- [4] L. Mirsky, The spread of a matrix, Mathematika 3 (1956), 127–130.