

# On the extremal values of the number of vertices with an interval spectrum on the set of proper edge colorings of the graph of the $n$ -dimensional cube

A.M. Khachatryan<sup>1</sup>, R.R. Kamalian<sup>2</sup>

<sup>1</sup>Ijevan Branch of Yerevan State University, e-mail: khachatryanarpine@gmail.com

<sup>2</sup>The Institute for Informatics and Automation Problems of NAS RA,  
e-mail: rrkamalian@yahoo.com

## Abstract

For an undirected, simple, finite, connected graph  $G$ , we denote by  $V(G)$  and  $E(G)$  the sets of its vertices and edges, respectively. A function  $\varphi : E(G) \rightarrow \{1, \dots, t\}$  is called a proper edge  $t$ -coloring of a graph  $G$ , if adjacent edges are colored differently and each of  $t$  colors is used. The least value of  $t$  for which there exists a proper edge  $t$ -coloring of a graph  $G$  is denoted by  $\chi'(G)$ . For any graph  $G$ , and for any integer  $t$  satisfying the inequality  $\chi'(G) \leq t \leq |E(G)|$ , we denote by  $\alpha(G, t)$  the set of all proper edge  $t$ -colorings of  $G$ . Let us also define a set  $\alpha(G)$  of all proper edge colorings of a graph  $G$ :

$$\alpha(G) \equiv \bigcup_{t=\chi'(G)}^{|E(G)|} \alpha(G, t).$$

An arbitrary nonempty finite subset of consecutive integers is called an interval. If  $\varphi \in \alpha(G)$  and  $x \in V(G)$ , then the set of colors of edges of  $G$  which are incident with  $x$  is denoted by  $S_G(x, \varphi)$  and is called a spectrum of the vertex  $x$  of the graph  $G$  at the proper edge coloring  $\varphi$ . If  $G$  is a graph and  $\varphi \in \alpha(G)$ , then define  $f_G(\varphi) \equiv |\{x \in V(G) / S_G(x, \varphi) \text{ is an interval}\}|$ .

For a graph  $G$  and any integer  $t$ , satisfying the inequality  $\chi'(G) \leq t \leq |E(G)|$ , we define:

$$\mu_1(G, t) \equiv \min_{\varphi \in \alpha(G, t)} f_G(\varphi), \quad \mu_2(G, t) \equiv \max_{\varphi \in \alpha(G, t)} f_G(\varphi).$$

For any graph  $G$ , we set:

$$\begin{aligned} \mu_{11}(G) &\equiv \min_{\chi'(G) \leq t \leq |E(G)|} \mu_1(G, t), & \mu_{12}(G) &\equiv \max_{\chi'(G) \leq t \leq |E(G)|} \mu_1(G, t), \\ \mu_{21}(G) &\equiv \min_{\chi'(G) \leq t \leq |E(G)|} \mu_2(G, t), & \mu_{22}(G) &\equiv \max_{\chi'(G) \leq t \leq |E(G)|} \mu_2(G, t). \end{aligned}$$

For any positive integer  $n$ , the exact values of the parameters  $\mu_{11}$ ,  $\mu_{12}$ ,  $\mu_{21}$  and  $\mu_{22}$  are found for the graph of the  $n$ -dimensional cube.

Keywords:  $n$ -dimensional cube, proper edge coloring, interval spectrum, game.

Math. Classification: 05C15

We consider finite, undirected, connected graphs without loops and multiple edges containing at least one edge. For any graph  $G$ , we denote by  $V(G)$  and  $E(G)$  the sets of vertices and edges of  $G$ , respectively. For any  $x \in V(G)$ ,  $d_G(x)$  denotes the degree of the vertex  $x$  in  $G$ . For a graph  $G$ ,  $\delta(G)$  and  $\Delta(G)$  denote the minimum and maximum degrees of vertices in  $G$ , respectively. For a graph  $G$ , and for any  $V_0 \subseteq V(G)$ , we denote by  $G[V_0]$  the subgraph of the graph  $G$  induced [1] by the subset  $V_0$  of its vertices. For arbitrary graphs  $G$  and  $H$ ,  $G \square H$  denotes their cartesian product [1].

For any positive integer  $n$ , we denote by  $Q_n$  the graph of the  $n$ -dimensional cube [2]. Clearly, for any positive integer  $n$ ,  $Q_n$  is a regular bipartite graph with  $|V(Q_n)| = 2^n$ ,  $|E(Q_n)| = n \cdot 2^{n-1}$ ,  $\Delta(Q_n) = n$ .

An arbitrary nonempty finite subset of consecutive integers is called an interval. An interval with the minimum element  $p$  and the maximum element  $q$  is denoted by  $[p, q]$ .

A function  $\varphi : E(G) \rightarrow [1, t]$  is called a proper edge  $t$ -coloring of a graph  $G$ , if each of  $t$  colors is used, and adjacent edges are colored differently.

The minimum value of  $t$  for which there exists a proper edge  $t$ -coloring of a graph  $G$  is denoted by  $\chi'(G)$  [3].

For any graph  $G$ , and for any  $t \in [\chi'(G), |E(G)|]$ , we denote by  $\alpha(G, t)$  the set of all proper edge  $t$ -colorings of  $G$ .

Let us also define a set  $\alpha(G)$  of all proper edge colorings of a graph  $G$ :

$$\alpha(G) \equiv \bigcup_{t=\chi'(G)}^{|E(G)|} \alpha(G, t).$$

If  $\varphi \in \alpha(G)$  and  $x \in V(G)$ , then the set  $\{\varphi(e)/e \in E(G), e \text{ is incident with } x\}$  is called a spectrum of the vertex  $x$  of the graph  $G$  at the proper edge coloring  $\varphi$  and is denoted by  $S_G(x, \varphi)$ ; if  $S_G(x, \varphi)$  is an interval, we say that  $\varphi$  is interval in  $x$ .

If  $G$  is a graph,  $\varphi \in \alpha(G)$ ,  $R \subseteq V(G)$ , then we say, that  $\varphi$  is interval on  $R$  iff for  $\forall x \in R$ ,  $\varphi$  is interval in  $x$ . We say, that a subset  $R$  of vertices of a graph  $G$  has an  $i$ -property iff there exists  $\varphi \in \alpha(G)$  interval on  $R$ . If  $G$  is a graph, and a subset  $R$  of its vertices has an  $i$ -property, we denote by  $w_R(G)$  and  $W_R(G)$  (omiting the index in these notations in the peculiar case of  $R = V(G)$ ) the minimum and the maximum value of  $t$ , respectively, for which  $\exists \varphi \in \alpha(G, t)$  interval on  $R$  [4, 5, 6] (see also [20]).

If  $G$  is a graph,  $\varphi \in \alpha(G)$ , then set  $V_{int}(G, \varphi) \equiv \{x \in V(G)/S_G(x, \varphi) \text{ is an interval}\}$  and  $f_G(\varphi) \equiv |V_{int}(G, \varphi)|$ . A proper edge coloring  $\varphi \in \alpha(G)$  is called an interval edge coloring [4, 5, 6] of the graph  $G$  iff  $f_G(\varphi) = |V(G)|$ . The set of all graphs having an interval edge coloring is denoted by  $\mathfrak{N}$ . The simplest example of the graph which doesn't belong to  $\mathfrak{N}$  is  $K_3$ . The terms and concepts which are not defined can be found in [1].

For a graph  $G$ , and for any  $t \in [\chi'(G), |E(G)|]$ , we set [7]:

$$\mu_1(G, t) \equiv \min_{\varphi \in \alpha(G, t)} f_G(\varphi), \quad \mu_2(G, t) \equiv \max_{\varphi \in \alpha(G, t)} f_G(\varphi).$$

For any graph  $G$ , we set [7]:

$$\begin{aligned} \mu_{11}(G) &\equiv \min_{\chi'(G) \leq t \leq |E(G)|} \mu_1(G, t), & \mu_{12}(G) &\equiv \max_{\chi'(G) \leq t \leq |E(G)|} \mu_1(G, t), \\ \mu_{21}(G) &\equiv \min_{\chi'(G) \leq t \leq |E(G)|} \mu_2(G, t), & \mu_{22}(G) &\equiv \max_{\chi'(G) \leq t \leq |E(G)|} \mu_2(G, t). \end{aligned}$$

Clearly, the parameters  $\mu_{11}$ ,  $\mu_{12}$ ,  $\mu_{21}$  and  $\mu_{22}$  are correctly defined for an arbitrary graph.

Let us note that exact values of the parameters  $\mu_{12}$  and  $\mu_{21}$  have certain game interpretations. Suppose that all edges of a graph  $G$  are colored in the game of Alice and Bob with antagonistic interests and asymmetric distribution of roles. Alice determines the number  $t$  of colors in the future proper edge coloring  $\varphi$  of the graph  $G$ , satisfying the condition  $t \in [\chi'(G), |E(G)|]$ , Bob colors edges of  $G$  with  $t$  colors.

When Alice aspires to maximize, Bob aspires to minimize the value of the function  $f_G(\varphi)$ , and both players choose their best strategies, then at the finish of the game exactly  $\mu_{12}(G)$  vertices of  $G$  will receive an interval spectrum.

When Alice aspires to minimize, Bob aspires to maximize the value of the function  $f_G(\varphi)$ , and both players choose their best strategies, then at the finish of the game exactly  $\mu_{21}(G)$  vertices of  $G$  will receive an interval spectrum.

The exact values of the parameters  $\mu_{11}$ ,  $\mu_{12}$ ,  $\mu_{21}$  and  $\mu_{22}$  are found for simple paths, simple cycles and simple cycles with a chord [8, 9], "Möbius ladders" [7, 22], complete graphs [11], complete bipartite graphs [12, 13], prisms [14, 22],  $n$ -dimensional cubes [14, 15] and the Petersen graph [23]. The exact values of  $\mu_{11}$  and  $\mu_{22}$  for trees are found in [16]. The exact value of  $\mu_{12}$  for an arbitrary tree is found in [17] (see also [18, 19]).

In this paper, for any positive integer  $n$ , we determine the exact values of the parameters  $\mu_{11}$ ,  $\mu_{12}$ ,  $\mu_{21}$  and  $\mu_{22}$  for the graph  $Q_n$ .

First we recall some known results.

**Lemma 1.** [4, 5, 6] *If  $G \in \mathfrak{N}$ , then  $\chi'(G) = \Delta(G)$ .*

**Corollary 1.** *If  $G \in \mathfrak{N}$ , then  $\Delta(G) = \chi'(G) \leq w(G) \leq W(G) \leq |E(G)|$ .*

**Corollary 2.** *If  $G$  is a regular graph, then  $G \in \mathfrak{N}$  iff  $\chi'(G) = \Delta(G)$ .*

**Corollary 3.** *If  $G$  is a regular graph with  $\chi'(G) = \Delta(G)$ , then  $G \in \mathfrak{N}$  and  $w(G) = \chi'(G) = \Delta(G)$ .*

**Corollary 4.** *If  $G$  is a regular bipartite graph, then  $G \in \mathfrak{N}$  and  $w(G) = \chi'(G) = \Delta(G)$ .*

**Corollary 5.** *If  $G \in \mathfrak{N}$ , and a subset  $R \subseteq V(G)$  has an  $i$ -property, then  $\Delta(G) = \chi'(G) \leq w_R(G) \leq w(G) \leq W(G) \leq W_R(G) \leq |E(G)|$ .*

**Lemma 2.** [4, 5, 6] *If  $R$  is the set of all vertices of an arbitrary part of a bipartite graph  $G$ , then:*

- 1)  $R$  has an  $i$ -property,
- 2)  $W_R(G) = |E(G)|$ ,
- 3) for any  $t \in [w_R(G), W_R(G)]$ , there exists  $\varphi_t \in \alpha(G, t)$  interval on  $R$ .

**Corollary 6.** *If  $R$  is the set of all vertices of an arbitrary part of a regular bipartite graph  $G$ , then:*

- 1)  $R$  has an  $i$ -property,
- 2)  $W_R(G) = |E(G)|$ ,
- 3)  $w_R(G) = \Delta(G)$ ,
- 4) for any  $t \in [w_R(G), W_R(G)]$ , there exists  $\varphi_t \in \alpha(G, t)$  interval on  $R$ .

**Lemma 3.** [21] If  $G$  is a graph with  $\delta(G) \geq 2$ ,  $\varphi \in \alpha(G, |E(G)|)$ ,  $V_{int}(G, \varphi) \neq \emptyset$ , then  $G[V_{int}(G, \varphi)]$  is a forest each connected component of which is a simple path.

**Lemma 4.** [8, 9] If  $G$  is a regular graph with  $\chi'(G) = \Delta(G)$ , then  $\mu_{12}(G) = |V(G)|$ .

**Proof.** Let us note that for an arbitrary  $\varphi \in \alpha(G, \Delta(G))$ ,  $f_G(\varphi) = |V(G)|$ . Consequently,  $|V(G)| = \mu_1(G, \Delta(G)) \leq \max_{t \in [\chi'(G), |E(G)|]} \mu_1(G, t) = \mu_{12}(G) \leq |V(G)|$ . Hence,  $\mu_{12}(G) = |V(G)|$ .

**The Lemma is proved.**

**Lemma 5.** [8, 9] For an arbitrary graph  $G$ , the following inequalities hold:  $\mu_{11}(G) \leq \mu_{12}(G) \leq \mu_{22}(G)$ ,  $\mu_{11}(G) \leq \mu_{21}(G) \leq \mu_{22}(G)$ .

**Proof.** Clearly,

$$\mu_{11}(G) = \min_{\chi'(G) \leq t \leq |E(G)|} \mu_1(G, t) \leq \max_{\chi'(G) \leq t \leq |E(G)|} \mu_1(G, t) = \mu_{12}(G).$$

and

$$\mu_{21}(G) = \min_{\chi'(G) \leq t \leq |E(G)|} \mu_2(G, t) \leq \max_{\chi'(G) \leq t \leq |E(G)|} \mu_2(G, t) = \mu_{22}(G).$$

The inequality  $\mu_{12}(G) \leq \mu_{22}(G)$  is provided by the existence of an integer  $t' \in [\chi'(G), |E(G)|]$ , for which the relation

$$\begin{aligned} \mu_{12}(G) &= \max_{\chi'(G) \leq t \leq |E(G)|} \mu_1(G, t) = \mu_1(G, t') = \min_{\varphi \in \alpha(G, t')} f_G(\varphi) \leq \\ &\leq \max_{\varphi \in \alpha(G, t')} f_G(\varphi) = \mu_2(G, t') \leq \max_{\chi'(G) \leq t \leq |E(G)|} \mu_2(G, t) = \mu_{22}(G) \end{aligned}$$

is true.

The inequality  $\mu_{11}(G) \leq \mu_{21}(G)$  is provided by the existence of an integer  $t'' \in [\chi'(G), |E(G)|]$ , for which the relation

$$\begin{aligned} \mu_{21}(G) &= \min_{\chi'(G) \leq t \leq |E(G)|} \mu_2(G, t) = \mu_2(G, t'') = \max_{\varphi \in \alpha(G, t'')} f_G(\varphi) \geq \\ &\geq \min_{\varphi \in \alpha(G, t'')} f_G(\varphi) = \mu_1(G, t'') \geq \min_{\chi'(G) \leq t \leq |E(G)|} \mu_1(G, t) = \mu_{11}(G) \end{aligned}$$

is true.

**The Lemma is proved.**

**Remark 1.** [8, 9] There are graphs  $G$  with  $\mu_{21}(G) < \mu_{12}(G)$ . There are graphs  $G$  with  $\mu_{12}(G) < \mu_{21}(G)$ .

**Lemma 6.** [10, 24] If a subset  $V_0$  of the set of vertices of the graph  $Q_3$  contains at least 6 vertices, then at least one of the following two statements is true:

- 1) there exist such vertices  $x_1, x_2, x_3, x_4$  in  $V_0$  that  $Q_3[\{x_1, x_2, x_3, x_4\}] \cong K_{3,1}$ ,
- 2) there exist such vertices  $y_1, y_2, y_3, y_4, y_5, y_6$  in  $V_0$  that  $Q_3[\{y_1, y_2, y_3, y_4, y_5, y_6\}] \cong C_6$ .

**Lemma 7.** [10, 24] If a subset  $V_0$  of the set of vertices of the graph  $Q_n$  ( $n \geq 4$ ) contains at least  $2^{n-1} + 1$  vertices, then at least one of the following two statements is true:

- 1) there exist such vertices  $x_1, x_2, x_3, x_4$  in  $V_0$  that  $Q_n[\{x_1, x_2, x_3, x_4\}] \cong K_{3,1}$ ,

2) there exist such vertices  $y_1, y_2, y_3, y_4, y_5, y_6, y_7, y_8$  in  $V_0$  that  $Q_n[\{y_1, y_2, y_3, y_4, y_5, y_6, y_7, y_8\}] \cong C_8$ .

If  $G$  is a graph with  $\chi'(G) = \Delta(G)$ ,  $t \in [\Delta(G), |E(G)|]$ ,  $\xi \in \alpha(G, t)$ , then for any  $j \in [1, t]$ , we denote by  $E(G, \xi, j)$  the set of all edges of  $G$  colored by the color  $j$  at the proper edge  $t$ -coloring  $\xi$ ;  $\xi$  is called a harmonic [12] edge  $t$ -coloring of the graph  $G$ , if for any  $i \in [1, \Delta(G)]$ , the set

$$\bigcup_{1 \leq j \leq t, j \equiv i \pmod{\Delta(G)}} E(G, \xi, j)$$

is a matching in  $G$ .

Suppose that  $G$  is a graph with  $\chi'(G) = \Delta(G)$ ,  $t \in [\Delta(G), |E(G)|]$ ,  $\xi$  is a harmonic edge  $t$ -coloring of  $G$ . Let us define a sequence  $\xi_0^*, \xi_1^*, \dots, \xi_{t-\chi'(G)}^*$  of proper edge colorings of the graph  $G$ .

Set  $\xi_0^* \equiv \xi$ .

**Case 1.**  $t = \chi'(G)$ . The desired sequence mentioned above is already constructed.

**Case 2.**  $t \in [\chi'(G) + 1, |E(G)|]$ . Suppose that  $j \in [1, t - \chi'(G)]$ , and the proper edge colorings  $\xi_0^*, \dots, \xi_{j-1}^*$  of the graph  $G$  are already constructed. Let us define  $\xi_j^*$ . For an arbitrary  $e \in E(G)$ , set:

$$\xi_j^*(e) \equiv \begin{cases} \xi_{j-1}^*(e), & \text{if } \xi_{j-1}^*(e) \neq \max(\{\xi_{j-1}^*(e)/e \in E(G)\}) \\ \xi_{j-1}^*(e) - \Delta(G), & \text{if } \xi_{j-1}^*(e) = \max(\{\xi_{j-1}^*(e)/e \in E(G)\}). \end{cases}$$

**Remark 2.** Suppose that  $G$  is a graph with  $\chi'(G) = \Delta(G)$ ,  $t \in [\Delta(G), |E(G)|]$ ,  $\xi$  is a harmonic edge  $t$ -coloring of  $G$ . All proper edge colorings  $\xi_0^*, \xi_1^*, \dots, \xi_{t-\chi'(G)}^*$  of the graph  $G$  are determined definitely.

**Remark 3.** Suppose that  $G$  is a graph with  $\chi'(G) = \Delta(G) < |E(G)|$ ,  $t \in [1 + \chi'(G), |E(G)|]$ ,  $\xi$  is a harmonic edge  $t$ -coloring of  $G$ . It is not difficult to see, that for any  $j \in [1, t - \chi'(G)]$ ,  $\xi_j^*$  is a harmonic edge  $(t - j)$ -coloring of the graph  $G$ .

**Remark 4.** Suppose that  $G$  is a graph with  $\chi'(G) = \Delta(G) < |E(G)|$ ,  $t \in [1 + \chi'(G), |E(G)|]$ ,  $\xi$  is a harmonic edge  $t$ -coloring of  $G$ . Assume that  $\xi$  is interval in some vertex  $z_0 \in V(G)$  with  $d_G(z_0) = \Delta(G)$ . Then, for any  $j \in [1, t - \chi'(G)]$ ,  $\xi_j^*$  is interval in  $z_0$ .

From results of [8, 9] we have

**Proposition 1.**  $\mu_{11}(Q_1) = 2$ ,  $\mu_{11}(Q_2) = 1$ ,  $\mu_{21}(Q_1) = 2$ ,  $\mu_{21}(Q_2) = 3$ .

**Lemma 8.** If an  $r$ -regular ( $r \geq 3$ ) graph  $G$  has  $\varphi \in \alpha(G, r + 1)$  with  $f_G(\varphi) = 0$ , then the graph  $G \square K_2$  has  $\psi \in \alpha(G \square K_2, r + 2)$  with  $f_{G \square K_2}(\psi) = 0$ .

**Proof** is evident.

**Lemma 9.** There exists  $\varphi \in \alpha(Q_3, 4)$  with  $f_{Q_3}(\varphi) = 0$ . There exists  $\psi \in \alpha(Q_3, 12)$  with  $f_{Q_3}(\psi) = 5$ .

**Proof.** Let  $V(Q_3) = \{x_1, x_2, x_3, x_4, y_1, y_2, y_3, y_4\}$ ,  $E(Q_3) = \{(x_1, x_2), (x_2, x_3), (x_3, x_4), (x_1, x_4), (y_1, y_2), (y_2, y_3), (y_3, y_4), (y_1, y_4), (x_1, y_1), (x_2, y_2), (x_3, y_3), (x_4, y_4)\}$ .

Set:  $\varphi((x_1, x_2)) = \varphi((x_3, x_4)) = \varphi((y_1, y_2)) = \varphi((y_3, y_4)) = 1$ ,  $\varphi((x_1, x_4)) = \varphi((x_2, x_3)) = 2$ ,  $\varphi((y_1, y_4)) = \varphi((y_2, y_3)) = 3$ ,  $\varphi((x_1, y_1)) = \varphi((x_2, y_2)) = \varphi((x_3, y_3)) = \varphi((x_4, y_4)) = 4$ .

It is not difficult to see that  $\varphi \in \alpha(Q_3, 4)$  and  $f_{Q_3}(\varphi) = 0$ .

Set:  $\psi((x_1, y_1)) = 1$ ,  $\psi((y_1, y_2)) = 2$ ,  $\psi((y_1, y_4)) = 3$ ,  $\psi((y_3, y_4)) = 4$ ,  $\psi((x_4, y_4)) = 5$ ,  $\psi((x_1, x_4)) = 6$ ,  $\psi((x_3, x_4)) = 7$ ,  $\psi((x_3, y_3)) = 8$ ,  $\psi((x_2, x_3)) = 9$ ,  $\psi((x_2, y_2)) = 10$ ,  $\psi((x_1, x_2)) = 11$ ,  $\psi((y_2, y_3)) = 12$ .

It is not difficult to see that  $\psi \in \alpha(Q_3, 12)$  and  $f_{Q_3}(\psi) = 5$ .

**The Lemma is proved.**

From Lemmas 8 and 9 we obtain

**Proposition 2.** *For any integer  $n \geq 3$ ,  $\mu_{11}(Q_n) = 0$ .*

From Propositions 1 and 2 we obtain

**Proposition 3.** *For any positive integer  $n$ ,  $\mu_{11}(Q_n) = 3 - \min\{3, n\}$ .*

From Lemma 4 we obtain

**Proposition 4.** *For any positive integer  $n$ ,  $\mu_{12}(Q_n) = 2^n$ .*

From Proposition 4 and Lemma 5 we obtain

**Proposition 5.** *For any positive integer  $n$ ,  $\mu_{22}(Q_n) = 2^n$ .*

**Lemma 10.** *For any integer  $n \geq 4$ ,  $\mu_2(Q_n, n \cdot 2^{n-1}) \geq 2^{n-1}$ .*

**Proof** follows from Lemma 2.

**Lemma 11.** *For any integer  $n \geq 4$ ,  $\mu_2(Q_n, n \cdot 2^{n-1}) \leq 2^{n-1}$ .*

**Proof.** Assume the contrary. Then there exist an integer  $n_0 \geq 4$  and  $\varphi_0 \in \alpha(Q_{n_0}, n_0 \cdot 2^{n_0-1})$ , for which  $f_{Q_{n_0}}(\varphi_0) \geq 2^{n_0-1} + 1$ . By Lemma 3,  $Q_{n_0}[V_{int}(Q_{n_0}, \varphi_0)]$  is a forest, each connected component of which is a simple path. But it is incompatible with Lemma 7.

**The Lemma is proved.**

From Lemmas 10 and 11 we obtain

**Lemma 12.** *For any integer  $n \geq 4$ ,  $\mu_2(Q_n, n \cdot 2^{n-1}) = 2^{n-1}$ .*

**Lemma 13.** *For arbitrary integers  $n$  and  $t$ , satisfying the conditions  $n \geq 4$ ,  $n \leq t \leq n \cdot 2^{n-1}$ , the inequality  $\mu_2(Q_n, t) \geq 2^{n-1}$  is true.*

**Proof.** Choose an arbitrary integer  $n_0 \geq 4$ . It follows from Corollary 6, that for any integer  $t$ , satisfying the inequality  $n_0 \leq t \leq n_0 \cdot 2^{n_0-1}$ , there exists  $\varphi_t \in \alpha(Q_{n_0}, t)$  with  $f_{Q_{n_0}}(\varphi_t) \geq 2^{n_0-1}$ . It means that for any integer  $t$ , satisfying the inequality  $n_0 \leq t \leq n_0 \cdot 2^{n_0-1}$ , we also have  $\mu_2(Q_{n_0}, t) \geq 2^{n_0-1}$ .

**The Lemma is proved.**

From Lemmas 12 and 13 we obtain

**Proposition 6.** *For any integer  $n \geq 4$ ,  $\mu_{21}(Q_n) = 2^{n-1}$ .*

From Lemma 9 we obtain

**Lemma 14.**  $\mu_2(Q_3, 12) \geq 5$ .

**Lemma 15.**  $\mu_2(Q_3, 12) \leq 5$ .

**Proof.** Assume the contrary:  $\mu_2(Q_3, 12) \geq 6$ . It means that there exists  $\xi \in \alpha(Q_3, 12)$  with  $f_{Q_3}(\xi) \geq 6$ . By Lemma 3,  $Q_3[V_{int}(Q_3, \xi)]$  is a forest each connected component of which is a simple path. But it is incompatible with Lemma 6.

**The Lemma is proved.**

From Lemmas 14 and 15 we obtain

**Lemma 16.**  $\mu_2(Q_3, 12) = 5$ .

**Lemma 17.** For any integer  $t$ , satisfying the condition  $3 \leq t \leq 12$ , the inequality  $\mu_2(Q_3, t) \geq 5$  is true.

**Proof.** It is not difficult to check that the proper edge 12-coloring  $\psi$  of the graph  $Q_3$  constructed for the proof of Lemma 9 is a harmonic edge 12-coloring of  $Q_3$ . From Remark 4 it follows that for any integer  $t$  satisfying the inequality  $3 \leq t \leq 11$ , there exists  $\psi_t \in \alpha(Q_3, t)$  with  $V_{int}(Q_3, \psi) \subseteq V_{int}(Q_3, \psi_t)$ . Hence, for the same values of  $t$ , the inequality  $\mu_2(Q_3, t) \geq f_{Q_3}(\psi) = 5$  is true.

**The Lemma is proved.**

From Lemmas 16 and 17 we obtain

**Proposition 7.**  $\mu_{21}(Q_3) = 5$ .

From Propositions 1, 6 and 7 we obtain

**Proposition 8.** For any positive integer  $n$ ,

$$\mu_{21}(Q_n) = 2^{n-1} + \left\lceil 1 - \frac{\min\{4, n\}}{4} \right\rceil.$$

From Propositions 3 - 5 and 8 we obtain

**Theorem 1.** [14, 15] For any positive integer  $n$ , the following equalities hold:

$$\mu_{11}(Q_n) = 3 - \min\{3, n\}, \quad \mu_{12}(Q_n) = \mu_{22}(Q_n) = 2^n,$$

$$\mu_{21}(Q_n) = 2^{n-1} + \left\lceil 1 - \frac{\min\{4, n\}}{4} \right\rceil.$$

## References

- [1] D.B. West, *Introduction to Graph Theory*, Prentice-Hall, New Jersey, 1996.
- [2] F.R.K. Chung, Z. Füredi, R.L. Graham, P. Seymour, *On induced subgraphs of the cube*, Journal of Comb. Theory, Series A, 49 (1988), pp. 180–187.
- [3] V.G. Vizing, *The chromatic index of a multigraph*, Kibernetika 3 (1965), pp. 29–39.
- [4] A.S. Asratian, R.R. Kamalian, *Interval colorings of edges of a multigraph*, Appl. Math. 5 (1987), Yerevan State University, pp. 25–34 (in Russian).
- [5] A.S. Asratian, R.R. Kamalian, *Investigation of interval edge-colorings of graphs*, Journal of Combinatorial Theory. Series B 62 (1994), no.1, pp. 34–43.

- [6] R.R. Kamalian, *Interval Edge Colorings of Graphs*, Doctoral dissertation, the Institute of Mathematics of the Siberian Branch of the Academy of Sciences of USSR, Novosibirsk, 1990 (in Russian).
- [7] N.N. Davtyan, R.R. Kamalian, *On boundaries of extremums of the number of vertices with an interval spectrum among the set of proper edge colorings of "Möbius ladders" with  $t$  colors under variation of  $t$* , Proc. of the 3<sup>th</sup> Ann. Sci. Conf. (December 5–10, 2008) of the RAU, Yerevan, 2009, pp. 81–84 (in Russian).
- [8] N.N. Davtyan, R.R. Kamalian, *On properties of the number of vertices with an interval spectrum in proper edge colorings of some graphs*, the Herald of the RAU, №2, Yerevan, 2009, pp. 33–42.
- [9] N.N. Davtyan, R.R. Kamalian, *Some properties of the number of vertices with an interval spectrum in proper edge colorings of graphs*, The Collection "Akunq" of Scientific Papers of Ijevan Branch of Yerevan State University, Yerevan, 2012, pp. 18–27.
- [10] R.R. Kamalian, A.M. Khachatryan, *On a property of the  $n$ -dimensional cube*, Mathematical Problems of Computer Science, Vol. 38, Yerevan, 2012, pp. 66–67.
- [11] A.M. Khachatryan, *On boundaries of extremums of the number of vertices with an interval spectrum among the set of proper edge colorings of complete graphs with  $t$  colors under variation of  $t$* , Proc. of the 5<sup>th</sup> Ann. Sci. Conf. (December 6–10, 2010) of the RAU, Yerevan, 2011, pp. 268–272 (in Russian).
- [12] A.M. Khachatryan, *On the parameters  $\mu_{11}$ ,  $\mu_{12}$  and  $\mu_{22}$  of complete bipartite graphs*, the Herald of the RAU, №1, Yerevan, 2011, pp. 76–83 (in Russian).
- [13] R.R. Kamalian, A.M. Khachatryan, *On the sharp value of the parameter  $\mu_{21}$  of complete bipartite graphs*, the Herald of the RAU, №2, Yerevan, 2011, pp. 19–25 (in Russian).
- [14] R.R. Kamalian, A.M. Khachatryan, *On properties of a number of vertices with an interval spectrum among the set of proper edge colorings of some regular graphs*, Proc. of the 6<sup>th</sup> Ann. Sci. Conf. (December 5–9, 2011) of the RAU, Yerevan, 2012, pp. 62–65 (in Russian).
- [15] A.M. Khachatryan, R.R. Kamalian, *On the  $\mu$ -parameters of the graph of the  $n$ -dimensional cube*, Book of abstracts of the International Mathematical Conference on occasion to the 70th year anniversary of Professor Vladimir Kirichenko, June 13-19 (2012), Mykolaiv, Ukraine, pp. 38–39.
- [16] N.N. Davtyan, *On the least and the greatest possible numbers of vertices with an interval spectrum on the set of proper edge colorings of a tree*, Math. Problems of Computer Science, Vol. 32, Yerevan, 2009, pp. 107–111 (in Russian).
- [17] N.N. Davtyan, R.R. Kamalian, *On the parameter  $\mu_{12}$  of a tree*, Proc. of the 4<sup>th</sup> Ann. Sci. Conf. (November 30 – December 4, 2009) of the RAU, Yerevan, 2010, pp. 149–151 (in Russian).
- [18] N.N. Davtyan, R.R. Kamalian, *On an algorithm of evaluation of the exact value of the parameter  $\mu_{12}$  of an arbitrary tree*, the Herald of the RAU, №1, Yerevan, 2011, pp. 57–63 (in Russian).
- [19] N.N. Davtyan, *On a property of the parameter  $\mu_{12}$  of trees of special kind*, the Herald of the RAU, №2, Yerevan, 2010, pp. 77–82 (in Russian).



- [20] R.R. Kamalian, *On one-sided interval colorings of bipartite graphs*, the Herald of the RAU, №2, Yerevan, 2010, pp. 3–11 (in Russian).
- [21] N.N. Davtyan, A.M. Khachatryan, R.R. Kamalian, *On a subgraph induced at a labeling of a graph by the subset of vertices with an interval spectrum*, Book of abstracts of the 8<sup>th</sup> International Algebraic Conference in Ukraine. July 5–12(2011), Lugansk, pp. 61–62.
- [22] N.N. Davtyan, A.M. Khachatryan, R.R. Kamalian, *On Boundaries of Extrema of the Number of Vertices with an Interval Spectrum on the Sets of Proper Edge  $t$ -colorings of Some Cubic Graphs under Variation of  $t$* , International Mathematical Forum, Vol. 8, 2013, no. 24, pp. 1195 - 1198, <http://dx.doi.org/10.12988/imf.2013.3491>.
- [23] N.N. Davtyan, *Private information* [26.06.2013]
- [24] R.R. Kamalian, A.M. Khachatryan, *On a property of the  $n$ -dimensional cube*, Intellectual Archive Bulletin, Abstracts and the descriptions of works in Art and Science submitted to [www.IntellectualArchive.com](http://www.IntellectualArchive.com), Toronto, July, 2012.