

Symmetric Reduction and Hamilton-Jacobi Equation of Rigid Spacecraft with a Rotor

Hong Wang

School of Mathematical Sciences and LPMC,
Nankai University, Tianjin 300071, P.R.China

E-mail: hongwang@nankai.edu.cn

August 17, 2013

Abstract. In this paper, we consider the rigid spacecraft with an internal rotor as a regular point reducible regular controlled Hamiltonian (RCH) system. In the cases of coincident and non-coincident centers of buoyancy and gravity, we give explicitly the motion equation and Hamilton-Jacobi equation of reduced spacecraft-rotor system on a symplectic leaf by calculation in detail, respectively, which show the effect on controls in regular symplectic reduction and Hamilton-Jacobi theory.

Keywords: rigid spacecraft with a rotor, regular controlled Hamiltonian system, non-coincident center, regular point reduction, Hamilton-Jacobi equation.

AMS Classification: 70H33, 70H20, 70Q05.

Contents

1	Introduction	1
2	The Rigid Spacecraft with a Rotor	4
2.1	The Spacecraft-Rotor System with Coincident Centers	4
2.2	The Spacecraft-Rotor System with Non-coincident Centers	5
3	Symmetric Reduction of the Rigid Spacecraft with a Rotor	6
3.1	Symmetric Reduction of Spacecraft-Rotor System with Coincident Centers . . .	6
3.2	Symmetric Reduction of Spacecraft-Rotor System with Non-coincident Centers .	8
4	Hamilton-Jacobi Equation of the Rigid Spacecraft with a Rotor	12
4.1	H-J Equation of Spacecraft-Rotor System with Coincident Centers	13
4.2	H-J Equation of Spacecraft-Rotor System with Non-coincident Centers	14

1 Introduction

It is well-known that the theory of controlled mechanical systems has formed an important subject in recent years. Its research gathers together some separate areas of research such as mechanics, differential geometry and nonlinear control theory, etc., and the emphasis of this research on geometry is motivated by the aim of understanding the structure of equations of

motion of the system in a way that helps both analysis and design. There is a natural two-fold method in the study of controlled mechanical systems. First, it is a special class of nonlinear control systems considered, and one can study it by using the feedback control and optimal control method. The second, it is also a special class of mechanical systems, one can study it combining with the analysis of dynamic systems and the geometric reduction theory of Hamiltonian and Lagrangian systems. Thus, the theory of controlled mechanical systems presents a challenging and promising research area between the study of classical mechanics and that of modern nonlinear geometric control theory, and a lot of researchers are absorbed to pour into the area and there have been a lot of interesting results. Such as, Bloch et al in [4–7], studied the symmetry and control as well as referred to the use of feedback control to realize a modification to the structure of a given mechanical system; Nijmeijer and Van der Schaft in [25], studied the nonlinear dynamical control systems as well as the use of feedback control to stabilize mechanical systems, and Van der Schaft in [29, 30] referred to the reduction and control of implicit (port) Hamiltonian systems.

In particular, we note that in Marsden et al [22], the authors studied regular reduction theory of controlled Hamiltonian systems with symplectic structure and symmetry, as an extension of regular symplectic reduction theory of Hamiltonian systems under regular controlled Hamiltonian equivalence conditions. Wang in [31] generalized the work in [22] to study the singular reduction theory of regular controlled Hamiltonian systems, and Wang and Zhang in [34] generalized the work in [22] to study optimal reduction theory of controlled Hamiltonian systems with Poisson structure and symmetry by using optimal momentum map and reduced Poisson tensor (or reduced symplectic form), and Ratiu and Wang in [28] studied the Poisson reduction of controlled Hamiltonian system by controllability distribution. These research work not only gave a variety of reduction methods for controlled Hamiltonian systems, but also showed a variety of relationships of controlled Hamiltonian equivalence of these systems.

At the same time, we note also that Hamilton-Jacobi theory is an important part of classical mechanics. On the one hand, it provides a characterization of the generating functions of certain time-dependent canonical transformations, such that a given Hamiltonian system in such a form that its solutions are extremely easy to find by reduction to the equilibrium, see Abraham and Marsden [1], Arnold [2] and Marsden and Ratiu [21]. On the other hand, it is possible in many cases that Hamilton-Jacobi theory provides an immediate way to integrate the equation of motion of system, even when the problem of Hamiltonian system itself has not been or cannot be solved completely. In addition, the Hamilton-Jacobi equation is also fundamental in the study of the quantum-classical relationship in quantization, and it also plays an important role in the development of numerical integrators that preserve the symplectic structure and in the study of stochastic dynamical systems, see Woodhouse [35], Ge and Marsden [10], Marsden and West [23] and Lázaro-Camí and Ortega [13]. For these reasons it is described as a useful tools in the study of Hamiltonian system theory, and has been extensively developed in past many years and become one of the most active subjects in the study of modern applied mathematics and analytical mechanics, which absorbed a lot of researchers to pour into it and a lot of deep and beautiful results have been obtained, see Cariñena et al [8] and [9], Iglesias et al [11], León et al [3, 14, 15] Ohsawa and Bloch [26] for more details.

Just as we have known that Hamilton-Jacobi theory from the variational point of view is originally developed by Jacobi in 1866, which state that the integral of Lagrangian of a system along the solution of its Euler-Lagrange equation satisfies the Hamilton-Jacobi equation. The

classical description of this problem from the geometrical point of view is given by Abraham and Marsden in [1], and it is developed in the context of time-dependent Hamiltonian system by Marsden and Ratiu in [21]. The Hamilton-Jacobi equation may be regarded as a nonlinear partial differential equation for some generating function S , and the problem is become how to choose a time-dependent canonical transformation $\Psi : T^*Q \times \mathbb{R} \rightarrow T^*Q \times \mathbb{R}$, which transforms the dynamical vector field of a time-dependent Hamiltonian system to equilibrium, such that the generating function S of Ψ satisfies the time-dependent Hamilton-Jacobi equation. In particular, for the time-independent Hamiltonian system, we may look for a symplectic map as the canonical transformation. This work offers a important idea that one can use the dynamical vector field of a Hamiltonian system to describe Hamilton-Jacobi equation. Moreover, assume that $\gamma : Q \rightarrow T^*Q$ is a closed one-form on smooth manifold Q , and define that $X_H^\gamma = T\pi_Q \cdot X_H \cdot \gamma$, where X_H is the dynamical vector field of Hamiltonian system (T^*Q, ω, H) . Then the fact that X_H^γ and X_H are γ -related, that is, $T\gamma \cdot X_H^\gamma = X_H \cdot \gamma$ is equivalent that $\mathbf{d}(H \cdot \gamma) = 0$, which is given in Cariñena et al [8] and Iglesias et al [11]. Since the Hamilton-Jacobi theory is developed based on the Hamiltonian picture of dynamics, motivated by the above research work, Wang in [32] uses the dynamical vector field of Hamiltonian system and the regular reduced Hamiltonian system to describe the Hamilton-Jacobi theory for these systems. Moreover, Wang in [33] extends the Hamilton-Jacobi theory to the regular controlled Hamiltonian system and its regular reduced systems, and describes the relationship between the RCH-equivalence for RCH systems and the solutions of corresponding Hamilton-Jacobi equations.

Now, it is a natural problem if there is a practical controlled Hamiltonian system and how to show the effect on controls in regular symplectic reduction and Hamilton-Jacobi theory of the system. In this paper, as an application of the regular point symplectic reduction and Hamilton-Jacobi theory of RCH system with symmetry, we consider that rigid spacecraft with an internal rotor is a regular point reducible RCH system given in Marsden et al [22], where the rigid spacecraft with an internal rotor is modeled as a Hamiltonian system with control, which is given in Bloch and Leonard [5], Bloch, Leonard and Marsden [7]. In the cases of coincident and non-coincident centers of buoyancy and gravity, we give explicitly the motion equation and Hamilton-Jacobi equation of reduced spacecraft-rotor system on a symplectic leaf by calculation in detail, respectively. These equations are more complex than that of Hamiltonian system without control and describe explicitly the effect on controls in regular symplectic reduction and Hamilton-Jacobi theory.

A brief of outline of this paper is as follows. In the second section, we first review some relevant definitions and basic facts about rigid spacecraft with an internal rotor, which will be used in subsequent sections. As an application of the theoretical result of symplectic reduction of RCH system given by Marsden et al [22], in the third section we regard the rigid spacecraft with an internal rotor as a regular point reducible RCH system on the generalization of rotation group $SO(3) \times S^1$ and on the generalization of Euclidean group $SE(3) \times S^1$, respectively, and in the cases of coincident and non-coincident centers of buoyancy and gravity, we give explicitly the motion equations of their reduced RCH systems on the symplectic leaves by calculation in detail. Moreover, as an application of the theoretical result of Hamilton-Jacobi theory of regular reduced RCH system given by Wang [33], in the fourth section, we give the Hamilton-Jacobi equations of the reduced rigid spacecraft-rotor systems on the symplectic leaves by calculation in detail, respectively, in the cases of coincident and non-coincident centers of buoyancy and gravity. These research work develop the application of symplectic reduction and Hamilton-Jacobi theory of RCH systems with symmetry and make us have much deeper understanding

and recognition for the structure of Hamiltonian systems and RCH systems.

2 The Rigid Spacecraft with a Rotor

In this paper, our goal is to give the regular point reduction and Hamilton-Jacobi theorem of rigid spacecraft with an internal rotor. In order to do these, in this section, we first review some relevant definitions and basic facts about rigid spacecraft with an internal rotor, which will be used in subsequent sections. We shall follow the notations and conventions introduced in Bloch and Leonard [5], Bloch, Leonard and Marsden [7], Marsden [18], Marsden and Ratiu [21], and Marsden et al [22]. In this paper, we assume that all manifolds are real, smooth and finite dimensional and all actions are smooth left actions. For convenience, we also assume that all controls appearing in this paper are the admissible controls.

2.1 The Spacecraft-Rotor System with Coincident Centers

We consider a rigid spacecraft (to be called the carrier body) carrying an internal rotor, and assume that the only external forces and torques acting on the spacecraft-rotor system are due to buoyancy and gravity. In general, it is possible that the spacecraft's center of buoyancy may not be coincident with its center of gravity. But, in this subsection we assume that the spacecraft is symmetric and to have uniformly distributed mass, and the center of buoyancy and the center of gravity are coincident. Denote by O the center of mass of the system in the body frame and at O place a set of (orthonormal) body axes. Assume that the body coordinate axes are aligned with principal axes of the carrier body, and the rotor is aligned along the third principal axis, see Bloch and Leonard [5] and Bloch, Leonard and Marsden [7]. The rotor spins under the influence of a torque u acting on the rotor. If translations are ignored and only rotations are considered, in this case, then the configuration space is $Q = \text{SO}(3) \times S^1$, with the first factor being the attitude of rigid spacecraft and the second factor being the angle of rotor. The corresponding phase space is the cotangent bundle $T^*Q = T^*\text{SO}(3) \times T^*S^1$, where $T^*S^1 \cong T^*\mathbb{R} \cong \mathbb{R} \times \mathbb{R}$, with the canonical symplectic form.

Let $I = \text{diag}(I_1, I_2, I_3)$ be the moment of inertia of the carrier body in the principal body-fixed frame, and J_3 be the moment of inertia of rotor around its rotation axis. Let J_{3k} , $k = 1, 2$, be the moments of inertia of the rotor around the k th principal axis with $k = 1, 2$, and denote by $\bar{I}_k = I_k + J_{3k}$, $k = 1, 2$, $\bar{I}_3 = I_3$. Let $\Omega = (\Omega_1, \Omega_2, \Omega_3)$ be the vector of body angular velocities computed with respect to the axes fixed in the body and $(\Omega_1, \Omega_2, \Omega_3) \in \mathfrak{so}(3)$. Let α be the relative angle of rotor and $\dot{\alpha}$ the rotor relative angular velocity about the third principal axis with respect to a carrier body fixed frame. Consider the Lagrangian of the system $L(A, \Omega, \alpha, \dot{\alpha}) : TQ \cong \text{SO}(3) \times \mathfrak{so}(3) \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$, which is the total kinetic energy of the rigid spacecraft plus the kinetic energy of rotor, given by

$$L(A, \Omega, \alpha, \dot{\alpha}) = \frac{1}{2}[\bar{I}_1\Omega_1^2 + \bar{I}_2\Omega_2^2 + \bar{I}_3\Omega_3^2 + J_3(\Omega_3 + \dot{\alpha})^2],$$

where $A \in \text{SO}(3)$, $\Omega = (\Omega_1, \Omega_2, \Omega_3) \in \mathfrak{so}(3)$, $\alpha \in \mathbb{R}$, $\dot{\alpha} \in \mathbb{R}$. If we introduce the conjugate angular momentum, given by $\Pi_k = \frac{\partial L}{\partial \Omega_k} = \bar{I}_k\Omega_k$, $k = 1, 2$, $\Pi_3 = \frac{\partial L}{\partial \Omega_3} = \bar{I}_3\Omega_3 + J_3(\Omega_3 + \dot{\alpha})$, $l = \frac{\partial L}{\partial \dot{\alpha}} = J_3(\Omega_3 + \dot{\alpha})$, and by the Legendre transformation $FL : \text{SO}(3) \times \mathfrak{so}(3) \times \mathbb{R} \times \mathbb{R} \rightarrow \text{SO}(3) \times \mathfrak{so}^*(3) \times \mathbb{R} \times \mathbb{R}$, $(A, \Omega, \alpha, \dot{\alpha}) \rightarrow (A, \Pi, \alpha, l)$, where $\Pi = (\Pi_1, \Pi_2, \Pi_3) \in \mathfrak{so}^*(3)$, $l \in \mathbb{R}$,

we have the Hamiltonian $H(A, \Pi, \alpha, l) : T^*Q \cong \text{SO}(3) \times \mathfrak{so}^*(3) \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ given by

$$\begin{aligned} H(A, \Pi, \alpha, l) &= \Omega \cdot \Pi + \dot{\alpha} \cdot l - L(A, \Omega, \alpha, \dot{\alpha}) \\ &= \frac{1}{2} \left[\frac{\Pi_1^2}{\bar{I}_1} + \frac{\Pi_2^2}{\bar{I}_2} + \frac{(\Pi_3 - l)^2}{\bar{I}_3} + \frac{l^2}{J_3} \right]. \end{aligned} \quad (2.1)$$

In order to give the motion equation of spacecraft-rotor system, in the following we need to consider the symmetry and reduced symplectic structure of the configuration space $Q = \text{SO}(3) \times S^1$.

2.2 The Spacecraft-Rotor System with Non-coincident Centers

Since it is possible that the spacecraft's center of buoyancy may not be coincident with its center of gravity, in this subsection then we consider the spacecraft-rotor system with non-coincident centers of buoyancy and gravity. We fix an orthonormal coordinate frame to the carrier body with origin located at the center of buoyancy and axes aligned with the principal axes of the carrier body, and the rotor is aligned along the third principal axis, see Bloch and Leonard [5], Bloch, Leonard and Marsden [7], and Leonard and Marsden [16]. The rotor spins under the influence of a torque u acting on the rotor. When the carrier body is oriented so that the body-fixed frame is aligned with the inertial frame, the third principal axis aligns with the direction of gravity. The vector from the center of buoyancy to the center of gravity with respect to the body-fixed frame is $h\chi$, where χ is an unit vector on the line connecting the two centers which is assumed to be aligned along the third principal axis, and h is the length of this segment. The mass of the carrier body is denoted m and the magnitude of gravitational acceleration is g , and let Γ be the unit vector viewed by an observer moving with the body. In this case, the configuration space is $Q = \text{SO}(3) \circledast \mathbb{R}^3 \times S^1 \cong \text{SE}(3) \times S^1$, with the first factor being the attitude of rigid spacecraft and the drift of spacecraft in the rotational process and the second factor being the angle of rotor. The corresponding phase space is the cotangent bundle $T^*Q = T^*\text{SE}(3) \times T^*S^1$, where $T^*S^1 \cong T^*\mathbb{R} \cong \mathbb{R} \times \mathbb{R}$, with the canonical symplectic form.

Consider the Lagrangian of the system $L(A, c, \Omega, \Gamma, \alpha, \dot{\alpha}) : TQ \cong \text{SE}(3) \times \mathfrak{se}(3) \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$, which is the total kinetic energy of the rigid spacecraft plus the kinetic energy of rotor minus potential energy of the system, given by

$$L(A, c, \Omega, \Gamma, \alpha, \dot{\alpha}) = \frac{1}{2} [\bar{I}_1 \Omega_1^2 + \bar{I}_2 \Omega_2^2 + \bar{I}_3 \Omega_3^2 + J_3 (\Omega_3 + \dot{\alpha})^2] - mgh\Gamma \cdot \chi,$$

where $(A, c) \in \text{SE}(3)$, $\Omega = (\Omega_1, \Omega_2, \Omega_3) \in \mathfrak{so}(3)$, $\Gamma \in \mathbb{R}^3$, $(\Omega, \Gamma) \in \mathfrak{se}(3)$, $\alpha \in \mathbb{R}$, $\dot{\alpha} \in \mathbb{R}$.

If we introduce the conjugate angular momentum, given by $\Pi_k = \frac{\partial L}{\partial \Omega_k} = \bar{I}_k \Omega_k$, $k = 1, 2$, $\Pi_3 = \frac{\partial L}{\partial \Omega_3} = \bar{I}_3 \Omega_3 + J_3 (\Omega_3 + \dot{\alpha})$, $l = \frac{\partial L}{\partial \dot{\alpha}} = J_3 (\Omega_3 + \dot{\alpha})$, and by the Legendre transformation

$$FL : \text{SE}(3) \times \mathfrak{se}(3) \times \mathbb{R} \times \mathbb{R} \rightarrow \text{SE}(3) \times \mathfrak{se}^*(3) \times \mathbb{R} \times \mathbb{R}, \quad (A, c, \Omega, \Gamma, \alpha, \dot{\alpha}) \rightarrow (A, c, \Pi, \Gamma, \alpha, l),$$

where $\Pi = (\Pi_1, \Pi_2, \Pi_3) \in \mathfrak{so}^*(3)$, $(\Pi, \Gamma) \in \mathfrak{se}^*(3)$, $l \in \mathbb{R}$, we have the Hamiltonian $H(A, c, \Pi, \Gamma, \alpha, l) : T^*Q \cong \text{SE}(3) \times \mathfrak{se}^*(3) \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ given by

$$\begin{aligned} H(A, c, \Pi, \Gamma, \alpha, l) &= \Omega \cdot \Pi + \dot{\alpha} \cdot l - L(A, c, \Omega, \Gamma, \alpha, \dot{\alpha}) \\ &= \frac{1}{2} \left[\frac{\Pi_1^2}{\bar{I}_1} + \frac{\Pi_2^2}{\bar{I}_2} + \frac{(\Pi_3 - l)^2}{\bar{I}_3} + \frac{l^2}{J_3} \right] + mgh\Gamma \cdot \chi. \end{aligned} \quad (2.2)$$

In order to give the motion equation of spacecraft-rotor system, in the following we need to consider the symmetry and reduced symplectic structure of the configuration space $Q = \text{SE}(3) \times S^1$.

3 Symmetric Reduction of the Rigid Spacecraft with a Rotor

In the following we regard the rigid spacecraft with an internal rotor as a regular point reducible RCH system on the generalization of rotation group $\text{SO}(3) \times S^1$ and on the generalization of Euclidean group $\text{SE}(3) \times S^1$, respectively, and give the motion equations of their reduced RCH systems on the symplectic leaves by calculation in detail. It is worthy note that it is different from the symmetric reduction of Hamiltonian system in Bloch and Leonard [5], Bloch, Leonard and Marsden [7], Marsden [18], the reductions in this paper are all the controlled Hamiltonian reductions, that is, the symmetric reductions of (regular) controlled Hamiltonian systems, see Marsden et al [22]. We also follow the notations and conventions introduced in Marsden et al [19, 20], Marsden and Ratiu [21], Libermann and Marle [17], Ortega and Ratiu [27].

3.1 Symmetric Reduction of Spacecraft-Rotor System with Coincident Centers

We first give the regular point reduction of spacecraft-rotor system with coincident centers of buoyancy and gravity. Assume that Lie group $G = \text{SO}(3)$ acts freely and properly on $Q = \text{SO}(3) \times S^1$ by the left translation on the first factor $\text{SO}(3)$, and the trivial action on the second factor S^1 . By using the left trivialization of $T^*\text{SO}(3) = \text{SO}(3) \times \mathfrak{so}^*(3)$, then the action of $\text{SO}(3)$ on phase space $T^*Q = T^*\text{SO}(3) \times T^*S^1$ is by cotangent lift of left translation on $\text{SO}(3)$ at the identity, that is, $\Phi : \text{SO}(3) \times T^*\text{SO}(3) \times T^*S^1 \cong \text{SO}(3) \times \text{SO}(3) \times \mathfrak{so}^*(3) \times \mathbb{R} \times \mathbb{R} \rightarrow \text{SO}(3) \times \mathfrak{so}^*(3) \times \mathbb{R} \times \mathbb{R}$, given by $\Phi(B, (A, \Pi, \alpha, l)) = (BA, \Pi, \alpha, l)$, for any $A, B \in \text{SO}(3)$, $\Pi \in \mathfrak{so}^*(3)$, $\alpha, l \in \mathbb{R}$, which is also free and proper, and the orbit space $(T^*Q)/\text{SO}(3)$ is a smooth manifold and $\pi : T^*Q \rightarrow (T^*Q)/\text{SO}(3)$ is a smooth submersion. Since $\text{SO}(3)$ acts trivially on $\mathfrak{so}^*(3)$ and \mathbb{R} , it follows that $(T^*Q)/\text{SO}(3)$ is diffeomorphic to $\mathfrak{so}^*(3) \times \mathbb{R} \times \mathbb{R}$.

We know that $\mathfrak{so}^*(3)$ is a Poisson manifold with respect to its rigid body Lie-Poisson bracket defined by

$$\{F, K\}_{\mathfrak{so}^*(3)}(\Pi) = -\Pi \cdot (\nabla_{\Pi} F \times \nabla_{\Pi} K), \quad \forall F, K \in C^{\infty}(\mathfrak{so}^*(3)), \quad \Pi \in \mathfrak{so}^*(3). \quad (3.1)$$

For $\mu \in \mathfrak{so}^*(3)$, the coadjoint orbit $\mathcal{O}_{\mu} \subset \mathfrak{so}^*(3)$ has the induced orbit symplectic form $\omega_{\mathcal{O}_{\mu}}^{-}$, which is coincide with the restriction of the Lie-Poisson bracket on $\mathfrak{so}^*(3)$ to the coadjoint orbit \mathcal{O}_{μ} . From the Symplectic Stratification theorem we know that the coadjoint orbits $(\mathcal{O}_{\mu}, \omega_{\mathcal{O}_{\mu}}^{-})$, $\mu \in \mathfrak{so}^*(3)$, form the symplectic leaves of the Poisson manifold $(\mathfrak{so}^*(3), \{\cdot, \cdot\}_{\mathfrak{so}^*(3)})$. Let $\omega_{\mathbb{R}}$ be the canonical symplectic form on $T^*\mathbb{R} \cong \mathbb{R} \times \mathbb{R}$, which is given by

$$\omega_{\mathbb{R}}((\theta_1, \lambda_1), (\theta_2, \lambda_2)) = \langle \lambda_2, \theta_1 \rangle - \langle \lambda_1, \theta_2 \rangle, \quad (3.2)$$

where $(\theta_i, \lambda_i) \in \mathbb{R} \times \mathbb{R}$, $i = 1, 2$, $\langle \cdot, \cdot \rangle$ is the standard inner product on $\mathbb{R} \times \mathbb{R}$. It induces a canonical Poisson bracket $\{\cdot, \cdot\}_{\mathbb{R}}$ on $T^*\mathbb{R}$, which is given by

$$\{F, K\}_{\mathbb{R}}(\theta, \lambda) = \frac{\partial F}{\partial \theta} \frac{\partial K}{\partial \lambda} - \frac{\partial K}{\partial \theta} \frac{\partial F}{\partial \lambda}. \quad (3.3)$$

Thus, we can induce a symplectic form $\tilde{\omega}_{\mathcal{O}_{\mu} \times \mathbb{R} \times \mathbb{R}}^{-} = \pi_{\mathcal{O}_{\mu}}^* \omega_{\mathcal{O}_{\mu}}^{-} + \pi_{\mathbb{R}}^* \omega_{\mathbb{R}}$ on the smooth manifold $\mathcal{O}_{\mu} \times \mathbb{R} \times \mathbb{R}$, where the maps $\pi_{\mathcal{O}_{\mu}} : \mathcal{O}_{\mu} \times \mathbb{R} \times \mathbb{R} \rightarrow \mathcal{O}_{\mu}$ and $\pi_{\mathbb{R}} : \mathcal{O}_{\mu} \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R} \times \mathbb{R}$ are

canonical projections, and induce a Poisson bracket $\{\cdot, \cdot\}_- = \pi_{\mathfrak{so}^*(3)}^* \{\cdot, \cdot\}_{\mathfrak{so}^*(3)} + \pi_{\mathbb{R}}^* \{\cdot, \cdot\}_{\mathbb{R}}$ on the smooth manifold $\mathfrak{so}^*(3) \times \mathbb{R} \times \mathbb{R}$, where the maps $\pi_{\mathfrak{so}^*(3)} : \mathfrak{so}^*(3) \times \mathbb{R} \times \mathbb{R} \rightarrow \mathfrak{so}^*(3)$ and $\pi_{\mathbb{R}} : \mathfrak{so}^*(3) \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R} \times \mathbb{R}$ are canonical projections, and such that $(\mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R}, \tilde{\omega}_{\mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R}}^-)$ is a symplectic leaf of the Poisson manifold $(\mathfrak{so}^*(3) \times \mathbb{R} \times \mathbb{R}, \{\cdot, \cdot\}_-)$.

On the other hand, from $T^*Q = T^*\text{SO}(3) \times T^*S^1$ we know that there is a canonical symplectic form $\omega_Q = \pi_{\text{SO}(3)}^* \omega_0 + \pi_{S^1}^* \omega_{S^1}$ on T^*Q , where ω_0 is the canonical symplectic form on $T^*\text{SO}(3)$ and the maps $\pi_{\text{SO}(3)} : Q = \text{SO}(3) \times S^1 \rightarrow \text{SO}(3)$ and $\pi_{S^1} : Q = \text{SO}(3) \times S^1 \rightarrow S^1$ are canonical projections. Then the cotangent lift of left $\text{SO}(3)$ -action $\Phi : \text{SO}(3) \times T^*Q \rightarrow T^*Q$ is also symplectic, and admits an associated Ad^* -equivariant momentum map $\mathbf{J}_Q : T^*Q \rightarrow \mathfrak{so}^*(3)$ such that $\mathbf{J}_Q \cdot \pi_{\text{SO}(3)}^* = \mathbf{J}_{\text{SO}(3)}$, where $\mathbf{J}_{\text{SO}(3)} : T^*\text{SO}(3) \rightarrow \mathfrak{so}^*(3)$ is a momentum map of left $\text{SO}(3)$ -action on $T^*\text{SO}(3)$, and $\pi_{\text{SO}(3)}^* : T^*\text{SO}(3) \rightarrow T^*Q$. If $\mu \in \mathfrak{so}^*(3)$ is a regular value of \mathbf{J}_Q , then $\mu \in \mathfrak{so}^*(3)$ is also a regular value of $\mathbf{J}_{\text{SO}(3)}$ and $\mathbf{J}_Q^{-1}(\mu) \cong \mathbf{J}_{\text{SO}(3)}^{-1}(\mu) \times \mathbb{R} \times \mathbb{R}$. Denote by $\text{SO}(3)_\mu = \{g \in \text{SO}(3) \mid \text{Ad}_g^* \mu = \mu\}$ the isotropy subgroup of coadjoint $\text{SO}(3)$ -action at the point $\mu \in \mathfrak{so}^*(3)$. It follows that $\text{SO}(3)_\mu$ acts also freely and properly on $\mathbf{J}_Q^{-1}(\mu)$, the regular point reduced space $(T^*Q)_\mu = \mathbf{J}_Q^{-1}(\mu) / \text{SO}(3)_\mu \cong (T^*\text{SO}(3))_\mu \times \mathbb{R} \times \mathbb{R}$ of (T^*Q, ω_Q) at μ , is a symplectic manifold with symplectic form ω_μ uniquely characterized by the relation $\pi_\mu^* \omega_\mu = i_\mu^* \omega_Q = i_\mu^* \pi_{\text{SO}(3)}^* \omega_0 + i_\mu^* \pi_{S^1}^* \omega_{S^1}$, where the map $i_\mu : \mathbf{J}_Q^{-1}(\mu) \rightarrow T^*Q$ is the inclusion and $\pi_\mu : \mathbf{J}_Q^{-1}(\mu) \rightarrow (T^*Q)_\mu$ is the projection. Because from Abraham and Marsden [1], we know that $((T^*\text{SO}(3))_\mu, \omega_\mu)$ is symplectically diffeomorphic to $(\mathcal{O}_\mu, \omega_{\mathcal{O}_\mu}^-)$, and hence we have that $((T^*Q)_\mu, \omega_\mu)$ is symplectically diffeomorphic to $(\mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R}, \tilde{\omega}_{\mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R}}^-)$, which is a symplectic leaf of the Poisson manifold $(\mathfrak{so}^*(3) \times \mathbb{R} \times \mathbb{R}, \{\cdot, \cdot\}_-)$.

From the expression (2.1) of the Hamiltonian, we know that $H(A, \Pi, \alpha, l)$ is invariant under the left $\text{SO}(3)$ -action $\Phi : \text{SO}(3) \times T^*Q \rightarrow T^*Q$. For the case $\mu \in \mathfrak{so}^*(3)$ is the regular value of \mathbf{J}_Q , we have the reduced Hamiltonian $h_\mu(\Pi, \alpha, l) : \mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R} (\subset \mathfrak{so}^*(3) \times \mathbb{R} \times \mathbb{R}) \rightarrow \mathbb{R}$ given by $h_\mu(\Pi, \alpha, l) = \pi_\mu(H(A, \Pi, \alpha, l)) = H(A, \Pi, \alpha, l)|_{\mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R}}$. From the rigid body Poisson bracket on $\mathfrak{so}^*(3)$ and the Poisson bracket on $T^*\mathbb{R}$, we can get the Poisson bracket on $\mathfrak{so}^*(3) \times \mathbb{R} \times \mathbb{R}$, that is, for $F, K : \mathfrak{so}^*(3) \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$, we have that

$$\{F, K\}_-(\Pi, \alpha, l) = -\Pi \cdot (\nabla_\Pi F \times \nabla_\Pi K) + \{F, K\}_{\mathbb{R}}(\alpha, l). \quad (3.4)$$

See Krishnaprasad and Marsden [12]. In particular, for $F_\mu, K_\mu : \mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$, we have that $\tilde{\omega}_{\mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R}}^-(X_{F_\mu}, X_{K_\mu}) = \{F_\mu, K_\mu\}_-|_{\mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R}}$. Moreover, for reduced Hamiltonian $h_\mu(\Pi, \alpha, l) : \mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$, we have the Hamiltonian vector field $X_{h_\mu}(K_\mu) = \{K_\mu, h_\mu\}_-|_{\mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R}}$, and hence we have that

$$\begin{aligned} \frac{d\Pi}{dt} &= X_{h_\mu}(\Pi)(\Pi, \alpha, l) = \{\Pi, h_\mu\}_-(\Pi, \alpha, l) \\ &= -\Pi \cdot (\nabla_\Pi \Pi \times \nabla_\Pi h_\mu) + \left(\frac{\partial \Pi}{\partial \alpha} \frac{\partial h_\mu}{\partial l} - \frac{\partial h_\mu}{\partial \alpha} \frac{\partial \Pi}{\partial l} \right) = -\nabla_\Pi \Pi \cdot (\nabla_\Pi h_\mu \times \Pi) = \Pi \times \Omega, \end{aligned}$$

$$\begin{aligned} \frac{d\alpha}{dt} &= X_{h_\mu}(\alpha)(\Pi, \alpha, l) = \{\alpha, h_\mu\}_-(\Pi, \alpha, l) \\ &= -\Pi \cdot (\nabla_\Pi \alpha \times \nabla_\Pi h_\mu) + \left(\frac{\partial \alpha}{\partial \alpha} \frac{\partial h_\mu}{\partial l} - \frac{\partial h_\mu}{\partial \alpha} \frac{\partial \alpha}{\partial l} \right) = -\frac{(\Pi_3 - l)}{I_3} + \frac{l}{J_3}, \end{aligned}$$

$$\begin{aligned}\frac{dl}{dt} &= X_{h_\mu}(l)(\Pi, \alpha, l) = \{l, h_\mu\}_-(\Pi, \alpha, l) \\ &= -\Pi \cdot (\nabla_\Pi l \times \nabla_\Pi h_\mu) + \left(\frac{\partial l}{\partial \alpha} \frac{\partial h_\mu}{\partial l} - \frac{\partial h_\mu}{\partial \alpha} \frac{\partial l}{\partial l} \right) = 0,\end{aligned}$$

since $\nabla_\Pi \Pi = 1$, $\nabla_\Pi \alpha = \nabla_\Pi l = 0$, $\nabla_\Pi h_\mu = \Omega$, and $\frac{\partial \Pi}{\partial \alpha} = \frac{\partial l}{\partial \alpha} = \frac{\partial h_\mu}{\partial \alpha} = 0$. If we consider the rigid spacecraft-rotor system with a control torque $u : T^*Q \rightarrow T^*Q$ acting on the rotor, and $u \in \mathbf{J}_Q^{-1}(\mu)$ is invariant under the left $SO(3)$ -action, and its reduced control torque $u_\mu : \mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R} \rightarrow \mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R}$ is given by $u_\mu(\Pi, \alpha, l) = \pi_\mu(u(A, \Pi, \alpha, l)) = u(A, \Pi, \alpha, l)|_{\mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R}}$, where $\pi_\mu : \mathbf{J}_Q^{-1}(\mu) \rightarrow \mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R}$. Thus, in the case of coincident centers of buoyancy and gravity, the equations of motion for reduced spacecraft-rotor system with the control torque u acting on the rotor are given by

$$\begin{cases} \frac{d\Pi}{dt} = \Pi \times \Omega, \\ \frac{d\alpha}{dt} = -\frac{(\Pi_3 - l)}{I_3} + \frac{l}{J_3}, \\ \frac{dl}{dt} = \text{vlift}(u_\mu), \end{cases} \quad (3.5)$$

where $\text{vlift}(u_\mu) = \text{vlift}(u_\mu)X_{h_\mu} \in T(\mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R})$. Note that $\text{vlift}(u_\mu)X_{h_\mu}$ is the vertical lift of vector field X_{h_μ} under the action of u_μ along fibers, that is,

$$\text{vlift}(u_\mu)X_{h_\mu}(\Pi, \alpha, l) = \text{vlift}((Tu_\mu X_{h_\mu})(u_\mu(\Pi, \alpha, l)), (\Pi, \alpha, l)) = (Tu_\mu X_{h_\mu})_\sigma^v(\Pi, \alpha, l),$$

where σ is a geodesic in $\mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R}$ connecting $u_\mu(\Pi, \alpha, l)$ and (Π, α, l) , and $(Tu_\mu X_{h_\mu})_\sigma^v(\Pi, \alpha, l)$ is the parallel displacement of the vertical vector $(Tu_\mu X_{h_\mu})^v(\Pi, \alpha, l)$ along the geodesic σ from $u_\mu(\Pi, \alpha, l)$ to (Π, α, l) , see Marsden et al [22] and Wang [33]. To sum up the above discussion, we have the following theorem.

Theorem 3.1 *In the case of coincident centers of buoyancy and gravity, the spacecraft-rotor system with the control torque u acting on the rotor, that is, the 5-tuple $(T^*Q, SO(3), \omega_Q, H, u)$, where $Q = SO(3) \times S^1$, is a regular point reducible RCH system. For a point $\mu \in \mathfrak{so}^*(3)$, the regular value of the momentum map $\mathbf{J}_Q : SO(3) \times \mathfrak{so}^*(3) \times \mathbb{R} \times \mathbb{R} \rightarrow \mathfrak{so}^*(3)$, the regular point reduced system is the 4-tuple $(\mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R}, \tilde{\omega}_{\mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R}}^-, h_\mu, u_\mu)$, where $\mathcal{O}_\mu \subset \mathfrak{so}^*(3)$ is the coadjoint orbit, $\tilde{\omega}_{\mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R}}^-$ is orbit symplectic form on $\mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R}$, $h_\mu(\Pi, \alpha, l) = \pi_\mu(H(A, \Pi, \alpha, l)) = H(A, \Pi, \alpha, l)|_{\mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R}}$, $u_\mu(\Pi, \alpha, l) = \pi_\mu(u(A, \Pi, \alpha, l)) = u(A, \Pi, \alpha, l)|_{\mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R}}$, and its equations of motion are given by (3.5).*

Remark 3.2 *When the rigid spacecraft does not carry any internal rotor, in this case the configuration space is $Q = G = SO(3)$, the motion of rigid spacecraft is just the rotation motion of a rigid body, the above symmetric reduction of spacecraft-rotor system is just the Marsden-Weinstein reduction of a rigid body at a regular value of momentum map, and the motion equation (3.5) of reduced spacecraft-rotor system becomes the motion equation of a reduced rigid body on a coadjoint orbit of Lie group $SO(3)$. See Marsden and Ratiu [21].*

3.2 Symmetric Reduction of Spacecraft-Rotor System with Non-coincident Centers

In the following we shall give the regular point reduction of spacecraft-rotor system with non-coincident centers of buoyancy and gravity. Because the drift in the direction of gravity breaks

the symmetry and the spacecraft-rotor system is no longer $\text{SO}(3)$ invariant. In this case, its physical phase space is $T^*\text{SO}(3) \times T^*S^1$ and the symmetry group is S^1 , regarded as rotations about the third principal axis, that is, the axis of gravity. By the semidirect product reduction theorem, see Marsden et al [19], we know that the reduction of $T^*\text{SO}(3)$ by S^1 gives a space which is symplectically diffeomorphic to the reduced space obtained by the reduction of $T^*\text{SE}(3)$ by left action of $\text{SE}(3)$, that is the coadjoint orbit $\mathcal{O}_{(\mu,a)} \subset \mathfrak{se}^*(3) \cong T^*\text{SE}(3)/\text{SE}(3)$. In fact, in this case, we can identify the phase space $T^*\text{SO}(3)$ with the reduction of the cotangent bundle of the special Euclidean group $\text{SE}(3) = \text{SO}(3) \ltimes \mathbb{R}^3$ by the Euclidean translation subgroup \mathbb{R}^3 and identifies the symmetry group S^1 with isotropy group $G_a = \{A \in \text{SO}(3) \mid Aa = a\} = S^1$, which is Abelian and $(G_a)_{\mu_a} = G_a = S^1$, $\forall \mu_a \in \mathfrak{g}_a^*$, where a is a vector aligned with the direction of gravity and where $\text{SO}(3)$ acts on \mathbb{R}^3 in the standard way.

Assume that Lie group $G = \text{SE}(3)$ acts freely and properly on $Q = \text{SE}(3) \times S^1$ by the left translation on the first factor $\text{SE}(3)$, and the trivial action on the second factor S^1 . By using the left trivialization of $T^*\text{SE}(3) = \text{SE}(3) \times \mathfrak{se}^*(3)$, then the action of $\text{SE}(3)$ on phase space $T^*Q = T^*\text{SE}(3) \times T^*S^1$ is by cotangent lift of left translation on $\text{SE}(3)$ at the identity, that is, $\Phi : \text{SE}(3) \times T^*\text{SE}(3) \times T^*S^1 \cong \text{SE}(3) \times \text{SE}(3) \times \mathfrak{se}^*(3) \times \mathbb{R} \times \mathbb{R} \rightarrow \text{SE}(3) \times \mathfrak{se}^*(3) \times \mathbb{R} \times \mathbb{R}$, given by $\Phi((B, b)(A, c, \Pi, \Gamma, \alpha, l)) = (BA, c, \Pi, \Gamma, \alpha, l)$, for any $A, B \in \text{SO}(3)$, $\Pi \in \mathfrak{so}^*(3)$, $b, c, \Gamma \in \mathbb{R}^3$, $\alpha, l \in \mathbb{R}$, which is also free and proper, and the orbit space $(T^*Q)/\text{SE}(3)$ is a smooth manifold and $\pi : T^*Q \rightarrow (T^*Q)/\text{SE}(3)$ is a smooth submersion. Since $\text{SE}(3)$ acts trivially on $\mathfrak{se}^*(3)$ and \mathbb{R} , it follows that $(T^*Q)/\text{SE}(3)$ is diffeomorphic to $\mathfrak{se}^*(3) \times \mathbb{R} \times \mathbb{R}$.

We know that $\mathfrak{se}^*(3)$ is a Poisson manifold with respect to its heavy top Lie-Poisson bracket defined by

$$\{F, K\}_{\mathfrak{se}^*(3)}(\Pi, \Gamma) = -\Pi \cdot (\nabla_{\Pi} F \times \nabla_{\Pi} K) - \Gamma \cdot (\nabla_{\Pi} F \times \nabla_{\Gamma} K - \nabla_{\Pi} K \times \nabla_{\Gamma} F), \quad (3.6)$$

where $F, K \in C^\infty(\mathfrak{se}^*(3))$, $(\Pi, \Gamma) \in \mathfrak{se}^*(3)$. For $(\mu, a) \in \mathfrak{se}^*(3)$, the coadjoint orbit $\mathcal{O}_{(\mu,a)} \subset \mathfrak{se}^*(3)$ has the induced orbit symplectic form $\omega_{\mathcal{O}_{(\mu,a)}}^-$, which is coincide with the restriction of the Lie-Poisson bracket on $\mathfrak{se}^*(3)$ to the coadjoint orbit $\mathcal{O}_{(\mu,a)}$, and the coadjoint orbits $(\mathcal{O}_{(\mu,a)}, \omega_{\mathcal{O}_{(\mu,a)}}^-)$, $(\mu, a) \in \mathfrak{se}^*(3)$, form the symplectic leaves of the Poisson manifold $(\mathfrak{se}^*(3), \{\cdot, \cdot\}_{\mathfrak{se}^*(3)})$. Let $\omega_{\mathbb{R}}$ be the canonical symplectic form on $T^*\mathbb{R} \cong \mathbb{R} \times \mathbb{R}$ given by (3.2), and it induces a canonical Poisson bracket $\{\cdot, \cdot\}_{\mathbb{R}}$ on $T^*\mathbb{R}$ given by (3.3). Thus, we can induce a symplectic form $\tilde{\omega}_{\mathcal{O}_{(\mu,a)} \times \mathbb{R} \times \mathbb{R}}^- = \pi_{\mathcal{O}_{(\mu,a)}}^* \omega_{\mathcal{O}_{(\mu,a)}}^- + \pi_{\mathbb{R}}^* \omega_{\mathbb{R}}$ on the smooth manifold $\mathcal{O}_{(\mu,a)} \times \mathbb{R} \times \mathbb{R}$, where the maps $\pi_{\mathcal{O}_{(\mu,a)}} : \mathcal{O}_{(\mu,a)} \times \mathbb{R} \times \mathbb{R} \rightarrow \mathcal{O}_{(\mu,a)}$ and $\pi_{\mathbb{R}} : \mathcal{O}_{(\mu,a)} \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R} \times \mathbb{R}$ are canonical projections, and induce a Poisson bracket $\{\cdot, \cdot\}_- = \pi_{\mathfrak{se}^*(3)}^* \{\cdot, \cdot\}_{\mathfrak{se}^*(3)} + \pi_{\mathbb{R}}^* \{\cdot, \cdot\}_{\mathbb{R}}$ on the smooth manifold $\mathfrak{se}^*(3) \times \mathbb{R} \times \mathbb{R}$, where the maps $\pi_{\mathfrak{se}^*(3)} : \mathfrak{se}^*(3) \times \mathbb{R} \times \mathbb{R} \rightarrow \mathfrak{se}^*(3)$ and $\pi_{\mathbb{R}} : \mathfrak{se}^*(3) \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R} \times \mathbb{R}$ are canonical projections, and such that $(\mathcal{O}_{(\mu,a)} \times \mathbb{R} \times \mathbb{R}, \tilde{\omega}_{\mathcal{O}_{(\mu,a)} \times \mathbb{R} \times \mathbb{R}}^-)$ is a symplectic leaf of the Poisson manifold $(\mathfrak{se}^*(3) \times \mathbb{R} \times \mathbb{R}, \{\cdot, \cdot\}_-)$.

On the other hand, from $T^*Q = T^*\text{SE}(3) \times T^*S^1$ we know that there is a canonical symplectic form $\omega_Q = \pi_{\text{SE}(3)}^* \omega_1 + \pi_{S^1}^* \omega_{S^1}$ on T^*Q , where ω_1 is the canonical symplectic form on $T^*\text{SE}(3)$ and the maps $\pi_{\text{SE}(3)} : Q = \text{SE}(3) \times S^1 \rightarrow \text{SE}(3)$ and $\pi_{S^1} : Q = \text{SE}(3) \times S^1 \rightarrow S^1$ are canonical projections. Then the cotangent lift of left $\text{SE}(3)$ -action $\Phi : \text{SE}(3) \times T^*Q \rightarrow T^*Q$ is also symplectic, and admits an associated Ad^* -equivariant momentum map $\mathbf{J}_Q : T^*Q \rightarrow \mathfrak{se}^*(3)$ such that $\mathbf{J}_Q \cdot \pi_{\text{SE}(3)}^* = \mathbf{J}_{\text{SE}(3)}$, where $\mathbf{J}_{\text{SE}(3)} : T^*\text{SE}(3) \rightarrow \mathfrak{se}^*(3)$ is a momentum map of left $\text{SE}(3)$ -action on $T^*\text{SE}(3)$, and $\pi_{\text{SE}(3)}^* : T^*\text{SE}(3) \rightarrow T^*Q$. If $(\mu, a) \in \mathfrak{se}^*(3)$ is a regular value of \mathbf{J}_Q , then

$(\mu, a) \in \mathfrak{se}^*(3)$ is also a regular value of $\mathbf{J}_{\text{SE}(3)}$ and $\mathbf{J}_Q^{-1}(\mu, a) \cong \mathbf{J}_{\text{SE}(3)}^{-1}(\mu, a) \times \mathbb{R} \times \mathbb{R}$. Denote by $\text{SE}(3)_{(\mu, a)} = \{g \in \text{SE}(3) \mid \text{Ad}_g^*(\mu, a) = (\mu, a)\}$ the isotropy subgroup of coadjoint $\text{SE}(3)$ -action at the point $(\mu, a) \in \mathfrak{se}^*(3)$. It follows that $\text{SE}(3)_{(\mu, a)}$ acts also freely and properly on $\mathbf{J}_Q^{-1}(\mu, a)$, the regular point reduced space $(T^*Q)_{(\mu, a)} = \mathbf{J}_Q^{-1}(\mu, a)/\text{SE}(3)_{(\mu, a)} \cong (T^*\text{SE}(3))_{(\mu, a)} \times \mathbb{R} \times \mathbb{R}$ of (T^*Q, ω_Q) at (μ, a) , is a symplectic manifold with symplectic form $\omega_{(\mu, a)}$ uniquely characterized by the relation $\pi_{(\mu, a)}^* \omega_{(\mu, a)} = i_{(\mu, a)}^* \omega_Q = i_{(\mu, a)}^* \pi_{\text{SE}(3)}^* \omega_1 + i_{(\mu, a)}^* \pi_{S^1}^* \omega_{S^1}$, where the map $i_{(\mu, a)} : \mathbf{J}_Q^{-1}(\mu, a) \rightarrow T^*Q$ is the inclusion and $\pi_{(\mu, a)} : \mathbf{J}_Q^{-1}(\mu, a) \rightarrow (T^*Q)_{(\mu, a)}$ is the projection. Because from Abraham and Marsden [1], we know that $((T^*\text{SE}(3))_{(\mu, a)}, \omega_{(\mu, a)})$ is symplectically diffeomorphic to $(\mathcal{O}_{(\mu, a)}, \omega_{\mathcal{O}_{(\mu, a)}}^-)$, and hence we have that $((T^*Q)_{(\mu, a)}, \omega_{(\mu, a)})$ is symplectically diffeomorphic to $(\mathcal{O}_{(\mu, a)} \times \mathbb{R} \times \mathbb{R}, \tilde{\omega}_{\mathcal{O}_{(\mu, a)} \times \mathbb{R} \times \mathbb{R}}^-)$, which is a symplectic leaf of the Poisson manifold $(\mathfrak{se}^*(3) \times \mathbb{R} \times \mathbb{R}, \{\cdot, \cdot\}_-)$.

From the expression (2.2) of the Hamiltonian, we know that $H(A, c, \Pi, \Gamma, \alpha, l)$ is invariant under the left $\text{SE}(3)$ -action $\Phi : \text{SE}(3) \times T^*Q \rightarrow T^*Q$. For the case $(\mu, a) \in \mathfrak{se}^*(3)$ is the regular value of \mathbf{J}_Q , we have the reduced Hamiltonian $h_{(\mu, a)}(\Pi, \Gamma, \alpha, l) : \mathcal{O}_{(\mu, a)} \times \mathbb{R} \times \mathbb{R} (\subset \mathfrak{se}^*(3) \times \mathbb{R} \times \mathbb{R}) \rightarrow \mathbb{R}$ given by $h_{(\mu, a)}(\Pi, \Gamma, \alpha, l) = \pi_{(\mu, a)}(H(A, c, \Pi, \Gamma, \alpha, l)) = H(A, c, \Pi, \Gamma, \alpha, l)|_{\mathcal{O}_{(\mu, a)} \times \mathbb{R} \times \mathbb{R}}$. From the heavy top Poisson bracket on $\mathfrak{se}^*(3)$ and the Poisson bracket on $T^*\mathbb{R}$, we can get the Poisson bracket on $\mathfrak{se}^*(3) \times \mathbb{R} \times \mathbb{R}$, that is, for $F, K : \mathfrak{se}^*(3) \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$, we have that

$$\begin{aligned} \{F, K\}_- (\Pi, \Gamma, \alpha, l) &= -\Pi \cdot (\nabla_{\Pi} F \times \nabla_{\Pi} K) \\ &\quad - \Gamma \cdot (\nabla_{\Pi} F \times \nabla_{\Gamma} K - \nabla_{\Pi} K \times \nabla_{\Gamma} F) + \{F, K\}_{\mathbb{R}}(\alpha, l). \end{aligned} \quad (3.7)$$

See Krishnaprasad and Marsden [12]. In particular, for $F_{(\mu, a)}, K_{(\mu, a)} : \mathcal{O}_{(\mu, a)} \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$, we have that $\tilde{\omega}_{\mathcal{O}_{(\mu, a)} \times \mathbb{R} \times \mathbb{R}}(X_{F_{(\mu, a)}}, X_{K_{(\mu, a)}}) = \{F_{(\mu, a)}, K_{(\mu, a)}\}_-|_{\mathcal{O}_{(\mu, a)} \times \mathbb{R} \times \mathbb{R}}$. Moreover, for reduced Hamiltonian $h_{(\mu, a)}(\Pi, \Gamma, \alpha, l) : \mathcal{O}_{(\mu, a)} \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$, we have the Hamiltonian vector field $X_{h_{(\mu, a)}}(K_{(\mu, a)}) = \{K_{(\mu, a)}, h_{(\mu, a)}\}_-|_{\mathcal{O}_{(\mu, a)} \times \mathbb{R} \times \mathbb{R}}$, and hence we have that

$$\begin{aligned} \frac{d\Pi}{dt} &= X_{h_{(\mu, a)}}(\Pi)(\Pi, \Gamma, \alpha, l) = \{\Pi, h_{(\mu, a)}\}_- (\Pi, \Gamma, \alpha, l) = -\Pi \cdot (\nabla_{\Pi} \Pi \times \nabla_{\Pi} h_{(\mu, a)}) \\ &\quad - \Gamma \cdot (\nabla_{\Pi} \Pi \times \nabla_{\Gamma} h_{(\mu, a)} - \nabla_{\Pi} h_{(\mu, a)} \times \nabla_{\Gamma} \Pi) + \left(\frac{\partial \Pi}{\partial \alpha} \frac{\partial h_{(\mu, a)}}{\partial l} - \frac{\partial h_{(\mu, a)}}{\partial \alpha} \frac{\partial \Pi}{\partial l} \right) \\ &= \Pi \times \Omega - mgh\chi \times \Gamma = \Pi \times \Omega + mgh\Gamma \times \chi, \end{aligned}$$

$$\begin{aligned} \frac{d\Gamma}{dt} &= X_{h_{(\mu, a)}}(\Gamma)(\Pi, \Gamma, \alpha, l) = \{\Gamma, h_{(\mu, a)}\}_- (\Pi, \Gamma, \alpha, l) = -\Pi \cdot (\nabla_{\Pi} \Gamma \times \nabla_{\Pi} h_{(\mu, a)}) \\ &\quad - \Gamma \cdot (\nabla_{\Pi} \Gamma \times \nabla_{\Gamma} h_{(\mu, a)} - \nabla_{\Pi} h_{(\mu, a)} \times \nabla_{\Gamma} \Gamma) + \left(\frac{\partial \Gamma}{\partial \alpha} \frac{\partial h_{(\mu, a)}}{\partial l} - \frac{\partial h_{(\mu, a)}}{\partial \alpha} \frac{\partial \Gamma}{\partial l} \right) \\ &= \nabla_{\Gamma} \Gamma \cdot (\Gamma \times \nabla_{\Pi} h_{(\mu, a)}) = \Gamma \times \Omega, \end{aligned}$$

$$\begin{aligned} \frac{d\alpha}{dt} &= X_{h_{(\mu, a)}}(\alpha)(\Pi, \Gamma, \alpha, l) = \{\alpha, h_{(\mu, a)}\}_- (\Pi, \Gamma, \alpha, l) = -\Pi \cdot (\nabla_{\Pi} \alpha \times \nabla_{\Pi} h_{(\mu, a)}) \\ &\quad - \Gamma \cdot (\nabla_{\Pi} \alpha \times \nabla_{\Gamma} h_{(\mu, a)} - \nabla_{\Pi} h_{(\mu, a)} \times \nabla_{\Gamma} \alpha) + \left(\frac{\partial \alpha}{\partial \alpha} \frac{\partial h_{(\mu, a)}}{\partial l} - \frac{\partial h_{(\mu, a)}}{\partial \alpha} \frac{\partial \alpha}{\partial l} \right) \\ &= -\frac{(\Pi_3 - l)}{\bar{I}_3} + \frac{l}{J_3}, \end{aligned}$$

$$\begin{aligned} \frac{dl}{dt} &= X_{h(\mu,a)}(l)(\Pi, \Gamma, \alpha, l) = \{l, h(\mu,a)\}_-(\Pi, \Gamma, \alpha, l) = -\Pi \cdot (\nabla_{\Pi} l \times \nabla_{\Pi} h(\mu,a)) \\ &\quad - \Gamma \cdot (\nabla_{\Pi} l \times \nabla_{\Gamma} h(\mu,a) - \nabla_{\Pi} h(\mu,a) \times \nabla_{\Gamma} l) + \left(\frac{\partial l}{\partial \alpha} \frac{\partial h(\mu,a)}{\partial l} - \frac{\partial h(\mu,a)}{\partial \alpha} \frac{\partial l}{\partial l} \right) = 0, \end{aligned}$$

since $\nabla_{\Pi} \Pi = \nabla_{\Gamma} \Gamma = 1$, $\nabla_{\Gamma} \Pi = \nabla_{\Pi} \Gamma = \nabla_{\Pi} \alpha = \nabla_{\Pi} l = \nabla_{\Gamma} \alpha = \nabla_{\Gamma} l = 0$, $\nabla_{\Pi} h(\mu,a) = \Omega$, and $\frac{\partial \Pi}{\partial \alpha} = \frac{\partial \Gamma}{\partial \alpha} = \frac{\partial l}{\partial \alpha} = \frac{\partial h(\mu,a)}{\partial \alpha} = 0$. If we consider the rigid spacecraft-rotor system with a control torque $u : T^*Q \rightarrow T^*Q$ acting on the rotor, and $u \in \mathbf{J}_Q^{-1}(\mu, a)$ is invariant under the left $SE(3)$ -action, and its reduced control torque $u_{(\mu,a)} : \mathcal{O}_{(\mu,a)} \times \mathbb{R} \times \mathbb{R} \rightarrow \mathcal{O}_{(\mu,a)} \times \mathbb{R} \times \mathbb{R}$ is given by $u_{(\mu,a)}(\Pi, \Gamma, \alpha, l) = \pi_{(\mu,a)}(u(A, c, \Pi, \Gamma, \alpha, l)) = u(A, c, \Pi, \Gamma, \alpha, l)|_{\mathcal{O}_{(\mu,a)} \times \mathbb{R} \times \mathbb{R}}$, where $\pi_{(\mu,a)} : \mathbf{J}_Q^{-1}(\mu, a) \rightarrow \mathcal{O}_{(\mu,a)} \times \mathbb{R} \times \mathbb{R}$. Thus, in the case of non-coincident centers of buoyancy and gravity, the equations of motion for reduced spacecraft-rotor system with the control torque u acting on the rotor are given by

$$\begin{cases} \frac{d\Pi}{dt} = \Pi \times \Omega + mgh\Gamma \times \chi, \\ \frac{d\Gamma}{dt} = \Gamma \times \Omega, \\ \frac{d\alpha}{dt} = -\frac{(\Pi_3 - l)}{I_3} + \frac{l}{J_3}, \\ \frac{dl}{dt} = \text{vlift}(u_{(\mu,a)}). \end{cases} \quad (3.8)$$

where $\text{vlift}(u_{(\mu,a)}) = \text{vlift}(u_{(\mu,a)})X_{h(\mu,a)} \in T(\mathcal{O}_{(\mu,a)} \times \mathbb{R} \times \mathbb{R})$. Note that $\text{vlift}(u_{(\mu,a)})X_{h(\mu,a)}$ is the vertical lift of vector field $X_{h(\mu,a)}$ under the action of $u_{(\mu,a)}$ along fibers, that is,

$$\begin{aligned} \text{vlift}(u_{(\mu,a)})X_{h(\mu,a)}(\Pi, \Gamma, \alpha, l) &= \text{vlift}((Tu_{(\mu,a)}X_{h(\mu,a)})(u_{(\mu,a)}(\Pi, \Gamma, \alpha, l)), (\Pi, \Gamma, \alpha, l)) \\ &= (Tu_{(\mu,a)}X_{h(\mu,a)})_{\sigma}^v(\Pi, \Gamma, \alpha, l), \end{aligned}$$

see Marsden et al [22] and Wang [33]. To sum up the above discussion, we have the following theorem.

Theorem 3.3 *In the case of non-coincident centers of buoyancy and gravity, the spacecraft-rotor system with the control torque u acting on the rotor, that is, the 5-tuple $(T^*Q, SE(3), \omega_Q, H, u)$, where $Q = SE(3) \times S^1$, is a regular point reducible RCH system. For a point $(\mu, a) \in \mathfrak{se}^*(3)$, the regular value of the momentum map $\mathbf{J}_Q : SE(3) \times \mathfrak{se}^*(3) \times \mathbb{R} \times \mathbb{R} \rightarrow \mathfrak{se}^*(3)$, the regular point reduced system is the 4-tuple $(\mathcal{O}_{(\mu,a)} \times \mathbb{R} \times \mathbb{R}, \tilde{\omega}_{\mathcal{O}_{(\mu,a)} \times \mathbb{R} \times \mathbb{R}}^-, h_{(\mu,a)}, u_{(\mu,a)})$, where $\mathcal{O}_{(\mu,a)} \subset \mathfrak{se}^*(3)$ is the coadjoint orbit, $\tilde{\omega}_{\mathcal{O}_{(\mu,a)} \times \mathbb{R} \times \mathbb{R}}^-$ is orbit symplectic form on $\mathcal{O}_{(\mu,a)} \times \mathbb{R} \times \mathbb{R}$, $h_{(\mu,a)}(\Pi, \Gamma, \alpha, l) = \pi_{(\mu,a)}(H(A, c, \Pi, \Gamma, \alpha, l)) = H(A, c, \Pi, \Gamma, \alpha, l)|_{\mathcal{O}_{(\mu,a)} \times \mathbb{R} \times \mathbb{R}}$, and $u_{(\mu,a)}(\Pi, \Gamma, \alpha, l) = \pi_{(\mu,a)}(u(A, c, \Pi, \Gamma, \alpha, l)) = u(A, c, \Pi, \Gamma, \alpha, l)|_{\mathcal{O}_{(\mu,a)} \times \mathbb{R} \times \mathbb{R}}$, and its equations of motion are given by (3.8).*

Remark 3.4 *When the rigid spacecraft does not carry any internal rotor, in this case the configuration space is $Q = G = SE(3)$, the motion of rigid spacecraft is just the rotation motion with drift of a rigid body, the above symmetric reduction of spacecraft-rotor system is just the Marsden-Weinstein reduction of a heavy top at a regular value of momentum map, and the motion equation (3.8) of reduced spacecraft-rotor system becomes the motion equation of a reduced heavy top on a coadjoint orbit of Lie group $SE(3)$. See Marsden and Ratiu [21].*

4 Hamilton-Jacobi Equation of the Rigid Spacecraft with a Rotor

It is well-known that Hamilton-Jacobi theory provides a characterization of the generating functions of certain time-dependent canonical transformations, such that a given Hamiltonian system in such a form that its solutions are extremely easy to find by reduction to the equilibrium, see Abraham and Marsden [1], Arnold [2] and Marsden and Ratiu [21]. In general, we know that it is not easy to find the solutions of Hamilton's equation. But, if we can get a solution of Hamilton-Jacobi equation of the Hamiltonian system, by using the relationship between Hamilton's equation and Hamilton-Jacobi equation, it is easy to give a special solution of Hamilton's equation. Thus, it is very important to give explicitly the Hamilton-Jacobi equation of a Hamiltonian system. Recently, the author in [33] proved the following Hamilton-Jacobi theorem for regular point reducible RCH system on the generalization of a Lie group.

Theorem 4.1 *For the regular point reducible RCH system $(T^*Q, G, \omega_Q, H, F, W)$ on the generalization of a Lie group $Q = G \times V$, where G is a Lie group and V is a k -dimensional vector space, assume that $\gamma : Q \rightarrow T^*Q$ is an one-form on Q , and $\gamma^* : T^*T^*Q \rightarrow T^*Q$ is symplectic, where there is an induced symplectic form $\pi_Q^* \omega_Q$ on T^*T^*Q , $\pi_Q^* : T^*Q \rightarrow T^*T^*Q$, $\pi_Q : T^*Q \rightarrow Q$, and γ is closed with respect to $T\pi_Q : TT^*Q \rightarrow TQ$, and $\tilde{X}^\gamma = T\pi_Q \cdot \tilde{X} \cdot \gamma$, where $\tilde{X} = X_{(T^*Q, G, \omega_Q, H, F, u)}$ is the dynamical vector field of the regular point reducible RCH system $(T^*Q, G, \omega_Q, H, F, W)$ with a control law u . Moreover, assume that $\mu \in \mathfrak{g}^*$ is the regular reducible point of the RCH system, and $\text{Im}(\gamma) \subset \mathbf{J}_Q^{-1}(\mu)$, and it is G_μ -invariant, and $\bar{\gamma} = \pi_\mu(\gamma) : Q \rightarrow \mathcal{O}_\mu \times V \times V^*$, where G_μ is the isotropy subgroup of coadjoint action at μ , and $\pi_\mu : \mathbf{J}_Q^{-1}(\mu) \rightarrow (T^*Q)_\mu \cong \mathcal{O}_\mu \times V \times V^*$. Then the following two assertions are equivalent: (i) \tilde{X}^γ and \tilde{X}_μ are $\bar{\gamma}$ -related, where $\tilde{X}_\mu = X_{(\mathcal{O}_\mu \times V \times V^*, \tilde{\omega}_{\mathcal{O}_\mu \times V \times V^*}, h_\mu, f_\mu, u_\mu)}$ is the dynamical vector field of regular point reduced RCH system $(\mathcal{O}_\mu \times V \times V^*, \tilde{\omega}_{\mathcal{O}_\mu \times V \times V^*}, h_\mu, f_\mu, u_\mu)$; (ii) $X_{h_\mu \cdot \bar{\gamma}} + \text{vlift}(f_\mu \cdot \bar{\gamma}) + \text{vlift}(u_\mu \cdot \bar{\gamma}) = 0$, or $X_{h_\mu \cdot \bar{\gamma}} + \text{vlift}(f_\mu \cdot \bar{\gamma}) + \text{vlift}(u_\mu \cdot \bar{\gamma}) = \tilde{X}^\gamma$. Moreover, γ is a solution of the Hamilton-Jacobi equation $X_{H \cdot \gamma} + \text{vlift}(F \cdot \gamma) + \text{vlift}(u \cdot \gamma) = 0$, if and only if $\bar{\gamma}$ is a solution of the Hamilton-Jacobi equation $X_{h_\mu \cdot \bar{\gamma}} + \text{vlift}(f_\mu \cdot \bar{\gamma}) + \text{vlift}(u_\mu \cdot \bar{\gamma}) = 0$.*

The maps involved in the above theorem are shown in Diagram-1.

$$\begin{array}{ccccc}
 \mathbf{J}^{-1}(\mu) & \xrightarrow{i_\mu} & T^*Q & \xrightarrow{\pi_Q^*} & T^*T^*Q & \xleftarrow{\pi_\mu^*} & T^*(T^*Q)_\mu \\
 & & \downarrow \pi_Q & & \downarrow \gamma^* & \swarrow \bar{\gamma}^* & \\
 & & Q & \xrightarrow{\gamma} & T^*Q & \xrightarrow{\pi_\mu} & (T^*Q)_\mu \\
 & & \downarrow \tilde{X}^\gamma & & \downarrow \tilde{X} & & \downarrow \tilde{X}_\mu \\
 & & TQ & \xrightarrow{T\gamma} & T(T^*Q) & \xrightarrow{T\pi_\mu} & T(T^*Q)_\mu
 \end{array}$$

Diagram-1

As an application of the theoretical result, in this section we regard the rigid spacecraft with an internal rotor as a regular point reducible RCH system on the generalization of rotation group $\text{SO}(3) \times S^1$ and on the generalization of Euclidean group $\text{SE}(3) \times S^1$, respectively, and give the Hamilton-Jacobi equations of their reduced RCH systems on the symplectic leaves by calculation in detail, which show the effect on controls in Hamilton-Jacobi theory. We shall follow the notations and conventions introduced in Marsden and Ratiu [21], Marsden [18], Marsden et al [22], and Wang [33].

4.1 H-J Equation of Spacecraft-Rotor System with Coincident Centers

In the following we first give the Hamilton-Jacobi equation for regular point reduced spacecraft-rotor system with coincident centers of buoyancy and gravity. From the expression (2.1) of the Hamiltonian, we know that $H(A, \Pi, \alpha, l)$ is invariant under the left $\text{SO}(3)$ -action. For the case $\mu \in \mathfrak{so}^*(3)$ is the regular value of \mathbf{J}_Q , we have the reduced Hamiltonian $h_\mu(\Pi, \alpha, l) : \mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$, which is given by $h_\mu(\Pi, \alpha, l) = \pi_\mu(H(A, \Pi, \alpha, l)) = H(A, \Pi, \alpha, l)|_{\mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R}}$, and we have the reduced Hamiltonian vector field $X_{h_\mu}(K_\mu) = \{K_\mu, h_\mu\}_-|_{\mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R}}$.

Assume that $\gamma : \text{SO}(3) \times S^1 \rightarrow T^*(\text{SO}(3) \times S^1)$ is an one-form on $\text{SO}(3) \times S^1$, and $\gamma^* : T^*T^*(\text{SO}(3) \times S^1) \rightarrow T^*(\text{SO}(3) \times S^1)$ is symplectic, and γ is closed with respect to $T\pi_{\text{SO}(3) \times S^1} : TT^*(\text{SO}(3) \times S^1) \rightarrow T(\text{SO}(3) \times S^1)$, and $\text{Im}(\gamma) \subset \mathbf{J}_Q^{-1}(\mu)$, and it is $\text{SO}(3)_\mu$ -invariant, and $\bar{\gamma} = \pi_\mu(\gamma) : \text{SO}(3) \times S^1 \rightarrow \mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R}$. Denote by $\bar{\gamma}(A, \alpha) = (\bar{\gamma}_1, \bar{\gamma}_2, \bar{\gamma}_3, \bar{\gamma}_4, \bar{\gamma}_5)(A, \alpha) \in \mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R} (\subset \mathfrak{so}^*(3) \times \mathbb{R} \times \mathbb{R})$, and $(\bar{\gamma}_1, \bar{\gamma}_2, \bar{\gamma}_3)(A, \alpha) \in \mathcal{O}_\mu$, then $h_\mu \cdot \bar{\gamma} : \text{SO}(3) \times S^1 \rightarrow \mathbb{R}$ is given by

$$h_\mu \cdot \bar{\gamma}(A, \alpha) = H \cdot \bar{\gamma}(A, \alpha)|_{\mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R}} = \frac{1}{2} \left[\frac{\bar{\gamma}_1^2}{\bar{I}_1} + \frac{\bar{\gamma}_2^2}{\bar{I}_2} + \frac{(\bar{\gamma}_3 - \bar{\gamma}_5)^2}{\bar{I}_3} + \frac{\bar{\gamma}_5^2}{\bar{J}_3} \right], \quad (4.1)$$

and the vector field

$$\begin{aligned} X_{h_\mu \cdot \bar{\gamma}}(\Pi) &= \{\Pi, h_\mu \cdot \bar{\gamma}(A, \alpha)\}_-|_{\mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R}} \\ &= -\Pi \cdot (\nabla_\Pi \Pi \times \nabla_\Pi (h_\mu \cdot \bar{\gamma})) + \{\Pi, h_\mu \cdot \bar{\gamma}(A, \alpha)\}_\mathbb{R}|_{\mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R}} \\ &= -\nabla_\Pi \Pi \cdot (\nabla_\Pi (h_\mu \cdot \bar{\gamma}) \times \Pi) + \left(\frac{\partial \Pi}{\partial \alpha} \frac{\partial (h_\mu \cdot \bar{\gamma})}{\partial l} - \frac{\partial (h_\mu \cdot \bar{\gamma})}{\partial \alpha} \frac{\partial \Pi}{\partial l} \right) \\ &= (\Pi_1, \Pi_2, \Pi_3) \times \left(\frac{\bar{\gamma}_1}{\bar{I}_1}, \frac{\bar{\gamma}_2}{\bar{I}_2}, \frac{(\bar{\gamma}_3 - \bar{\gamma}_5)}{\bar{I}_3} \right) \\ &= \left(\frac{\bar{I}_2 \Pi_2 (\bar{\gamma}_3 - \bar{\gamma}_5) - \bar{I}_3 \Pi_3 \bar{\gamma}_2}{\bar{I}_2 \bar{I}_3}, \frac{\bar{I}_3 \Pi_3 \bar{\gamma}_1 - \bar{I}_1 \Pi_1 (\bar{\gamma}_3 - \bar{\gamma}_5)}{\bar{I}_3 \bar{I}_1}, \frac{\bar{I}_1 \Pi_1 \bar{\gamma}_2 - \bar{I}_2 \Pi_2 \bar{\gamma}_1}{\bar{I}_1 \bar{I}_2} \right), \end{aligned}$$

since $\nabla_\Pi \Pi = 1$, and $\nabla_{\Pi_j} (h_\mu \cdot \bar{\gamma}) = \bar{\gamma}_j / \bar{I}_j$, $j = 1, 2$, $\nabla_{\Pi_3} (h_\mu \cdot \bar{\gamma}) = (\bar{\gamma}_3 - \bar{\gamma}_5) / \bar{I}_3$, and $\frac{\partial \Pi}{\partial \alpha} = \frac{\partial (h_\mu \cdot \bar{\gamma})}{\partial \alpha} = 0$.

$$\begin{aligned} X_{h_\mu \cdot \bar{\gamma}}(\alpha) &= \{\alpha, h_\mu \cdot \bar{\gamma}(A, \alpha)\}_-|_{\mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R}} \\ &= -\Pi \cdot (\nabla_\Pi \alpha \times \nabla_\Pi (h_\mu \cdot \bar{\gamma})) + \{\alpha, h_\mu \cdot \bar{\gamma}(A, \alpha)\}_\mathbb{R}|_{\mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R}} \\ &= -\nabla_\Pi \alpha \cdot (\nabla_\Pi (h_\mu \cdot \bar{\gamma}) \times \Pi) + \left(\frac{\partial \alpha}{\partial \alpha} \frac{\partial (h_\mu \cdot \bar{\gamma})}{\partial l} - \frac{\partial (h_\mu \cdot \bar{\gamma})}{\partial \alpha} \frac{\partial \alpha}{\partial l} \right) \\ &= -\frac{(\bar{\gamma}_3 - \bar{\gamma}_5)}{\bar{I}_3} + \frac{\bar{\gamma}_5}{\bar{J}_3}, \end{aligned}$$

$$\begin{aligned} X_{h_\mu \cdot \bar{\gamma}}(l) &= \{l, h_\mu \cdot \bar{\gamma}(A, \alpha)\}_-|_{\mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R}} \\ &= -\Pi \cdot (\nabla_\Pi l \times \nabla_\Pi (h_\mu \cdot \bar{\gamma})) + \{l, h_\mu \cdot \bar{\gamma}(A, \alpha)\}_\mathbb{R}|_{\mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R}} \\ &= -\nabla_\Pi l \cdot (\nabla_\Pi (h_\mu \cdot \bar{\gamma}) \times \Pi) + \left(\frac{\partial l}{\partial \alpha} \frac{\partial (h_\mu \cdot \bar{\gamma})}{\partial l} - \frac{\partial (h_\mu \cdot \bar{\gamma})}{\partial \alpha} \frac{\partial l}{\partial l} \right) = 0, \end{aligned}$$

since $\nabla_\Pi \alpha = 0$, $\nabla_\Pi l = 0$, and $\frac{\partial l}{\partial \alpha} = \frac{\partial \alpha}{\partial l} = \frac{\partial (h_\mu \cdot \bar{\gamma})}{\partial \alpha} = 0$. If we consider the rigid spacecraft-rotor system with a control torque $u : T^*Q \rightarrow T^*Q$ acting on the rotor, and $u \in \mathbf{J}_Q^{-1}(\mu)$ is invariant

under the left $SO(3)$ -action, and its reduced control torque $u_\mu : \mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R} \rightarrow \mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R}$ is given by $u_\mu(\Pi, \alpha, l) = \pi_\mu(u(A, \Pi, \alpha, l)) = u(A, \Pi, \alpha, l)|_{\mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R}}$, where $\pi_\mu : \mathbf{J}_Q^{-1}(\mu) \rightarrow \mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R}$. The dynamical vector field of regular point reduced spacecraft-rotor system $(\mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R}, \tilde{\omega}_{\mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R}}, h_\mu, u_\mu)$ is given by

$$X_{(\mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R}, \tilde{\omega}_{\mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R}}, h_\mu, u_\mu)} = X_{h_\mu} + \text{vlift}(u_\mu),$$

where $\text{vlift}(u_\mu) = \text{vlift}(u_\mu)X_{h_\mu} \in T(\mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R})$. Assume that

$$\text{vlift}(u_\mu \cdot \bar{\gamma})X_{h_\mu}(A, \alpha) = (U_1, U_2, U_3, U_4, U_5)(A, \alpha) \in T(\mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R}).$$

Thus, in the case of coincident centers of buoyancy and gravity, the Hamilton-Jacobi equations for reduced spacecraft-rotor system with the control torque u acting on the rotor are given by

$$\begin{cases} \bar{I}_2 \Pi_2 (\bar{\gamma}_3 - \bar{\gamma}_5) - \bar{I}_3 \Pi_3 \bar{\gamma}_2 + \bar{I}_2 \bar{I}_3 U_1 = 0, \\ \bar{I}_3 \Pi_3 \bar{\gamma}_1 - \bar{I}_1 \Pi_1 (\bar{\gamma}_3 - \bar{\gamma}_5) + \bar{I}_3 \bar{I}_1 U_2 = 0, \\ \bar{I}_1 \Pi_1 \bar{\gamma}_2 - \bar{I}_2 \Pi_2 \bar{\gamma}_1 + \bar{I}_1 \bar{I}_2 U_3 = 0, \\ -J_3 (\bar{\gamma}_3 - \bar{\gamma}_5) + \bar{I}_3 \bar{\gamma}_5 + \bar{I}_3 J_3 U_4 = 0, \\ U_5 = 0. \end{cases} \quad (4.2)$$

To sum up the above discussion, we have the following theorem.

Theorem 4.2 *In the case of coincident centers of buoyancy and gravity, for a point $\mu \in \mathfrak{so}^*(3)$, the regular value of the momentum map $\mathbf{J}_Q : SO(3) \times \mathfrak{so}^*(3) \times \mathbb{R} \times \mathbb{R} \rightarrow \mathfrak{so}^*(3)$, the regular point reduced system of spacecraft-rotor system with the control torque u acting on the rotor $(T^*Q, SO(3), \omega_Q, H, u)$, where $Q = SO(3) \times S^1$, is the 4-tuple $(\mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R}, \tilde{\omega}_{\mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R}}, h_\mu, u_\mu)$, where $\mathcal{O}_\mu \subset \mathfrak{so}^*(3)$ is the coadjoint orbit, $\tilde{\omega}_{\mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R}}$ is orbit symplectic form on $\mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R}$, $h_\mu(\Pi, \alpha, l) = H(A, \Pi, \alpha, l)|_{\mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R}}$, $u_\mu(\Pi, \alpha, l) = u(A, \Pi, \alpha, l)|_{\mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R}}$. Assume that $\gamma : SO(3) \times S^1 \rightarrow T^*(SO(3) \times S^1)$ is an one-form on $SO(3) \times S^1$, and $\gamma^* : T^*T^*(SO(3) \times S^1) \rightarrow T^*(SO(3) \times S^1)$ is symplectic, and γ is closed with respect to $T\pi_Q : TT^*Q \rightarrow TQ$, and $\text{Im}(\gamma) \subset \mathbf{J}_Q^{-1}(\mu)$, and it is $SO(3)_\mu$ -invariant, where $SO(3)_\mu$ is the isotropy subgroup of coadjoint $SO(3)$ -action at the point $\mu \in \mathfrak{so}^*(3)$, and $\bar{\gamma} = \pi_\mu(\gamma) : SO(3) \times S^1 \rightarrow \mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R}$. Then $\bar{\gamma}$ is a solution of either Hamilton-Jacobi equation of reduced spacecraft-rotor system given by (4.2), or the equation $X_{h_\mu \cdot \bar{\gamma}} + \text{vlift}(u_\mu \cdot \bar{\gamma}) = \tilde{X}^\gamma$, if and only if \tilde{X}^γ and \tilde{X}_μ are $\bar{\gamma}$ -related, where $\tilde{X}^\gamma = T\pi_Q \cdot \tilde{X} \cdot \gamma$, $\tilde{X} = X_{(T^*Q, SO(3), \omega_Q, H, u)}$, and $\tilde{X}_\mu = X_{(\mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R}, \tilde{\omega}_{\mathcal{O}_\mu \times \mathbb{R} \times \mathbb{R}}, h_\mu, u_\mu)}$.*

Remark 4.3 *When the rigid spacecraft does not carry any internal rotor, in this case the configuration space is $Q = G = SO(3)$, the above Hamilton-Jacobi equation (4.2) of reduced spacecraft-rotor system is just the Lie-Poisson Hamilton-Jacobi equation of Marsden-Weinstein reduced Hamiltonian system $(\mathcal{O}_\mu, \omega_{\mathcal{O}_\mu}, h_{\mathcal{O}_\mu})$ on the coadjoint orbit of Lie group $SO(3)$, since the symplectic structure on the coadjoint orbit \mathcal{O}_μ is induced by the (-)-Lie-Poisson brackets on $\mathfrak{so}^*(3)$. See Marsden and Ratiu [21], Ge and Marsden [10], and Wang [32].*

4.2 H-J Equation of Spacecraft-Rotor System with Non-coincident Centers

In the following we shall give the Hamilton-Jacobi equation for regular point reduced spacecraft-rotor system with non-coincident centers of buoyancy and gravity. From the expression (2.2) of the Hamiltonian, we know that $H(A, c, \Pi, \Gamma, \alpha, l)$ is invariant under the left $SE(3)$ -action.

For the case $(\mu, a) \in \mathfrak{se}^*(3)$ is the regular value of \mathbf{J}_Q , we have the reduced Hamiltonian $h_{(\mu,a)}(\Pi, \Gamma, \alpha, l) : \mathcal{O}_{(\mu,a)} \times \mathbb{R} \times \mathbb{R} (\subset \mathfrak{se}^*(3) \times \mathbb{R} \times \mathbb{R}) \rightarrow \mathbb{R}$, which is given by $h_{(\mu,a)}(\Pi, \Gamma, \alpha, l) = \pi_{(\mu,a)}(H(A, c, \Pi, \Gamma, \alpha, l)) = H(A, c, \Pi, \Gamma, \alpha, l)|_{\mathcal{O}_{(\mu,a)} \times \mathbb{R} \times \mathbb{R}}$, and we have the reduced Hamiltonian vector field $X_{h_{(\mu,a)}}(K_{(\mu,a)}) = \{K_{(\mu,a)}, h_{(\mu,a)}\}_-|_{\mathcal{O}_{(\mu,a)} \times \mathbb{R} \times \mathbb{R}}$.

Assume that $\gamma : \text{SE}(3) \times S^1 \rightarrow T^*(\text{SE}(3) \times S^1)$ is an one-form on $\text{SE}(3) \times S^1$, and $\gamma^* : T^*T^*(\text{SE}(3) \times S^1) \rightarrow T^*(\text{SE}(3) \times S^1)$ is symplectic, and γ is closed with respect to $T\pi_{\text{SE}(3) \times S^1} : TT^*(\text{SE}(3) \times S^1) \rightarrow T(\text{SE}(3) \times S^1)$, and $\text{Im}(\gamma) \subset \mathbf{J}^{-1}((\mu, a))$, and it is $\text{SE}(3)_{(\mu,a)}$ -invariant, and $\bar{\gamma} = \pi_{(\mu,a)}(\gamma) : \text{SE}(3) \times S^1 \rightarrow \mathcal{O}_{(\mu,a)} \times \mathbb{R} \times \mathbb{R}$. Denote by $\bar{\gamma}(A, c, \alpha) = (\bar{\gamma}_1, \bar{\gamma}_2, \bar{\gamma}_3, \Gamma_1, \Gamma_2, \Gamma_3, \bar{\gamma}_4, \bar{\gamma}_5)(A, c, \alpha) \in \mathcal{O}_{(\mu,a)} \times \mathbb{R} \times \mathbb{R} (\subset \mathfrak{se}^*(3) \times \mathbb{R} \times \mathbb{R})$, then $h_{(\mu,a)} \cdot \bar{\gamma} : \text{SE}(3) \times S^1 \rightarrow \mathbb{R}$ is given by

$$\begin{aligned} h_{(\mu,a)} \cdot \bar{\gamma}(A, c, \alpha) &= H \cdot \bar{\gamma}(A, c, \alpha)|_{\mathcal{O}_{(\mu,a)} \times \mathbb{R} \times \mathbb{R}} \\ &= \frac{1}{2} \left[\frac{\bar{\gamma}_1^2}{\bar{I}_1} + \frac{\bar{\gamma}_2^2}{\bar{I}_2} + \frac{(\bar{\gamma}_3 - \bar{\gamma}_5)^2}{\bar{I}_3} + \frac{\bar{\gamma}_5^2}{J_3} \right] + mgh\Gamma \cdot \chi, \end{aligned} \quad (4.3)$$

and the vector field

$$\begin{aligned} X_{h_{(\mu,a)} \cdot \bar{\gamma}}(\Pi) &= \{\Pi, h_{(\mu,a)} \cdot \bar{\gamma}(A, c, \alpha)\}_-|_{\mathcal{O}_{(\mu,a)} \times \mathbb{R} \times \mathbb{R}} \\ &= -\Pi \cdot (\nabla_{\Pi}\Pi \times \nabla_{\Pi}(h_{(\mu,a)} \cdot \bar{\gamma})) - \Gamma \cdot (\nabla_{\Pi}\Pi \times \nabla_{\Gamma}(h_{(\mu,a)} \cdot \bar{\gamma}) - \nabla_{\Pi}(h_{(\mu,a)} \cdot \bar{\gamma}) \times \nabla_{\Gamma}\Pi) \\ &\quad + \{\Pi, h_{(\mu,a)} \cdot \bar{\gamma}(A, c, \alpha)\}_{\mathbb{R}}|_{\mathcal{O}_{(\mu,a)} \times \mathbb{R} \times \mathbb{R}} \\ &= -\nabla_{\Pi}\Pi \cdot (\nabla_{\Pi}(h_{(\mu,a)} \cdot \bar{\gamma}) \times \Pi) - \nabla_{\Pi}\Pi \cdot (\nabla_{\Gamma}(h_{(\mu,a)} \cdot \bar{\gamma}) \times \Gamma) \\ &\quad + \left(\frac{\partial \Pi}{\partial \alpha} \frac{\partial (h_{(\mu,a)} \cdot \bar{\gamma})}{\partial l} - \frac{\partial (h_{(\mu,a)} \cdot \bar{\gamma})}{\partial \alpha} \frac{\partial \Pi}{\partial l} \right) \\ &= (\Pi_1, \Pi_2, \Pi_3) \times \left(\frac{\bar{\gamma}_1}{\bar{I}_1}, \frac{\bar{\gamma}_2}{\bar{I}_2}, \frac{(\bar{\gamma}_3 - \bar{\gamma}_5)}{\bar{I}_3} \right) + mgh(\Gamma_1, \Gamma_2, \Gamma_3) \times (\chi_1, \chi_2, \chi_3) \\ &= \left(\frac{\bar{I}_2\Pi_2(\bar{\gamma}_3 - \bar{\gamma}_5) - \bar{I}_3\Pi_3\bar{\gamma}_2}{\bar{I}_2\bar{I}_3} + mgh(\Gamma_2\chi_3 - \Gamma_3\chi_2), \right. \\ &\quad \left. \frac{\bar{I}_3\Pi_3\bar{\gamma}_1 - \bar{I}_1\Pi_1(\bar{\gamma}_3 - \bar{\gamma}_5)}{\bar{I}_3\bar{I}_1} + mgh(\Gamma_3\chi_1 - \Gamma_1\chi_3), \right. \\ &\quad \left. \frac{\bar{I}_1\Pi_1\bar{\gamma}_2 - \bar{I}_2\Pi_2\bar{\gamma}_1}{\bar{I}_1\bar{I}_2} + mgh(\Gamma_1\chi_2 - \Gamma_2\chi_1) \right), \end{aligned}$$

since $\nabla_{\Pi}\Pi = 1$, $\nabla_{\Gamma}\Pi = 0$, $\frac{\partial \Pi}{\partial \alpha} = \frac{\partial \Pi}{\partial l} = 0$, $\Gamma = (\Gamma_1, \Gamma_2, \Gamma_3)$, $\chi = (\chi_1, \chi_2, \chi_3)$, and $\nabla_{\Pi_j}(h_{(\mu,a)} \cdot \bar{\gamma}) = \bar{\gamma}_j/\bar{I}_j$, $j = 1, 2$, $\nabla_{\Pi_3}(h_{(\mu,a)} \cdot \bar{\gamma}) = (\bar{\gamma}_3 - \bar{\gamma}_5)/\bar{I}_3$.

$$\begin{aligned} X_{h_{(\mu,a)} \cdot \bar{\gamma}}(\Gamma) &= \{\Gamma, h_{(\mu,a)} \cdot \bar{\gamma}(A, c, \alpha)\}_-|_{\mathcal{O}_{(\mu,a)} \times \mathbb{R} \times \mathbb{R}} \\ &= -\Pi \cdot (\nabla_{\Pi}\Gamma \times \nabla_{\Pi}(h_{(\mu,a)} \cdot \bar{\gamma})) - \Gamma \cdot (\nabla_{\Pi}\Gamma \times \nabla_{\Gamma}(h_{(\mu,a)} \cdot \bar{\gamma}) - \nabla_{\Pi}(h_{(\mu,a)} \cdot \bar{\gamma}) \times \nabla_{\Gamma}\Gamma) \\ &\quad + \{\Gamma, h_{(\mu,a)} \cdot \bar{\gamma}(A, c, \alpha)\}_{\mathbb{R}}|_{\mathcal{O}_{(\mu,a)} \times \mathbb{R} \times \mathbb{R}} \\ &= \nabla_{\Gamma}\Gamma \cdot (\Gamma \times \nabla_{\Pi}(h_{(\mu,a)} \cdot \bar{\gamma})) + \left(\frac{\partial \Gamma}{\partial \alpha} \frac{\partial (h_{(\mu,a)} \cdot \bar{\gamma})}{\partial l} - \frac{\partial (h_{(\mu,a)} \cdot \bar{\gamma})}{\partial \alpha} \frac{\partial \Gamma}{\partial l} \right) \\ &= (\Gamma_1, \Gamma_2, \Gamma_3) \times \left(\frac{\bar{\gamma}_1}{\bar{I}_1}, \frac{\bar{\gamma}_2}{\bar{I}_2}, \frac{(\bar{\gamma}_3 - \bar{\gamma}_5)}{\bar{I}_3} \right) \\ &= \left(\frac{\bar{I}_2\Gamma_2(\bar{\gamma}_3 - \bar{\gamma}_5) - \bar{I}_3\Gamma_3\bar{\gamma}_2}{\bar{I}_2\bar{I}_3}, \frac{\bar{I}_3\Gamma_3\bar{\gamma}_1 - \bar{I}_1\Gamma_1(\bar{\gamma}_3 - \bar{\gamma}_5)}{\bar{I}_3\bar{I}_1}, \frac{\bar{I}_1\Gamma_1\bar{\gamma}_2 - \bar{I}_2\Gamma_2\bar{\gamma}_1}{\bar{I}_1\bar{I}_2} \right), \end{aligned}$$

since $\nabla_{\Gamma}\Gamma = 1$, $\nabla_{\Pi}\Gamma = 0$, $\frac{\partial \Gamma}{\partial \alpha} = \frac{\partial \Gamma}{\partial l} = 0$.

$$\begin{aligned}
X_{h_{(\mu,a)} \cdot \bar{\gamma}}(\alpha) &= \{\alpha, h_{(\mu,a)} \cdot \bar{\gamma}(A, c, \alpha)\} - |_{\mathcal{O}_{(\mu,a)} \times \mathbb{R} \times \mathbb{R}} \\
&= -\Pi \cdot (\nabla_{\Pi} \alpha \times \nabla_{\Pi}(h_{(\mu,a)} \cdot \bar{\gamma})) - \Gamma \cdot (\nabla_{\Pi} \alpha \times \nabla_{\Gamma}(h_{(\mu,a)} \cdot \bar{\gamma}) - \nabla_{\Pi}(h_{(\mu,a)} \cdot \bar{\gamma}) \times \nabla_{\Gamma} \alpha) \\
&\quad + \{\alpha, h_{(\mu,a)} \cdot \bar{\gamma}(A, c, \alpha)\} |_{\mathcal{O}_{(\mu,a)} \times \mathbb{R} \times \mathbb{R}} \\
&= \left(\frac{\partial \alpha}{\partial \alpha} \frac{\partial (h_{(\mu,a)} \cdot \bar{\gamma})}{\partial l} - \frac{\partial (h_{(\mu,a)} \cdot \bar{\gamma})}{\partial \alpha} \frac{\partial \alpha}{\partial l} \right) = -\frac{(\bar{\gamma}_3 - \bar{\gamma}_5)}{\bar{I}_3} + \frac{\bar{\gamma}_5}{J_3},
\end{aligned}$$

$$\begin{aligned}
X_{h_{(\mu,a)} \cdot \bar{\gamma}}(l) &= \{l, h_{(\mu,a)} \cdot \bar{\gamma}(A, c, \alpha)\} - |_{\mathcal{O}_{(\mu,a)} \times \mathbb{R} \times \mathbb{R}} \\
&= -\Pi \cdot (\nabla_{\Pi} l \times \nabla_{\Pi}(h_{(\mu,a)} \cdot \bar{\gamma})) - \Gamma \cdot (\nabla_{\Pi} l \times \nabla_{\Gamma}(h_{(\mu,a)} \cdot \bar{\gamma}) - \nabla_{\Pi}(h_{(\mu,a)} \cdot \bar{\gamma}) \times \nabla_{\Gamma} l) \\
&\quad + \{l, h_{(\mu,a)} \cdot \bar{\gamma}(A, c, \alpha)\} |_{\mathcal{O}_{(\mu,a)} \times \mathbb{R} \times \mathbb{R}} \\
&= \left(\frac{\partial l}{\partial \alpha} \frac{\partial (h_{(\mu,a)} \cdot \bar{\gamma})}{\partial l} - \frac{\partial (h_{(\mu,a)} \cdot \bar{\gamma})}{\partial \alpha} \frac{\partial l}{\partial l} \right) = 0,
\end{aligned}$$

since $\nabla_{\Pi} \alpha = \nabla_{\Gamma} \alpha = 0$, $\nabla_{\Pi} l = \nabla_{\Gamma} l = 0$, $\frac{\partial \alpha}{\partial l} = \frac{\partial l}{\partial \alpha} = 0$, and $\frac{\partial (h_{(\mu,a)} \cdot \bar{\gamma})}{\partial \alpha} = 0$. If we consider the spacecraft-rotor system with a control torque $u : T^*Q \rightarrow T^*Q$ acting on the rotor, and $u \in \mathbf{J}_Q^{-1}((\mu, a))$ is invariant under the left $SE(3)$ -action, and its reduced control torque $u_{(\mu,a)} : \mathcal{O}_{(\mu,a)} \times \mathbb{R} \times \mathbb{R} \rightarrow \mathcal{O}_{(\mu,a)} \times \mathbb{R} \times \mathbb{R}$ is given by $u_{(\mu,a)}(\Pi, \Gamma, \alpha, l) = \pi_{(\mu,a)}(u(A, c, \Pi, \Gamma, \alpha, l)) = u(A, c, \Pi, \Gamma, \alpha, l) |_{\mathcal{O}_{(\mu,a)} \times \mathbb{R} \times \mathbb{R}}$, where $\pi_{(\mu,a)} : \mathbf{J}_Q^{-1}((\mu, a)) \rightarrow \mathcal{O}_{(\mu,a)} \times \mathbb{R} \times \mathbb{R}$. The dynamical vector field of regular point reduced spacecraft-rotor system $(\mathcal{O}_{(\mu,a)} \times \mathbb{R} \times \mathbb{R}, \tilde{\omega}_{\mathcal{O}_{(\mu,a)} \times \mathbb{R} \times \mathbb{R}}^-, h_{(\mu,a)}, u_{(\mu,a)})$ is given by

$$X_{(\mathcal{O}_{(\mu,a)} \times \mathbb{R} \times \mathbb{R}, \tilde{\omega}_{\mathcal{O}_{(\mu,a)} \times \mathbb{R} \times \mathbb{R}}^-, h_{(\mu,a)}, u_{(\mu,a)})} = X_{h_{(\mu,a)}} + \text{vlift}(u_{(\mu,a)}),$$

where $\text{vlift}(u_{(\mu,a)}) = \text{vlift}(u_{(\mu,a)})X_{h_{(\mu,a)}} \in T(\mathcal{O}_{(\mu,a)} \times \mathbb{R} \times \mathbb{R})$. Assume that

$$\text{vlift}(u_{(\mu,a)} \cdot \bar{\gamma})X_{h_{(\mu,a)}}(A, c, \alpha) = (U_1, U_2, U_3, U_4, U_5, U_6, U_7, U_8)(A, c, \alpha) \in T(\mathcal{O}_{(\mu,a)} \times \mathbb{R} \times \mathbb{R}).$$

Thus, in the case of non-coincident centers of buoyancy and gravity, the Hamilton-Jacobi equations for reduced spacecraft-rotor system with the control torque u acting on the rotor are given by

$$\begin{cases}
\bar{I}_2 \Pi_2 (\bar{\gamma}_3 - \bar{\gamma}_5) - \bar{I}_3 \Pi_3 \bar{\gamma}_2 + mgh \bar{I}_2 \bar{I}_3 (\Gamma_2 \chi_3 - \Gamma_3 \chi_2) + \bar{I}_2 \bar{I}_3 U_1 = 0, \\
\bar{I}_3 \Pi_3 \bar{\gamma}_1 - \bar{I}_1 \Pi_1 (\bar{\gamma}_3 - \bar{\gamma}_5) + mgh \bar{I}_3 \bar{I}_1 (\Gamma_3 \chi_1 - \Gamma_1 \chi_3) + \bar{I}_3 \bar{I}_1 U_2 = 0, \\
\bar{I}_1 \Pi_1 \bar{\gamma}_2 - \bar{I}_2 \Pi_2 \bar{\gamma}_1 + mgh \bar{I}_1 \bar{I}_2 (\Gamma_1 \chi_2 - \Gamma_2 \chi_1) + \bar{I}_1 \bar{I}_2 U_3 = 0, \\
\bar{I}_2 \Gamma_2 (\bar{\gamma}_3 - \bar{\gamma}_5) - \bar{I}_3 \Gamma_3 \bar{\gamma}_2 + \bar{I}_2 \bar{I}_3 U_4 = 0, \\
\bar{I}_3 \Gamma_3 \bar{\gamma}_1 - \bar{I}_1 \Gamma_1 (\bar{\gamma}_3 - \bar{\gamma}_5) + \bar{I}_3 \bar{I}_1 U_5 = 0, \\
\bar{I}_1 \Gamma_1 \bar{\gamma}_2 - \bar{I}_2 \Gamma_2 \bar{\gamma}_1 + \bar{I}_1 \bar{I}_2 U_6 = 0, \\
-J_3 (\bar{\gamma}_3 - \bar{\gamma}_5) + \bar{I}_3 \bar{\gamma}_5 + \bar{I}_3 J_3 U_7 = 0, \\
U_8 = 0.
\end{cases} \tag{4.4}$$

To sum up the above discussion, we have the following proposition.

Theorem 4.4 *In the case of non-coincident centers of buoyancy and gravity, for a point $(\mu, a) \in \mathfrak{se}^*(3)$, the regular value of the momentum map $\mathbf{J}_Q : SE(3) \times \mathfrak{se}^*(3) \times \mathbb{R} \times \mathbb{R} \rightarrow \mathfrak{se}^*(3)$, the regular point reduced system of spacecraft-rotor system with the control torque u acting on the rotor $(T^*Q, SE(3), \omega_Q, H, u)$, where $Q = SE(3) \times S^1$, is the 4-tuple $(\mathcal{O}_{(\mu,a)} \times \mathbb{R} \times \mathbb{R}, \tilde{\omega}_{\mathcal{O}_{(\mu,a)} \times \mathbb{R} \times \mathbb{R}}^-, h_{(\mu,a)}, u_{(\mu,a)})$,*

where $\mathcal{O}_{(\mu,a)} \subset \mathfrak{se}^*(3)$ is the coadjoint orbit, $\tilde{\omega}_{\mathcal{O}_{(\mu,a)} \times \mathbb{R} \times \mathbb{R}}^-$ is orbit symplectic form on $\mathcal{O}_{(\mu,a)} \times \mathbb{R} \times \mathbb{R}$, $h_{(\mu,a)}(\Pi, \Gamma, \theta, l) = H(A, v, \Pi, \Gamma, \theta, l)|_{\mathcal{O}_{(\mu,a)} \times \mathbb{R} \times \mathbb{R}}$, and $u_{(\mu,a)}(\Pi, \Gamma, \theta, l) = u(A, v, \Pi, \Gamma, \theta, l)|_{\mathcal{O}_{(\mu,a)} \times \mathbb{R} \times \mathbb{R}}$. Assume that $\gamma : SE(3) \times S^1 \rightarrow T^*(SE(3) \times S^1)$ is an one-form on $SE(3) \times S^1$, and $\gamma^* : T^*T^*(SE(3) \times S^1) \rightarrow T^*(SE(3) \times S^1)$ is symplectic, and γ is closed with respect to $T\pi_Q : TT^*Q \rightarrow TQ$, and $\text{Im}(\gamma) \subset \mathbf{J}^{-1}(\mu, a)$, and it is $SE(3)_{(\mu,a)}$ -invariant, where $SE(3)_{(\mu,a)}$ is the isotropy subgroup of coadjoint $SE(3)$ -action at the point $(\mu, a) \in \mathfrak{se}^*(3)$, and $\bar{\gamma} = \pi_{(\mu,a)}(\gamma) : SE(3) \times S^1 \rightarrow \mathcal{O}_{(\mu,a)} \times \mathbb{R} \times \mathbb{R}$. Then $\bar{\gamma}$ is a solution of either the Hamilton-Jacobi equation of reduced spacecraft-rotor system given by (4.4), or the equation $X_{h_{\mathcal{O}_{(\mu,a)}} \cdot \bar{\gamma}} + \text{vlift}(u_{(\mu,a)} \cdot \bar{\gamma}) = \tilde{X}^\gamma$, if and only if \tilde{X}^γ and $\tilde{X}_{(\mu,a)}$ are $\bar{\gamma}$ -related, where $\tilde{X}^\gamma = T\pi_Q \cdot \tilde{X} \cdot \gamma$, $\tilde{X} = X_{(T^*Q, SE(3), \omega_Q, H, u)}$, and $\tilde{X}_{(\mu,a)} = X_{(\mathcal{O}_{(\mu,a)} \times \mathbb{R} \times \mathbb{R}, \tilde{\omega}_{\mathcal{O}_{(\mu,a)} \times \mathbb{R} \times \mathbb{R}}^-, h_{(\mu,a)}, u_{(\mu,a)})}$.

Remark 4.5 When the rigid spacecraft does not carry any internal rotor, in this case the configuration space is $Q = G = SE(3)$, the above Hamilton-Jacobi equation (4.4) of reduced spacecraft-rotor system is just the Lie-Poisson Hamilton-Jacobi equation of Marsden-Weinstein reduced Hamiltonian system $(\mathcal{O}_{(\mu,a)}, \omega_{\mathcal{O}_{(\mu,a)}}^-, h_{\mathcal{O}_{(\mu,a)}})$ on the coadjoint orbit of Lie group $SE(3)$, since the symplectic structure on the coadjoint orbit $\mathcal{O}_{(\mu,a)}$ is induced by the (-)-Lie-Poisson brackets on $\mathfrak{se}^*(3)$. See Marsden and Ratiu [21], Ge and Marsden [10], and Wang [32].

The theory of controlled mechanical system is a very important subject, following the theoretical and applied development of geometric mechanics, a lot of important problems about this subject are being explored and studied. In this paper, as an application of the symplectic reduction and Hamilton-Jacobi theory of regular controlled Hamiltonian systems with symmetry, in the cases of coincident and non-coincident centers of buoyancy and gravity, we give explicitly the motion equation and Hamilton-Jacobi equation of reduced spacecraft-rotor system on a symplectic leaf by calculation in detail, respectively, which show the effect on controls in regular symplectic reduction and Hamilton-Jacobi theory. We also note that in [24], the authors study the dynamics of a rigid spacecraft under the influence of gravity torques and give the dynamical equations in a first-order form with a special coefficient matrix. Moreover, we hope to study the stabilization of rigid spacecraft with an internal rotor and to describe the action of controls of the system. On the other hand, if we define a controlled Hamiltonian system on the cotangent bundle T^*Q by using a Poisson structure, see Wang and Zhang in [34] and Ratiu and Wang in [28], and the way of symplectic reduction for regular controlled Hamiltonian system cannot be used, what and how we could do? This is a problem worthy to be considered in detail. In addition, we also note that there have been a lot of beautiful results of reduction theory of Hamiltonian systems in celestial mechanics, hydrodynamics and plasma physics. Thus, it is an important topic to study the application of reduction theory of controlled Hamiltonian systems in celestial mechanics, hydrodynamics and plasma physics. These are our goals in future research.

References

- [1] R. Abraham and J.E. Marsden, Foundations of Mechanics, second ed., Addison-Wesley, Reading, MA, 1978.
- [2] V.I. Arnold, Mathematical Methods of Classical Mechanics, second ed., In: Graduate Texts in Mathematics, Vol. 60, Springer-Verlag, 1989.

- [3] M. Barbero-Liñán, M. de León, J.C. Marrero and D. Martín de Diego and M.C. Muñoz-Lecanda, Kinematic reduction and the Hamilton-Jacobi equation, *J. Geom. Mech.* 4(2012), No.3, 207-237.
- [4] A.M. Bloch, P.S. Krishnaprasad, J.E. Marsden, and G. Sánchez de Alvarez, Stabilization of rigid body dynamics by internal and external torques, *Automatica*, 28(1992), No.4, 745-756.
- [5] A.M. Bloch and N.E. Leonard, Symmetries, conservation laws, and control, In “Geometry, Mechanics and Dynamics, Volume in Honor of the 60th Birthday of J.E. Marsden” (eds. P. Newton, P. Holmes and A. Weinstein), Springer, New York, 2002, 431-460.
- [6] A.M. Bloch, N.E. Leonard and J.E. Marsden, Controlled Lagrangian and the stabilization of mechanical systems I: the first matching theorem, *IEEE Trans, Automatic Control*, 45(2000), 2253–2270.
- [7] A.M. Bloch, N.E. Leonard and J.E. Marsden, Controlled Lagrangians and the stabilization of Euler-Poincaré mechanical systems, *Int. J. Nonlinear and Robust Control*, 11(2001), 191-214.
- [8] J.F. Cariñena, X. Gràcia, G. Marmo, E. Martínez, M. Muñoz-Lecanda and N. Román-Roy: Geometric Hamilton-Jacobi theory, *Int. J. Geom. Methods Mod. Phys.* 3(2006), 1417-1458.
- [9] J.F. Cariñena, X. Gràcia, G. Marmo, E. Martínez, M. Muñoz-Lecanda and N. Román-Roy: Geometric Hamilton-Jacobi theory for nonholonomic dynamical systems, *Int. J. Geom. Methods Mod. Phys.* 7(2010), 431-454.
- [10] Z. Ge and J.E. Marsden, Lie-Poisson integrators and Lie-Poisson Hamilton-Jacobi theory, *Phys. Lett. A*, 133(1988), 134-139.
- [11] D. Iglesias-Ponte, M. de León and D. Martín de Diego: Towards a Hamilton-Jacobi theory for nonholonomic mechanical systems, *J. Phys. A: Math. Theor.* 41(2008), 1-14.
- [12] P.S. Krishnaprasad and J.E. Marsden, Hamiltonian structure and stability for rigid bodies with flexible attachments, *Arch. Rat. Mech. An.* 98(1987), 137–158.
- [13] J-A Lázaro-Camí and J-P Ortega, The stochastic Hamilton-Jacobi equation, *J. Geom. Mech.* 1(2009), 295-315.
- [14] M. de León, J.C. Marrero and D. Martín de Diego, A geometric Hamilton-Jacobi theory for classical field theories, In: *Variations, Geometry and Physics*, Nova Sci. Publ., New York, 2009, 129-140.
- [15] M. de León, J.C. Marrero and D. Martín de Diego, Linear almost Poisson structures and Hamilton-Jacobi equation, Applications to nonholonomic mechanics, *J. Geom. Mech.* 2(2010), 159-198.
- [16] N.E. Leonard and J.E. Marsden, Stability and drift of underwater vehicle dynamics: mechanical systems with rigid motion symmetry, *Physica D*, 105(1997), 130–162.
- [17] P. Libermann, C.M. Marle, *Symplectic Geometry and Analytical Mechanics*, Kluwer Academic Publishers, 1987.
- [18] J.E. Marsden, *Lectures on Mechanics*, In: *London Mathematical Society Lecture Notes Series*, Vol. 174, Cambridge University Press, 1992.

- [19] J.E. Marsden, G. Misiolek, J.P. Ortega, M. Perlmutter, T.S. Ratiu, Hamiltonian Reduction by Stages, In: Lecture Notes in Mathematics, Vol. 1913, Springer, 2007.
- [20] J.E. Marsden, R. Montgomery, T.S. Ratiu, Reduction, Symmetry and Phases in Mechanics, In: Memoirs of the American Mathematical Society, Vol. 88, American Mathematical Society, Providence, Rhode Island, 1990.
- [21] J.E. Marsden, T.S. Ratiu, Introduction to Mechanics and Symmetry, second ed., In: Texts in Applied Mathematics, Vol. 17, Springer-Verlag, New York, 1999.
- [22] J.E. Marsden, H. Wang, Z.X. Zhang, Regular reduction of controlled Hamiltonian system with symplectic structure and symmetry, (2010, arXiv: 1202.3564, To submit to Diff. Geom. Appl.).
- [23] J.E. Marsden and M. West, Discrete mechanics and variational integrators, Acta Numerica, (2001) 357-514.
- [24] C. Mladenova and I. Mladenov, Spacecraft dynamics under the influence of gravity torques, Jour. Theo. Appl. Mech., 38(2008) 3-22.
- [25] H. Nijmeijer and A.J. van der Schaft, Nonlinear Dynamical Control Systems, Springer-Verlag, New York, 1990.
- [26] T. Ohsawa and A.M. Bloch, Nonholonomic Hamilton-Jacobi equation and integrability, J. Geom. Mech., 1 (2009), 461-481.
- [27] J.P. Ortega, T.S. Ratiu, Momentum Maps and Hamiltonian Reduction, In: Progress in Mathematics, Vol. 222, Birkhäuser, 2004.
- [28] T.S. Ratiu and H. Wang, Poisson reduction of controlled Hamiltonian system by controllability distribution, (2012a).
- [29] A.J. van der Schaft, L_2 -Gain and Passivity Techniques in Nonlinear Control, 2nd revised and enlarged edition, Comm. Control Engrg. Ser., Springer-Verlag, London, 2000.
- [30] A.J. van der Schaft, Port-Hamiltonian systems: an introductory survey, In: Proceedings of the International Congress of Mathematicians, Madrid, Spain, (2006), 1339–1365.
- [31] H. Wang, Singular reduction of regular controlled Hamiltonian system with symmetry, (2012b).
- [32] H. Wang, Hamilton-Jacobi theorem for regular reducible Hamiltonian system on a cotangent bundle, (2013a, arXiv: 1303.5840, To submit to Jour. Geom. Phys.).
- [33] H. Wang, Hamilton-Jacobi theorems for regular controlled Hamiltonian system and its reductions, (2013d, arXiv: 1305.3457, To submit to J. Geom. Mech.).
- [34] H. Wang and Z.X. Zhang, Optimal reduction of controlled Hamiltonian system with Poisson structure and symmetry, Jour. Geom. Phys., 62 (5)(2012), 953-975.
- [35] N.M.J. Woodhouse, Geometric Quantization, second ed., Clarendon Press, Oxford, 1992.