

IF B AND $f(B)$ ARE BROWNIAN MOTIONS, THEN f IS AFFINE

MICHAEL R. TEHRANCHI

ABSTRACT. It is shown that if the processes B and $f(B)$ are both Brownian motions (without a random time change) then f must be an affine function. As a by-product of the proof, it is shown that the only functions which are solutions to both the Laplace equation and the eikonal equation are affine.

1. STATEMENT OF RESULTS

Suppose that the process B is a Brownian motion and that the function f is affine. Then the process $f(B)$ is again a Brownian motion. This short note proves the converse: if both B and $f(B)$ are Brownian motions, then f must be affine.

To be precise, we will use the following definition of Brownian motion:

Definition 1. The continuous process $B = (B_t)_{t \geq 0}$ is called a n -dimensional Brownian motion in a filtration $\mathcal{F} = (\mathcal{F}_t)_{t \geq 0}$ iff there exists a n -dimensional vector b and $n \times n$ non-negative definite matrix A such that for all $0 \leq s \leq t$ the conditional distribution of the increment $B_t - B_s$ given \mathcal{F}_s is normal with mean $(t - s)b$ and covariance matrix $(t - s)A$.

A Brownian motion is standard iff $B_0 = 0$, $b = 0$ and A is the $n \times n$ identity matrix.

The main result of this note is this theorem:

Theorem 2. *Suppose B is an n -dimensional Brownian motion in the filtration \mathcal{F} with non-singular diffusion matrix A . Suppose the process $f(B) = (f(B_t))_{t \geq 0}$ is an m -dimensional Brownian motion in the same filtration \mathcal{F} for a measurable function $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$. Then*

$$f(x) = Px + q$$

for some $m \times n$ matrix P and $q \in \mathbb{R}^m$.

There are a number of similar results already in the literature. For instance, Dudley [2] showed that if B is a one-dimensional standard Brownian and $f : \mathbb{R} \rightarrow \mathbb{R}$ is a continuous function such that law of the process $f(B)$ is absolutely continuous with respect to the law of B , then necessarily $f(x) = x$ or $f(x) = -x$. This implies our Theorem 2 in the case $n = 1$.

When B is an n -dimensional standard Brownian motion, Bernard, Campbell & Davie [1] studied functions $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ such that $f(B)$ is a standard Brownian motion up to a random time change. For instance, it is easy to see by the Dambis–Dubins–Schwarz theorem (see, for instance, Section 3.4.B of Karatzas & Shreve’s book [4]) that in the case $m = 1$, it is sufficient that f is harmonic with $f(0) = 0$. In particular, we do not allow time change in our Theorem 2, and hence more structure is imposed on the function f .

Letac & Pradines proved that if $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is such that $f(x + \sqrt{t}Z)$ has the normal distribution for all $x \in \mathbb{R}^n$ and $t \geq 0$, where Z is an n -dimensional standard normal random vector, then f is necessarily equal to an affine function almost everywhere. At first look, it would seem that Letac & Pradines’s result would imply our Theorem 2 since if $f(B)$ is a Brownian motion then $f(B_t)$ is normally distributed for all $t \geq 0$. However, the implication is not entirely obvious, thanks to the following (perhaps surprising) result:

Theorem 3. *Let B be an n -dimensional standard Brownian motion with $n \geq 2$. There exists a continuous non-linear function $g : \mathbb{R}^n \rightarrow \mathbb{R}^n$ such that the random vectors $g(B_t)$ and B_t have the same law for each $t \geq 0$.*

Date: September 17, 2021.

Key words and phrases: Brownian motion, harmonic function, Laplace equation, eikonal equation.

Indeed, the reason that Letac & Pradines's result does not contradict Theorem 3 above is that they impose normality for all $x \in \mathbb{R}^n$, whereas the mean is fixed at $x = B_0 = 0$ in Theorem 3.

The idea of the proof of Theorem 2 is simply an application of the following form of Jensen's inequality: if G is strictly convex and $\int G(x) d\mu = G(\int x d\mu)$ for a probability measure μ , then μ is a point mass. A similar argument yields a related theorem. We will use the notation $\|\cdot\|$ for the Euclidean norm and $\langle \cdot, \cdot \rangle$ the Euclidean inner product on \mathbb{R}^n .

Theorem 4. *Let $D \subseteq \mathbb{R}^n$ be an open, connected set, and suppose $u : D \rightarrow \mathbb{R}$ is a classical solution to both the Laplace equation*

$$\Delta u = 0$$

and the eikonal equation

$$\|\nabla u\| = 1.$$

Then $u(x) = \langle p, x \rangle + q$ for some constants $p \in \mathbb{R}^n$ and $q \in \mathbb{R}$, where $\|p\| = 1$.

Theorem 4 is contained in Lemma 4.1 of the recent paper of Garnica, Palmas & Ruiz-Hernandez [3]. Their proof appeals to methods of differential geometry, while the proof given below only uses Jensen's inequality.

Remark 5. There is little loss in assuming that u is a classical solution to the Laplace equation. Indeed, if u is only assumed to be locally integrable and a solution to the Laplace equation in the sense of distributions, then u is automatically infinitely differentiable, and in particular, a classical solution to the Laplace equation. See Section 9.3 of Lieb & Loss's textbook [6].

2. PROOFS

In this section, we prove the results presented above.

Proof of Theorem 2. Since every component of a vector-valued Brownian motion is a scalar Brownian motion, it is sufficient to consider the case $m = 1$.

First we show that f is smooth. Now since the conditional distribution of $f(B_t)$ given \mathcal{F}_0 is normal, we can conclude that

$$\mathbb{E}[|f(B_t)| | \mathcal{F}_0] < \infty$$

a.s. for all $t \geq 0$. In particular, we have the growth bound

$$(*) \quad x \mapsto f(x)e^{-\epsilon\|x\|^2} \text{ is Lebesgue integrable on } \mathbb{R}^n$$

for all $\epsilon > 0$. Now since $f(B)$ is a Brownian motion, there is a constant $\mu \in \mathbb{R}$ such that

$$\mathbb{E}[f(B_t) | \mathcal{F}_s] = (t - s)\mu + f(B_s),$$

and hence we have the representation

$$f(x) = -\tau\mu + \int f(y)\phi(\tau, x, y)dy$$

where

$$\phi(\tau, x, y) = (2\pi\tau)^{-n/2} \det(A)^{-1/2} \exp\left(-\frac{1}{2\tau}\langle y - b\tau - x, A^{-1}(y - b\tau - x) \rangle\right)$$

is the Brownian transition density. But by the boundedness property (*) and the smoothness of $x \mapsto \phi(t, x, y)$ combined with the dominated convergence theorem, the function f is differentiable. Furthermore, its gradient ∇f has the representation

$$\nabla f(x) = \int \nabla f(y)\phi(\tau, x, y)dy$$

and also satisfies the boundedness property (*). By iterating this argument, we see that f is infinitely differentiable.

Now we show that f must satisfy an eikonal equation. Note that Itô's formula says

$$df(B_t) = \langle \nabla f(B_t), dB_t \rangle + \frac{1}{2}\Delta f(B_t)dt.$$

Since $f(B)$ is a Brownian motion, the quadratic variation is

$$[f(B)]_t = \int_0^t \|\nabla f(B_s)\|^2 ds = \sigma^2 t$$

for some constant $\sigma \geq 0$. Hence ∇f is a solution of the eikonal equation

$$\|\nabla f\| = \sigma.$$

almost everywhere. But since f is smooth, it solves the eikonal equation everywhere. Now note that

$$\sigma^2 = \|\nabla f(x)\|^2 = \int \|\nabla f(y)\|^2 \phi(\tau, x, y) dy.$$

Since the squared Euclidean norm is strictly convex, Jensen's inequality says that for every x there exists a vector $p_x \in \mathbb{R}^n$, possibly depending on x , such that $\nabla f(y) = p_x$ a.e $y \in \mathbb{R}^n$. Since ∇f is continuous, we must have $\nabla f(y) = p$ for all y and for some constant vector p . Hence $f(y) = \langle p, y \rangle + q$ as claimed. \square

We now proceed to the proof of the Theorem 4. It follows the same pattern, but it differs in a few details which we spell out for completeness.

Proof of Theorem 4. We will show that there is a unit vector p such that $\nabla u(x) = p$ everywhere in D . Below we will use the notation $B = \{x \in \mathbb{R}^n : \|x\| < 1\}$ to denote the open unit ball in \mathbb{R}^n , and hence $x + rB$ denotes the ball of radius $r \geq 0$ centred at the point $x \in \mathbb{R}^n$.

Since u is harmonic, it is well known (again, see Section 9.3 of [6]) that u has the mean-value property: for every constant $r > 0$ such that $x + rB \subseteq D$ we have

$$u(x) = \frac{1}{r^n V} \int_{rB} u(x+y) dy$$

where

$$V = \frac{\pi^{n/2}}{\Gamma(n/2)}$$

denotes the Lebesgue measure of the unit ball B . Since u is continuously differentiable in D , the gradient ∇u is bounded on compact sets, so the dominated convergence theorem allows us to differentiate both sides of the above equation, yielding

$$\nabla u(x) = \frac{1}{r^n V} \int_{rB} \nabla u(x+y) dy$$

Now for each $x \in \mathbb{R}^n$, note that

$$1 = \|\nabla u(x)\|^2 = \frac{1}{r^n V} \int_{rB} \|\nabla u(x+y)\|^2 dy.$$

Again, since the squared Euclidean norm is strictly convex, Jensen's inequality says that there is a vector p_x , possibly depending on x , such that $\nabla u(z) = p_x$ a.e $z \in x + rB$ and $\|p_x\| = 1$. Since ∇u is continuous, we must have $\nabla u(z) = p_x$ for all z such that $\|x - z\| \leq r$.

Now fix two points x and x' in D . Since D is open and connected, there exists a path $C \subseteq D$ connecting them. Hence there exists a finite number of points $x = x_1, \dots, x_N = x' \in D$ and radii $r_1, \dots, r_N > 0$ such that $\{x_i + r_i B\}_{i=1}^N$ is a cover of the compact set $C \subseteq D$. In particular, the $p_x = p_{x'}$ and hence ∇u is constant on D as claimed. \square

Finally, we construct an example of the function g claimed to exist in Theorem 3.

Proof of Theorem 3. Let $S^{n-1} = \{x \in \mathbb{R}^n : \|x\| = 1\}$ be the unit $(n-1)$ -dimensional sphere and let λ be the uniform probability measure on S^{n-1} . Let $h : S^{n-1} \rightarrow S^{n-1}$ be a continuous λ -preserving transformation. Finally, let $g(0) = 0$ and

$$g(x) = \|x\| h\left(\frac{x}{\|x\|}\right)$$

when $x \neq 0$.

Fix a bounded and measurable function $\varphi : \mathbb{R}^n \rightarrow \mathbb{R}$ and $t \geq 0$. Using the assumption that the transformation h preserves the measure λ , we have

$$\begin{aligned} \mathbb{E}[\varphi \circ g(B_t)] &= \int_{\mathbb{R}^n} \varphi \left[\sqrt{t} \|x\| h \left(\frac{x}{\|x\|} \right) \right] \frac{e^{-\|x\|^2/2}}{(2\pi)^{n/2}} dx \\ &= \int_0^\infty \int_{S^{n-1}} \varphi \left[\sqrt{tr} h(u) \right] \frac{r^{n-1} e^{-r^2/2}}{2^{n/2-1} \Gamma(n/2)} \lambda(du) dr \\ &= \int_0^\infty \int_{S^{n-1}} \varphi(\sqrt{tr}u) \frac{r^{n-1} e^{-r^2/2}}{2^{n/2-1} \Gamma(n/2)} \lambda(du) dr \\ &= \int_{\mathbb{R}^n} \varphi(\sqrt{t}x) \frac{e^{-\|x\|^2/2}}{(2\pi)^{n/2}} dx \\ &= \mathbb{E}[\varphi(B_t)] \end{aligned}$$

where we have used the polar coordinates $x = ru$ with $r \geq 0$ and $u \in S^{n-1}$. Hence $g(B_t)$ and B_t have the same law for each $t \geq 0$.

To show that there exist at least one function h which is non-linear, it is sufficient to consider the case $n = 2$ since we may restrict attention to the first two coordinates of B . Now let $h : S^1 \rightarrow S^1$ be defined by $h(\cos(\theta), \sin(\theta)) = (\cos(2\theta), \sin(2\theta))$. It is well-known that this transformation h is measure preserving. Explicitly, the function g in this case is

$$g(x_1, x_2) = \left(\frac{x_1^2 - x_2^2}{\sqrt{x_1^2 + x_2^2}}, \frac{2x_1x_2}{\sqrt{x_1^2 + x_2^2}} \right)$$

when $x \neq 0$. □

Acknowledgment. I would like to thank Pat Fitzsimmons for feedback on an early draft of this note, and in particular, for alerting me to several papers in the literature. I would also like to thank Chris Rogers and Arun Thillaisundaram for interesting discussions of this problem, and the members of the Laboratoire de Probabilités et Modèles Aléatoires at Université Paris 6 for their hospitality during much of the writing of this note. Finally, I would like to thank an anonymous referee for helpful comments.

REFERENCES

- [1] A. Bernard, E.A. Campbell and A.M. Davie. Brownian motion and generalized analytic and inner functions. *Annales de l'Institut Fourier (Grenoble)* **29**(1): xvi, 207-228 (1979)
- [2] R. Dudley. Non-linear equivalence transformations of Brownian motion. *Zeitschrift für Wahrscheinlichkeitstheorie und Verwandte Gebiete* **20**: 249-258 (1971)
- [3] E. Garnica, O. Palmas and G. Ruiz-Hernandez. Classification of constant angle hypersurfaces in warped products via eikonal functions. *Sociedad Matemática Mexicana. Boletín. Tercera Serie* **18**(1): 29-41 (2012)
- [4] I. Karatzas and S. Shreve. *Brownian Motion and Stochastic Calculus*. Graduate Texts in Mathematics **113**. Springer (1991)
- [5] G. Letac and J. Pradines. Seules les affinités préservent les lois normales. *Comptes Rendus Hebdomadaires des Séances de l'Académie des Sciences, Série A*. **286**(8): 399-402 (1978)
- [6] E. Lieb and M. Loss. *Analysis*. Graduate Studies in Mathematics **14**. American Mathematics Association (2001)