

Comparison of modules of wild type and mutant huntingtin and TP53 protein interaction networks : implications in biological processes and functions

Mahashweta Basu¹, Nitai P. Bhattacharyya², Pradeep K. Mohanty^{1*}

1 Theoretical Condensed Matter Physics Division,
Saha Institute of Nuclear Physics, 1/AF Bidhan Nagar, Kolkata 700064,
India

2 Crystallography and Molecular Biology Division,
Saha Institute of Nuclear Physics, 1/AF Bidhan Nagar, Kolkata 700064,
India

* E-mail: pk.mohanty@saha.ac.in

Abstract

Disease-causing mutations usually change the interacting partners of mutant proteins. In this article, we propose that the biological consequences of mutation are directly related to the alteration of corresponding protein protein interaction networks (PPIN). Mutation of Huntingtin (HTT) which causes Huntington's disease (HD) and mutations to TP53 which is associated with different cancers are studied as two example cases. We construct the PPIN of wild type and mutant proteins separately and identify the structural modules of each of the networks. The functional role of these modules are then assessed by Gene Ontology (GO) enrichment analysis for biological processes (BPs). We find that a large number of significantly enriched ($p < 0.0001$) GO terms in mutant PPIN were absent in the wild type PPIN indicating the gain of BPs due to mutation. Similarly some of the GO terms enriched in wild type PPIN cease to exist in the modules of mutant PPIN, representing the loss. GO terms common in modules of mutant and wild type networks indicate both loss and gain of BPs. We further assign relevant biological function(s) to each module by classifying the enriched GO terms associated with it. It turns out that most of these biological functions in HTT networks are already known to be altered in HD and those of TP53 networks are altered in cancers. We argue that gain of BPs, and the corresponding biological functions, are due to new interacting partners acquired by mutant proteins. The methodology we adopt here could be applied to genetic diseases where mutations alter the ability of the protein to interact with other proteins.

Introduction

Cellular functions are carried out by proteins interacting with other proteins and macromolecules like DNA, RNA, etc. It is believed [1] that the modular organization of cellular functions are related to the underlying modular structure of the protein protein interaction network (PPIN). Understanding PPIN would elucidate how such interactions execute

basic functions in cells and may explain the abnormalities arising from mutations in genes. In particular, mutation at the binding site of a protein may lead to loss of its ability to function together with existing interacting partner(s). On the other hand, mutation may also create regions where new protein partners can bind. Therefore, loss or gain of interaction due to mutation may contribute to causation, progression or modulation of disease. It has been reported recently [2] that out of 119 mutations in 65 distinct diseases, 95 mutations result in loss of function (LOF), 17 mutations result in gain of function (GOF) and 4 mutations changes the preferences for interaction. Based on this experimentally validated data, it has been predicted that 1428 mutations might be related to interaction defect. Using the structural information at atomic levels either through crystallography or homology modeling, it has been shown that 21,716 mutations in 624 genes either alter amino acid sequences or produce truncated proteins. Among 12,059 mutations that alter amino acid sequences, 7833 mutations are located in the interface of interaction with other proteins. Such mutations at interfaces of interactions may disrupt or enhance the interactions with the partners. This study also emphasizes the role of loss or gain of interactions of mutant proteins in human diseases. However, for such analysis, it is necessary to have structural information at atomic levels, which may be achieved if 3-dimensional structures of the proteins or their homologs are known. But, for the most of the protein protein interactions such information is not available [3]. Moreover, very little is known about the role of such altered interactions in corresponding pathological conditions. It remains a challenge to relate genetic mutation data to PPIN and to understand molecular cause of disease. In the present communication, we probe whether gain or loss of interactions of mutant Huntingtin protein (HTT) that causes Huntingtons disease (HD) can explain functional abnormalities observed in HD. We have also used the same approach to find how loss or gain of interactions of mutant TP53 in cancers may result in alterations of functions.

Analysis and Results

Mutation in HTT protein

Huntingtons disease (OMIM ID: 143100) is a rare autosomal dominant progressive degenerative neurological disease caused by expansion of normally polymorphic CAG repeats beyond 36 at the exon1 of the gene Huntingtin (HTT) [4]. Over the years, various cellular processes/conditions like excitotoxicity, oxidative stress, mitochondrial dysfunction, endoplasmic reticulum stress, axonal transport, ubiquitin proteasome system, autophagy, transcriptional deregulation and apoptosis have been implicated in HD pathology [5, 6]. Even though GOF was inferred initially from the autosomal dominant nature of transmittance of the disease, the underlying molecular details still remain largely unknown. Inverse correlations between age at onset and number of CAG repeat beyond 36 in HTT gene, increased aggregates of mutant HTT (mHTT) and apoptosis, correlation of CAG

repeat numbers in HTT gene with levels of ATP/ADP and altered expression of few genes [4,7–10] suggest toxic GOF of mutant protein that disrupts normal cellular functions and causes neuronal death. Mutant HTT preferentially interacts with DNA sequences, alters conformation of DNA facilitating binding of other transcription factors to the specific sequences and modulates transcription of genes. This result also indicates a dominant GOF of mHTT [11]. Wild type HTT (wHTT) is known to be involved in protection of apoptosis [12–15], regulation of gene expression [16,17], mitosis and neurogenesis [18], neuronal development [19] and maintenance of body weight [20]; all these processes are altered in HD [5,6]. These results indicate that loss of one of the alleles in HD could contribute to increased apoptosis and altered gene expressions observed in HD. LOF of wild type protein may thus contribute, at least partially, to HD pathology [21]. There are also several experimental evidences available against simple LOF(s) of wild type HTT [22–25].

Construction of HTT-interacting protein network

We have collected the HTT interacting proteins from published data and find that 17 proteins preferentially interact with wHTT, while 37 proteins are either identified in aggregates of mHTT only or interact preferentially with mHTT (the references for each of the observations are provided in [Dataset S1 \(sheet 1\)](#) and in [Supporting Text \(Text 1\)](#)). These 17 and 37 proteins are referred to as the primary interactors of wHTT and mHTT respectively. Next, we assimilate interacting partners of these primary interactors from BioGrid (Version 3.1.88, May 2012), a public database that contains genetic and protein protein interaction data for humans and other organisms [26]. In the present study, we have considered both physical and genetic interactions (refer to the section ‘Robustness analysis’ for details). It turns out that there are 288 secondary interactors of wHTT (proteins which interact with the 17 primary interactors), whereas there are 1504 secondary proteins which interact with 37 primary interactors of mHTT. The PPIN of wHTT interacting proteins is then constructed by considering all these 306 proteins (wHTT + 17 primary + 288 secondary interactors of wHTT) as nodes of the network; two nodes are connected if corresponding pair of proteins are found to be interacting partners of each other in BioGrid. Altogether there are 1397 interactions in wHTT network which are listed in [Dataset S1 \(sheet 2\)](#). Similarly the PPIN of mHTT is constructed with 1542 nodes (mHTT + 37 primary + 1504 secondary interactors of mHTT) which has 13142 interactions from BioGrid ([Dataset S1 \(sheet 3\)](#)). We have used Cytoscape [27] for visual presentation of the wHTT and mHTT networks, which are shown in [Supporting Text \(Fig. S1\)](#). Both the networks are densely interconnected and the nodes are too tangled there to find any apparent or obvious modular structures.

Characteristics of networks

A quantifiable description of these networks can be obtained by using graph theory, which provides several measures for comparison and characterization of complex networks. The




Figure 1. Construction and modularization of wild type and mutant HTT networks.

(a) Proteins involved in the wHTT and mHTT networks: wHTT (mHTT) protein (red square) has 17 (37) primary and 288 (1504) secondary interactors, shown schematically as the inner and outer circles. Of the 17 primary interactors of wHTT, 8 proteins (deep green) become secondary interactors of mHTT. Among the 288 secondary interactors of wHTT, 107 (shaded) proteins remain as the secondary interactor of mHTT whereas 10 proteins (deep blue) becomes the primary interactors of mHTT. (b) and (c) Modules of the wHTT and mHTT networks from NGM algorithm, which yields 7 ($W1, W2, \dots, W7$) and 8 modules ($M1, M2, \dots, M8$) respectively are shown along with the relevant biological functions (obtained GO term enrichment analysis from GeneCodis3). Significant functions associated with the modules are also shown. Details of the GO terms are shown in [Table S2](#) and [Table S3](#) of the [Supporting Text](#), respectively for wHTT and mHTT.

most elementary characteristic of a node is its degree, k , which represents the number of other nodes (proteins) it is connected with. The degree distribution, $P(k)$, gives the probability that a randomly selected node has exactly k links. We find that both the wild and mutant PPINs follow a power law degree distribution, $P(k) \sim k^{-\gamma}$ ([Fig. S3](#) in [Supporting Text](#)) with exponents $\gamma = 1.99, 1.95$ and average degrees $\langle k \rangle = 9.13, 17.05$ respectively. Another important quantity is the clustering coefficient which characterizes how connected are the neighbors of a given node. It is observed that the average clustering coefficient $C = 0.361$ for mHTT network is lower compared to $C = 0.436$ for wHTT PPIN. This indicates that, the former network is less compact and the interacting partners of the proteins are poorly connected among themselves. We have also calculated the average shortest path length L , and the network diameter D (listed in [Supporting Text](#) ([Table. S2](#))), which describe the structural properties of the network. The detailed definitions of C , D and L along with their evaluation procedure is illustrated in [Supporting Text](#) ([Text 2](#)).

Gain and Loss of interactions due to mutation

A closer look at PPINs of wHTT and mHTT reveals that among the 17 primary interactors of wHTT, 8 proteins still appear in PPIN of mHTT as secondary interactors, *i.e.* they interact with some of the primary interactors of mHTT. Again, among 288 secondary interactors of wHTT, 107 proteins are secondary interactors of mHTT, 10 proteins interact directly with mHTT and the rest 171 proteins do not take part in PPIN of mHTT (see Fig. 1(a)). Evidently, the mutant HTT network has gained several new interactions, 27 proteins as primary interactors and 1389 proteins as secondary interactors. This result is shown schematically in Fig. 1(a) and the detailed list of these proteins is given in [Supporting Text \(Text 1 and Table S1\)](#). Since mutation of HTT has changed the PPIN substantially one expects a significant change in its functions.

Modules of wHTT and mHTT networks

There are several methods for obtaining natural modules of a network (or partitions of a graph) [28]. We adopt Newman-Girvans modularization (NGM) algorithm [29], a commonly used method, to detect the modules of wHTT and mHTT networks. This algorithm partitions the network in a way that the intra-module connections between nodes are maximized in comparison to the inter-module connections. To find the modules, Newman and Girvan [30] proposed a score called modularity Q for every possible partition of a network; the maximum value of Q corresponds to the best partition. The details of the NGM algorithm for maximization of Q is described in [Supporting Text \(Text 2\)](#). The NGM algorithm modularizes the PPIN of wHTT into 7 modules of sizes (18, 66, 79, 18, 82, 8 and 35) (see [Table S2 in Supporting Text](#)), with modularity $Q = 0.415$, whereas PPIN of mHTT is partitioned into 8 modules of sizes (643, 3, 377, 2, 485, 7, 22 and 3) with $Q = 0.302$. Modules of wHTT and mHTT networks are denoted by W and M respectively. Figures 1(b) and (c) represent the modularized networks; all proteins belonging to a given module are shown in same color. Clearly, the mHTT network is visibly more complex than that of wHTT, which is consistent with the fact that it has a lower Q value [31].

Similarity between the modules

Once the wild type and mutant networks are modularized, it is important to ask how similar is a module of wild type network with that of mutant network, in terms of their protein constituents. Mutant and wild type HTT networks have 125 proteins common between them. After both the networks are modularized, these common proteins are distributed among the pair of wHTT- mHTT modules. For example, the module $M5$ (485 proteins) has 49 proteins in common with $W5$ (485 proteins), whereas it has only one common protein in $W2$ (out of 66 proteins) and two common proteins in $W1$ (18

proteins). The detailed distribution of common proteins among wild and mutant modules of HTT are shown in Fig. 2.




Figure 2. Similarity between modules of wHTT and mHTT networks. The figure describes pictorially the closeness between the modules of wHTT and mHTT PPIN; the modules having common protein or common GO terms are joined with edges (numerical value written on the edge as : common proteins).

To calculate the similarity among modules, first we construct a unique set of proteins from combining the proteins involved in the wild and mutant networks. This set consists of 1723 proteins in case of HTT. Now every module of wHTT and mHTT are considered as a unique 1723 dimensional vector as follows. Each protein is identified with a specific position in the vector; presence (or absence) of a specific protein in a module, say w , is mapped on to a corresponding vector \vec{R}_w by inserting 1 (or 0) at respective position. A similarity measure between a pair of modules w and m is well represented by the angle $\theta(w, m)$ between the corresponding vectors \vec{R}_w and \vec{R}_m ,

$$\theta(w, m) = \cos^{-1} \frac{\vec{R}_w \cdot \vec{R}_m}{|\vec{R}_w| |\vec{R}_m|}$$

It is rather simpler to use $\cos(\theta(w, m))$ as the similarity measure as *cosine* function is monotonic in the range $(0, \pi)$. It is easy to see that if the modules have N_w and N_m proteins individually and N_{wm} protein in common, the similarity measure is

$$\sigma(w, m) \equiv \cos(\theta(w, m)) = N_{wm} / \sqrt{N_w N_m}. \quad (1)$$

Clearly $\sigma(w, m)$ varies in the range $(0, 1)$ with maximum value 1 corresponding to the fact that the modules are identical, *i.e.* they have same set of proteins.

In Fig. 2, we represent the similarity among modules of mHTT and wHTT as a bipartite network with links having thickness proportional to σ_{wm} . The thickest link between $M5$ and $W5$ indicates that these modules are significantly similar. For examples, the module $W5$ has 82 proteins and $M5$ has 485 proteins; 49 proteins are common among the proteins in these 2 modules; thus the protein similarity index for $W5$ - $M5$ pair is $\sigma(W5, M5) = 0.246$. Similarly among 66 proteins in $W2$ and 643 proteins in $M1$, 26 proteins are common (corresponding $\sigma(W2, M1) = 0.126$).

Enrichment of GO terms for biological process

It has been observed that the proteins identified in a particular complex are involved in similar functions [32]. From network perspective, these complexes are represented by modules and they appear as distinct group of nodes which are highly interconnected with each other but have only a few connections with the nodes outside of the module. It is important to ask, if such a structural partition relates to any functional enrichment. Among many bioinformatics tools available for such analysis [34] we utilize GeneCodis3 [33] (explained in [Supporting Text \(Text 3\)](#)) to obtain the possible Biological processes enriched by the proteins in a given module. Given a query set of proteins GeneCodis3 provides the enriched biological process, molecular functions, and cellular components as defined by the Gene ontology. Biological process in Gene ontology is described as a series of events carried out by one or more ordered assemblies of molecular functions [35]. The proteins in each module are used as input to GeneCodis3 [33] and significantly enriched GO terms for BPs obtained using p -values calculated through Hypergeometric analysis corrected for false discovery rate (FDR). Results of enrichment analyses for 7 modules of wHTT and 8 modules of mHTT network are shown in [Datasets S2](#) and [S3](#) respectively.

Since many proteins are known to be involved in a particular BP, and a given protein may also contribute to multiple BPs, it is likely that proteins in different modules in wHTT and mHTT network participate in a specific BP due to either overlap in proteins or BPs. To identify the overlaps of BPs between modules in wHTT and mHTT networks, we separately identify the common GO terms between the wHTT and the mHTT modules. It is evident from [Dataset S4 \(sheet 2\)](#) that 390 unique GO terms are being enriched ($p < 0.0001$) due to proteins in modules of mHTT network, while 129 GO terms are enriched with proteins in the modules of wHTT network ([Dataset S2 \(sheet 1\)](#)). Among the GO terms present in wHTT and mHTT network, 65 are common. As a result due to mutation, 325 GO terms are gained by mHTT and 64 GO terms are lost by wHTT. The common 65 GO terms represents both gain and loss.

For convenience, we clubbed the the GO terms in a given module to broadly assign one or more appropriate biological function(s). For example, GO:0010506 (regulation of autophagy), GO:0016559 (peroxisome fission), GO:0031929 (TOR signaling cascade), GO:0000045 (autophagic vacuole assembly), GO:0006897 (endocytosis) in module *M1* are bought under a single biological function “Autophagy”. Similarly in module *W4* GO:0043507 (positive regulation of JUN kinase activity), GO:0072383 (plus-end-directed vesicle transport along microtubule), GO:0046330 (positive regulation of JNK cascade), GO:0046328 (regulation of JNK cascade) are clubbed under “Signaling”. The assigned biological functions for modules of wHTT and mHTT are shown in Fig. 1(b) and (c) (details are given in [Dataset S4](#)).

Gain and loss of biological process in HTT networks

Comparison of enriched BPs in the modules of wHTT and mHTT reveal that the mHTT

network has acquired several new BPs which were absent in wHTT, indicating gain of biological processes. Similarly enriched BPs of wHTT which are not present in mHTT are lost. Hence biological functions carried out by the BPs which are gained or lost in mHTT networks may result in functional gain or loss due to mutation in HTT.

Gain of biological process : The unique GO terms enriched in the modules of mHTT networks are listed in [Dataset S4 \(sheet 2\)](#) and in [Supporting Text \(Table S3\)](#). The GO terms in module *M1* are related to cell cycle (4 GO terms), signaling (30), transcription processes and regulation (5), apoptosis (11), DNA damage and repair (6), Immunological (7), protein folding (7), autophagy (5), translation (3), metabolism (1), development and differentiation (4), cell migration and shape (4), proteasomal degradation (14), Protein complex/membrane assembly/stabilization (9) and others (4). It is known that many of these processes are involved in HD pathogenesis [36]. In *M3*, the enriched GO terms are assigned to DNA repair (17), Transcription processes and regulation (5), DNA replication (12), cell cycle (12) and others (5). Note that, it has been shown recently that DNA repair, replication and cell cycle are involved in HD. In fact, activation of DNA synthesis and cell cycle increase apoptosis in terminally differentiated neuronal cells, instead of increasing cell division [37, 38]. Besides, recent studies have explored the role of DNA repair in neurodegenerative disease [39] and show that interaction of mHTT with Ku70/ XRCC6 impairs repair activity [40]. A large number of GO terms related to development and differentiation (57 GO term), transcription process and regulation (31), cell cycle (5), DNA damage and repair (4), Carbohydrate/Glucose transport/metabolism (4), Cell growth (7), signaling (31) and others (6) are enriched in module *M5*. The role of development and differentiation in HD is not clear. However recent studies in HD [19, 41] indicate that neurogenesis is possibly altered and differentiation/development could be defective. Deregulation of transcription is considered to be one of the most important abnormalities in HD [42]. GO terms related to differentiation are also enriched with proteins in module *M6*, although the terms are distinct from that in module *M5*. All 4 GO terms enriched in *M7* are related to transcription by RNA polymerase III. It is known that both tRNA and some miRNAs [43] are synthesized by RNA polymerase III, however their role in HD is unknown. Thus it is evident that the the protein interactions gained in mHTT network result in enrichment of the biological processes in its modules.

Loss of biological process : The unique GO terms enriched in the modules of wHTT which are absent in the modules of mHTT network represent the loss of functions due to mutation in HTT protein. The 5 GO terms in *W1* include gene silencing, micro RNA processing and translational regulation. The GO terms relating to proteasomal degradation (1 GO term), cell cycle (1), apoptosis (1) and circadian rhythm (1) are present in *W2*. Similarly, signaling (17 GO terms), synaptic transmission, neuronal activities (12) transport (ion/sugar) (5) and others (4) are associated with module *W3*; glucose/carbohydrate transport and metabolism (5), cell cycle (3) and protein/transmembrane transport (4) with *W6*. In *W7* only one GO term describing transcription processes and regulation is enriched. The GO terms and the associated BPs that are lost due to mutation are

provided in [Dataset S4 \(sheet 1\)](#) and in [Supporting Text \(Table S3\)](#) respectively.

We have clubbed the relevant GO terms to represent signaling, transcription process and regulation, apoptosis, cell cycle etc. (refer to [Table S3 of Supporting Text](#)). For example, GO terms (GO:0000088) and (GO:0000236, GO:0000087, GO:0007091) which are enriched in *W2* and *W6* respectively relates to cell cycle. Similarly the 21 GO terms which are enriched in *M1(4)*, *M3(12)*, and *M5(5)* ([Dataset S4 \(sheet 2\)](#)) are also associated to cell cycle. Although cell cycle is enriched in both wHTT and mHTT modules, no GO terms are common among them. Thus, the loss of interaction with wHTT may result in loss of above 4 GO terms in wild type network resulting in LOF, whereas the gain of interaction with mHTT may be associated with gain of these 21 GO terms relating to GOF of cell cycle.

It is interesting to note (from [Table S3 of Supporting Text](#)) that the GO terms related to DNA replication, protein folding, autophagy, cell growth are only observed in the modules of mHTT networks. So these processes are gained due to new interaction with mHTT. Similarly, GO terms related to gene silencing/ microRNA processing/ translation, transport (ion/protein/sugar etc) are observed in wHTT network only. Therefore, loss of interaction with wHTT may result in the loss of these BPs in HD.

Both loss and gain of biological process : Modules in wHTT and mHTT networks have several proteins or GO terms common among them, which indicate loss as well as gain of functions and support the notion that both loss and gain may occur due to mutation in HTT [21]. For example, modules (*W1, W5, W7*) and (*M3, M5, M6*) have 17 enriched GO terms related to transcription processes and regulation. Similarly, modules (*W2, W5*) and (*M1, M3, M5*) share 12 enriched common GO terms related to apoptosis and 4 common GO terms relating to cell cycle. Thus, the general function of transcription and apoptosis could arise from loss as well as gain of interactions of mHTT protein. The details of the functions associated with the 65 GO terms (common between wHTT and mHTT) are presented in [Dataset S4 \(sheet 3\)](#) and in [Supporting Text \(Table S3\)](#), they correspond to the gain and loss of functions in the HD.

From the above analysis we observe that most of the functions that are enriched in the modules of wHTT and mHTT networks are altered in the pathogenesis of HD. The post transcriptional regulation of genes, associated with module *W1* of wHTT network, can be related to negative regulation of gene expression by the non-coding RNAs like micro RNAs, which are well documented [44]. Role of apoptosis [5,36], synaptic transmission [45], JNK pathway [46], transcription deregulation [42], glucose transport [47,48], estrogen [49] and various types of epigenetic changes including histone modifications in different neurological diseases [50] in HD pathogenesis have also been reported.

In summary, many new BPs (GO terms) appear in the mHTT network and some of the BPs present in wHTT network are lost; a few are found to be common between modules of wHTT and mHTT. As a result some biological functions involving the enriched GO terms are gained by mHTT and a few are lost from the modules of wHTT. This provides molecular mechanism of the gain and/or loss of functions observed in HD pathogenesis.

Mutation in TP53 protein

TP53 protein, initially identified as an oncogene, is now established as a tumor suppressor gene which participates in diverse cellular functions like transcription regulation, DNA repair, apoptosis, and genome stability, and many others. Mutation to TP53 is identified in more than 50% of the tumors. It is evident from COSMIC database [51] that R175H, R273H and R248W mutations of TP53 are the most prevalent ones. Since TP53 is a tumor suppressor gene, it is expected that its mutations might result in the LOF of the wild type protein. Some mutations of TP53 are also known to attain new function(s) [52,53]. For example, exogenous expression of mutant TP53 (R273H and others) in mouse cells devoid of endogenous TP53 results in several cellular phenotypes of cancers [54–56]. To understand the underlying molecular mechanism of GOF of mutant TP53, it was recently shown [57] that nardilysin (NRD1) protein, which does not interact with wild type TP53 but interacts only with mutant TP53 (R273H), may contribute to the metastatic properties of this mutant protein.

PPIN of wTP53 and R273H mutant TP53 (mTP53)

In a recent study [57], it has been shown that 17 proteins preferentially interact with the wild type TP53 (wTP53) and 30 other proteins interact exclusively with mutant TP53 (mTP53). To construct the protein interaction networks we take these primary interacting proteins of wTP53 and mTP53 and consider their interacting partners existing in BioGrid database [26]. The detailed protein interaction data are given in the [Dataset S5](#). The PPIN is constructed separately for wTP53 and mTP53, as described for HTT. It turns out that wTP53 has 601 secondary interactors whereas mTP53 has only 547. Thus the PPIN of wTP53 and mTP53 are constructed taking 619 proteins (wTP53 + 17 primary + 601 secondary) and 578 proteins (mTP53 + 30 primary + 547 secondary) respectively. Both the networks (shown in [Supporting Text \(Fig. S2\)](#)) are found to be densely packed with similar structural properties. Their degree distributions are scale free ($P(k) \sim k^{-\gamma}$) with the exponents $\gamma = 2.04$ (wTP53) and 1.89 (mTP53) ([Fig. S3 in Supporting Text](#)) and average degree $\langle k \rangle = 12.07, 12.29$. The other network properties, like the average clustering coefficient $C = 0.452, 0.406$, the diameter of the networks $D = 4, 4$ are also comparable (listed in [Supporting Text \(Table S2\)](#)).

The change in interactions and the interacting partners due to mutation of TP53 is shown schematically in [Fig. 3\(a\)](#). Of 17 primary interactors of wTP53, only 5 proteins remain involved in mutant network as secondary interactors of mTP53 and the remaining 12 do not interact with mTP53. Among the 601 secondary interactors of wTP53, 111 proteins remain as a secondary interactor of mTP53 and 7 of them interact directly, *i.e.* 7 secondary interactors of wTP53 become primary interactors of mTP53. Lists of these proteins are given in [Supporting Text \(Text 1 and Table S1\)](#).




Figure 3. Construction and modularization of wild type and mutant TP53 networks.

(a) Proteins in wTP53 and mTP53 networks : wTP53 (mTP53) protein (red square) has 17 (30) primary and 601 (547) secondary interactors, represented by the inner and outer circles. Only 5 (7) primary (secondary) proteins of wTP53 interact with mTP53 as secondary (primary) interactors. Again 111 secondary proteins of wTP53 remain as secondary interactors of mTP53. (b) and (c) shows the modules of wTP53 and mTP53 network along with few plausible candidate BPs. Details of the GO terms are shown for wTP53 and mTP53 respectively in [Table S6](#) and [Table S7](#) of the [Supporting Text](#).

Modules of wTP53 and mTP53 networks

In order to identify the modules of the wTP53 and mTP53 networks, we use NGM algorithm [29]. It turns out that PPIN of wTP53 is modularized into 4 modules of size 204, 151, 183 and 81, whereas mTP53 network gives 5 modules of size 193, 127, 25, 111 and 122. The corresponding modularity values are $Q = 0.331$ and 0.338 . Figures 3(b) and (c) show the modules of wTP53 and mTP53 with different colours. Each module of wTP53 or mTP53 has unique set of protein. However, there is a large overlap of secondary interactors (proteins which do not interact directly with TP53) in the wTP53 and mTP53 networks, which is distributed among different modules (in total 123). We observe that among 123 common proteins, 34 belong to module $W3 - M5$, whereas module pairs $M1 - W1$ (and $W4 - M2$) have 23 (and 10) common proteins. One can define a similarity measure σ_{mn} using Eq. (1) for every pair of wTP53-mTP53 modules. Taking the similarity indices σ_{wm} as weights (or thickness) of the link we have constructed a bipartite network which is shown in Fig. 4; the number of proteins is written beside each of the modules and the number of common proteins is specified along the links.

Enrichment of biological processes for the proteins present in every module of wTP53 and mTP53 PPIN using GeneCodis3 are presented in [Dataset S6](#) and [S7](#) respectively, where only the GO terms with $p < 0.0001$ are considered. The number of enriched GO

terms in modules of wTP53 PPIN are $W1(30)$, $W2(17)$, $W3(63)$, $W4(36)$ and those for mTP53 are $M1(52)$, $M2(71)$, $M4(1)$, $M5(67)$. Note that module $M3$ has no GO terms enriched with $p < 0.0001$.




Figure 4. Similarity between modules of wTP53 and mTP53 networks. The bipartite network constructed with the modules of wTP53 and mTP53; the common proteins present between a pair of wild and mutant module is written on respective link. The number of proteins that constitute the modules are written beside it.

Loss and Gain of biological processes in TP53 networks

Enrichment analysis of proteins in modules of wTP53 and mTP53 using GeneCodis3 reveals that respectively 127 and 172 GO terms (or biological processes) are enriched significantly ($p < 0.0001$). Among 127 GO terms of wTP53 57 GO terms do not appear in the mTP53 representing loss of the corresponding biological processes. Again the mTP53 network has 102 new GO terms (which were absent in wTP53). Besides, 70 enriched GO terms are found to be common in modules of mTP53 and wTP53 networks. We further associate each of the enriched GO terms with a relevant function. Loss and gain of these broadly classified functions are discussed below.

Gain of biological processes : The biological processes related to 102 new GO terms of mTP53 are gained due to mutation. The functions enriched in module $M1$ of mTP53 network are cell-cell communication (no of GO terms 5), signaling (13), protein complex/membrane assembly/stabilization (4), proteasomal degradation (2), cell cycle (3), DNA damage and repair (1) and others (2). GO terms related to DNA replication (11), DNA damage and repair (14), cell cycle (4), immunological functions (3), proteasomal degradation (3) and signaling (1) are enriched in $M2$. Similarly GO terms related to differentiation and development (11), signaling (7), transcription (6), cell proliferation (4), apoptosis (1), cell cycle (1) and DNA damage (2) and others (4) are enriched with proteins in module $M5$. The extensive list of the GOF is given in [Dataset S8 \(sheet 2\)](#) and in [Supporting Text \(Table S4\)](#). Thus new functions carried out by these biological processes are due to gain of interaction.

Loss of biological processes : On the other hand some of the enriched GO terms of wTP53 are absent in the mutant network. Corresponding biological processes are lost due to mutation in TP53. Altogether 57 unique GO terms are enriched with proteins in modules of wTP53 networks which are classified into broad class of functions (see [Dataset S8 \(sheet 1\)](#) and [Supporting Text \(Table S4\)](#)). The resulting loss of biological functions in various modules are, *W1* : signaling (8), proteasomal degradation (1), translation (1), cell migration and movement (2) and others (1); *W2* : signaling (4), apoptosis (2) and immunological (3); *W3* : cell cycle (3), signaling (1), transcription process and regulation (13), DNA replication (3), DNA damage and repair (1); *W4* : transcription process and regulation (2), proteasomal degradation (2), translation (5) and others (4); *W7* : transcription process and regulation (1).

Both loss and gain of biological processes : The 70 GO terms common between wTP53 and mTP53 networks are related to the functions, cell cycle (14 GO terms), transcription (15), DNA damage and repair (10), cell growth (2) and apoptosis (4), signaling (9), DNA replication (3), proteasomal degradation (3), immunological (2), development and differentiation (1), metabolism (2) and others (5). Thus these functions are possibly enriched due to both gain and loss of interactions (details are shown in [Dataset S8 \(sheet 3\)](#) and in [Supporting Text \(Table S4\)](#)).

Analysis of proteins in different modules using tool GeneDecks

Recently metastasis has been shown as the GOF as R273H cells attain metastatic property in cell model [57]. Since metastasis is not described as a “biological process” in Gene Ontology term, we have used another tool, GeneDecks [68], which provides a similarity metric by highlighting shared descriptors between genes, based on annotation within the GeneCards compendium of human genes (see [Text 4 in Supporting Text](#) for details). Taking the proteins of the modules of wTP53 and mTP53 separately as a query field, we look for “metastasis” in the attribute “disorder” among many other descriptors which are enriched for different types of cancers ([Dataset S9](#)). It is observed that the descriptor “metastasis” is enriched with the protein modules *W1*, *W2*, *W3* of wTP53 network and all the modules (*M1*, *M2*, *M3*, *M4*, *M5*) of mTP53 network. Thus, the loss of interactions of proteins in the modules *W1*, *W2*, *W3* of wTP53 due to mutation may result in the LOFs related to metastasis. Similarly, the gain of interactions of proteins in all the modules of mTP53 may result in the GOFs related to metastasis.

That LOF of wTP53 and GOF of mTP53 may contribute to invasion and metastasis, is reviewed recently [69]. TP53 mutations at the DNA binding domain are common and such mutations suppress expression of target genes. It is supported by several experiments [69] that suppression of transcriptional program for genes involved in epithelial-mesenchymal transition (EMT) may contribute to induction of EMT resulting in metastasis. Further, it is ascertained that loss of functions in wTP53 lead to increased cell motility in various cell types, and increased expression of fibronectin, collagens and extracellular matrix

(ECM) proteins. Enhanced expression of these proteins potentially increase the interaction between cells and ECM. LOF in wTP53 also activate Rho GTPases and modulates cell migration [69].

Role of mTP53 in metastasis has been established in many other studies. Mutant TP53 (R175H) is involved in TGF mediated invasion and metastasis in breast cancer cells through TP63 and SMAD3 [55]. Note that, in our analysis, SMAD is present in module *M5* of mTP53 network. It is known that mutant TP53 (R175H and R273H) increases endocytic recycling of adhesion molecule integrin and EGFR promoting and metastasis [56, 70]. Mutation in TP53 also activate EGFR/PI3K/AKT pathways and thereby increases invasion [71]. Various other mechanisms of increased metastasis by the mutant TP53 have also been studied [69]. Thus the gain of biological processes obtained from the analysis of mTP53 protein networks provides an explanation of GOFs observed in cancers.

Robustness analysis

In general, the modularization methods partition the network into communities of proteins which are densely connected. Thus in a large network it is quite expected that deletion of small fraction of links, whether selected methodically or randomly, does not alter the overall structure significantly. In fact, the degree distributions of all four networks studied here (namely PPIN of wHTT, mHTT, wTP53 and mTP53) are scale free (see Fig. S3 of Supporting Text), and it is known that such scale free networks are robust against random removal of nodes or links, but they could be fragile against targeted attack [72].

Table 1. Change in the total number of proteins and the interactions after excluding (a) genetic interactions and then (b) excluding interactions which are validated by only one Y2H experiment.

	Total no. of interactions	(a)Excluding genetic(%)	(b)Excluding genetic & Y2H(%)	Total no. of proteins	Excluding genetic & Y2H(%)
PPIN(human)	59027	58927(99.84%)	47244(80.04%)	12515	11630(9.29%)
wHTT	1380	1375(99.64%)	1231(89.20%)	306	292(9.54%)
mHTT	13105	13058(99.64%)	11478(87.58%)	1542	1486(9.64%)
wTP53	3718	3716(99.95%)	3205(86.20%)	619	590(9.53%)
mTP53	3521	3515(99.83%)	3136(89.07%)	578	551(9.53%)

Again, since several databases of protein interactions largely overlap [73] in their contents, it is natural to expect that the broadly classified biological functions [obtained here for HTT and TP53 networks would not differ substantially. In this study we used Biogrid [26] for creating the differential PPIN of the wild type and mutant HTT and TP53

Table 2. Comparison of number of proteins and GO terms in the modules of mHTT with respective of ‘most similar module’ of the network (a) after excluding genetic and Y2H experiments and (b) after deletion of 10% links.

	mHTT Module	(a) Excluding genetic & Y2H		(b) Random deletion of 10% links	
		MSM	Common (%)	MSM	Common (%)
M1	No. of Proteins : 643	656	542 (84.29%)	612	521 (81.03%)
	No. of GO terms : 161	161	147 (91.30%)	163	147 (91.30%)
M3	No. of Proteins : 377	287	211 (55.97%)	309	209 (55.44%)
	No. of GO terms : 78	95	64 (82.05%)	80	59 (75.64%)
M5	No. of Proteins : 485	442	423 (87.22%)	424	397 (81.86%)
	No. of GO terms : 198	186	173 (87.37%)	174	162 (81.82%)
M6	No. of Proteins : 7	5	5 (71.43%)	5	5 (71.43%)
	No. of GO terms : 12	16	11 (91.67%)	16	11 (91.67%)
M7	No. of Proteins : 22	7	7 (31.82%)	612	7 (31.82%)
	No. of GO terms : 5	8	5 (100.0%)	7	4 (80.00%)

proteins by connecting every pair of proteins which are listed in BioGrid as interacting partner of each other. This includes experimentally validated genetic and physical interactions. To check the robustness of our analysis, first let us remove all genetic interactions listed in BioGrid. This reduces the total number of protein interactions of BioGrid to 99.84%, whereas the interactions of wHTT, mHTT, wTP53 and mTP53 are reduced to 99.64%, 99.64%, 99.95% and 99.83% respectively (see Table 1). Among the other experiments considered in BioGrid, Yeast 2 Hybrid (Y2H) assay results in larger false positives [74]. Thus we further remove all the interactions which are identified only once by Y2H. This stringent criterion consequently reduces both the number of interactions and the number of proteins by $\sim 10\%$. The total number of interactions of BioGrid is, however, reduced by 20%. Since the wild type and mutant networks are altered only a little compared to the expected value 20%, one expects that deletion of a small fraction of interactions will not change the network properties significantly.

To demonstrate this explicitly, we reconstruct the PPIN of mHTT keeping only the reduced set of interactions and then identify the protein modules using Newman Girvan algorithm. The enriched GO terms ($p < 0.0001$) from GeneCodis3 shows that every module of mHTT (*M1*, *M3*, *M5*, *M6* and *M7*) has significant protein overlap with only ‘one distinct module’ of the reduced network, which is referred to as the ‘most similar module’ (MSM) henceforth. The number of overlapping proteins and GO terms between the modules of mHTT and their corresponding MSM in the reduced network are listed in Table 2. Evidently, in all cases, about 90% of the GO terms are retained. Thus, the loss, gain and loss/gain of biological processes obtained from BioGrid are quite robust.

For completeness, we also removed randomly 10% links of mHTT network and repeat the above analysis which is summarized in Table 2. Again, we find that about 90% of the GO terms enriched in this network are identical to those obtained for mHTT. Thus, in general, the enriched biological processes obtained through this analysis are quite robust.

Discussion and Conclusion

Mutation in protein may change its preference for binding with other proteins and alter the corresponding PPIN substantially. We use a graph theory based modularization approach to identify the modules of PPINs, and provide a comparative study of these differential networks using two examples; one for HD and another for cancers. The general philosophy of this analysis is depicted schematically in Fig. 5. In this figure, the wild type protein interacts with many other proteins forming a complex interaction network. Broadly, the schematic wild type network has three subgraphs or modules (A , B and C); proteins in each module are marked there with identical colours. The mutant protein loses some proteins as interacting partners (marked as pink) and gains some new ones (marked as orange, blue and violet). The network of the mutated protein has a revised modular structure A' , B' and D . Module A' and B' are re-structured and they have some proteins from other modules and some new proteins. Module D is gained by the mutation as most of proteins in this module were not present in the wild type network, and module C is lost. Correspondingly, the biological processes (GO terms) which are enriched in module D are gained and those enriched in module C are lost. We argue that this loss or gain of BPs lead to loss or gain of functions in the pathogenesis of the mutation induced disease.

In this article we explained the general idea of ‘obtaining the loss and gain of functions from the loss and gain of BPs enriched in protein modules’ using two examples; one for HD and another for cancers. Our analysis predict a set of broadly classified biological processes (from the the GO terms enriched in the modules of HTT and TP53 networks) which could be involved in the pathogenesis of HD and cancers respectively. In HD, the broadly classified BPs, like post transcriptional regulation of genes, apoptosis, synaptic transmission, JNK pathway, transcription deregulation, glucose transport, histone modifications etc are enriched with the proteins in modules of wHTT and mHTT networks. These BPs are already known to be altered in HD pathogenesis. Similarly, the gain and loss of BPs mTP53 results in the metastatic properties, which have been observed recently.

Although, we demonstrated the plausible loss and gain of biological processes in two examples where mutation alters protein interaction networks of wild type protein, the methodology discussed here can be adopted and applied to study differential PPIN in general. In particular, knowing the changes in the protein interaction network, either due to mutations that modify the structure of the protein at the binding surface or due to the change in interaction environments, one can predict what alteration might occur in the biological processes and functions. Such analysis may help understanding the loss or

gain of biological processes/functions in genetic diseases caused by mutations. This may in future lead to better design of disease intervention through targeting the biological processes/functions of specific modules.




Figure 5. Loss and gain of functions from differential network studies. The general philosophy of the current work is described here for a schematic protein interaction network, where the wild type and the mutant protein have 52 and 54 interactors respectively. There are three modules in wild type network (*A*, *B*, and *C*); all proteins in a given module are marked with same colour. After mutation the protein loses some interactors (marked as pink) and gain some new ones (marked as orange, blue and violet). The PPIN of mutant protein has three modules *A'*, *B'* and *D*. Module *A'*, which primarily contains proteins of module *A*, has some proteins from module *B* (green) and some new proteins (violet). Most of the proteins in module *D* are new interactors and thus this module is gained by the mutation. Similarly proteins of module *C* have lost their interactions. Correspondingly, the BPs which are enriched in module *D* are gained and those enriched in module *C* are lost. This loss or gain of BPs lead to loss or gain of functions in the pathogenesis of the mutation causing disease.

Supporting Information

Supporting Text : Text 1: Differential interaction due to mutation in HTT and TP53.

Text 2: Analysis of network structure. Text 3: Enriched biological processes in modules.

Text 4: Enrichment of metastasis from GeneDeck. (PDF)

Dataset S1 : Differential interaction of the wHTT and mHTT protein. (XLS)

Dataset S2 : The proteins belonging to different modules of wHTT network and their GO term enrichment analysis. (XLS)

Dataset S3 : The proteins belonging to different modules of mHTT network and their GO term enrichment analysis. (XLS)

Dataset S4 : The list of LOF,GOF and GOF/LOF for wHTT and mHTT networks. (XLS)

Dataset S5 : Differential interaction of the wTP53 and mTP53 protein. (XLS)

Dataset S6 : The proteins belonging to different modules of wTP53 network and their GO term enrichment analysis. (XLS)

Dataset S7 : The proteins belonging to different modules of mTP53 network and their GO term enrichment analysis. (XLS)

Dataset S8 : The list of LOF,GOF and GOF/LOF for wTP53 and mTP53 networks. (XLS)

Dataset S9 : The GeneDeck analysis of the proteins in the modules of wTP53 and mTP53 networks and enrichment of metastasis. (XLS)

Acknowledgments

The authors acknowledge Saikat Mukhopadhyay for his technical help and Urna Basu for careful reading of the manuscript.

References

1. Barabási A-L, Oltvai ZN (2004) Network biology: understanding the cell's functional organization. Nat Rev Genet 5:101113.
2. Schuster-Bckler B, Bateman A (2008) Protein interactions in human genetic diseases. Genome Biol 9:R9.

3. Wang X, Wei X, Thijssen B, Das J, Lipkin SM, et. al. (2012) Three-dimensional reconstruction of protein networks provides insight into human genetic disease. *Nat Biotechnol* 30:159-164.
4. The Huntington's Disease Collaborative Research Group (1993) A novel gene containing a trinucleotide repeat that is expanded and unstable on Huntington's disease chromosomes. *Cell* 72:971-983.
5. Imarisio S, J. Carmichael, V. Korolchuk, Chen CW, Saiki S, et. al. (2008) Huntington's disease: from pathology and genetics to potential therapies. *Biochem J* 412:191-209.
6. Ross CA, Tabrizi SJ (2011) Huntington's disease: from molecular pathogenesis to clinical treatment. *Lancet Neurol* 10:83-98.
7. Snell RG, MacMillan JC, Cheadle JP, Fenton I, Lazarou LP, et. al. (1993) Relationship between trinucleotide repeat expansion and phenotypic variation in Huntington's disease. *Nat Genet* 4:393-397.
8. Seong IS, Ivanova E, Lee JM, Choo YS, Fossale E, et. al. (2005) HD CAG repeat implicates a dominant property of huntingtin in mitochondrial energy metabolism. *Hum Mol Genet* 14:2871-2880.
9. Cowan CM (2006) Raymond Selective neuronal degeneration in Huntington's disease. *Curr Top Dev Biol* 75:25-71.
10. Jacobsen JC, Gregory GC, Woda JM, Thompson MN, Coser KR, et. al. (2011) HD CAG-correlated gene expression changes support a simple dominant gain of function. *Hum Mol Genet* 20:2846-2860.
11. Benn CL, Sun T, Sadri-Vakili G, McFarland KN, DiRocco DP, et. al. (2008) Huntingtin modulates transcription, occupies gene promoters in vivo, and binds directly to DNA in a polyglutamine-dependent manner. *J Neurosci* 28:10720-33.
12. Duyao MP, Auerbach AB, Ryan A, Persichetti F, Barnes GT, et. al. (1995) Inactivation of the mouse Huntingtons disease gene homolog Hdh. *Science* 269:407410.
13. Nasir J, Floresco SB, O'Kusky JR, Diewert VM, Richman JM, et. al. (1995) Targeted disruption of the Huntingtons disease gene results in embryonic lethality and behavioral and morphological changes in heterozygotes. *Cell* 81:811823.
14. S. Zeitlin, J. P. Liu, D. L. Chapman, V. E. Papaioannou, A. Efstratiadis (1995) Increased apoptosis and early embryonic lethality in mice nullizygous for the Huntington's disease gene homologue. *Nat Genet* 11:155-163.

15. Rigamonti D, Bauer JH, De-Fraja C, Conti L, Sipione S, et. al. (2000) Wild-type huntingtin protects from apoptosis upstream of caspase-3. *J. Neurosci.* 20:3705-3713; Rigamonti D, et. al. (2001) Huntingtin neuroprotective activity, occurs via inhibition of pro-caspase 9 processing. *J Biol Chem* 276:14545-14548.
16. Kegel KB, Meloni AR, Yi Y, Kim YJ, Doyle E, et. al. (2002) Huntingtin is present in the nucleus, interacts with the transcriptional corepressor C-terminal binding protein, and represses transcription. *J Biol Chem* 277:7466-7476.
17. Zhang H, Das S, Li QZ, Dragatsis I, Repa J, et. al. (2008) Elucidating a normal function of huntingtin by functional and microarray analysis of huntingtin-null mouse embryonic fibroblasts. *BMC Neurosci* 9:38.
18. Godin JD, Colombo K, Molina-Calavita M, Keryer G, Zala D, et. al. (2010) Huntingtin is required for mitotic spindle orientation and mammalian neurogenesis. *Neuron* 67:392-406.
19. Tong Y, Ha TJ, Liu L, Nishimoto A, Reiner A, et. al. (2011) Spatial and temporal requirements for huntingtin (Htt) in neuronal migration and survival during brain development. *J Neurosci* 31:14794-14799.
20. Van Raamsdonk JM, Gibson WT, Pearson J, Murphy Z, Lu G, et. al. (2006) Body weight is modulated by levels of full-length huntingtin. *Hum Mol Genet* 15:1513-1523.
21. Cattaneo E, Rigamonti D, Goffredo D, Zuccato C, Squitieri F, et. al. (2001) Loss of normal huntingtin function: new developments in Huntington's disease research. *Trends Neurosci* 24:182-188.
22. Ambrose CM, Duyao MP, Barnes G, Bates GP, Lin CS, et al. (1994) Structure and expression of the Huntington's disease gene: evidence against simple inactivation due to an expanded CAG repeat. *Somat. Cell Mol. Genet* 20:27-38.
23. Gottfried M, Lavine L, Roessmann U (1981) Neuropathological findings in Wolf-Hirschhorn (4p-) syndrome. *Acta Neuropathol* 55:163-165.
24. Wexler NS, Young AB, Tanzi RE, Travers H, Starosta-Rubinstein S, et al. (1987) Homozygotes for Huntington's disease. *Nature* 326:194-197.
25. Myers RH, Leavitt J, Farrer LA, Jagadeesh J, McFarlane H, et al. (1989) Homozygote for Huntington disease. *Am J Hum Genet* 45:615-618.
26. Stark C, Breitkreutz BJ, Chatr-Aryamontri A, Boucher L, Oughtred R, et al. The BioGRID Interaction Database: 2011 update, *Nucleic Acids Res* 39 (Database issue):D698-D704.

27. Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, et al. (2003) Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome Res* 13:2498-2504.
28. Fortunato S (2010) Community detection in graphs. *Phys Rep* 486:75-174.
29. Newman MEJ (2006) Modularity and community structure in networks. *Proc Natl Acad Sci USA* 103:8577-8582.
30. Newman MEJ, Girvan M (2004) Finding and evaluating community structure in networks. *Phys Rev E*. 69, 026113.
31. Sun S, Dong X, Fu Y, Tian W (2011) An iterative network partition algorithm for accurate identification of dense network modules. *Nucleic Acids Res.* 40(3):e18.
32. Spirin V, Leonid A, Mirny LA (2003) Protein complexes and functional modules in molecular networks. *Proc Natl Acad Sci USA* 100:12123-12128.
33. Tabas-Madrid D, Nogales-Cadenas R, Pascual-Montano A (2012) GeneCodis3: a non-redundant and modular enrichment analysis tool for functional genomics. *Nucleic Acids Res.* 40:W478-W483.
34. Huang DW, Sherman BT, Lempicki RA (2009) Bioinformatics enrichment tools: paths toward the comprehensive functional analysis of large gene lists. *Nucleic Acids Res.* 37:1-13.
35. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, et al. (2000) Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. *Nat Genet.* 25:25-29.
36. Bano D, Zanetti F, Mende Y, Nicotera P (2011) Neurodegenerative processes in Huntington's disease. *Cell Death Dis* 2, e228. doi: 10.1038/cddis.2011.112.
37. Pelegrí C, Duran-Vilaregut J, del Valle J, Crespo-Biel N, Ferrer I, et al. (2008) Cell cycle activation in striatal neurons from Huntington's disease patients and rats treated with 3-nitropropionic acid. *Int J Dev Neurosci.*, 26:665-671
38. Olch J, Junyent F, Verdaguer E, Auladell C, Pizarro JG, et al. (2012) Role of Cell Cycle Re-Entry in Neurons: A Common Apoptotic Mechanism of Neuronal Cell Death. *Neurotox Res* 22:195-207.
39. Jeppesen DK, Bohr VA, Stevnsner T (2011) DNA repair deficiency in neurodegeneration. *Prog Neurobiol* 94:166-200.
40. Enokido Y, Tamura T, Ito H, Arumughan A, Komuro A, et al. (2010) Mutant huntingtin impairs Ku70-mediated DNA repair, *J Cell Biol.* 189:425-443.

41. Humbert S (2010) Is Huntington disease a developmental disorder? *EMBO Rep.* 11(12):899.
42. Seredenina T, Luthi-Carter R (2012) What have we learned from gene expression profiles in Huntington's disease? *Neurobiol Dis* 45:83-98.
43. Borchert GM, Lanier W, Davidson BL (2006) RNA polymerase III transcribes human microRNA. *Nat Struct Mol Biol* 13:1097-1101.
44. Sinha M, Mukhopadhyay S, Bhattacharyya NP (2012) Mechanism(s) of Alteration of Micro RNA Expressions in Huntington's Disease and Their Possible Contributions to the Observed Cellular and Molecular Dysfunctions in the Disease. *Neuro-molecular Med* [Epub ahead of print] PMID: 22581158.
45. Milnerwood AJ, Raymond LA (2010) Early synaptic pathophysiology in neurodegeneration: insights from Huntington's disease. *Trends Neurosci* 33:513-523.
46. Perrin V, Dufour N, Raoul C, Hassig R, Brouillet E, et. al. (2009) Implication of the JNK pathway in a rat model of Huntington's disease. *Exp Neurol* 215:191-200.
47. Gamberino WC, Brennan WA Jr (1994) Glucose transporter isoform expression in Huntington's disease brain. *J Neurochem* 63:1392-1397.
48. Ciarmiello A, Giovacchini G, Orobello S, Bruselli L, Elifani F, et. al. (2012) 18F-FDG PET uptake in the pre-Huntington disease caudate affects the time-to-onset independently of CAG expansion size. *Eur J Nucl Med Mol Imaging*, 39, 1030-1036.
49. Bode FJ, Stephan M, Suhling H, Pabst R, Straub RH, et. al. (2008) Sex differences in a transgenic rat model of Huntington's disease: decreased 17beta-estradiol levels correlate with reduced numbers of DARPP32+ neurons in males. *Hum Mol Genet* 17:2595-2609.
50. Urdinguio RG, Sanchez-Mut JV, Esteller M (2009) Epigenetic mechanisms in neurological diseases: genes, syndromes, and therapies. *Lancet Neurol* 8:1056-1072.
51. Forbes SA, Bindal N, Bamford S, Cole C, Kok CY, et. al. (2011) COSMIC: mining complete cancer genomes in the Catalogue of Somatic Mutations in Cancer. *Nucleic Acids Res* 39(Database issue):D945-D950.
52. Freed-Pastor WA, Prives C (2012) Mutant p53: one name, many proteins. *Genes Dev* 26:1268-1286.
53. Oren M, Rotter V (2010) Mutant p53 Gain-of-Function in Cancer. *Cold Spring Harb Perspect Biol* 2(2):a001107.

54. Dittmer D, Pati S, Zambetti G, Chu S, Teresky AK, et. al.(1993) Gain of function mutations in p53. *Nat Genet* 4:4246.
55. Adorno M, Cordenonsi M, Montagner M, Dupont S, Wong C, et. al. (2009) A mutant-p53/Smad complex opposes p63 to empower TGFbeta-induced metastasis. *Cell* 137:8798.
56. Muller PA, Caswell PT, Doyle B, Iwanicki MP, Tan EH, et. al. (2009) Mutant p53 drives invasion by promoting integrin recycling. *Cell* 139:1327-1341.
57. Coffill CR, Muller PA, Oh HK, Neo SP, Hogue KA, et. al. (2012) Mutant p53 interactome identifies nardilysin as a p53R273H-specific binding partner that promotes invasion. *EMBO Rep.*13:638-644.
58. Shen Y, White E (2001) p53-dependent apoptosis pathways. *Adv Cancer Res* 82:55-84.
59. Sengupta S, Harris CC (2005) p53: traffic cop at the crossroads of DNA repair and recombination. *Nat Rev Mol Cell Biol* 6:44-55.
60. Laptenko O, Prives C (2006) Transcriptional regulation by p53: one protein, many possibilities. *Cell Death Differ* 13:951-961.
61. Levine AJ, Oren M (2009) The first 30 years of p53: growing ever more complex. *Nature Rev* 9:749-758.
62. Meek DW (2009) Tumour suppression by p53: a role for the DNA damage response? *Nat Rev Cancer* 9:714-723.
63. Riley T, Sontag E, Chen P, Levine A (2009) Transcriptional control of human p53-regulated genes. *Nat Rev Mol Cell Biol* 9:402-412.
64. Vousden KH, Ryan KM (2009) p53 and metabolism. *Nat Rev Cancer* 9:691-700.
65. Jain AK, Barton MC (2010) Making sense of ubiquitin ligases that regulate p53. *Cancer Biol Ther* 7:665-672.
66. Ozaki T, Nakagawara A (2011) Ap53: The Attractive Tumor Suppressor in the Cancer Research Field. *J Biomed Biotechnol Article ID* 603925.
67. Goh AM, Coffill CR, Lane DP (2011) The role of mutant p53 in human cancer. *J Pathol* 223:116-126.
68. Safran M, Dalah I, Alexander J, Rosen N, Iny Stein T, et. al. (2010) GeneCards Version 3: the human gene integrator, Database (Oxford). 2010 Aug 5;2010:baq020.

69. Muller PA, Vousden KH, Norman JC (2011) p53 and its mutants in tumor cell migration and invasion. *J Cell Biol* 192:209-218.
70. Selivanova G, Ivaska J (2009) Integrins and Mutant p53 on the Road to Metastasis. *Cell* 139:1220-1222.
71. Dong P, Xu Z, Jia N, Li D, Feng Y (2009) Elevated expression of p53 gain-of-function mutation R175H in endometrial cancer cells can increase the invasive phenotypes by activation of the EGFR/PI3K/AKT pathway. *Mol Cancer* 8, 103.
72. Albert R, Jeong H, Barabaši AL (2000) Error and attack tolerance of complex networks. *Nature* 406:378-382
73. Mathivanan S, Periaswamy B, Gandhi TKB, Kandasamy K, Suresh S, et. al. (2006) An evaluation of human protein-protein interaction data in the public domain. *BMC Bioinformatics* 7:S19.
74. Von Mering C, Krause R, Snel B, Cornell M, Oliver SG, et. al. (2002) Comparative assessment of large-scale data sets of protein-protein interactions. *Nature* 417:399-403.

Supporting Information

Mahashweta Basu¹, Nitai P. Bhattacharyya², P. K. Mohanty¹

**1 Theoretical Condensed Matter Physics Division,
Saha Institute of Nuclear Physics, 1/AF Bidhan Nagar, Kolkata 700064,
India**

**2 Crystallography and Molecular Biology Division,
Saha Institute of Nuclear Physics, 1/AF Bidhan Nagar, Kolkata 700064,
India**

Text 1 : Differential interaction due to mutation of HTT and TP53

Huntingtin protein

The mutated huntingtin protein mHTT preferentially interact with a set of different proteins compared to the wild type HTT (wHTT). The data for this differential interactions are curated from literature as follows. Primarily we have collected 328 HTT-interacting proteins identified by yeast 2 hybrid (Y2H) assays [1–3], affinity pull down followed by mass spectrometry assay [3] and proteins in the aggregates of mutant HTT [4]. Since Y2H assay results in false positives [5], we consider only 154 HTT interacting proteins that are validated by (i) second method like co-immunoprecipitation or (ii) the interacting proteins are tested for their implications in HD pathogenesis in cell or animal models of HD. Out of these sorted 154 HTT interacting proteins 17 proteins are seen to interact with wHTT, 12 proteins are identified in the aggregates of mutant HTT and 25 proteins interact preferentially with mutant HTT. Thus in totality, wHTT interact with 17 proteins whereas mHTT interacts with completely new 37 proteins. The list of these protein along with other subsidiary informations are presented in [Dataset S1 \(sheet 1\)](#).

The PPIN of wHTT is constructed by taking 306 proteins (wHTT, its 17 primary and 288 secondary interactors) and interactions among them, whereas the PPIN of mHTT has 1542 proteins ([see Dataset S1 \(sheet 2 and sheet 3\)](#)). Corresponding networks, drawn using Cytoscape [6], are shown in [Fig. S1](#).

Common proteins in wHTT and mHTT networks

Out of the 17 primary interactors of wHTT, 8 proteins namely (DLG4, CASP3, CTBP1, EIF2C2, REST, ZDHHC17, MEF2D, HIP1) change their direct interaction with wHTT and get involved in mHTT PPIN as secondary interactors. Whereas out of 288 secondary interactors in wHTT PPIN, 10 proteins now directly interact with mHTT. The list of these proteins are (PARK2, CASP10, TP53, CREBBP, SIN3A, CASP2, NFYC, CASP8, SP1, TBP). Also it is to be noted that, there are 107 proteins remain as secondary interactors




Figure S1. The interactome of wHHT and mHHT involves 306 and 1542 proteins respectively. If a pair of proteins present in a particular set has interaction between them, then a link is constructed between these two proteins(nodes). Exhausting every pair of proteins for both the sets we end up in constructing the interactome of the wHHT (left) and mHHT (right) using Cytoscape software [6].

in both the wHHT and mHHT PPIN (list given in [Table S1](#)).

TP53 protein

TP53 is a tumor suppressor gene. The mutation of this gene causes cancer. In a recent work Coffill *et. al.* [7] have studied the biological effects resulting from the missense mutations in TP53 (p53R273H). Their experimental results indicate that wTP53 and mTP53 differentially interact with 17 and 30 proteins; list of these proteins are given [Dataset S5 \(sheet 1\)](#).

The PPIN of wTP53 is constructed by taking 619 proteins (wTP53, its 17 primary and 601 secondary interactors) and interactions among them, whereas the PPIN of mTP53 has 578 proteins (see [Dataset S5 \(sheet 2 and sheet 3\)](#)). The corresponding networks, drawn using Cytoscape [6], are shown in [Fig. S2](#).




Figure S2. The PPIN of wTP53 and mTP53 interacting proteins are constructed using the Cytoscape software [6]. The network in the left is for the wild TP53 and in right is for mutant TP53.

Common proteins in wTP53 and mTP53 networks

Similar to the previous case the mutation of TP53 causes change in interacting partners of TP53, altogether resulting in different secondary interactors. Wild TP53 primarily interact with 17 proteins, among them 5 proteins (YWHAG, BAG6, TUBB3, RPL5, MDM2) lose their interaction with wTP53 and get involved in mTP53 PPIN as secondary interactor. Again these 7 proteins (PARD3, PLK1, KHDRBS1, CSNK2A1, PSMD2, CSNK2B, PSMC1) which were secondary interactors in wTP53 PPIN now interact with mTP53 directly. Apart from these there are 111 proteins common to both wild and mutant TP53 network, and are secondary interactors for both the networks (listed in [Table S1](#)).

Text 2 : Analysis of network structure

In recent years theory of complex networks has been used enormously in various fields [8]. The social systems (friendship, mobile or citation networks), information systems (World Wide Web and router networks), biological systems (neural, protein folding and gene regulatory networks) are a few to mention. Network theory provides a quantifiable description for comparison and characterization of complex networks. Here we shall focus on some robust measures of network topology : the degree distribution, the clustering coefficient, the average path length and the diameter.

Degree Distribution: The most elementary characteristic of a node (vertex) is its degree. The degree of a vertex k in a network is defined as the number of edges connected to that vertex. The degree distribution, $P(k)$, gives the probability that a selected node has exactly k links. $P(k)$ is obtained by counting the number of nodes with $k = 1, 2, \dots$ links and then normalizing by the number of nodes N . The degree distribution of many real world networks, including PPINs [9] are commonly scale free, *i.e.*

$$P(k) \sim k^{-\gamma}.$$

The degree distribution $P(k)$ for all four PPINs, *i.e.* wild and mutant networks of HTT and TP53, are found to be scale free (see [Fig. S3](#)), with the exponent $\gamma \simeq 2$. The average degree and our best estimate of the exponent γ are listed in [Table S2](#).

Clustering coefficient: In a highly dense network, the neighbours of a given node are very likely to be connected among themselves. The clustering coefficient

$$c_i = \frac{2E_i}{k_i(k_i - 1)}$$

gives a measure of how connected are the neighbours of node i . Here E_i the actual number of links which connects all the k_i neighbours of node i and $k_i(k_i - 1)/2$ is the maximum

number of possible links. Clearly the maximum value of c_i is unity. The average clustering coefficient of a network of N nodes is

$$C = \frac{1}{N} \sum_i^N c_i,$$

which gives a notion of how dense is the network, with maximum value 1 corresponding to a clique (*i.e.* all N nodes are connected to each other).




Figure S3. Degree distribution of wild and mutant HTT and TP53 network. All the four network follows scale free degree distribution. The corresponding power-law exponent is written in the graph itself.

Average path length : In a connected network, a given node i can be reached from any other node j by several alternative paths. The shortest path $l(i, j)$ is the path with the smallest number of links between nodes i and j . The average path length of a network of N nodes, thus $N(N - 1)/2$ vertex pairs, is

$$L = \frac{\sum_{i \neq j}^N 2l(i, j)}{N(N - 1)}.$$

The network diameter : The network diameter D is the largest of the shortest paths among all the vertex pairs. In other words, D is the length of the ‘longest’ shortest path in the network. The average path length and diameter for the networks of wHTT , mHTT, wTP53 and mTP53 are calculated and their values are listed in the [Table S2](#).

Modular structure of network : Other than these network parameters, detection of modules or community structures is widely used to reveal the organizational principles of nodes within the context of the entire system. Modules are specific groups of nodes which has larger density of links within the group than outside it. Each of such modules

can be regarded as a discrete entity whose function or properties are in some way separable from other modules. A large variety of community detection techniques [10] have been developed based on centrality measures, link density, percolation theory etc.

The most obvious way of finding groups in a network is to minimize the number of edges connecting the groups. Recently, Newman *et.al.* [11] proposed a method of finding community structure of a network by associating a score called modularity Q for each possible partition of a network; a good partition of a network can be obtained by maximizing Q . It is argued that simply rearranging the network, such that only few edges exist between the communities, is not enough. Rather one must rearrange it in a way that the communities are connected with fewer than expected edges. Based on this, Newman *et. al.* defined a score called modularity Q which is proportional to the number of edges present within the groups minus the expected number in an equivalent random network.

To be more precise, let us consider an undirected network of N nodes labeled by $i = 1, \dots, N$, having an adjacency matrix A with boolean elements,

$$A_{ij} = \begin{cases} 1, & \text{if nodes } i \text{ and } j \text{ are connected} \\ 0, & \text{otherwise.} \end{cases}$$

Thus, the degree of the node i is $k_i = \sum_j A_{ij}$, and the total number of links $m = \frac{1}{2} \sum_i k_i = \frac{1}{2} \sum_{ij} A_{i,j}$. If the network is to be partitioned into groups, the modularity is defined by

$$Q = \frac{1}{2m} \sum_{ij} \left(A_{ij} - \frac{k_i k_j}{2m} \right) \kappa_{i,j}, \quad (1)$$

where $\kappa_{i,j} = 1$ (or 0) if nodes i and j belong to same (different) partitions. Note that $k_i k_j / 2m$ is the expected number of links between nodes i and j , if edges were placed at random and it assures that Q is maximum when two groups are connected by smaller than expected number of links. A positive values of Q indicates the possible presence of community structure(s) and one need to look for a partition for which the modularity is preferably large and positive. In the following we apply this procedure to obtain modular structure of wHTT, mHTT, wTP53 and mTP53 networks.

Modularizing the wHTT and mHTT network we see that wHTT network contains 7 and mHTT network contains 8 modules. Also the wTP53 and mTP53 networks divides into 4 and 5 modules. The size of each modules *i.e.* number proteins present in each modules are given in the [Table S2](#). The names of the proteins that constitute the modules of wild and mutant HTT network are provided in the [Dataset S2 \(sheet 1\)](#) and [S3 \(sheet 1\)](#) respectively. Similarly for TP53, the name of the proteins present in the modules of the wild and mutant TP53 networks are given in the [Dataset S6 \(sheet 1\)](#) and [S7 \(sheet 1\)](#) respectively.

Text 3 : Enriched biological processes in modules

To find the possible function(s) of the modules, we utilized bioinformatics enrichment tools. There are many tools available for such enrichment analysis, we mainly used GeneCodis3 [12] for the convenience.

GeneCodis is a web-based tool for interpretation of large scale data integrating from different sources like Gene Ontology (GO, pathways in Kyoto Encyclopedia of Genes and Genomes (KEGG), PANTHER pathways, Online Mendelian inheritance in Man and others). Given a set of genes in the query field, GeneCodis associates GO terms for each gene and calculate whether the fraction of genes in a particular GO term among the input gene list is over represented compared to the background frequency (the total number of genes involved in that GO term over the total number gene in the organism). It further corrects the significance level for multiple testing. When the list of proteins present in a given module is feed to GeneCodis3 as the query set, it gives a list of associated biological processes (GO terms) and their respective significance values. In this study we considered only those GO terms which have $p < 0.0001$.

HTT PPIN : The set of proteins present in each of the 7 and 8 modules of wHTT and mHTT PPINs are analyzed with GeneCodis3. The GO terms associated with each of the modules are enlisted in [Dataset S2 and S3](#) according to high to low significance. The GO terms with corrected significance value greater than 0.0001 are discarded. Thus the number of GO terms belonging to each of the modules of wHTT PPIN are $W1(7)$, $W2(24)$, $W3(47)$, $W4(7)$, $W5(32)$, $W6(13)$, $W7(3)$ and for mHTT PPIN are $M1(161)$, $M3(78)$, $M5(198)$, $M6(12)$, $M7(5)$. For the modules $M2$, $M4$ and $M8$, all the GO terms that are obtained after GeneCodis analysis have p -value greater than 0.0001, so these GO terms are not considered for further analysis. Accumulating all the GO terms involved in the modules of wHTT and mHTT PPIN, one gets 129 and 389 GO terms respectively. (Note that merely adding the number of GO terms for each modules separately for wHTT and mHTT networks gives numbers greater than that mentioned; reason for this is that a single GO term can occur simultaneously in two or more modules of wHTT or mHTT network.) A detailed study reveals that both these set of GO terms obtained for wHTT and mHTT modules has 65 GO terms in common. Thus one can conclude that the unique GO terms that are involved only in wHTT and mHTT network are $64(= 129 - 65)$ and $324(= 389 - 65)$. The Biological processes relating to these 64 GO terms in wHTT network are lost due to mutation-LOF, whereas the BPs corresponding to 324 GO terms are GOF, and BPs for the common ones 65 are GOF/LOF both. We group the similar GO terms under a single broad BP. A clear data demonstrating these LOF, GOF and GOF/LOF (see [Table S3](#)) is given [Dataset S4](#).

TP53 PPIN : A similar study for the module of wild and mutant TP53 PPIN modules is being done to obtain the LOF, GOF and GOF/LOF separately. The total number of GO terms (considering GO terms with $p < 0.0001$ for all the modules of wTP53 and mTP53 separately) that are present for wild and mutant TP53 networks are 127 and 172

respectively. These two sets has 70 GO terms in common. Thus the unique GO terms present for the wild and mutant TP53 are 57 and 102; these are responsible for LOF and GOF function respectively. For the details of the loss and gain of functions refer to [Table S4](#) and [Dataset S8](#).

Text 4 : Enrichment of metastasis from GeneDeck

Metastasis, which is shown as GOF in mutation of TP53 [7], is not described for “biological process” in Gene Ontology term. We have used another tool, GeneDecks [13], which provides a similarity metric by highlighting shared descriptors between genes, based on annotation within the GeneCards compendium of human genes. Given a set of genes as query to the field Set Distiller in GeneDecks (Version 3), an online analysis tool that provides output of various descriptors like various cancers and diseases with the attribute Disorder taking data from different resources like Alma Knowledge Server, PharmGKB, The Breast Cancer Gene Database, Tumor Gene Database, GeneTests (formerly GeneClinics), OMIM, Swiss-Prot and Genatlas. The proteins present in the modules of wTP53 and mTP53 are analyzed using GeneDeck in order to get the information about the presence of metastasis in the corresponding modules. It is seen that the almost all the modules of wTP53 (except module *W4*) and mTP53 shows enriched metastasis (refer to [Dataset S9](#)).

References

1. Faber PW, Barnes GT, Srinidhi J, Chen J, Gusella JF, et. al. (1998) Huntingtin interacts with a family of WW domain proteins. *Hum Mol Genet* 7:1463-1474.
2. Goehler H, Lalowski M, Stelzl U, Waelter S, Stroedicke M, et. al. (2004) A protein inter- action network links GIT1, an enhancer of huntingtin aggregation, to Huntingtons disease. *Mol Cell* 15:853-65.
3. Kaltenbach LS, Romero E, Becklin RR, Chettier R, Bell R, et. al. (2007) Huntingtin interacting proteins are genetic modifiers of neurodegeneration. *PLoS Genet* 3, e82.
4. Mitsui K, Nakayama H, Akagi T, Nekooki M, Ohtawa K, et. al. (2002) Purification of polyglutamine aggregates and identification of elongation factor-1alpha and heat shock protein 84 as aggregate-interacting proteins. *J Neurosci* 22:9267-9277.
5. Von Mering C, Krause R, Snel B, Cornell M, Oliver SG, et. al. (2002) Comparative assessment of large-scale data sets of protein-protein interactions. *Nature* 417:399-403.

6. Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, et. al. (2003) Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome Res* 13:2498-2504.
7. Coffill CR, Muller PA, Oh HK, Neo SP, Hogue KA, et. al. (2012) Mutant p53 interactome identifies nardilysin as a p53R273H-specific binding partner that promotes invasion. *EMBO Rep* 13:638-644.
8. R. Albert, A. Barabási (2002) A Statistical mechanics of complex networks. *Rev of Mod Phys* 74:47-97.
9. R. Albert (2005) Scale-free networks in cell biology. *J Cell Sci.* 118:4947-4957.
10. Fortunato S (2010) Community detection in graphs. *Phys Rep* 486:75-174.
11. Newman MEJ, Girvan M (2004) Finding and evaluating community structure in networks. *Phys Rev E* 69, 026113.
12. Huang Da W, Sherman BT, Lempicki RA (2009) Bioinformatics enrichment tools: paths toward the comprehensive functional analysis of large gene lists. *Nucleic Acids Res.* 37:1-13.
13. Safran M, Dalah I, Alexander J, Rosen N, Iny Stein T, et. al. (2010) GeneCards Version 3: the human gene integrator, Database (Oxford). 2010 Aug 5;2010:baq020.

Table S1. List of the secondary interacting proteins that are common in to both wild and mutant PPIN of HTT and TP53 separately.

Common	Protein name
wHTT-mHTT(107)	YWHAQ, NFATC2, BCL3, HIST3H3, SMARCA4, TBK1, MAPK1, BTRC, SMARCC2, APC, PRKCA, CTTN, SMARCE1, UBC, MEN1, WAS, TCF7L2, KDM1A, NEUROD1, CASP6, RBBP5, GTF2E2, CBL, ZEB1, NUP98, AKT1, NUB1, IFT57, DHX58, BLM, TUBG1, EHMT1, COPS5, EHMT2, KDM5C, MAPK14, PSMC3, CFLAR, CARM1, NCOR1, CASK, MDM2, PRMT5, NCOA3, HSPB1, CDC42, BRCA1, HTR2C, HSPD1, FZD7, TBL1X, UBE2I, SENP3, HDAC1, HDAC2, HEY2, HDAC3, HDAC4, SUB1, SREBF2, HDAC5, TP73, THAP11, EP300, BID, HDAC9, TCF4, GTF2B, WIPF1, EGFR, KAT2A, SNCA, KAT2B, MLL, PFDN1, GRIK5, SERPINH1, RPS10, DNMI, SUMO1, SUMO2, USO1, KAT8, CHD3, RAC1, SIN3B, IKZF1, HIC1, DBNL, GRIN2B, IKZF4, SOX2, EIF6, YWHAE, DICER1, MAP3K14, BIRC2, MECOM, TGIF1, KHDRBS1, BIRC5, CABIN1, SATB1, HGS, ABL1, BCL2, XIAP
wTP53-mTP53(111)	UBC, TANK, RBBP4, RBBP7, PPP4C, MAGED1, SETDB1, SMAD1, UIMC1, CDH1, TERF1, NOLC1, PSMD10, CCDC85B, UBE2I, USP42, IKBKB, WRAP73, DCAF7, E2F1, MGMT, NCL, HSP90AA1, PRMT1, SRRM1, SRRM2, CHEK2, FBXW11, ITCH, PIK3R1, RB1, KIF5C, YWHAZ, STUB1, ARRB1, EP300, AR, SNAI1, RPS6KB1, MAP3K5, ESR1, UCHL1, UCHL5, STAU1, PCK1, MDM4, RPA1, RPA2, RPA1, BCL3, CDC5L, RASSF8, HSPA1A, PSMA3, NCOR2, PSMA7, TP53BP1, NDRG1, EXO1, CASP8, ARAF, H2AFX, PRPF40A, PELP1, EIF2C3, PIN1, NR3C1, CLSPN, RAD21, NEDD4, NEDD8, IMMT, CHAF1A, TSC1, MTA1, PUF60, CTBP1, MLLT4, MAPK6, TNFRSF1A, MEPCE, ARR3, SRGAP2, CREBBP, SRC, PRKCZ, SFN, CDK9, ATM, WNK1, RIPK1, SUMO1, SUMO2, HDGF, BTRC, LRIF1, CDKN1A, MDC1, RNPS1, PPP2R2D, HIST1H3A, HDAC1, HDAC3, DISC1, HDAC5, HDAC6, TP73, TAF1, PSMC4, PSMC5, C5ORF25

Table S2. Comparison of network properties between wild-type and mutant PPIN of HTT and TP53.

network	node	edge ¹	$\langle k \rangle$	γ	C	d	L	No. of Modules:	Module sizes	Q
wHTT	306	1380+17	9.13	1.95	0.436	4	2.418	7:	18, 66, 79, 18, 82, 8, 35	0.415
mHTT	1542	13105+37	17.05	1.99	0.361	4	2.349	8:	643, 3, 377, 2, 485, 7, 22, 3	0.302
wTP53	619	3718+17	12.07	2.04	0.452	4	2.232	4:	204, 151, 183, 81	0.331
mTP53	578	3521+30	12.29	1.89	0.406	4	2.282	5:	193, 127, 25, 111, 122	0.338

Table S3. LOF, GOF, LOF/GOF occurred due to mutation of HTT.

GO terms related to	Modules of mHTT network (unique GO terms)	Modules of wHTT network (unique GO terms)	Modules of wHTT-mHTT network (unique GO terms)
Cell cycle	M1(4),M3(12),M5(5)	W2(1),W6(3)	(W2,W5)-(M1,M3,M5)(4)
Signaling	M1(30),M5(31),M6(2)	W3(17),W4(4)	(W3,W4,W5,W7)-(M1,M5)(9)
Transcription processes and regulation	M1(5), M3(5), M5(31)	W7(1)	(W1,W5,W7)-(M3,M5,M6(17)
Apoptosis	M1(11)	W2(1)	(W2,W5)-(M1,M3,M5)(12)
DNA replication	M3(12)	-	-
DNA damage and repair	M1(6),M3(17),M5(4)	-	(W3,W5)-(M1,M3,M5)(4)
Immunological	M1(7)	-	-
Protein folding	M1(7)	-	-
Autophagy	M1(5)	-	-
Translation	M1(3)	-	-
Metabolism	M1(1)	-	-
Development and differentiation	M1(4),M5(57),M6(8)	-	(W3,W5)-M5(5)
Cell migration and shape	M1(4)	-	-
Proteasomal degradation	M1(14)	W2(1)	(W2,W5)-(M1,M3)(3)
Carbohydrate/Glucose transport/metabolism	M5(4)	W6(5)	-
Protein complex/membrane assembly/stabilization	M1(9)	-	-
Cell growth	M5(7)	-	-
Transcription from RNA polymerase III	M7(4)	-	-
Others	M1(4),M3(5),M5(6)	W3(4)	(W2,W3,W4,W5,W6)-(M1,M5)(6)
Gene silencing/micro RNA and RNA processing /translation	-	W1(5)	-
Synaptic transmission and neuronal activity	-	W3(12)	W3-M1(2)
Transport (ion/sugar)	-	W3(5)	-
Circadian rhythm	-	W2(1)	-
Protein /transmembrane transport	-	W6(4)	-
NFKB1 regulation	-	-	W2-M1(2)
Proliferation/growth	-	-	W5-(M1,M3,M5)(1)

Table S4. LOF, GOF, LOF/GOF occurred due to mutation of TP53.

GO terms related to	Modules of mTP53 network (unique GO terms)	Modules of wTP53 network (unique GO terms)	Modules of wTP53-mTP53 network (unique GO terms)
Cell cycle	M1(3),M2(4),M5(1)	W3(3)	(W1,W3,W4)-(M1,M2,M5) (14)
Signaling	M1(13), M5(7), M2(1)	W1(8), W3(1), W2(4)	(W1,W2,W3)-(M1,M4,M5) (9)
Transcription processes and regulation	M5(6)	W3(13), W4(2), W7(1)	(W1,W2,W3,W4)-(M1,M2,M5) (15)
Apoptosis	M5(1)	W2(2)	(W1,W2,W3,W4)-(M1,M2,M5) (4)
DNA replication	M2(11)	W3(3)	W3-(M2,M5) (3)
DNA damage and repair	M1(1), M2(14), M5(2)	W3(1)	(W1,W2,W3,W4)-(M1,M2,M5) (10)
Immunological	M2(3)	W2(3)	(W2,W4)-(M1.M2) (2)
Development and differentiation	M5(11)	-	W3-M5 (1)
Proteasomal degradation	M1(2),M2(3)	W1(1), W4(2)	(W4)-(M1.M2) (3)
Cell-cell communication	M1(5)	-	-
Cell proliferation/growth	M5(4)	-	W3-M5 (2)
Protein complex/membrane assembly/stabilization	M1(4)	-	-
Others	M1(2), M5(4)	W1(1), W4(4)	(W1,W2,W3,W4)-(M1,M2,M5) (5)
Translation	-	W1(1), W4(5)	-
Cell Migration and movement	-	W1(2)	-
Circadian rhythm	-	W4(1)	-
Metabolism	-	-	W4-M2 (2)