

# EVALUATIONS OF $q$ -WATSON TYPE, $q$ -DIXON TYPE AND $q$ -WHIPPLE TYPE SERIES

<sup>A,B</sup>CHUANAN WEI\*, <sup>C</sup>XIAOXIA WANG

<sup>A</sup>Department of Mathematics

Shanghai Normal University, Shanghai 200234, China

<sup>B</sup>Department of Information Technology

Hainan Medical College, Haikou 571199, China

<sup>C</sup>Department of Mathematics

Shanghai University, Shanghai 200444, China

**ABSTRACT.** According to the method of series rearrangement, we establish several extensions of  $q$ -Watson formulas with two extra integer parameters. Then they and Sear's transformation formula are used to derive some generalizations of  $q$ -Dixon formulas and  $q$ -Whipple formulas with two extra integer parameters. As the special cases of these results, many interesting evaluations of  $q$ -Watson type,  $q$ -Dixon type and  $q$ -Whipple type series are displayed.

## 1. INTRODUCTION

For two complex numbers  $x$  and  $q$ , define the  $q$ -shifted factorial by

$$(x; q)_0 = 1 \quad \text{and} \quad (x; q)_n = \prod_{k=0}^{n-1} (1 - xq^k) \quad \text{when } n \in \mathbb{N}.$$

The fraction form of it reads as

$$\left[ \begin{matrix} a, & b, & \cdots, & c \\ \alpha, & \beta, & \cdots, & \gamma \end{matrix} \middle| q \right]_n = \frac{(a; q)_n (b; q)_n \cdots (c; q)_n}{(\alpha; q)_n (\beta; q)_n \cdots (\gamma; q)_n}.$$

Following Gasper and Rahman [8], define the basic hypergeometric series by

$${}_1\phi_1 \left[ \begin{matrix} a_0, & a_1, & \cdots, & a_r \\ b_1, & b_2, & \cdots, & b_s \end{matrix} \middle| q; z \right] = \sum_{k=0}^{\infty} \left[ \begin{matrix} a_0, & a_1, & \cdots, & a_r \\ b_1, & b_2, & \cdots, & b_s \end{matrix} \middle| q \right]_k z^k$$

where  $\{a_i\}_{i \geq 0}$  and  $\{b_j\}_{j \geq 1}$  are complex parameters such that no zero factors appear in the denominators of the summand on the right hand side. Then Sear's transformation formula (cf. [8, Equation (2.10.4)]) can be expressed as

$${}_4\phi_3 \left[ \begin{matrix} q^{-n}, a, b, c \\ d, e, q^{1-n}abc/de \end{matrix} \middle| q; q \right] = \left[ \begin{matrix} d/a, de/bc \\ d, de/abc \end{matrix} \middle| q \right] {}_4\phi_3 \left[ \begin{matrix} q^{-n}, a, e/b, e/c \\ e, de/bc, q^{1-n}a/d \end{matrix} \middle| q; q \right]. \quad (1)$$

There are a lot of interesting summation formulas for basic hypergeometric series in the literature. The  $q$ -Watson formula due to Andrews [1] and the  $q$ -Watson formula due to Jain [9, Equation (3.17)] read, respectively, as

$${}_4\phi_3 \left[ \begin{matrix} q^{-n}, q^{1+n}a, \sqrt{c}, -\sqrt{c} \\ q\sqrt{a}, -q\sqrt{a}, c \end{matrix} \middle| q; q \right] = c^{n/2} \left[ \begin{matrix} q, q^2a/c \\ q^2a, qc \end{matrix} \middle| q^2 \right]_{\frac{n}{2}} \chi(n) \quad (2)$$

---

2010 *Mathematics Subject Classification*: Primary 05A19 and Secondary 33D15.

*Key words and phrases.*  $q$ -Watson formula;  $q$ -Dixon formula;  $q$ -Whipple formula.

\*Corresponding author. Email address: weichuanan78@163.com (C. Wei).

where  $\chi(n) = \{1 + (-1)^n\}/2$ ,

$${}_4\phi_3 \left[ \begin{matrix} a, c, q^{-n}, -q^{-n} \\ \sqrt{qac}, -\sqrt{qac}, q^{-2n} \end{matrix} \middle| q; q \right] = \left[ \begin{matrix} qa, qc \\ q, qac \end{matrix} \middle| q^2 \right]_n. \quad (3)$$

The  $q$ -Bailey-Dixon formula (cf. [8, p. 58]) can be stated as

$${}_4\phi_3 \left[ \begin{matrix} q^{-n}, a, c, -q^{1-n}/ac \\ q^{1-n}/a, q^{1-n}/c, -ac \end{matrix} \middle| q; q \right] = \left[ \begin{matrix} -a, c^2 \\ -ac, c \end{matrix} \middle| q \right]_n \left[ \begin{matrix} q, q^n a^2 c^2 \\ qc^2, q^n a^2 \end{matrix} \middle| q^2 \right]_n \chi(n). \quad (4)$$

By specifying the parameters in (1), we have the relation

$${}_4\phi_3 \left[ \begin{matrix} a, c, q^{-n}, -q^{1+n}a/c \\ qa/c, q^{1+n}a, -q^{-n}c \end{matrix} \middle| q; q \right] = \left[ \begin{matrix} q^{1+n}, -qa/c \\ q^{1+n}a, -q/c \end{matrix} \middle| q \right]_n {}_4\phi_3 \left[ \begin{matrix} a, qa/c^2, q^{-n}, -q^{-n} \\ qa/c, -qa/c, q^{-2n} \end{matrix} \middle| q; q \right].$$

The combination of the last formula and (3) creates another  $q$ -Dixon formula

$${}_4\phi_3 \left[ \begin{matrix} a, c, q^{-n}, -q^{1+n}a/c \\ qa/c, q^{1+n}a, -q^{-n}c \end{matrix} \middle| q; q \right] = \left[ \begin{matrix} q^{1+n}, -qa/c \\ q^{1+n}a, -q/c \end{matrix} \middle| q \right]_n \left[ \begin{matrix} qa, q^2 a/c^2 \\ q, q^2 a^2/c^2 \end{matrix} \middle| q^2 \right]_n. \quad (5)$$

The  $q$ -Whipple formula due to Andrews [1] and the  $q$ -Whipple formula due to Jain [9, Equation (3.19)] read, respectively, as

$${}_4\phi_3 \left[ \begin{matrix} q^{-n}, q^{1+n}, \sqrt{qac}, -\sqrt{qac} \\ -q, qa, qc \end{matrix} \middle| q; q \right] = \frac{(q^{1-n}a; q^2)_n (q^{1-n}c; q^2)_n}{(qa; q)_n (qc; q)_n} q^{\binom{n+1}{2}}, \quad (6)$$

$${}_4\phi_3 \left[ \begin{matrix} a, q/a, q^{-n}, -q^{-n} \\ c, q^{1-2n}/c, -q \end{matrix} \middle| q; q \right] = \frac{(ac; q^2)_n (qc/a; q^2)_n}{(c; q)_{2n}}. \quad (7)$$

Let  $u$  and  $v$  are both integers throughout the paper. Inspired by the work of [4], [6] and [7], we shall establish summation formulae for the following six series:

$$\begin{aligned} & {}_4\phi_3 \left[ \begin{matrix} q^{-n}, q^{1+u+v}a, \sqrt{c}, -q^v\sqrt{c} \\ q\sqrt{a}, -q^{1+u}\sqrt{a}, q^v c \end{matrix} \middle| q; q \right], & & {}_4\phi_3 \left[ \begin{matrix} q^u a, c, q^{-n}, -q^{v-n} \\ \sqrt{qac}, -q^u\sqrt{qac}, q^{v-2n} \end{matrix} \middle| q; q \right]; \\ & {}_4\phi_3 \left[ \begin{matrix} q^{-n}, a, c, -q^{1+u+v-n}/ac \\ q^{1+u-n}/a, q^{1+v-n}/c, -ac \end{matrix} \middle| q; q \right], & & {}_4\phi_3 \left[ \begin{matrix} a, c, q^{-n}, -q^{1+u+v+n}a/c \\ q^{1+u}a/c, q^{1+v+n}a, -q^{-n}c \end{matrix} \middle| q; q \right]; \\ & {}_4\phi_3 \left[ \begin{matrix} q^{-n}, q^{1+u+v} \\ -q, q^{1+u+v}a, qc \end{matrix} \middle| q; q \right], & & {}_4\phi_3 \left[ \begin{matrix} a, q^{1+u}/a, q^{-n}, -q^{v-n} \\ c, q^{1+u+v-2n}/c, -q \end{matrix} \middle| q; q \right]. \end{aligned}$$

## 2. EXTENSIONS OF $q$ -WATSON FORMULAS

### 2.1. Extensions of $q$ -Andrews-Watson formulas.

**Proposition 1.** *For two complex numbers  $\{a, c\}$  and a nonnegative integer  $m$ , there holds*

$$\begin{aligned} & {}_4\phi_3 \left[ \begin{matrix} q^{-n}, q^{1+n}a, \sqrt{c}, -q^m\sqrt{c} \\ q\sqrt{a}, -q\sqrt{a}, q^m c \end{matrix} \middle| q; q \right] \\ &= \sum_{j=0}^m q^{(m+n)j - \binom{j}{2}} c^{n/2} \left[ \begin{matrix} q^{-m}, q^{-n}, q^{1+n}a, \sqrt{c}, -\sqrt{c} \\ q, q\sqrt{a}, -q\sqrt{a}, q^m c, q^{j-1}c \end{matrix} \middle| q \right]_j \\ &\times \left[ \begin{matrix} q, q^2 a/c \\ q^{2+2j} a, q^{1+2j} c \end{matrix} \middle| q^2 \right]_{\frac{n-j}{2}} \chi(n-j). \end{aligned}$$

*Proof.* Letting  $a \rightarrow c/q$ ,  $b \rightarrow q^{-k}$ ,  $c \rightarrow -\sqrt{c}$  in the  ${}_6\phi_5$ -series identity (cf. [8, p. 42]):

$${}_6\phi_5 \left[ \begin{matrix} a, q\sqrt{a}, -q\sqrt{a}, b, c, q^{-m} \\ \sqrt{a}, -\sqrt{a}, qa/b, qa/c, q^{1+m}a \end{matrix} \middle| q; \frac{q^{1+m}a}{bc} \right] = \left[ \begin{matrix} qa, qa/bc \\ qa/b, qa/c \end{matrix} \middle| q \right]_m, \quad (8)$$

we get the equation

$$\begin{aligned} & \sum_{j=0}^m q^{mj + \binom{j}{2}} c^{j/2} \frac{1 - q^{2j-1}c}{1 - q^{j+m-1}c} \left[ \begin{matrix} q^{-m} \\ q \end{matrix} \middle| q \right]_j \left[ \begin{matrix} -\sqrt{c} \\ q^{j-1}c \end{matrix} \middle| q \right]_m \\ & \times \frac{(q^{k-j+1}; q)_j (q^{k+j}c; q)_{m-j}}{(-q^k\sqrt{c}; q)_m} = 1. \end{aligned}$$

Then there is the following relation

$$\begin{aligned} {}_4\phi_3 \left[ \begin{matrix} q^{-n}, q^{1+n}a, \sqrt{c}, -q^m\sqrt{c} \\ q\sqrt{a}, -q\sqrt{a}, q^m c \end{matrix} \middle| q; q \right] &= \sum_{k=0}^n \left[ \begin{matrix} q^{-n}, q^{1+n}a, \sqrt{c}, -q^m\sqrt{c} \\ q, q\sqrt{a}, -q\sqrt{a}, q^m c \end{matrix} \middle| q \right]_k q^k \\ &= \sum_{k=0}^n \left[ \begin{matrix} q^{-n}, q^{1+n}a, \sqrt{c}, -q^m\sqrt{c} \\ q, q\sqrt{a}, -q\sqrt{a}, q^m c \end{matrix} \middle| q \right]_k q^k \sum_{j=0}^m q^{mj+(j)} c^{j/2} \frac{1-q^{2j-1}c}{1-q^{j+m-1}c} \\ &\quad \times \left[ \begin{matrix} q^{-m} \\ q \end{matrix} \middle| q \right]_j \left[ \begin{matrix} -\sqrt{c} \\ q^{j-1}c \end{matrix} \middle| q \right]_m \frac{(q^{k-j+1}; q)_j (q^{k+j}c; q)_{m-j}}{(-q^k\sqrt{c}; q)_m}. \end{aligned}$$

Interchange the summation order for the last double sum to achieve

$$\begin{aligned} {}_4\phi_3 \left[ \begin{matrix} q^{-n}, q^{1+n}a, \sqrt{c}, -q^m\sqrt{c} \\ q\sqrt{a}, -q\sqrt{a}, q^m c \end{matrix} \middle| q; q \right] &= \sum_{j=0}^m q^{mj+(j)} c^{j/2} \frac{1-q^{2j-1}c}{1-q^{j+m-1}c} \\ &\quad \times \left[ \begin{matrix} q^{-m} \\ q \end{matrix} \middle| q \right]_j \left[ \begin{matrix} -\sqrt{c} \\ q^{j-1}c \end{matrix} \middle| q \right]_m \\ &\quad \times \sum_{k=j}^n \left[ \begin{matrix} q^{-n}, q^{1+n}a, \sqrt{c}, -q^m\sqrt{c} \\ q, q\sqrt{a}, -q\sqrt{a}, q^m c \end{matrix} \middle| q \right]_k q^k \frac{(q^{k-j+1}; q)_j (q^{k+j}c; q)_{m-j}}{(-q^k\sqrt{c}; q)_m}. \end{aligned}$$

Shifting the index  $k \rightarrow i+j$  for the sum on the last line, the result reads as

$$\begin{aligned} {}_4\phi_3 \left[ \begin{matrix} q^{-n}, q^{1+n}a, \sqrt{c}, -q^m\sqrt{c} \\ q\sqrt{a}, -q\sqrt{a}, q^m c \end{matrix} \middle| q; q \right] &= \sum_{j=0}^m q^{mj+(j+1)} c^{j/2} \left[ \begin{matrix} q^{-m}, q^{-n}, q^{1+n}a, \sqrt{c}, -\sqrt{c} \\ q, q\sqrt{a}, -q\sqrt{a}, q^m c, q^{j-1}c \end{matrix} \middle| q \right]_j \\ &\quad \times {}_4\phi_3 \left[ \begin{matrix} q^{j-n}, q^{1+n+j}a, q^j\sqrt{c}, -q^j\sqrt{c} \\ q^{1+j}\sqrt{a}, -q^{1+j}\sqrt{a}, q^{2j}c \end{matrix} \middle| q; q \right]. \end{aligned} \tag{9}$$

Calculating the  ${}_4\phi_3$ -series on the last line by (2), we attain Proposition 1 to complete the proof.  $\square$

**Example 1** ( $m = 1$  in Proposition 1).

$$\begin{aligned} {}_4\phi_3 \left[ \begin{matrix} q^{-n}, q^{1+n}a, \sqrt{c}, -q\sqrt{c} \\ q\sqrt{a}, -q\sqrt{a}, qc \end{matrix} \middle| q; q \right] &= \begin{cases} c^s \left[ \begin{matrix} q, q^2a/c \\ q^2a, qc \end{matrix} \middle| q^2 \right]_s, & n = 2s; \\ c^{\frac{1}{2}+s} \left[ \begin{matrix} q \\ qc \end{matrix} \middle| q^2 \right]_{1+s} \left[ \begin{matrix} q^2a/c \\ q^2a \end{matrix} \middle| q^2 \right]_s, & n = 1 + 2s. \end{cases} \end{aligned}$$

**Theorem 2.** For two complex numbers  $\{a, c\}$  and two nonnegative integers  $\{\ell, m\}$ , there holds

$$\begin{aligned} {}_4\phi_3 \left[ \begin{matrix} q^{-n}, q^{1+\ell+n}a, \sqrt{c}, -q^m\sqrt{c} \\ q\sqrt{a}, -q^{1+\ell}\sqrt{a}, q^m c \end{matrix} \middle| q; q \right] &= \left[ \begin{matrix} q^{1+\ell+n}a, q^{1+n}\sqrt{a} \\ q^{1+\ell+2n}a, q\sqrt{a} \end{matrix} \middle| q \right]_\ell \\ &= \sum_{i=0}^{\ell} \sum_{j=0}^m q^{2i+(m+n-i)j-(j)} \frac{c^{(n-i)/2}}{a^{i/2}} \frac{1-q^{1+2\ell+2n-2i}a}{1-q^{1+2\ell+2n}a} \left[ \begin{matrix} q^{-\ell}, q^{-n}, q^{-2\ell-2n-1}/a \\ q, q^{-\ell-2n}/a, q^{-2\ell-n}/a \end{matrix} \middle| q \right]_i \\ &\quad \times \left[ \begin{matrix} q^{-m}, q^{i-n}, q^{1+2\ell+n-i}a, \sqrt{c}, -\sqrt{c} \\ q, q^{1+\ell}\sqrt{a}, -q^{1+\ell}\sqrt{a}, q^m c, q^{j-1}c \end{matrix} \middle| q \right]_j \left[ \begin{matrix} q, q^{2+2\ell}a/c \\ q^{2+2\ell+2j}a, q^{1+2j}c \end{matrix} \middle| q^2 \right]_{\frac{n-i-j}{2}} \chi(n-i-j). \end{aligned}$$

*Proof.* Performing the replacements  $m \rightarrow \ell$ ,  $a \rightarrow q^{-2\ell-2n-1}/a$ ,  $b \rightarrow q^{k-n}$ ,  $c \rightarrow q^{-\ell-n}/\sqrt{a}$  in (8), we obtain the equation

$$\begin{aligned} &\left[ \begin{matrix} q^{1+\ell+n}a, q^{1+n}\sqrt{a} \\ q^{1+\ell+2n}a, q^{1+k}\sqrt{a} \end{matrix} \middle| q \right]_\ell \sum_{i=0}^{\ell} \frac{q^{2i}}{a^{i/2}} \frac{1-q^{1+2\ell+2n-2i}a}{1-q^{1+2\ell+2n}a} \left[ \begin{matrix} q^{-\ell}, q^{k-n}, q^{-2\ell-2n-1}/a \\ q, q^{-\ell-2n}/a, q^{-2\ell-n}/a \end{matrix} \middle| q \right]_i \\ &\quad \times \left[ \begin{matrix} q^{\ell+n+k+1}a \\ q^{\ell+n+1}a \end{matrix} \middle| q \right]_{\ell-i} = 1. \end{aligned}$$

Then there exists the following relation

$$\begin{aligned}
{}_4\phi_3 \left[ \begin{matrix} q^{-n}, q^{1+\ell+n}a, \sqrt{c}, -q^v\sqrt{c} \\ q\sqrt{a}, -q^{1+\ell}\sqrt{a}, q^v c \end{matrix} \middle| q; q \right] &= \sum_{k=0}^n \left[ \begin{matrix} q^{-n}, q^{1+\ell+n}a, \sqrt{c}, -q^v\sqrt{c} \\ q, q\sqrt{a}, -q^{1+\ell}\sqrt{a}, q^v c \end{matrix} \middle| q \right]_k q^k \\
&= \sum_{k=0}^n \left[ \begin{matrix} q^{-n}, q^{1+\ell+n}a, \sqrt{c}, -q^v\sqrt{c} \\ q, q\sqrt{a}, -q^{1+\ell}\sqrt{a}, q^v c \end{matrix} \middle| q \right]_k q^k \left[ \begin{matrix} q^{1+\ell+n}a, q^{1+n}\sqrt{a} \\ q^{1+\ell+2n}a, q^{1+k}\sqrt{a} \end{matrix} \middle| q \right]_\ell \\
&\times \sum_{i=0}^{\ell} \frac{q^{2i}}{a^{i/2}} \frac{1-q^{1+2\ell+2n-2i}a}{1-q^{1+2\ell+2n}a} \left[ \begin{matrix} q^{-\ell}, q^{k-n}, q^{-2\ell-2n-1}/a \\ q, q^{-\ell-2n}/a, q^{-2\ell-n}/a \end{matrix} \middle| q \right]_i \left[ \begin{matrix} q^{\ell+n+k+1}a \\ q^{\ell+n+1}a \end{matrix} \middle| q \right]_{\ell-i} \\
&= \left[ \begin{matrix} q^{1+\ell+n}a, q^{1+n}\sqrt{a} \\ q^{1+\ell+2n}a, q\sqrt{a} \end{matrix} \middle| q \right]_\ell \sum_{i=0}^{\ell} \frac{q^{2i}}{a^{i/2}} \frac{1-q^{1+2\ell+2n-2i}a}{1-q^{1+2\ell+2n}a} \left[ \begin{matrix} q^{-\ell}, q^{-2\ell-2n-1}/a \\ q, q^{-\ell-2n}/a, q^{-2\ell-n}/a \end{matrix} \middle| q \right]_i \\
&\times \sum_{k=0}^n \left[ \begin{matrix} q^{-n}, q^{1+\ell+n}a, \sqrt{c}, -q^v\sqrt{c} \\ q, q\sqrt{a}, -q^{1+\ell}\sqrt{a}, q^v c \end{matrix} \middle| q \right]_k q^k \frac{(q\sqrt{a}; q)_\ell (q^{k-n}; q)_i}{(q^{1+k}\sqrt{a}; q)_\ell} \left[ \begin{matrix} q^{\ell+n+k+1}a \\ q^{\ell+n+1}a \end{matrix} \middle| q \right]_{\ell-i}.
\end{aligned}$$

After some routine simplification, the result reads as

$$\begin{aligned}
{}_4\phi_3 \left[ \begin{matrix} q^{-n}, q^{1+\ell+n}a, \sqrt{c}, -q^v\sqrt{c} \\ q\sqrt{a}, -q^{1+\ell}\sqrt{a}, q^v c \end{matrix} \middle| q; q \right] &= \left[ \begin{matrix} q^{1+\ell+n}a, q^{1+n}\sqrt{a} \\ q^{1+\ell+2n}a, q\sqrt{a} \end{matrix} \middle| q \right]_\ell \\
&\times \sum_{i=0}^{\ell} \frac{q^{2i}}{a^{i/2}} \frac{1-q^{1+2\ell+2n-2i}a}{1-q^{1+2\ell+2n}a} \left[ \begin{matrix} q^{-\ell}, q^{-n}, q^{-2\ell-2n-1}/a \\ q, q^{-\ell-2n}/a, q^{-2\ell-n}/a \end{matrix} \middle| q \right]_i \\
&\times {}_4\phi_3 \left[ \begin{matrix} q^{i-n}, q^{1+2\ell+n-i}a, \sqrt{c}, -q^v\sqrt{c} \\ q^{1+\ell}\sqrt{a}, -q^{1+\ell}\sqrt{a}, q^v c \end{matrix} \middle| q; q \right]. \tag{10}
\end{aligned}$$

Setting  $v = m$  in (10) and evaluating the  ${}_4\phi_3$ -series on the right hand side by Proposition 1, we gain Theorem 2 to finish the proof.  $\square$

**Example 2** ( $\ell = 1, m = 0$  in Theorem 2).

$$\begin{aligned}
&{}_4\phi_3 \left[ \begin{matrix} q^{-n}, q^{2+n}a, \sqrt{c}, -\sqrt{c} \\ q\sqrt{a}, -q^2\sqrt{a}, c \end{matrix} \middle| q; q \right] \\
&= \begin{cases} \frac{(1+q\sqrt{a})c^s}{1+q^{1+2s}\sqrt{a}} \left[ \begin{matrix} q, q^4a/c \\ q^2a, qc \end{matrix} \middle| q^2 \right]_s, & n = 2s; \\ \frac{(q^2-q)\sqrt{a}c^s}{(1-q\sqrt{a})(1+q^{2+2s}\sqrt{a})} \left[ \begin{matrix} q^3, q^4a/c \\ q^4a, qc \end{matrix} \middle| q^2 \right]_s, & n = 1 + 2s. \end{cases}
\end{aligned}$$

**Example 3** ( $\ell = 1, m = 1$  in Theorem 2).

$$\begin{aligned}
&{}_4\phi_3 \left[ \begin{matrix} q^{-n}, q^{2+n}a, \sqrt{c}, -q\sqrt{c} \\ q\sqrt{a}, -q^2\sqrt{a}, qc \end{matrix} \middle| q; q \right] \\
&= \begin{cases} \frac{(1+q\sqrt{a})(\sqrt{c}+q^{1+2s}\sqrt{a})c^s}{(\sqrt{c}+q\sqrt{a})(1+q^{1+2s}\sqrt{a})} \left[ \begin{matrix} q, q^2a/c \\ q^2a, qc \end{matrix} \middle| q^2 \right]_s, & n = 2s; \\ \frac{(1-q)(\sqrt{c}-q\sqrt{a})(1+q^{2+2s}\sqrt{a}c)c^s}{(1-q\sqrt{a})(1-qc)(1+q^{2+2s}\sqrt{a})} \left[ \begin{matrix} q^3, q^4a/c \\ q^4a, q^3c \end{matrix} \middle| q^2 \right]_s, & n = 1 + 2s. \end{cases}
\end{aligned}$$

Replacing  $\sqrt{a}$  by  $-q^{-\ell}\sqrt{a}$  in Theorem 2, we get the equation

$$\begin{aligned}
&{}_4\phi_3 \left[ \begin{matrix} q^{-n}, q^{1-\ell+n}a, \sqrt{c}, -q^m\sqrt{c} \\ q\sqrt{a}, -q^{1-\ell}\sqrt{a}, q^m c \end{matrix} \middle| q; q \right] = \left[ \begin{matrix} q^{-n}/a, -q^{-n}/\sqrt{a} \\ q^{-2n}/a, -1/\sqrt{a} \end{matrix} \middle| q \right]_\ell \\
&= \sum_{i=0}^{\ell} \sum_{j=0}^m (-1)^i q^{(\ell+2)i+(m+n-i)j-\binom{j}{2}} \frac{c^{(n-i)/2}}{a^{i/2}} \frac{1-q^{1+2n-2i}a}{1-q^{1+2n}a} \left[ \begin{matrix} q^{-\ell}, q^{-n}, q^{-2n-1}/a \\ q, q^{-n}/a, q^{\ell-2n}/a \end{matrix} \middle| q \right]_i \\
&\times \left[ \begin{matrix} q^{-m}, q^{i-n}, q^{1+n-i}a, \sqrt{c}, -\sqrt{c} \\ q, q\sqrt{a}, -q\sqrt{a}, q^m c, q^{j-1}c \end{matrix} \middle| q \right]_j \left[ \begin{matrix} q, q^2a/c \\ q^{2+2j}a, q^{1+2j}c \end{matrix} \middle| q^2 \right]_{\frac{n-i-j}{2}} \chi(n-i-j). \tag{11}
\end{aligned}$$

Employing the substitution  $\sqrt{c} \rightarrow -q^{-m}\sqrt{c}$  in Theorem 2, we achieve the formula

$$\begin{aligned} {}_4\phi_3 & \left[ \begin{matrix} q^{-n}, q^{1+\ell+n}a, \sqrt{c}, -q^{-m}\sqrt{c} \\ q\sqrt{a}, -q^{1+\ell}\sqrt{a}, q^{-m}c \end{matrix} \middle| q; q \right] = \left[ \begin{matrix} q^{1+\ell+n}a, q^{1+n}\sqrt{a} \\ q^{1+\ell+2n}a, q\sqrt{a} \end{matrix} \middle| q \right]_{\ell} \\ &= \sum_{i=0}^{\ell} \sum_{j=0}^m (-1)^j q^{2i+(n-i-j)(j-m)+\binom{j+1}{2}} \frac{c^{(n-i)/2}}{a^{i/2}} \frac{1-q^{1+2\ell+2n-2i}a}{1-q^{1+2\ell+2n}a} \\ &\quad \times \left[ \begin{matrix} q^{-\ell}, q^{-n}, q^{-2\ell-2n-1}/a \\ q, q^{-\ell-2n}/a, q^{-2\ell-n}/a \end{matrix} \middle| q \right]_i \left[ \begin{matrix} q^{-m}, q^{i-n}, q^{1+2\ell+n-i}a, q^{-m}\sqrt{c}, -q^{-m}\sqrt{c} \\ q, q^{1+\ell}\sqrt{a}, -q^{1+\ell}\sqrt{a}, q^{-m}c, q^{j-2m-1}c \end{matrix} \middle| q \right]_j \\ &\quad \times \left[ \begin{matrix} q, q^{2+2\ell+2m}a/c \\ q^{2+2\ell+2j}a, q^{1-2m+2j}c \end{matrix} \middle| q^2 \right]_{\frac{n-i-j}{2}} \chi(n-i-j). \end{aligned} \quad (12)$$

Letting  $\sqrt{a} \rightarrow -q^{-\ell}\sqrt{a}$ ,  $\sqrt{c} \rightarrow -q^{-m}\sqrt{c}$  in Theorem 2, we attain the result

$$\begin{aligned} {}_4\phi_3 & \left[ \begin{matrix} q^{-n}, q^{1-\ell+n}a, \sqrt{c}, -q^{-m}\sqrt{c} \\ q\sqrt{a}, -q^{1-\ell}\sqrt{a}, q^{-m}c \end{matrix} \middle| q; q \right] = \left[ \begin{matrix} q^{-n}/a, -q^{-n}/\sqrt{a} \\ q^{-2n}/a, -1/\sqrt{a} \end{matrix} \middle| q \right]_{\ell} \\ &= \sum_{i=0}^{\ell} \sum_{j=0}^m (-1)^{i+j} q^{(\ell+2)i+(n-i-j)(j-m)+\binom{j+1}{2}} \frac{c^{(n-i)/2}}{a^{i/2}} \frac{1-q^{1+2n-2i}a}{1-q^{1+2n}a} \\ &\quad \times \left[ \begin{matrix} q^{-\ell}, q^{-n}, q^{-2n-1}/a \\ q, q^{-n}/a, q^{\ell-2n}/a \end{matrix} \middle| q \right]_i \left[ \begin{matrix} q^{-m}, q^{i-n}, q^{1+n-i}a, q^{-m}\sqrt{c}, -q^{-m}\sqrt{c} \\ q, q\sqrt{a}, -q\sqrt{a}, q^{-m}c, q^{j-2m-1}c \end{matrix} \middle| q \right]_j \\ &\quad \times \left[ \begin{matrix} q, q^{2+2m}a/c \\ q^{2+2j}a, q^{1-2m+2j}c \end{matrix} \middle| q^2 \right]_{\frac{n-i-j}{2}} \chi(n-i-j). \end{aligned} \quad (13)$$

**Remark:** Theorem 2, (11), (12) and (13) are equivalent with each other, but the corresponding hypergeometric series identities of them are essentially different.

## 2.2. Extensions of $q$ -Jain-Watson formula.

**Proposition 3.** For two complex numbers  $\{a, c\}$  and a nonnegative integer  $m$  with  $m \leq n$ , there holds

$$\begin{aligned} {}_4\phi_3 & \left[ \begin{matrix} a, c, q^{-n}, -q^{m-n} \\ \sqrt{qac}, -\sqrt{qac}, q^{m-2n} \end{matrix} \middle| q; q \right] \\ &= \sum_{j=0}^m q^{(m-n)j+\binom{j+1}{2}} \left[ \begin{matrix} q^{-m}, q^{-n}, -q^{-n}, a, c \\ q, \sqrt{qac}, -\sqrt{qac}, q^{m-2n}, q^{j-2n-1} \end{matrix} \middle| q \right]_j \\ &\quad \times \left[ \begin{matrix} q^{1+j}a, q^{1+j}c \\ q, q^{1+2j}ac \end{matrix} \middle| q^2 \right]_{n-j}. \end{aligned}$$

*Proof.* Performing the replacement  $a \rightarrow q^{-1-n}a$  in (9), we obtain

$$\begin{aligned} {}_4\phi_3 & \left[ \begin{matrix} q^{-n}, a, \sqrt{c}, -q^{m}\sqrt{c} \\ \sqrt{q^{1-n}a}, -\sqrt{q^{1-n}a}, q^m c \end{matrix} \middle| q; q \right] \\ &= \sum_{j=0}^m q^{mj+\binom{j+1}{2}} c^{j/2} \left[ \begin{matrix} q^{-m}, q^{-n}, a, \sqrt{c}, -\sqrt{c} \\ q, \sqrt{q^{1-n}a}, -\sqrt{q^{1-n}a}, q^m c, q^{j-1}c \end{matrix} \middle| q \right]_j \\ &\quad \times {}_4\phi_3 \left[ \begin{matrix} q^{j-n}, q^j a, q^j \sqrt{c}, -q^j \sqrt{c} \\ q^j \sqrt{q^{1-n}a}, -q^j \sqrt{q^{1-n}a}, q^{2j} c \end{matrix} \middle| q; q \right]. \end{aligned} \quad (14)$$

Replacing  $c$  and  $q^{-n}$  by  $q^{-2n}$  and  $c$ , respectively, in (14), the result reads as

$$\begin{aligned} {}_4\phi_3 & \left[ \begin{matrix} a, c, q^{-n}, -q^{m-n} \\ \sqrt{qac}, -\sqrt{qac}, q^{m-2n} \end{matrix} \middle| q; q \right] \\ &= \sum_{j=0}^m q^{(m-n)j+\binom{j+1}{2}} \left[ \begin{matrix} q^{-m}, q^{-n}, -q^{-n}, a, c \\ q, \sqrt{qac}, -\sqrt{qac}, q^{m-2n}, q^{j-2n-1} \end{matrix} \middle| q \right]_j \\ &\quad \times {}_4\phi_3 \left[ \begin{matrix} q^j a, q^j c, q^{j-n}, -q^{j-n} \\ q^j \sqrt{qac}, -q^j \sqrt{qac}, q^{2j-2n} \end{matrix} \middle| q; q \right]. \end{aligned}$$

Calculating the  ${}_4\phi_3$ -series on the last line by (3), we gain Proposition 3 to complete the proof.  $\square$

**Example 4** ( $m = 1$  in Proposition 3:  $n \geq 1$ ).

$${}_4\phi_3 \left[ \begin{matrix} a, c, q^{-n}, -q^{1-n} \\ \sqrt{qac}, -\sqrt{qac}, q^{1-2n} \end{matrix} \middle| q; q \right] = \left[ \begin{matrix} qa, qc \\ q, qac \end{matrix} \middle| q^2 \right]_n + q^n \left[ \begin{matrix} a, c \\ q, qac \end{matrix} \middle| q^2 \right]_n.$$

**Proposition 4.** For two complex numbers  $\{a, c\}$  and a nonnegative integer  $m$ , there holds

$$\begin{aligned} & {}_4\phi_3 \left[ \begin{matrix} a, c, q^{-n}, -q^{-m-n} \\ \sqrt{qac}, -\sqrt{qac}, q^{-m-2n} \end{matrix} \middle| q; q \right] \\ &= \sum_{j=0}^m (-1)^j q^{\binom{j+1}{2}-nj} \left[ \begin{matrix} q^{-m}, q^{-m-n}, -q^{-m-n}, a, c \\ q, \sqrt{qac}, -\sqrt{qac}, q^{-m-2n}, q^{j-2m-2n-1} \end{matrix} \middle| q \right]_j \\ &\times \left[ \begin{matrix} q^{1+j}a, q^{1+j}c \\ q, q^{1+2j}ac \end{matrix} \middle| q^2 \right]_{m+n-j}. \end{aligned}$$

*Proof.* Employing the substitution  $\sqrt{c} \rightarrow -q^{-m}\sqrt{c}$  in (14), we get

$$\begin{aligned} & {}_4\phi_3 \left[ \begin{matrix} q^{-n}, a, \sqrt{c}, -q^{-m}\sqrt{c} \\ \sqrt{q^{1-n}a}, -\sqrt{q^{1-n}a}, q^{-m}c \end{matrix} \middle| q; q \right] \\ &= \sum_{j=0}^m (-1)^j q^{\binom{j+1}{2}} c^{j/2} \left[ \begin{matrix} q^{-m}, q^{-n}, a, q^{-m}\sqrt{c}, -q^{-m}\sqrt{c} \\ q, \sqrt{q^{1-n}a}, -\sqrt{q^{1-n}a}, q^{-m}c, q^{j-2m-1}c \end{matrix} \middle| q \right]_j \\ &\times {}_4\phi_3 \left[ \begin{matrix} q^{j-n}, q^j a, q^{j-m}\sqrt{c}, -q^{j-m}\sqrt{c} \\ q^j \sqrt{q^{1-n}a}, -q^j \sqrt{q^{1-n}a}, q^{2j-2m}c \end{matrix} \middle| q; q \right]. \end{aligned}$$

Letting  $c \rightarrow q^{-2n}$ ,  $q^{-n} \rightarrow c$  in the last relation, the result reads as

$$\begin{aligned} & {}_4\phi_3 \left[ \begin{matrix} a, c, q^{-n}, -q^{-m-n} \\ \sqrt{qac}, -\sqrt{qac}, q^{-m-2n} \end{matrix} \middle| q; q \right] \\ &= \sum_{j=0}^m (-1)^j q^{\binom{j+1}{2}-nj} \left[ \begin{matrix} q^{-m}, q^{-m-n}, -q^{-m-n}, a, c \\ q, \sqrt{qac}, -\sqrt{qac}, q^{-m-2n}, q^{j-2m-2n-1} \end{matrix} \middle| q \right]_j \\ &\times {}_4\phi_3 \left[ \begin{matrix} q^j a, q^j c, q^{j-m-n}, -q^{j-m-n} \\ q^j \sqrt{qac}, -q^j \sqrt{qac}, q^{2j-2m-2n} \end{matrix} \middle| q; q \right]. \end{aligned}$$

Evaluating the  ${}_4\phi_3$ -series on the last line by (3), we achieve Proposition 4 to finish the proof.  $\square$

**Example 5** ( $m = 1$  in Proposition 4).

$$\begin{aligned} & {}_4\phi_3 \left[ \begin{matrix} a, c, q^{-n}, -q^{-1-n} \\ \sqrt{qac}, -\sqrt{qac}, q^{-1-2n} \end{matrix} \middle| q; q \right] \\ &= \left[ \begin{matrix} qa, qc \\ q, qac \end{matrix} \middle| q^2 \right]_{n+1} - q^{n+1} \left[ \begin{matrix} a, c \\ q, qac \end{matrix} \middle| q^2 \right]_{n+1}. \end{aligned}$$

**Theorem 5.** For two complex numbers  $\{a, c\}$  and two nonnegative integers  $\{\ell, m\}$  with  $m \leq n$ , there holds

$$\begin{aligned} & {}_4\phi_3 \left[ \begin{matrix} q^\ell a, c, q^{-n}, -q^{m-n} \\ \sqrt{qac}, -q^\ell \sqrt{qac}, q^{m-2n} \end{matrix} \middle| q; q \right] = \left[ \begin{matrix} q^\ell a, \sqrt{qa/c} \\ q^\ell a/c, \sqrt{qac} \end{matrix} \middle| q \right]_\ell \\ &\times \sum_{i=0}^{\ell} \sum_{j=0}^m q^{\frac{5i}{2}+(m-n)j+\binom{j+1}{2}} \frac{1}{(ac)^{i/2}} \frac{1-q^{2\ell-2i}a/c}{1-q^{2\ell}a/c} \left[ \begin{matrix} q^{-\ell}, c, q^{-2\ell}c/a \\ q, q^{1-2\ell}/a, q^{1-\ell}c/a \end{matrix} \middle| q \right]_i \\ &\times \left[ \begin{matrix} q^{-m}, q^{-n}, -q^{-n}, q^{2\ell-i}a, q^i c \\ q, q^\ell \sqrt{qac}, -q^\ell \sqrt{qac}, q^{m-2n}, q^{j-2n-1} \end{matrix} \middle| q \right]_j \left[ \begin{matrix} q^{1+2\ell-i+j}a, q^{1+i+j}c \\ q, q^{1+2\ell+2j}ac \end{matrix} \middle| q^2 \right]_{n-j}. \end{aligned}$$

*Proof.* Performing the replacement  $a \rightarrow q^{-n-1}a$  in (10), we attain

$$\begin{aligned} {}_4\phi_3 & \left[ \begin{matrix} q^{-n}, q^\ell a, \sqrt{c}, -q^v \sqrt{c} \\ \sqrt{q^{1-n}a}, -q^\ell \sqrt{q^{1-n}a}, q^v c \end{matrix} \middle| q; q \right] = \left[ \begin{matrix} q^\ell a, \sqrt{q^{1+n}a} \\ q^{\ell+n} a, \sqrt{q^{1-n}a} \end{matrix} \middle| q \right]_\ell \\ & \times \sum_{i=0}^{\ell} \frac{q^{(5+n)i/2}}{a^{i/2}} \frac{1 - q^{2\ell+n-2i}a}{1 - q^{2\ell+n}a} \left[ \begin{matrix} q^{-\ell}, q^{-n}, q^{-2\ell-n}/a \\ q, q^{1-\ell-n}/a, q^{1-2\ell}/a \end{matrix} \middle| q \right]_i \\ & \times {}_4\phi_3 \left[ \begin{matrix} q^{i-n}, q^{2\ell-i} a, \sqrt{c}, -q^v \sqrt{c} \\ q^\ell \sqrt{q^{1-n}a}, -q^\ell \sqrt{q^{1-n}a}, q^v c \end{matrix} \middle| q; q \right]. \end{aligned}$$

Replacing  $q^{-n}$  and  $c$  by  $c$  and  $q^{-2n}$ , respectively, in the last relation, the result reads as

$$\begin{aligned} {}_4\phi_3 & \left[ \begin{matrix} q^\ell a, c, q^{-n}, -q^{v-n} \\ \sqrt{qac}, -q^\ell \sqrt{qac}, q^{v-2n} \end{matrix} \middle| q; q \right] = \left[ \begin{matrix} q^\ell a, \sqrt{qa/c} \\ q^\ell a/c, \sqrt{qac} \end{matrix} \middle| q \right]_\ell \\ & \times \sum_{i=0}^{\ell} \frac{q^{5i/2}}{(ac)^{i/2}} \frac{1 - q^{2\ell-2i}a/c}{1 - q^{2\ell}a/c} \left[ \begin{matrix} q^{-\ell}, c, q^{-2\ell}c/a \\ q, q^{1-2\ell}/a, q^{1-\ell}c/a \end{matrix} \middle| q \right]_i \\ & \times {}_4\phi_3 \left[ \begin{matrix} q^{2\ell-i} a, q^i c, q^{-n}, -q^{v-n} \\ q^\ell \sqrt{qac}, -q^\ell \sqrt{qac}, q^{v-2n} \end{matrix} \middle| q; q \right]. \end{aligned} \quad (15)$$

Taking  $v = m$  in (15) and calculating the  ${}_4\phi_3$ -series on the right hand side by Proposition 3, we obtain Theorem 5 to complete the proof.  $\square$

**Example 6** ( $\ell = 1, m = 0$  in Theorem 5).

$$\begin{aligned} {}_4\phi_3 & \left[ \begin{matrix} qa, c, q^{-n}, -q^{-n} \\ \sqrt{qac}, -\sqrt{q^3ac}, q^{-2n} \end{matrix} \middle| q; q \right] \\ & = \frac{1 - qa}{(1 - \sqrt{qac})(1 + \sqrt{qa/c})} \left[ \begin{matrix} q^3 a, qc \\ q, q^3 ac \end{matrix} \middle| q^2 \right]_n \\ & + \frac{1 - c}{(1 - \sqrt{qac})(1 + \sqrt{c/qa})} \left[ \begin{matrix} q^2 a, q^2 c \\ q, q^3 ac \end{matrix} \middle| q^2 \right]_n. \end{aligned}$$

**Example 7** ( $\ell = 1, m = 1$  in Theorem 5:  $n \geq 1$ ).

$$\begin{aligned} {}_4\phi_3 & \left[ \begin{matrix} qa, c, q^{-n}, -q^{1-n} \\ \sqrt{qac}, -\sqrt{q^3ac}, q^{1-2n} \end{matrix} \middle| q; q \right] \\ & = \frac{(1 - \sqrt{q^{1+2n}ac})(1 + \sqrt{q^{1+2n}a/c})}{(1 - \sqrt{qac})(1 + \sqrt{qa/c})} \left[ \begin{matrix} qa, qc \\ q, q^3 ac \end{matrix} \middle| q^2 \right]_n \\ & + \frac{(1 - \sqrt{q^{1+2n}ac})(q^n + \sqrt{qa/c})}{(1 - \sqrt{qac})(1 + \sqrt{qa/c})} \left[ \begin{matrix} q^2 a, c \\ q, q^3 ac \end{matrix} \middle| q^2 \right]_n. \end{aligned}$$

Employing the substitution  $\sqrt{a} \rightarrow -q^{-\ell}\sqrt{a}$  in Theorem 5, we gain the equation

$$\begin{aligned} {}_4\phi_3 & \left[ \begin{matrix} q^{-\ell} a, c, q^{-n}, -q^{m-n} \\ \sqrt{qac}, -q^{-\ell} \sqrt{qac}, q^{m-2n} \end{matrix} \middle| q; q \right] = \left[ \begin{matrix} q/a, -\sqrt{qc/a} \\ qc/a, -\sqrt{q/ac} \end{matrix} \middle| q \right]_\ell \\ & \times \sum_{i=0}^{\ell} \sum_{j=0}^m (-1)^i q^{\frac{(1+2\ell)i}{2} + (m-n)j + \binom{j+1}{2}} \frac{1}{(ac)^{i/2}} \frac{1 - q^{2i}c/a}{1 - c/a} \left[ \begin{matrix} q^{-\ell}, c, c/a \\ q, q/a, q^{1+\ell}c/a \end{matrix} \middle| q \right]_i \\ & \times \left[ \begin{matrix} q^{-m}, q^{-n}, -q^{-n}, q^{-i} a, q^i c \\ q, \sqrt{qac}, -\sqrt{qac}, q^{m-2n}, q^{j-2n-1} \end{matrix} \middle| q \right]_j \left[ \begin{matrix} q^{1-i+j} a, q^{1+i+j} c \\ q, q^{1+2j} ac \end{matrix} \middle| q^2 \right]_{n-j}. \end{aligned} \quad (16)$$

Setting  $v = -m$  in (15) and evaluating the  ${}_4\phi_3$ -series on the right hand side by Proposition 4, we get the following theorem.

**Theorem 6.** For two complex numbers  $\{a, c\}$  and two nonnegative integers  $\{\ell, m\}$ , there holds

$$\begin{aligned} {}_4\phi_3 \left[ \begin{matrix} q^\ell a, c, q^{-n}, -q^{-m-n} \\ \sqrt{qac}, -q^\ell \sqrt{qac}, q^{-m-2n} \end{matrix} \middle| q; q \right] &= \left[ \begin{matrix} q^\ell a, \sqrt{qa/c} \\ q^\ell a/c, \sqrt{qac} \end{matrix} \middle| q \right]_\ell \\ &\times \sum_{i=0}^{\ell} \sum_{j=0}^m (-1)^j q^{\frac{5i}{2}-nj+\binom{j+1}{2}} \frac{1}{(ac)^{i/2}} \frac{1-q^{2\ell-2i}a/c}{1-q^{2\ell}a/c} \left[ \begin{matrix} q^{-\ell}, c, q^{-2\ell}c/a \\ q, q^{1-2\ell}/a, q^{1-\ell}c/a \end{matrix} \middle| q \right]_i \\ &\times \left[ \begin{matrix} q^{-m}, q^{-m-n}, -q^{-m-n}, q^{2\ell-i}a, q^i c \\ q, q^\ell \sqrt{qac}, -q^\ell \sqrt{qac}, q^{-m-2n}, q^{j-2m-2n-1} \end{matrix} \middle| q \right]_j \left[ \begin{matrix} q^{1+2\ell-i+j}a, q^{1+i+j}c \\ q, q^{1+2\ell+2j}ac \end{matrix} \middle| q^2 \right]_{m+n-j}. \end{aligned}$$

**Example 8** ( $\ell = 1, m = 1$  in Theorem 6).

$$\begin{aligned} {}_4\phi_3 \left[ \begin{matrix} qa, c, q^{-n}, -q^{-1-n} \\ \sqrt{qac}, -\sqrt{q^3ac}, q^{-1-2n} \end{matrix} \middle| q; q \right] &= \frac{(1 + \sqrt{q^{3+2n}ac})(1 - \sqrt{q^{3+2n}a/c})}{(1 - \sqrt{qac})(1 + \sqrt{qa/c})} \left[ \begin{matrix} qa, qc \\ q, q^3ac \end{matrix} \middle| q^2 \right]_{1+n} \\ &- \frac{(1 + \sqrt{q^{3+2n}ac})(q^{1+n} - \sqrt{qa/c})}{(1 - \sqrt{qac})(1 + \sqrt{qa/c})} \left[ \begin{matrix} q^2a, c \\ q, q^3ac \end{matrix} \middle| q^2 \right]_{1+n}. \end{aligned}$$

Letting  $\sqrt{a} \rightarrow -q^{-\ell}\sqrt{a}$  in Theorem 6, we achieve the formula

$$\begin{aligned} {}_4\phi_3 \left[ \begin{matrix} q^{-\ell}a, c, q^{-n}, -q^{-m-n} \\ \sqrt{qac}, -q^{-\ell}\sqrt{qac}, q^{-m-2n} \end{matrix} \middle| q; q \right] &= \left[ \begin{matrix} q/a, -\sqrt{qc/a} \\ qc/a, -\sqrt{q/ac} \end{matrix} \middle| q \right]_\ell \\ &\times \sum_{i=0}^{\ell} \sum_{j=0}^m (-1)^{i+j} q^{\frac{(1+2\ell)i}{2}-nj+\binom{j+1}{2}} \frac{1}{(ac)^{i/2}} \frac{1-q^{2i}c/a}{1-c/a} \left[ \begin{matrix} q^{-\ell}, c, c/a \\ q, q/a, q^{1+\ell}c/a \end{matrix} \middle| q \right]_i \\ &\times \left[ \begin{matrix} q^{-m}, q^{-m-n}, -q^{-m-n}, q^{-i}a, q^i c \\ q, \sqrt{qac}, -\sqrt{qac}, q^{-m-2n}, q^{j-2m-2n-1} \end{matrix} \middle| q \right]_j \left[ \begin{matrix} q^{1-i+j}a, q^{1+i+j}c \\ q, q^{1+2j}ac \end{matrix} \middle| q^2 \right]_{m+n-j}. \quad (17) \end{aligned}$$

It should be pointed out that the corresponding hypergeometric series identities of Theorem 5 and (16) are different. The same discuss is adapt to Theorem 6 and (17).

### 3. EXTENSIONS OF $q$ -DIXON FORMULAS

#### 3.1. Extensions of $q$ -Bailey-Dixon formula.

**Theorem 7.** For two complex numbers  $\{a, c\}$  and two nonnegative integers  $\{\ell, m\}$ , there holds

$$\begin{aligned} {}_4\phi_3 \left[ \begin{matrix} q^{-n}, a, c, -q^{1+\ell+m-n}/ac \\ q^{1+\ell-n}/a, q^{1+m-n}/c, -ac \end{matrix} \middle| q; q \right] &= \left[ \begin{matrix} q^{1+\ell-n}/a^2, -q/a \\ q^{1+\ell}/a^2, -q^{1-n}/a \end{matrix} \middle| q \right]_\ell \left[ \begin{matrix} -a, q^{-m}c^2 \\ -ac, q^{-m}c \end{matrix} \middle| q \right]_n \\ &\times \sum_{i=0}^{\ell} \sum_{j=0}^m (-1)^{i+j} q^{ni+(n-i-j)(j-m)+\binom{j+1}{2}} a^i c^{n-i} \frac{1-q^{2i-2\ell-1}a^2}{1-q^{-2\ell-1}a^2} \\ &\times \left[ \begin{matrix} q^{-\ell}, q^{-n}, q^{-2\ell-1}a^2 \\ q, q^{n-2\ell}a^2, q^{-\ell}a^2 \end{matrix} \middle| q \right]_i \left[ \begin{matrix} q^{-m}, q^{i-n}, q^{1+2\ell-n-i}/a^2, q^{-m}c, -q^{-m}c \\ q, q^{1+\ell-n}/a, -q^{1+\ell-n}/a, q^{-m}c^2, q^{j-2m-1}c^2 \end{matrix} \middle| q \right]_j \\ &\times \left[ \begin{matrix} q, q^{2+2\ell+2m-2n}/a^2c^2 \\ q^{2+2\ell-2n+2j}/a^2, q^{1-2m+2j}c^2 \end{matrix} \middle| q^2 \right]_{n-i-j} \chi(n-i-j). \end{aligned}$$

*Proof.* In terms of (1), we attain the relation

$$\begin{aligned} {}_4\phi_3 \left[ \begin{matrix} q^{-n}, a, c, -q^{1+u+v-n}/ac \\ q^{1+u-n}/a, q^{1+v-n}/c, -ac \end{matrix} \middle| q; q \right] &= \left[ \begin{matrix} -a, q^{-v}c^2 \\ -ac, q^{-v}c \end{matrix} \middle| q \right]_n \\ &\times {}_4\phi_3 \left[ \begin{matrix} q^{-n}, q^{1+u-n}/a^2, c, -q^{-v}c \\ q^{1+u-n}/a, -q^{1-n}/a, q^{-v}c^2 \end{matrix} \middle| q; q \right]. \quad (18) \end{aligned}$$

Taking  $u = \ell$ ,  $v = m$  in (18) and calculating the  ${}_4\phi_3$ -series on the right hand side by (12), we obtain Theorem 7 to finish the proof.  $\square$

**Example 9** ( $\ell = 0, m = 1$  in Theorem 7).

$$\begin{aligned} {}_4\phi_3 & \left[ \begin{matrix} q^{-n}, a, c, -q^{2-n}/ac \\ q^{1-n}/a, q^{2-n}/c, -ac \end{matrix} \middle| q; q \right] \\ &= \begin{cases} \left[ \begin{matrix} -a, c^2/q \\ -ac, c/q \end{matrix} \right]_{2s} \left[ \begin{matrix} q, q^{2s-2}a^2c^2 \\ c^2/q, q^{2s}a^2 \end{matrix} \right]_s, & n = 2s; \\ \frac{(q-1)c}{q-c^2} \left[ \begin{matrix} -a, c^2/q \\ -ac, c/q \end{matrix} \right]_{1+2s} \left[ \begin{matrix} q^3, q^{2s}a^2c^2 \\ qc^2, q^{2+2s}a^2 \end{matrix} \right]_s, & n = 1 + 2s. \end{cases} \end{aligned}$$

**Example 10** ( $\ell = 1, m = 1$  in Theorem 7).

$$\begin{aligned} {}_4\phi_3 & \left[ \begin{matrix} q^{-n}, a, c, -q^{3-n}/ac \\ q^{2-n}/a, q^{2-n}/c, -ac \end{matrix} \middle| q; q \right] \\ &= \begin{cases} \frac{(q^2+ac)(q^2-q^{2s}a^2)}{(q^2-a^2)(q^2+q^{2s}ac)} \left[ \begin{matrix} -a/q, c^2/q \\ -ac, c/q \end{matrix} \right]_{2s} \left[ \begin{matrix} q, q^{2s-2}a^2c^2 \\ c^2/q, q^{2s-2}a^2 \end{matrix} \right]_s, & n = 2s; \\ \frac{(a+c)(q-1)(q-q^{2s}ac)}{(q^2-a^2)(1-c^2/q^2)} \left[ \begin{matrix} -a/q, c^2/q^2 \\ -ac, c/q \end{matrix} \right]_{1+2s} \left[ \begin{matrix} q^3, q^{2s-2}a^2c^2 \\ c^2/q, q^{2s}a^2 \end{matrix} \right]_s, & n = 1 + 2s. \end{cases} \end{aligned}$$

Setting  $u = \ell$ ,  $v = -m$  in (18) and evaluating the  ${}_4\phi_3$ -series on the right hand side by Theorem 2, we gain the following theorem.

**Theorem 8.** For two complex numbers  $\{a, c\}$  and two nonnegative integers  $\{\ell, m\}$ , there holds

$$\begin{aligned} {}_4\phi_3 & \left[ \begin{matrix} q^{-n}, a, c, -q^{1+\ell-m-n}/ac \\ q^{1+\ell-n}/a, q^{1-m-n}/c, -ac \end{matrix} \middle| q; q \right] = \left[ \begin{matrix} q^{1+\ell-n}/a^2, -q/a \\ q^{1+\ell}/a^2, -q^{1-n}/a \end{matrix} \right]_\ell \left[ \begin{matrix} -a, q^m c^2 \\ -ac, q^m c \end{matrix} \right]_n \\ & \times \sum_{i=0}^{\ell} \sum_{j=0}^m (-1)^i q^{ni+(m+n-i)j-\binom{j}{2}} a^i c^{n-i} \frac{1-q^{2i-2\ell-1}a^2}{1-q^{-2\ell-1}a^2} \\ & \times \left[ \begin{matrix} q^{-\ell}, q^{-n}, q^{-2\ell-1}a^2 \\ q, q^{1-2\ell}a^2, q^{-\ell}a^2 \end{matrix} \right]_i \left[ \begin{matrix} q^{-m}, q^{i-n}, q^{1+2\ell-n-i}/a^2, c, -c \\ q, q^{1+\ell-n}/a, -q^{1+\ell-n}/a, q^m c^2, q^{j-1}c^2 \end{matrix} \right]_j \\ & \times \left[ \begin{matrix} q, q^{2+2\ell-2n}/a^2c^2 \\ q^{2+2\ell-2n+2j}/a^2, q^{1+2j}c^2 \end{matrix} \right]_{\frac{n-i-j}{2}} \chi(n-i-j). \end{aligned}$$

**Example 11** ( $\ell = 0, m = 1$  in Theorem 8).

$$\begin{aligned} {}_4\phi_3 & \left[ \begin{matrix} q^{-n}, a, c, -q^{-n}/ac \\ q^{1-n}/a, q^{-n}/c, -ac \end{matrix} \middle| q; q \right] \\ &= \begin{cases} \left[ \begin{matrix} -a, qc^2 \\ -ac, qc \end{matrix} \right]_{2s} \left[ \begin{matrix} q, q^{2s}a^2c^2 \\ qc^2, q^{2s}a^2 \end{matrix} \right]_s, & n = 2s; \\ \frac{(1-q)c}{1-qc^2} \left[ \begin{matrix} -a, qc^2 \\ -ac, qc \end{matrix} \right]_{1+2s} \left[ \begin{matrix} q^3, q^{2+2s}a^2c^2 \\ q^3c^2, q^{2+2s}a^2 \end{matrix} \right]_s, & n = 1 + 2s. \end{cases} \end{aligned}$$

**Example 12** ( $\ell = 1, m = 1$  in Theorem 8).

$$\begin{aligned} {}_4\phi_3 & \left[ \begin{matrix} q^{-n}, a, c, -q^{1-n}/ac \\ q^{2-n}/a, q^{-n}/c, -ac \end{matrix} \middle| q; q \right] \\ &= \begin{cases} \frac{(q^2-a^2c^2)(1-q^{2s-2}a^2)}{(q^2-a^2)(1-q^{2s-2}a^2c^2)} \left[ \begin{matrix} -a/q, qc^2 \\ -ac/q, qc \end{matrix} \right]_{2s} \left[ \begin{matrix} q, q^{2s-2}a^2c^2 \\ qc^2, q^{2s-2}a^2 \end{matrix} \right]_s, & n = 2s; \\ \frac{(1-q)(qc-a)}{(q-a)(1-qc)} \left[ \begin{matrix} -a, qc^2 \\ -ac, q^2c \end{matrix} \right]_{2s} \left[ \begin{matrix} q^3, q^{2s}a^2c^2 \\ qc^2, q^{2s}a^2 \end{matrix} \right]_s, & n = 1 + 2s. \end{cases} \end{aligned}$$

Taking  $u = -\ell$ ,  $v = -m$  in (18) and calculating the  ${}_4\phi_3$ -series on the right hand side by (11), we get the following theorem.

**Theorem 9.** *For two complex numbers  $\{a, c\}$  and two nonnegative integers  $\{\ell, m\}$ , there holds*

$$\begin{aligned} {}_4\phi_3 \left[ \begin{matrix} q^{-n}, a, c, -q^{1-\ell-m-n}/ac \\ q^{1-\ell-n}/a, q^{1-m-n}/c, -ac \end{matrix} \middle| q; q \right] &= \left[ \begin{matrix} a, q^n a^2 \\ a^2, q^n a \end{matrix} \middle| q \right]_{\ell} \left[ \begin{matrix} -a, q^m c^2 \\ -ac, q^m c \end{matrix} \middle| q \right]_n \\ &\times \sum_{i=0}^{\ell} \sum_{j=0}^m q^{(\ell+n)i+(m+n-i)j-\binom{j}{2}} a^i c^{n-i} \frac{1-q^{2i-1}a^2}{1-q^{-1}a^2} \\ &\times \left[ \begin{matrix} q^{-\ell}, q^{-n}, q^{-1}a^2 \\ q, q^n a^2, q^\ell a^2 \end{matrix} \middle| q \right]_i \left[ \begin{matrix} q^{-m}, q^{i-n}, q^{1-n-i}/a^2, c, -c \\ q, q^{1-n}/a, -q^{1-n}/a, q^m c^2, q^{j-1} c^2 \end{matrix} \middle| q \right]_j \\ &\times \left[ \begin{matrix} q, q^{2-2n}/a^2 c^2 \\ q^{2-2n+2j}/a^2, q^{1+2j} c^2 \end{matrix} \middle| q^2 \right]_{\frac{n-i-j}{2}} \chi(n-i-j). \end{aligned}$$

**Example 13** ( $\ell = 1, m = 1$  in Theorem 9).

$$\begin{aligned} {}_4\phi_3 \left[ \begin{matrix} q^{-n}, a, c, -q^{-1-n}/ac \\ q^{-n}/a, q^{-n}/c, -ac \end{matrix} \middle| q; q \right] &= \begin{cases} \left[ \begin{matrix} -qa, qc^2 \\ -qac, qc \end{matrix} \middle| q \right]_{2s} \left[ \begin{matrix} q, q^{2s+2} a^2 c^2 \\ qc^2, q^{2s+2} a^2 \end{matrix} \middle| q^2 \right]_s, & n = 2s; \\ \frac{a+c}{1+ac} \left[ \begin{matrix} -qa, qc^2 \\ -qac, qc \end{matrix} \middle| q \right]_{1+2s} \left[ \begin{matrix} q, q^{2s+2} a^2 c^2 \\ qc^2, q^{2s+2} a^2 \end{matrix} \middle| q^2 \right]_{1+s}, & n = 1 + 2s. \end{cases} \end{aligned}$$

### 3.2. Extensions of another $q$ -Dixon formula.

**Theorem 10.** *For two complex numbers  $\{a, c\}$  and two nonnegative integers  $\{\ell, m\}$ , there holds*

$$\begin{aligned} {}_4\phi_3 \left[ \begin{matrix} a, c, q^{-n}, -q^{1+\ell+m+n}a/c \\ q^{1+\ell}a/c, q^{1+m+n}a, -q^{-n}c \end{matrix} \middle| q; q \right] &= \left[ \begin{matrix} a, -q^{-\ell}c \\ q^{-\ell-1}c^2, -qa/c \end{matrix} \middle| q \right]_{\ell} \left[ \begin{matrix} q^{1+m+n}, -qa/c \\ q^{1+m+n}a, -q/c \end{matrix} \middle| q \right]_n \\ &\times \sum_{i=0}^{\ell} \sum_{j=0}^m (-1)^{i+j} q^{(\binom{j+1}{2}-nj)} \left( \frac{c}{a} \right)^i \frac{1-q^{1+2i}/c^2}{1-q/c^2} \left[ \begin{matrix} q^{-\ell}, q^{1+\ell}a/c^2, q/c^2 \\ q, q^{1-\ell}/a, q^{2+\ell}/c^2 \end{matrix} \middle| q^2 \right]_i \\ &\times \left[ \begin{matrix} q^{-m}, q^{-m-n}, -q^{-m-n}, q^{\ell-i}a, q^{1+\ell+i}a/c^2 \\ q, q^{1+\ell}a/c, -q^{1+\ell}a/c, q^{-m-2n}, q^{j-2m-2n-1} \end{matrix} \middle| q \right]_j \\ &\times \left[ \begin{matrix} q^{1+\ell-i+j}a, q^{2+\ell+i+j}a/c^2 \\ q, q^{2+2\ell+2j}a^2/c^2 \end{matrix} \middle| q^2 \right]_{m+n-j}. \end{aligned}$$

*Proof.* By means of (1), we achieve the relation

$$\begin{aligned} {}_4\phi_3 \left[ \begin{matrix} a, c, q^{-n}, -q^{1+u+v+n}a/c \\ q^{1+u}a/c, q^{1+v+n}a, -q^{-n}c \end{matrix} \middle| q; q \right] &= \left[ \begin{matrix} q^{1+v+n}, -qa/c \\ q^{1+v+n}a, -q/c \end{matrix} \middle| q \right]_n \\ &\times {}_4\phi_3 \left[ \begin{matrix} a, q^{1+u}a/c^2, q^{-n}, -q^{-v-n} \\ q^{1+u}a/c, -qa/c, q^{-v-2n} \end{matrix} \middle| q; q \right]. \quad (19) \end{aligned}$$

Setting  $u = \ell$ ,  $v = m$  in (19) and evaluating the  ${}_4\phi_3$ -series on the right hand side by Theorem 6, we attain Theorem 10 to complete the proof.  $\square$

**Example 14** ( $\ell = 0, m = 1$  in Theorem 10).

$$\begin{aligned} {}_4\phi_3 \left[ \begin{matrix} a, c, q^{-n}, -q^{2+n}a/c \\ qa/c, q^{2+n}a, -q^{-n}c \end{matrix} \middle| q; q \right] &= \left[ \begin{matrix} q^{2+n}, -qa/c \\ q^{2+n}a, -q/c \end{matrix} \middle| q \right]_n \left[ \begin{matrix} qa, q^2 a/c^2 \\ q, q^2 a^2/c^2 \end{matrix} \middle| q^2 \right]_{1+n} \\ &- \left[ \begin{matrix} q^{2+n}, -qa/c \\ q^{2+n}a, -q/c \end{matrix} \middle| q \right]_n \left[ \begin{matrix} a, qa/c^2 \\ q, q^2 a^2/c^2 \end{matrix} \middle| q^2 \right]_{1+n} q^{1+n}. \end{aligned}$$

**Example 15** ( $\ell = 1, m = 0$  in Theorem 10).

$$\begin{aligned} {}_4\phi_3 & \left[ \begin{matrix} a, c, q^{-n}, -q^{2+n}a/c \\ q^2a/c, q^{1+n}a, -q^{-n}c \end{matrix} \middle| q; q \right] \\ &= \frac{qc(1-a)}{(qa+c)(q-c)} \left[ \begin{matrix} q^{1+n}, -qa/c \\ q^{1+n}a, -q/c \end{matrix} \middle| q \right]_n \left[ \begin{matrix} q^2a, q^3a/c^2 \\ q, q^4a^2/c^2 \end{matrix} \middle| q^2 \right]_n \\ &+ \frac{q^2a - c^2}{(qa+c)(q-c)} \left[ \begin{matrix} q^{1+n}, -qa/c \\ q^{1+n}a, -q/c \end{matrix} \middle| q \right]_n \left[ \begin{matrix} qa, q^4a/c^2 \\ q, q^4a^2/c^2 \end{matrix} \middle| q^2 \right]_n. \end{aligned}$$

**Example 16** ( $\ell = 1, m = 1$  in Theorem 10).

$$\begin{aligned} {}_4\phi_3 & \left[ \begin{matrix} a, c, q^{-n}, -q^{3+n}a/c \\ q^2a/c, q^{2+n}a, -q^{-n}c \end{matrix} \middle| q; q \right] \\ &= \frac{q(1+q^n)c(c-q^{2+n}a)}{(qa+c)(q-c)} \left[ \begin{matrix} q^{2+n}, -qa/c \\ q^{2+n}a, -q/c \end{matrix} \middle| q \right]_n \left[ \begin{matrix} a, q^3a/c^2 \\ q, q^4a^2/c^2 \end{matrix} \middle| q^2 \right]_{1+n} \\ &- \frac{(c+q^{2+n})(c-q^{2+n}a)}{(qa+c)(q-c)} \left[ \begin{matrix} q^{2+n}, -qa/c \\ q^{2+n}a, -q/c \end{matrix} \middle| q \right]_n \left[ \begin{matrix} qa, q^2a/c^2 \\ q, q^4a^2/c^2 \end{matrix} \middle| q^2 \right]_{1+n}. \end{aligned}$$

Taking  $u = \ell, v = -m$  in (19) and calculating the  ${}_4\phi_3$ -series on the right hand side by Theorem 5, we obtain the following theorem.

**Theorem 11.** For two complex numbers  $\{a, c\}$  and two nonnegative integers  $\{\ell, m\}$  with  $m \leq n$ , there holds

$$\begin{aligned} {}_4\phi_3 & \left[ \begin{matrix} a, c, q^{-n}, -q^{1+\ell-m+n}a/c \\ q^{1+\ell}a/c, q^{1-m+n}a, -q^{-n}c \end{matrix} \middle| q; q \right] = \left[ \begin{matrix} a, -q^{-\ell}c \\ q^{-\ell-1}c^2, -qa/c \end{matrix} \middle| q \right]_\ell \left[ \begin{matrix} q^{1-m+n}, -qa/c \\ q^{1-m+n}a, -q/c \end{matrix} \middle| q \right]_n \\ &\times \sum_{i=0}^{\ell} \sum_{j=0}^m (-1)^i q^{(m-n)j + \binom{j+1}{2}} \left( \frac{c}{a} \right)^i \frac{1-q^{1+2i}/c^2}{1-q/c^2} \left[ \begin{matrix} q^{-\ell}, q^{1+\ell}a/c^2, q/c^2 \\ q, q^{1-\ell}/a, q^{2+\ell}/c^2 \end{matrix} \middle| q \right]_i \\ &\times \left[ \begin{matrix} q^{-m}, q^{-n}, -q^{-n}, q^{\ell-i}a, q^{1+\ell+i}a/c^2 \\ q, q^{1+\ell}a/c, -q^{1+\ell}a/c, q^{m-2n}, q^{j-2n-1} \end{matrix} \middle| q \right]_j \left[ \begin{matrix} q^{1+\ell-i+j}a, q^{2+\ell+i+j}a/c^2 \\ q, q^{2+2\ell+2j}a^2/c^2 \end{matrix} \middle| q^2 \right]_{n-j}. \end{aligned}$$

**Example 17** ( $\ell = 0, m = 1$  in Theorem 11:  $n \geq 1$ ).

$$\begin{aligned} {}_4\phi_3 & \left[ \begin{matrix} a, c, q^{-n}, -q^n a/c \\ qa/c, q^n a, -q^{-n}c \end{matrix} \middle| q; q \right] \\ &= \left[ \begin{matrix} q^n, -qa/c \\ q^n a, -q/c \end{matrix} \middle| q \right]_n \left[ \begin{matrix} qa, q^2 a/c^2 \\ q, q^2 a^2/c^2 \end{matrix} \middle| q^2 \right]_n \\ &+ \left[ \begin{matrix} q^n, -qa/c \\ q^n a, -q/c \end{matrix} \middle| q \right]_n \left[ \begin{matrix} a, qa/c^2 \\ q, q^2 a^2/c^2 \end{matrix} \middle| q^2 \right]_n q^n. \end{aligned}$$

**Example 18** ( $\ell = 1, m = 1$  in Theorem 11:  $n \geq 1$ ).

$$\begin{aligned} {}_4\phi_3 & \left[ \begin{matrix} a, c, q^{-n}, -q^{1+n}a/c \\ q^2a/c, q^n a, -q^{-n}c \end{matrix} \middle| q; q \right] \\ &= \frac{q - q^n c}{q - c} \left[ \begin{matrix} q^n, -q^2 a/c \\ q^n a, -q/c \end{matrix} \middle| q \right]_n \left[ \begin{matrix} a, q^3 a/c^2 \\ q, q^4 a^2/c^2 \end{matrix} \middle| q^2 \right]_n \\ &+ \frac{q^{1+n} - c}{q - c} \left[ \begin{matrix} q^n, -q^2 a/c \\ q^n a, -q/c \end{matrix} \middle| q \right]_n \left[ \begin{matrix} qa, q^2 a/c^2 \\ q, q^4 a^2/c^2 \end{matrix} \middle| q^2 \right]_n. \end{aligned}$$

Setting  $u = -\ell, v = m$  in (19) and evaluating the  ${}_4\phi_3$ -series on the right hand side by (17), we gain the following theorem.

**Theorem 12.** For two complex numbers  $\{a, c\}$  and two nonnegative integers  $\{\ell, m\}$ , there holds

$$\begin{aligned} {}_4\phi_3 \left[ \begin{matrix} a, c, q^{-n}, -q^{1-\ell+m+n}a/c \\ q^{1-\ell}a/c, q^{1+m+n}a, -q^{-n}c \end{matrix} \middle| q; q \right] &= \left[ \begin{matrix} a, c \\ q^{\ell-1}c^2, q^{1-\ell}a/c \end{matrix} \right]_{\ell} \left[ \begin{matrix} q^{1+m+n}, -qa/c \\ q^{1+m+n}a, -q/c \end{matrix} \middle| q \right]_n \\ &\times \sum_{i=0}^{\ell} \sum_{j=0}^m (-1)^j q^{\ell i - nj + \binom{j+1}{2}} \left( \frac{c}{a} \right)^i \frac{1 - q^{1-2\ell+2i}/c^2}{1 - q^{1-2\ell}/c^2} \left[ \begin{matrix} q^{-\ell}, q^{1-\ell}a/c^2, q^{1-2\ell}/c^2 \\ q, q^{1-\ell}/a, q^{2-\ell}/c^2 \end{matrix} \middle| q \right]_i \\ &\times \left[ \begin{matrix} q^{-m}, q^{-m-n}, -q^{-m-n}, q^{\ell-i}a, q^{1-\ell+i}a/c^2 \\ q, qa/c, -qa/c, q^{-m-2n}, q^{j-2m-2n-1} \end{matrix} \middle| q \right]_j \left[ \begin{matrix} q^{1+\ell-i+j}a, q^{2-\ell+i+j}a/c^2 \\ q, q^{2+2j}a^2/c^2 \end{matrix} \middle| q^2 \right]_{m+n-j}. \end{aligned}$$

**Example 19** ( $\ell = 1, m = 0$  in Theorem 12).

$$\begin{aligned} {}_4\phi_3 \left[ \begin{matrix} a, c, q^{-n}, -q^n a/c \\ a/c, q^{1+n}a, -q^{-n}c \end{matrix} \middle| q; q \right] &= \frac{(1-a)c}{(1+c)(c-a)} \left[ \begin{matrix} q^{1+n}, -qa/c \\ q^{1+n}a, -q/c \end{matrix} \right]_n \left[ \begin{matrix} q^2a, qa/c^2 \\ q, q^2a^2/c^2 \end{matrix} \middle| q^2 \right]_n \\ &+ \frac{c^2 - a}{(1+c)(c-a)} \left[ \begin{matrix} q^{1+n}, -qa/c \\ q^{1+n}a, -q/c \end{matrix} \right]_n \left[ \begin{matrix} qa, q^2a/c^2 \\ q, q^2a^2/c^2 \end{matrix} \middle| q^2 \right]_n. \end{aligned}$$

**Example 20** ( $\ell = 1, m = 1$  in Theorem 12).

$$\begin{aligned} {}_4\phi_3 \left[ \begin{matrix} a, c, q^{-n}, -q^{1+n}a/c \\ a/c, q^{2+n}a, -q^{-n}c \end{matrix} \middle| q; q \right] &= \frac{(qa+c)(1-q^{1+n}c)}{(1+c)(c-a)} \left[ \begin{matrix} q^{2+n}, -q^2a/c \\ q^{2+n}a, -q/c \end{matrix} \right]_n \left[ \begin{matrix} a, qa/c^2 \\ q, q^2a^2/c^2 \end{matrix} \middle| q^2 \right]_{1+n} \\ &+ \frac{(qa+c)(c-q^{1+n})}{(1+c)(c-a)} \left[ \begin{matrix} q^{2+n}, -q^2a/c \\ q^{2+n}a, -q/c \end{matrix} \right]_n \left[ \begin{matrix} qa, a/c^2 \\ q, q^2a^2/c^2 \end{matrix} \middle| q^2 \right]_{1+n}. \end{aligned}$$

Taking  $u = -\ell$ ,  $v = -m$  in (19) and calculating the  ${}_4\phi_3$ -series on the right hand side by (16), we get the following theorem.

**Theorem 13.** For two complex numbers  $\{a, c\}$  and two nonnegative integers  $\{\ell, m\}$  with  $m \leq n$ , there holds

$$\begin{aligned} {}_4\phi_3 \left[ \begin{matrix} a, c, q^{-n}, -q^{1-\ell-m+n}a/c \\ q^{1-\ell}a/c, q^{1-m+n}a, -q^{-n}c \end{matrix} \middle| q; q \right] &= \left[ \begin{matrix} a, c \\ q^{\ell-1}c^2, q^{1-\ell}a/c \end{matrix} \right]_{\ell} \left[ \begin{matrix} q^{1-m+n}, -qa/c \\ q^{1-m+n}a, -q/c \end{matrix} \middle| q \right]_n \\ &\times \sum_{i=0}^{\ell} \sum_{j=0}^m q^{\ell i + (m-n)j + \binom{j+1}{2}} \left( \frac{c}{a} \right)^i \frac{1 - q^{1-2\ell+2i}/c^2}{1 - q^{1-2\ell}/c^2} \left[ \begin{matrix} q^{-\ell}, q^{1-\ell}a/c^2, q^{1-2\ell}/c^2 \\ q, q^{1-\ell}/a, q^{2-\ell}/c^2 \end{matrix} \middle| q \right]_i \\ &\times \left[ \begin{matrix} q^{-m}, q^{-n}, -q^{-n}, q^{\ell-i}a, q^{1-\ell+i}a/c^2 \\ q, qa/c, -qa/c, q^{m-2n}, q^{j-2n-1} \end{matrix} \middle| q \right]_j \left[ \begin{matrix} q^{1+\ell-i+j}a, q^{2-\ell+i+j}a/c^2 \\ q, q^{2+2j}a^2/c^2 \end{matrix} \middle| q^2 \right]_{n-j}. \end{aligned}$$

**Example 21** ( $\ell = 1, m = 1$  in Theorem 13:  $n \geq 1$ ).

$$\begin{aligned} {}_4\phi_3 \left[ \begin{matrix} a, c, q^{-n}, -q^{n-1}a/c \\ a/c, q^n a, -q^{-n}c \end{matrix} \middle| q; q \right] &= \left[ \begin{matrix} q^n, -a/c \\ q^n a, -1/c \end{matrix} \right]_n \left[ \begin{matrix} qa, a/c^2 \\ q, a^2/c^2 \end{matrix} \middle| q^2 \right]_n \\ &+ \frac{1+q^n c}{1+c} \left[ \begin{matrix} q^n, -a/c \\ q^n a, -q/c \end{matrix} \right]_n \left[ \begin{matrix} a, qa/c^2 \\ q, a^2/c^2 \end{matrix} \middle| q^2 \right]_n. \end{aligned}$$

#### 4. EXTENSIONS OF $q$ -WHIPPLE FORMULAS

##### 4.1. Extensions of $q$ -Andrews-Whipple formula.

**Theorem 14.** For two complex numbers  $\{a, c\}$  and two nonnegative integers  $\{\ell, m\}$ , there holds

$$\begin{aligned} {}_4\phi_3 \left[ \begin{matrix} q^{-n}, q^{1+\ell+n}, \sqrt{qac}, -q^m \sqrt{qac} \\ -q, q^{1+\ell+m}a, qc \end{matrix} \middle| q; q \right] &= \left[ \begin{matrix} q^{-n} \sqrt{qa/c}, -q^{m-n} \sqrt{qa/c}, q^{1+\ell+n} \\ q^{1+\ell+m}a, q^{-n}/c, -q \end{matrix} \middle| q \right]_n \\ &\times \left[ \begin{matrix} q^{m-n}a, \sqrt{qac} \\ q^m ac, q^{-n} \sqrt{qa/c} \end{matrix} \middle| q \right]_m \sum_{i=0}^{\ell} \sum_{j=0}^m (-1)^i q^{(n+\frac{5}{2})j-n+i+1} \binom{i+1}{2} \left( \frac{c}{a} \right)^{\frac{j}{2}} \frac{1-q^{2m-2j}ac}{1-q^{2m}ac} \\ &\times \left[ \begin{matrix} q^{-\ell}, q^{-\ell-n}, -q^{-\ell-n}, q^{2m-n-j}a, q^{j-n}/c \\ q, q^{m-n} \sqrt{qa/c}, -q^{m-n} \sqrt{qa/c}, q^{-\ell-2n}, q^{i-2\ell-2n-1} \end{matrix} \middle| q \right]_i \left[ \begin{matrix} q^{-m}, q^{-n}/c, q^{-2m}/ac \\ q, q^{1-2m+n}/a, q^{1-m}/ac \end{matrix} \middle| q \right]_j \\ &\times \left[ \begin{matrix} q^{1+2m-n+i-j}a, q^{1-n+i+j}/c \\ q, q^{1+2m-2n+2i}a/c \end{matrix} \middle| q^2 \right]_{\ell+n-i}. \end{aligned}$$

*Proof.* The iteration of (1) produces the transformation formula:

$${}_4\phi_3 \left[ \begin{matrix} q^{-n}, a, b, c \\ d, e, q^{1-n}abc/de \end{matrix} \middle| q; q \right] = \left[ \begin{matrix} c, de/ac, de/bc \\ d, e, de/abc \end{matrix} \middle| q \right]_n {}_4\phi_3 \left[ \begin{matrix} q^{-n}, d/c, e/c, de/abc \\ de/ac, de/bc, q^{1-n}/c \end{matrix} \middle| q; q \right].$$

In terms of the last identity, we achieve the relation

$$\begin{aligned} {}_4\phi_3 \left[ \begin{matrix} q^{-n}, q^{1+n+u}, \sqrt{qac}, -q^v \sqrt{qac} \\ -q, q^{1+u+v}a, qc \end{matrix} \middle| q; q \right] \\ = \left[ \begin{matrix} q^{-n} \sqrt{qa/c}, -q^{v-n} \sqrt{qa/c}, q^{1+n+u} \\ q^{1+u+v}a, q^{-n}/c, -q \end{matrix} \middle| q \right]_n \\ \times {}_4\phi_3 \left[ \begin{matrix} q^{v-n}a, q^{-n}/c, q^{-n}, -q^{-u-n} \\ q^{-n} \sqrt{qa/c}, -q^{v-n} \sqrt{qa/c}, q^{-u-2n} \end{matrix} \middle| q; q \right]. \end{aligned} \quad (20)$$

Setting  $u = \ell$ ,  $v = m$  in (20) and evaluating the  ${}_4\phi_3$ -series on the right hand side by Theorem 6, we attain Theorem 14 to finish the proof.  $\square$

**Example 22** ( $\ell = 0, m = 1$  in Theorem 14).

$$\begin{aligned} {}_4\phi_3 \left[ \begin{matrix} q^{-n}, q^{1+n}, \sqrt{qac}, -\sqrt{q^3ac} \\ -q, q^2a, qc \end{matrix} \middle| q; q \right] \\ = \frac{q^{(\frac{1+n}{2})}}{(1 + \sqrt{qac})(1 - \sqrt{qa/c})} \frac{(q^{1-n}a; q^2)_{1+n} (q^{1-n}c; q^2)_n}{(q^2a; q)_n (qc; q)_n} \\ + \frac{q^{(\frac{1+n}{2})}}{(1 + \sqrt{qac})(1 - \sqrt{c/q})} \frac{(q^{2-n}a; q^2)_n (q^{-n}c; q^2)_{1+n}}{(q^2a; q)_n (qc; q)_n}. \end{aligned}$$

**Example 23** ( $\ell = 1, m = 0$  in Theorem 14).

$$\begin{aligned} {}_4\phi_3 \left[ \begin{matrix} q^{-n}, q^{2+n}, \sqrt{qac}, -\sqrt{qac} \\ -q, q^2a, qc \end{matrix} \middle| q; q \right] \\ = \frac{q^{(\frac{2+n}{2})}}{(qa - c)(1 - q^{1+n})} \frac{(q^{1-n}a; q^2)_{1+n} (q^{-1-n}c; q^2)_{1+n}}{(q^2a; q)_n (qc; q)_n} \\ - \frac{q^{(\frac{2+n}{2})}}{(qa - c)(1 - q^{1+n})} \frac{(q^{-n}a; q^2)_{1+n} (q^{-n}c; q^2)_{1+n}}{(q^2a; q)_n (qc; q)_n}. \end{aligned}$$

**Example 24** ( $\ell = 1, m = 1$  in Theorem 14).

$$\begin{aligned} {}_4\phi_3 \left[ \begin{matrix} q^{-n}, q^{2+n}, \sqrt{qac}, -\sqrt{q^3ac} \\ -q, q^3a, qc \end{matrix} \middle| q; q \right] \\ = \frac{q^{(\frac{2+n}{2})}(1 - \sqrt{q^{-1-2n}ac})}{(\sqrt{qa} - \sqrt{c})(\sqrt{q^3a} - \sqrt{c})(1 + \sqrt{qac})(1 - q^{1+n})} \frac{(q^{2-n}a; q^2)_{1+n} (q^{-n}c; q^2)_{1+n}}{(q^3a; q)_n (qc; q)_n} \\ - \frac{q^{(\frac{2+n}{2})}(1 - \sqrt{q^{3+2n}ac})}{(\sqrt{qa} - \sqrt{c})(\sqrt{q^3a} - \sqrt{c})(1 + \sqrt{qac})(1 - q^{1+n})} \frac{(q^{1-n}a; q^2)_{1+n} (q^{-1-n}c; q^2)_{1+n}}{(q^3a; q)_n (qc; q)_n}. \end{aligned}$$

Taking  $u = -\ell$ ,  $v = m$  in (20) and calculating the  ${}_4\phi_3$ -series on the right hand side by Theorem 5, we obtain the following theorem.

**Theorem 15.** For two complex numbers  $\{a, c\}$  and two nonnegative integers  $\{\ell, m\}$  with  $\ell \leq n$ , there holds

$$\begin{aligned} {}_4\phi_3 & \left[ \begin{matrix} q^{-n}, q^{1-\ell+n}, \sqrt{qac}, -q^m \sqrt{qac} \\ -q, q^{1-\ell+m}a, qc \end{matrix} \middle| q; q \right] = \left[ \begin{matrix} q^{-n}\sqrt{qa/c}, -q^{m-n}\sqrt{qa/c}, q^{1-\ell+n} \\ q^{1-\ell+m}a, q^{-n}/c, -q \end{matrix} \middle| q \right]_n \\ & \times \left[ \begin{matrix} q^{m-n}a, \sqrt{qac} \\ q^m ac, q^{-n}\sqrt{qa/c} \end{matrix} \middle| q \right]_m \sum_{i=0}^{\ell} \sum_{j=0}^m q^{(\ell-n)i + (\frac{i+1}{2}) + (n+\frac{5}{2})j} \left( \frac{c}{a} \right)^{\frac{j}{2}} \frac{1 - q^{2m-2j}ac}{1 - q^{2m}ac} \\ & \times \left[ \begin{matrix} q^{-\ell}, q^{-n}, -q^{-n}, q^{2m-n-j}a, q^{j-n}/c \\ q, q^{m-n}\sqrt{qa/c}, -q^{m-n}\sqrt{qa/c}, q^{\ell-2n}, q^{i-2n-1} \end{matrix} \middle| q \right]_i \left[ \begin{matrix} q^{-m}, q^{-n}/c, q^{-2m}/ac \\ q, q^{1-2m+n}/a, q^{1-m}/ac \end{matrix} \middle| q \right]_j \\ & \times \left[ \begin{matrix} q^{1+2m-n+i-j}a, q^{1-n+i+j}/c \\ q, q^{1+2m-2n+2i}a/c \end{matrix} \middle| q^2 \right]_{n-i}. \end{aligned}$$

**Example 25** ( $\ell = 1, m = 0$  in Theorem 15).

$$\begin{aligned} {}_4\phi_3 & \left[ \begin{matrix} q^{-n}, q^n, \sqrt{qac}, -\sqrt{qac} \\ -q, a, qc \end{matrix} \middle| q; q \right] \\ & = \frac{q^{(\frac{1+n}{2})}}{1+q^n} \frac{(q^{1-n}a; q^2)_n (q^{1-n}c; q^2)_n}{(a; q)_n (qc; q)_n} \\ & + \frac{q^{(\frac{1+n}{2})}}{1+q^n} \frac{(q^{-n}a; q^2)_n (q^{2-n}c; q^2)_n}{(a; q)_n (qc; q)_n}. \end{aligned}$$

**Example 26** ( $\ell = 1, m = 1$  in Theorem 15).

$$\begin{aligned} {}_4\phi_3 & \left[ \begin{matrix} q^{-n}, q^n, \sqrt{qac}, -\sqrt{q^3ac} \\ -q, qa, qc \end{matrix} \middle| q; q \right] \\ & = \frac{q^{(\frac{1+n}{2})}(1 + \sqrt{q^{1+2n}ac})}{(1 + q^n)(1 + \sqrt{qac})} \frac{(q^{1-n}a; q^2)_n (q^{1-n}c; q^2)_n}{(qa; q)_n (qc; q)_n} \\ & + \frac{q^{(\frac{1+n}{2})}(1 + \sqrt{q^{1-2n}ac})}{(1 + q^n)(1 + \sqrt{qac})} \frac{(q^{2-n}a; q^2)_n (q^{2-n}c; q^2)_n}{(qa; q)_n (qc; q)_n}. \end{aligned}$$

#### 4.2. Extensions of $q$ -Jain-Whipple formula.

**Theorem 16.** For two complex numbers  $\{a, c\}$  and two nonnegative integers  $\{\ell, m\}$  with  $m \leq n$ , there holds

$$\begin{aligned} {}_4\phi_3 & \left[ \begin{matrix} a, q^{1+\ell}/a, q^{-n}, -q^{m-n} \\ c, q^{1+\ell+m-2n}/c, -q \end{matrix} \middle| q; q \right] = \left[ \begin{matrix} q^2/ac, -q/a \\ q^2/a^2, -q/c \end{matrix} \middle| q \right]_{\ell} \left[ \begin{matrix} q^{1-m+n}, -q^{-\ell}c \\ q^{n-m-\ell}c, -q \end{matrix} \middle| q \right]_n \\ & \times \sum_{i=0}^{\ell} \sum_{j=0}^m q^{(\ell+1)i + (m-n)j + (\frac{j+1}{2})} \frac{(-1)^i}{c^i} \frac{1 - q^{1+2i}/a^2}{1 - q/a^2} \left[ \begin{matrix} q^{-\ell}, c/a, q/a^2 \\ q, q^2/ac, q^{2+\ell}/a^2 \end{matrix} \middle| q \right]_i \\ & \times \left[ \begin{matrix} q^{-m}, q^{-n}, -q^{-n}, q^{-i-1}ac, q^i c/a \\ q, c, -c, q^{m-2n}, q^{j-2n-1} \end{matrix} \middle| q \right]_j \left[ \begin{matrix} q^{j-i}ac, q^{1+i+j}c/a \\ q, q^{2j}c^2 \end{matrix} \middle| q^2 \right]_{n-j}. \end{aligned}$$

*Proof.* By means of (1), we gain the relation

$$\begin{aligned} {}_4\phi_3 & \left[ \begin{matrix} a, q^{1+u}/a, q^{-n}, -q^{v-n} \\ c, q^{1+u+v-2n}/c, -q \end{matrix} \middle| q; q \right] = \left[ \begin{matrix} q^{1-v+n}, -q^{-u}c \\ q^{n-u-v}c, -q \end{matrix} \middle| q \right]_n \\ & \times {}_4\phi_3 \left[ \begin{matrix} q^{-u-1}ac, c/a, q^{-n}, -q^{v-n} \\ c, -q^{-u}c, q^{v-2n} \end{matrix} \middle| q; q \right]. \quad (21) \end{aligned}$$

Setting  $u = \ell, v = m$  in (21) and evaluating the  ${}_4\phi_3$ -series on the right hand side by (16), we get Theorem 16 to complete the proof.  $\square$

**Example 27** ( $\ell = 0, m = 1$  in Theorem 16).

$$\begin{aligned} {}_4\phi_3 & \left[ \begin{matrix} a, q/a, q^{-n}, -q^{1-n} \\ c, q^{2-2n}/c, -q \end{matrix} \middle| q; q \right] \\ &= \frac{1}{1+q^n} \frac{(ac; q^2)_n (qc/a; q^2)_n}{(c; q)_n (q^{n-1}c; q)_n} \\ &+ \frac{q^n}{1+q^n} \frac{(ac/q; q^2)_n (c/a; q^2)_n}{(c; q)_n (q^{n-1}c; q)_n}. \end{aligned}$$

**Example 28** ( $\ell = 1, m = 0$  in Theorem 16).

$$\begin{aligned} {}_4\phi_3 & \left[ \begin{matrix} a, q^2/a, q^{-n}, -q^{-n} \\ c, q^{2-2n}/c, -q \end{matrix} \middle| q; q \right] \\ &= \frac{(q-c)(q^2-ac)}{(q-a)(q^2-q^{2n}c^2)} \frac{(ac; q^2)_n (qc/a; q^2)_n}{(c/q; q)_{2n}} \\ &+ \frac{q(q-c)(c-a)}{(q-a)(q^2-q^{2n}c^2)} \frac{(ac/q; q^2)_n (q^2c/a; q^2)_n}{(c/q; q)_{2n}}. \end{aligned}$$

**Example 29** ( $\ell = 1, m = 1$  in Theorem 16).

$$\begin{aligned} {}_4\phi_3 & \left[ \begin{matrix} a, q^2/a, q^{-n}, -q^{1-n} \\ c, q^{3-2n}/c, -q \end{matrix} \middle| q; q \right] \\ &= \frac{q-q^n a}{(q-a)(1+q^n)} \frac{(ac/q^2; q^2)_n (qc/a; q^2)_n}{(c; q)_n (q^{n-2}c; q)_n} \\ &+ \frac{q^{1+n}-a}{(q-a)(1+q^n)} \frac{(ac/q; q^2)_n (c/a; q^2)_n}{(c; q)_n (q^{n-2}c; q)_n}. \end{aligned}$$

Taking  $u = \ell, v = -m$  in (21) and calculating the  ${}_4\phi_3$ -series on the right hand side by (17), we achieve the following theorem.

**Theorem 17.** For two complex numbers  $\{a, c\}$  and two nonnegative integers  $\{\ell, m\}$ , there holds

$$\begin{aligned} {}_4\phi_3 & \left[ \begin{matrix} a, q^{1+\ell}/a, q^{-n}, -q^{-m-n} \\ c, q^{1+\ell-m-2n}/c, -q \end{matrix} \middle| q; q \right] = \left[ \begin{matrix} q^2/ac, -q/a \\ q^2/a^2, -q/c \end{matrix} \right]_\ell \left[ \begin{matrix} q^{1+m+n}, -q^{-\ell}c \\ q^{n+m-\ell}c, -q \end{matrix} \right]_n \\ & \times \sum_{i=0}^{\ell} \sum_{j=0}^m q^{(\ell+1)i-nj+{j+1 \choose 2}} \frac{(-1)^{i+j}}{c^i} \frac{1-q^{1+2i}/a^2}{1-q/a^2} \left[ \begin{matrix} q^{-\ell}, c/a, q/a^2 \\ q, q^2/ac, q^{2+\ell}/a^2 \end{matrix} \right]_i \\ & \times \left[ \begin{matrix} q^{-m}, q^{-m-n}, -q^{-m-n}, q^{-i-1}ac, q^i c/a \\ q, c, -c, q^{-m-2n}, q^{j-2m-2n-1} \end{matrix} \right]_j \left[ \begin{matrix} q^{j-i}ac, q^{1+i+j}c/a \\ q, q^{2j}c^2 \end{matrix} \right]_{m+n-j}. \end{aligned}$$

**Example 30** ( $\ell = 0, m = 1$  in Theorem 17).

$$\begin{aligned} {}_4\phi_3 & \left[ \begin{matrix} a, q/a, q^{-n}, -q^{-1-n} \\ c, q^{-2n}/c, -q \end{matrix} \middle| q; q \right] \\ &= \frac{1}{(1-q^{1+n})(1+q^n)c} \frac{(ac; q^2)_{1+n} (qc/a; q^2)_{1+n}}{(c; q)_{1+2n}} \\ &- \frac{q^{1+n}}{(1-q^{1+n})(1+q^n)c} \frac{(ac/q; q^2)_{1+n} (c/a; q^2)_{1+n}}{(c; q)_{1+2n}}. \end{aligned}$$

**Example 31** ( $\ell = 1, m = 1$  in Theorem 17).

$$\begin{aligned} {}_4\phi_3 & \left[ \begin{matrix} a, q^2/a, q^{-n}, -q^{-1-n} \\ c, q^{1-2n}/c, -q \end{matrix} \middle| q; q \right] \\ &= \frac{q^2(1+q^n)a}{(q-a)(1+q^n)c(q+q^n)(1-q^{1+n})} \frac{(ac/q^2; q^2)_{1+n} (qc/a; q^2)_{1+n}}{(c; q)_{2n}} \\ &- \frac{q(a+q^{2+n})}{(q-a)(1+q^n)c(q+q^n)(1-q^{1+n})} \frac{(ac/q; q^2)_{1+n} (c/a; q^2)_{1+n}}{(c; q)_{2n}}. \end{aligned}$$

Setting  $u = -\ell, v = m$  in (21) and evaluating the  ${}_4\phi_3$ -series on the right hand side by Theorem 5, we attain the following theorem.

**Theorem 18.** For two complex numbers  $\{a, c\}$  and two nonnegative integers  $\{\ell, m\}$  with  $m \leq n$ , there holds

$$\begin{aligned} {}_4\phi_3 \left[ \begin{matrix} a, q^{1-\ell}/a, q^{-n}, -q^{m-n} \\ c, q^{1-\ell+m-2n}/c, -q \end{matrix} \middle| q; q \right] &= \left[ \begin{matrix} a, q^{\ell-1}ac \\ c, q^{\ell-1}a^2 \end{matrix} \right]_{\ell} \left[ \begin{matrix} q^{1-m+n}, -q^{\ell}c \\ q^{n-m+\ell}c, -q \end{matrix} \right]_n \\ &\times \sum_{i=0}^{\ell} \sum_{j=0}^m q^{i+(m-n)j+{j+1 \choose 2}} \frac{1}{c^i} \frac{1-q^{1-2\ell+2i}/a^2}{1-q^{1-2\ell}/a^2} \left[ \begin{matrix} q^{-\ell}, c/a, q^{1-2\ell}/a^2 \\ q, q^{2-2\ell}/ac, q^{2-\ell}/a^2 \end{matrix} \right]_i \\ &\times \left[ \begin{matrix} q^{-m}, q^{-n}, -q^{-n}, q^{2\ell-i-1}ac, q^i c/a \\ q, q^{\ell}c, -q^{\ell}c, q^{m-2n}, q^{j-2n-1} \end{matrix} \right]_j \left[ \begin{matrix} q^{2\ell-i+j}ac, q^{1+i+j}c/a \\ q, q^{2\ell+2j}c^2 \end{matrix} \right]_{n-j}. \end{aligned}$$

**Example 32** ( $\ell = 1, m = 0$  in Theorem 18).

$$\begin{aligned} {}_4\phi_3 \left[ \begin{matrix} a, 1/a, q^{-n}, -q^{-n} \\ c, q^{-2n}/c, -q \end{matrix} \middle| q; q \right] &= \frac{1-ac}{1+a} \frac{(q^2 ac; q^2)_n (qc/a; q^2)_n}{(c; q)_{1+2n}} \\ &+ \frac{a-c}{1+a} \frac{(qac; q^2)_n (q^2 c/a; q^2)_n}{(c; q)_{1+2n}}. \end{aligned}$$

**Example 33** ( $\ell = 1, m = 1$  in Theorem 18).

$$\begin{aligned} {}_4\phi_3 \left[ \begin{matrix} a, 1/a, q^{-n}, -q^{1-n} \\ c, q^{1-2n}/c, -q \end{matrix} \middle| q; q \right] &= \frac{1+aq^n}{(1+a)(1+q^n)} \frac{(ac; q^2)_n (qc/a; q^2)_n}{(c; q)_{2n}} \\ &+ \frac{a+q^n}{(1+a)(1+q^n)} \frac{(qac; q^2)_n (c/a; q^2)_n}{(c; q)_{2n}}. \end{aligned}$$

Taking  $u = -\ell$ ,  $v = -m$  in (21) and calculating the  ${}_4\phi_3$ -series on the right hand side by Theorem 6, we obtain the following theorem.

**Theorem 19.** For two complex numbers  $\{a, c\}$  and two nonnegative integers  $\{\ell, m\}$ , there holds

$$\begin{aligned} {}_4\phi_3 \left[ \begin{matrix} a, q^{1-\ell}/a, q^{-n}, -q^{-m-n} \\ c, q^{1-\ell-m-2n}/c, -q \end{matrix} \middle| q; q \right] &= \left[ \begin{matrix} a, q^{\ell-1}ac \\ c, q^{\ell-1}a^2 \end{matrix} \right]_{\ell} \left[ \begin{matrix} q^{1+m+n}, -q^{\ell}c \\ q^{\ell+m+n}c, -q \end{matrix} \right]_n \\ &\times \sum_{i=0}^{\ell} \sum_{j=0}^m q^{i-nj+{j+1 \choose 2}} \frac{(-1)^j}{c^i} \frac{1-q^{1-2\ell+2i}/a^2}{1-q^{1-2\ell}/a^2} \left[ \begin{matrix} q^{-\ell}, c/a, q^{1-2\ell}/a^2 \\ q, q^{2-2\ell}/ac, q^{2-\ell}/a^2 \end{matrix} \right]_i \\ &\times \left[ \begin{matrix} q^{-m}, q^{-m-n}, -q^{-m-n}, q^{2\ell-i-1}ac, q^i c/a \\ q, q^{\ell}c, -q^{\ell}c, q^{-m-2n}, q^{j-2m-2n-1} \end{matrix} \right]_j \left[ \begin{matrix} q^{2\ell-i+j}ac, q^{1+i+j}c/a \\ q, q^{2\ell+2j}c^2 \end{matrix} \right]_{m+n-j}. \end{aligned}$$

**Example 34** ( $\ell = 1, m = 1$  in Theorem 19).

$$\begin{aligned} {}_4\phi_3 \left[ \begin{matrix} a, 1/a, q^{-n}, -q^{-1-n} \\ c, q^{-1-2n}/c, -q \end{matrix} \middle| q; q \right] &= \frac{1-aq^{1+n}}{(1+a)(1-q^{1+n})} \frac{(ac; q^2)_{1+n} (qc/a; q^2)_{1+n}}{(c; q)_{2+2n}} \\ &+ \frac{a-q^{1+n}}{(1+a)(1-q^{1+n})} \frac{(qac; q^2)_{1+n} (c/a; q^2)_{1+n}}{(c; q)_{2+2n}}. \end{aligned}$$

With the change of the parameters  $\ell$  and  $m$ , Theorems 2 and 5-19 can produce more concrete formulas. Due to the limit of space, the corresponding results will not be laid out in the paper.

### Acknowledgments

The work is supported by the Natural Science Foundations of China (Nos. 11301120, 11201241 and 11201291).

## REFERENCES

- [1] G.E. Andrews, On  $q$ -analogues of the Watson and Whipple summations, SIAM J. Math. Anal. 7 (1976), 332-336.
- [2] G.E. Andrews, R. Askey, R. Roy, Special Functions, Cambridge University Press, Cambridge, 2000.
- [3] W. Chu, Abel's method on summation by parts and balanced  $q$ -series identities, Appl. Anal. Discrete Math. 4 (2010), 54-65.
- [4] W. Chu, Analytical formulae for extended  ${}_3F_2$ -series of Watson-Whipple-Dixon with two extra integer parameters, Math. Comp. 81 (2012), 467-479.
- [5] V.J.W. Guo, Elementary proofs of some  $q$ -identities of Jackson and Andrews-Jain, Discrete Math. 295 (2005), 63-74.
- [6] J. L. Lavoie, Some summation formulas for the series  ${}_3F_2(1)$ , Math. Comp. 49 (1987), 269-274.
- [7] J. L. Lavoie, F. Grondin, A.K. Rathie, Generalizatins of Watson's theorem on the sum of a  ${}_3F_2$ , Indian J. Math. 34 (1992), 23-32.
- [8] G. Gasper, M. Rahman, Basic Hypergeometric Series (2nd edition), Cambridge Univercity Press, Cambridge, 2004.
- [9] V.K. Jain, Some transformations of basic hypergeometric functions. II, SIAM J. MATH. Anal. 12(1981), 957-961.