

LOCAL UNIVERSALITY OF ZEROES OF RANDOM POLYNOMIALS

TERENCE TAO AND VAN VU

ABSTRACT. In this paper, we establish some local universality results concerning the correlation functions of the zeroes of random polynomials with independent coefficients. More precisely, consider two random polynomials $f = \sum_{i=1}^n c_i \xi_i z^i$ and $\tilde{f} = \sum_{i=1}^n c_i \tilde{\xi}_i z^i$, where the ξ_i and $\tilde{\xi}_i$ are iid random variables that match moments to second order, the coefficients c_i are deterministic, and the degree parameter n is large. Our results show, under some light conditions on the coefficients c_i and the tails of $\xi_i, \tilde{\xi}_i$, that the correlation functions of the zeroes of f and \tilde{f} are approximately the same. As an application, we give some answers to the classical question

“How many zeroes of a random polynomials are real ?”

for several classes of random polynomial models.

Our analysis relies on a general *replacement principle*, motivated by some recent work in random matrix theory. This principle enables one to compare the correlation functions of two random functions f and \tilde{f} if their log magnitudes $\log |f|, \log |\tilde{f}|$ are close in distribution, and if some non-concentration bounds are obeyed.

1991 *Mathematics Subject Classification.* 15A52.

T. Tao is partially supported by a Simons Investigator award from the Simons Foundation and by NSF grant DMS-0649473.

V. Vu is supported by research grants from NSF and Airforce.

CONTENTS

| | |
|-------------------------------------------------------------------------|----|
| 1. Introduction | 3 |
| 2. Replacement principle, complex case | 9 |
| 3. Replacement principle: real case | 11 |
| 4. Guaranteeing the assumptions in the replacement principle | 13 |
| 5. Universality of the correlation functions of the classical ensembles | 16 |
| 6. Proof of the replacement principle, complex case | 23 |
| 7. Proof of the replacement principle, real case | 31 |
| 8. Non-clustering via sharp concentration | 36 |
| 9. Assumption verification: Proof of Lemma 13 | 38 |
| 10. Assumption verification: Proof of Theorem 16 | 40 |
| 11. Assumption verification: repulsion bounds | 44 |
| 12. Universality for flat polynomials | 47 |
| 13. Universality for elliptic polynomials | 54 |
| 14. Universality for Kac polynomials | 59 |
| References | 66 |

1. INTRODUCTION

1.1. Models of random polynomials. In this paper we study the distribution of the zeroes of a random polynomial $f = f_n$ when the degree parameter n is large (or goes asymptotically to infinity). For sake of exposition, we will focus on a simple model of random polynomials in which the coefficients are independent and derived from a common atom distribution, although several of our results extend to more general models.

Definition 1 (Random polynomials). Let n be a positive integer, let c_0, \dots, c_n be deterministic complex numbers, and let ξ be a complex random variable (which we call the *atom distribution*) of mean zero and finite non-zero variance. Given the coefficients c_0, \dots, c_n and atom distribution ξ , we associate the random polynomial $f = f_n = f_{n,\xi} : \mathbb{C} \rightarrow \mathbb{C}$ defined by the formula

$$f(z) := \sum_{i=0}^n c_i \xi_i z^i,$$

where ξ_0, \dots, ξ_n are jointly independent copies of ξ .

In practice, we will usually normalize the atom distribution ξ to have unit variance; note that this normalization clearly does not affect the zeroes of f . In some literature, one replaces either the bottom coefficient ξ_0 or the top coefficient ξ_n with the constant 1. This generally has a negligible impact on the distribution of the zeroes in the large n limit; however, we shall avoid such normalizations here (although this has the consequence that the polynomial f may occasionally have degree less than n , or even vanish entirely). Our focus in this paper will primarily be on the *universality phenomenon* in the context of zeroes of such random polynomials, which roughly speaking asserts that the (appropriately normalized) asymptotic behavior of these zeroes as $n \rightarrow \infty$ should become independent of the choice of atom distribution.

We isolate three specific choices of coefficients c_i that have been studied for a long time:

- (i) *Flat polynomials* or *Weyl polynomials* are polynomials associated to the coefficients $c_i := \sqrt{\frac{1}{i!}}$.
- (ii) *Elliptic polynomials* or *binomial polynomials* are polynomials associated to the coefficients $c_i := \sqrt{\binom{n}{i}}$.
- (iii) *Kac polynomials* are polynomials associated to the coefficients $c_i := 1$.

One can view Kac polynomials as the special case $L = 1$ of *hyperbolic polynomials* in which $c_i := \sqrt{\frac{L(L+1)\cdots(L+i-1)}{i!}}$ for some parameter $L > 0$, but for simplicity we will focus on the classical Kac polynomial case as a proxy for the more general hyperbolic case.

These polynomials have been intensively studied, particularly in the case when the atom distribution ξ is either the real gaussian $N(0, 1)_{\mathbb{R}}$ or the complex gaussian $N(0, 1)_{\mathbb{C}}$; see [25, 19]. As we shall recall later, the situation with the Kac polynomials is somewhat special when compared to the other models described above (its zeroes tend to cluster around the unit circle, instead of being distributed throughout a two-dimensional region in the plane).

If $f = f_n$ is a random polynomial of the form described in Definition 1, then f has degree at most n . If f is not identically zero, then from the fundamental theorem of algebra it has $\deg(f)$ zeroes in the complex plane \mathbb{C} (counting multiplicity). We adopt the convention that f also has $n - \deg(f)$ zeroes at infinity, and when f is identically zero we adopt the convention that f has n zeroes at infinity and no zeroes in \mathbb{C} . With these (admittedly artificial) conventions, f thus always has n zeroes ζ_1, \dots, ζ_n (ordered in some arbitrarily chosen fashion, e.g. lexicographically) in the Riemann sphere $\mathbb{C} \cup \{\infty\}$ (counting multiplicity), so that $\{\zeta_1, \dots, \zeta_n\}$ may be viewed as a point process in $\mathbb{C} \cup \{\infty\}$. We will sometimes refer to the set $\{\zeta_1, \dots, \zeta_n\}$ as the *spectrum* of f .

1.2. Number of real zeroes. With $f = f_n$ as above, and any subset Ω of \mathbb{C} , write

$$N_{\Omega} := |\{1 \leq i \leq n : \zeta_i \in \Omega\}|$$

for the number of zeroes of f in Ω (counting multiplicity); in particular, $N_{\mathbb{R}}$ is the number of real zeroes. This is a random variable taking values in $\{0, \dots, n\}$. The issue of understanding the typical size of $N_{\mathbb{R}}$ was already raised by Waring as far back as 1782 ([60, page 618], [29]), and has generated a huge amount of literature, of which we now pause to give a (incomplete and brief) survey. The statistic $N_{\mathbb{R}}$ is of interest primarily in the case when the atom distribution ξ and the coefficients c_i are both real-valued, since in the genuinely complex case one usually expects that none of the zeroes of the associated polynomial f will be real.

Most earlier works focused on the case of Kac polynomials, which are easier to analyze but have an atypical behavior compared to other random polynomial models. One of the first results in this context is by Bloch and Pólya [4], who studied the case of Kac polynomials with ξ uniformly distributed in $\{-1, 0, 1\}$, and established the somewhat weak upper bound

$$\mathbf{E}N_{\mathbb{R}} \ll n^{1/2}$$

where we use the usual asymptotic notation $X = O(Y)$ or $X \ll Y$ to denote the bound $|X| \leq CY$ where C is independent of Y . This bound is not sharp, and Kac polynomials actually have a remarkably small number of real zeroes. Indeed, in a series of papers [33, 34, 35, 36], Littlewood and Offord proved that for Kac polynomials with many basic atom distributions (such as gaussian, Bernoulli or uniform on $[-1, 1]$), one has the bounds

$$\frac{\log n}{\log \log \log n} \ll N_{\mathbb{R}} \ll \log^2 n$$

with probability $1 - o(1)$, where we use $o(1)$ to denote a quantity that goes to 0 as $n \rightarrow \infty$.

Later, Kac [26] found an exact formula for $\mathbf{E}N_{\mathbb{R}}$ in the case that ξ_i are real gaussians, and showed that

$$\mathbf{E}N_{\mathbb{R}} = \left(\frac{2}{\pi} + o(1) \right) \log n$$

in this case (see [61], [12] for more precise asymptotics).

This asymptotic has been extended to Kac polynomials with more general atom distributions. In a subsequent paper [27], Kac extended the result to the case when ξ has the uniform distribution on $[-1, 1]$. Erdős and Offord [14] extended the result to the Bernoulli distribution case (i.e. when ξ is uniform on $\{-1, +1\}$). Stevens [55] extended the asymptotics for a wide class of distributions, and finally Ibragimov and Maslova [20, 21] extended the result to all mean-zero distributions in the domain of attraction of the normal law, with the extra assumption that $\mathbf{P}(\xi = 0) = 0$. Maslova [39, 40] also proved that if $\mathbf{P}(\xi = 0) = 0$ and $\mathbf{E}|\xi|^{2+\varepsilon} < \infty$ for some constant $\varepsilon > 0$ then the variance of $N_{\mathbb{R}}$ is $(\frac{4}{\pi}(1 - \frac{2}{\pi}) + o(1)) \log n$, and furthermore established a central limit theorem for $N_{\mathbb{R}}$. In [10], Dembo et. al. computed the probability that $N_{\mathbb{R}} = k$ for any fixed k . There are also some non-trivial deterministic bounds for the maximum value of $N_{\mathbb{R}}$ (when the coefficients are, say, drawn from $\{-1, 0, 1\}$) which we will not describe in detail here, but see e.g. [13] for some recent results in this direction.

For non-Kac models such as the flat or elliptic polynomial ensembles, the behavior of $N_{\mathbb{R}}$ changes considerably. These types of random polynomial models were already studied to some extent in the classical papers of Littlewood and Offord, but most of the work on these models appeared later, partially motivated by connections to physics [5] and random analytic functions [19] or problems in numerical analysis and computation theory [29, 52]. In particular, many researchers consider the elliptic (or binomial) polynomial the most “natural” random polynomial [12, Section 1], [29, 52]; one reason for this is that in the case when the atom distribution ξ is the complex Gaussian $N(0, 1)_{\mathbb{C}}$, the distribution of the zeroes of the associated random polynomial is invariant with respect to rotations of the Riemann sphere $\mathbb{C} \cup \{\infty\}$ (see e.g. [19, Proposition 2.3.4]).

It is known that when the atom distribution ξ is the real Gaussian $N(0, 1)_{\mathbb{R}}$, one has

$$\mathbf{E}N_{\mathbb{R}} = \left(\frac{2}{\pi} + o(1) \right) \sqrt{n}$$

for flat (Weyl) polynomials and

$$\mathbf{E}N_{\mathbb{R}} = \sqrt{n}$$

for elliptic (binomial) polynomials; see [12] for a nice geometric proof of these facts. Thus one has substantially more zeroes for such polynomials than in the Kac case when n is large. However, unlike the situation with Kac polynomials, the extension of these results to more general (non-gaussian) distributions was not fully understood. The reader is referred to [22, 23, 37, 38, 62, 7, 56] and the books [3, 15] for several results and further discussion.

1.3. Distribution of zeroes: Correlation functions. We now turn to a popular way to study the distribution of zeroes of random polynomials, namely by investigating their *correlation functions*. To define these functions, let us first consider the complex case in which the coefficients c_i and the atom distribution ξ are not required to be real valued. In this case the point process $\{\zeta_1, \dots, \zeta_n\}$ of zeroes of a random polynomial $f = f_n$ can be described using the (complex) k -point *correlation functions* $\rho^{(k)} = \rho_f^{(k)} : \mathbb{C}^k \rightarrow \mathbb{R}^+$, defined for any fixed natural number k by requiring that

$$(1) \quad \mathbf{E} \sum_{i_1, \dots, i_k \text{ distinct}} \varphi(\zeta_{i_1}, \dots, \zeta_{i_k}) = \int_{\mathbb{C}^k} \varphi(z_1, \dots, z_k) \rho^{(k)}(z_1, \dots, z_k) dz_1 \dots dz_k$$

for any continuous, compactly supported, test function $\varphi : \mathbb{C}^k \rightarrow \mathbb{C}$, with the convention that $\varphi(\infty) = 0$; see e.g. [19, 1]. This definition of $\rho^{(k)}$ is clearly independent of the choice of ordering ζ_1, \dots, ζ_n of the zeroes. Note that if the random polynomial f has a discrete law rather than a continuous one, then $\rho_f^{(k)}$ needs to be interpreted as a measure¹ rather than as a function.

Remark 2. When ξ has a continuous complex distribution, the c_i are non-zero, then the zeroes are almost surely simple. In this case if z_1, \dots, z_k are distinct, and one can interpret $\rho^{(k)}(z_1, \dots, z_k)$ as the unique quantity such that the probability that there is a zero in each of the disks $B(z_i, \varepsilon)$ for $i = 1, \dots, k$ is $(\rho^{(k)}(z_1, \dots, z_k) + o_{\varepsilon \rightarrow 0}(1))(\pi\varepsilon^2)^k$ in the limit $\varepsilon \rightarrow 0$.

When the random polynomials f have real coefficients, the zeroes ζ_1, \dots, ζ_n are symmetric around the real axis, and one expects several of the zeroes to lie on this axis. Because of this, it is not as natural to work with the complex k -point correlation functions $\rho_f^{(k)}$, as they are likely to become singular on the real axis. Instead, we divide the complex plane \mathbb{C} into three pieces $\mathbb{C} = \mathbb{R} \cup \mathbb{C}_+ \cup \mathbb{C}_-$, with $\mathbb{C}_+ := \{z \in \mathbb{C} : \text{Im}(z) > 0\}$ being the upper half-plane and $\mathbb{C}_- := \{z \in \mathbb{C} : \text{Im}(z) < 0\}$ being the lower half-plane. By the aforementioned symmetry, we may restrict attention to the zeroes in \mathbb{R} and \mathbb{C}_+ only. For any natural numbers $k, l \geq 0$, we then define the *mixed (k, l) -correlation function* $\rho_f^{(k, l)} = \rho_f^{(k, l)} : \mathbb{R}^k \times (\mathbb{C}_+ \cup \mathbb{C}_-)^l \rightarrow \mathbb{R}^+$ of a random polynomial f to be the function defined by the formula

$$(2) \quad \mathbf{E} \sum_{i_1, \dots, i_k \text{ distinct}} \sum_{j_1, \dots, j_l \text{ distinct}} \varphi(\zeta_{i_1, \mathbb{R}}, \dots, \zeta_{i_k, \mathbb{R}}, \zeta_{j_1, \mathbb{C}_+}, \dots, \zeta_{j_l, \mathbb{C}_+}) \\ = \int_{\mathbb{R}^k} \int_{\mathbb{C}_+^l} \varphi(x_1, \dots, x_k, z_1, \dots, z_l) \rho_f^{(k, l)}(x_1, \dots, x_k, z_1, \dots, z_l) dz_1 \dots dz_l dx_1 \dots dx_k$$

for any continuous, compactly supported test function $\varphi : \mathbb{R}^k \times \mathbb{C}^l \rightarrow \mathbb{C}$ (note that we do not require φ to vanish at the boundary of \mathbb{C}_+^l), $\zeta_{i, \mathbb{R}}$ runs over an arbitrary

¹We point out one subtlety in the discrete case: the summation (1) requires the *indices* i_1, \dots, i_k to be distinct, but allows the *zeroes* $\zeta_{i_1}, \dots, \zeta_{i_k}$ to be repeated; thus for instance if f is the deterministic polynomial z^n then $\rho^{(k)}$ is $\frac{n!}{(n-k)!}$ times the Dirac mass at the origin in \mathbb{C}^k . This convention allows for identities such as $(n-k)\rho^{(k)}(z_1, \dots, z_k) = \int_{\mathbb{C} \cup \infty} \rho^{(k+1)}(z_1, \dots, z_{k+1}) dz_{k+1}$ to be extended to the discrete setting (after being interpreted in an appropriate distributional sense); it also ensures that the distribution functions $\rho^{(k)}$ vary continuously (in the vague topology) with respect to perturbations of law of the random polynomial f (again measured in the vague topology). Of course, when f has a continuous distribution, the zeroes are almost surely simple, and this subtlety becomes irrelevant.

enumeration of the real zeroes of f_n , and ζ_{j, \mathbb{C}_+} runs over an arbitrary enumeration of the zeroes of f_n in \mathbb{C}_+ . This defines $\rho^{(k,l)}$ (in the sense of distributions, at least) for $x_1, \dots, x_k \in \mathbb{R}$ and $z_1, \dots, z_l \in \mathbb{C}_+$; we then extend $\rho^{(k,l)}(x_1, \dots, x_k, z_1, \dots, z_l)$ to all other values of $x_1, \dots, x_k \in \mathbb{R}$ and $z_1, \dots, z_l \in \mathbb{C}_+ \cup \mathbb{C}_-$ by requiring that $\rho^{(k,l)}$ is symmetric with respect to conjugation of any or all of the z_1, \dots, z_l parameters. Again, we permit $\rho^{(k,l)}$ to be a measure² instead of a function when the random polynomial f_n has a discrete distribution.

In the case $l = 0$, the correlation functions $\rho^{(k,0)}$ for $k \geq 1$ provide (in principle, at least) all the essential information about the distribution of the real zeroes, which as mentioned previously, was the original motivation of the very first papers studying random polynomials. For instance, one easily verifies the identity

$$(3) \quad \mathbf{E}N_{\mathbb{R}} = \int_{\mathbb{R}} \rho^{(1,0)}(x) dx$$

and similarly

$$(4) \quad \mathbf{Var}N_{\mathbb{R}} = \int_{\mathbb{R}} \int_{\mathbb{R}} \rho^{(2,0)}(x, y) - \rho^{(1,0)}(x)\rho^{(1,0)}(y) dx dy + \int_{\mathbb{R}} \rho^{(1,0)}(x) dx$$

Remark 3. When ξ has a continuous real distribution, the c_i are non-zero real, then the zeroes are almost surely simple. If the $x_1, \dots, x_k \in \mathbb{R}$ are distinct, and the $z_1, \dots, z_l \in \mathbb{C}_+$ are distinct, then one can interpret $\rho^{(k,l)}(x_1, \dots, x_k, z_1, \dots, z_l)$ as the unique quantity such that the probability that there is a zero in each of the intervals $[x_i - \varepsilon, x_i + \varepsilon]$ and disks $B(z_j, \varepsilon)$ for $i = 1, \dots, k$ and $j = 1, \dots, l$ is $(\rho^{(k,l)}(x_1, \dots, x_k, z_1, \dots, z_l) + o_{\varepsilon \rightarrow 0}(1))(2\varepsilon)^k (\pi\varepsilon^2)^l$ in the limit $\varepsilon \rightarrow 0$.

Remark 4. In principle, one could express the complex correlation functions in a distributional sense in terms of the real correlation functions, for instance we have

$$\rho^{(1)}(z) = \rho^{(0,1)}(z) + \rho^{(1,0)}(\operatorname{Re}z)\delta(\operatorname{Im}z)$$

in the sense of distributions, where δ is the Dirac distribution at 0, with similar (but significantly more complicated) identities for $\rho^{(k)}$ when $k > 1$, reflecting the many combinatorial possibilities for k complex zeroes to lie on the real line, or to be complex conjugates of each other. We will however not use such identities in this paper.

1.4. Universality. In the case when the atom distribution ξ is a real or complex gaussian, the correlation functions $\rho^{(k,l)}$ (in the real case) or $\rho^{(k)}$ (in the complex case) can be computed explicitly using tools such as the Kac-Rice formula; see [19] or Lemma 34 below. When the atom distribution is not gaussian, the Kac-Rice formula is still available, but is considerably less tractable. Nevertheless, it has been widely believed that the asymptotic behavior of the correlation functions in the non-gaussian case should match that of the gaussian case once one has performed

²As in the complex case, we allow the real zeros $\zeta_{i_1, \mathbb{R}}, \dots, \zeta_{i_k, \mathbb{R}}$ or the complex zeroes $\zeta_{j_1, \mathbb{C}_+}, \dots, \zeta_{j_l, \mathbb{C}_+}$ to have multiplicity; it is only the indices $i_1, \dots, i_k, j_1, \dots, j_l$ that are required to be distinct. In particular, in the discrete case it is possible for $\rho^{(0,2)}(z_1, z_2)$ (say) to have non-zero mass on the diagonal $z_1 = z_2$ or the conjugate diagonal $z_1 = \overline{z_2}$, if f has a repeated complex eigenvalue with positive probability.

appropriate normalizations, at least if the atom distribution ξ is sufficiently short-tailed. This type of meta-conjecture is commonly referred to as the *universality phenomenon*.

At macroscopic (or *global*) scales (comparable to the diameter of the bulk of the set of zeroes), universality results for polynomials given by Definition 1 were established recently by Kabluchko and Zaporozhets [25]. For instance, they established the analogue of the circular law for Weyl polynomials given only a mild log-integrability condition for the atom distribution (see [25, Theorem 2.3]), as well as many other results of this nature; see [25] for full details.

In this paper we will be concerned primarily with universality of correlation functions at the microscopic (or *local*) scale, comparable to the mean spacing between zeroes; through formulae such as (3), this also can lead to some partial universality results for quantities such as $N_{\mathbb{R}}$. At such microscopic scales, the most general previous result we found concerning universality is due to Bleher and Di [7, Theorem 7.2], who considered binomial polynomials in which the atom distribution ξ was real-valued, with unit variance, and was sufficiently smooth and rapidly decaying (see [7, Theorem 7.2] for the precise technical conditions required on ξ). With these hypotheses, they showed that the pointwise limit of the normalized correlation function $n^{-k/2} \rho^{(k,0)}(a + \frac{x_1}{\sqrt{n}}, \dots, a + \frac{x_k}{\sqrt{n}})$ for any fixed k, a, x_1, \dots, x_k (with $a \neq 0$) as $n \rightarrow \infty$ was independent of the choice of ξ (with an explicit formula for the limiting distribution). Again, see [7, Theorem 7.2] for a precise statement. One of the main tools used in that argument was the Kac-Rice formula.

In this paper, we first introduce a new method to prove universality, which makes no distinction between continuous and discrete random variables (and in particular, avoids the use of the Kac-Rice formula, except when verifying a certain technical level repulsion estimate in the real gaussian case). As a matter of fact, we will only require some bounded moment assumption on the atom distribution. This approach relies on a general *replacement principle*, which we will present in the next section. This reduces the task of establishing universality for zeroes of a random polynomial f to that of establishing universality for the log-magnitude $\log |f(z)|$ of that polynomial evaluated at various points z , together with that of verifying some technical eigenvalue repulsion bounds. This principle was implicitly introduced in our previous paper [58] in the context where f was the characteristic polynomial of a random matrix (and is thus can be viewed as a microscopic analogue to the macroscopic replacement principle in [57, Theorem 2.1] to establish the circular law for various ensembles of random matrices), but applies for more general random matrix models, and is in fact particularly easy to use for the models in Definition 1 since $f(z)$ is just the sum of independent random variables for each given z in this case. As applications of this principle we will establish universality results for all the classical ensembles listed above. We would like to emphasize here that while in this paper we focus on random polynomials with independent coefficients, our replacement principle *does not* require this assumption. For example, it can be applied to characteristic polynomials of random matrices [58].

1.5. Notation. We use 1_E to denote the indicator of E , thus 1_E equals 1 when E is true and 0 when E is false. We also write $1_\Omega(x)$ for $1_{x \in \Omega}$.

We use $\sqrt{-1}$ to denote the unit imaginary, in order to free up the symbol i as an index of summation. As we will be using two-dimensional integration on the complex plane $\mathbb{C} := \{z = x + \sqrt{-1}y : x, y \in \mathbb{R}\}$ far more often than we will be using contour integration, we use $dz := dx dy$ to denote Lebesgue measure on the complex numbers, rather than the complex line element $dx + \sqrt{-1}dy$. For $z \in \mathbb{C}$ and $r > 0$, we use $B(z, r) := \{w \in \mathbb{C} : |z - w| < r\}$ to denote the open disk of radius r centered at z .

If $G : \mathbb{R}^k \rightarrow \mathbb{C}$ is a function and $a \geq 0$, we use $\nabla^a G$ to denote the tensor $(\frac{\partial^a}{\partial x_{i_1} \dots \partial x_{i_a}} G)_{1 \leq i_1, \dots, i_a \leq k}$; in particular,

$$|\nabla^a G| := \left(\sum_{1 \leq i_1, \dots, i_a \leq k} \left| \frac{\partial^a}{\partial x_{i_1} \dots \partial x_{i_a}} G \right|^2 \right)^{1/2}.$$

Following [59], we say that two complex random variables ξ, ξ' *match moments to order m* if one has

$$\mathbf{E} \operatorname{Re}(\xi)^a \operatorname{Im}(\xi)^b = \mathbf{E} \operatorname{Re}(\xi')^a \operatorname{Im}(\xi')^b$$

for all natural numbers $a, b \geq 0$ with $a + b \leq m$.

2. REPLACEMENT PRINCIPLE, COMPLEX CASE

Our replacement principle asserts, roughly speaking, that the k -correlation functions of the zeroes of two random polynomials f and \tilde{f} are asymptotically the same provided that

- (i) (Comparability of log-magnitudes) The joint distribution of $\log |f|$ at a few values is close to the joint distribution of $\log |\tilde{f}|$ at those same values; and
- (ii) (Non-clustering property) f and \tilde{f} do not have too many zeroes concentrating in a small region.

We will also need a mild non-degeneracy condition that prevents f or \tilde{f} from vanishing identically too often, but this hypothesis is easily verified in practice.

Moreover, we can show that the non-clustering property holds if the variables $\log |f(z)|$ and $\log |\tilde{f}(z)|$ are strongly concentrated around a suitable deterministic function $G(z)$ (see Proposition 12 below). So, in order to compare the distribution of the zeroes of f and \tilde{f} , all we need is to study the distribution of the log-magnitude functions $\log |f(z)|$ and $\log |\tilde{f}(z)|$ for various choices of parameter z .

When the random polynomials f and \tilde{f} have real coefficients, we can prove a similar replacement principle for the mixed (k, l) -point correlation functions involving k real numbers and l strictly complex numbers, provided we assume an additional

level repulsion estimate on at least one of f, \tilde{f} . In practice, this estimate will be easy to verify for many random polynomials with real gaussian coefficients.

We now give the formal statement of the replacement principle in the complex case.

Theorem 5 (Replacement principle, complex case). *Let $C, r_0 \geq 1 \geq c_0 > 0$ be real constants and $k, a_0 \geq 1$ be integer constants, and set*

$$(5) \quad A := \frac{100ka_0}{c_0}.$$

Let $n \geq 1$ be a natural number, and let $f = f_n, \tilde{f} = \tilde{f}_n$ be random polynomials of degree at most n (not necessarily of the form in Definition 1) and z_1, \dots, z_k be complex numbers that are allowed to depend on n . Assume the following axioms.

- (i) (Non-degeneracy) *With probability at least $1 - Cn^{-A}$, f is not identically zero, and similarly for \tilde{f} .*
- (ii) (Non-clustering property) *For $r \geq 1$, one has $N_{B(z_i, r)}(f) \leq Cn^{1/A}r^2$ with probability at least $1 - Cn^{-A}$. Similarly for \tilde{f} .*
- (iii) (Comparability of log-magnitudes) *Given any $1 \leq k' \leq n^{c_0}$, any complex numbers $z'_1, \dots, z'_{k'} \in \bigcup_{i=1}^k B(z_i, 20r_0)$, and any smooth function $F : \mathbb{C}^{k'} \rightarrow \mathbb{C}$ obeying the derivative bounds*

$$|\nabla^a F(w)| \leq n^{c_0}$$

for all $0 \leq a \leq a_0$ and $w \in \mathbb{C}^{k'}$, we have

$$(6) \quad \left| \mathbf{E} \left(F(\log |f(z'_1)|, \dots, \log |f(z'_{k'})|) - F(\log |\tilde{f}(z'_1)|, \dots, \log |\tilde{f}(z'_{k'})|) \right) \right| \leq Cn^{-c_0}$$

with the convention that F vanishes when one or more of its arguments are undefined.

Let $G : \mathbb{C}^k \rightarrow \mathbb{C}$ be a smooth function supported on the polydisc $B(0, r_0)^k$ that obeys the bounds

$$(7) \quad |\nabla^a G(w)| \leq M$$

for all $0 \leq a \leq a_0 + 2k + 1$, all $w \in \mathbb{C}^k$, and some $M > 0$. Then

$$\left| \int_{\mathbb{C}^k} G(w_1, \dots, w_k) \rho_f^{(k)}(z_1 + w_1, \dots, z_k + w_k) dw_1 \dots dw_k - \int_{\mathbb{C}^k} G(w_1, \dots, w_k) \rho_{\tilde{f}}^{(k)}(z_1 + w_1, \dots, z_k + w_k) dw_1 \dots dw_k \right| \leq \tilde{C}Mn^{-c_0/4},$$

where \tilde{C} depends only on the quantities C, r_0, c_0, k, a_0 .

We will prove this theorem in Section 6.

Remark 6. In the applications in this paper, we will always take $a_0 = 3$, because our statistics will ultimately only depend on the first two moments of the atom distribution, and this can be exploited by Taylor expansions with a third order error. In applications to random matrices such as [58], it is more convenient to take $a_0 = 5$, because random matrix statistics may be sensitive to the first four

moments of the atom distribution (thanks to the *Four Moment Theorem*), which require Taylor expansions with a fifth order error to exploit.

Remark 7. One can view the above result as a local version of the replacement principle in [57, Theorem 2.1], which assumed a much weaker non-clustering bound (which, in the context of characteristic polynomials of random matrices, was formulated as a Frobenius norm bound on the relevant random matrices) and which assumed asymptotic comparability of $\frac{1}{n} \log |f(z)|$ and $\frac{1}{n} \log |\tilde{f}(z)|$ for each complex number z , rather than local comparability (the relationship between the two is roughly analogous to the relationship between the law of large numbers and the central limit theorem), but only gave conclusions about the global distribution of zeroes, rather than the local correlation functions. Versions of this latter principle were used in the recent work of Kabluchko and Zaporozhets [25] on global universality for random polynomials.

Remark 8. The theorem requires some smoothness bounds (7) on the test function G , but if one is willing to replace the quantitative bound $\tilde{C}Mn^{-c_0/4}$ in the conclusion of the theorem by weaker upper and lower bounds with error terms that go to zero as $n \rightarrow \infty$, one can extend the result to functions G that are merely assumed to be continuous rather than smooth, by using tools such as the Stone-Weierstrass theorem to approximate continuous G above and below by smooth G ; we omit the details.

Remark 9. Theorem 18 is adapted to the situation in which the mean spacing between zeroes is expected to be comparable to 1 (so that the correlation functions $\rho^{(k)}$ are also expected to have average magnitude comparable to 1). In practice, we may employ a rescaling in order to allow Theorem 18 to meaningfully apply to settings in which the mean spacing is at some other scale (e.g. $1/\sqrt{n}$ or $1/n$). One could also develop more general versions of Theorem 18 in which the mean spacing near each reference point z_j varies with j , but we will not detail such generalizations here in order to simplify the exposition.

3. REPLACEMENT PRINCIPLE: REAL CASE

Now we give the analogue of Theorem 5 in the case of polynomials with real coefficients, which is slightly more complicated and has slightly worse constants, but is otherwise very similar to the complex replacement principle.

Theorem 10 (Replacement principle, real case). *Let $C, r_0 \geq 1 \geq c_0 > 0$ be real constants, and $a_0 \geq 1$ and $k, l \geq 0$ be integer constants with $k + l > 0$, and set*

$$(8) \quad A := \frac{200(k+l)^2(a_0+2)}{c_0}.$$

Let $n \geq 1$ be a natural number, let $f = f_n, \tilde{f} = \tilde{f}_n$ be random polynomials of degree at most n with real coefficients (not necessarily of the form in Definition 1) and let $x_1, \dots, x_k \in \mathbb{R}$ and $z_1, \dots, z_l \in \mathbb{C}$ be numbers that are allowed to depend on n . Assume the following axioms.

- (i) (Non-degeneracy) *With probability at least $1 - Cn^{-A}$, f is not identically zero, and similarly for \tilde{f} .*

- (ii) (*Non-clustering property*) For $r \geq 1$, we have $N_{B(x_i, r)}(f), N_{B(z_j, r)}(f) \leq Cn^{1/A}r^2$ for $1 \leq i \leq k, 1 \leq j \leq l$ with probability at least $1 - Cn^{-A}$, and similarly for \tilde{f}_n .
- (iii) (*Comparability of log-magnitudes*) Given any $1 \leq k' \leq n^{c_0}$, any complex numbers

$$z'_1, \dots, z'_{k'} \in \bigcup_{i=1}^k B(x_i, 100r_0) \cup \bigcup_{j=1}^l B(z_j, 100r_0),$$

and any smooth function $F : \mathbb{C}^{k'} \rightarrow \mathbb{C}$ obeying the derivative bounds

$$|\nabla^a F(w)| \leq n^{c_0}$$

for all $0 \leq a \leq a_0$, we have

$$\left| \mathbf{E} \left(F(\log |f(z'_1)|, \dots, \log |f(z'_{k'})|) - F(\log |\tilde{f}(z'_1)|, \dots, \log |\tilde{f}(z'_{k'})|) \right) \right| \leq Cn^{-c_0}.$$

- (iv) (*Weak level repulsion*) For x, y real, z complex in the region

$$\bigcup_{i=1}^k B(x_i, 100r_0) \cup \bigcup_{j=1}^l B(z_j, 100r_0)$$

with $|x - y|, |\operatorname{Im}(z)| \leq 1/C$, we have the pointwise bounds

$$(9) \quad \rho_{\tilde{f}}^{(2,0)}(x, y) \leq C$$

and

$$(10) \quad \rho_{\tilde{f}}^{(0,1)}(z) \leq C.$$

Let $G : \mathbb{R}^k \times \mathbb{C}^l \rightarrow \mathbb{C}$ be a smooth function supported on $[-r_0, r_0]^k \times B(0, r_0)^l$ that obeys the bounds

$$(11) \quad |\nabla^a G(w)| \leq M$$

for all $0 \leq a \leq a_0 + 2(k+l) + 1$, all $w \in \mathbb{R}^k \times \mathbb{C}^l$, and some $M > 0$. Then

$$\begin{aligned} & \left| \int_{\mathbb{R}^k} \int_{\mathbb{C}^l} G(y_1, \dots, y_k, w_1, \dots, w_l) \right. \\ & \quad \rho_f^{(k,l)}(x_1 + y_1, \dots, x_k + y_k, z_1 + w_1, \dots, z_l + w_l) dw_1 \dots dw_l dy_1 \dots dy_l \\ & \quad - \int_{\mathbb{R}^k} \int_{\mathbb{C}^l} G(y_1, \dots, y_k, w_1, \dots, w_l) \\ & \quad \left. \rho_{\tilde{f}}^{(k,l)}(x_1 + y_1, \dots, x_k + y_k, z_1 + w_1, \dots, z_l + w_l) dw_1 \dots dw_l dy_1 \dots dy_l \right| \\ & \leq \tilde{C} M n^{-\frac{c_0}{200(a_0+2)(k+l)}} \end{aligned}$$

where \tilde{C} depends only on $C, r_0, c_0, k, l, r_0, a_0$.

We prove this theorem in Section 7.

Remark 11. Notice that we require the weak repulsion (9), (10) just for \tilde{f} and not for f . In practice, we can often choose \tilde{f} to have gaussian coefficients and verify this axiom by a direct (although not entirely trivial) computation (using tools such as the Kac-Rice formula), while f is permitted to have a discrete distribution (which

would be very unlikely to obey (9), (10) in a pointwise sense). In our applications one can usually establish a stronger level repulsion bound than (9), (10), namely a bound which decays linearly in $|x - y|$ or $|\operatorname{Im}z|$, but we will not need this stronger bound here. The reader may notice the difference in the exponent in the final bound, compared to the complex case ($\frac{c_0}{200(a_0+2)(k+l)}$ instead of $\frac{c_0}{4}$). This is due to the fact that we will need to apply the result in the complex case for a function G with derivatives that can be polynomially large. We make no attempt to optimize these constants whatsoever.

4. GUARANTEEING THE ASSUMPTIONS IN THE REPLACEMENT PRINCIPLE

We now present some tools to verify the various axioms in the replacement principle (Theorem 5 or Theorem 10). In order to use asymptotic notation such as $O()$ and $o()$, it will be convenient to phrase these tools in the asymptotic setting in which n is going to infinity (rather than being large and fixed), although one could easily rewrite the propositions below in the non-asymptotic language of a single fixed n if desired.

Let us say that an event depending on n occurs *with overwhelming probability* if it occurs with probability $1 - O(n^{-A})$ for any fixed A (independent of n), where the implied constant is allowed to depend on A .

We now give a general result (which implicitly appears in our previous paper [58]; see also [8] for a closely related argument) that guarantees the non-degeneracy and non-clustering axioms (i), (ii) in Theorem 5 or Theorem 10 if one can obtain concentration result for the log-magnitude $\log |f|$.

Proposition 12 (Criterion for non-clustering). *Let $n \geq 1$ be a natural number, and let $f = f_n$ be a random polynomial of degree at most n . Let z_0 be a complex number depending on n , and let $0 < c \leq r$ be quantities that are permitted to depend on n , with the polynomial size bounds $r \ll n^{O(1)}$ and $c \gg n^{-O(1)}$. Assume the following axiom:*

(i) (*Concentration of logarithm*) For any $z \in B(z_0, r+c) \setminus B(z_0, r-c)$, one has

$$\log |f(z)| = G(z) + O(n^{o(1)})$$

with overwhelming probability, where $G : \mathbb{C} \rightarrow \mathbb{R}$ is a (deterministic) smooth function (that can depend on n) obeying the polynomial size bound

$$(12) \quad \sup_{z \in B(z_0, r+c) \setminus B(z_0, r-c)} |G(z)| \ll n^{O(1)},$$

and we adopt the convention $\log |0| = -\infty$.

Then one has with overwhelming probability that f is non-vanishing and obeys the bound

$$(13) \quad N_{B(z_0, r)}(f) = \frac{1}{2\pi} \int_{B(z_0, r)} \Delta G(z) dz + O(n^{o(1)} c^{-1} r) + O \left(\int_{B(z_0, r+c) \setminus B(z_0, r-c)} |\Delta G(z)| dz \right).$$

Furthermore, the implied constants in the conclusions depend only on the implied constants in the hypotheses.

This proposition can be viewed as a variant of the classical Jensen formula linking the zeroes of a holomorphic function to a certain integral of the log-magnitude of this function. We will prove it in Section 8. To use this proposition, we now specialize to the case of polynomials $f = f_\xi$ of the form of Definition 1. We will normalize the atom distribution ξ to have unit variance. A short computation then reveals that for any complex number z , the random variable $f(z)$ has mean zero and variance

$$(14) \quad V(z) := \mathbf{E}(f(z)\bar{f}(z)) = \sum_{i=0}^n |c_i|^2 |z|^{2i}.$$

Note that this quantity is independent of the atom distribution ξ (once it has been normalized as above). It is then natural to expect the concentration result

$$\log |f(z)| = \frac{1}{2} \log V(z) + O(n^{o(1)})$$

with overwhelming probability, which would give the hypothesis of Proposition 12 with $G(z) := \frac{1}{2} \log V(z)$. The following proposition makes this prediction rigorous, provided that the coefficients c_i contain a sufficiently long and non-trivial lacunary subsequence:

Lemma 13. *Let $n \geq 1$, and let $f = f_n$ be a random polynomial of the type in Definition 1 whose atom distribution ξ has mean zero and variance one; suppose further that we have the moment condition $\mathbf{E}|\xi|^{2+\varepsilon} \leq M$ for some $\varepsilon > 0$ and $M < \infty$. Let z be a complex number (that can depend on n), and let $V(z)$ be defined by (14). Assume that there are indices $i_1, \dots, i_m \in \{0, \dots, n\}$ for some $m = \omega(\log n)$ (thus $m \geq C(n) \log n$ for some $C(n)$ that goes to infinity as $n \rightarrow \infty$) such that we have the lacunarity property*

$$|c_{i_j} z^{i_j}| \geq 2 |c_{i_{j+1}} z^{i_{j+1}}|$$

for all $1 \leq j < m$, and the lower bound

$$|c_{i_m} z^{i_m}| \geq V(z)^{1/2} \exp(-n^{o(1)}).$$

Then with overwhelming probability we have

$$\log |f(z)| = \frac{1}{2} \log V(z) + O(n^{o(1)}).$$

The implied constants in the conclusion depend on those in the hypotheses, and also on ε and M , but are otherwise uniform in ξ .

We establish this lemma in Section 9.

The following simple lemma is useful in proving the existence of the subsequence i_j in the above lemma.

Lemma 14. *Assume that $b_0 \geq b_1 \geq \dots \geq b_l > 0$ and $b_i/b_{i+1} \leq C$ for some $C \geq 2$ then the sequence b_i contains a subsequence b_{i_1}, \dots, b_{i_m} of length $m \gg \log_C b_0/b_l$ that obeys the lacunarity property $b_{i_j} \geq 2b_{i_{j+1}}$ for all $1 \leq j < m$.*

Proof. This is immediate from the greedy algorithm. \square

Remark 15. Proposition 12 combined with Lemma 13 suggests that the first intensity $\rho^{(1)}(z)$ of a random polynomial f of the form in Definition 1 should be approximately equal to $\frac{1}{4\pi}\Delta \log V(z)$ in some weak sense. In the case that the atom distribution ξ was complex gaussian, this approximation was in fact shown to be exact in [12] (see also [53]); this can also be derived from the Kac-Rice formula. The results in [25] can be viewed as a verification of this approximation $\rho^{(1)}(z) \approx \frac{1}{4\pi}\Delta \log V(z)$ at global scales.

4.1. Comparability of log-magnitudes. Next, we present a two moment theorem for the log-magnitude, which assures assumption (iii) of the replacement principle.

Theorem 16 (Two moment theorem for log-magnitude). *Let $\xi, \tilde{\xi}$ be two complex random variables of mean zero, variance one, which match moments to second order, and which obey the moment bound $\mathbf{E}|\xi|^{2+\varepsilon}, \mathbf{E}|\tilde{\xi}|^{2+\varepsilon} < M$ for some finite ε, M . Let $n \geq 1$, and suppose that $f_{n,\xi}, f_{n,\tilde{\xi}}$ are random polynomials of the form in Definition 1 with atom distributions $\xi, \tilde{\xi}$ respectively, and some choices of coefficients c_0, \dots, c_n . Let $k \geq 1$ be a natural number with $k \leq n^{\alpha_0}$ for some $\alpha_0 > 0$, and let z_1, \dots, z_k be complex numbers obeying the delocalization bounds*

$$(15) \quad |c_i z_j^i| \leq n^{-\alpha_1} V(z_j)^{1/2}.$$

for $0 \leq i \leq n$ and $1 \leq j \leq k$, where V is defined in (14). Let $G : \mathbb{C}^k \rightarrow \mathbb{C}$ be a smooth function (possibly depending on n) obeying the derivative bounds

$$|\nabla^a G(w)| \leq n^{\alpha_0}$$

for all $0 \leq a \leq 3$. Then if α_0 is sufficiently small depending only on α_1, ε , we have

$$(16) \quad \mathbf{E}G(\log |f_{n,\xi}(z_1)|, \dots, \log |f_{n,\xi}(z_k)|) - G(\log |f_{n,\tilde{\xi}}(z_1)|, \dots, \log |f_{n,\tilde{\xi}}(z_k)|) = O(n^{-\alpha_0})$$

where the implied constant depends only on $\alpha_0, \alpha_1, \varepsilon, M$.

We prove this theorem in Section 10. The $2+\varepsilon$ moment bound is needed to obtain a polynomial decay rate $O(n^{-\alpha_0})$ in (16), which in turn is needed in our version of the replacement principle. It may however be possible through a more careful analysis to obtain local universality results for polynomials that do not obey this bound, at the cost of replacing $O(n^{-\alpha_0})$ type error terms in the final universality bounds with qualitative decay terms $o(1)$. We will not pursue this issue here. Note that if $\xi, \tilde{\xi}$ are both real valued, then the hypothesis of matching moments to second order is automatic since the $\xi, \tilde{\xi}$ are normalized to have mean zero and variance one. The leading idea is to use Lindeberg replacement trick, originated in [32] (see also [45, 6] for more recent discussions). The arguments we will use follow the spirit of [59, 58], where characteristic polynomials of random matrices were considered.

4.2. A sufficient condition for the repulsion bounds. Finally, we give a lemma for verifying the repulsion axiom (iv) of the real replacement principle in

the case when the atom distribution is gaussian. We use the usual exterior product $\wedge : \mathbb{C}^{n+1} \times \mathbb{C}^{n+1} \rightarrow \wedge^2 \mathbb{C}^{n+1}$ on the vector space \mathbb{C}^{n+1} , in particular

$$|v \wedge w| = \left(\sum_{0 \leq i < j \leq n} |v_i w_j - v_j w_i|^2 \right)^{1/2}$$

for any $v = (v_0, \dots, v_n)$ and $w = (w_0, \dots, w_n)$ in \mathbb{C}^{n+1} .

Lemma 17 (Repulsion of zeroes). *Let $n \geq 1$, and let $f = f_n$ be a random polynomial of the type in Definition 1, with real coefficients c_0, \dots, c_n and with atom distribution ξ given by the real gaussian $N(0, 1)_{\mathbb{R}}$. Let x be a real number, let $C > 1$ and $r_0 > 0$. Let $R : B(x, r_0) \rightarrow \mathbb{C}$ be a holomorphic function that is nonvanishing in $B(x, r_0)$, and let $v : \mathbb{C} \rightarrow \mathbb{C}^{n+1}$ be the vector valued holomorphic function*

$$v(z) := (R(z)c_i z^i)_{i=0}^n.$$

Assume the axiom

$$(17) \quad |v(z)| \leq C$$

for all $z \in B(x, r_0)$, as well as the axiom

$$(18) \quad \left| v(x) \wedge \frac{d}{dx} v(x) \right| \geq C^{-1}.$$

Then, if δ is sufficiently small depending on r_0 and C , one has the real repulsion estimate

$$(19) \quad \rho_f^{(2,0)}(x, x + \delta) = O(\delta).$$

and the complex repulsion estimate

$$(20) \quad \rho^{(0,1)}(x + \sqrt{-1}\delta) = O(\delta).$$

Here the implied constants are allowed to depend on C and r_0 .

We prove this lemma in Section 11. Our main tools will be the Kac-Rice formula [26], [28], [47], the Cauchy integral formula, and certain geometric arguments. The holomorphic factor $R(z)$ should be viewed as a normalization factor that one is free to choose in order to make the two hypotheses (17), (18) of the lemma hold simultaneously.

5. UNIVERSALITY OF THE CORRELATION FUNCTIONS OF THE CLASSICAL ENSEMBLES

We now specialize the above results to the classical ensembles mentioned in the introduction, namely the flat, elliptic, hyperbolic, and Kac polynomials.

5.1. Flat polynomials. We begin with the case of flat polynomials (or Weyl polynomials), i.e. random polynomials of the form in Definition 1 with $c_i := \frac{1}{\sqrt{i!}}$. Under extremely mild assumptions on the atom distribution³ ξ , it was shown by Kabluchko and Zaporozhets [25, Theorem 2.3] that the zeroes of such polynomials obeyed the circular law, thus for instance for any Jordan-measurable subset Ω of the complex plane (e.g. a ball or a rectangle), one has

$$\frac{1}{n}N_\Omega \rightarrow \int_\Omega \frac{1}{2\pi} 1_{B(0,1)}(z/\sqrt{n}) dz$$

both in probability and in the almost sure sense as $n \rightarrow \infty$ (assuming the atom distribution ξ is independent of n). Thus, in particular, the bulk of the zeroes should lie inside the disk $B(0, \sqrt{n})$ and be uniformly distributed within that disk at global scales (i.e. at scales comparable to \sqrt{n}). As such, we expect the mean eigenvalue spacing to be comparable to 1.

We now can present our main universality results for flat polynomials at local scales.

Theorem 18 (Two moment theorem for flat polynomials; complex case). *Let $k \geq 1$, $\varepsilon > 0$, and $C > 0$ be constants. Let n be a natural number, Let $f_{n,\xi}, f_{n,\tilde{\xi}}$ be flat polynomials with atom distributions $\xi, \tilde{\xi}$ being complex random variables of mean zero and variance one, matching moments to second order and also obeying the bounds $\mathbf{E}|\xi|^{2+\varepsilon}, \mathbf{E}|\tilde{\xi}|^{2+\varepsilon} \leq C$. Let $z_1, \dots, z_k \in \mathbb{C}$ be quantities depending on n with $n^\varepsilon \leq |z_i| \leq \sqrt{n} + C$ for all $i = 1, \dots, k$.*

Let $G : \mathbb{C}^k \rightarrow \mathbb{C}$ be a smooth function supported on the polydisc $B(0, C)^k$ that obeys the bounds

$$|\nabla^a G(w)| \leq C$$

for all $0 \leq a \leq 5k + 1$ and all $w \in \mathbb{C}^k$. Then

$$\left| \int_{\mathbb{C}^k} G(w_1, \dots, w_k) \rho_{f_{n,\xi}}^{(k)}(z_1 + w_1, \dots, z_k + w_k) dw_1 \dots dw_k \right. \\ \left. - \int_{\mathbb{C}^k} G(w_1, \dots, w_k) \rho_{f_{n,\tilde{\xi}}}^{(k)}(z_1 + w_1, \dots, z_k + w_k) dw_1 \dots dw_k \right| \leq \tilde{C} n^{-c_0}$$

for some \tilde{C} depending only on k, ε, C , and some $c_0 > 0$ depending only on ε .

Informally, this theorem establishes local universality of the zeroes in the bulk and edge of the spectrum, except when one is near the origin. Note that we do not expect universality near the origin, since $\mathbf{P}(f_{n,\xi}(0) = 0) = \mathbf{P}(\xi = 0)$ is clearly not universal in ξ ; see [7] for further discussion of this issue (in the context of elliptic polynomials rather than flat ones). Away from the disk $B(0, \sqrt{n})$, one expects very few zeroes, which suggests that Theorem 18 should also hold in this case, but our methods do not cover this regime.

³In [25], ξ does not even need to have finite mean or variance; the hypothesis $\mathbf{E} \log(1+|\xi|) < \infty$ suffices. It is unlikely however that weak hypotheses continue to suffice for local universality. For instance, in [37] it was shown that the number of real zeroes of a Kac polynomial changes significantly when one takes ξ to be drawn from the Cauchy distribution rather than from a distribution of finite variance; see also [24] for some stronger and more general results in this direction.

We also remark that a result similar to Theorem 18 has recently been established (by a rather different method) by Ledoan, Merkli, and Starr [31]. In our language, the results in [31] establish universality for the distribution of the random variable

$$\sum_{i=1}^n \phi(\zeta_i - z)$$

where ϕ is a continuous, compactly supported function independent of n , and z is close to the boundary of the spectrum. The main idea is to establish a central limit theorem for a normalized partial Taylor series expansion of f around z . Their argument is simpler than the one given here, but does not appear to give a uniform polynomial rate of convergence as in Theorem 18 (or Theorem 19 below), which is needed in some of our applications.

In the case when the coefficients are real, we obtain the following two moment theorem for the mixed correlation functions.

Theorem 19 (Two moment theorem for flat polynomials; real case). *Let $k, l \geq 0$, $\varepsilon > 0$, and $C > 0$ be constants with $k + l > 0$. Let n be a natural number, Let $f_{n,\xi}, f_{n,\tilde{\xi}}$ be flat polynomials with atom distributions $\xi, \tilde{\xi}$ being real random variables of mean zero and variance one obeying the bounds $\mathbf{E}|\xi|^{2+\varepsilon}, \mathbf{E}|\tilde{\xi}|^{2+\varepsilon} \leq C$. Let $x_1, \dots, x_k \in \mathbb{R}$ and $z_1, \dots, z_l \in \mathbb{C}$ be quantities depending on n with $n^\varepsilon \leq |x_i|, |z_j| \leq \sqrt{n} + C$ for all $i = 1, \dots, k, j = 1, \dots, l$. Let $G : \mathbb{R}^k \times \mathbb{C}^l \rightarrow \mathbb{C}$ be a smooth function supported on $[-C, C]^k \times B(0, C)^l$ that obeys the bounds*

$$|\nabla^a G(w)| \leq C$$

for all $0 \leq a \leq 5(k + l) + 1$ and all $w \in \mathbb{R}^k \times \mathbb{C}^l$. Then

$$\begin{aligned} & \left| \int_{\mathbb{R}^k} \int_{\mathbb{C}^l} G(y_1, \dots, y_k, w_1, \dots, w_l) \right. \\ & \quad \rho_{f_{n,\xi}}^{(k,l)}(x_1 + y_1, \dots, x_k + y_k, z_1 + w_1, \dots, z_l + w_l) dw_1 \dots dw_l dy_1 \dots dy_l \\ & \quad - \int_{\mathbb{R}^k} \int_{\mathbb{C}^l} G(y_1, \dots, y_k, w_1, \dots, w_l) \\ & \quad \left. \rho_{f_{n,\tilde{\xi}}}^{(k,l)}(x_1 + y_1, \dots, x_k + y_k, z_1 + w_1, \dots, z_l + w_l) dw_1 \dots dw_l dy_1 \dots dy_l \right| \\ & \leq \tilde{C} n^{-c_0} \end{aligned}$$

where \tilde{C} depends only on C, k, l , and $c_0 > 0$ depends only on k, l .

We prove these theorems as consequences of the previously stated results in Section 12. As an application of these results we are able to establish some new results about the number $N_{\mathbb{R}}$ of real eigenvalues of flat polynomials:

Theorem 20 (Number of real zeroes of polynomials). *Let n be a natural number, Let $f_{n,\xi}$ be a flat polynomial with atom distributions ξ being a real random variables of mean zero and variance one obeying the bound $\mathbf{E}|\xi|^{2+\varepsilon} \leq C$ for some $C, \varepsilon > 0$. Then one has*

$$\mathbf{E}N_{\mathbb{R}} = \frac{2}{\pi} \sqrt{n} + O(n^{1/2-c}),$$

where the implied constant in the $O()$ notation depends only on C, ε . More generally, for any interval $I \subset \mathbb{R}$, one has

$$\mathbf{E}N_I = \frac{1}{\pi} |I \cap [-\sqrt{n}, \sqrt{n}]| + O(n^{1/2-c}).$$

We establish this result in Section 12 also. With some additional calculation that we will sketch in that section, one can also obtain the bound $\mathbf{Var}N_I = O(n^{1-c})$, which by Chebyshev's inequality then tells us that

$$N_I = \frac{1}{\pi} |I \cap [-\sqrt{n}, \sqrt{n}]| + O(n^{1/2-c})$$

with probability $1 - O(n^{-c})$. Informally, this asserts that a global scales, the real zeroes of a flat real polynomial are asymptotically uniformly distributed in $[-\sqrt{n}, \sqrt{n}]$ with intensity $1/\pi$. As a matter of fact, our local universality results allow us to consider the number of real zeros in intervals of length $O(1)$.

5.2. Elliptic polynomials. We turn now to the elliptic polynomials

$$f'_{n,\xi}(z) = \sum_{i=0}^n \sqrt{\binom{n}{i}} \xi_i z^i,$$

where we normalize ξ to have mean zero and variance one. As shown in [25], the majority of the zeroes of this polynomial have norm $O(1)$ asymptotically almost surely. As a matter of fact, the limiting density function is $\frac{n}{\pi}(1 + |z|^2)^{-2}$; see [25]. In particular, we expect the typical separation between zeroes to be of the order of $1/\sqrt{n}$, in contrast to the flat case. In order to renormalize the typical separation between zeroes to be comparable to one (which is the scale to which the replacement principle is adapted), we replace the polynomial $f'_{n,\xi}(z) = \sum_{i=0}^n \sqrt{\binom{n}{i}} \xi_i z^i$ by the *rescaled* version

$$(21) \quad f_{n,\xi}(z) := \sum_{i=0}^n \sqrt{\binom{n}{i}} n^{-i} \xi_i z^i.$$

It is clear that if z is a zero of the non-scaled polynomial then \sqrt{z} is a zero of the rescaled one. In the following theorems (and their proofs), f_n is the rescaled polynomial. By the results of [25], the limiting density function for the rescaled elliptic functions is given by the formula

$$(22) \quad \rho_\varepsilon(z) := \frac{1}{\pi} (1 + |z|^2/n)^{-2}.$$

This can be compared with the limiting density $\frac{1}{\pi} \mathbf{1}_{B(0, \sqrt{n})}(z)$ for flat polynomials.

We can now give the analogues of Theorems 18, 19.

Theorem 21 (Two moment theorem for elliptic polynomials; complex case). *Let $k \geq 1$, $\varepsilon > 0$, and $C > 0$ be constants. Let n be a natural number and $f_{n,\xi}, f_{n,\tilde{\xi}}$ be rescaled elliptic polynomials with atom distributions $\xi, \tilde{\xi}$ being complex random variables of mean zero and variance one, matching moments to second order and*

also obeying the bounds $\mathbf{E}|\xi|^{2+\varepsilon}, \mathbf{E}|\tilde{\xi}|^{2+\varepsilon} \leq C$. Let $z_1, \dots, z_k \in \mathbb{C}$ be quantities depending on n with $n^\varepsilon \leq |z_i| \leq C\sqrt{n}$ for all $i = 1, \dots, k$.

Let $G : \mathbb{C}^k \rightarrow \mathbb{C}$ be a smooth function supported on the polydisc $B(0, C)^k$ that obeys the bounds

$$|\nabla^a G(w)| \leq C$$

for all $0 \leq a \leq 5k + 1$ and all $w \in \mathbb{C}^k$. Then

$$\left| \int_{\mathbb{C}^k} G(w_1, \dots, w_k) \rho_{f_{n,\xi}}^{(k)}(z_1 + w_1, \dots, z_k + w_k) dw_1 \dots dw_k \right. \\ \left. - \int_{\mathbb{C}^k} G(w_1, \dots, w_k) \rho_{f_{n,\tilde{\xi}}}^{(k)}(z_1 + w_1, \dots, z_k + w_k) dw_1 \dots dw_k \right| \leq \tilde{C}n^{-c_0}$$

for some \tilde{C} depending only on k, ε, C , and some $c_0 > 0$ depending only on ε .

Note that we allow our reference points z_1, \dots, z_k in the spectrum to have magnitude as large as $C\sqrt{n}$, as compared to the flat case where we can only establish universality up to magnitude $\sqrt{n} + C$. This reflects the different nature of the spectrum in the elliptic case, which does not have an 'edge' at $\{|z| = \sqrt{n}\}$ in contrast to the flat case. Note that the reflected polynomial $\tilde{f}_{n,\xi}(z) := z^n f_{n,\xi}(1/z)$ has the same distribution as $f_{n,\xi}$, so the law of the zeroes of $f_{n,\xi}$ is invariant with respect to the inversion map $z \mapsto 1/z$. Because of this, one can invert Theorem 21 (and Theorem 22 below) to give universality results in the region $\sqrt{n}/C \leq |z| \leq n^{1-\varepsilon}$ as well (albeit with some additional Jacobian factors that are powers of the $|z_i|$). We omit the details.

Theorem 22 (Two moment theorem for elliptic polynomials; real case). *Let $k, l \geq 0$, $\varepsilon > 0$, and $C > 0$ be constants with $k + l > 0$. Let n be a natural number and $f_{n,\xi}, f_{n,\tilde{\xi}}$ be rescaled elliptic polynomials with atom distributions $\xi, \tilde{\xi}$ being real random variables of mean zero and variance one obeying the bounds $\mathbf{E}|\xi|^{2+\varepsilon}, \mathbf{E}|\tilde{\xi}|^{2+\varepsilon} \leq C$. Let $x_1, \dots, x_k \in \mathbb{R}$ and $z_1, \dots, z_l \in \mathbb{C}$ be quantities depending on n with $n^\varepsilon \leq |x_i|, |z_j| \leq C\sqrt{n}$ for all $i = 1, \dots, k, j = 1, \dots, l$. Let $G : \mathbb{R}^k \times \mathbb{C}^l \rightarrow \mathbb{C}$ be a smooth function supported on $[-C, C]^k \times B(0, C)^l$ that obeys the bounds*

$$|\nabla^a G(w)| \leq C$$

for all $0 \leq a \leq 5(k + l) + 1$ and all $w \in \mathbb{R}^k \times \mathbb{C}^l$. Then

$$\left| \int_{\mathbb{R}^k} \int_{\mathbb{C}^l} G(y_1, \dots, y_k, w_1, \dots, w_l) \right. \\ \left. \rho_{f_{n,\xi}}^{(k,l)}(x_1 + y_1, \dots, x_k + y_k, z_1 + w_1, \dots, z_l + w_l) dw_1 \dots dw_l dy_1 \dots dy_l \right. \\ \left. - \int_{\mathbb{R}^k} \int_{\mathbb{C}^l} G(y_1, \dots, y_k, w_1, \dots, w_l) \right. \\ \left. \rho_{f_{n,\tilde{\xi}}}^{(k,l)}(x_1 + y_1, \dots, x_k + y_k, z_1 + w_1, \dots, z_l + w_l) dw_1 \dots dw_l dy_1 \dots dy_l \right| \\ \leq \tilde{C}n^{-c_0}$$

where \tilde{C} depends only on C, k, l , and $c_0 > 0$ depends only on k, l .

Finally, we can give an analogue of Theorem 20:

Theorem 23 (Real zeroes). *Let $f_{n,\xi}$ be a rescaled elliptic polynomial with ξ being a real random variable of mean zero and variance one obeying the bound $\mathbf{E}|\xi|^{2+\varepsilon} \leq C$ for some $C, \varepsilon > 0$. Then one has*

$$\mathbf{E}N_{\mathbb{R}} = \sqrt{n} + O(n^{1/2-c})$$

for some $c > 0$ depending only on ε , where the implied constant depends only on C, ε . More generally, one has

$$\mathbf{E}N_I = \int_I \frac{1}{\pi} \frac{dx}{1+x^2/n} + O(n^{1/2-c})$$

for any interval $I \subset \mathbb{R}$.

We establish these results in Section 13.

5.3. Kac polynomials. We now turn to the situation of Kac polynomials $f(z) = f_{n,\xi}(z) = \sum_{i=0}^n \xi_i z^i$. In the case that the atom distribution ξ is a complex Gaussian $N(0, 1)_{\mathbb{C}}$, then the first intensity $\rho_n^{(1)}$ can be computed explicitly from either the Kac-Rice formula [26], [28], [47] or the formula of Edelman and Kostlan [12] as

$$\rho_n^{(1)}(z) = \frac{1}{4\pi} \Delta \log \sum_{i=0}^n |z|^{2i}$$

which can be shown to be $(1+o(1))n^2 F(a)$ if $||z| - 1| = \frac{a}{n}$ for constant $a > 0$, with

$$F(a) := \frac{1 - (a/\sinh a)^2}{4\pi a^2};$$

see [24].

In particular, this shows that the zeroes concentrate uniformly around the unit circle.

It turns out that there are some additional technical difficulties when using the methods of this paper to study Kac polynomials instead of elliptic or flat polynomials. The singular nature of the limit first intensity at the unit circle is the most obvious such difficulty, but a less obvious difficulty is the partial breakdown of concentration of the log-magnitude $\log|f(z)|$ when z is a root of unity. For instance, consider the log-magnitude

$$\log|f_{n,\xi}(1)| = \left| \sum_{i=0}^n \xi_i \right|$$

at 1. If the atom distribution ξ is Bernoulli and n is odd, then the RHS equals 0 with probability $\frac{\binom{n+1}{2}}{2^{n+1}} = \Theta(n^{-1/2})$. Therefore, the logarithm diverges to $-\infty$ with probability $\Theta(n^{-1/2})$, which is not strong enough for the purposes of Proposition 12. In a similar spirit, the log-magnitude $\log|f_{n,\xi}(e^{2\pi\sqrt{-1}a/b})|$ when a, b are coprime integers with b bounded can be shown in the Bernoulli case to diverge to $-\infty$ with probability $\Theta(n^{-b/2})$; we omit the details. To overcome this difficulty, we make use of recent results in both inverse Littlewood-Offord theory (see [44]) and quantitative versions of Gromov's theorem (see [48]). With these tools, we are able to show that the roots of unity are essentially the only new obstruction to this

concentration, allowing the rest of the theory to go through without much further modification. In particular, we can establish the following local universality results for Kac polynomials;

Theorem 24 (Two moment theorem for Kac polynomials; complex case). *Let $k \geq 1$, $\varepsilon > 0$, and $C > 0$ be constants. Let n be a natural number, Let $f_{n,\xi}, f_{n,\tilde{\xi}}$ be Kac polynomials with atom distributions $\xi, \tilde{\xi}$ being complex random variables of mean zero and variance one, matching moments to second order and also obeying the bounds $\mathbf{E}|\xi|^{2+\varepsilon}, \mathbf{E}|\tilde{\xi}|^{2+\varepsilon} \leq C$. Let $1/n \leq r \leq n^{-\varepsilon}$ be a radius, and let $z_1, \dots, z_k \in \mathbb{C}$ be quantities depending on n with*

$$r \leq \frac{1}{n} + ||z_i| - 1| \leq 2r$$

for all $i = 1, \dots, k$.

Let $G : \mathbb{C}^k \rightarrow \mathbb{C}$ be a smooth function supported on the polydisc $B(0, 10^{-3})^k$ that obeys the bounds

$$|\nabla^a G(w)| \leq C$$

for all $0 \leq a \leq 5k + 1$ and all $w \in \mathbb{C}^k$. Then

$$\left| \int_{\mathbb{C}^k} G(w_1, \dots, w_k) r^{2k} \rho_{f_{n,\xi}}^{(k)}(z_1 + rw_1, \dots, z_k + rw_k) dw_1 \dots dw_k - \int_{\mathbb{C}^k} G(w_1, \dots, w_k) r^{2k} \rho_{f_{n,\tilde{\xi}}}^{(k)}(z_1 + rw_1, \dots, z_k + rw_k) dw_1 \dots dw_k \right| \leq \tilde{C} n^{-c_0}$$

for some \tilde{C} depending only on k, ε, C , and some $c_0 > 0$ depending only on ε .

Note that the reference points z_1, \dots, z_k are required to remain at essentially the same distance r from the unit circle. It is possible to use the methods of this paper to obtain more general local universality results when the z_1, \dots, z_k are at widely differing distances from the unit circle, but this requires the generalization of Theorem 5 alluded to in Remark 9, and we omit this generalization in order to simplify the exposition.

As usual, we have an analogue of the above local universality result in the real case:

Theorem 25 (Two moment theorem for Kac polynomials; real case). *Let $k, l \geq 0$, $\varepsilon > 0$, and $C > 0$ be constants with $k + l > 0$. Let n be a natural number, Let $f_{n,\xi}, f_{n,\tilde{\xi}}$ be Kac polynomials with atom distributions $\xi, \tilde{\xi}$ being real random variables of mean zero and variance one obeying the bounds $\mathbf{E}|\xi|^{2+\varepsilon}, \mathbf{E}|\tilde{\xi}|^{2+\varepsilon} \leq C$. Let $1/n \leq r \leq n^{-\varepsilon}$ be a radius, and let $x_1, \dots, x_k \in \mathbb{R}$ and $z_1, \dots, z_l \in \mathbb{C}$ be quantities depending on n with*

$$r \leq \frac{1}{n} + ||x_i| - 1|, \frac{1}{n} + ||z_j| - 1| \leq 2r$$

for all $i = 1, \dots, k$, $j = 1, \dots, l$. Let $G : \mathbb{R}^k \times \mathbb{C}^l \rightarrow \mathbb{C}$ be a smooth function supported on $[-10^{-3}, 10^{-3}]^k \times B(0, 10^{-3})^l$ that obeys the bounds

$$|\nabla^a G(w)| \leq C$$

for all $0 \leq a \leq 5(k+l) + 1$ and all $w \in \mathbb{R}^k \times \mathbb{C}^l$. Then

$$\begin{aligned} & \left| \int_{\mathbb{R}^k} \int_{\mathbb{C}^l} G(y_1, \dots, y_k, w_1, \dots, w_l) \right. \\ & \quad \left. r^{2k+l} \rho_{f_{n,\xi}}^{(k,l)}(x_1 + ry_1, \dots, x_k + ry_k, z_1 + rw_1, \dots, z_l + rw_l) dw_1 \dots dw_l dy_1 \dots dy_l \right. \\ & \quad \left. - \int_{\mathbb{R}^k} \int_{\mathbb{C}^l} G(y_1, \dots, y_k, w_1, \dots, w_l) \right. \\ & \quad \left. r^{2k+l} \rho_{f_{n,\xi}}^{(k,l)}(x_1 + ry_1, \dots, x_k + ry_k, z_1 + rw_1, \dots, z_l + rw_l) dw_1 \dots dw_l dy_1 \dots dy_l \right| \\ & \leq \tilde{C} n^{-c_0} \end{aligned}$$

where \tilde{C} depends only on C, k, l , and $c_0 > 0$ depends only on k, l .

Notice that a rescaling already took place in the conclusion of the theorems, so we do not need to rescale f here. We establish these results in Theorem 14. As far as real roots are concerned, our results yield statements about the distribution of number of real roots in short l intervals (where the expectation of the number of real roots is $\Theta(1)$). To our best knowledge, such results have not been obtained anywhere else for general Kac polynomials. On the other hand, on the global scale, we do not obtain a better estimate than Ibragimov-Maslova bound.

5.4. General polynomials. Our result applies for general random polynomials of the form $f_{n,\xi} = \sum_{i=0}^n c_i \xi_i z^i$ where the (deterministic) coefficients c_i need to satisfy some mild conditions, but otherwise can be fairly arbitrary. Thus, one can use our result to derive information about the zeroes (in particular the real zeroes) of these polynomials.

As an example, the expectation of the number of real zeroes of $f_{n,\xi}$ in an interval can be computed using Kac formula or Edelman-Kostlan formula when ξ_i are iid standard real gaussian. Our universality result (for the real case) would should that this expectation remains asymptotically the same when the atom variable ξ is Bernoulli. As far as we know, prior to this paper, no general method has been available to prove such a result. The reader is invited to work out a few examples.

6. PROOF OF THE REPLACEMENT PRINCIPLE, COMPLEX CASE

In this section we establish Theorem 5. We will use the approach developed in [58, §6].

Fix k, C, r_0, c_0, a_0 as in Theorem 5; all implied constants in the $O()$ notation will be allowed to depend on these parameters. Let A be defined by (5), let n be a natural number, and let z_1, \dots, z_k be complex numbers and f, \tilde{f} be random polynomials obeying the hypotheses of the theorem. We may assume that n is sufficiently large depending on the parameters k, C, r_0, c_0, a_0 , as the claim is trivial otherwise.

By conditioning out the event that f or \tilde{f} vanish identically (which by the non-degeneracy axiom (i) only occurs with probability $O(n^{-A})$), we may assume that f and \tilde{f} are non-vanishing almost surely, as this conditioning does not significantly alter the hypotheses (i)-(iii) or conclusion of the theorem (after adjusting C by a multiplicative constant). This conditioning might destroy any independence properties enjoyed by the coefficients of the f, \tilde{f} , but this will not be an issue as such independence properties are not directly assumed in Theorem 5.

The first observation to use Fourier analysis to reduce to proving the following variant of the conclusion of Theorem 5: we have the bound

$$(23) \quad \left| \int_{\mathbb{C}^k} G(w_1, \dots, w_k) \rho_f^{(k)}(z_1 + w_1, \dots, z_k + w_k) dw_1 \dots dw_k \right. \\ \left. - \int_{\mathbb{C}^k} G(w_1, \dots, w_k) \rho_{\tilde{f}}^{(k)}(z_1 + w_1, \dots, z_k + w_k) dw_1 \dots dw_k \right| \ll n^{-c_0/4},$$

whenever G is a function of the form

$$(24) \quad G(w_1, \dots, w_k) = G_1(w_1) \dots G_k(w_k)$$

for some smooth $G_1, \dots, G_k : \mathbb{C} \rightarrow \mathbb{C}$ supported in $B(0, 10r_0)$ and such that

$$(25) \quad |\nabla^a G_j(w)| \ll 1$$

for all $0 \leq a \leq a_0$ and $1 \leq j \leq k$.

Indeed, suppose we had the bound (23). Now let G be a function of the form required for Theorem 5. We view $B(0, r_0)$ as a subset of the square $[-1.1r_0, 1.1r_0]^2$, which in turn we can identify with the torus $(\mathbb{R}/2.2r_0\mathbf{Z})^2$. Thus G can be viewed as a smooth function on the torus $(\mathbb{R}/(2.2r_0)\mathbf{Z})^{2k}$. We can then expand G as a Fourier series

$$G(w) = \sum_{b,c \in \mathbf{Z}^k} g_{b,c} e^{2\pi\sqrt{-1}(b \cdot \operatorname{Re}(w) + c \cdot \operatorname{Im}(w))/(2.2r_0)}$$

in $[-1.1r_0, 1.1r_0]^2$, where the Fourier coefficients $g_{b,c}$ are given by the formula

$$g_{b,c} := (2.2r_0)^{-2k} \int_{B(0, r_0)^k} e^{2\pi\sqrt{-1}(b \cdot \operatorname{Re}(w) + c \cdot \operatorname{Im}(w))/(r_0/4)} G(w) dw.$$

Let $\eta : \mathbb{R} \rightarrow \mathbb{R}$ be a function supported on $[-1.1r_0, 1.1r_0]$ that equals one on $[-r_0, r_0]$. We can then write

$$G(w) = \sum_{b,c \in \mathbf{Z}^k} G_{b,c}(w)$$

for all $w \in \mathbb{C}^k$, where

$$G_{b,c}(w) := g_{b,c} \prod_{i=1}^k \psi_{b,c,i}(w_i)$$

and

$$\psi_{b,c,i}(w_i) := \prod_{i=1}^k e^{2\pi\sqrt{-1}(b_i \operatorname{Re}(w_i) + c_i \operatorname{Im}(w_i))/(2.2r_0)} \eta(\operatorname{Re}(w_i)) \eta(\operatorname{Im}(w_i)).$$

Observe that $\psi_{b,c,i}$ is supported on $B(0, 10r_0)$ and that

$$|\nabla^a G_{b,c}(w)| \ll (1 + |b| + |c|)^{a_0} |g_{b,c}|$$

for all $w \in \mathbb{C}^k$ and $0 \leq a \leq a_0$. From (23) and the triangle inequality, we conclude that

$$\begin{aligned} & \left| \int_{\mathbb{C}^k} G(w_1, \dots, w_k) \rho_f^{(k)}(z_1 + w_1, \dots, z_k + w_k) dw_1 \dots dw_k \right. \\ & \quad \left. - \int_{\mathbb{C}^k} G(w_1, \dots, w_k) \rho_{\tilde{f}}^{(k)}(z_1 + w_1, \dots, z_k + w_k) dw_1 \dots dw_k \right| \\ & \ll n^{-c_0/4} \sum_{b, c \in \mathbf{Z}^k} |g_{b, c}| (1 + |b| + |c|)^{a_0}. \end{aligned}$$

On the other hand, from (7) and integration by parts we have

$$|g_{b, c}| \ll (1 + |b| + |c|)^{-(a_0 + 2k + 1)} M$$

and Theorem 5 follows.

Now let G be of the form (24). For any $\alpha > 0$, we call a statistic $X(f) \in \mathbb{C}$ of a random polynomial f α -insensitive if one has

$$(26) \quad \mathbf{E}|X(f) - X(\tilde{f})| = O(n^{-\alpha}).$$

Thus, for instance, the comparability axiom (iii) tells us that the statistic

$$F(\log |f(z'_1)|, \dots, \log |f(z'_{k'})|)$$

is c_0 -insensitive for all $1 \leq k' \leq n^{c_0}$, $z'_1, \dots, z'_{k'} \in \bigcup_{i=1}^k B(z_i, 20r_0)$, and smooth $F : \mathbb{C}^{k'} \rightarrow \mathbb{C}$ obeying the derivative bounds

$$|\nabla^a F(w)| \leq n^{c_0}$$

for all $w \in \mathbb{C}^{k'}$ and $0 \leq a \leq a_0$.

It now suffices to show that the statistic

$$(27) \quad \int_{\mathbb{C}^k} G(w_1, \dots, w_k) \rho_n^{(k)}(z_1 + w_1, \dots, z_k + w_k) dw_1 \dots dw_k$$

is $c_0/4$ -insensitive.

Let ζ_1, \dots, ζ_n denote the zeroes of f . By (1), the quantity (27) is equal to

$$(28) \quad \mathbf{E} \sum_{i_1, \dots, i_k \text{ distinct}} G(\zeta_{i_1} - z_1, \dots, \zeta_{i_k} - z_k)$$

By the inclusion-exclusion formula, we may decompose this expression as

$$(29) \quad \mathbf{E} \prod_{j=1}^k X_{z_j, G_j}$$

plus a bounded number of lower order terms which are of the form (29) for a smaller value of k (and different choices of G_j , and a subset of the $\{z_1, \dots, z_k\}$), where $X_{z_j, G_j} = X_{z_j, G_j}(f)$ denotes the linear statistic

$$(30) \quad X_{z_j, G_j} := \sum_{i=1}^n G_j(\zeta_i - z_j).$$

For instance, in the $k = 2$ case, we have

$$\begin{aligned} \sum_{1 \leq i \neq j \leq n} G_1(\zeta_i - z_1) G_2(\zeta_j - z_2) &= \left[\sum_{i=1}^n G_1(\zeta_i - z_1) \right] \left[\sum_{j=1}^n G_2(\zeta_j - z_2) \right] \\ &\quad - \sum_{i=1}^n G_1(\zeta_i - z_1) G_2(\zeta_i - z_2) \\ &= X_{z_1, G_1} X_{z_2, G_2} - X_{z_1, G_3}, \end{aligned}$$

where

$$G_3(\zeta) := G_1(\zeta) G_2(\zeta - z_2 + z_1).$$

Note that G_3 obeys similar bounds (25) to G_1, G_2 , though with a slightly different choice of implied constant. Clearly, similar decompositions are also available for more general values of k .

By induction on k , it thus suffices to show that the expression (29) is $c_0/4$ -insensitive. By the non-clustering hypothesis, we have $X_{z_j, G_j} = O(n^{1/A})$ with probability at least $1 - O(n^{-A})$ for each $1 \leq j \leq k$, while from the pointwise bounds (25) we have the crude deterministic bound $X_{z_j, G_j} = O(n)$. To use these bounds, we introduce a smooth approximation $P(\zeta_1, \dots, \zeta_k)$ of the product $\zeta_1 \dots \zeta_k$ such that

- (i) $P(\zeta_1, \dots, \zeta_k) = \zeta_1 \dots \zeta_k$ on $B(0, n^{2/A})^k$;
- (ii) P is supported on $B(0, 2n^{2/A})^k$; and
- (iii) P obeys the derivative bounds

$$(31) \quad |\nabla^a P(\zeta_1, \dots, \zeta_k)| \ll n^{2k/A} = n^{\frac{c_0}{50a_0}}$$

for all $0 \leq a \leq a_0$ and $\zeta_1, \dots, \zeta_k \in \mathbb{C}$.

For instance, we may define P explicitly by the formula

$$P(\zeta_1, \dots, \zeta_k) := \prod_{i=1}^k \zeta_i \phi(|\zeta_i|/n^{2/A})$$

where ϕ is a smooth function supported on $[-2, 2]$ that equals 1 on $[-1, 1]$; it is easy to see that this choice of P obeys all the axioms claimed above.

Using the non-clustering axiom (ii), we have

$$\prod_{j=1}^k X_{z_j, G_j} = P(X_{z_1, G_1}, \dots, X_{z_k, G_k})$$

with probability $1 - O(n^{-A})$, and we have the crude deterministic bound

$$\prod_{j=1}^k X_{z_j, G_j} = P(X_{z_1, G_1}, \dots, X_{z_k, G_k}) + O(n^k)$$

outside of this event. Taking expectations, we conclude that

$$\mathbf{E} \prod_{j=1}^k X_{z_j, G_j} = \mathbf{E} P(X_{z_1, G_1}, \dots, X_{z_k, G_k}) + O(n^{-A} n^k).$$

By (5), we certainly have $n^{-A} n^k = O(n^{-c_0/4})$. It thus suffices to show that the expression

$$(32) \quad \mathbf{E} P(X_{z_1, G_1}, \dots, X_{z_k, G_k})$$

is $c_0/4$ -insensitive.

From the fundamental theorem of algebra we have

$$(33) \quad \log |f(z)| = a_n + \sum_{i: \zeta_i \neq \infty} \log |\zeta_i - z|$$

for all $z \in \mathbb{C}$ and some almost surely finite quantity a_n independent of z . (Here we are using the previous reduction that f almost surely does not vanish identically.) By Green's theorem, (30), and the smooth compactly supported nature of G_j , we conclude that

$$X_{z_j, G_j} = \int_{\mathbb{C}} \log |f(z)| H_j(z) dz$$

where

$$H_j(z) := -\frac{1}{2\pi} \Delta G_j(z - z_j)$$

and $\Delta = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}$ is the Laplacian on \mathbb{C} . Note that H_j is a bounded smooth function supported on $B(z_j, C)$.

We now recall a standard sampling lemma from [58, Lemma 38]:

Lemma 26 (Monte Carlo sampling lemma). *Let (X, μ) be a probability space, and let $F : X \rightarrow \mathbb{C}$ be a square-integrable function. Let $m \geq 1$, let x_1, \dots, x_m be drawn independently at random from X with distribution μ , and let S be the empirical average*

$$S := \frac{1}{m} (F(x_1) + \dots + F(x_m)).$$

Then S has mean $\int_X F d\mu$ and variance $\int_X (F - \int_X F d\mu)^2 d\mu$. In particular, by Chebyshev's inequality, one has

$$\mathbf{P}(|S - \int_X F d\mu| \geq \lambda) \leq \frac{1}{m\lambda^2} \int_X (F - \int_X F d\mu)^2 d\mu$$

for any $\lambda > 0$. Equivalently, for any $\delta > 0$ one has the bound

$$|S - \int_X F d\mu| \leq \frac{1}{\sqrt{m\delta}} \left(\int_X (F - \int_X F d\mu)^2 d\mu \right)^{1/2}$$

with probability at least $1 - \delta$.

Proof. The random variables $F(x_i)$ for $i = 1, \dots, m$ are jointly independent with mean $\int_X F d\mu$ and variance $\frac{1}{m} \int_X (F - \int_X F d\mu)^2 d\mu$. Averaging these variables, we obtain the claim. \square

Ideally, we would like to use the Markov sampling method (Lemma 26) to approximate $\int_{\mathbb{C}} \log |f(z)| H_j(z) dz$. However, there is an obstacle: as f can have many zeroes far from z_j , the error term given in Lemma 26 can be too large. To overcome this difficulty, we introduce a method to reduce the variance by exploiting the cancellation properties of the function H_j . Indeed, note that H_j is the Laplacian of a smooth compactly supported function, thus it is orthogonal to any (affine, real-) linear function by integration by parts. To exploit this, we define a (random) affine real-linear function $L_j : \mathbb{C} \rightarrow \mathbb{C}$ by first selecting a reference complex number $w_{j,0}$ drawn uniformly at random from $B(z_j, 1)$ (independently of all previous random quantities), and defining $L_j(z)$ to be the random affine real-linear function of $\operatorname{Re}(z), \operatorname{Im}(z)$ that equals $\log |f(z)|$ when $z = w_{j,0}, w_{j,0} + 1, w_{j,0} + \sqrt{-1}$. More explicitly, we have

$$(34) \quad \begin{aligned} L(z) &:= \log |f(w_{j,0})| \\ &+ (\log |f(w_{j,0} + 1)| - \log |f(w_{j,0})|) \operatorname{Re}(z - w_{j,0}) \\ &+ (\log |f(w_{j,0} + \sqrt{-1})| - \log |f(w_{j,0})|) \operatorname{Im}(z - w_{j,0}). \end{aligned}$$

By the above observation,

$$\int_{\mathbb{C}} L_j(z) H_j(z) dz = 0$$

so we can write

$$X_{z_j, G_j} = \int_{\mathbb{C}} K_j(z) dz$$

where

$$K_j(z) := (\log |f(z)| - L_j(z)) H_j(z).$$

The point now is that with high probability, K_j has reasonably small L^2 norm:

Lemma 27. *For any constant $\delta > 0$, we have*

$$(35) \quad \|K_j\|_{L^2} \leq n^\delta$$

for all $1 \leq j \leq k$ with probability at least $1 - O(n^{-\delta+2/A}) - O(n^{-A+1})$.

Proof. We follow the proof of [58, Lemma 39]. Notice that by the union bound, it suffices to prove the claim for a single j . We split $K_j = \sum_{i: \zeta_i \neq \infty} K_{j,i}(z)$, where

$$K_{j,i}(z) := (\log |z - \zeta_i| - L_{j,i}(z)) H_j(z)$$

and $L_{j,i} : \mathbb{C} \rightarrow \mathbb{C}$ is the random linear function that equals $\log |z - \zeta_i|$ when $z = w_{j,0}, w_{j,0} + 1, w_{j,0} + \sqrt{-1}$. By the triangle inequality, we thus have

$$\|K_j\|_{L^2} \leq \sum_{i: \zeta_i \neq \infty} \|K_{j,i}\|_{L^2}.$$

By the non-clustering axiom, for each z_j and $r \geq 1$, one has

$$N_{B(z_j, r)} \ll n^{1/A} r^2$$

with probability at least $1 - O(n^{-A})$. By taking r of the form $r = 2^i, 0 \leq i \leq \log_2 \sqrt{n}$ and using the union bound, we can conclude that

$$(36) \quad N_{B(z_j, r)} \ll n^{1/A} r^2,$$

for all z_j and any $r \geq 1$, with probability at least $1 - O(n^{-A+1})$. (Notice that if $r \geq \sqrt{n}$ the bound holds trivially as there are at most n zeroes overall.)

We may now condition on the polynomial f and assume it obeys (36). The only remaining source of randomness are the $w_{j,0}$'s. In particular, the zeroes ζ_i are now deterministic. By Markov's inequality, it suffices to show that

$$(37) \quad \mathbf{E} \|K_j\|_{L^2} \ll n^{2/A}.$$

(The expectation is with respect to the $w_{j,0}$, of course.)

Recall that H_j is supported in $B(z_j, 10r_0)$. If $1 \leq i \leq n$ is such that $\zeta_i \in B(z_j, 20r_0)$, then a short computation (based on the square-integrability of the logarithm function) shows that the expected value of $\|K_{j,i}\|_{L^2}$ (averaged over all choices of $w_{j,0}$) is $O(1)$. By (36), there are $O(n^{1/A})$ indices i in this case. Thus, the total contribution from this case is $O(n^{1/A})$, which is acceptable.

Now, we consider the more delicate case when $\zeta_i \notin B(z_j, 20r_0)$. Let us write $z := w_{j,0} + x + \sqrt{-1}y$ and Taylor expand $\log |z - \zeta_i|$ around the point $w_{j,0} - \zeta_i$. Since we only care about $z \in B(z_j, 10r_0)$, we have $|x|, |y| = O(1)$ in this neighborhood and so

$$(38) \quad \begin{aligned} \log |z - \zeta_i| &= \log |x + \sqrt{-1}y + w_{j,0} - \zeta_i| \\ &= \log |w_{j,0} - \zeta_i| + \frac{\operatorname{Re}(w_{j,0} - \zeta_i)}{|w_{j,0} - \zeta_i|^2} x + \frac{\operatorname{Im}(w_{j,0} - \zeta_i)}{|w_{j,0} - \zeta_i|^2} y \\ &\quad + O\left(\frac{1}{|w_{j,0} - \zeta_i|^2}\right). \end{aligned}$$

Under the new notation, we can write $L_{j,i}$ as

$$(39) \quad \begin{aligned} L_{j,i}(z) &:= \log |w_{j,0} - \zeta_i| \\ &\quad + (\log |w_{j,0} + 1 - \zeta_i| - \log |w_{j,0} - \zeta_i|) x \\ &\quad + (\log |w_{j,0} + \sqrt{-1} - \zeta_i| - \log |w_{j,0} - \zeta_i|) y. \end{aligned}$$

The point here that this almost cancels out the linear part in (38). Indeed, by considering the Taylor expansion of

$$\log |w_{j,0} + 1 - \zeta_i| - \log |w_{j,0} - \zeta_i|$$

and

$$\log |w_{j,0} + \sqrt{-1} - \zeta_i| - \log |w_{j,0} - \zeta_i|$$

we easily see that the difference between $L_{j,i}$ and the linear part of (38) is at most $O(\frac{1}{|w_{j,0} - \zeta_i|^2})$. Thus, we conclude that the (conditional) expectation of $\|K_{j,i}\|_{L^2}$ (with respect to the random choice of $w_{j,0}$) is only $O(\frac{1}{|w_{j,0} - \zeta_i|^2})$. As $C \geq 1$, we can

replace it by a more convenient bound

$$O\left(\frac{1}{1 + |w_{j,0} - \zeta_i|^2}\right) = O\left(\frac{1}{1 + |\zeta_i - z_j|^2}\right),$$

which also holds for ζ_i close to z_j .

Summing over i , we see that the (conditional) expected value of $\|K_j\|_{L^2}$ is at most

$$O\left(\sum_{i:\zeta_i \neq \infty} \frac{1}{1 + |\zeta_i - z_j|^2}\right).$$

By (36), the number of ζ_i such that $2^l < |\zeta_i - z_j| \leq 2^{l+1}$ is $O(n^{1/A}4^l)$, for all $0 \leq l \leq \log_2 \sqrt{n}$. Furthermore, there are $O(n^{1/A})$ indices such that $|\zeta_i - z_j| \leq 1$, and there are trivially at most n indices for which $|\zeta_i - z_j| \geq \sqrt{n}$. Thus, the above sum is

$$O\left(\sum_{l=0}^{\log_2 \sqrt{n}} \frac{n^{1/A}4^l}{4^l} + \frac{n}{1 + (\sqrt{n})^2}\right) = O(n^{2/A}),$$

proves (37) and hence the lemma. \square

Let $\gamma_0, \gamma_1, \gamma_2$ be positive constants to be determined later (they will end up being constant multiples of c_0). Set $m := \lfloor n^{\gamma_0} \rfloor$, and for each $1 \leq j \leq k$ let $w_{j,1}, \dots, w_{j,m}$ be drawn uniformly at random from $B(z_j, 10r_0)$, independently of f and the $w_{j,0}$.

From Lemma 27, we have that with probability $1 - O(n^{-\gamma_1+2/A} + n^{-A+1})$, we have $\|K_j\|_{L^2} \leq n^{\gamma_1}$. If we condition on this event and use Lemma 26 (with respect to the sample points $w_{j,1}, \dots, w_{j,m}$), then we have the estimate

$$X_{z_j, F_j} = \frac{\pi(10r_0)^2}{m} \sum_{i=1}^m K_j(w_{j,i}) + O\left(\frac{1}{\sqrt{mn^{-\gamma_2}}} n^{\gamma_1/2}\right)$$

with probability $1 - O(n^{-\gamma_2})$.

Putting all this together, we conclude that

$$(40) \quad X_{z_j, F_j} = \frac{\pi(10r_0)^2}{m} \sum_{i=1}^m K_j(w_{j,i}) + O\left(n^{-\frac{\gamma_0 - \gamma_1 - \gamma_2}{2}}\right)$$

with probability at least $1 - O(n^{-\gamma_1+2/A} + n^{-\gamma_2} + n^{-A+1})$.

Notice that if (40) holds, then by (31) we have

$$P(X_{z_1, F_1}, \dots, X_{z_k, F_k}) = P \left(\left(\frac{\pi(10r_0)^2}{m} \sum_{i=1}^m K_j(w_{j,i}) \right)_{1 \leq j \leq k} \right) + O(n^{-\frac{\gamma_0 - (\gamma_1 + \gamma_2)}{2}} n^{\frac{c_0}{50a_0}}).$$

Now we can estimate the expectation of P as

$$\begin{aligned} \mathbf{E}P(X_{z_1, F_1}, \dots, X_{z_k, F_k}) &= \mathbf{E}P \left(\left(\frac{\pi(10r_0)^2}{m} \sum_{i=1}^m K_j(w_{j,i}) \right)_{1 \leq j \leq k} \right) \\ &+ O(n^{-\frac{\gamma_0 - (\gamma_1 + \gamma_2)}{2}} n^{\frac{c_0}{50a_0}}) \\ &+ O(n^{-\gamma_1 + 2/A} + n^{-\gamma_2} + n^{-A+1}) n^{\frac{c_0}{50a_0}}. \end{aligned}$$

If we set $\gamma_0 = 0.99c_0$, $\gamma_1 = \gamma_2 = 0.3c_0$ (say) and use (5), then it is easy to see that the two error terms on the RHS are of size $O(n^{-c_0/4})$. Furthermore, with these choices of $\gamma_0, \gamma_1, \gamma_2$, one sees (using (31)) that the statistic

$$\mathbf{E}P \left(\left(\frac{\pi(10r_0)^2}{m} \sum_{i=1}^m K_j(w_{j,i}) \right)_{1 \leq j \leq k} \right)$$

obeys the hypotheses required for the comparability axiom (iii) and is thus c_0 -insensitive, uniformly for all deterministic choices of $w_{j,0} \in B(z_j, 1)$ and $w_{j,l} \in B(z_j, C)$; $l = 1, \dots, m$. It follows that $\mathbf{E}P(X_{z_1, F_1}, \dots, X_{z_k, F_k})$ is $c_0/4$ -insensitive, concluding the proof of the theorem.

7. PROOF OF THE REPLACEMENT PRINCIPLE, REAL CASE

We now prove Theorem 10. Let k, l, C, c_0, r_0, a_0 be as in that theorem; all implied constants in the $O()$ notation will be allowed to depend on these parameters. Let A be defined by (8). Let n be a natural number, and let $x_1, \dots, x_k, z_1, \dots, z_l$, and $f = f_n, \tilde{f} = \tilde{f}_n$ obeying the hypotheses of the theorem. We may assume that n is sufficiently large depending on k, l, C, c_0, r_0, a_0 , as the claim is trivial otherwise.

As in the proof of Theorem 5, we may assume that f_n and \tilde{f}_n are almost surely non-vanishing, and that m and M are equal to 1.

Write $c_1 := \frac{c_0}{100(a_0+2)(k+l)}$. By the Fourier-analytic arguments of the previous section, it will suffice to show that the quantity

$$(41) \quad \int_{\mathbb{R}^k} \int_{C^l} G(y_1, \dots, y_k, w_1, \dots, w_l) \rho_f^{(k,l)}(x_1 + y_1, \dots, x_k + y_k, z_1 + w_1, \dots, z_l + w_l) dw_1 \dots dw_l dy_1 \dots dy_l$$

is c_1 -insensitive, whenever G takes the form

$$G(y_1, \dots, y_k, w_1, \dots, w_l) = F_1(y_1) \dots F_k(y_k) G_1(w_1) \dots G_l(w_l)$$

where $F_i : \mathbb{R} \rightarrow \mathbb{C}$ and $G_j : \mathbb{C} \rightarrow \mathbb{C}$ are smooth functions supported on $[-10r_0, 10r_0]$ and $B(0, 10r_0)$ respectively, such that

$$|\nabla^a F_i(x)|, |\nabla^a G_j(z)| \ll 1$$

for all $1 \leq i \leq k; 1 \leq j \leq l, 0 \leq a \leq a_0$, and $x \in \mathbb{R}, z \in \mathbb{C}$.

By repeating the inclusion-exclusion arguments in the complex case, by separating the spectrum into contributions from $\mathbb{R}, \mathbb{C}_+, \mathbb{C}_-$ (and increasing C as necessary), it suffices to show that the quantity

$$(42) \quad \mathbf{E} \left(\prod_{i=1}^{\tilde{k}} X_{\tilde{x}_i, F_i, \mathbb{R}} \right) \left(\prod_{j=1}^{\tilde{l}} X_{\tilde{z}_j, G_j, \mathbb{C}_+} \right) \left(\prod_{j'=1}^{\tilde{l}'} X_{\tilde{z}'_{j'}, G'_{j'}, \mathbb{C}_-} \right)$$

is c_1 -insensitive, where $\tilde{k} \leq k$ and $\tilde{l} + \tilde{l}' \leq l$, $\tilde{x}_1, \dots, \tilde{x}_{\tilde{k}} \in \{x_1, \dots, x_k\}$ and $\tilde{z}_1, \dots, \tilde{z}_{\tilde{l}}, \tilde{z}'_1, \dots, \tilde{z}'_{\tilde{l}'} \in \{z_1, \dots, z_l\}$, and

$$X_{x, F, \mathbb{R}} := \sum_{i: \zeta_i \in \mathbb{R}} F(\zeta_i - x)$$

and

$$X_{z, G, \mathbb{C}_\pm} := \sum_{i: \zeta_i \in \mathbb{C}_\pm} G(\zeta_i - z),$$

and the $F_i : \mathbb{R} \rightarrow \mathbb{C}, G_j : \mathbb{C} \rightarrow \mathbb{C}, G'_{j'} : \mathbb{C} \rightarrow \mathbb{C}$ are smooth functions supported on $B(0, 10r_0)$ obeying the bounds

$$|\nabla^a F_i(x)|, |\nabla^a G_j(z)|, |\nabla^a G'_{j'}(z)| \ll 1$$

for all $0 \leq a \leq a_0, x \in \mathbb{R}, z \in \mathbb{C}$, and ζ_i enumerates the zeroes of f .

As the zeroes of f are symmetric around the real axis (and $f(\bar{z}) = \overline{f(z)}$), one has

$$X_{z, G, \mathbb{C}_-} = X_{\bar{z}, \tilde{G}, \mathbb{C}_+}$$

where $\tilde{G}(z) := G(\bar{z})$. Thus we may concatenate the G_j with the $G'_{j'}$, and assume without loss of generality that $\tilde{l}' = 0$, at the cost of placing $\tilde{z}_1, \dots, \tilde{z}_{\tilde{l}}$ in $\{z_1, \dots, z_l, \bar{z}_1, \dots, \bar{z}_l\}$ rather than $\{z_1, \dots, z_l\}$. Thus we are now seeking to establish the c_1 -insensitivity of

$$(43) \quad \mathbf{E} \left(\prod_{i=1}^{\tilde{k}} X_{\tilde{x}_i, F_i, \mathbb{R}} \right) \left(\prod_{j=1}^{\tilde{l}} X_{\tilde{z}_j, G_j, \mathbb{C}_+} \right).$$

On the other hand, by repeating the remainder of the arguments for the complex case with essentially no changes, we can show that the quantity

$$(44) \quad \mathbf{E} \prod_{p=1}^m X_{z'_p, H_p}$$

is $c_0/4$ -insensitive for any $m \leq k + l$, any complex numbers z'_1, \dots, z'_m in

$$\bigcup_{i=1}^k B(x_i, 20r_0) \cup \bigcup_{j=1}^l B(z_j, 20r_0) \cup B(\bar{z}_j, 20r_0),$$

and any smooth $H_p : \mathbb{C} \rightarrow \mathbb{C}$ supported in $B(0, 20r_0)$ and obeying the bounds

$$|\nabla^a H_p(z)| \leq 1$$

for all $0 \leq a \leq a_0$ and $z \in \mathbb{C}$, where

$$X_{z,H} := \sum_{i:\zeta_i \neq \infty} H(\zeta_i - z).$$

(Here we use the trivial remark that $\log |f(\bar{z})| = \log |f(z)|$, so that one can freely replace $\{z_1, \dots, z_l\}$ by $\{z_1, \dots, z_l, \bar{z}_1, \dots, \bar{z}_l\}$ in the comparability axiom (iii).)

We are going to deduce the c_1 -insensitivity of (43) from the $c_0/4$ -insensitivity of (44). The main idea is to extend a real function to a complex one without changing the value of the expectation in (44) by too much. This will be the place where we make an essential use of the weak repulsion axiom (iv).

Notice that from the non-clustering axiom and (8) that

$$(45) \quad \mathbf{E}|X_{\tilde{x}_i, F_i, \mathbb{R}}|^{\tilde{k}+\tilde{l}}, \mathbf{E}|X_{\tilde{z}_j, G_j, \mathbb{C}_+}|^{\tilde{k}+\tilde{l}} \ll n^{(\tilde{k}+\tilde{l})/A} + n^{-A+\tilde{k}+\tilde{l}} \\ \ll n^{(k+l)/A},$$

for all $1 \leq i \leq \tilde{k}$ and $1 \leq j \leq \tilde{l}$.

In the next, and critical, lemma, we use the weak repulsion hypotheses (9), (10) to show that there are very few complex zeroes near the real line.

Lemma 28 (Level repulsion). *Let x be a real number in the set*

$$\bigcup_{i=1}^{\tilde{k}} B(\tilde{x}_i, 50r_0) \cup \bigcup_{j=1}^{\tilde{l}} B(\tilde{z}_j, 50r_0).$$

Let γ be a quantity with $\gamma \geq n^{-\frac{c_0}{20(a_0+2)}}$. Then we have

$$(46) \quad \mathbf{P}(N_{B(x, 10\gamma)} \geq 2) \ll \gamma^2$$

for both f and \tilde{f} .

Proof. We can assume that $\gamma \leq \frac{1}{100C}$ (say) as the claim is trivially true otherwise. Let H be a non-negative bump function supported on $B(x, 20\gamma)$ that equals one on $B(x, 10\gamma)$. Observe that $X_{x,H}^2 - X_{x,H^2}$ is always non-negative, and is at least 2 when $N_{B(x, 10\gamma)} \geq 2$. Thus by Markov's inequality, it suffices to show that

$$\mathbf{E}X_{x,H}^2 - X_{x,H^2} \ll \gamma^2$$

for both f and \tilde{f} . By construction we see that the first a_0 derivatives of H and H^2 are less than $n^{c_0/8}$, so by Theorem 5 we have

$$\mathbf{E}X_{x,H}^2(f) = \mathbf{E}X_{x,H}^2(\tilde{f}) + O(n^{-c_0/8})$$

and similarly for X_{x,H^2} . Since $O(n^{-c_0/8}) = O(\gamma^2)$, we conclude that it will suffice to establish the claim for \tilde{f} :

$$\mathbf{E}X_{x,H}^2(\tilde{f}) - X_{x,H^2}(\tilde{f}) \ll \gamma^2.$$

Next, observe that the expression $X_{x,H}^2(\tilde{f}) - X_{x,H^2}(\tilde{f})$ vanishes if \tilde{f} has at most one zero in $B(x, 20\gamma) \cap \mathbb{R}$ and no zeroes in $B(x, 20\gamma) \cap \mathbb{C}_+$. Thus it suffices to show that

$$(47) \quad \mathbf{P}(N_{B(x, 20\gamma) \cap \mathbb{C}_+}(\tilde{f}) \geq 1) \ll \gamma^2$$

and

$$(48) \quad \mathbf{P}(N_{B(x, 20\gamma) \cap \mathbb{R}}(\tilde{f}) \geq 2) \ll \gamma^2.$$

This will follow from the bounds

$$\int_{B(x, 20\gamma) \cap \mathbb{C}_+} \rho_{\tilde{f}}^{(0,1)}(z) dz \ll \gamma^2$$

and

$$\int_{B(x, 20\gamma) \cap \mathbb{R}} \int_{B(x, 20\gamma) \cap \mathbb{R}} \rho_{\tilde{f}}^{(2,0)}(y, y') dy dy' \ll \gamma^2$$

respectively; but these are immediate from (10), (9). \square

Remark 29. If one had some additional decay on the right-hand sides of (10), (9) as $|x - y|$ or $\text{Im}z$ went to zero, then one could improve the powers of γ in the bound (46). For instance, the level repulsion bounds provided by Lemma 17 allow one to improve the bound from $O(\gamma^2)$ to $O(\gamma^3)$. But for the argument here, any bound on this probability which decays as $O(\gamma^c)$ for some $c > 1$ will suffice.

Set

$$(49) \quad \gamma := n^{-\frac{c_0}{20(a_0+2)}},$$

and for any real number x , let $E_{x,\gamma}$ be the event that there are two zeroes ζ_i, ζ_j of f in the strip $S_{x,\gamma} := \{z \in B(x, 20r_0) : \text{Im}(z) \leq \gamma\}$ with $i \neq j$ such that $|\zeta_i - \zeta_j| \leq 2\gamma$. Then by Lemma 28 and a covering argument, we have $\mathbf{P}(E_{x,\gamma}) = O(\gamma)$ whenever $x \in \bigcup_{i=1}^k B(\tilde{x}_i, 10r_0) \cup \bigcup_{j=1}^l B(\tilde{z}_j, 10r_0)$.

From the symmetry of the spectrum, we observe that if $E_{x,\gamma}$ does not hold, then there cannot be any strictly complex zero ζ_i in the strip $S_{x,\gamma}$, since in that case $\bar{\zeta}_i$ would be distinct zero in the strip at a distance at most 2γ from $\lambda_i(M_n)$. In particular, we see that

$$(50) \quad \mathbf{P}(N_{S_{x,\gamma} \setminus [x-10r_0, x+10r_0]} = 0) = 1 - O(\gamma)$$

whenever $x \in \bigcup_{i=1}^k B(\tilde{x}_i, C) \cup \bigcup_{j=1}^l B(\tilde{z}_j, C)$.

We can use (50) to simplify the expression (43) in two ways. First we may “thicken” each factor $X_{\tilde{x}_i, F_i, \mathbb{R}}$ by replacing it with $X_{\tilde{x}_i, \tilde{F}_i}$, where $\tilde{F}_i : \mathbb{C} \rightarrow \mathbb{C}$ is a smooth extension of F_i that is supported on the strip $\{z : |\text{Im}(z)| \leq \gamma\}$, and more specifically

$$\tilde{F}_i(z) := F_i(\text{Re}(z))\varphi(\text{Im}(z)/\gamma)$$

where $\varphi : \mathbb{R} \rightarrow \mathbb{R}$ is a smooth function supported on $[-1, 1]$ that equals one at the origin. From (50) and the non-clustering axiom (iii), we see that

$$X_{\tilde{x}_i, F_i, \mathbb{R}} = X_{\tilde{x}_i, \tilde{F}_i} + D_i,$$

where

- $D_i = 0$ with probability $1 - O(\gamma)$;
- $|D_i| \ll n^{1/A}$ with probability $1 - n^{-A}$; and
- $|D_i| \ll n$ with probability 1.

In particular, from (8) we have

$$(51) \quad \mathbf{E}|X_{\tilde{x}_i, F_i, \mathbb{R}} - X_{\tilde{x}_i, \tilde{F}_i}|^{\tilde{k} + \tilde{l}} \ll \gamma n^{(\tilde{k} + \tilde{l})/A} + n^{\tilde{k} + \tilde{l} - A} \\ \ll \gamma n^{(k+l)/A}.$$

Furthermore, by performing a smooth truncation, we have the derivative bounds $\nabla^a \tilde{F}_i = O(\gamma^{-a_0})$ for $0 \leq a \leq a_0$.

In a similar vein, Similarly, we replace each of the G_j in (42) with a function \tilde{G}_j that vanishes on the half-plane $\{z - z_j : \text{Im}(z) \leq \gamma/2\}$; more explicitly we set

$$\tilde{G}_j(z) := G_j(z)\eta(\text{Im}(z + z_j)/\gamma)$$

where $\eta : \mathbb{R} \rightarrow \mathbb{R}$ is a smooth function supported on $[1/2, \infty)$ that equals one on $[1, \infty)$. Then we have

$$X_{\tilde{z}_j, G_j, \mathbb{C}_+} = X_{\tilde{z}_j, \tilde{G}_j} + H_j,$$

where H_j has properties similar to D_i . In particular we have

$$(52) \quad \mathbf{E}|X_{\tilde{z}_j, G_j, \mathbb{C}_+} - X_{\tilde{z}_j, \tilde{G}_j}|^{\tilde{k} + \tilde{l}} \ll \gamma n^{(k+l)/A}.$$

By telescoping the difference

$$\left(\prod_{i=1}^{\tilde{k}} X_{\tilde{x}_i, F_i, \mathbb{R}} \right) \left(\prod_{j=1}^{\tilde{l}} X_{\tilde{z}_j, G_j, \mathbb{C}_+} \right) - \left(\prod_{i=1}^{\tilde{k}} X_{\tilde{x}_i, \tilde{F}_i} \right) \left(\prod_{j=1}^{\tilde{l}} X_{\tilde{z}_j, \tilde{G}_j} \right)$$

and applying Hölder's inequality followed by (45), (51), (52), we see that

$$\mathbf{E} \left| \left(\prod_{i=1}^{\tilde{k}} X_{\tilde{x}_i, F_i, \mathbb{R}} \right) \left(\prod_{j=1}^{\tilde{l}} X_{\tilde{z}_j, G_j, \mathbb{C}_+} \right) - \left(\prod_{i=1}^{\tilde{k}} X_{\tilde{x}_i, \tilde{F}_i} \right) \left(\prod_{j=1}^{\tilde{l}} X_{\tilde{z}_j, \tilde{G}_j} \right) \right| \ll \gamma n^{(k+l)^2/A}.$$

From (8) and (49) we see that the right-hand side is $O(n^{-c_1})$. Thus, to show the c_1 -insensitivity of (43), it suffices to show that the quantity

$$\mathbf{E} \left(\prod_{i=1}^{\tilde{k}} X_{\tilde{x}_i, \tilde{F}_i} \right) \left(\prod_{j=1}^{\tilde{l}} X_{\tilde{z}_j, \tilde{G}_j} \right)$$

is c_1 -insensitive. However, from the $c_0/4$ -insensitivity of (44) and the derivative bounds on \tilde{F}_i, \tilde{G}_j (and homogeneity) we see that this quantity changes by at most

$$O\left(n^{-c_0/4}(\gamma^{-a_0})^{\tilde{k} + \tilde{l}}\right)$$

when one replaces f with \tilde{f} . From (49) this quantity is $O(n^{-c_1})$, and the claim follows.

8. NON-CLUSTERING VIA SHARP CONCENTRATION

In this section we prove Proposition 12. Let n, f_n, z_0, c, r, G be as in that proposition. Since the condition $\log |f(z)| = G(z) + O(n^{o(1)})$ can only hold when f is non-vanishing, we see from the concentration axiom that f is non-vanishing with overwhelming probability. We now condition to the event that f is non-vanishing, noting that this does not significantly impact the hypothesis or conclusion of the proposition, and so we assume henceforth that f is almost surely non-vanishing.

We first prove the upper bound
(53)

$$N_{B(z_0, r)}(f) \leq \frac{1}{2\pi} \int_{B(z_0, r)} \Delta G(z) dz + O(n^{o(1)}c^{-1}r) + O\left(\int_{B(z_0, r+c) \setminus B(z_0, r-c)} |\Delta G(z)| dz\right)$$

with overwhelming probability, and then explain how to modify the argument to obtain the matching lower bound at the end of this section.

Let φ_+ be a smooth function supported on $B(z_0, r+c)$ which equals 1 on $B(z_0, r)$, is bounded between 0 and 1 on the annulus $B(z_0, r+c) \setminus B(z_0, r)$, and has the second derivative bound $|\nabla^2 \varphi_+| = O(c^{-2})$ on this annulus; such a function is easily constructed since $0 < c \leq r$. Then

$$N_{B(z_0, r)}(f) \leq \sum_{i=1}^n \varphi_+(\zeta_i)$$

where ζ_1, \dots, ζ_n are the zeroes of f . Applying Green's theorem as in the proof of Theorem 5, we have the identity

$$\sum_{i=1}^n \varphi_+(\zeta_i) = \frac{1}{2\pi} \int_{\mathbb{C}} (\Delta \varphi_+(z)) \log |f_n(z)| dz.$$

Meanwhile, from another application of Green's theorem we have

$$\begin{aligned} \int_{\mathbb{C}} (\Delta \varphi_+(z)) G(z) dz &= \int_{\mathbb{C}} \varphi_+(z) \Delta G(z) dz \\ &= \int_{B(z_0, r)} \Delta G(z) dz + O\left(\int_{B(z_0, r+c) \setminus B(z_0, r)} |\Delta G(z)| dz\right). \end{aligned}$$

Set $H(z) := |\log |f_n(z)| - G(z)|$; by the triangle inequality, we thus have

$$N_{B(z_0, r)}(f) \leq \frac{1}{2\pi} \int_{B(z_0, r)} \Delta G(z) dz + O\left(\int_{B(z_0, r+c) \setminus B(z_0, r)} |\Delta G(z)| dz\right) + O\left(\int_{\mathbb{C}} |\Delta \varphi_+(z)| H(z) dz\right).$$

Since $\Delta \varphi_+$ is supported on $B(z_0, r+c) \setminus B(z_0, r)$ and has magnitude $O(c^{-2})$, it thus suffices by the triangle inequality to establish the upper bound

$$\int_{B(z_0, r+c) \setminus B(z_0, r)} H(z) dz \ll n^{o(1)} cr.$$

We first observe a crude polynomial bound

$$(54) \quad \int_{B(z_0, r+c) \setminus B(z_0, r)} |H(z)|^2 dz \ll n^{O(1)}$$

with overwhelming probability. To see this, first observe from (12) and the polynomial size bound on r (and hence on c) that

$$\int_{B(z_0, r+c) \setminus B(z_0, r)} |G(z)|^2 dz \ll n^{O(1)}$$

and so it suffices to show that

$$\int_{B(z_0, r+c) \setminus B(z_0, r)} |\log |f_n(z)||^2 dz \ll n^{O(1)}.$$

Let z_1 be any element of $B(z_0, r+c) \setminus B(z_0, r)$. By the hypotheses, we have $\log f_n(z_1) = O(n^{O(1)})$ with overwhelming probability, so it suffices by the triangle inequality again to show that

$$\int_{B(z_0, r+c) \setminus B(z_0, r)} |\log |f_n(z)| - \log |f_n(z_1)||^2 dz \ll n^{O(1)}.$$

But as

$$\log |f_n(z)| - \log |f_n(z_1)| = \sum_{1 \leq i \leq n: \zeta_i \neq \infty} \log |z - \zeta_i| - \log |z_1 - \zeta_i|,$$

the claim follows from yet another application of the triangle inequality, together with a direct calculation using the square-integrability of the log function $\log |z|$.

Now we apply Lemma 26. To use this lemma, let $m := n^A$ for some large fixed A to be chosen later, and let z_1, \dots, z_m be drawn uniformly at random from the annulus $B(z_0, r+c) \setminus B(z_0, r)$, independently of each other and of f (and hence of H). After temporarily conditioning H to be fixed, applying Lemma 26 to the normalised measure on the annulus $B(z_0, r+c) \setminus B(z_0, r)$, and then undoing the conditioning, we see from (54) that one has

$$\int_{B(z_0, r+c) \setminus B(z_0, r)} H(z) dz = |B(z_0, r+c) \setminus B(z_0, r)| \left(\frac{1}{m} \sum_{i=1}^m H(z_i) + O(n^{O(1)-A/4}) \right)$$

with probability $1 - O(n^{-A/2})$. On the other hand, we have $|B(z_0, r+c) \setminus B(z_0, r)| \ll cr$, and from the hypothesis of concentration of the log-magnitude and the union bound (and after temporarily conditioning the z_1, \dots, z_m to be fixed) we see that with overwhelming probability, one has $H(z_i) = O(n^{o(1)})$ for all $i = 1, \dots, m$. We conclude that

$$\int_{B(z_0, r+c) \setminus B(z_0, r)} H(z) dz \ll n^{o(1)} cr + O(n^{O(1)-A/4})$$

with probability $1 - O(n^{-A/2})$, and the claim then follows by diagonalising in A (and using the polynomial size of c, r).

This concludes the proof of the upper bound (53) with overwhelming probability. To prove the matching lower bound

$$N_{B(z_0, r)}(f) \geq \frac{1}{2\pi} \int_{B(z_0, r)} \Delta G(z) dz - O(n^{\rho(1)} c^{-1} r) - O\left(\int_{B(z_0, r+c) \setminus B(z_0, r-c)} |\Delta G(z)| dz\right),$$

one performs a similar argument but with φ_+ replaced by a test function φ_- that equals 1 on $B(z_0, r - c)$ and 0 outside of $B(z_0, r)$; we leave the details to the interested reader.

Remark 30. The above argument also establishes the following variant of Proposition 12; if one is willing to weaken the conclusion of Proposition 12 from holding with overwhelming probability to that of holding with probability $1 - O(n^{-A})$ for some fixed A , then one may also weaken the hypothesis in (i) from holding with overwhelming probability to that of holding with probability $1 - O(n^{-B})$ for some B depending on A .

9. ASSUMPTION VERIFICATION: PROOF OF LEMMA 13

We now prove Lemma 13. We first need an elementary lemma of Paley-Zygmund type.

Lemma 31 (Paley-Zygmund type lemma). *Let ξ be a random variable of mean zero and variance one, and obeying the bound $\mathbf{E}|\xi|^{2+\varepsilon} \leq C$ for some $\varepsilon, C > 0$. Then one can find $A > 1$ depending only on ε, C such that*

$$\mathbf{P}(A^{-1} \leq |\xi - \xi'| \leq A) \geq A^{-1}$$

where ξ' is an independent copy of ξ .

Proof. Let A be sufficiently large depending on ε, C, δ . From Chebyshev's inequality, we see that

$$\mathbf{P}(|\xi| \leq A/2), \mathbf{P}(|\xi'| \leq A/2) \geq 1 - 4/A^2$$

and hence by the triangle inequality

$$\mathbf{P}(|\xi - \xi'| \leq A) \geq 1 - 8/A^2.$$

It thus suffices (for A large enough) to show that

$$\mathbf{P}(|\xi - \xi'| \geq A^{-1}) \geq A^{-1}.$$

Suppose this were not the case, then

$$\mathbf{P}(|\xi - \xi'| \leq A^{-1}) \geq 1 - A^{-1}.$$

By conditioning on ξ' , there thus exists a complex number z_0 such that

$$\mathbf{P}(|\xi - z_0| \leq A^{-1}) \geq 1 - A^{-1}.$$

From Cauchy-Schwarz one has

$$\mathbf{E}\xi \leq \mathbf{P}(|\xi - z_0| \leq A^{-1})(z_0 + O(A^{-1})) + \mathbf{P}(|\xi - z_0| > A^{-1})^{1/2}(\mathbf{E}|\xi|^2)^{1/2};$$

since ξ has mean zero and variance one, we conclude that

$$z_0 = O(A^{-1/2})$$

and thus

$$\mathbf{P}(|\xi| \leq C_0 A^{-1/2}) \geq 1 - A^{-1}$$

for some absolute constant $C_0 > 0$. From Hölder's inequality, we thus have

$$\mathbf{E}|\xi|^2 \ll A^{-1} + \mathbf{P}(|\xi| \geq C_0 A^{-1/2})^{\varepsilon/(2+\varepsilon)} (\mathbf{E}|\xi|^{2+\varepsilon})^{2/(2+\varepsilon)};$$

since ξ has variance one and second moment bounded by C , we conclude that

$$1 \ll A^{-1} + A^{-\varepsilon/(2+\varepsilon)}$$

which leads to a contradiction if A is large enough. \square

Using this lemma, we can obtain the following result of ‘‘Littlewood-Offord’’ type.

Lemma 32 (Small ball probability for lacunary steps). *Let v_1, \dots, v_n be complex numbers, and suppose there is a subsequence $v_{i_1}, v_{i_2}, \dots, v_{i_m}$ with the property that*

$$|v_{i_j}| \geq 2|v_{i_{j+1}}|$$

for all $j = 1, \dots, m-1$. Let ξ_1, \dots, ξ_n be iid complex random variables whose common distribution ξ has mean zero and variance one, and obeys the bound $\mathbf{E}|\xi|^{2+\varepsilon} \leq C$ for some $\varepsilon, C > 0$. Then one has the non-concentration inequality

$$\sup_{z \in \mathbb{C}} \mathbf{P}(|\xi_1 v_1 + \dots + \xi_n v_n - z| \leq |v_{i_m}|) \leq C' \exp(-cm)$$

for some $C', c > 0$ depending only on ε, C .

Proof. In order to set up a conditioning argument later, we will introduce some additional sources of randomness. Let ξ'_1, \dots, ξ'_n be independent copies of ξ_1, \dots, ξ_n , let $\epsilon_1, \dots, \epsilon_n \in \{-1, 1\}$ be independent Bernoulli variables (independent of both ξ_i and ξ'_i , and let $\tilde{\xi}_i$ be the random variable that equals ξ_i when $\epsilon_i = +1$ and ξ'_i when $\epsilon_i = -1$. Then $\tilde{\xi}_1, \dots, \tilde{\xi}_n$ has the same joint distribution as ξ_1, \dots, ξ_n , so it suffices to obtain the bound

$$\sup_{z \in \mathbb{C}} \mathbf{P}(|\tilde{\xi}_1 v_1 + \dots + \tilde{\xi}_n v_n - z| \leq |v_{i_m}|) \leq C' \exp(-cm)$$

Next, let ξ' be an independent copy of ξ . By Lemma 31 we may find $A > 1$ depending only on ε, C such that

$$\mathbf{P}(A^{-1} < |\xi - \xi'| < A) > A^{-1}.$$

In particular

$$(55) \quad \mathbf{P}(A^{-1} < |\xi_i - \xi'_i| < A) > A^{-1}$$

for all $1 \leq i \leq n$.

Next, we may refine the sequence i_1, \dots, i_m to a subsequence $\tilde{i}_1, \dots, \tilde{i}_{\tilde{m}}$ with

$$\tilde{m} \gg m - O(1)$$

and

$$(56) \quad |v_{\tilde{i}_j}| \geq 4A^2 |v_{\tilde{i}_{j+1}}|$$

and

$$(57) \quad |v_{\tilde{i}_{\tilde{m}}}| \geq 4A |v_{i_m}|.$$

Let $J \subset \{1, \dots, \tilde{m}\}$ be the set of indices j for which

$$(58) \quad A \geq |\xi_{i_j} - \xi'_{i_j}| \geq A^{-1}.$$

From (55) and the Chernoff (or Hoeffding) inequality, one has

$$(59) \quad |J| \geq c\tilde{m}$$

with probability at least $1 - O(\exp(-c'\tilde{m})) = 1 - O(\exp(-c''m))$ for some quantities $c, c', c'' > 0$ depending only on A , where implied constants in the $O()$ notation may depend on A .

We now on the event that (59) occurs (we even fix all values of ξ, ξ'), and then further fix the signs ϵ_i for $i \notin J$. After this conditioning, the only remaining source of randomness comes from the signs ϵ_{i_j} for $j \in J$. We also fix the complex number z . Observe from (58) that each reversal of a sign ϵ_{i_j} alters the sum $\tilde{\xi}_1 v_1 + \dots + \tilde{\xi}_n v_n - z$ by a quantity of magnitude between $\varepsilon |v_{i_j}|$ and $A |v_{i_j}|$. Using (56), (57) and the triangle inequality, we conclude that if we modify a non-zero number of signs ϵ_{i_j} for $j \in J$, then the above sum is altered by more than $2|v_{i_m}|$. In particular, of the $2^{|J|}$ possible choices of these signs, at most one of them can lead to the sum having magnitude bounded by $|v_{i_m}|$. This gives an upper bound of $2^{-|J|} = O(\exp(-c'''m))$ for this event for some $c''' > 0$ depending only on A , and the claim follows. \square

We are now ready to prove Lemma 13.

Proof of Lemma 13. By Markov's inequality (or Chebyshev's inequality), we have with overwhelming probability

$$|f(z)| \leq V(z)^{1/2} \exp(\log^2 n) = V(z)^{1/2} \exp(n^{o(1)}).$$

Thus with overwhelming probability we have the upper bound

$$(60) \quad \log |f(z)| \leq \frac{1}{2} \log V(z) + n^{o(1)}.$$

Meanwhile, the lower bound

$$(61) \quad \log |f(z)| \geq \frac{1}{2} \log V(z) - n^{o(1)},$$

with overwhelming probability is immediate from Lemma 32. \square

10. ASSUMPTION VERIFICATION: PROOF OF THEOREM 16

In this section we establish Theorem 16. We begin by proving a variant of Theorem 16 in which the logarithms in (16) are removed:

Proposition 33. *Let $\xi, \tilde{\xi}, \varepsilon, M, n, f_{n,\xi}, f_{n,\tilde{\xi}}, c_0, \dots, c_n, k, z_1, \dots, z_k, \alpha_0, \alpha_1, V$ be as in Theorem 16. Assume that α_0 is sufficiently small depending on α_1, ε , and that $V(z_1), \dots, V(z_k) > 0$. Then for smooth function $H : \mathbb{C}^k \rightarrow \mathbb{C}$ obeying the derivative bounds*

$$(62) \quad |\nabla^a H(\zeta_1, \dots, \zeta_k)| \ll n^{\alpha_0}, 0 \leq a \leq 3,$$

we have

$$(63) \quad \begin{aligned} & |\mathbf{E}H(V(z_1)^{-1/2}f_n(z_1), \dots, V(z_k)^{-1/2}f_n(z_k)) \\ & - H(V(z_1)^{-1/2}\tilde{f}_n(z_1), \dots, V(z_k)^{-1/2}\tilde{f}_n(z_k))| \ll n^{-\alpha_0}, \end{aligned}$$

where the implied constants depend on $\varepsilon, M, \alpha_0, \alpha_1$.

Proof. We use the Lindeberg swapping argument. Let ξ_0, \dots, ξ_n be iid copies of ξ , and $\tilde{\xi}_0, \dots, \tilde{\xi}_n$ be iid copies of $\tilde{\xi}$ that are independent of ξ_0, \dots, ξ_n . We introduce the intermediate polynomials

$$f_{n,i_0}(z) := \sum_{0 \leq i < i_0} c_i \tilde{\xi}_i z^i + \sum_{i_0 \leq i \leq n} c_i \xi_i z^i$$

for $0 \leq i_0 \leq n+1$, and the random variables

$$Y_{j,i_0} := V(z_j)^{-1/2} f_{n,i_0}(z_j)$$

for $0 \leq i_0 \leq n+1$ and $1 \leq j \leq k$. We can then write (63) as

$$|\mathbf{E}H(Y_{1,0}, \dots, Y_{k,0}) - H(Y_{1,n+1}, \dots, Y_{k,n+1})| \ll n^{-\alpha_0},$$

and so by telescoping series it will suffice to show that

$$(64) \quad \sum_{i_0=1}^n |\mathbf{E}H(Y_{1,i_0}, \dots, Y_{k,i_0}) - H(Y_{1,i_0+1}, \dots, Y_{k,i_0+1})| \ll n^{-\alpha_0}.$$

Fix i_0 . We can then write

$$f_{n,i_0}(z) = \hat{f}_{n,i_0}(z) + c_{i_0} \xi_{i_0} z^{i_0}$$

and

$$f_{n,i_0+1}(z) = \hat{f}_{n,i_0}(z) + c_{i_0} \tilde{\xi}_{i_0} z^{i_0}$$

for any z , where

$$\hat{f}_{n,i_0}(z) := \sum_{0 \leq i < i_0} c_i \tilde{\xi}_i z^i + \sum_{i_0 < i \leq n} c_i \xi_i z^i.$$

In particular we have

$$Y_{j,i_0} = \tilde{Y}_{j,i_0} + a_{j,i_0} \xi_{i_0}$$

and

$$Y_{j,i_0+1} = \tilde{Y}_{j,i_0} + a_{j,i_0} \tilde{\xi}_{i_0}$$

where

$$\tilde{Y}_{j,i_0} := V(z_j)^{-1/2} \hat{f}_{n,i_0}(z_j)$$

and

$$(65) \quad a_{j,i_0} := \frac{c_{i_0} z_j^{i_0}}{V(z_j)^{1/2}}.$$

Now let us condition all the $\xi_i, \tilde{\xi}_i$ for $i \neq i_0$ to be fixed, leaving only ξ_{i_0} and $\tilde{\xi}_{i_0}$ as sources of randomness; in particular, the \tilde{Y}_{j,i_0} are now deterministic. We consider the conditional expectation

$$|\mathbf{E}_{\xi_{i_0}, \tilde{\xi}_{i_0}} H(Y_{1,i_0}, \dots, Y_{k,i_0}) - H(Y_{1,i_0+1}, \dots, Y_{k,i_0+1})|.$$

We can write

$$H(Y_{1,i_0}, \dots, Y_{k,i_0}) = H(\tilde{Y}_{1,i_0} + a_{1,i_0}\xi_{i_0}, \dots, \tilde{Y}_{k,i_0} + a_{k,i_0}\xi_{i_0}).$$

From (62), the bound $k \leq n^{\alpha_0}$, and Taylor expansion with remainder, we have

$$H(Y_{1,i_0}, \dots, Y_{k,i_0}) = H_{0,0} + H_{1,0}\text{Re}\xi_{i_0} + H_{0,1}\text{Im}\xi_{i_0} + O(a_{i_0}^2 n^{4\alpha_0} |\xi_{i_0}|^2)$$

and

(66)

$$\begin{aligned} H(Y_{1,i_0}, \dots, Y_{k,i_0}) &= H_{0,0} + H_{1,0}\text{Re}\xi_{i_0} + H_{0,1}\text{Im}\xi_{i_0} \\ &\quad + H_{2,0}(\text{Re}\xi_{i_0})^2 + H_{1,1}\text{Re}\xi_{i_0}\text{Im}\xi_{i_0} + H_{0,2}(\text{Im}\xi_{i_0})^2 + O(a_{i_0}^3 n^{4\alpha_0} |\xi_{i_0}|^3) \end{aligned}$$

(say), where

$$H_{r,s} := \frac{1}{r!s!} \frac{\partial^{r+s}}{(\partial x)^r (\partial y)^s} H(\tilde{Y}_{1,i_0} + a_{1,i_0}\xi_{i_0}, \dots, \tilde{Y}_{k,i_0} + a_{k,i_0}(x + \sqrt{-1}y))|_{x=y=0}$$

and

$$(67) \quad a_{i_0} := \left(\sum_{j=1}^k |a_{j,i_0}|^2 \right)^{1/2}.$$

One can verify that

$$H_{2,0}, H_{1,1}, H_{0,2} = O(a_{i_0}^2 n^{4\alpha_0} |\xi_{i_0}|^2)$$

and so the error term in (66) is both $O(a_{i_0}^2 n^{O(\alpha_0)} |\xi_{i_0}|^2)$ and $O(a_{i_0}^3 n^{O(\alpha_0)} |\xi_{i_0}|^3)$. Interpolating, we see that

$$\begin{aligned} H(Y_{1,i_0}, \dots, Y_{k,i_0}) &= H_{0,0} + H_{1,0}\text{Re}\xi_{i_0} + H_{0,1}\text{Im}\xi_{i_0} + H_{2,0}(\text{Re}\xi_{i_0})^2 + H_{1,1}\text{Re}\xi_{i_0}\text{Im}\xi_{i_0} \\ &\quad + H_{0,2}(\text{Im}\xi_{i_0})^2 + O(a_{i_0}^{2+\varepsilon} n^{4\alpha_0} |\xi_{i_0}|^{2+\varepsilon}). \end{aligned}$$

Similarly for $H(Y_{1,i_0+1}, \dots, Y_{k,i_0+1})$ and $\tilde{\xi}_{i_0}$. Taking expectations in $\xi_{i_0}, \tilde{\xi}_{i_0}$ and using the bounded moment assumption, and the fact that $\xi, \tilde{\xi}$ match moments to second order, we conclude that

$$|\mathbf{E}_{\xi_{i_0}, \tilde{\xi}_{i_0}} H(Y_{1,i_0}, \dots, Y_{k,i_0}) - H(Y_{1,i_0+1}, \dots, Y_{k,i_0+1})| \ll a_{i_0}^{2+\varepsilon} n^{4\alpha_0}.$$

Integrating out all the other variables, we see that we may bound the left-hand side of (64) by

$$n^{4\alpha_0} \sum_{i_0=1}^n a_{i_0}^{2+\varepsilon}.$$

From (14), (65), (67) we have

$$\sum_{i_0=1}^n a_{i_0}^2 = k \leq n^{\alpha_0}$$

and from (15) we have

$$\sup_{1 \leq i_0 \leq n} a_{i_0} \leq kn^{-\alpha_1}$$

and the claim (64) now follows if α_0 is sufficiently small depending on ε, α_1 . \square

Now we can reinstate the logarithms and complete the proof of Theorem 16. Let the notation and hypotheses be as in that theorem. If one of the $V(z_j)$ vanishes then $f(z_j)$ and $\tilde{f}(z_j)$ are almost surely zero and the claim is vacuously true thanks to our conventions, so we may assume that $V(z_j) > 0$ for all j .

As the conclusions of the theorem are transitive in $\xi, \tilde{\xi}$, we may assume without loss of generality that one of these distributions, say ξ , has a gaussian distribution (whose real and imaginary part have the same covariance matrix as that of ξ , in particular having mean zero and variance one)

Using a smooth partition of unity, we can split $G = G_1 + G_2$, where G_1 is supported on those ζ_1, \dots, ζ_k with $\inf_{1 \leq i \leq k} \zeta_i \leq -50\alpha_0 \log n$, and G_2 is supported on those ζ_1, \dots, ζ_k with $\inf_{1 \leq i \leq k} \zeta_i \geq -50\alpha_0 \log n - 1$, and with the bounds

$$|\nabla^a G_i(x_1, \dots, x_k)| \ll n^{5\alpha_0}$$

(say) for $0 \leq a \leq 3$, $i = 1, 2$, and all $x_1, \dots, x_k \in \mathbb{R}$. (The constants 5, 10, 50, 100 are rather arbitrary and generous.)

We first show that the contribution coming from G_1 is negligible. Indeed,

$$|\mathbf{E}G_1(\log |Y_1|, \dots, \log |Y_k|)| \leq \mathbf{E}H_1(Y_1, \dots, Y_k)$$

for some smooth function $H_1 : \mathbb{C}^k \rightarrow \mathbb{R}^+$ supported on the region $\{(\zeta_1, \dots, \zeta_k) \in \mathbb{C}^k : \inf_{1 \leq i \leq k} |\zeta_i| \ll n^{-50\alpha_0}\}$ obeying the derivative bounds (62) (but with α_0 replaced by a constant multiple of itself). By Proposition 33 (and reducing α_0 as necessary), we have

$$\mathbf{E}H_1(Y_1, \dots, Y_k) \leq \mathbf{E}H_1(\tilde{Y}_1, \dots, \tilde{Y}_k) + O(n^{-\alpha_0}).$$

But as the $\tilde{Y}_1, \dots, \tilde{Y}_k$ are independent gaussian with mean zero and variance one, the support of H_1 has measure $O(kn^{-50\alpha_0}) = O(n^{-49\alpha_0})$ with respect to the product gaussian measure (regardless of the structure of the covariance matrix). Furthermore, by assumption $|H_1| \leq n^{10\alpha_0}$. This implies

$$|\mathbf{E}G_1(\log |Y_1|, \dots, \log |Y_k|)| = O(n^{-39\alpha_0}) = o(n^{-\alpha_0}).$$

With \tilde{Y}_i , we can argue similarly, and without using Proposition 33.

To conclude the proof, it suffices to show that

$$\mathbf{E}G_2(\log |Y_1|, \dots, \log |Y_k|) - G_2(\log |\tilde{Y}_1|, \dots, \log |\tilde{Y}_k|) = O(n^{-\alpha_0}).$$

We can rewrite this as

$$\mathbf{E}H_2(Y_1, \dots, Y_k) - H_2(\tilde{Y}_1, \dots, \tilde{Y}_k) = O(n^{-\alpha_0}),$$

where

$$H_2(\zeta_1, \dots, \zeta_k) := G_2(\log |\zeta_1|, \dots, \log |\zeta_k|).$$

From the derivative and support hypotheses on G_2 , we see from the chain rule that H_2 obeys the derivative bounds (62) (but with α_0 replaced by a constant multiple of itself), and the claim now follows from Proposition 33 (again reducing α_0 as necessary).

11. ASSUMPTION VERIFICATION: REPULSION BOUNDS

In this section we prove Lemma 17. Let $n, f, c_0, \dots, c_n, R, v, x, \delta$ be as in that lemma. Our primary tool will be the following (vector-valued) version of the well-known Kac-Rice formula [26], [28], [47]:

Lemma 34 (Kac-Rice formula). *Let f be as above. Let $k, l, n \geq 0$ be integers with $k + 2l \leq n$. Let $x_1, \dots, x_k \in \mathbb{R}$ be distinct real numbers, and $z_1, \dots, z_l \in \mathbb{C}_+$ be distinct complex numbers in the upper half-plane. Then we have*

$$\begin{aligned} & \rho^{(k,l)}(x_1, \dots, x_k, z_1, \dots, z_l) = \\ & p_{\mathbb{R}^k \times \mathbb{C}^l}((f(x_1), \dots, f(x_k), f(z_1), \dots, f(z_l)) = (0, \dots, 0)) \\ & \quad \times \mathbf{E}(|f'(x_1)| \dots |f'(x_k)| |f'(z_1)|^2 \dots |f'(z_l)|^2 | (f(x_1), \dots, f(x_k), f(z_1), \dots, f(z_l)) = (0, \dots, 0)). \end{aligned}$$

where $p_{\mathbb{R}^k \times \mathbb{C}^l}((f(x_1), \dots, f(x_k), f(z_1), \dots, f(z_l)) = (0, \dots, 0))$ denotes the probability density function of the random variable $(f(x_1), \dots, f(x_k), f(z_1), \dots, f(z_l))$ (viewed as taking values in $\mathbb{R}^k \times \mathbb{C}^l$) at the origin $(0, \dots, 0)$.

Specialising the above lemma to the cases $(k, l) = (2, 0), (0, 1)$ and $n \geq 2$, we see that

$$(68) \quad \begin{aligned} & \rho^{(2,0)}(x, x + \delta) = p_{\mathbb{R}^2}((f(x), f(x + \delta)) = (0, 0)) \\ & \quad \times \mathbf{E}(|f'(x)| |f'(x + \delta)| | (f(x), f(x + \delta)) = (0, 0)) \end{aligned}$$

and

$$(69) \quad \begin{aligned} & \rho^{(0,1)}(x + \sqrt{-1}\delta) = p_{\mathbb{C}}(f(x + \sqrt{-1}\delta) = 0) \\ & \quad \times \mathbf{E}(|f'(x + \sqrt{-1}\delta)|^2 | f(x + \sqrt{-1}\delta) = 0). \end{aligned}$$

Observe that random variables such as

$$f(x), f(y), \operatorname{Re}f(z), \operatorname{Im}f(z), f'(x), f'(y), \operatorname{Re}f'(z), \operatorname{Im}f'(z)$$

can be written in the form $X \cdot v$, where $X \in \mathbb{R}^{n+1}$ is the random real gaussian vector $X := (\xi_0, \dots, \xi_n)$, and $v \in \mathbb{R}^{n+1}$ is a deterministic vector depending on x, y , or z . For computing the quantities in (68), (69), we observe the following identities:

Lemma 35 (Gaussian identities). *Let $1 \leq m \leq n$, and let v_1, \dots, v_m be linearly independent (deterministic) vectors in \mathbb{R}^{n+1} , and let $X \in \mathbb{R}^{n+1}$ be a random real gaussian vector. Then*

$$p_{\mathbb{R}^m}((X \cdot v_1, \dots, X \cdot v_m) = (0, \dots, 0)) = (2\pi)^{-m/2} |v_1 \wedge \dots \wedge v_m|^{-1}.$$

Furthermore, if v is another vector in \mathbb{R}^{n+1} , then

$$(70) \quad \mathbf{E}(|X \cdot v|^2 | (X \cdot v_1, \dots, X \cdot v_m) = (0, \dots, 0)) = \operatorname{dist}(v, \operatorname{span}(v_1, \dots, v_m))^2.$$

and similarly

$$(71) \quad \mathbf{E}(|X \cdot v| | (X \cdot v_1, \dots, X \cdot v_m) = (0, \dots, 0)) = \sqrt{\frac{2}{\pi}} \operatorname{dist}(v, \operatorname{span}(v_1, \dots, v_m)).$$

Proof. We can assume that v does not belong to the span of v_1, \dots, v_m , as otherwise both sides of (70) and (71) are zero. By applying an invertible linear transformation to the v_1, \dots, v_m , we may reduce to the case when the v_1, \dots, v_m are an orthonormal system. As the distribution of the gaussian random vector X is rotation invariant, we may then assume without loss of generality that v_1, \dots, v_m are the first m vectors e_1, \dots, e_m of the standard basis e_1, \dots, e_{n+1} . Since we may subtract any linear combination of v_1, \dots, v_m from v without affecting either side of (70), we may assume without loss of generality that v is orthogonal to e_1, \dots, e_m ; by rotating and rescaling we may then normalize $v = e_{m+1}$. The claims then follow by direct computation. \square

11.1. **Estimating $\rho^{(2,0)}(x, y)$.** We apply this lemma to obtain the bound (19). By Cauchy-Schwarz, we have

$$\begin{aligned} \mathbf{E}(|f'(x)| |f'(x + \delta)| | (f(x), f(x + \delta)) = (0, 0)) &\leq \mathbf{E}(|f'(x)|^2 | (f(x), f(x + \delta)) = (0, 0))^{1/2} \\ &\quad \times \mathbf{E}(|f'(x + \delta)|^2 | (f(x), f(x + \delta)) = (0, 0))^{1/2} \end{aligned}$$

and hence by (68) and Lemma 35 we have

$$(72) \quad \rho^{(2,0)}(x, x + \delta) \ll |\mathbf{v}_x \wedge \mathbf{v}_{x+\delta}|^{-1} \operatorname{dist}(\mathbf{w}_x, \operatorname{span}(\mathbf{v}_x, \mathbf{v}_{x+\delta})) \operatorname{dist}(\mathbf{w}_{x+\delta}, \operatorname{span}(\mathbf{v}_x, \mathbf{v}_{x+\delta}))$$

where $\mathbf{v}_x, \mathbf{v}_{x+\delta}, \mathbf{w}_x, \mathbf{w}_{x+\delta}$ are the vectors

$$(73) \quad \mathbf{v}_x := (c_i x^i)_{i=0}^n$$

$$(74) \quad \mathbf{v}_{x+\delta} := (c_i (x + \delta)^i)_{i=0}^n$$

$$(75) \quad \mathbf{w}_x := (i c_i x^{i-1})_{i=0}^n$$

$$(76) \quad \mathbf{w}_{x+\delta} := (i c_i (x + \delta)^{i-1})_{i=0}^n.$$

Note from the quotient rule and the hypotheses on v, R in Lemma 17 that

$$\begin{aligned} \mathbf{v}_x &= \frac{v(x)}{R(x)} \\ \mathbf{v}_{x+\delta} &= \frac{v(x + \delta)}{R(x + \delta)} \\ \mathbf{w}_x &= \frac{1}{R(x)} v_x(x) - \frac{R'(x)}{R(x)} \mathbf{v}_x \\ \mathbf{w}_{x+\delta} &= \frac{1}{R(x + \delta)} v'(x + \delta) - \frac{R'(x + \delta)}{R(x + \delta)} \mathbf{v}_{x+\delta} \end{aligned}$$

where v' is the complex derivative of the holomorphic function v , and similarly for R . One can thus write the right-hand side of (72) as

$$|v(x) \wedge v(x + \delta)|^{-1} \operatorname{dist}(v'(x), \operatorname{span}(v(x), v(x + \delta))) \operatorname{dist}(v_x(x + \delta), \operatorname{span}(v(x), v(x + \delta))).$$

To obtain the desired bound (19), it will thus suffice to establish the bounds

$$(77) \quad |v(x) \wedge v(x + \delta)| \gg \delta$$

$$(78) \quad \text{dist}(v'(x), \text{span}(v(x), v(x + \delta))) \ll \delta$$

$$(79) \quad \text{dist}(v'(x + \delta), \text{span}(v(x), v(x + \delta))) \ll \delta.$$

From (17) and the Cauchy integral formula we have the bounds

$$(80) \quad \left| \frac{d^k}{dz^k} v(z) \right| \ll 1$$

for all $z \in B(x, \delta)$ and $k = 0, 1, 2$ if δ is sufficiently small depending on r_0 (recall that implied constants are allowed to depend on r_0, C). Using this and Taylor's theorem with remainder, we see that

$$|v(x + \delta) - v(x) - \delta v'(x)| \ll \delta^2$$

and thus

$$|v(x) \wedge v(x + \delta) - \delta v(x) \wedge v'(x)| \ll \delta^2$$

which together with (18) gives (77) for δ sufficiently small. Also, from (80) and Taylor's theorem with remainder we have

$$v(x + \delta) = v(x) + \delta v'(x) + O(\delta^2)$$

and so

$$v'(x) = \frac{1}{\delta} v(x + \delta) - \frac{1}{\delta} v(x) + O(\delta)$$

and (78) follows. A similar argument gives (79), and (19) follows.

11.2. Estimating $\rho^{(0,1)}(z)$. We now establish the bound (20).

Applying (69), splitting into real and imaginary parts, and then using Lemma 35, we see that

$$(81) \quad \rho^{(0,1)}(z) \ll |\text{Re} \mathbf{v}_{x+\sqrt{-1}\delta} \wedge \text{Im} \mathbf{v}_{x+\sqrt{-1}\delta}|^{-1} \\ (\text{dist}(\text{Re} \mathbf{w}_{x+\sqrt{-1}\delta}, \text{span}(\text{Re} \mathbf{v}_{x+\sqrt{-1}\delta}, \text{Im} \mathbf{v}_{x+\sqrt{-1}\delta}))^2 \\ + \text{dist}(\text{Im} \mathbf{w}_{x+\sqrt{-1}\delta}, \text{span}(\text{Re} \mathbf{v}_{x+\sqrt{-1}\delta}, \text{Im} \mathbf{v}_{x+\sqrt{-1}\delta}))^2)$$

where

$$(82) \quad \mathbf{v}_{x+\sqrt{-1}\delta} := (c_i(x + \sqrt{-1}\delta))_{i=0}^n$$

$$(83) \quad \mathbf{w}_{x+\sqrt{-1}\delta} := (ic_i(x + \sqrt{-1}\delta)^{i-1})_{i=0}^n,$$

distances and span are computed over the reals rather than over the complex numbers, and we adopt the convention that the real or imaginary part of a complex vector is computed by taking the real or imaginary part of each of its coefficients separately. Again, the quotient rule gives

$$\mathbf{v}_{x+\sqrt{-1}\delta} = v(x + \sqrt{-1}\delta)/R(x + \sqrt{-1}\delta) \\ \mathbf{w}_{x+\sqrt{-1}\delta} = \frac{1}{R(x + \sqrt{-1}\delta)} v'(x + \sqrt{-1}\delta) - \frac{R'(x + \sqrt{-1}\delta)}{R(x + \sqrt{-1}\delta)} \mathbf{v}_{x+\sqrt{-1}\delta}.$$

Thus we may rewrite the right-hand side of (72) as

$$\begin{aligned} & |\operatorname{Re}v(x + \sqrt{-1}\delta) \wedge \operatorname{Im}v(x + \sqrt{-1}\delta)|^{-1} \\ & \quad \times (\operatorname{dist}(\operatorname{Re}v'(x + \sqrt{-1}\delta), \operatorname{span}(\operatorname{Re}v(x + \sqrt{-1}\delta), \operatorname{Im}v(x + \sqrt{-1}\delta)))^2 \\ & \quad + \operatorname{dist}(\operatorname{Im}v'(x + \sqrt{-1}\delta), \operatorname{span}(\operatorname{Re}v(x + \sqrt{-1}\delta), \operatorname{Im}v(x + \sqrt{-1}\delta)))^2). \end{aligned}$$

Since

$$\operatorname{Re}v(x + \sqrt{-1}\delta) = \frac{v(x + \sqrt{-1}\delta) + v(x - \sqrt{-1}\delta)}{2}$$

and

$$\operatorname{Im}v(x + \sqrt{-1}\delta) = \frac{v(x + \sqrt{-1}\delta) - v(x - \sqrt{-1}\delta)}{2\sqrt{-1}}$$

It thus suffices to establish the bounds

$$(84) \quad |v(x + \sqrt{-1}\delta) \wedge \operatorname{Im}v(x - \sqrt{-1}\delta)| \gg \delta$$

$$(85) \quad \operatorname{dist}(v_y(x + \sqrt{-1}\delta), \operatorname{span}(v(x + \sqrt{-1}\delta), v(x - \sqrt{-1}\delta))) \ll \delta$$

$$(86) \quad \operatorname{dist}(v_y(x - \sqrt{-1}\delta), \operatorname{span}(v(x + \sqrt{-1}\delta), v(x - \sqrt{-1}\delta))) \ll \delta,$$

where the notions of distance and span are now over the complex numbers rather than the reals. But these bounds can be achieved by adapting the proofs of (77), (78), (79) (inserting factors of $\sqrt{-1}$ at various stages of the argument; we leave the details to the interested reader.

12. UNIVERSALITY FOR FLAT POLYNOMIALS

In this section we establish our main results for flat polynomials, namely Theorems 18, 19, 20. This will largely be accomplished by invoking the results obtained in previous sections.

The first basic result we will need is a concentration result for the log-magnitude $\log|f(z)|$ of a flat polynomial:

Lemma 36 (Concentration for log-magnitude). *Let $C, \varepsilon > 0$ be constants, let n be a natural number, and let z be a complex number with*

$$n^\varepsilon \leq |z| \leq Cn^{1/2}.$$

Let $f = f_{n,\xi}$ be a flat polynomial whose atom distribution ξ has mean zero and variance one with $\mathbf{E}|\xi|^{2+\varepsilon} \leq C$. Then with overwhelming probability, one has

$$\log|f(z)| = \frac{1}{2}|z|^2 + O(n^{o(1)})$$

when $n^\varepsilon \leq |z| \leq n^{1/2}$, and

$$\log|f(z)| = n \log|z| - \frac{1}{2}n \log n + \frac{1}{2}n + O(n^{o(1)})$$

when $n^{1/2} \leq |z| \leq Cn^{1/2}$. The implied constants in the asymptotic notation can depend on C, ε .

Note that some lower bound on $|z|$ is necessary here, because $\log |f(0)|$ has the distribution of $\log |\xi|$ and this does not need to concentrate at the origin if ξ is discrete (and in particular, ξ could equal zero with non-zero probability).

Proof. We first compute the quantity $V(z)$ from (14). In the flat case we have

$$V(z) = \sum_{i=0}^n \frac{|z|^{2i}}{i!}.$$

A standard application of Taylor expansion or Stirling approximation (see e.g. [58, Lemma 64]) shows that

$$\log V(z) = |z|^2 + O(n^{o(1)})$$

for $|z| \leq n^{1/2}$ and

$$\log V(z) = 2n \log |z| - n \log n + n + O(n^{o(1)})$$

for $|z| \geq n^{1/2}$.

We now apply Lemma 13. Comparing that lemma with the current situation, we see that it will suffice to find indices $i_1, \dots, i_m \in \{0, \dots, n\}$ for some $m = \omega(\log n)$ such that we have the lacunarity property

$$|z^{i_j}/\sqrt{i_j!}| \geq 2|z^{i_{j+1}}/\sqrt{i_{j+1}!}|$$

for all $1 \leq j < m$, and the lower bound

$$|z^{i_m}/\sqrt{i_m!}| \geq V(z)^{1/2} \exp(-n^{o(1)}).$$

Observe that the sequence $i \mapsto |z^i/\sqrt{i!}|$ is increasing for $i < |z|^2$ and decreasing for $i > |z|^2$, with its largest value being at least $(V(z)/(n+1))^{1/2}$. Also, the ratio between adjacent elements of this sequence is $O(1)$ when i is comparable to $|z|^2$. If $n^\varepsilon \leq |z| \leq \sqrt{n}$, then the desired indices i_0, \dots, i_m can then be obtained by applying Lemma 14 to the (reversal of the) subsequence of the $|z^i/\sqrt{i!}|$ for which $|z|^2/2 \leq i \leq |z|^2$ (note that the ratio between the largest and smallest elements of this sequence is at least $\exp(c|z|^2) \geq \exp(cn^{2\varepsilon})$ for some $c > 0$). Similarly, if $\sqrt{n} \leq |z| \leq C\sqrt{n}$, the claim follows by applying Lemma 14 to the (reversal of the) subsequence of the $|z^i/\sqrt{i!}|$ for which $n/2 \leq i \leq n$. \square

Note that if we let $G : \mathbb{C} \rightarrow \mathbb{R}$ be the function defined by

$$G(z) := \frac{1}{2}|z|^2$$

for $|z| \leq \sqrt{n}$ and

$$G(z) := n \log |z| - \frac{1}{2}n \log n + \frac{1}{2}n$$

then a short computation shows that

$$\Delta G(z) = 21_{B(0, \sqrt{n})}(z)$$

in the sense of distributions. Applying Proposition 12 (after performing an infinitesimal regularization of G at the boundary of $B(0, \sqrt{n})$ to make it smooth), we conclude that for any $n^{-C} \leq c \leq r \leq C\sqrt{n}/3$ and $z_0 \in B(0, C\sqrt{n}/3)$ with the

property that $B(z_0, r+c) \setminus B(z_0, r-c)$ is disjoint from $B(0, n^\varepsilon)$, with overwhelming probability f is non-vanishing and obeys the *local circular law* (87)

$$N_{B(z_0, r)}(f) = \int_{B(z_0, r)} \frac{1}{\pi} 1_{B(0, \sqrt{n})}(z) dz + O(n^{o(1)} c^{-1} r) + O\left(\int_{B(z_0, r+c) \setminus B(z_0, r-c)} 1_{B(0, \sqrt{n})}(z) dz\right).$$

This already gives axiom (i) for Theorems 5, 10. The formula (87) leads to two further consequences of importance to us. First, for any $z_0 \in B(0, C\sqrt{n}/3)$ and $r \geq 1$, one has with overwhelming probability that

$$(88) \quad N_{B(z_0, r)} \ll n^{o(1)} r^2.$$

Indeed, this claim is trivial for $r \geq \sqrt{n}/3$ (say) from the trivial bound $N_{B(z_0, r)} \leq n$, and for $1 \leq r \leq \sqrt{n}/3$ the claim follows from (87) with $c = 1$ and bounding $O(\int_{B(z_0, r+1) \setminus B(z_0, r-1)} 1_{B(0, \sqrt{n})}(z) dz)$ by $O(r)$, after enlarging r as necessary in order to avoid the ball $B(0, n^\varepsilon)$. (This incurs a loss of $n^{\varepsilon+o(1)}$ rather than $n^{o(1)}$, but the gain of $n^{o(1)}$ can then be recovered by diagonalizing in ε .) In particular, this gives axiom (ii) for Theorems 5, 10.

We will also apply (87) in the case when $z_0 = 0$, $r = \sqrt{n} + n^{1/2-\varepsilon}$, and $c := n^{1/2-\varepsilon}$, leading to the bound

$$(89) \quad N_{B(0, \sqrt{n} + n^{1/2-\varepsilon})} = n + O(n^{\varepsilon+o(1)})$$

with overwhelming probability. In other words, with overwhelming probability, all but $O(n^{\varepsilon+o(1)})$ of the zeroes of f lie inside the disk $B(0, \sqrt{n} + n^{1/2-\varepsilon})$.

Next, we establish the comparability of log magnitudes required for axioms (iii) of Theorems 5, 10.

Proposition 37 (Comparability of log-magnitudes). *Let $C, \varepsilon > 0$ be constants, and let $c_0 > 0$ be sufficiently small depending on ε . Let n be a natural number, let $1 \leq k \leq n^{c_0}$ be another natural number, and let z_1, \dots, z_k be complex numbers such that*

$$n^\varepsilon \leq |z| \leq n^{1/2} + C.$$

Let $f_{n, \xi}, \tilde{f}_{n, \tilde{\xi}}$ be flat polynomials whose atom distributions $\xi, \tilde{\xi}$ have mean zero and variance one matching moments to second order with $\mathbf{E}|\xi|^{2+\varepsilon}, |\tilde{\xi}|^{2+\varepsilon} \leq C$. Let $F : \mathbb{C}^k \rightarrow \mathbb{C}$ be a smooth function obeying the bounds

$$|\nabla^a F(z)| \leq C$$

for all $0 \leq a \leq 3$. Then, if c_0 is sufficiently small, one has

$$\mathbf{E}\left(F(\log |f(z'_1)|, \dots, \log |f(z'_{k'})|) - F(\log |\tilde{f}(z'_1)|, \dots, \log |\tilde{f}(z'_{k'})|)\right) = O(n^{-c_0}),$$

where the implied constant in the $O()$ notation depends on C, ε, c_0 .

Proof. We may assume that n is sufficiently large depending on C, ε, c_0 , as the claim is trivial otherwise. We may also take ε to be small (e.g. $\varepsilon < 1/4$).

We use Theorem 16. Inspecting the hypotheses and conclusion of that theorem, we see that it will suffice to verify the delocalization bound

$$(90) \quad |z_j^i/\sqrt{i!}| \ll n^{-\alpha_1} V(z_j)^{1/2}$$

for all $1 \leq j \leq n$ and some $\alpha_1 > 0$ that can depend on ε but is independent of c_0 .

Fix j . As observed previously, the sequence $|z_j^i/\sqrt{i!}|$ is increasing for $i < |z_j|^2$ and decreasing for $i > |z_j|^2$. A routine application of Stirling's formula reveals that the magnitudes $|z_j^i/\sqrt{i!}|$ are comparable to each other for $i = |z_j|^2 + O(|z_j|)$, which in the regime $n^\varepsilon \leq |z_j| \leq \sqrt{n} + C$ occupies at least $\gg n^\varepsilon$ of the indices i in $\{0, \dots, n\}$, including the index i that maximizes $|z_j^i/\sqrt{i!}|$. The claim (90) then follows with $\alpha_1 := \varepsilon/2$. \square

As we have now verified all three axioms (i)-(iii) of Theorem 5, we obtain Theorem 18 as an immediate consequence. To establish Theorem 19, we see from Theorem 10 (and comparing both real atom distributions $\xi, \tilde{\xi}$ to the real gaussian distribution $N(0, 1)_{\mathbb{R}}$), it suffices to establish axiom (iv) of Theorem 10 in the case that $\tilde{\xi}$ has the distribution of $N(0, 1)_{\mathbb{R}}$. More precisely, it suffices to establish the following estimate (which is actually a little stronger than we need):

Proposition 38 (Level repulsion). *Let $\varepsilon > 0$, and let $C > 1$ be a sufficiently large constant. Let n be a natural number, and let $x, y \in \mathbb{R}$ and $z \in \mathbb{C}$ be such that*

$$(91) \quad n^\varepsilon \leq |x|, |y|, |z| \leq \sqrt{n} + C$$

and

$$|x - y|, |\operatorname{Im} z| \leq 1/C.$$

Let $f = f_{n, \xi}$ be a flat polynomial whose atom distribution ξ is drawn from the real gaussian ensemble $N(0, 1)_{\mathbb{R}}$. Then we have the pointwise bounds

$$(92) \quad \rho_f^{(2,0)}(x, y) \ll |x - y|$$

and

$$(93) \quad \rho_f^{(0,1)}(z) \ll |\operatorname{Im} z|,$$

where the implied constants depend on C .

A modification of the calculations below in fact show that the bounds (92), (93) continue to hold without the hypothesis (91), but we will only need the bounds under the hypothesis (91).

Proof. We will apply Lemma 17 with $R(z) := e^{-z^2/2}$. Thus, it suffices to establish the bounds

$$(94) \quad |v(z)| \ll 1$$

$$(95) \quad |v(x) \wedge v'(x)| \gg 1$$

$$(96)$$

for all $z \in B(x_0, 1)$, where

$$v(z) := e^{-z^2/2} (z^i / \sqrt{i!})_{i=0}^n.$$

We begin with the proof of (94). We have

$$|v(z)|^2 = |e^{-z^2}| \sum_{i=0}^n \frac{|z|^{2i}}{i!}.$$

Note that when $z \in B(x_0, 1)$, one has

$$|e^{-z^2}| \ll e^{-|z|^2}$$

while from Taylor series one has

$$\sum_{i=0}^n \frac{|z|^{2i}}{i!} \leq \sum_{i=0}^{\infty} \frac{|z|^{2i}}{i!} = e^{|z|^2}$$

and the claim (94) follows.

Now we prove (95). Observe that

$$v'(x) = e^{-x^2/2} \left(\frac{i-x^2}{x} \frac{x^i}{\sqrt{i!}} \right)_{i=0}^n$$

and so

$$|v(x) \wedge v'(x)|^2 = e^{-2x^2} \sum_{0 \leq i < j \leq n} \frac{|i-j|^2 x^{2i} x^{2j}}{x^2 i! j!}.$$

From Stirling's approximation we see that $\frac{x^{2i}}{i!}$ is comparable to $x^{-1}e^{x^2}$ when $i = x^2 + O(x)$, and the claim (95) easily follows (noting that $x^2 \leq n + O(x)$ when $x \leq \sqrt{n} + C$). \square

As all of the hypotheses (i)-(iv) of Theorem 10 are obeyed, Theorem 19 is now established.

Finally, we are able to establish Theorem 20. Let $\varepsilon, n, f = f_{n,\xi}$ be as in that theorem. From (89) we see that with overwhelming probability, there are at most $O(n^{1/4+o(1)})$ real zeroes outside the interval $[-\sqrt{n}-n^{1/4}, \sqrt{n}+n^{1/4}]$. Meanwhile, by covering the intervals $[-\sqrt{n}-n^{1/4}, -\sqrt{n}+n^{1/4}]$, $[-n^{1/4}, n^{1/4}]$, and $[\sqrt{n}-n^{1/4}, \sqrt{n}+n^{1/4}]$ by $O(n^{1/4})$ disks of radius 1 and applying (88) and the union bound, we see that with overwhelming probability, there are also $O(n^{1/4+o(1)})$ zeroes in these intervals. In view of these facts, it suffices to show that for any interval $I \subset [-\sqrt{n}+n^{1/4}, -n^{1/4}] \cup [n^{1/4}, \sqrt{n}-n^{1/4}]$, one has

$$\mathbf{E}N_I = \frac{1}{\pi}|I| + O(n^{1/2-c})$$

with probability $1 - O(n^{1/2-c})$.

By approximating the indicator function 1_I above and below by smooth functions, it will suffice to show that

$$\mathbf{E} \sum_{1 \leq i \leq n: \zeta_i \in \mathbb{R}} F(\zeta_i) = \frac{1}{\pi} \int_{\mathbb{R}} F(x) dx + O(n^{1/2-c})$$

for any smooth function $F : \mathbb{R} \rightarrow \mathbb{R}$ supported in $\{x \in \mathbb{R} : n^{1/4}/2 \leq |x| \leq \sqrt{n} - n^{1/4}/2\}$ which obeys the derivative bounds

$$|F^{(a)}(x)| \ll 1$$

for all $0 \leq a \leq 100$ (say).

Fix F . By (2), we may rewrite the above claim as the bound

$$(97) \quad \int_{\mathbb{R}} F(x) \rho_{f_{n,\xi}}^{(1,0)}(x) dx = \frac{1}{\pi} \int_{\mathbb{R}} F(x) dx + O(n^{1/2-c})$$

Now let $f_{n,\tilde{\xi}}$ be a flat polynomial whose atom distribution $\tilde{\xi}$ is given by the real gaussian $N(0,1)_{\mathbb{R}}$. By smoothly decomposing F into $O(n^{1/2})$ components each supported on an interval $[x-1, x+1]$ and applying Theorem 19 repeatedly, we see that

$$\int_{\mathbb{R}} F(x) \rho_{f_{n,\xi}}^{(1,0)}(x) dx = \int_{\mathbb{R}} F(x) \rho_{f_{n,\tilde{\xi}}}^{(1,0)}(x) dx + O(n^{1/2-c})$$

Thus, it suffices to establish the analogue of (97) for the real gaussian flat polynomial $f_{n,\tilde{\xi}}$. Such a bound can be implicitly extracted from the work of Edelman and Kostlan [12], but for the sake of completeness we give a proof of this bound here.

Using the Kac-Rice formula (Lemma 34) we have

$$\rho_{f_{n,\tilde{\xi}}}^{(1,0)}(x) = p_{\mathbb{R}}(f_{n,\tilde{\xi}}(x) = 0) \mathbf{E} \left(|f'_{n,\tilde{\xi}}(x)| \mid f_{n,\tilde{\xi}}(x) = 0 \right)$$

for any real x . By Lemma 35, we can write the right-hand side as

$$\frac{1}{\pi} \frac{|\text{dist}(\mathbf{w}_x, \mathbf{v}_x)|}{|\mathbf{v}_x|}$$

where

$$\mathbf{v}_x := \left(\frac{x^i}{\sqrt{i!}} \right)_{i=0}^n$$

and

$$\mathbf{w}_x := \left(\frac{i}{x} \frac{x^i}{\sqrt{i!}} \right)_{i=0}^n.$$

for any non-zero x . We can rearrange this as

$$\frac{1}{\pi} \frac{|\text{dist}(v'(x), v(x))|}{|v(x)|}$$

where

$$v(x) := e^{-x^2/2} \left(\frac{x^i}{\sqrt{i!}} \right)_{i=0}^n$$

and

$$v'(x) := e^{-x^2/2} \left(\frac{i-x^2}{x} \frac{x^i}{\sqrt{i!}} \right)_{i=0}^n.$$

We can expand

$$\begin{aligned} |v(x)|^2 &= e^{-x^2} \sum_{i=0}^n \frac{x^{2i}}{i!} \\ v(x) \cdot v'(x) &= e^{-x^2} \sum_{i=0}^n \frac{i - x^2}{x} \frac{x^{2i}}{i!} \\ |v'(x)|^2 &= e^{-x^2} \sum_{i=0}^n \left(\frac{i - x^2}{x} \right)^2 \frac{x^{2i}}{i!}. \end{aligned}$$

By differentiating the identity

$$e^{x^2} = \sum_{i=0}^{\infty} \frac{x^{2i}}{i!}$$

twice, we obtain the identities⁴

$$\sum_{i=0}^{\infty} \frac{i - x^2}{x} \frac{x^{2i}}{i!} = 0$$

and

$$\sum_{i=0}^{\infty} \left(\frac{i - x^2}{x} \right)^2 \frac{x^{2i}}{i!} = 1.$$

For x in the support of F , we have $n^\varepsilon/2 \leq |x| \leq n^{1/2} - n^{1/4}/2$, and if we truncate the above infinite sums to n using Stirling's approximation we conclude that

$$|v(x)|^2, |v'(x)|^2 = 1 + O(n^{-\varepsilon+o(1)})$$

and

$$v(x) \cdot v'(x) = O(n^{-\varepsilon+o(1)})$$

so that

$$\rho_{f_{n,\xi}}^{(1,0)}(x) = \frac{1}{\pi} + O(n^{-\varepsilon+o(1)}),$$

which gives (97) for $\tilde{\xi}$ and hence for ξ . This concludes the proof of Theorem 20.

12.1. The variance bound. As asserted in the introduction, one can extend these calculations to obtain a variance bound $\mathbf{Var} N_I = O(n^{1-c})$. We sketch the argument as follows. As before, we may assume that I is contained in the region $\{x : n^{1/4} \leq |x| \leq \sqrt{n} - n^{1/4}\}$. In addition to the bound (97) just established, one needs to establish the additional bound

$$(98) \quad \int_{\mathbb{R}} \int_{\mathbb{R}} F(x) F(y) \rho_{f_{n,\xi}}^{(2,0)}(x, y) dx = \left(\frac{1}{\pi} \int_{\mathbb{R}} F(x) dx \right)^2 + O(n^{1-c}).$$

Using Theorem 19 as before, we may replace ξ by $\tilde{\xi}$. We can then apply the Kac-Rice formula and Lemma 35 to conclude that

$$\rho_{f_{n,\tilde{\xi}}(x,y)}^{(2,0)} = \frac{1}{2\pi} |v(x) \wedge v(y)|^{-1} \mathbf{E}(|W_x| |W_y| | (V_x, V_y) = (0, 0))$$

⁴These are also the identities for the mean and variance of a Poisson random variable.

where V_x, V_y, W_x, W_y are real gaussian random variables with mean zero and covariance matrix

$$\mathbf{E} \begin{pmatrix} V_x^2 & V_x V_y & V_x W_x & V_x W_y \\ V_y V_x & V_y^2 & V_y W_x & V_y W_y \\ W_x V_x & W_x V_y & W_x^2 & W_x W_y \\ W_y V_x & W_y V_y & W_y W_x & W_y^2 \end{pmatrix} = \begin{pmatrix} |v(x)|^2 & v(x) \cdot v(y) & v(x) \cdot v_x(x) & v(x) \cdot v_x(y) \\ v(y) \cdot v(x) & |v(y)|^2 & v(y) \cdot v_x(x) & v(y) \cdot v_x(y) \\ v_x(x) \cdot v(x) & v_x(x) \cdot v(y) & |v_x(x)|^2 & v_x(x) \cdot v_x(y) \\ v_x(y) \cdot v(x) & v_x(y) \cdot v(y) & v_x(y) \cdot v_x(x) & |v_x(y)|^2 \end{pmatrix}.$$

A rather tedious calculation along the lines of those used in the proof of Theorem 20 reveals that this covariance matrix differs from the identity matrix by $O(\exp(-|x-y|^2/10)) + O(n^{-\varepsilon+o(1)})$ (say) in the region $n^{1/4}/2 \leq |x|, |y| \leq \sqrt{n} - n^{1/4}/2$, which implies that

$$\rho_{f_{n,\tilde{\xi}}^{(2,0)}}(x,y) = \frac{1}{\pi^2} + O(\exp(-|x-y|^2/10)) + O(n^{-\varepsilon+o(1)})$$

which gives (98) for $\tilde{\xi}$. We omit the details.

13. UNIVERSALITY FOR ELLIPTIC POLYNOMIALS

In this section we establish our main results for elliptic polynomials, namely Theorems 21, 22, 23. Our arguments here will be closely analogous to those for flat polynomials in the previous section.

This will largely be accomplished by invoking the results obtained in previous sections. Again, our starting point is the concentration of log-magnitudes.

Lemma 39 (Concentration for log-magnitude). *Let $C, \varepsilon > 0$ be constants, let n be a natural number, and let z be a complex number with*

$$n^\varepsilon \leq |z| \leq n^{1-\varepsilon}.$$

Let $f = f_{n,\xi}$ be a rescaled elliptic polynomial whose atom distribution ξ has mean zero and variance one with $\mathbf{E}|\xi|^{2+\varepsilon} \leq C$. Then with overwhelming probability, one has

$$\log |f(z)| = \frac{1}{2}n \log\left(1 + \frac{|z|^2}{n}\right) + O(n^{o(1)}).$$

The implied constants in the asymptotic notation can depend on C, ε .

Proof. As before, we first compute the quantity $V(z)$ from (14). This quantity is given by

$$V(z) = \sum_{i=0}^n \frac{|z|^{2i}}{n^i} \binom{n}{i} = (1 + |z|^2/n)^n.$$

In particular, we have

$$\log V(z) = n \log\left(1 + \frac{|z|^2}{n}\right).$$

Applying Lemma 13 as before, indices $i_1, \dots, i_m \in \{0, \dots, n\}$ for some $m = \omega(\log n)$ such that we have the lacunarity property

$$\left| \sqrt{\binom{n}{i_j} n^{-i_j} z^{i_j} / \sqrt{i_j!}} \right| \geq 2 \left| \sqrt{\binom{n}{i_{j+1}} n^{-i_{j+1}} z^{i_{j+1}}} \right|$$

for all $1 \leq j < m$, and the lower bound

$$\left| \sqrt{\binom{n}{i_m}} n^{-i_m} z^{i_m} \right| \geq V(z)^{1/2} \exp(-n^{o(1)}).$$

The sequence $i \mapsto \sqrt{\binom{n}{i}} n^{-i} z^i$ is increasing for $i < \frac{|z|^2}{1+|z|^2/n}$ and decreasing for $i > \frac{|z|^2}{1+|z|^2/n}$, with its largest value being at least $(V(z)/(n+1))^{1/2}$. If $n^\varepsilon \leq |z| \leq \sqrt{n}$, the ratio between adjacent elements of this sequence is $O(1)$ in the range

$$\frac{1}{2} \frac{|z|^2}{1+|z|^2/n} \leq i \leq \frac{|z|^2}{1+|z|^2/n},$$

and the claim then follows by applying Lemma 14 to the (reversal of) this subsequence. Conversely, if $\sqrt{n} \leq |z| \leq n^{1-\varepsilon}$, then the ratio between adjacent elements of this sequence is $O(1)$ in the range

$$\frac{1}{2} \frac{n}{1+|z|^2/n} \leq n-i \leq \frac{n}{1+|z|^2/n},$$

and the claim follows by Lemma 14 to this subsequence. \square

If we set

$$G(z) := \frac{1}{2} n \log\left(1 + \frac{|z|^2}{n}\right)$$

then a short computation shows that

$$\Delta G(z) = \frac{2}{(1+|z|^2/n)^2}$$

Applying Proposition 12, we conclude that for any $n^{-C} \leq c \leq r \leq n^{1-\varepsilon}/3$ and $z_0 \in B(0, n^{1-\varepsilon}/3)$ with the property that $B(z_0, r+c) \setminus B(z_0, r-c)$ is disjoint from $B(0, n^\varepsilon)$, with overwhelming probability f is non-vanishing and obeys

$$(99) \quad N_{B(z_0, r)}(f) = \int_{B(z_0, r)} \frac{1}{\pi} (1+|z|^2/n)^{-2} dz + O(n^{o(1)} c^{-1} r) + O\left(\int_{B(z_0, r+c) \setminus B(z_0, r-c)} (1+|z|^2/n)^{-2} dz\right).$$

This already gives axiom (i) for Theorems 5, 10. Setting $c = 2/r$, we conclude the non-concentration estimate

$$(100) \quad N_{B(z_0, r)} \ll n^{o(1)} r^2$$

with overwhelming probability for any $z_0 \in B(0, C\sqrt{n})$ and any $r \geq 1$ such that $B(z_0, r+c) \setminus B(z_0, r-c)$ is disjoint from $B(0, n^\varepsilon)$ (note that the claim is trivial for say $r \geq \sqrt{n}/3$); this gives axiom (ii) for Theorems 5, 10.

As before, the next stage is to establish axiom (iii) for Theorems 5, 10.

Proposition 40 (Comparability of log-magnitudes). *Let $C, \varepsilon > 0$ be constants, and let $c_0 > 0$ be sufficiently small depending on ε . Let n be a natural number, let $1 \leq k \leq n^{c_0}$ be another natural number, and let z_1, \dots, z_k be complex numbers such that*

$$n^\varepsilon \leq |z| \leq Cn^{1/2}.$$

Let $f_{n,\xi}, f_{n,\tilde{\xi}}$ be normalized elliptic polynomials whose atom distributions $\xi, \tilde{\xi}$ have mean zero and variance one matching moments to second order with $\mathbf{E}|\xi|^{2+\varepsilon}, |\tilde{\xi}|^{2+\varepsilon} \leq C$. Let $F : \mathbb{C}^k \rightarrow \mathbb{C}$ be a smooth function obeying the bounds

$$|\nabla^a F(z)| \leq C$$

for all $0 \leq a \leq 3$. Then, if c_0 is sufficiently small, one has

$$\mathbf{E}\left(F(\log |f(z'_1)|, \dots, \log |f(z'_{k'})|) - F(\log |\tilde{f}(z'_1)|, \dots, \log |\tilde{f}(z'_{k'})|)\right) = O(n^{-c_0}),$$

where the implied constant in the $O()$ notation depends on C, ε, c_0 .

This is proven in exact analogy with Proposition 37 (with the quantity $|z_j|^2$ being replaced by $\frac{|z_j|^2}{1+|z_j|^2/n}$); we leave the details to the interested reader.

Applying Theorem 5, we now obtain Theorem 21 as an immediate consequence. To similarly use Theorem 10 to deduce Theorem 22, we need the following analogue of Proposition 38:

Proposition 41 (Level repulsion). *Let $\varepsilon > 0$, and let $C > 1$ be a sufficiently large constant. Let n be a natural number, and let $x, y \in \mathbb{R}$ and $z \in \mathbb{C}$ be such that*

$$(101) \quad n^\varepsilon \leq |x|, |y|, |z| \leq C\sqrt{n}$$

and

$$|x - y|, |\operatorname{Im}z| \leq 1/C.$$

Let $f = f_{n,\xi}$ be a rescaled elliptic polynomial whose atom distribution is drawn from the real gaussian ensemble $N(0, 1)_{\mathbb{R}}$. Then we have the pointwise bounds

$$(102) \quad \rho_{\tilde{f}}^{(2,0)}(x, y) \ll |x - y|$$

and

$$(103) \quad \rho_{\tilde{f}}^{(0,1)}(z) \ll |\operatorname{Im}z|,$$

where the implied constants depend on C .

Proof. Applying Lemma 17 with the holomorphic function $R(z) := (1 + z^2/n)^{-n/2}$ on $B(x, 1)$ (noting that we are well away from the poles $\pm\sqrt{-1}\sqrt{n}$ of this function), it suffices to establish the bounds

$$(104) \quad |v(z)| \ll 1$$

$$(105) \quad |v(x) \wedge v'(x)| \gg 1$$

for all $x \in \mathbb{R}$ with $n^\varepsilon/2 < |x| \leq 2C\sqrt{n}$ and all $z \in B(x, 1)$.

To prove (104), we compute

$$\begin{aligned} |v(z)|^2 &:= |1 + z^2/n|^{-n} \sum_{i=0}^n \binom{n}{i} n^{-i} |z|^{2i} \\ &= \left(\frac{1 + |z|^2/n}{|1 + z^2/n|} \right)^n. \end{aligned}$$

But one can compute that $|1 + z^2/n| = 1 + |z|^2/n + O(1/n)$, and the claim (104) follows.

To obtain (105), we compute

$$v'(x) = (1 + x^2/n)^{-n/2} \left(\frac{i - x^2/(1 + x^2/n)}{x} \sqrt{\binom{n}{i} n^{-i} x^i} \right)_{i=0}^n$$

and so

$$|v(x) \wedge v'(x)|^2 = (1 + x^2/n)^{-2n} \sum_{0 \leq i < j \leq n} \frac{|i - j|^2}{x^2} \binom{n}{i} n^{-i} x^{2i} \binom{n}{j} n^{-j} x^{2j}.$$

The expression on the right-hand side can in fact be computed exactly, but for the purposes of establishing the lower bound (105), we may appeal instead to Stirling's formula, which reveals that

$$\binom{n}{i} n^{-i} x^{2i} \gg \frac{1}{x} (1 + x^2/n)^n$$

when $i = \frac{x^2}{1+x^2/n} + O(x)$, and the claim follows much as in the analogous computation for flat polynomials in Proposition 38. \square

As all of the hypotheses (i)-(iv) of Theorem 10 are obeyed, Theorem 22 is now established.

Finally, we establish Theorem 23. We need to show

$$(106) \quad \mathbf{E}N_I = \int_I \frac{1}{\pi} \frac{dx}{1 + x^2/n} + O(n^{1/2-c})$$

for all intervals I .

It will be convenient to use inversion symmetry to work in the region $I \subset [-\sqrt{n}, \sqrt{n}]$. Observe that if

$$f_{n,\xi}(z) = \sum_{i=0}^n \xi_i \sqrt{\binom{n}{i} n^{-i} z^i}$$

is a rescaled elliptic polynomial, then

$$n^{-n/2} z^n f_{n,\xi}(n/z) = \sum_{i=0}^n \xi_{n-i} \sqrt{\binom{n}{i} n^{-i} z^i}$$

is also a rescaled elliptic polynomial with the same distribution as $f_{n,\xi}$. Thus the distribution of the zeroes of $f_{n,\xi}$ are invariant with respect to the inversion map $z \mapsto n/z$. Among other things, this implies that if (106) holds for an interval I avoiding the origin, then it also holds for the inverse interval $\{n/x : x \in I\}$. From this (and (100)) it thus suffices to establish (106) in the case $I \subset [-\sqrt{n}, \sqrt{n}]$.

Covering the interval $[-n^{1/4}, n^{1/4}]$ by $O(n^{1/4})$ balls of unit radius and then applying (100), we see that this interval has $O(n^{1/4+o(1)})$ zeroes with overwhelming

probability. This establishes (106) when $I \subset [-n^{1/4}, n^{1/4}]$, so we may assume without loss of generality that $I \subset [-\sqrt{n}, -n^{1/4}]$ or $I \subset [n^{1/4}, \sqrt{n}]$.

By using Theorem 22 (and upper and lower bounding 1_I by smooth functions), it suffices to establish (106) in the case when $I \subset \{x : n^{1/4}/2 \leq |x| \leq 2\sqrt{n}\}$ and ξ has the real gaussian distribution $N(0, 1)_{\mathbb{R}}$. By the Kac-Rice formula as in the previous section, we have

$$\mathbf{E}N_I = \int_I \rho^{(1,0)}(x) dx$$

where

$$\rho^{(1,0)}(x) = \frac{1}{\pi} \frac{|\text{dist}(v'(x), v(x))|}{|v(x)|},$$

$$v(x) := (1 + x^2/n)^{-n/2} \left(\sum_{i=0}^n \binom{n}{i} n^{-i} x^i \right)$$

and

$$v'(x) := (1 + x^2/n)^{-n/2} \left(\frac{i - \frac{x^2}{1+x^2/n}}{x} \sqrt{\binom{n}{i} n^{-i} x^i} \right)_{i=0}^n.$$

We have

$$\begin{aligned} |v(x)|^2 &= (1 + x^2/n)^{-n} \sum_{i=0}^n \binom{n}{i} n^{-i} x^{2i} \\ &= 1 \end{aligned}$$

and thus on differentiation

$$v(x) \cdot v'(x) = 0.$$

We also have

$$|v'(x)|^2 = (1 + x^2/n)^{-n} \sum_{i=0}^n \frac{(i - \frac{x^2}{1+x^2/n})^2}{x^2} \binom{n}{i} n^{-i} x^{2i}.$$

We can differentiate the binomial identity

$$(1 + x^2/n)^n = \sum_{i=0}^n \binom{n}{i} n^{-i} x^{2i}$$

to obtain

$$\frac{x^2}{1 + x^2/n} (1 + x^2/n)^n = \sum_{i=0}^n i \binom{n}{i} n^{-i} x^{2i}$$

and

$$\frac{x^4 + 1}{(1 + x^2/n)^2} (1 + x^2/n)^n = \sum_{i=0}^n i^2 \binom{n}{i} n^{-i} x^{2i}$$

(these are also the formulae for the mean and variance of a binomial random variable) and so

$$|v'(x)|^2 = \frac{1}{(1 + x^2/n)^2}.$$

This implies that

$$\rho^{(1,0)}(x) = \frac{1}{\pi} \frac{1}{1 + x^2/n}$$

and (106) follows (indeed, the formula is even exact in this case). As a matter of fact, we can improve the error term $O(n^{1/2-\epsilon})$ to $O(|I|n^{-\epsilon})$ in this case.

14. UNIVERSALITY FOR KAC POLYNOMIALS

We now prove Theorems 24 and 25. As before, the first step is to obtain concentration results for the log-magnitude $\log |f(z)|$. Here, a new difficulty arises: when z is a root of unity, then $f(z)$ can vanish with polynomially small probability. For instance, if $z = 1$, n is odd, and ξ has the Bernoulli distribution (thus $P(\xi = +1) = P(\xi = -1) = 1/2$), then $f(1) = \xi_0 + \dots + \xi_n$ vanishes with probability comparable to $1/\sqrt{n}$, as can be easily verified using Stirling's formula. The key new idea in our proof is to show that this is essentially the only obstruction to concentration of the log magnitude, provided that z stays away from zero and from infinity.

Lemma 42 (Concentration for log-magnitude). *Let $C, A, \epsilon > 0$ be constants, let n be a natural number, and let z be a complex number with*

$$\epsilon \leq |z| \leq 1/\epsilon.$$

Let $f = f_{n,\xi}$ be a Kac polynomial whose atom distribution ξ has mean zero and variance one with $\mathbf{E}|\xi|^{2+\epsilon} \leq C$. Then one of the following holds:

- (i) $\epsilon \leq |z| \leq 1$, and one has $\log |f(z)| = O(n^{o(1)})$ with probability $1 - O(n^{-A})$.
- (ii) $1 \leq |z| \leq 1/\epsilon$, and one has $\log |f(z)| = n \log |z| + O(n^{o(1)})$ with probability $1 - O(n^{-A})$.
- (iii) One has $z = \omega + O(n^{-A})$, where ω is a root of unity with $\omega^k = 1$ for some $k = O(1)$.

The implied constants in the asymptotic notation can depend on C, ϵ, A .

Proof. We may assume n sufficiently large depending on C, ϵ, A , as the claim is trivial otherwise.

Note that $z \mapsto z^n f(1/z)$ is a Kac polynomial with the same distribution as f , so the claims in this lemma for $|z| \geq 1$ will follow from the claims when $|z| \leq 1$. Thus we may assume without loss of generality that $\epsilon \leq |z| \leq 1$.

If $\epsilon \leq |z| \leq 1 - \frac{\log^2 n}{n}$ (say) then the claim (i) follows easily from Lemma 13 (and Lemma 14), so we may assume that

$$(107) \quad 1 - \frac{\log^2 n}{n} \leq |z| \leq 1.$$

The random variable $f(z)$ has mean zero and variance

$$\mathbf{E}|f(z)|^2 = \sum_{i=0}^n |z|^{2i} \leq n$$

and so by Chebyshev's inequality we certainly have the upper bound

$$\log |f(z)| \leq n^{o(1)}$$

with overwhelming probability. If we have

$$\log |f(z)| \geq -\log^3 n$$

(say) with probability at least $1 - n^{-A}$ we are done, so suppose instead that

$$\mathbf{P}(\log |f(z)| \geq -\log^2 n) < 1 - n^{-A},$$

thus

$$(108) \quad \mathbf{P}(|f(z)| < \exp(-\log^3 n)) > n^{-A}.$$

The quantity $f(z) = \sum_{i=0}^n \xi_i z^i$ is a sum of independent random variables, and so (108) is an assertion that the *small ball probability* of this random sum is large. We can use this to constrain the coefficients z^i of $f(z)$ by means of *inverse Littlewood-Offord theorems*. There are many such theorems in the literature; we will use [44, Theorem 2.9]. We first note from Lemma 31 that

$$\mathbf{P}(B^{-1} \leq |\xi - \xi'| \leq B) \gg 1$$

for some $B = O(1)$, if ξ' is an independent copy of ξ . This is essentially⁵ the hypothesis in [44, (7)] up to some rescalings. If one applies [44, Theorem 2.9], one can then complex numbers v_1, \dots, v_r for some $r = O(1)$ with the property that for all but at most \sqrt{n} (say) of the numbers $z^i, 0 \leq i \leq n$, one has a representation of the form

$$(109) \quad z^i = a_{i,1}v_1 + \dots + a_{i,r}v_r + O(n^{O(1)} \exp(-\log^3 n))$$

where $a_{i,1}, \dots, a_{i,r}$ are integers of magnitude $O(n^{O(1)})$; in particular, by the pigeon-hole principle we can find $0 \leq i_0 \leq n - \sqrt{n}$ such that one has a representation (109) for all $i_0 \leq i \leq i_0 + \sqrt{n}$. Actually, the results in [44] provide significantly more precise results than this, but these bounds will suffice for our purposes.

It will be convenient to ensure that the generators v_1, \dots, v_r are approximately linearly independent in a certain sense. Observe that if we have an approximate linear relation between the v_1, \dots, v_r of the form

$$(110) \quad b_1v_1 + \dots + b_rv_r = O(n^{O(1)} \exp(-\log^3 n))$$

for some integers $b_1, \dots, b_r = O(n^{O(1)})$, not all zero, then after clearing denominators we can eliminate one of the v_i from the basis v_1, \dots, v_r and divide all the other elements by integers of size $O(n^{O(1)})$ and obtain a new basis of $r - 1$ elements for which one still has representations of the form (109) (with worse values of implied constants in the $O()$ notation) for all $i_0 \leq i \leq i_0 + \sqrt{n}$. Iterating this observation

⁵In [44] the lower bound on $\mathbf{P}(B^{-1} \leq |\xi - \xi'| \leq B)$ is $1/2$ rather than $\gg 1$, but one can verify that the arguments in that paper are not significantly changed if one alters the lower bound, provided of course one allows all subsequent constants to depend on this new lower bound.

at most r times, we may assume without loss of generality that there is no⁶ linear relation of the form (110).

Among other things, this approximate linear independence shows (if the $O()$ notation is suitably interpreted) that the r -tuple $\vec{a}_i := (a_{i,1}, \dots, a_{i,r}) \in \mathbf{Z}^r$ appearing in (109) is uniquely defined for each $i_0 \leq i \leq i_0 + \sqrt{n}$. From (107) and the approximate linear independence we also see that the \vec{a}_i are all non-zero.

Note that for any $i_0 \leq i_1 \leq i_2 \leq i_0 + \sqrt{n}$, the linear span V_{i_1, i_2} of the vectors \vec{a}_i for $i_1 \leq i \leq i_2$ has dimension between 1 and r , and is non-decreasing in i_2 and non-increasing in i_1 . By the pigeonhole principle, one can thus find

$$i_0 + 0.1\sqrt{n} \leq i_1 \leq i_2 \leq i_0 + 0.9\sqrt{n}$$

(say) such that

$$V_{i_1, i_2} = V_{i_1 - n^{1/4}, i_2 + n^{1/4}}$$

(say). In particular, we have $\vec{a}_i \in V_{i_1, i_2}$ for all $i_1 - n^{1/4} \leq i \leq i_2 + n^{1/4}$.

By the Steinitz exchange lemma, we can find a sequence $\vec{a}_{j_1}, \dots, \vec{a}_{j_d}$ for $i_1 \leq j_1 < \dots < j_d \leq i_2$ and some $d = O(1)$ that span V_{i_1} . Then for any integer m with $-n^{1/4} \leq m \leq n^{1/4}$, the vectors $\vec{a}_{j_1+m}, \dots, \vec{a}_{j_d+m}$ are linear combinations of $\vec{a}_{j_1}, \dots, \vec{a}_{j_d}$, thus we have

$$(111) \quad \begin{pmatrix} \vec{a}_{j_1+m} \\ \dots \\ \vec{a}_{j_d+m} \end{pmatrix} = T_m \begin{pmatrix} \vec{a}_{j_1} \\ \dots \\ \vec{a}_{j_d} \end{pmatrix}$$

for some (unique) $d \times d$ matrix T_m with integer entries. From Cramer's rule we see that all entries of T_m have magnitude $O(n^{O(1)})$ for any $-n^{1/4} \leq m \leq n^{1/4}$.

Clearly T_0 is the identity matrix. We claim that

$$(112) \quad T_{m+1} = T_m T_1$$

for any $-n^{1/4} \leq m \leq n^{1/4} - 1$, which implies by induction that T_1 is invertible (so in particular T_1 lies in $\mathrm{SL}_d(\mathbf{Z})$) and $T_m = T_1^m$ for all $-n^{1/4} \leq m \leq n^{1/4}$. To see this, we note from (111) and (109) that

$$(113) \quad \begin{pmatrix} z^{j_1+m} \\ \dots \\ z^{j_d+m} \end{pmatrix} = T_m \begin{pmatrix} z^{j_1} \\ \dots \\ z^{j_d} \end{pmatrix} + O(n^{O(1)} \exp(-\log^3 n)).$$

⁶Strictly speaking, one has to take some care with the asymptotic notation $O()$ in order to make this statement rigorous. There are several (essentially equivalent) ways in which this can be achieved. One is to reformulate the current argument (which is written in the context of a fixed n) in asymptotic fashion, involving a sequence of values of n tending to infinity, with $O(1)$ now referring to a quantity that is bounded uniformly in n , at which point there is no difficulty interpreting the argument here rigorously. Another approach, which we will not detail here, is to reformulate the argument in the language of nonstandard analysis via the device of forming an ultraproduct, so that n is now a nonstandard natural number rather than a standard one, and the $O()$ notation again has a precise interpretation. If instead one wishes to stay in the context of a fixed (standard) n , then one interprets (110) as the claim that there are no integers b_1, \dots, b_r of magnitude at most $F(C)n^{F(C)}$, not all zero, for which $|b_1 v_1 + \dots + b_r v_r| \leq F(C)n^{F(C)} \exp(-\log^3 n)$, where C bounds all the implied constants in previous usages of asymptotic notation (such as (109)) and $F: \mathbb{R}^+ \rightarrow \mathbb{R}^+$ is a sufficiently rapidly growing function (not depending on n) to be chosen later.

Multiplying by z , we see that

$$\begin{pmatrix} z^{j_1+m+1} \\ \dots \\ z^{j_d+m+1} \end{pmatrix} = T_m \begin{pmatrix} z^{j_1+1} \\ \dots \\ z^{j_d+1} \end{pmatrix} + O(n^{O(1)} \exp(-\log^3 n))$$

and by comparing this with (113) with m replaced by $m+1$ and by 1, we conclude that

$$T_{m+1} \begin{pmatrix} z^{j_1} \\ \dots \\ z^{j_d} \end{pmatrix} = T_m T_1 \begin{pmatrix} z^{j_1} \\ \dots \\ z^{j_d} \end{pmatrix} + O(n^{O(1)} \exp(-\log^3 n)).$$

Using (109) and the approximate linear independence of the v_1, \dots, v_r , we conclude that

$$T_{m+1} \begin{pmatrix} \vec{a}_{j_1} \\ \dots \\ \vec{a}_{j_d} \end{pmatrix} = T_m T_1 \begin{pmatrix} \vec{a}_{j_1} \\ \dots \\ \vec{a}_{j_d} \end{pmatrix}$$

and from the linear independence of the $\vec{a}_{j_1}, \dots, \vec{a}_{j_d}$ we conclude (112).

We now see that T_1 is a matrix in $\mathrm{SL}_d(\mathbf{Z})$ with the property that T_1^m has magnitude $O(n^{O(1)})$ for all $-n^{1/4} \leq m \leq n^{1/4}$. This is a polynomial growth condition on T_1 , and one can use results related⁷ to quantitative versions [48] of Gromov's theorem to then force T_1 to be virtually unipotent. Indeed, if we apply [48, Proposition 13.1], we conclude that there exists a non-zero vector $\vec{c} \in \mathbf{Z}^d$ and a natural number $k = O(1)$ such that $T_1^k \vec{c} = \vec{c}$. From Cramer's rule we can take $\vec{c} = (c_1, \dots, c_d)$ to have magnitude $O(n^{O(1)})$. From (111) and (109) we see that

$$(114) \quad \sum_{l=1}^d c_l z^{j_l+k} = \sum_{l=1}^d c_l z^{j_l} + O(n^C \exp(-\log^3 n))$$

for some $C = O(1)$. If we had

$$\left| \sum_{l=1}^d c_l z^{j_l} \right| \leq n^{A+C} \exp(-\log^3 n)$$

then by (109) this would contradict the approximate linear independence of the v_1, \dots, v_r , the actual linear independence of the $\vec{a}_{j_1}, \dots, \vec{a}_{j_d}$, and the non-zero nature of \vec{c} , so we have

$$\left| \sum_{l=1}^d c_l z^{j_l} \right| > n^{A+C} \exp(-\log^3 n)$$

and hence from (114) one has $z^k = 1 + O(n^{-A})$, which gives the conclusion (iii). \square

As the roots of unity are fairly sparse, they can be avoided for the purposes of obtaining non-concentration bounds on zeroes:

⁷The situation here does not require the full strength of Gromov's theorem [18] (or quantitative versions thereof), and is actually closer to the older work of Milnor [42] and Wolf [63] treating polynomial growth in solvable groups.

Lemma 43 (Non-clustering bounds). *Let $C, \varepsilon > 0$ be constants, and let n be a natural number. Let $f = f_{n, \xi}$ be a Kac polynomial whose atom distribution ξ has mean zero and variance one with $\mathbf{E}|\xi|^{2+\varepsilon} \leq C$. Let $B(z_0, r)$ be a ball in the complex plane. Then one has*

$$N_{B(z_0, r)} \ll n^{o(1)}(1 + nr)$$

with overwhelming probability. If in addition $B(z_0, 2r)$ is disjoint from the unit circle $\{z \in \mathbb{C} : |z| = 1\}$, one can improve this bound to

$$N_{B(z_0, r)} \ll n^{o(1)}$$

with overwhelming probability. Furthermore, f is non-vanishing with overwhelming probability.

Proof. We apply Proposition 12 with the function $G : \mathbb{C} \rightarrow \mathbb{R}$ defined by setting $G(z) := 0$ for $|z| \leq 1$, and $G(z) := n \log |z|$ for $|z| > 1$. Strictly speaking, G is not smooth enough for Proposition 12 to apply as stated, but this technical difficulty can be overcome by a routine infinitesimal mollification which we omit here. One can compute from Green's theorem that

$$\frac{1}{2\pi} \Delta G = n d\sigma$$

in the sense of distributions, where $d\sigma$ is the uniform probability measure on the unit circle $\{z : |z| = 1\}$. From Proposition 12 (with Remark 30) and Lemma 42, we conclude that f is non-vanishing with overwhelming probability, and for any fixed $A > 0$, any ball $B(z_0, r)$, and any $0 < c \leq r$ with $r \ll n^{O(1)}$ and $c \gg n^{-O(1)}$, one has

$$N_{B(z_0, r)}(f) \ll n \int_{B(z_0, r+c)} d\sigma + O(n^{o(1)}c^{-1}r)$$

whenever the region $B(z_0, r+c) \setminus B(z_0, r)$ lies in the annulus $\{z : 1/10 \leq |z| \leq 10\}$ (say) and also avoids the disks $B(\omega, n^{-A})$ whenever $\omega^k = 1$ for some $k \leq C_A$, where C_A depends only on A . In particular, under these hypotheses we have

$$N_{B(z_0, r)}(f) \ll n^{o(1)}(nr + c^{-1}r)$$

together with the stronger bound

$$N_{B(z_0, r)}(f) \ll n^{o(1)}c^{-1}r$$

when $B(z_0, r+c)$ is disjoint from the unit circle.

From these estimates, we see that

$$N_{B(0, 0.9)}(f) \ll n^{o(1)}$$

(say) with probability $1 - O(n^{-A})$, and by setting $c = r$ we also see that

$$N_{B(z_0, (1-|z_0|)/10)}(f) \ll n^{o(1)}$$

with probability $1 - O(n^{-A})$ for any z_0 with $1/2 \leq |z_0| \leq 1 - 1/n$. By letting z_0 lie on the unit circle, setting c to be a small multiple of r , and enlarging r as necessary in order for $B(z_0, r+c) \setminus B(z_0, r)$ to avoid the disks $B(\omega, n^{-A})$, we also conclude that

$$N_{B(z_0, r)} \ll n^{o(1)}(1 + nr)$$

with probability $1 - O(n^{-A})$ for any $0 < r < 1/10$ (say) and z_0 on the unit circle. By diagonalization in A , these events in fact hold with overwhelming probability. From these bounds and covering argument one obtains the required bounds for the contribution of the zeroes on or inside the unit circle; the contribution of the zeroes outside the unit circle can then be handled by exploiting the invariance of the distribution of the zeroes with respect to the transformation $z \mapsto 1/z$. \square

Following our treatment of the flat and elliptic polynomials, the next step is to obtain comparability of log-magnitudes.

Proposition 44 (Comparability of log-magnitudes). *Let $C, \varepsilon > 0$ be constants, and let $c_0 > 0$ be sufficiently small depending on ε . Let n be a natural number, let $1 \leq k \leq n^{c_0}$ be another natural number, and let z_1, \dots, z_k be complex numbers such that*

$$1 - n^{-\varepsilon} \leq |z_j| \leq 1 + n^{-\varepsilon}$$

for $j = 1, \dots, k$. Let $f_{n,\xi}, f_{n,\tilde{\xi}}$ be Kac polynomials whose atom distributions $\xi, \tilde{\xi}$ have mean zero and variance one matching moments to second order with $\mathbf{E}|\xi|^{2+\varepsilon}, |\tilde{\xi}|^{2+\varepsilon} \leq C$. Let $F : \mathbb{C}^k \rightarrow \mathbb{C}$ be a smooth function obeying the bounds

$$|\nabla^a F(z)| \leq C$$

for all $0 \leq a \leq 3$. Then, if c_0 is sufficiently small, one has

$$\mathbf{E}\left(F(\log |f(z'_1)|, \dots, \log |f(z'_{k'})|) - F(\log |\tilde{f}(z'_1)|, \dots, \log |\tilde{f}(z'_{k'})|)\right) = O(n^{-c_0}),$$

where the implied constant in the $O()$ notation depends on C, ε, c_0 .

Proof. This is immediate from Theorem 16, noting that for $1 - n^{-\varepsilon} \leq |z_j| \leq 1$ one has $V(z_j)^{1/2} \gg n^\varepsilon$ and for $1 \leq |z_j| \leq 1 + n^{-\varepsilon}$ one has $V(z_j)^{1/2} \gg |z_j|^{n^\varepsilon}$. \square

Theorem 24 now follows from Theorem 5 (with $r_0 := 1$ and $a_0 := 3$), after first performing the rescaling

$$f'(z) := f(10^{-3}rz)$$

and replacing the z_j by $z_j/(10^{-3}r)$; note that the correlation functions rescale according to the law

$$\rho_{f'}^{(k)}(z_1, \dots, z_k) := (10^{-3}r)^{2k} \rho_f^{(k)}(10^{-3}rz_1, \dots, 10^{-3}rz_k).$$

Finally, to establish Theorem 25, we need a rescaled level repulsion estimate:

Proposition 45 (Level repulsion). *Let $\varepsilon > 0$, and let $C > 1$ be a sufficiently large constant. Let n be a natural number, let r be a radius with*

$$\frac{1}{n} \leq r \leq n^{-\varepsilon},$$

and let $x, y \in \mathbb{R}$ and $z \in \mathbb{C}$ be such that

$$(115) \quad r \leq \frac{1}{n} + ||x| - 1|, \frac{1}{n} + ||y| - 1|, \frac{1}{n} + ||z| - 1| \leq 2r$$

and and

$$|x - y|, |\operatorname{Im}z| \leq r/C.$$

Let $f = f_{n,\xi}$ be a Kac polynomial whose atom distribution is drawn from the real gaussian ensemble $N(0, 1)_{\mathbb{R}}$. Then we have the pointwise bounds

$$(116) \quad \rho_{\tilde{f}}^{(2,0)}(x, y) \ll |x - y|/r^3$$

and

$$(117) \quad \rho_{\tilde{f}}^{(0,1)}(z) \ll |\operatorname{Im}z|/r^3,$$

where the implied constants depend on C .

Proof. We will work in the regime when $|x|, |y|, |z| \leq 1 + \frac{1}{n}$; the opposing case $|x|, |y|, |z| \geq 1 - \frac{1}{n}$ can be treated similarly, and in any event is essentially equivalent to the former case after using the symmetry $z \mapsto 1/z$ of the distribution of the zeroes of a Kac polynomial.

We work with the rescaled polynomials $\tilde{f}'(z) := \tilde{f}(rz)$, and note that it suffices to show that

$$(118) \quad \rho_{\tilde{f}'}^{(2,0)}(x', y') \ll |x' - y'|$$

and

$$(119) \quad \rho_{\tilde{f}'}^{(0,1)}(z') \ll |\operatorname{Im}z'|$$

where $x' := x/r$, $y' := y/r$, $z' := z/r$.

Applying Lemma 17, and then undoing the rescaling, it suffices to show that

$$(120) \quad |v(z)| \ll 1$$

$$(121) \quad |v(x) \wedge v'(x)| \gg 1/r$$

whenever $z \in B(x, \frac{1}{Cr})$, $-1 - \frac{1}{n} \leq x \leq 1 + \frac{1}{n}$ is such that

$$r \leq \frac{1}{n} + ||x| - 1| \leq 2r,$$

and

$$v(z) := R(z)(z^i)_{i=0}^n$$

where we can for instance take

$$R(z) := (1 - z^2 + \frac{100}{n})^{1/2}.$$

To prove (120), we expand

$$|v(z)|^2 \leq |1 - z^2 + \frac{100}{n}| \sum_{i=0}^n |z|^{2i}.$$

Bounding $|1 - z^2 + \frac{100}{n}| \ll 1 - |z|^2$ and

$$\sum_{i=0}^n |z|^{2i} \leq \sum_{i=0}^{\infty} |z|^{2i} = \frac{1}{1 - |z|^2}$$

we obtain (120). To obtain (121), we compute

$$v'(x) = R(x) \left(\left(\frac{i}{x} - \frac{x}{1 - x^2 + \frac{100}{n}} \right) x^i \right)_{i=0}^n$$

and so

$$|v(x) \wedge v'(x)|^2 = \left|1 - x^2 + \frac{100}{n}\right|^2 \sum_{0 \leq i, j \leq n} \frac{|i-j|^2}{x^2} x^{2i} x^{2j}.$$

But $1 - x^2 + \frac{100}{n}$ is comparable to r , and x^{2i} is comparable to 1 when $i = O(1/r)$, and the claim (121) follows. \square

Theorem 25 then follows from Theorem 10 after using the same rescaling used to establish Theorem 24.

Acknowledgement. We would like to thank O. Nguyen, M. Krishnapur and T. Reddy for useful comments.

REFERENCES

- [1] G. Anderson, A. Guionnet, O. Zeitouni, An introduction to random matrices. Cambridge Studies in Advanced Mathematics, 118. Cambridge University Press, Cambridge, 2010.
- [2] R. Adler, J. Taylor, Random fields and geometry. Springer Monographs in Mathematics. Springer, New York, 2007.
- [3] A. T. Bharucha-Reid, M. Sambandham, Random polynomials. Probability and Mathematical Statistics. Academic Press, Inc., Orlando, Fla., 1986.
- [4] A. Bloch, G. Polya, On the roots of certain algebraic equations, *Proc. London Math. Soc.* **33**(1932), 102–114.
- [5] E. Bogomolny, O. Bohias and P. Leboeuf, Distribution of roots of polynomials, *Physical Review Letters*, **68** (1992), 2726–2729.
- [6] S. Chatterjee, A simple invariance theorem, unpublished.
- [7] P. Bleher, X. Di, Correlations between zeros of non-Gaussian random polynomials, *Int. Math. Res. Not.*, **46** (2004), 2443–2484.
- [8] P. Bourgade, H.-T. Yau and J. Yin, Local Circular Law for Random Matrices, preprint, [arXiv:1206.1449](https://arxiv.org/abs/1206.1449).
- [9] P. Bourgade, H.-T. Yau and J. Yin, The local circular law II: the edge case, preprint, [arXiv:1206.3187](https://arxiv.org/abs/1206.3187).
- [10] A. Dembo, B. Poonen, Q.-M. Shao, O. Zeitouni, Random polynomials having few or no real zeros, *J. Amer. Math. Soc.* **15** (2002), no. 4, 857–892.
- [11] I. Emiris, A. Gallago, E. Tsigaridas, Random polynomials and expected complexity of bisection methods for real solving, *ISSAC '10 Proceedings of the 2010 International Symposium on Symbolic and Algebraic Computation*, 235–242.
- [12] A. Edelman, E. Kostlan, How many zeros of a random polynomial are real?, *Bull. Amer. Math. Soc. (N.S.)* **32** (1995), 1–37. Erratum: *Bull. Amer. Math. Soc. (N.S.)* **33** (1996), 325.
- [13] T. Erdélyi, Extensions of the Bloch-Pólya theorem on the number of real zeroes of polynomials, *J. Théor. Nombres Bordeaux* **20** (2008), no. 2, 281–287.
- [14] P. Erdős, A. C. Offord, On the number of real roots of a random algebraic equation, *Proc. London Math. Soc.* **6** (1956), 139–160.
- [15] K. Farahmand, Topics in random polynomials. Pitman research notes in mathematics series 393. Longman, Harlow, 1998.
- [16] P. J. Forrester, G. Honner, Exact statistical properties of the zeros of complex random polynomials, *J. Phys. A*, **32** (1999), 2961–2981.
- [17] J. Ginibre, Statistical ensembles of complex, quaternion, and real matrices, *Journal of Mathematical Physics* **6** (1965), 440–449.
- [18] M. Gromov, Groups of polynomial growth and expanding maps, *Inst. Hautes Études Sci. Publ. Math.* **53** (1981), 53–73.
- [19] J. Hough, M. Krishnapur, Y. Peres, B. Virág, Zeros of Gaussian Analytic Functions and Determinantal Point Processes, University Lecture Series, American Mathematical Society, 2009.

- [20] I. A. Ibragimov, N. B. Maslova, The average number of zeros of random polynomials, *Vestnik Leningrad. Univ.* **23** (1968), 171–172.
- [21] I. A. Ibragimov, N. B. Maslova, The mean number of real zeros of random polynomials. I. Coefficients with zero mean, *Theor. Probability Appl.* **16** (1971), 228–248.
- [22] I. A. Ibragimov, N. B. Maslova, The mean number of real zeros of random polynomials. II. Coefficients with a nonzero mean., *Theor. Probability Appl.* **16** (1971), 485–493.
- [23] I. A. Ibragimov, N. B. Maslova, The average number of real roots of random polynomials, *Soviet Math. Dokl.* **12** (1971), 1004–1008.
- [24] I. A. Ibragimov and O. Zeitouni, On roots of random polynomials, *Trans. Amer. Math. Soc.* **349** (1997), no. 6, 2427–2441.
- [25] Z. Kabluchko, D. Zaporozhets, *Universality for zeros of random analytic functions*, preprint. [arXiv:1205.5355](https://arxiv.org/abs/1205.5355)
- [26] M. Kac, On the average number of real roots of a random algebraic equation, *Bull. Amer. Math. Soc.* **49** (1943) 314–320.
- [27] M. Kac, On the average number of real roots of a random algebraic equation. II. *Proc. London Math. Soc.* 50, (1949), 390–408.
- [28] M. Kac, Probability and related topics in physical sciences. Lectures in Applied Mathematics. Proceedings of the Summer Seminar, Boulder, Colo., 1957, Vol. I Interscience Publishers, London-New York, 1959.
- [29] E. Kostlan, On the distribution of roots of random polynomials, Chapter 38, *From Topology to Computation: Proceeding of the Samefest*, edited by M. W. Hirsch, J.E. , Marsden and M. Shub, Springer-Verlag, NY 1993.
- [30] M. Krishnapur, B. Virág, The Ginibre ensemble and Gaussian analytic functions, preprint. [arXiv:1112.2457](https://arxiv.org/abs/1112.2457)
- [31] A. Ledoan, M. Merkli, and S. Starr, A universality property of Gaussian Analytic Functions. *J. Theor. Probab.*, **25** (2012), 496–504.
- [32] J. W. Lindeberg, Eine neue Herleitung des Exponentialgesetzes in der Wahrscheinlichkeit-srechnung. *Math. Z.* 15 (1922), no. 1, 211–225.
- [33] J. E. Littlewood, A. C. Offord, On the distribution of the zeros and a-values of a random integral function. I., *J. Lond. Math. Soc.*, **20** (1945), 120–136.
- [34] J. E. Littlewood and A. C. Offord, On the number of real roots of a random algebraic equation. II. *Proc. Cambridge Philos. Soc.* **35**, (1939), 133–148.
- [35] J. E. Littlewood and A. C. Offord, On the number of real roots of a random algebraic equation. III. *Rec. Math. [Mat. Sbornik] N.S.* **54**, (1943), 277–286.
- [36] J. E. Littlewood, A. C. Offord, On the distribution of the zeros and a-values of a random integral function. II., *Ann. Math.* **49** (1948), 885–952. Errata, **50** (1949), 990–991.
- [37] B. F. Logan, L. A. Shepp, Real zeros of random polynomials. *Proc. London Math. Soc.* **18** (1968), 29–35.
- [38] B. F. Logan, L. A. Shepp, *Real zeros of random polynomials. II.* *Proc. London Math. Soc.* **18** (1968), 308–314.
- [39] N. B. Maslova, The variance of the number of real roots of random polynomials. *Teor. Veroyatnost. i Primenen.* **19** (1974), 36–51.
- [40] N. B. Maslova, The distribution of the number of real roots of random polynomials. *Theor. Probability Appl.* **19** (1974), 461–473
- [41] M. Mehta, *Random matrices*. Third edition. Pure and Applied Mathematics (Amsterdam), 142. Elsevier/Academic Press, Amsterdam, 2004.
- [42] J. Milnor, Growth of finitely generated solvable groups, *J. Diff. Geom.* **2** (1968), 447–449.
- [43] F. Nazarov, M. Sodin, Correlation functions for random complex zeroes: strong clustering and local universality, *Comm. Math. Phys.* **310** (2012), no. 1, 75–98.
- [44] H. Nguyen, V. Vu, Optimal Inverse Littlewood-Offord theorems, *Adv. Math.* **226** (2011), no. 6, 5298–5319.
- [45] V. Paulauskas and A. Raskauskas, *Approximation theory in central limit theorem*, Kluwer Academic Publishers, 2009.
- [46] R. E. A. C. Paley, N. Wiener, *Fourier transforms in the complex domain*. AMS Colloquium Publications, 1934.
- [47] S. O. Rice, Mathematical analysis of random noise. *Bell System Tech. J.* **23** (1944), 282–332; *Ibid* **24** (1945), 46–156.

- [48] Y. Shalom, T. Tao, A finitary version of Gromov’s polynomial growth theorem, *GAF* **20** (2010), no. 6, 1502–1547.
- [49] L. Shepp, R. J. Vanderbei, The complex zeros of random polynomials. *Trans. AMS*, 347:4365–4383, 1995.
- [50] B. Shiffman, S. Zelditch, Distribution of zeros of random and quantum chaotic sections of positive line bundles, *Comm. Math. Phys.*, **200** (1999), 661–683.
- [51] B. Shiffman, S. Zelditch, Equilibrium distribution of zeros of random polynomials, *Int. Math. Res. Not.* **1** (2003), 25–49.
- [52] M. Shub, S. Smale, Complexity of Bezout’s theorem II: Volumes and Probabilities, in Computational Algebraic Geometry, edited by F. Eyssette and A. Galligo, Progress in Mathematics, 109, Birkhauser, 1993, 267–285.
- [53] M. Sodin, Zeroes of Gaussian analytic functions, *Math. Res. Lett.* **7** (2000), no. 4, 371–381.
- [54] S. Starr, Universality of correlations for random analytic functions, *Entropy and the quantum II*, 135144, Contemp. Math., 552, Amer. Math. Soc., Providence, RI, 2011.
- [55] Stevens, D. C. The average number of real zeros of a random polynomial. *Comm. Pure Appl. Math.* **22** (1969), 457–477.
- [56] B. Shiffman, S. Zelditch, Equilibrium distribution of zeros of random polynomials. *Int. Math. Res. Not.* 2003, no. 1, 25–49.
- [57] T. Tao and V. Vu, Random matrices: universality of ESDs and the circular law. With an appendix by Manjunath Krishnapur, *Ann. Probab.* **38** (2010), no. 5, 2023–2065.
- [58] T. Tao and V. Vu, *Random matrices: universality of local spectral statistics of non-Hermitian matrices, to appear in Annals of Probability.*
- [59] T. Tao and V. Vu, *Random matrices: universality of local spectral statistics*, Acta Math. **206** (2011), no. 1, 127–204.
- [60] Todhunter, I. A history of the mathematical theory of probability. Stechert, New York, 1931.
- [61] Y. Wang, Bounds on the average number of real roots of a random algebraic equation. *Chinese Ann. Math. Ser. A* **4** (1983), 601–605.
- [62] J. E. Wilkins, *An asymptotic expansion for the expected number of real zeros of a random polynomial.* Proc. Amer. Math. Soc. **103** (1988), 1249–1258.
- [63] J. Wolf, *Growth of finitely generated solvable groups and curvature of Riemannian manifolds*, J. Diff. Geom. **2** (1968), 421–446.

DEPARTMENT OF MATHEMATICS, UCLA, LOS ANGELES CA 90095-1555

E-mail address: tao@math.ucla.edu

DEPARTMENT OF MATHEMATICS, YALE, NEW HAVEN, CT 06520

E-mail address: van.vu@yale.edu