# On the strong metric generators of strong product graphs

Dorota Kuziak<sup>(1)</sup>, Ismael G. Yero<sup>(2)</sup> and Juan A. Rodríguez-Velázquez<sup>(1)</sup>

 $(1)$ Departament d'Enginyeria Informàtica i Matemàtiques,

Universitat Rovira i Virgili, Av. Pa¨ısos Catalans 26, 43007 Tarragona, Spain.

dorota.kuziak@urv.cat, juanalberto.rodriguez@urv.cat

 $(2)$ Departamento de Matemáticas, Escuela Politécnica Superior de Algeciras

Universidad de Cádiz, Av. Ramón Puyol s/n, 11202 Algeciras, Spain.

ismael.gonzalez@uca.es

#### Abstract

Let G be a connected graph. A vertex  $w \in V(G)$  strongly resolves two vertices  $u, v \in V(G)$ if there exists some shortest  $u-w$  path containing v or some shortest  $v-w$  path containing u. A set S of vertices is a strong metric generator for G if every pair of vertices of G is strongly resolved by some vertex of S. The smallest cardinality of a strong metric generator for G is called the strong metric dimension of  $G$ . It is well known that the problem of computing this invariant is NP-hard. In this paper we study the problem of finding exact values or sharp bounds for the strong metric dimension of strong product graphs and express these in terms of invariants of the factor graphs.

*Keywords:* Strong metric dimension; strong metric basis; strong metric generator; strong product graphs.

*AMS Subject Classification Numbers:* 05C12; 05C69; 05C76.

## 1 Introduction

A *generator* of a metric space is a set S of points in the space with the property that every point of the space is uniquely determined by its distances from the elements of S. Given a simple and connected graph  $G = (V, E)$ , we consider the metric  $d_G : V \times V \to \mathbb{R}^+$ , where  $d_G(x, y)$  is the length of a shortest path between x and y.  $(V, d_G)$  is clearly a metric space. A vertex  $v \in V$  is said to distinguish two vertices x and y if  $d_G(v, x) \neq d_G(v, y)$ . A set  $S \subset V$  is said to be a *metric generator* for G if any pair of vertices of G is distinguished by some element of S. A minimum generator is called a *metric basis*, and its cardinality the *metric dimension* of G. Motivated by the problem of uniquely determining the location of an intruder in a network, the concept of metric dimension of a graph was introduced by Slater in [\[10\]](#page-11-0), where the metric generators were called *locating sets*. The concept of metric dimension of a graph was introduced independently by Harary and Melter in [\[4\]](#page-11-1), where metric generators were called *resolving sets*.

In [\[9\]](#page-11-2) a more restricted invariant than the metric dimension is introduced. A vertex  $w \in V(G)$ *strongly resolves* two vertices  $u, v \in V(G)$  if  $d_G(w, u) = d_G(w, v) + d_G(v, u)$  or  $d_G(w, v) = d_G(w, u) + d_G(w, u)$ 

 $d_G(u, v)$ , *i.e.*, there exists some shortest  $w - u$  path containing v or some shortest  $w - v$  path containing u. A set S of vertices in a connected graph G is a *strong metric generator* for G if every two vertices of  $G$  are strongly resolved by some vertex of  $S$ . The smallest cardinality of a strong metric generator of G is called the *strong metric dimension* and is denoted by  $dim_s(G)$ . A *strong metric basis* of G is a strong metric generator for G of cardinality  $\dim_s(G)$ .

Given a simple graph  $G = (V, E)$ , we denote two adjacent vertices u, v by  $u \sim v$  and, in this case, we say that uv is an edge of G, *i.e.*,  $uv \in E$ . For a vertex  $v \in V$ , the set  $N(v) = \{u \in V :$  $u \sim v$ } is the open neighborhood of v and the set  $N[v] = N(v) \cup \{v\}$  is the closed neighborhood of v. The diameter of G is defined as  $D(G) = \max_{u,v \in V} \{d(u,v)\}\)$ . The vertex  $x \in V$  is diametral in G if there exists  $y \in V$  such that  $d_G(x, y) = D(G)$ . We say that G is 2-antipodal if for each vertex  $x \in V$  there exists exactly one vertex  $y \in V$  such that  $d_G(x, y) = D(G)$ .

A set S of vertices of G is a *vertex cover* of G if every edge of G is incident with at least one vertex of S. The *vertex cover number* of G, denoted by  $\alpha(G)$ , is the smallest cardinality of a vertex cover of G. We refer to an  $\alpha(G)$ -set in a graph G as a vertex cover set of cardinality  $\alpha(G)$ . A vertex u of G is *maximally distant* from v if for every  $w \in N_G(u)$ ,  $d_G(v, w) \leq d_G(u, v)$ . If u is maximally distant from v and v is maximally distant from u, then we say that u and v are *mutually maximally distant.* The *boundary* of  $G = (V, E)$  is defined as

 $\partial(G) = \{u \in V : \text{ exists } v \in V \text{ such that } u, v \text{ are mutually maximally distant}\}.$ 

In [\[6\]](#page-11-3) was first presented a relationship between the boundary of a graph and its strong metric generators. Also, in  $[6]$  the authors defined the concept of *strong resolving graph*  $G_{SR}$  of a graph G like a graph with vertex set  $V(G_{SR}) = V$  and two vertices u, v are adjacent in  $G_{SR}$  if and only if  $u$  and  $v$  are mutually maximally distant in  $G$ .

We recall that the *Cartesian product* of two graphs  $G = (V_1, E_1)$  and  $H = (V_2, E_2)$  is the graph  $G\Box H = (V, E)$ , such that  $V = V_1 \times V_2$  and two vertices  $(a, b) \in V$  and  $(c, d) \in V$  are adjacent in  $G\Box H$  if and only if either

- $a = c$  and  $bd \in E_2$ , or
- $ac \in E_1$  and  $b = d$ .

The *strong product* of two graphs  $G = (V_1, E_1)$  and  $H = (V_2, E_2)$  is the graph  $G \boxtimes H = (V, E)$ , such that  $V = V_1 \times V_2$  and two vertices  $(a, b) \in V$  and  $(c, d) \in V$  are adjacent in  $G \boxtimes H$  if and only if either

- $a = c$  and  $bd \in E_2$ , or
- $ac \in E_1$  and  $b = d$ , or
- $ac \in E_1$  and  $bd \in E_2$ .

The *lexicographic product* of two graphs  $G = (V_1, E_1)$  and  $H = (V_2, E_2)$  is the graph  $G \circ H$ with the vertex set  $V = V_1 \times V_2$  and two vertices  $(a, b) \in V$  and  $(c, d)$  are adjacent if either

- $ac \in E_1$ , or
- $a = c$  and  $bd \in E_2$ .

The *Cartesian sum* of two graphs  $G = (V_1, E_1)$  and  $H = (V_2, E_2)$ , denoted by  $G \oplus H$ , has as the vertex set  $V = V_1 \times V_2$  and two vertices  $(a, b) \in V$  and  $(c, d) \in V$  are adjacent in  $G \oplus H$  if and only if  $ac \in E_1$  or  $bd \in E_2$ . This notion of graph product was introduced by Ore [\[7\]](#page-11-4). The Cartesian sum is also known as the disjunctive product [\[8\]](#page-11-5).

Let  $G = (V, E)$  and  $G' = (V', E')$  be two graphs. If  $V' \subseteq V$  and  $E' \subseteq E$ , then G' is a subgraph of G and we denote that by  $G' \sqsubseteq G$ .

In this article we are interested in the study of strong metric generators of strong product graphs. It was shown in [\[6\]](#page-11-3) that the problem of computing  $dim_s(G)$  is NP-hard. This suggests obtaining exact values of the strong metric dimension for special classes of strong product graphs or finding sharp bounds on this invariant.

#### 2 Results

*-*

Oellermann and Peters-Fransen [\[6\]](#page-11-3) showed that the problem of finding the strong metric dimension of a graph G can be transformed into the problem of computing the vertex cover number of  $G_{SR}$ .

<span id="page-2-0"></span>Theorem 1. [\[6\]](#page-11-3) *For any connected graph* G*,*

$$
dim_s(G) = \alpha(G_{SR}).
$$

Recall that the largest cardinality of a set of vertices of G, no two of which are adjacent, is called the *independence number* of G and is denoted by  $\beta(G)$ . We refer to an  $\beta(G)$ -set in a graph G as an independent set of cardinality  $\beta(G)$ . The following well-known result, due to Gallai, states the relationship between the independence number and the vertex cover number of a graph.

<span id="page-2-1"></span>Theorem 2. (Gallai's theorem) *For any graph* G *of order* n*,*

$$
\alpha(G) + \beta(G) = n.
$$

Thus, for any graphs G and H of order  $n_1$  $n_1$  and  $n_2$ , respectively, by using Theorems 1 and [2,](#page-2-1) we immediately obtain that

<span id="page-2-3"></span>
$$
dim_s(G \boxtimes H) = n_1 \cdot n_2 - \beta((G \boxtimes H)_{SR})
$$
\n(1)

The following basic remark leads to a corollary about the neighborhood of a vertex in the strong product graph  $G \boxtimes H$ , which will be useful to present our results.

**Remark 3.** Let G and H be two graphs. For every  $u \in V(G)$  and  $v \in V(H)$ 

$$
N_{G\boxtimes H}[(u,v)] = N_G[u] \times N_H[v].
$$

<span id="page-2-2"></span>**Corollary 4.** Let G and H be two graphs and let  $u, u' \in V(G)$  and  $v, v' \in V(H)$ . The following *assertion hold.*

- (i) If  $(u', v') \in N_{G \boxtimes H}(u, v)$ , then  $u' \in N_G[u]$  and  $v' \in N_G[v]$ .
- (ii) If  $u' \in N_G(u)$  and  $v' \in N_G(v)$ , then  $(u', v') \in N_{G \boxtimes H}(u, v)$ .

The following result about the boundary of strong product graphs was presented in [\[1\]](#page-11-6). Nevertheless in such a paper the authors are more interested into the cardinality of the boundary  $\partial(G \boxtimes H)$  than into how the subgraph induced by boundary looks like.

**Theorem 5.** [\[1\]](#page-11-6) *For any graphs* G and H,  $\partial(G \boxtimes H) = (\partial(G) \times V(H)) \cup (V(G) \times \partial(H))$ *.* 

In the next lemma we pretend to describe the structure of the strong resolving graph of  $G \boxtimes H$ .

<span id="page-3-0"></span>Lemma 6. *Let* G *and* H *be two connected nontrivial graphs. Let* u, x *be two vertices of* G *and let* v, y be two vertices of H. Then  $(u, v)$  and  $(x, y)$  are mutually maximally distant vertices in  $G \boxtimes H$ *if and only if one of the following conditions holds:*

- (i) u, x *are mutually maximally distant in* G *and* v, y *are mutually maximally distant in* H*;*
- (ii) u, x are mutually maximally distant in G and  $v = y$ ;
- (iii) v, y are mutually maximally distant in H and  $u = x$ ;
- (iv) u, x are mutually maximally distant in G and  $d_G(u, x) > d_H(v, y)$ ;
- (v) v, y are mutually maximally distant in H and  $d_G(u, x) < d_H(v, y)$ .

*Proof.* (Sufficiency) Let  $(u', v') \in N_{G \boxtimes H}(u, v)$  and  $(x', y') \in N_{G \boxtimes H}(x, y)$ . By Corollary [4](#page-2-2) we have  $u' \in N_G[u], x' \in N_G[x], v' \in N_H[v]$  and  $y' \in N_H[y]$ .

(i) If  $u, x$  are mutually maximally distant in G and  $v, y$  are mutually maximally distant in H, then

$$
d_{G\boxtimes H}((u',v'),(x,y)) = \max\{d_G(u',x),d_H(v',y)\} \le \max\{d_G(u,x),d_H(v,y)\} = d_{G\boxtimes H}((u,v),(x,y))
$$

and

$$
d_{G\boxtimes H}((u, v), (x, 'y')) = \max\{d_G(u, x'), d_H(v, y')\} \le \max\{d_G(u, x), d_H(v, y)\} = d_{G\boxtimes H}((u, v), (x, y)).
$$

Thus,  $(u, v)$  and  $(x, y)$  are mutually maximally distant vertices in  $G \boxtimes H$ .

(ii) If  $u, x$  are mutually maximally distant in G and  $v = y$ , then

$$
d_{G\boxtimes H}((u',v'),(x,y)) = \max\{d_G(u',x),d_H(v',y)\} = d_G(u',x) \leq d_G(u,x) = d_{G\boxtimes H}((u,v),(x,y))
$$

and

$$
d_{G\boxtimes H}((u, v), (x', y')) = \max\{d_G(u, x'), d_H(v, y')\} = d_G(u, x') \leq d_G(u, x) = d_{G\boxtimes H}((u, v), (x, y)).
$$

Thus,  $(u, v)$  and  $(x, y)$  are mutually maximally distant vertices in  $G \boxtimes H$ .

(iii) By using analogous procedure to (ii) we can show that if  $u = x$  and  $v, y$  are mutually maximally distant in G, then  $(u, v)$  and  $(x, y)$  are mutually maximally distant vertices in  $G \boxtimes H$ .

(iv) If u, x are mutually maximally distant in G and  $d_G(u, x) > d_H(v, y)$ , then

$$
d_{G \boxtimes H}((u', v'), (x, y)) = \max\{d_G(u', x), d_H(v', y)\}
$$
  
\n
$$
\leq \max\{d_G(u, x), d_H(v, y) + 1\}
$$
  
\n
$$
= \max\{d_G(u, x), d_H(v, y)\}
$$
  
\n
$$
= d_{G \boxtimes H}((u, v), (x, y))
$$

$$
d_{G \boxtimes H}((u, v), (x, 'y')) = \max\{d_G(u, x'), d_H(v, y')\}
$$
  
\n
$$
\leq \max\{d_G(u, x), d_H(v, y) + 1\}
$$
  
\n
$$
= \max\{d_G(u, x), d_H(v, y)\}
$$
  
\n
$$
= d_{G \boxtimes H}((u, v), (x, y)).
$$

Thus,  $(u, v)$  and  $(x, y)$  are mutually maximally distant vertices in  $G \boxtimes H$ .

(v) By using analogous procedure as in (iv) we can show that if  $v, y$  are mutually maximally distant in H and  $d_G(u, x) < d_H(v, y)$ , then  $(u, v)$  and  $(x, y)$  are mutually maximally distant vertices in  $G \boxtimes H$ .

(Necessity) Let  $(u, v)$  and  $(x, y)$  be two mutually maximally distant vertices in  $G \boxtimes H$ . Let  $u' \in N_G(u), x' \in N_G(x), v' \in N_H(v)$  and  $y' \in N_H(y)$ . Notice that, by Corollary [4](#page-2-2)  $(u', v') \in$  $N_{G\boxtimes H}(u, v)$  and  $(x', y') \in N_{G\boxtimes H}(x, y)$ . So, we have that

$$
d_{G\boxtimes H}((u,v),(x,y)) \geq d_{G\boxtimes H}((u',v'),(x,y))
$$

and

$$
d_{G\boxtimes H}((u, v), (x, y)) \ge d_{G\boxtimes H}((u, v), (x', y')).
$$

We differentiate two cases.

Case 1.  $d_G(u, x) \ge d_H(v, y)$ . Hence,  $d_{G \boxtimes H}(u, v), (x, y) = \max\{d_G(u, x), d_H(v, y)\} = d_G(u, x)$ . Thus,

$$
d_G(u, x) \ge \max\{d_G(u', x), d_H(v', y)\}
$$

and

 $d_G(u, x) \ge \max\{d_G(u, x'), d_H(v, y')\}.$ 

So, we obtain four inequalities:

<span id="page-4-0"></span>
$$
d_G(u, x) \ge d_G(u', x),\tag{2}
$$

<span id="page-4-2"></span>
$$
d_G(u,x) \ge d_H(v',y),\tag{3}
$$

<span id="page-4-1"></span>
$$
d_G(u, x) \ge d_G(u, x'),\tag{4}
$$

<span id="page-4-3"></span>
$$
d_G(u, x) \ge d_H(v, y'). \tag{5}
$$

From [\(2\)](#page-4-0) and [\(4\)](#page-4-1) we have, that u and x are mutually maximally distant in G. If v and y are mutually maximally distant in  $H$ , then (i) holds and, if  $v = y$ , then (ii) holds. Suppose that there exists a vertex  $v'' \in N_H(v)$  such that  $d_H(v'', y) > d_H(v, y)$  or there exists a vertex  $y'' \in N_H(y)$ such that  $d_H(v, y'') > d_H(v, y)$ . In such a case,

$$
d_H(v'', y) \ge d_H(v, y) + 1 \tag{6}
$$

or

$$
d_H(v, y'') \ge d_H(v, y) + 1. \tag{7}
$$

Since  $v'' \in N_H(v)$ , for any  $u'' \in N_G(u)$  we have  $(u'', v'') \in N_{G \boxtimes H}(u, v)$  and following the above procedure, taking  $(u'', v'')$  instead of  $(u', v')$  we obtain two inequalities equivalent to [\(3\)](#page-4-2) and [\(5\)](#page-4-3). Thus,

$$
d_G(u, x) \ge d_H(v'', y) > d_H(v, y)
$$
\n(8)

and

$$
d_G(u, x) \ge d_H(v, y'') > d_H(v, y). \tag{9}
$$

So, u, x are mutually maximally distant in G and  $d_G(u, x) > d_H(v, y)$ . Hence, (iv) is satisfied.

Case 2.  $d_G(u, x) < d_H(v, y)$ . By using analogous procedure we can prove that  $v, y$  are mutually maximally distant in H and  $u = x$  or  $d_G(u, x) < d_H(v, y)$ , showing that (iii) and (v) hold. Therefore, the result follows.  $\Box$ 

Notice that Lemma [6](#page-3-0) leads to the following relationship.

<span id="page-5-4"></span>Theorem 7. *For any connected graphs* G *and* H*,*

$$
G_{SR} \boxtimes H_{SR} \sqsubseteq (G \boxtimes H)_{SR} \sqsubseteq G_{SR} \oplus H_{SR}.
$$

*Proof.* Notice that  $V(G_{SR} \boxtimes H_{SR}) = V((G \boxtimes H)_{SR}) = V(G_{SR} \oplus H_{SR}) = V_1 \times V_2$ . Let  $(u, v)$  and  $(x, y)$  be two vertices adjacent in  $G_{SR} \boxtimes H_{SR}$ . So, either

- $u = x$  and  $vy \in E(H_{SR})$ , or
- $ux \in E(G_{SR})$  and  $v = y$ , or
- $ux \in E(G_{SR})$  and  $vy \in E(H_{SR})$ .

Hence, by using respectively the condition (iii), (ii) and (i) of Lemma [6](#page-3-0) we have that  $(u, v)$  and  $(x, y)$  are also adjacent in  $(G \boxtimes H)_{SR}$ .

Now, let  $(u', v')$  and  $(x', y')$  be two vertices adjacent in  $(G \boxtimes H)_{SR}$ . From Lemma [6](#page-3-0) we obtain that  $u'x' \in E(G_{SR})$  or  $v'y' \in E(H_{SR})$ . Thus,  $(u', v')$  and  $(x', y')$  are also adjacent in  $G_{SR} \oplus H_{SR}$ .

<span id="page-5-1"></span>Corollary 8. *For any connected graphs* G *and* H*,*

$$
\beta(G_{SR} \boxtimes H_{SR}) \ge \beta((G \boxtimes H)_{SR}) \ge \beta(G_{SR} \oplus H_{SR}).
$$

In order to better understand how the strong resolving graph  $(G \boxtimes H)_{SR}$  looks like, by using Lemma [6,](#page-3-0) we prepare a kind of "graphical representation" of  $(G \boxtimes H)_{SR}$  which we present in Figure [1.](#page-6-0) According to the conditions (i), (ii) and (iii) of Lemma [6](#page-3-0) the solid lines represents those edges of  $(G \boxtimes H)_{SR}$  which always exists. Also, from the conditions (iv) and (v) of Lemma [6,](#page-3-0) two vertices belonging to different rounded rectangles with identically filled areas could be adjacent or not in  $(G \boxtimes H)_{SR}$ .

The following three known results will be useful for our purposes.

<span id="page-5-2"></span>Theorem 9. [\[5\]](#page-11-7) *For any graphs* G *and* H*,*

$$
\beta(G) \cdot \beta(H) \le \beta(G \boxtimes H) \le \beta(G \square H).
$$

<span id="page-5-3"></span>Theorem 10. (Vizing's theorem) *For any graphs* G *and* H*,*

 $\beta(G\Box H) \le \min\{\beta(G)|V(H)|, \beta(H)|V(G)|\}.$ 

<span id="page-5-0"></span>Theorem 11. [\[2\]](#page-11-8) *For any graphs* G *and* H*,*

$$
\beta(G \circ H) = \beta(G) \cdot \beta(H).
$$



<span id="page-6-0"></span>Figure 1: Sketch of a representation of a strong resolving graph  $(G \boxtimes H)_{SR}$ .

Next we present a lemma about the independence number of Cartesian sum graphs.

<span id="page-6-1"></span>Lemma 12. *For any graphs* G *and* H*,*

$$
\beta(G \oplus H) = \beta(G) \cdot \beta(H).
$$

*Proof.* Let A be a  $\beta(G)$ -set and let B be a  $\beta(H)$ -set. Let  $S = A \times B$ . We will show that S is an independent set in  $G \oplus H$ . Notice that if  $|S| = 1$ , then G, H and  $G \oplus H$  are complete graphs and the result follows. Now, we consider case  $|S| > 1$ . Let  $(u, v), (x, y) \in S$ . Suppose that  $(u, v)$  and  $(x, y)$  are adjacent in  $G \oplus H$ . If  $u = x$ , then v and y are adjacent in H, which is a contradiction. If  $v = y$ , then analogously we have a contradiction. Now, if  $u \neq x$  and  $v \neq y$ , then u and x are adjacent in G or v and y are adjacent in H, which is a contradiction. Thus, S is independent and we have that  $\beta(G \oplus H) \geq \beta(G) \cdot \beta(H)$ .

On the other hand, from the definitions of Cartesian sum and lexicographic product we have that  $V(G \circ H) = V(G \oplus H)$  and  $G \circ H \subseteq G \oplus H$ . Thus,  $\beta(G \circ H) \geq \beta(G \oplus H)$  and by using Theorem [11](#page-5-0) we have that  $\beta(G \oplus H) \leq \beta(G) \cdot \beta(H)$ . Therefore, the result follows.  $\Box$ 

<span id="page-6-2"></span>**Theorem 13.** Let G and H be two connected nontrivial graphs of order  $n_1$ ,  $n_2$ , respectively. Then

 $\max\{n_2 \cdot dim_s(G), n_1 \cdot dim_s(H)\} \leq dim_s(G \boxtimes H) \leq n_2 \cdot dim_s(G) + n_1 \cdot dim_s(H) - dim_s(G) \cdot dim_s(H).$ 

*Proof.* By using Corollary [8](#page-5-1) we have that  $\beta(G_{SR} \boxtimes H_{SR}) \geq \beta((G \boxtimes H)_{SR})$ . Hence, from equality

[\(1\)](#page-2-3), Theorem [9](#page-5-2) and Theorem [10](#page-5-3) we obtain

$$
dim_s(G \boxtimes H) = n_1 \cdot n_2 - \beta((G \boxtimes H)_{SR})
$$
  
\n
$$
\ge n_1 \cdot n_2 - \beta(G_{SR} \boxtimes H_{SR})
$$
  
\n
$$
\ge n_1 \cdot n_2 - \beta(G_{SR} \Box H_{SR})
$$
  
\n
$$
\ge n_1 \cdot n_2 - \min\{n_2 \cdot \beta(G_{SR}), n_1 \cdot \beta(H_{SR})\}
$$
  
\n
$$
= \max\{n_2(n_1 - \beta(G_{SR})), n_1(n_2 - \beta(H_{SR}))\}
$$
  
\n
$$
= \max\{n_2 \cdot dim_s(G), n_1 \cdot dim_s(H)\}.
$$

On the other hand, from Corollary [8](#page-5-1) it follows  $\beta((G \boxtimes H)_{SR}) \geq \beta(G_{SR} \oplus H_{SR})$ . So, by using [\(1\)](#page-2-3) and Lemma [12](#page-6-1) we have

$$
dim_s(G \boxtimes H) = n_1 \cdot n_2 - \beta((G \boxtimes H)_{SR})
$$
  
\n
$$
\leq n_1 \cdot n_2 - \beta(G_{SR} \oplus H_{SR})
$$
  
\n
$$
= n_1 \cdot n_2 - \beta(G_{SR}) \cdot \beta(H_{SR})
$$
  
\n
$$
= n_1 \cdot n_2 - (n_1 - dim_s(G)) \cdot (n_2 - dim_s(H))
$$
  
\n
$$
= n_2 \cdot dim_s(G) + n_1 \cdot dim_s(H) - dim_s(G) \cdot dim_s(H).
$$

We define a  $\mathcal{C}\text{-}graph$  as a graph G whose vertex set can be partitioned into  $\beta(G)$  cliques. Notice that there are several graphs which are  $\mathcal{C}$ -graphs. For instance, we emphasize the following cases: complete graphs and cycles of even order. In order to prove the next result we also need to introduce the following notation. Given two graphs  $G = (V_1, E_1), H = (V_2, E_2)$  and a subset X of vertices of  $G \boxtimes H = (V, E)$ , the *projections* of X over the graphs G and H, respectively, are the following ones

$$
P_G(X) = \{ u \in V_1 : (u, v) \in X, \text{ for some } v \in V_2 \},
$$
  

$$
P_H(X) = \{ v \in V_2 : (u, v) \in X, \text{ for some } u \in V_1 \}.
$$

<span id="page-7-0"></span>Lemma 14. *For any* C*-graph* G *and any graph* H*,*

$$
\beta(G \boxtimes H) = \beta(G) \cdot \beta(H).
$$

*Proof.* Let  $A_1, A_2, ..., A_{\beta(G)}$  be a partition of  $V(G)$  such that  $A_i$  is a clique for every  $i \in \{1, 2, ..., \beta(G)\}.$ Let S be an  $\beta(G \boxtimes H)$ -set and let  $S_i = S \cap (A_i \times V_2)$  for  $i \in \{1, 2, ..., \beta(G)\}\)$ . First we will show that  $P_H(S_i)$  is an independent set in H. If  $|P_H(S_i)| = 1$ , then  $P_H(S_i)$  is an independent set in H. If  $|P_H(S_i)| \geq 2$ , then for any two vertices  $x, y \in P_H(S_i)$  there exist  $u, v \in A_i$  such that  $(u, x), (v, y) \in S_i$ . We suppose that  $x \sim y$ . If  $u = v$ , then  $(u, x) \sim (v, y)$ , which is a contradiction. Thus,  $u \neq v$ . Since  $(u, x) \nsim(v, y)$ , we have that  $u \nsim v$ , which is a contradiction with the fact that  $A_i$  is a clique. Therefore, for every  $i \in \{1, 2, ..., \beta(G)\}\$ the projection  $P_H(S_i)$  is an independent set in H and  $\beta(H) \geq |P_H(S_i)|$ .

Now, if  $|S_i| > |P_H(S_i)|$  for some  $i \in \{1, 2, ..., \beta(G)\}\$ , then there exists a vertex  $z \in P_H(S_i)$  and two different vertices  $a, b \in A_i$  such that  $(a, z), (b, z) \in S_i$ , and this is a contradiction with the facts that  $A_i$  is a clique and  $S_i$  is an independent set. Thus,  $|S_i| = |P_H(S_i)|$ ,  $i \in \{1, 2, ..., \beta(G)\}\$ , and we have the following

$$
\beta(G \boxtimes H) = |S| = \sum_{i=1}^{\beta(G)} |S_i| = \sum_{i=1}^{\beta(G)} |P_H(S_i)| \leq \beta(G) \cdot \beta(H).
$$

Therefore, by using Theorem [9](#page-5-2) we conclude the proof.

<span id="page-8-2"></span>**Theorem 15.** Let G and H be two connected nontrivial graphs of order  $n_1$ ,  $n_2$ , respectively. If GSR *is a* C*-graph, then*

$$
dim_s(G \boxtimes H) = n_2 \cdot dim_s(G) + n_1 \cdot dim_s(H) - dim_s(G) \cdot dim_s(H).
$$

*Proof.* By using Corollary [8](#page-5-1) we have that  $\beta(G_{SR} \boxtimes H_{SR}) \geq \beta((G \boxtimes H)_{SR})$ . Hence, from equality [\(1\)](#page-2-3) and Lemma [14](#page-7-0) we have

$$
dim_s(G \boxtimes H) = n_1 \cdot n_2 - \beta((G \boxtimes H)_{SR})
$$
  
\n
$$
\ge n_1 \cdot n_2 - \beta(G_{SR} \boxtimes H_{SR})
$$
  
\n
$$
= n_1 \cdot n_2 - \beta(G_{SR}) \cdot \beta(H_{SR})
$$
  
\n
$$
= n_1 \cdot n_2 - (n_1 - dim_s(G)) \cdot (n_2 - dim_s(H))
$$
  
\n
$$
= n_2 \cdot dim_s(G) + n_1 \cdot dim_s(H) - dim_s(G) \cdot dim_s(H).
$$

The result now follows from Theorem [13.](#page-6-2)

A *cut vertex* in a graph is a vertex whose removal increases the number of connected component and a *simplicial vertex* is a vertex v such that the subgraph induced by  $N[v]$  is isomorphic to a complete graph. Also, a *block* is a maximal biconnected subgraph of the graph. Now, let  $\mathfrak{F}$  be the family of sequences of connected graphs  $G_1, G_2, ..., G_k, k \geq 2$ , such that  $G_1$  is a complete graph  $K_{n_1}, n_1 \geq 2$ , and  $G_i, i \geq 2$ , is obtained recursively from  $G_{i-1}$  by adding a complete graph  $K_{n_i}$ ,  $n_i \geq 2$ , and identifying a vertex of  $G_{i-1}$  with a vertex in  $K_{n_i}$ .

From this point we will say that a connected graph G is a *generalized tree*<sup>[1](#page-8-0)</sup> if and only if there exists a sequence  $\{G_1, G_2, ..., G_k\} \in \mathfrak{F}$  such that  $G_k = G$  for some  $k \geq 2$ . Notice that in these generalized trees every vertex is either, a cut vertex or a simplicial vertex. Also, every complete graph used to obtain the generalized tree is a block of the graph. Note that if every  $G_i$ is isomorphic to  $K_2$ , then  $G_k$  is a tree, justifying the terminology used.

At next we give examples of graphs for which its strong resolving graphs are C-graphs.

- $(K_n)_{SR}$  is isomorphic to  $K_n$ .
- For any complete k-partite graph such that at least all but one  $p_i \geq 2$ ,  $i \in \{1, 2, ..., k\}$ ,  $(K_{p_1,p_2,\dots,p_k})_{SR}$  is isomorphic to the graph  $\bigcup_{i=1}^k K_{p_i}$ .
- If G is a generalized tree of order n and c cut vertices, then  $G_{SR}$  is isomorphic to the graph  $K_{n-c} \cup (\bigcup_{i=1}^{c} K_1).$
- For any [2](#page-8-1)-antipodal<sup>2</sup> graph G of order n,  $G_{SR}$  is isomorphic to the graph  $\bigcup_{i=1}^{\frac{n}{2}} K_2$ .
- For any grid graph,  $(P_n \Box P_r)_{SR}$  is isomorphic to the graph  $K_4 \cup (\bigcup_{i=1}^{n \cdot r-4} K_1)$ .

By using the above examples and Theorem [15](#page-8-2) we have the following corollary.

<span id="page-8-3"></span>**Corollary 16.** Let G and H be two connected nontrivial graphs of order  $n_1$  and  $n_2$ , respectively.

 $\Box$ 

<sup>&</sup>lt;sup>1</sup>In some works those graphs are called block graphs.

<span id="page-8-1"></span><span id="page-8-0"></span><sup>2</sup>Notice that for instance cycles of even order are 2-antipodal graphs.

- (i)  $\dim_s(K_{n_1} \boxtimes H) = n_2(n_1 1) + n_1 \cdot \dim_s(H) (n_1 1)\dim_s(H)$ .
- (ii) *If* G *is a complete* k*-partite graph, then*

$$
dim_s(G \boxtimes H) = n_2(n_1 - k) + n_1 \cdot dim_s(H) - (n_1 - k)dim_s(H).
$$

(iii) *If* G *is a generalized tree with* c *cut vertices, then*

$$
dim_s(G \boxtimes H) = n_2(n_1 - c - 1) + n_1 \cdot dim_s(H) - (n_1 - c - 1)dim_s(H).
$$

*Particularly, if* G *is a tree with* l(G) *leaves, then*

$$
dim_s(G \boxtimes H) = n_2(l(G) - 1) + n_1 \cdot dim_s(H) - (l(G) - 1)dim_s(H).
$$

(iv) *If* G *is a* 2*-antipodal graph, then*

$$
dim_s(G \boxtimes H) = \frac{n_2 \cdot n_1}{2} + n_1 \cdot dim_s(H) - \frac{n_1}{2} \cdot dim_s(H).
$$

(v) *If* G *is a grid graph, then*

$$
dim_s(G \boxtimes H) = 3n_2 + n_1 \cdot dim_s(H) - 3dim_s(H).
$$

Notice that Corollary [16](#page-8-3) (iv) gives the value for the strong metric dimension of  $C_r \boxtimes H$  for any graph H and r even. Next we study separately the strong product graphs  $C_r \boxtimes H$  for any graph  $H$  and  $r$  odd. In order to prove the next result we need to introduce the following notation. We define a  $C_1$ -graph as a graph G whose vertex set can be partitioned into  $\beta(G)$  cliques and one isolated vertex. Notice that cycles with odd order are  $C_1$ -graphs.

<span id="page-9-0"></span>**Lemma 17.** For any  $C_1$ -graph G and any graph H,

$$
\beta(G \boxtimes H) \le \beta(G)(\beta(H) + 1).
$$

*Proof.* Let  $A_1, A_2, ..., A_{\beta(G)}, B$  be a partition of  $V(G)$  such that  $A_i$  is a clique for every  $i \in$  $\{1, 2, ..., \beta(G)\}\$  and  $B = \{b\}$ , where b is isolated vertex. Let S be an  $\beta(G \boxtimes H)$ -set and let  $S_i = S \cap (A_i \times V_2)$  and  $i \in \{1, 2, ..., \beta(G)\}\)$ . Let  $S_B = S \cap (B \times V_2)$ . By using analogous procedures as in proof of Lemma [14](#page-7-0) we can show that for every  $i \in \{1, 2, ..., \beta(G)\}\$ ,  $P_H(S_i)$  is an independent set in H and  $|S_i| = |P_H(S_i)|$ . Moreover, since  $|B| = 1$  we have that  $P_H(S_B)$  is an independent set in H and  $|S_B| = |P_H(S_B)|$ . Thus, we obtain the following

$$
\beta(G \boxtimes H) = |S| = \sum_{i=1}^{\beta(G)} |S_i| + |S_B| = \sum_{i=1}^{\beta(G)} |P_H(S_i)| + |P_H(S_B)| \le \beta(G) \cdot \beta(H) + \beta(H) = \beta(G)(\beta(H) + 1).
$$

□

<span id="page-9-1"></span>**Theorem 18.** Let G and H be two connected nontrivial graphs of order  $n_1$ ,  $n_2$ , respectively. If  $G_{SR}$  *is a*  $C_1$ -graph, then

$$
dim_s(G \boxtimes H) \ge n_1(dim_s(H) - 1) + dim_s(G)(n_2 - dim_s(H) + 1).
$$

*Proof.* By using Corollary [8](#page-5-1) we have that  $\beta(G_{SR} \boxtimes H_{SR}) \geq \beta((G \boxtimes H)_{SR})$ . Hence, from equality [\(1\)](#page-2-3) and Lemma [17](#page-9-0) we have

$$
dim_s(G \boxtimes H) = n_1 \cdot n_2 - \beta((G \boxtimes H)_{SR})
$$
  
\n
$$
\ge n_1 \cdot n_2 - \beta(G_{SR} \boxtimes H_{SR})
$$
  
\n
$$
\ge n_1 \cdot n_2 - \beta(G_{SR})(\beta(H_{SR}) + 1)
$$
  
\n
$$
= n_1 \cdot n_2 - (n_1 - dim_s(G)) \cdot (n_2 - dim_s(H) + 1)
$$
  
\n
$$
= n_1(dim_s(H) - 1) + dim_s(G)(n_2 - dim_s(H) + 1).
$$

Since  $dim_s(C_{2r+1}) = r + 1$ , Theorems [13](#page-6-2) and [18](#page-9-1) lead to the following result.

**Theorem 19.** Let H be a connected nontrivial graphs of order n and  $r \geq 1$ . Then

$$
n(r + 1) + r(\dim_s(H) - 1) \le \dim_s(C_{2r+1} \boxtimes H) \le n(r + 1) + r \cdot \dim_s(H).
$$

The independence number of  $C_{2r+1} \boxtimes C_{2t+1}$  was studied in [\[3\]](#page-11-9). There was presented the following result.

<span id="page-10-0"></span>**Theorem 20.** [\[3\]](#page-11-9) *For*  $1 \leq r \leq t$ *,* 

$$
\beta(C_{2r+1} \boxtimes C_{2t+1}) = r \cdot t + \left\lfloor \frac{r}{2} \right\rfloor.
$$

By using the above result we obtain the following.

**Theorem 21.** *For*  $1 \leq r \leq t$ *,* 

$$
3rt + 2r + 2t + 1 - \left\lfloor \frac{r}{2} \right\rfloor \le \dim_s(C_{2r+1} \boxtimes C_{2t+1}) \le 3rt + 2r + 2t + 1.
$$

*Proof.* By using Theorem [7](#page-5-4) we have that  $G_{SR} \boxtimes H_{SR} \subseteq (G \boxtimes H)_{SR}$ . Thus,  $\beta(G_{SR} \boxtimes H_{SR}) \ge$  $\beta((G \boxtimes H)_{SR})$ . Hence, from equality [\(1\)](#page-2-3) and Theorem [20](#page-10-0) we have

$$
dim_s(C_{2r+1} \boxtimes C_{2t+1}) = (2r+1) \cdot (2t+1) - \beta((C_{2r+1} \boxtimes C_{2t+1})_{SR})
$$
  
\n
$$
\geq (2r+1) \cdot (2t+1) - \beta((C_{2r+1})_{SR} \boxtimes (C_{2t+1})_{SR})
$$
  
\n
$$
= (2r+1) \cdot (2t+1) - \beta(C_{2r+1} \boxtimes C_{2t+1})
$$
  
\n
$$
= (2r+1) \cdot (2t+1) - r \cdot t - \left\lfloor \frac{r}{2} \right\rfloor
$$
  
\n
$$
= 3rt + 2r + 2t + 1 - \left\lfloor \frac{r}{2} \right\rfloor.
$$

The upper bound is direct consequence of Theorem [18.](#page-9-1)

Notice that for  $r = 1$  the lower bound is equal to the upper bound in the above theorem. Thus,  $\dim_s(C_3 \boxtimes C_{2t+1}) = 5t + 3$  for every  $t \geq 1$ .

 $\Box$ 

 $\Box$ 

## <span id="page-11-6"></span>References

- [1] J. Cáceres, C. Hernando, M. Mora, I. M. Pelayo, and M. L. Puertas, Boundary-type sets and product operators in graphs, VII Jornadas de Matemática Discreta y Algorítmica, Castro Urdiales, Cantabria, July 2010. *<http://www.jmda2010.unican.es/actas/Caceres2.pdf>*
- <span id="page-11-9"></span><span id="page-11-8"></span>[2] D. Geller and S. Stahl, The chromatic number and other functions of the lexicographic product, *Journal of Combinatorial Theory* (B) 19 (1975) 87–95.
- <span id="page-11-1"></span>[3] R. S. Hales, Numerical invariants and the strong product of graphs, *Journal of Combinatorial Theory* (B) 15 (1973) 146–155.
- <span id="page-11-7"></span>[4] F. Harary and R. A. Melter, On the metric dimension of a graph, *Ars Combinatoria* 2 (1976) 191–195.
- <span id="page-11-3"></span>[5] P. K. Jha and G. Slutzki, Independence numbers of product graphs, *Applied Mathematics Letters* 7(4) (1994) 91–94.
- <span id="page-11-4"></span>[6] O. R. Oellermann and J. Peters-Fransen, The strong metric dimension of graphs and digraphs, *Discrete Applied Mathematics* 155 (2007) 356–364.
- <span id="page-11-5"></span>[7] O. Ore, Theory of Graphs, American Mathematical Society, 1962.
- <span id="page-11-2"></span>[8] E. Scheinerman and D. Ullman, Fractional Graph Theory, Wiley-Interscience Series in Discrete Mathematics and Optimization, 1997.
- [9] A. Seb˝o and E. Tannier, On metric generators of graphs, *Mathematics of Operations Research* 29 (2) (2004) 383–393.
- <span id="page-11-0"></span>[10] P. J. Slater, Leaves of trees, Proceeding of the 6th Southeastern Conference on Combinatorics, Graph Theory, and Computing, *Congressus Numerantium* 14 (1975) 549–559.