

ABELIAN VARIETIES IN BRILL–NOETHER LOCI

CIRO CILIBERTO, MARGARIDA MENDES LOPES, AND RITA PARDINI

ABSTRACT. In this paper, improving on results in [1, 7], we give the full classification of curves C of genus g such that a Brill–Noether locus $W_d^s(C)$, strictly contained in the jacobian $J(C)$ of C , contains a variety Z stable under translations by the elements of a positive dimensional abelian subvariety $A \subsetneq J(C)$ and such that $\dim(Z) = d - \dim(A) - 2s$, i.e., the maximum possible for such a Z .

CONTENTS

1. Introduction	1
2. Linear series on curves and related subvarieties of symmetric products	3
3. Abelian subvarieties of Brill–Noether loci	7
References	12

1. INTRODUCTION

In [1] the authors posed the problem of studying, and possibly classifying, situations like this:

- (*) C is a smooth, projective, complex curve of genus g , Z is an irreducible r -dimensional subvariety of a Brill–Noether locus $W_d^s(C) \subsetneq J^d(C)$, and Z is stable under translations by the elements of an abelian subvariety $A \subsetneq J(C)$ of dimension $a > 0$ (if so, we will say that Z is A -stable).

Actually in [1] the variety Z is the translate of a positive dimensional proper abelian subvariety of $J(C)$, while the above slightly more general formulation was given in [7].

The motivation for studying (*) resides, among other things, in a theorem of Faltings (see [9]) to the effect that if X is an abelian variety defined over a number field \mathbb{K} , and $Z \subsetneq X$ is a subvariety not containing any translate of a positive dimensional abelian subvariety of X , then the number of rational points of Z over \mathbb{K} is finite. The idea in [1] was to apply Faltings’ theorem to the d -fold symmetric product $C(d)$ of a curve C defined over a number field \mathbb{K} . If C has no positive dimensional linear series of degree d , then $C(d)$ is isomorphic to its Abel–Jacobi image $W_d(C)$ in $J^d(C)$. Thus $C(d)$ has finitely many rational points over \mathbb{K} if $W_d(C)$ does not contain any translate of a positive dimensional abelian subvariety of $J(C)$. The suggestion in [1] is that, if, by contrast, $W_d(C)$ contains the translate of a positive dimensional abelian subvariety of $J(C)$, then C should be *quite special*, e.g., it should admit a map to a curve of lower positive genus (curves of this kind clearly are in situation (*)). This idea was tested in [1], where a number of partial results were proven for low values of d .

Mathematics Subject Classification (2010). Primary 14H40. Secondary: 14H51

The first and third authors are members of GNSAGA of INdAM, and this research was partially supported by the MIUR–PRIN 2010 “Geometria delle varietà algebriche”. The second author is a member of the Center for Mathematical Analysis, Geometry and Dynamical Systems (IST/UTL). This research was partially supported by FCT (Portugal) through program POCTI/FEDER and Project PTDC/MAT/099275/2008.

The problem was taken up in [7], see also [8], where, among other things, it is proven that if (*) holds, then $r + a + 2s \leq d$, and, if in addition $d + r \leq g - 1$, then $r + a + 2s = d$ if and only if:

- (a) there is a degree 2 morphism $\varphi: C \rightarrow C'$, with C' a smooth curve of genus a , such that $A = \varphi^*(J(C'))$ and $Z = W_{d-2a-2s}(C) + \varphi^*(J^{a+s}(C'))$.

In [7] there is also the following example with $(d, s) = (g - 1, 0)$:

- (b) there is an (étale) degree 2 morphism $\varphi: C \rightarrow C'$, with C' a smooth curve of genus $g' = r + 1$, A is the Prym variety of φ and $Z \subset W_{g-1}(C)$ is the connected component of $\varphi_*^{-1}(K_{C'})$ consisting of divisor classes D with $h^0(\mathcal{O}_C(D))$ odd, where $\varphi_*: J^{g-1}(C) \rightarrow J^{g-1}(C')$ is the *norm map*. One has $Z \cong A$, hence $r = a$.

One more family of examples is the following (see Corollary 3.9 below):

- (c) C is hyperelliptic, there is a degree 2 morphism $\varphi: C \rightarrow C'$ with C' a smooth curve of genus a such that $g > 2a + 1$, $A = \varphi^*(J(C'))$, $0 < s < g - 1$ and $Z = \varphi^*(J^a(C')) + W_{d-2s-2a}(C) + W_{2s}^s(C)$ (notice that $W_{2s}^s(C)$ is a point).

The aforementioned result in [7] goes exactly in the direction indicated in [1]. The unfortunate feature of it is the hypothesis $d + r \leq g - 1$ which turns out to be *quite strong*. To understand how strong it is, consider the case $(d, s) = (g - 1, 0)$, which is indeed the crucial one (see [7, Proposition 3.3] and §3.2 below) and in which Debarre–Fahaloui’s theorem is void.

The aim of the present paper is to give the full classification of the cases in which (*) happens and $d = r + a + 2s$. What we prove (see Theorem 3.1 and Corollary 3.9) is that if (*) holds then, with no further assumption, either (a) or (b) or (c) occurs.

The idea of the proof is not so different, in principle, from the one proposed in [7] in the restricted situation considered there. Indeed, one uses the A -stability of Z and its maximal dimension to produce linear series on C which are not birational, in fact composed with a degree 2 irrational involution. The main tool in [7], inspired by [1], is a Castelnuovo’s type of analysis for the growth of the dimension of certain linear series.

Our approach also consists in producing a non birational linear series on C , but it is in a sense more direct. We consider (*) with $(d, s) = (g - 1, 0)$ and $a + r = g - 1$, i.e., the basic case (the others follow from this), in which Z is contained in $W_{g-1}(C)$, which is a translate of the theta divisor $\Theta \subset J(C)$. This immediately produces, restricting to Z the Gauss map of Θ , a base point free sublinear series L of dimension r of the canonical series of C . It turns out that Z is birational to an irreducible component of the variety $C(g - 1, L) \subset C(g - 1)$ consisting of all effective divisors of C of degree $g - 1$ contained in some divisor of L . The A -stability of Z implies that $C(g - 1, L)$ has some other component besides the one birational to Z , and this forces L to be non-birational. Once one knows this, a (rather subtle) analysis of the map determined by L and of its image leads to the conclusion.

As for the contents of the paper, §2 is devoted to a few general facts about the varieties $C(k, L) \subset C(k)$ of effective divisors of degree k contained in a divisor of a linear series L on a curve C . These varieties, as we said, play a crucial role in our analysis. Section 3 is devoted to the proof of our main result.

One final word about our own interest in this problem, which is quite different from the motivation of [1, 7]. It is in fact related to the study of irregular surfaces S of general type, where situation (*) presents itself in a rather natural way. For example, let $C \subset S$ be a smooth, irreducible curve, and assume that C corresponds to the general point of an irreducible component \mathcal{C} of the Hilbert scheme of curves on S which dominates $\text{Pic}^0(S)$. Recall that there is also only one irreducible component \mathcal{K} of the Hilbert scheme of curves algebraically equivalent to canonical curves on S which dominates $\text{Pic}^0(S)$ (this is called the *main paracanonical system*). The curves of \mathcal{C} cut out on C divisors which are residual, with respect to $|K_C|$, of divisors cut out by curves in \mathcal{K} . Consider now the one of the two systems \mathcal{C} and \mathcal{K} whose curves cut on C divisors of minimal

degree $d = \min\{C^2, K_S \cdot C\}$, and denote by s the dimension of the general fibre of this system over $\text{Pic}^0(S)$. Thus the image of the natural map $\text{Pic}^0(S) \rightarrow J^d(C)$ defined by $\eta \mapsto \mathcal{O}_C(C + \eta)$ (or by $\eta \mapsto \mathcal{O}_C(K_S + \eta)$) is a q -dimensional abelian variety contained in $W_d^s(C)$, which is what happens in (*). Thus understanding (*) would provide us with the understanding of (most) curves on irregular surfaces.

The results in this paper, even if restricted to the very special case of (*) in which Z has maximal dimension, turn out to be useful in surface theory. For example, if S is a minimal irregular surface of general type, then $K_S^2 \geq 2p_g$ (see [6]). Using the results in this paper we are able to classify irregular minimal surfaces for which $K_S^2 = 2p_g$. This classification is given in [5].

2. LINEAR SERIES ON CURVES AND RELATED SUBVARIETIES OF SYMMETRIC PRODUCTS

2.1. Generalities. Let C be a smooth, projective, irreducible curve of genus g . For an integer $k \geq 0$, we denote by $C(k)$ the k -th symmetric product of C (by convention, $C(0)$ is a point).

Let L be a base point free g_d^r on C . We denote the corresponding morphism by $\phi_L: C \rightarrow \bar{C} \subset \mathbb{P}^r$ and by C' the normalization of \bar{C} . We let $f: C \rightarrow C'$ be the induced morphism and L' the linear series on C' such that $L = f^*(L')$. We set $\deg(f) = \nu \geq 1$, so that $d = \delta\nu$, with $\delta = \deg(\bar{C})$. We say that L is *birational* if $\phi_L: C \rightarrow \bar{C}$ is birational.

Let $k \leq d$ be a positive integer. We consider the incidence correspondence

$$\mathcal{C}(k, L) = \{(D, H) \in C(k) \times L \mid D \leq H\}$$

with projections p_i (with $1 \leq i \leq 2$) to the first and second factor. Set $C(k, L) = p_1(\mathcal{C}(k, L))$, which has a natural scheme structure (see [4] or [2, p. 341]). Note the isomorphism

$$\mathfrak{s}: (D, H) \in \mathcal{C}(k, L) \rightarrow (H - D, H) \in \mathcal{C}(d - k, L).$$

Lemma 2.1. (i) *If $k \leq r$, then $C(k, L) = C(k)$.*

(ii) *If $k \geq r$, then $C(k, L)$ has pure dimension r . If V is an irreducible component of $C(k, L)$, there is a unique component \mathcal{V} of $\mathcal{C}(k, L)$ dominating V via p_1 and p_1 induces a birational morphism of \mathcal{V} onto V . Finally \mathcal{V} dominates L via p_2 .*

(iii) *If $\min\{k, d - k\} \geq r$ the isomorphism \mathfrak{s} induces a map $\mathfrak{r}: C(k, L) \dashrightarrow C(d - k, L)$, which is componentwise birational.*

(iv) *If L is birational, then $C(k, L)$ is irreducible.*

Proof. Part (i) is clear. The dimensionality assertion in (ii) follows from [4, § 1] or [2, Lemma (3.2), p. 342]. The rest of (ii) follows from these facts: the fibres of p_1 are isomorphic to linear subseries of L and p_2 is finite, so no component of $\mathcal{C}(k, L)$ has dimension larger than r . Part (iii) follows from (ii). Part (iv) follows from the Uniform Position Theorem (see [2, p. 112]). \square

Next we look at the case L non-birational and $k \geq r$. Consider the induced finite morphism $f_*: C(k, L) \rightarrow C'(k)$. For each partition $\underline{m} = (m_1, \dots, m_s)$ of k with $\delta \geq s \geq r$ and $1 \leq m_1 \leq m_2 \leq \dots \leq m_s \leq \nu$, we denote by $C(\underline{m}, L)$ the closure in $C(k, L)$ of the inverse image via f_* of the set of divisors of the form $m_1 y_1 + \dots + m_s y_s$, with $y_1 + \dots + y_s$ a reduced divisor in $C'(s, L')$. To denote a partition \underline{m} as above we may use the *exponential notation* $\underline{m} = (1^{\mu_1}, \dots, \nu^{\mu_\nu})$, meaning that i is repeated μ_i times, with $1 \leq i \leq \nu$, and we may omit i^{μ_i} if $\mu_i = 0$. Note that $\sum_{i=1}^{\nu} \mu_i = s \leq \delta$. Set $\mu_0 := \delta - s$. Then $\underline{m}^c = (1^{\mu_{\nu-1}}, \dots, (\nu-1)^{\mu_1}, \nu^{\mu_0})$ is a partition of $d - k$ which we call the *complementary partition* of \underline{m} .

Lemma 2.2. *In the above set up each irreducible component of $C(\underline{m}, L)$ has dimension r , hence it is an irreducible component of $C(k, L)$ and all irreducible components of $C(k, L)$ are of this type.*

Proof. Let V be an irreducible component of $C(k, L)$: by Lemma 2.1 there is a unique irreducible component \mathcal{V} of $\mathcal{C}(k, L)$ dominating it and $p_1|_{\mathcal{V}}: \mathcal{V} \rightarrow V$ is birational. Let $D \in V$ be a general point and let (D, H) be the unique point of \mathcal{V} mapping to D via p_1 , so that H is a general divisor in L (cf. Lemma 2.1, (ii)). Hence H consists of δ distinct fibres F_1, \dots, F_δ of f , each being a reduced divisor of degree ν on C . Then D consists of m_i points in F_i , for $1 \leq i \leq s \leq \delta$, where we may assume $1 \leq m_1 \leq m_2 \leq \dots \leq m_s \leq \nu$. Moreover, since $\dim(V) = \dim(\mathcal{V}) = r$ and p_2 is finite, one has $s \geq r$. Hence $D \in C(\underline{m}, L)$, with $\underline{m} = (m_1, \dots, m_s)$, i.e. $V \subseteq C(\underline{m}, L) \subseteq C(k, L)$ hence V is a component of $C(\underline{m}, L)$.

The above considerations and Lemma 2.1, (ii), applied to $C'(s, L')$, imply that the image of $C(\underline{m}, L)$ in $C'(k)$ has dimension r , so each component of $C(\underline{m}, L)$ has dimension r . \square

2.2. Abel–Jacobi images. We assume from now on that C has genus $g > 0$. For an integer k , we denote by $J^k(C) \subset \text{Pic}(C)$ the set of linear equivalence classes of divisors of degree k on C . So $J(C) := J^0(C)$ is the Jacobian of C , which is a principally polarised abelian variety whose theta divisor class we denote by Θ_C , or simply by Θ .

The abelian variety $J(C)$ acts via translation on $J^k(C)$ for all k . If $X \subseteq J^k(C)$ and $Y \subseteq J(C)$, we say that X is Y -stable, if for all $x \in X$ and for all $y \in Y$, one has $x + y \in X$.

For all integers k , fixing the class of a divisor of degree k determines an isomorphism $J^k(C) \cong J(C)$. Given a subvariety V of $J^k(C)$, one says that it *generates* $J^k(C)$ if the image of V via one of the above isomorphisms generates $J(C)$ as an abelian variety. This definition does not depend on the choice of the isomorphism $J^k(C) \cong J(C)$.

For every $k \geq 1$, we denote by $j_k: C(k) \rightarrow J^k(C) \cong J(C)$ (or simply by j) the Abel–Jacobi map. We denote by $W_k^s(C)$ the subscheme of $J^k(C)$ corresponding to classes of divisors D such that $h^0(\mathcal{O}_C(D)) \geq s + 1$ (these are the so-called *Brill–Noether loci*). One sets $W_k(C) := W_k^0(C) = \text{Im}(j_k)$ and $W_{g-1}(C)$ maps to a theta divisor of $J(C)$, so we may abuse notation and write $W_{g-1}(C) = \Theta_C$.

We denote by $\Gamma_C(k, L)$ [resp. $\Gamma_C(\underline{m}, L)$] the image in $W_k(C)$ of $C(k, L)$ [resp. $C(\underline{m}, L)$] (we may drop the subscript C if there is no danger of confusion). The expected dimension of $\Gamma(k, L)$ is $\min\{r, g\}$ (by dimension of a scheme we mean the maximum of the dimensions of its components).

We set $\rho_C(k, L) := \dim(\Gamma_C(k, L))$ (simply denoted by $\rho(k, L)$ or by ρ if no confusion arises). By Lemma 2.1, (i), one has:

$$(2.1) \quad \text{if } k \leq r \text{ then } \Gamma(k, L) = W_k(C), \text{ hence } \rho = \min\{k, g\}$$

So we will consider next the case $k > r$, in which $\rho \leq \dim(C(k, L)) = r$, by Lemma 2.1, (ii). Then the class $c(k, L)$ of $C(k, L)$ in the Chow ring of $C(k)$ is computed in [2, Lemma VIII.3.2]. If x is the class of $C(k-1) \subset C(k)$ and $\theta := j^*(\Theta)$, one has

$$(2.2) \quad c(k, L) = \sum_{s=0}^{k-r} \binom{d-g-r}{s} \frac{x^s \theta^{k-r-s}}{(k-r-s)!}.$$

Lemma 2.3. *Assume $k > r$ and $d - g - r \geq 0$. Then:*

- (i) *if $k - g \leq \min\{k - r, d - g - r\}$ one has $\rho = r \leq g$;*
- (ii) *if $k - g \geq \min\{k - r, d - g - r\} = k - r$, one has $\rho = g \leq r$;*
- (iii) *if $k - g \geq \min\{k - r, d - g - r\} = d - g - r$, one has $\rho = d - k \leq \min\{r, g\}$.*

Proof. Note that x^s is the class of $C(k-s) \subset C(k)$, for $1 \leq s \leq k$. Applying the projection formula (cf. [10], Example 8.1.7) to (2.2), we find the class $\gamma(k, L)$ of $\Gamma(k, L)$

$$(2.3) \quad \gamma(k, L) = \sum_{s=0}^{k-r} \binom{d-g-r}{s} \frac{w_{k-s} \Theta^{k-r-s}}{(k-r-s)!}$$

where w_i is the class of $W_i(C)$ for any $i \geq 0$. By Poincaré’s formula (cf. [3, §11.2]) one has

$$w_{k-s}\Theta^{k-r-s} = \begin{cases} \Theta^{k-r-s}, & \text{if } k-g \geq s \geq 0, \\ \frac{\Theta^{g-r}}{(g-k+s)!}, & \text{if } \max\{0, k-g\} \leq s \leq \min\{k-r, d-g-r\}, \end{cases}$$

whence the assertion follows. \square

Lemma 2.4. (i) If $\rho = g$ then $g \leq \min\{k, r\}$;
(ii) if $r \geq k \geq g$, then $\rho = g$;
(iii) if $k > r \geq g$ and $d \geq k + g$ then $\rho = g$;
(iv) $\rho = 0$ if and only if $k = d$.

Proof. Parts (i) and (ii) follow from Lemma 2.1.

(iii) In (2.3) one has the summand corresponding to the index $s = k - r > 0$, which is Θ^0 with the positive coefficient $\binom{d-g-r}{k-r}$, and no other summand in (2.3) cancels it.

(iv) If $k = d$ then $C(k, L) = L$ and clearly $\rho = 0$. Conversely, if $\rho = 0$ then in (2.3) the term Θ^g has to appear with non-zero coefficient and no other term Θ^i with $0 \leq i < g$ appears with non-zero coefficient. By looking at the proof of Lemma 2.3, we see that the summand Θ^g appears in (2.3) only if $0 \leq s = k - r - g$. Then $d \geq k \geq r + g$. So we may apply Lemma 2.3, and conclude that $\rho = 0$ occurs only in case (iii), if $k = d$. \square

Lemma 2.5. Let $A \subseteq J(C)$ be an abelian subvariety of dimension a and let $p: J^k(C) \rightarrow J' := J(C)/A$ be the map obtained by composing an isomorphism $J^k(C) \cong J(C)$ with the quotient map $J(C) \rightarrow J'$. Then

$$\dim(p(\Gamma(k, L))) = \min\{g - a, \rho\}.$$

Proof. If $\rho = g$ the statement is obvious, hence we assume $\rho < g$.

Consider first the case $k > r$. Assume by contradiction that $\dim(p(\Gamma(k, L))) < \min\{\rho, g - a\}$. Let ξ be the class of the pull back to $J(C)$ of an ample line bundle of J' . We have $\bar{\gamma}(k, L)\xi^\rho = 0$, where $\bar{\gamma}(k, L)$ is the ρ -dimensional part of $\gamma(k, L)$. By (2.3) one has $\bar{\gamma}(k, L) = \alpha\Theta^{g-\rho}$, where $\alpha \in \mathbb{Q}$ is positive because $\Gamma(k, L)$ is an effective non-zero cycle of dimension ρ . Hence $\bar{\gamma}(k, L)\xi^\rho = \alpha\Theta^{g-\rho}\xi^\rho > 0$, because Θ is ample. Thus we have a contradiction.

If $k \leq r$, then $\Gamma(k, L) = W_k(C)$, $\rho = k$ and $\gamma(k, L)$ is again a rational multiple of Θ^{g-k} (by Poincaré’s formula), so the proof proceeds as above. \square

Corollary 2.6. If $A \subseteq J(C)$ is an abelian subvariety of dimension $a > 0$ and $\Gamma(k, L)$ is A -stable, then the restriction of p to $\Gamma(k, L)$ is surjective onto $J' = J(C)/A$, hence $\Gamma(k, L) = J^k(C) = W_k(C)$, i.e., $\rho = g$.

Corollary 2.7. If $d - 1 \geq k \geq 1$, then $\Gamma(k, L)$ generates $J^k(C)$.

Proof. If $k \leq r$ then $\Gamma(k, L) = W_k(C)$ and the assertion is clear. Assume $k > r$. By Lemma 2.5, $\Gamma(k, L)$ generates $J^k(C)$ as soon as $\rho > 0$, which is the case by Lemma 2.4, (iv). \square

2.3. A useful lemma. Let L be a base point free g_d^1 , let $\phi_L: C \rightarrow \mathbb{P}^1$ be the corresponding map and denote by G_L the Galois group of ϕ_L .

Lemma 2.8. If L is a base point free g_4^1 , then one of the following occurs:

- (a) $C(2, L)$ is irreducible.
- (b) $C(2, L)$ has two components. This occurs if and only if $G_L \cong \mathbb{Z}_2, \mathbb{Z}_4$.
- (c) $C(2, L)$ has 3 components. This occurs if and only if $G_L \cong \mathbb{Z}_2^2$.

Proof. We argue as in the proof of [3, Lemma 12.7.1]. Let $\Delta \subset \mathbb{P}^1$ be the set of critical values of $\phi_L: C \rightarrow \mathbb{P}^1$, let $\rho: \pi_1(\mathbb{P}^1 \setminus \Delta) \rightarrow \mathfrak{S}_4$ be the monodromy representation and let $\Sigma := \text{Im}(\rho)$. The group Σ acts transitively on $\mathbb{I}_4 := \{1, 2, 3, 4\}$ (identified with the general divisor $x_1 + \dots +$

$x_4 \in L$) and its order s is divisible by 4. The irreducible components of $C(2, L)$ are in 1-to-1 correspondence with the Σ -orbits of the order two subsets of \mathbb{I}_4 . If Σ contains an element of order 3, then s is divisible by 12. Hence $\mathfrak{A}_4 \subseteq \Sigma$, thus the action is transitive, $C(2, L)$ is irreducible and (a) holds.

Assume Σ is a 2-group. If $s = 8$ then Σ is a 2-Sylow subgroup of \mathfrak{S}_4 , hence Σ is the dihedral group D_4 . Then C is obtained from a Σ -cover $C_0 \rightarrow \mathbb{P}^1$ by moding out by a reflection $\sigma \in \Sigma$. In the Σ -action on \mathbb{I}_4 we may assume that an element of order 4 acts by sending i to $i + 1$ modulo 4, for $1 \leq i \leq 4$. So the order 2 element in the center of Σ induces the involution ι of C that maps i to $i + 2$ modulo 4, for $1 \leq i \leq 4$, and ι generates $G_L \cong \mathbb{Z}_2$. There are two orbits for the Σ -action on the set of order two subsets of \mathbb{I}_4 : one of order 2 given by $\{\{1, 3\}, \{2, 4\}\}$, the other of order 4 given by $\{\{i, i + 1\}, \text{ for } 1 \leq i \leq 4\}$ (here i is taken modulo 4). These orbits respectively correspond to two components E_1, E_2 of $C(2, L)$ and we are in case (b).

Assume $s = 4$. Then ϕ_L is Galois with $G_L = \Sigma$. If $G_L \cong \mathbb{Z}_4$, then the Σ -orbits on the set of order two subsets of \mathbb{I}_4 are as in the previous case. If $G_L \cong \mathbb{Z}_2^2$, then one has $G_L = \{\text{Id}, (12)(34), (13)(24), (14)(32)\}$. There are then three orbits, corresponding to three components E_1, E_2, E_3 . \square

One can be more precise about the components E_i of $C(2, L)$ in Lemma 2.8, whose geometric genera we denote by g_i (with $1 \leq i \leq 2 + \epsilon$, and $\epsilon = 0$ in case (b), $\epsilon = 1$ in case (c)).

Lemma 2.9. *Same setting and notation as in Lemma 2.8 and its proof. Then:*

(i) *each component of $C(2, L)$ maps birationally to its image in $J^2(C)$ unless C is hyperelliptic and L is composed with the hyperelliptic involution \mathcal{L} : in this case one of the components of $\Gamma(2, L)$ is $\mathcal{L} = C(2, \mathcal{L})$, which is contracted to a point in $J^2(C)$;*

(ii) *in case (b) one has $E_1 \cong C/\iota$ (where ι is the non-trivial involution in G_L), which is hyperelliptic, and the abelian subvariety of $J^2(C) \cong J(C)$ generated by $j(E_1)$ is the ι -invariant part of $J(C)$. Moreover $2g_2 \geq g$;*

(iii) *In case (c) one has $E_i := C/\iota_i$ where ι_i are the three nonzero elements of G_L , for $1 \leq i \leq 3$.*

Proof. We prove the only non-trivial assertion, i.e., $2g_2 \geq g$ in part (ii).

First assume $G_L = \mathbb{Z}_4 = \langle \rho \rangle$. Consider in $C(2)$ the curves $E_1 = \{P + \rho^2(P) \mid P \in C\}$ and $E_2 = \{P + \rho(P) \mid P \in C\}$. One has $C(2, L) = E_1 \cup E_2$ and $E_1 \cong C/\rho^2$. The curve E_2 is the image in $C(2)$ of the graph of ρ , so $g_2 = g$.

Suppose now $G_L = \mathbb{Z}_2$ and $\Sigma = D_4$. Recall that C is obtained from a D_4 -Galois cover $f: C_0 \rightarrow \mathbb{P}^1$ by moding out by a reflection $\sigma \in D_4$ (see the proof of Lemma 2.8). Denote by g_0 the genus of C_0 . Let $\rho \in D_4$ be an element of order 4, so that $D_4 = \langle \sigma, \rho \rangle$. Let n be the number of points of C_0 fixed by σ , n' the number of points fixed by $\sigma\rho$, m the number of points fixed by ρ and $m + \varepsilon$ the number of points fixed by ρ^2 . The Hurwitz formula, applied to $C_0 \rightarrow \mathbb{P}^1$ and to $C_0 \rightarrow C = C_0/\sigma$, gives

$$(2.4) \quad g_0 = \frac{3}{2}m + n + n' + \frac{\varepsilon}{2} - 7, \quad \text{and} \quad g = \frac{n'}{2} + \frac{n}{4} + \frac{\varepsilon}{4} - \frac{3}{4}m - 3.$$

Claim 2.10. (i) m, n, n' and ε are even.

(ii) $n + m \equiv n' + m \equiv n + n' \equiv 0 \pmod{4}$, and at most one among m, n, n' can be 0.

Proof of the Claim. (i) The numbers n, n' and $m + \varepsilon$ are even because $\sigma, \sigma\rho$ and $\sigma\rho^2$ are involutions. If $P \in C_0$ is fixed by ρ , then $\sigma(P)$ is also fixed by ρ . Since the stabilizer of any point is cyclic, then $\sigma(P) \neq P$. This implies that m is even.

(ii) Consider the \mathbb{Z}_2^2 -cover $D := C_0/\rho^2 \rightarrow \mathbb{P}^1$. The cardinalities of the images in \mathbb{P}^1 of the fixed loci of the three involutions are $n/2, n'/2$ and $m/2$. Indeed, denote by γ_1 [resp. by γ_2] the image of σ [resp. of ρ] in $D_4/\rho^2 \cong \mathbb{Z}_2^2$. Let $Q \in \mathbb{P}^1$ be a branch point whose preimage in D is fixed by γ_1 . Then the preimage of Q in C_0 consists of 4 points, two of which fixed by σ and two by

$\sigma\rho^2$, so the number of such points Q is $(2n)/4 = n/2$. Similarly, the image in \mathbb{P}^1 of the set of points of D fixed by γ_2 has cardinality $m/2$ and the image of the set of points fixed by $\gamma_1\gamma_2$ has cardinality $n'/2$. Hurwitz formula for $D_1 := D/\gamma_1 \rightarrow \mathbb{P}^1$ gives

$$2g(D_1) - 2 = \frac{m}{2} + \frac{n'}{2} - 4,$$

hence $m + n' > 0$ is divisible by 4. Similarly $n + n'$ and $m + n$ are positive and divisible by 4. \square

We compute now the ramification of $f: C_0 \rightarrow \mathbb{P}^1$ and of L . Write the D_4 -orbit of $P \in C_0$ general as

$$(2.5) \quad \begin{array}{cccc} P & \rho(P) & \rho^2(P) & \rho^3(P) \\ \sigma(P) & \sigma\rho(P) & \sigma\rho^2(P) & \sigma\rho^3(P). \end{array}$$

Denote by Q_1, \dots, Q_4 the images in C of the points in the first row (or, what is the same, in the second row) of (2.5). The singular fibers of f occur when P has non trivial stabilizer, i.e., when:

- $P \in C_0$ is fixed by ρ . The fiber of f is $4(P + \sigma(P))$ and the corresponding divisor of L is $4Q_1$. There are $m/2$ divisors of L of this type;
- P is fixed by ρ^2 but not by ρ . The fiber of f is $2(P + \sigma(P) + \rho(P) + \sigma\rho(P))$. Then $Q_1 = Q_3$, $Q_2 = Q_4$, and the corresponding divisor of L is $2(Q_1 + Q_2)$. There are $\varepsilon/4$ such divisors;
- P is fixed by σ . The fibre of f is $2(P + \rho(P) + \rho^2(P) + \rho^3(P))$, and $Q_2 = Q_4$, while Q_1, Q_2 and Q_3 are distinct, so the corresponding divisor of L is $Q_1 + 2Q_2 + Q_3$ and there are $n/2$ such divisors;
- P is fixed by $\sigma\rho^2$. This is the same as the previous case;
- P is fixed by $\sigma\rho$. The fibre of f is again $2(P + \rho(P) + \rho^2(P) + \rho^3(P))$, and $Q_1 = Q_2$, $Q_3 = Q_4$, so the corresponding divisor of L is $2Q_1 + 2Q_3$ and there are $n'/2$ such fibers;
- P is fixed by $\sigma\rho^3$. This is the same as the previous case.

Denote by ι the involution of C induced by ρ^2 . Then $C(2, L)$ is the union of $E_1 = \{P + \iota(P) \mid P \in C\} = C/\iota$ and of the irreducible curve E_2 . Keeping the above notation, if $Q_1 + \dots + Q_4$ is the general divisor of L , then E_1 is described by the divisors $Q_1 + Q_3, Q_2 + Q_4$ and E_2 by $Q_1 + Q_2, Q_1 + Q_4, Q_2 + Q_3, Q_3 + Q_4$.

To compute g_2 , define a map $\phi: C_0 \rightarrow E_2$ by sending $P \in C_0$ to the image via $C_0 \rightarrow C$ of the divisor $P + \rho(P)$, i.e., $Q_1 + Q_2 \in E_2$.

Claim 2.11. (i) $\deg(\phi) = 2$ and E_2 is birational to $C_0/\sigma\rho$;

(ii) $g_2 = \frac{3}{4}m + \frac{n}{2} + \frac{n'}{4} + \frac{\varepsilon}{4} - 3$.

Proof of the Claim. (i) Let $Q + Q' \in E_2$ be a general point and let $P, \sigma(P)$ [resp. $P', \sigma(P')$] be the preimages of Q [resp. of Q'] on C_0 . Since $Q + Q'$ is of the form $Q_1 + Q_2, Q_1 + Q_4, Q_2 + Q_3$, or $Q_3 + Q_4$, we may assume that $P' = \rho(P)$, so that $\psi^{-1}(Q + Q') = \{P, \sigma\rho(P)\}$.

Part (ii) follows by applying Hurwitz formula. \square

Finally, suppose by contradiction that $2g_2 \leq g - 1$. Then, by (ii) of Claim 2.11 and by (2.4), we would have $3(m + n) + \varepsilon \leq 8$, hence $m + n \leq 2$, contradicting (ii) of Claim 2.10. \square

3. ABELIAN SUBVARIETIES OF BRILL–NOETHER LOCI

As in [1, 7], we consider $Z \subseteq W_d^s(C) \subsetneq J^d(C)$ an irreducible A -stable variety of dimension r , with $A \subsetneq J(C)$ an abelian subvariety of dimension $a > 0$. Note that $r \geq a$, with equality if and only if $Z \cong A$. Moreover, since $W_d^s(C) \subsetneq J^d(C)$, the general linear series $L \in Z$ is special, thus $s > d - g$ and $d \geq 2s$ by Clifford's theorem. From [7, Proposition 3.3], we have

$$(3.1) \quad r + a + 2s \leq d.$$

In this section, we classify the cases in which equality holds in (3.1), thus improving the partial results in [7] on this subject. Note that if equality holds in (3.1), then $Z \not\subseteq W_d^{s+1}(C)$ and A is a maximal abelian subvariety of $J(C)$ such that Z is A -stable.

3.1. The Theta divisor case. As in [7], we first consider the case $(d, s) = (g - 1, 0)$, i.e., $Z \subset W_{g-1}(C) = \Theta$ is an irreducible A -stable variety of dimension $r = g - 1 - a$. Then $Z \not\subseteq W_{g-1}^1(C) = \text{Sing}(\Theta)$, i.e., Z has a non-empty intersection with $\Theta_{\text{sm}} := \Theta - \text{Sing}(\Theta)$.

Theorem 3.1. *Let C be a curve of genus g . Let $A \subsetneq J(C)$ be an abelian variety of dimension $a > 0$ and $Z \subset \Theta$ an irreducible, A -stable variety of dimension $r = g - 1 - a$. Then there is a degree 2 morphism $\varphi: C \rightarrow C'$, with C' smooth of genus g' , such that one of the following occurs:*

- (a) $g' = a$, $A = \varphi^*(J(C'))$ and $Z = W_{g-1-2a}(C) + \varphi^*(J^a(C'))$;
- (b) $g' = r + 1$, φ is étale, A is the Prym variety of φ and $Z \subset W_{g-1}(C)$ is the connected component of $\varphi_*^{-1}(K_{C'})$ consisting of divisor classes D with $h^0(\mathcal{O}_C(D))$ odd, where $\varphi_*: J^{g-1}(C) \rightarrow J^{g-1}(C')$ is the norm map. In particular, $Z \cong A$ is an abelian variety if and only if either we are in case (a) and $g = 2a + 1$, or in case (b).

Remark 3.2. Cases (a) and (b) of Theorem 3.1 are not mutually exclusive. Indeed, if the curve C in case (b) is hyperelliptic, then the Abel-Prym map $C \rightarrow A$ induces a 2-to-1 map $\psi: C \rightarrow D$, where D is a smooth curve embedded into $A \cong J(D)$ by the Abel-Jacobi map. One can check that $A = \psi^*(J^r(D))$, namely this is also an instance of case (a) of Theorem 3.1.

The proof of Theorem 3.1 requires various preliminary lemmas. First, recall that the tangent space to $J(C)$ at 0 can be identified with $H^1(\mathcal{O}_C) \cong H^0(K_C)^*$. Denote by $T \subseteq H^0(K_C)^*$ the tangent space to A at 0 and by L the linear series $\mathbb{P}(T^\perp) \subseteq |K_C|$. One has $\dim(L) = g - 1 - a = r$.

Since $Z \cap \Theta_{\text{sm}} \neq \emptyset$, the Gauss map of Θ restricts to a rational map $\gamma: Z \dashrightarrow \mathbb{P} := |K_C|$.

Lemma 3.3. *One has $\gamma(Z) = L$.*

Proof. Since Z is A -stable, one has $\gamma(Z) \subseteq L$. A point of Θ_{sm} can be identified with a divisor D of degree $g - 1$ such that $h^0(\mathcal{O}_C(D)) = h^0(\mathcal{O}_C(K_C - D)) = 1$. The Gauss map sends $D \in \Theta_{\text{sm}}$ to the unique divisor of $|K_C|$ containing D . Then $\gamma^{-1}(\gamma(D))$ is finite. Hence $\dim(\gamma(Z)) = \dim(Z) = r = \dim(L)$. The assertion follows. \square

As in [7], formula (3.1) for $(d, s) = (g - 1, 0)$ follows from the argument in the proof of Lemma 3.3, and the general case follows from this (see §3.2 below).

Note the birational involution $\sigma: \Theta \dashrightarrow \Theta$, defined on Θ_{sm} , sending a divisor $D \in \Theta_{\text{sm}}$ to the unique effective divisor $D' \in |K_C - D|$. Then σ restricts on Z to a birational map (still denoted by σ) onto its image Z' , which is also A -stable.

Lemma 3.4. *The linear series L is base point free.*

Proof. Suppose $P \in C$ is a base point of L . If every $D \in Z$ contains P , then the map $D \rightarrow D - P$ defines an injection $Z \hookrightarrow W_{g-2}(C)$, contradicting (3.1). If the general $D \in Z$ does not contain P , then $\sigma(D)$ contains P for $D \in Z$ general, and we can apply the previous argument to Z' . \square

Lemma 3.5. *One has:*

- (i) Z is a component of $\Gamma(g - 1, L)$.
- (ii) $Z \subsetneq \Gamma(g - 1, L)$.
- (iii) *The linear series L is not birational.*

Proof. By Lemma 3.3, we have $Z \subseteq \Gamma(g - 1, L)$ and the dimension of $\Gamma(g - 1, L)$ is equal to r (by Lemma 2.3). So (i) holds. If $Z = \Gamma(g - 1, L)$, then $\Gamma(g - 1, L)$ is A -stable. By Corollary 2.6, we have $r = g$, a contradiction. This proves (ii). Then (iii) holds by Lemma 2.1, (iv). \square

Recall the notation $\phi_L: C \rightarrow \bar{C} \subset \mathbb{P}^r$, C' for the normalization of \bar{C} , $f: C \rightarrow C'$ the induced morphism and $\nu = \deg(f) > 1$. For any integer h one has the morphism $f^*: J^h(C') \rightarrow J^{h\nu}(C)$. Let g' be the genus of C' and L' the (birational) linear series on C' of dimension r such that $L = f^*(L')$, whose degree is $\delta = \frac{2g-2}{\nu}$.

Lemma 3.6. *Assume $\nu \leq 3$ and let $\underline{m} = (1^{\mu_1}, \dots, \nu^{\mu_\nu})$ be the partition of $g-1$ such that $Z \subseteq \Gamma(\underline{m}, L)$. Set $\mu_1 = \mu$, $\mu_2 = \mu'$. Then $\nu = 2$ and there are the following possibilities:*

- (i) $\mu = g-1$ (hence $\mu' = 0$);
- (ii) $\mu = g-1-2a$, $\mu' = a$, $A = f^*(J(C'))$ (hence $g' = a$) and $Z = W_\mu(C) + f^*(J^a(C'))$.

Proof. We first consider the case $\nu = 2$.

If $\mu = 0$, then $g-1 = 2\mu'$ and $Z = f^*(\Gamma_{C'}(\mu', L'))$, since $\Gamma_{C'}(\mu', L')$ is irreducible by Lemma 2.1. The general point of Z corresponds to a linearly isolated divisor, hence $\mu' \leq g'$. Since $f^*: J^{\mu'}(C') \rightarrow J^{g-1}(C)$ is finite, one has $r = \rho_{C'}(\mu', L') \leq \mu' \leq g'$. Since Z is A -stable, we have an isogeny $A \rightarrow \bar{A} \subseteq J(C')$ (so that $a \leq g'$) and $\Gamma_{C'}(\mu', L')$ is \bar{A} -stable. Hence by Corollary 2.6 one has $\Gamma_{C'}(\mu', L') = J^{\mu'}(C') = W_{\mu'}(C')$ and $g' \leq \min\{\mu', r\}$ (see Lemma 2.4, (i)). Then Z is $f^*(J(C'))$ -stable, and, since a is the maximal dimension of an abelian subvariety of $J(C)$ for which Z is stable and $a = \dim(\bar{A}) \leq g'$, it follows $a = g'$. In conclusion $\mu' = a = r = g'$, and we are in case (ii).

Assume now $\mu > 0$. The general point of Z (which is a component $\Gamma(\underline{m}, L)$) is smooth for Θ , hence it corresponds to a linearly isolated, effective divisor D of degree $g-1$, which is reduced (see Lemma 2.1, (ii)) and can be written in a unique way as $D = M + f^*(N)$, where M and N are effective divisors, with $\deg(M) = \mu$, $\deg(N) = \mu'$ and $M' := f_*(M)$ reduced. So there is a rational map $h: Z \dashrightarrow J^\mu(C)$ defined by $D = M + f^*(N) \mapsto M$.

Assume $\mu \leq r$. By Lemma 2.1 the image of h is $W_\mu(C)$. Identify A with its general translate inside Z . Then we have a morphism $h|_A: A \rightarrow J(C)$ whose image we denote by \bar{A} . Then $W_\mu(C)$ is \bar{A} -stable. Since $W_\mu(C)$ is birational to $C(\mu)$, which is of general type because $\mu < g$, then $\bar{A} = \{0\}$. It follows that each component of the general fibre of h is A -stable, in particular $r - \mu \geq a \geq 1$.

Take $M \in W_\mu(C)$ general and set $L'' := L'(-M')$. Since L' is birational and $M' \in W_\mu(C')$ is general, with $\mu < r = \dim(L')$, then L'' has dimension $r - \mu \geq 1$ and it is base point free. It is also birational as soon as $r - \mu \geq 2$.

Assume first $r - \mu \geq 2$. Then $C'(\mu', L'')$ is irreducible by Lemma 2.1, (iv), and there is a birational morphism $C'(\mu', L'') \rightarrow h^{-1}(M) \subset J^{g-1}(C)$ factoring through the map $C'(\mu', L'') \rightarrow J^{\mu'}(C') \hookrightarrow J^{g-1}(C)$, where the last inclusion is translation by M . Namely, up to a translation, $h^{-1}(M) = \Gamma(\mu', L'')$. In particular, $\dim(\Gamma(\mu', L'')) = \dim(C'(\mu', L'')) = r - \mu$, hence $\mu' \geq r - \mu$.

Remember that $h^{-1}(M) = \Gamma(\mu', L'')$ is A -stable for $M \in W_\mu(C)$ general. By Corollary 2.6 one has $\Gamma_{C'}(\mu', L'') = W_{\mu'}(C') = J^{\mu'}(C')$, so $g' \leq \min\{\mu', r - \mu\} = r - \mu$. On the other hand, since $h^0(\mathcal{O}_C(D)) = 1$, one has also $h^0(\mathcal{O}_C(f^*(N))) = 1$, hence $\mu' \leq g'$ and we conclude that $\mu' = g' = r - \mu$. The same argument as above yields $a = g'$ and we are again in case (ii).

If $r - \mu = 1$, then $a = 1$, $r = g - 2$, $\mu = g - 3$ and $\mu' = 1$. On the other hand L is cut out on the canonical image of C by the hyperplanes through the point p_A which is the projectivized tangent space to A at the origin. Then $\phi_L: C \rightarrow \bar{C}$ is the projection from p_A , \bar{C} it is a normal elliptic curve, and we are again in case (ii).

Assume now $\mu > r$ and keep the above notation. In this case the map $h: Z \dashrightarrow \bar{Z} := h(Z)$ is generically finite, \bar{A} is isogenous to A and \bar{Z} is \bar{A} -stable. By [7, Lemma 3.1], we have $g-1 \geq \mu \geq \dim(\bar{Z}) + \dim(\bar{A}) = r + a = g-1$, thus $\mu = g-1$, so we are in case (i).

Finally consider the case $\nu = 3$. Write the general $D \in \Gamma(\underline{m}, L)$ as $D = M_1 + M_2 + f^*(N)$, where M_1 is reduced of degree μ , $f_*(M_2) = 2M'_2$ with M'_2 reduced of degree μ' and, as above,

$\mu_3 = \deg(N) \leq g'$. Set $\tau = \mu + \mu'$ and consider the rational map $h: Z \dashrightarrow W_\tau(C) \subseteq J^\tau(C)$ defined by $D \mapsto M_1 + f^*(M'_2) - M_2$.

If $\tau \leq r$, arguing as above (and keeping a similar notation) one sees that $h(Z) = W_\tau(C)$, the general fibre of h is A -stable, hence $r - \tau \geq a$. However $a = 1$ and ϕ_L non-birational, forces, as we have seen, $\nu = 2$, which is not the case here. Hence we have $a \geq 2$. We consider now $L'' = L'(-f_*(M_1) - M'_2)$, which has dimension $r - \tau \geq 2$ and is base point free and birational, so $C'(\mu_3, L'')$ is irreducible. The general fiber $h^{-1}(h(D))$ of h is isomorphic to $\Gamma_{C'}(\mu_3, L'')$ and is A -stable. In particular $\dim(\Gamma_{C'}(\mu_3, L'')) = r - \tau$, hence $\mu_3 \geq r - \tau$. By Corollary 2.6, we have $\Gamma_{C'}(\mu_3, L'') \cong W_{\mu_3}(C') \cong J^{\mu_3}(C')$, so $g' \leq \min\{\mu_3, r - \tau\} = r - \tau$, hence $g' \leq r - \tau \leq \mu_3 \leq g'$, thus $\mu_3 = r - \tau = g'$. In addition, as above, we have $a = g'$, so that $A = f^*(J(C'))$. Then $g - 1 - a = r = \tau + \mu_3 = \mu + \mu' + \mu_3$. On the other hand $g - 1 = \mu + 2\mu' + 3\mu_3$. This yields $\mu' + 2\mu_3 = a = \mu_3$, hence $\mu' = \mu_3 = 0$, which is not possible.

If $\tau > r$ then $h: Z \dashrightarrow \bar{Z} := h(Z)$ is generically finite, \bar{A} is isogenous to A and \bar{Z} is \bar{A} -stable. By [7, Lemma 3.1], we have $g - 1 \geq \tau \geq \dim(\bar{Z}) + \dim(\bar{A}) = r + a = g - 1$, thus $\tau = g - 1$, contradicting $\tau \leq \deg(L') = \frac{2}{3}(g - 1)$. \square

Lemma 3.7. *If $\nu \geq 4$ then $\nu = 4$ and either*

- (i) *there is a degree 2 map $\psi: C \rightarrow E_1$ with E_1 a genus r hyperelliptic curve such that $Z = A = \psi^*(J^r(E_1))$; or*
- (ii) *there is a faithful \mathbb{Z}_2^2 -action on C with rational quotient; denoting by $f_i: C \rightarrow E_i$ (for $1 \leq i \leq 3$) the quotient map for the three non-trivial involutions of \mathbb{Z}_2^2 , with E_i of genus g_i , and $g_1 \geq g_2 \geq g_3$, then $g_1 = r + 1$, $g_2 + g_3 = r$ and $Z = A = f_2^*(J^{g_2}(E_2)) \times f_3^*(J^{g_3}(E_3))$ is the Prym variety associated to f_1 .*

Proof. Since L is base point free by Lemma 3.4 and it is not birational by Lemma 3.5, we have $\delta = \frac{2g-2}{\nu} \geq r \geq \frac{g-1}{2}$, hence $\nu \leq 4$.

Assume $\nu = 4$. Then \bar{C} is a curve of degree $\frac{g-1}{2}$ spanning a projective space of dimension $r \geq \frac{g-1}{2}$. Hence $r = a = \frac{g-1}{2}$, $Z = A$, and $\phi_L = f$ is the composition of a g_1^4 (that we denote by \mathcal{L}) with the degree r Veronese embedding $\mathbb{P}^1 \rightarrow \mathbb{P}^r$.

Let $\underline{m} = (1^{\mu_1}, \dots, 4^{\mu_4})$ be the partition of $2r$ such that $Z \subseteq \Gamma(\underline{m}, L)$.

Claim 3.8. *One has $\underline{m} = (2^r)$.*

Proof of the Claim. Assume by contradiction this is not the case, so that one among μ_1, μ_3, μ_4 is non-zero. We have $r = \dim(\Gamma(\underline{m}, L)) = \mu_1 + \dots + \mu_4$, because \bar{C} is a rational normal curve of degree r in \mathbb{P}^r , and $2r = g - 1 = \mu_1 + 2\mu_2 + 3\mu_3 + 4\mu_4$, hence $\mu_1 = \mu_3 + 2\mu_4$ and $\mu_1 > 0$. So we may write the general divisor $D \in Z$ as $D = M + N$, where $M' := f_*(M)$ is reduced of degree $\mu := \mu_1$. Then we proceed as in the proof of Lemma 3.6.

Consider the rational map $h: Z = A \dashrightarrow J^\mu(C)$ defined by $D = M + N \mapsto M$. It extends to a morphism and $\bar{A} := h(A)$ is an abelian variety contained in $W_\mu(C)$. Since $\mu \leq r = \frac{g-1}{2}$, one has $\bar{A} = W_\mu(C)$, which is impossible. \square

Let $E := C(2, L)$. Assume first C is not hyperelliptic. Then there is a birational (dominant) morphism $E(r) \rightarrow \Gamma((2^r), L)$ (see Lemma 2.9, (i)). If E is irreducible, then by Corollary 2.7 its image generates $J^2(C)$, hence also $\Gamma((2^r), L) = Z$ generates $J^{2r}(C)$ and we obtain a contradiction.

So E is reducible and we apply Lemma 2.8 and 2.9. Suppose we are in case (b) of Lemma 2.8 and (ii) of Lemma 2.9. Then $A = Z = \Gamma((2^r), L)$ is birational to $E_1(r_1) \times E_2(r_2)$, for non-negative integers r_1, r_2 . However, if $r_i > 0$, then $r_i = g_i$, for $i \in \{1, 2\}$. Since $2g_2 \geq g$, one has $r_2 = 0$ because $2r = g - 1 < g$. Hence A is birational to $E_1(g_1)$ and we are in case (i).

Consider now case (c) of Lemma 2.8 and (iii) of Lemma 2.9. We claim that $J(C)$ is isogenous to $J(E_1) \times J(E_2) \times J(E_3)$. Indeed, consider the representation of \mathbb{Z}_2^2 on $H^0(K_C)$. Since C/\mathbb{Z}_2^2 is

rational, we have $H^0(K_C) = V_{\chi_1} \oplus V_{\chi_2} \oplus V_{\chi_3}$, where for $i = 1, 2, 3$ the non trivial character χ_i of \mathbb{Z}_2^2 is orthogonal to the involution ι_i such that $C/\iota_i \cong E_i$, and \mathbb{Z}_2^2 acts on V_{χ_i} as multiplication by χ_i . Thus V_{χ_i} is the tangent space to $f_i^*(J(E_i))$.

Recall that there are three non–negative integers r_1, r_2, r_3 such that A is birational to $E_1(r_1) \times E_2(r_2) \times E_3(r_3)$. If $r_i > 0$, then $r_i = g_i$ (for $1 \leq i \leq 3$). Since $g_1 + g_2 + g_3 = g = 2r + 1$, at least one of the integers r_i is zero. If two of them are zero, we are again in case (i). So assume $r_1 = 0$ and r_2, r_3 non–zero. Then $r = g_2 + g_3$ and $g_1 = r + 1$. Moreover the involution ι_1 acts on A as multiplication by -1 , hence A is the Prym variety of $f_1: C \rightarrow E_1$, and we are in case (ii).

The case C hyperelliptic can be treated similarly. In case (b) of Lemma 2.8 and (ii) of Lemma 2.9, E_1 is rational (hence it is contracted to a point by j), and $A = Z = \Gamma((2^r), L)$ is birational to $E_2(r)$. Then $r = g_2$ and we reach a contradiction since $2g_2 \geq g$. In case (c) of Lemma 2.8 and (iii) of Lemma 2.9 one has $g_3 = 0$ and the above argument applies with no change. \square

Proof of Theorem 3.1. By Lemma 3.7, we may assume $\nu = 2$. Let \underline{m} be the partition of $2r$ such that $Z \subseteq \Gamma(\underline{m}, L)$. By Lemma 3.6, it is enough to consider the case $\nu = 2$ and $\underline{m} = (1^{g-1})$, i.e., case (i) of that lemma. Then Z is contained in the kernel P of $f_*: J^{g-1}(C) \rightarrow J^{g-1}(C')$ (namely P is the generalized Prym variety associated with f). The space $H^0(K_C)$ decomposes under the involution ι associated with f as $H^0(K_{C'}) \oplus V$, where V is the space of antiinvariant 1-forms. Hence V^* is the tangent space to P , the tangent space T to A is also contained in V^* and the linear series L is equal to $\mathbb{P}(T^\perp) \supseteq \mathbb{P}(H^0(K_{C'}))$. On the other hand, by construction ι acts trivially on L , hence $T = V^*$ and thus $A = P$. This implies that $Z = A = P$, $g = 2r + 1$ and f is unramified with $g' = r + 1$. \square

3.2. General Brill–Noether loci. The proof of the general formula (3.1) in [7] uses an argument which is useful to briefly recall. Let $Z \subseteq W_d^s(C) \subsetneq J^d(C)$ be A –stable and $L \in Z$ general, so L is a special g_d^s , which we may assume to be complete, so $d \geq 2s$. Let F_L be the fixed divisor of L and, if $s > 0$, let L' be the base point free residual linear series. Set $d' = \deg(L')$. Since L' is also special, we have $d' \geq 2s$. Then, by Lemma 2.8, (iii), one has the birational map $\tau: C(d' - s, L') \dashrightarrow C(s, L') \cong C(s)$.

Consider the morphism $j: C(d' - s, L') \times C(g - 1 - d + s) \rightarrow J^{g-1}(C)$ such that $j(D, D')$ is the class of $D + F_L + D'$ (if $s = 0$, we define $j: C(g - 1 - d) \rightarrow J^{g-1}(C)$ by $D' \mapsto F_L + D'$). If (D, D') is general in $C(d' - s, L') \times C(g - 1 - d + s)$, the divisor $D + F_L + D'$ is linearly isolated, hence j is generically finite onto its image Z_L , which therefore has dimension $g - 1 - d + 2s$. Consider the closure Z' of the union of the Z_L 's, with $L \in Z$ general, which is A –stable. One has $Z' \subseteq \Theta$ and the above discussion yields $r' := \dim(Z') = r + g - 1 - d + 2s$. Therefore $r' + a \leq g - 1$ if and only if (3.1) holds, with equality if and only if equality holds in (3.1).

Corollary 3.9. *Let C be a curve of genus g . Let $A \subsetneq J(C)$ be an abelian variety of dimension $a > 0$ and $Z \subseteq W_d^s(C) \subsetneq J^d(C)$ an irreducible, A –stable variety of dimension $r = d - 2s - a$. Assume $(d, s) \neq (g - 1, 0)$. Then there is a degree 2 morphism $\varphi: C \rightarrow C'$, with C' a smooth curve of genus a with $g > 2a + 1$, such that $A = \varphi^*(J(C'))$ and either*

- (i) $Z = W_{d-2a-2s}(C) + \varphi^*(J^{a+s}(C'))$, or
- (ii) C is hyperelliptic and $Z = \varphi^*(J^a(C')) + W_{d-2s-2a}(C) + W_{2s}^s(C)$.

Proof. We keep the same notation as above.

We apply Theorem 3.1 to Z' . Since $d = r + a + 2s \geq 2(a + s)$ and $(d, s) \neq (g - 1, 0)$, then $2a < g - 1$, hence case (b) of Theorem 3.1 does not occur for Z' . So we have a degree 2 morphism $\varphi: C \rightarrow C'$, with C' a smooth curve of genus a with $g > 2a + 1$, such that $A = \varphi^*(J(C'))$ and $Z' = W_{g-1-2a}(C) + \varphi^*(J^a(C'))$. The case $s = 0$ follows right away and we are in case (i). So we assume $s > 0$ from now on.

If L' is composed with the involution ι determined by φ , we are again in case (i). So assume L' is not composed with ι . The general $D \in C(d' - s, L')$ contains no fibre of φ and the same happens for the general $D' \in C(g - 1 - d + s)$. Since $D + F_L + D'$ corresponds to the general point of Z , and contains exactly a general fibres of φ , then F_L has to contain these a fibres, whose union we denote by F . Moreover, the description of Z implies that $D + D' + (F_L - F)$ is a general divisor of degree $g - 1 - 2a$, in particular D is a general divisor of degree $d' - s$. But D is also general in $C(d' - s, L')$, and this implies $d' - s \leq s$. On the other hand $d' \geq 2s$, hence $d' = 2s$. Then, either L' is the canonical series of C or C is hyperelliptic and L' is the s -multiple of the g_2^1 . However the former case does not occur since by construction $d' = \deg(L') \leq g - 1$, hence we are in case (ii). \square

REFERENCES

- [1] D. Abramovich, J. Harris, *Abelian varieties and curves in $W_d(C)$* , *Compositio Mathematica*, **78** (1991), 227–238.
- [2] E. Arbarello, M. Cornalba, P.A. Griffiths, J. Harris *Geometry of algebraic curves* Vol. I. Grundlehren der Mathematischen Wissenschaften, **267**, Springer-Verlag, New York, 1985.
- [3] C. Birkenhake, H. Lange, *Complex abelian varieties*, Second edition, Grundlehren in math. Wissens. **302** Springer-Verlag (2004).
- [4] C. Ciliberto, *On a proof of Torelli's theorem*, in Proceedings of the Conference “Problems in Algebraic Geometry”, Ravello, 1982, Springer Lecture Notes in Math. **997** (1983), 112-123.
- [5] C. Ciliberto, M. Mendes Lopes, R. Pardini, *The classification of minimal irregular surfaces of general type with $K^2 = 2p_g$* , preprint.
- [6] O. Debarre, *Inégalités numériques pour les surfaces de type général*, *Bull. Soc. Math. de France*, **110** (1982), 319–342.
- [7] O. Debarre, R. Fahlouai, *Abelian varieties in $W_d^r(C)$ and points of bounded degree on algebraic curves*, *Compositio Mathematica*, **88** (1993), 235–249.
- [8] O. Debarre, appendix to *Irreducibility, Brill-Noether loci and Vojta's inequality*, *Trans. Amer. Math. Soc.* **354**, no. 8 (2002), 3011–3029.
- [9] G. Faltings, *Diofantine approximation on abelian varieties*, *Ann. of Math.*, **133** (1991), 549–576.
- [10] W. Fulton, *Intersection theory*, second edition, Springer-Verlag Berlin Heidelberg (1998).

DIPARTIMENTO DI MATEMATICA, II UNIVERSITÀ DI ROMA, ITALY
E-mail address: `cilibert@xp.mat.uniroma2.it`

DEPARTAMENTO DE MATEMÁTICA, INSTITUTO SUPERIOR TÉCNICO, UNIVERSIDADE TÉCNICA DE LISBOA AV. ROVISCO PAIS, 1049-001 LISBOA, PORTUGAL
E-mail address: `mmlopes@math.ist.utl.pt`

DIPARTIMENTO DI MATEMATICA UNIVERSITÀ DI PISA, LARGO B. PONTECORVO, 5, 56127 PISA, ITALY
E-mail address: `pardini@dm.unipi.it`