

GROTHENDIECK'S THEOREM FOR ABSOLUTELY SUMMING MULTILINEAR OPERATORS IS OPTIMAL

DANIEL PELLEGRINO AND JUAN B. SEOANE-SEPÚLVEDA

ABSTRACT. Grothendieck's theorem asserts that every continuous linear operator from ℓ_1 to ℓ_2 is absolutely $(1; 1)$ -summing. In this note we prove that the optimal constant g_m so that every continuous m -linear operator from $\ell_1 \times \cdots \times \ell_1$ to ℓ_2 is absolutely $(g_m; 1)$ -summing is $\frac{2}{m+1}$. This result solves (in the positive) a conjecture posed by A.T. Bernardino in 2011.

1. INTRODUCTION

A celebrated result of Grothendieck asserts that every continuous linear operator from ℓ_1 to ℓ_2 is absolutely $(1; 1)$ -summing. It was recently proved [1] that this result can be lifted to multilinear operators in the following fashion:

Every continuous m -linear operator from $\ell_1 \times \cdots \times \ell_1$ to ℓ_2 is absolutely $\left(\frac{2}{m+1}; 1\right)$ -summing.

In the same paper the author conjectures that the value $\frac{2}{m+1}$ is optimal. In this note we remark that this statement is true. Our proof is inspired on ideas that date back to the classical work of Lindenstrauss and Pełczyński [6] and, later, explored in a series of papers (see, e.g., [2, 3, 4, 7]).

Throughout this note, X and Y shall stand for Banach spaces over $\mathbb{K} = \mathbb{R}$ or \mathbb{C} . The closed unit ball of X is denoted by B_X and the topological dual of X by X^* . Also, recall that a linear operator $u : X \rightarrow Y$ is absolutely $(q; 1)$ -summing (see [5]) if there exists $C \geq 0$ such that

$$\left(\sum_{j=1}^n \|u(x_j)\|^q \right)^{\frac{1}{q}} \leq C \sup_{\varphi \in B_{X^*}} \sum_{j=1}^n |\varphi(x_j)|$$

for every $n \in \mathbb{N}$ and $x_1, \dots, x_n \in X$. The nonlinear theory of absolutely summing operators was designed by Pietsch in 1983 ([8]) and since then has been intensively investigated. One of the possible polynomial generalizations of absolutely summing operators is the concept of absolutely summing polynomial. Given a positive integer m and $q \geq \frac{1}{m}$, an m -homogeneous polynomial $P : X \rightarrow Y$ is absolutely $(q; 1)$ -summing if there exists a constant $C \geq 0$ such that

$$\left(\sum_{j=1}^n \|P(x_j)\|^q \right)^{\frac{1}{q}} \leq C \left(\sup_{\varphi \in B_{X^*}} \sum_{j=1}^n |\varphi(x_j)| \right)^m$$

for every $n \in \mathbb{N}$ and $x_1, \dots, x_n \in X$. The infimum of the constants C satisfying the above inequality is a norm for the space of absolutely $(q, 1)$ -summing polynomials from X to Y , and it is denoted by $\pi_{q,1}$. For multilinear mappings the definition is similar:

An m -linear operator $T : X \times \cdots \times X \rightarrow Y$ is absolutely $(q; 1)$ -summing (with $q \geq \frac{1}{m}$) if there is a constant $C \geq 0$ such that

$$\left(\sum_{j=1}^n \|T(x_j^{(1)}, \dots, x_j^{(m)})\|^q \right)^{\frac{1}{q}} \leq C \prod_{k=1}^m \left(\sup_{\varphi \in B_{X^*}} \sum_{j=1}^n |\varphi(x_j^{(k)})| \right)$$

for every $n \in \mathbb{N}$ and $x_1^{(k)}, \dots, x_n^{(k)} \in X$, and $k = 1, \dots, m$.

2. THE PROOF OF THE OPTIMALITY OF $\frac{2}{m+1}$

Suppose that every continuous m -linear operator from $\ell_1 \times \cdots \times \ell_1$ to ℓ_2 is absolutely $(q_m; 1)$ -summing for certain $q_m < 2$. Then, from the very definitions it is plain that every continuous m -homogeneous polynomial from ℓ_1 to ℓ_2 is absolutely $(q_m; 1)$ -summing. The following argument is mentioned *en passant*, without proof, in [4], but since we have a more self-contained approach, we present the details for the sake of completeness. Let $n \in \mathbb{N}$ and $x_1, \dots, x_n \in \ell_1$ be given. Consider $x_1^*, \dots, x_n^* \in B_{\ell_\infty}$ so that $x_j^*(x_j) = \|x_j\|$ for every $j = 1, \dots, n$. Let a_1, \dots, a_n be scalars such that $\sum_{j=1}^n |a_j|^{\frac{2}{q_m}} = 1$ and define

$$P_n : \ell_1 \longrightarrow \ell_2, \quad P_n(x) = \sum_{j=1}^n |a_j|^{\frac{1}{q_m}} x_j^*(x)^m e_j,$$

where e_j is the j -th canonical vector of ℓ_2 . For every $x \in \ell_1$,

$$\|P_n(x)\| = \left(\sum_{j=1}^n \left| |a_j|^{\frac{1}{q_m}} x_j^*(x)^m \right|^2 \right)^{\frac{1}{2}} \leq \left(\sum_{j=1}^n |a_j|^{\frac{2}{q_m}} \right)^{\frac{1}{2}} \|x\|^m = \|x\|^m.$$

By the *Open Mapping Theorem* there exists $K > 0$ (not depending on n) such that $\pi_{q_m, 1}(P_n) \leq K \|P_n\| \leq K$. Note that for $k = 1, \dots, n$, we have

$$\|P_n(x_k)\| = \left\| \sum_{j=1}^n |a_j|^{\frac{1}{q_m}} x_j^*(x_k)^m e_j \right\| \geq |a_k|^{\frac{1}{q_m}} x_k^*(x_k)^m = |a_k|^{\frac{1}{q_m}} \|x_k\|^m.$$

To simplify the notation we write $\|(x_j)_{j=1}^n\|_{w, 1} := \sup_{\varphi \in B_{X^*}} \sum_{j=1}^n |\varphi(x_j^{(k)})|$. Thus, we have

$$\begin{aligned} \left(\sum_{j=1}^n \|x_j\|^{mq_m} |a_j| \right)^{\frac{1}{q_m}} &= \left(\sum_{j=1}^n \left(\|x_j\|^m |a_j|^{\frac{1}{q_m}} \right)^{q_m} \right)^{\frac{1}{q_m}} \\ &\leq \left(\sum_{j=1}^n \|P_n(x_j)\|^{q_m} \right)^{\frac{1}{q_m}} \\ &\leq \pi_{q_m, 1}(P_n) \|(x_j)_{j=1}^n\|_{w, 1}^m. \end{aligned}$$

Since this last inequality holds whenever $\sum_{j=1}^n |a_j|^{\frac{2}{q_m}} = 1$, we obtain

$$\begin{aligned} \left(\sum_{j=1}^n \|x_j\|^{mq_m \left(\frac{2}{q_m}\right)^*} \right)^{\frac{1}{\left(\frac{2}{q_m}\right)^*}} &\leq \sup \left\{ \sum_{j=1}^n |a_j| \|x_j\|^{mq_m}; \sum_{j=1}^n |a_j|^{\frac{2}{q_m}} = 1 \right\} \\ &\leq (\pi_{q_m,1}(P_n) \|(x_j)_{j=1}^n\|_{w,1}^m)^{q_m} \\ &\leq (K \|(x_j)_{j=1}^n\|_{w,1}^m)^{q_m} \end{aligned}$$

and, then,

$$\left(\sum_{j=1}^n \|x_j\|^{mq_m \left(\frac{2}{q_m}\right)^*} \right)^{\frac{1}{mq_m \left(\frac{2}{q_m}\right)^*}} \leq K^{\frac{1}{m}} \|(x_j)_{j=1}^n\|_{w,1}.$$

We conclude that id_{ℓ_1} is $\left(mq_m \left(\frac{2}{q_m}\right)^*; 1\right)$ -summing. Finally, from the *Dvoretzky-Rogers Theorem* we must have

$$\frac{2mq_m}{2 - q_m} = \left(\frac{2}{q_m}\right)^* mq_m \geq 2,$$

that is,

$$q_m \geq \frac{2}{m+1}.$$

REFERENCES

- [1] A.T. Bernardino, On cotype and a Grothendieck-type theorem for absolutely summing multilinear operators, *Quaest. Math.* **34** (2011), 513–519.
- [2] G. Botelho and D. Pellegrino, Scalar-valued dominated polynomials on Banach spaces, *Proc. Amer. Math. Soc.* **134** (2006), 1743–1751.
- [3] G. Botelho and D. Pellegrino, Absolutely summing polynomials on Banach spaces with unconditional basis, *J. Math. Anal. Appl.* **321** (2006), 50–58.
- [4] G. Botelho, D. Pellegrino, and P. Rueda, Cotype and absolutely summing linear operators. *Math. Z.* **267** (2011), no. 1–2, 1–7.
- [5] J. Diestel, H. Jarchow, and A. Tonge, *Absolutely summing operators*, Cambridge University Press, 1995.
- [6] J. Lindenstrauss and A. Pełczyński, Absolutely summing operators in \mathcal{L}_p spaces and their applications, *Studia Math.* **29** (1968), 275–326.
- [7] D. Pellegrino, Cotype and absolutely summing homogeneous polynomials in \mathcal{L}_p spaces, *Studia Math.* **157** (2003), 121–131.
- [8] A. Pietsch, *Ideals of Multilinear Functionals*. Proc. II Intem. Conf. on Operator Algebras, Ideals and Appl. in Theoretical Physics. Leipzig (1983) Teubner-Texte.

DEPARTAMENTO DE MATEMÁTICA,
UNIVERSIDADE FEDERAL DA PARAÍBA,
58.051-900 - JOÃO PESSOA, BRAZIL.
E-mail address: pellegrino@pq.cnpq.br and dmpellegrino@gmail.com

DEPARTAMENTO DE ANÁLISIS MATEMÁTICO,
FACULTAD DE CIENCIAS MATEMÁTICAS,
PLAZA DE CIENCIAS 3,
UNIVERSIDAD COMPLUTENSE DE MADRID,
MADRID, 28040, SPAIN.
E-mail address: jseoane@mat.ucm.es