ON COHOMOLOGICAL DIMENSION AND DEPTH UNDER LINKAGE

M. EGHBALI AND N. SHIRMOHAMMADI

ABSTRACT. Some relations between cohomological dimensions and depths of linked ideals are investigated and discussed by various examples.

1. INTRODUCTION

Throughout this paper, we assume that *R* is a commutative Noetherian ring and \mathfrak{a} is an ideal of *R*. Let *M* be an *R*-module. For an integer $i \in \mathbb{Z}$, let $H^i_{\mathfrak{a}}(M)$ denote the *i*th local cohomology module of *M* with respect to \mathfrak{a} as introduced by Grothendieck (cf. [12] and [2]).

One of the most interested subjects in commutative algebra and related topics is the cohomological dimension of an ideal. The cohomological dimension of a in R, cd(a, R) is defined as

$$cd(\mathfrak{a}, R) = \min\{i : H'_{\mathfrak{a}}(R) = 0 \text{ for all } j > i\}.$$

In the light of Grothendieck's Vanishing Theorem, $cd(\mathfrak{a}, R) \leq \dim R$ and the equality happens when (R, \mathfrak{m}) is a local ring and \mathfrak{a} is an \mathfrak{m} -primary ideal. The Hartshorne-Lichtenbaum Vanishing Theorem provides conditions for $cd(\mathfrak{a}, R) \leq \dim R - 1$, where \mathfrak{a} is not an \mathfrak{m} -primary ideal. The cohomological dimension has been studied by several authors; see for instance [10], [13], [17], [14], [16], [6] and [20].

Let (R, \mathfrak{m}) be a Gorenstein local ring. Two ideals \mathfrak{a} and \mathfrak{b} of R are linked by a Gorenstein ideal \mathfrak{c} if $\mathfrak{a} = \mathfrak{c} : \mathfrak{b}$ and $\mathfrak{b} = \mathfrak{c} : \mathfrak{a}$. Recall that an ideal \mathfrak{c} is a Gorenstein ideal if R/\mathfrak{c} is a Gorenstein ring. From now on we mean \mathfrak{a} and \mathfrak{b} are linked when they are linked by a Gorenstein ideal. Iterating the procedure, we say that an ideal \mathfrak{b} is in the linkage class of an ideal \mathfrak{a} if there are ideals $\mathfrak{c}_1, \ldots, \mathfrak{c}_n$ such that \mathfrak{b} is linked to $\mathfrak{c}_1, \mathfrak{c}_1$ is linked to \mathfrak{c}_2, \ldots , and \mathfrak{c}_n is linked to \mathfrak{a} . If, in addition, n is odd, we say that \mathfrak{b} is in the even linkage class of \mathfrak{a} , that is \mathfrak{b} is evenly linked to \mathfrak{a} . In case \mathfrak{a} is an unmixed ideal and $\mathfrak{c} \subset \mathfrak{a}$ is a Gorenstein ideal with ht $\mathfrak{a} = \mathfrak{ht} \mathfrak{c}$, then \mathfrak{a} and $\mathfrak{c} : \mathfrak{a}$ are linked by \mathfrak{c} (cf. [19, Proposition 2.2]).

Below, we recall some of the known results about the linkage that will be used in the course of the paper. We refer the reader to [19] or [21].

Lemma 1.1. Suppose that (R, \mathfrak{m}) is a Gorenstein local ring, and \mathfrak{a} and \mathfrak{b} are two linked ideals of R by a Gorenstein ideal \mathfrak{c} .

- (a) There exists a short exact sequence $0 \longrightarrow K_{R/\mathfrak{a}} \longrightarrow R/\mathfrak{c} \longrightarrow R/\mathfrak{b} \longrightarrow 0$, where $K_{R/\mathfrak{a}}$ denotes the canonical module of R/\mathfrak{a} .
- (b) If R/\mathfrak{a} is a Cohen-Macaulay ring, then so is R/\mathfrak{b} .

²⁰⁰⁰ Mathematics Subject Classification. 13D45, 13C14.

Key words and phrases. Cohomological dimension, depth, linkage, formal grade.

The first author is supported in part by a grant from IPM (No. 91130026).

One of the classical and long-standing problems in linkage theory is to determine what kind of certain properties of ideals which are preserved under linkage. The main purpose of this paper is to give answer to the following question.

Question 1.2. Suppose that (R, \mathfrak{m}) is a Gorenstein local ring, and \mathfrak{a} and \mathfrak{b} are two linked ideals of *R*. When do the equalities depth $R/\mathfrak{a} = \operatorname{depth} R/\mathfrak{b}$ and $\operatorname{cd}(\mathfrak{a}, R) = \operatorname{cd}(\mathfrak{b}, R)$ hold?

Schenzel [20] has studied the formal local cohomology modules $\mathfrak{F}^{i}_{\mathfrak{a}}(M) := \varprojlim_{n} H^{i}_{\mathfrak{m}}(M/\mathfrak{a}^{n}M)$ for a finitely generated module *M* over a local ring (R, \mathfrak{m}) and $i \in \mathbb{Z}$. It is called *i*th formal local cohomology of *M* with respect to \mathfrak{a} . In the case of a Gorenstein local ring

(1.1)
$$\mathfrak{F}^{i}_{\mathfrak{a}}(R) \cong \operatorname{Hom}_{R}(H^{\dim R-i}_{\mathfrak{a}}(R), E),$$

where *E* is the injective hull of R/\mathfrak{m} (cf. [20, Remark 3.6]). For more information on these kind of modules and their properties we refrer the reader to [1] and [7].

The first non-vanishing value of $\mathfrak{F}^{i}_{\mathfrak{a}}(M)$ is called the formal grade as follows

fgrade(
$$\mathfrak{a}, M$$
) := inf{ $i \in \mathbb{Z} : \mathfrak{F}^{i}_{\mathfrak{a}}(M) \neq 0$ }.

In case (R, \mathfrak{m}) is a Cohen-Macaulay local ring, then the equality

(1.2)
$$\operatorname{cd}(\mathfrak{a}, R) = \dim R - \operatorname{fgrade}(\mathfrak{a}, R)$$

holds, (cf. [1, Corollary 4.2]).

Recently formal local cohomology has been used as a technical tool to solve some problems, see for instance [8]. In this paper we use it to give information on the cohomological dimension of an ideal.

The outline of the paper is as follows. In Section 2, we consider the cohomological dimension of an ideal and a partial reverse statement of a result of Peskin-Szpiro. In Section 3, we give some answers to the Question 1.2 sorted below.

For the equality of cohomological dimensions:

- There is an example of a regular local ring possessing two Cohen-Macaulay linked ideals while their cohomological dimensions are not the same (cf. Example 3.1).
- For a regular local ring of positive characteristic with Cohen-Macaulay linked ideals, we give an affirmative answer (cf. Theorem 3.8(a)).
- Another positive answer happens for evenly linked squarefree monomial ideals in the ring *k*[*x*₁, ..., *x*_{*n*}]_{(*x*₁,...,*x*_{*n*})</sup> (cf. Theorem 3.8(c)).}

For the equality of depths:

- In a Gorenstein local ring, the depths of residue class rings of evenly linked ideals are the same(cf. Corollary 3.4).
- We give negative answers in Corollary 3.5 and Corollary 3.6.
- In a Gorenstein local ring, every 2-dimensional non Cohen-Macaulay linked ideals provide an affirmative answer (cf. Corollary 3.7).

2. Some remarks on cohomological dimension

Throughout this section, we assume that (R, \mathfrak{m}) is a local ring and \mathfrak{a} is an ideal of R. The first non-vanishing cohomological degree of the local cohomology modules $H^i_{\mathfrak{a}}(R)$ is well understood. It is the common length of maximal R-regular sequences in \mathfrak{a} . The last non-vanishing amount of local cohomology modules, instead, is more mysterious. In the light of Grothendieck's Vanishing Theorem, $cd(\mathfrak{a}, R) \leq \dim R$ and the equality happens when \mathfrak{a} is an \mathfrak{m} -primary ideal. In the following we characterize the last non-vanishing amount of local cohomology modules.

Proposition 2.1. Let a be an ideal of a local ring R and N be a finite R-module. Let t be a positive integer. *Then the following are equivalent:*

- (a) $H^i_{\mathfrak{a}}(N) = 0$ for all i > t,
- (b) $H^i_{\mathfrak{a}}(N)$ is finite for all i > t,
- (c) $\mathfrak{a} \subseteq \operatorname{Rad}(0:_R H^i_\mathfrak{a}(N))$ for all i > t.

Proof. The implications $(a) \implies (b)$ and $(b) \implies (c)$ are clear. In order to prove $(c) \implies (a)$, we argue by induction on $d := \dim N$. For d = 0, the claim is clear. Now assume that d > 0 and $\mathfrak{a} \subseteq \operatorname{Rad}(0:_R H^i_\mathfrak{a}(N))$ for all i > t. We are going to show that $H^i_\mathfrak{a}(N) = 0$ for all i > t. First note that $H^i_\mathfrak{a}(N/H^0_\mathfrak{a}(N)) \cong H^i_\mathfrak{a}(N)$ for all i > t(> 0) (cf. [2]). So, we may assume that $H^0_\mathfrak{a}(N) = 0$. Then there exists an *N*-regular element *r* in \mathfrak{a} . Consider the short exact sequence

$$0 \to N \xrightarrow{r} N \to N/rN \to 0.$$

which implies the following long exact sequence

(2.1)
$$\cdots \to H^i_{\mathfrak{a}}(N) \xrightarrow{r} H^i_{\mathfrak{a}}(N) \to H^i_{\mathfrak{a}}(N/rN) \to H^{i+1}_{\mathfrak{a}}(N) \to \cdots$$

for all *i*. In the light of 2.1 and the induction assumption, one can deduce that $H^i_{\mathfrak{a}}(N/rN) = 0$ for all i > t since dim N/rN = d - 1. Again using 2.1, we have $H^i_{\mathfrak{a}}(N) = rH^i_{\mathfrak{a}}(N)$ for all i > t. As $\mathfrak{a} \subseteq \operatorname{Rad}(0:_R H^i_{\mathfrak{a}}(N))$, one has $r^u H^i_{\mathfrak{a}}(N) = 0$ for some integer *u*, and so $H^i_{\mathfrak{a}}(N) = 0$ for all i > t.

It should be noted that it has been shown in [22, Proposition 3.1] that statements (a) and (b) in Proposition 2.1 are equivalent.

In the light of Proposition 2.1, we have

$$cd(\mathfrak{a}, R) = \sup\{i \mid H^{i}_{\mathfrak{a}}(R) \text{ is not finite}\}$$
$$= \sup\{i \mid \mathfrak{a} \not\subseteq \operatorname{Rad}(0:_{R} H^{i}_{\mathfrak{a}}(R))\}.$$

In their valuable paper, Peskine-Szpiro (cf. [18, Chap. IV]) have proved the following result.

Remark 2.2. Suppose (R, \mathfrak{m}) is a regular local ring of positive characteristic and t is a positive integer. If depth $R/\mathfrak{a} \ge t$, then $cd(\mathfrak{a}, R) \le \dim R - t$.

Proof. The claim is clear by virtue of [8, Remark 3.1] and the equality (1.2).

Next, we prove a slightly reverse statement of the above mentioned result of Peskin-Szpiro.

Lemma 2.3. Let $R = k[x_1, ..., x_n]$ be a polynomial ring in n variables over a field k with maximal ideal $\mathfrak{m} := (x_1, ..., x_n)$ and \mathfrak{a} be a squarefree monomial ideal of R. Assume that t is a non-negative integer. If $\operatorname{cd}(\mathfrak{a}, R) \leq n - t$, then depth $R/\mathfrak{a} \geq t$.

Proof. Since each of the modules is graded, so the issue of vanishing is unchanged under localization at m. Suppose that $cd(\mathfrak{a}, R) \leq n - t$. This means $H^i_{\mathfrak{a}}(R) = 0$ for all i > n - t. Hence, by (1.1) we have $\mathfrak{F}^j_{\mathfrak{a}}(R) = 0$ for all j < t. So by virtue of [8, Corollary 4.2], we have depth $R/\mathfrak{a} = \text{fgrade}(\mathfrak{a}, R) \geq t$.

Corollary 2.4. Suppose that $R = k[x_1, ..., x_n]_{(x_1,...,x_n)}$ is a polynomial ring over a field k of positive characteristic localized at $(x_1, ..., x_n)$ with a is a squarefree monomial ideal. Then

- (a) If $cd(\mathfrak{a}, R) \leq cd(\mathfrak{b}, R)$, then depth $R/\mathfrak{a} \geq depth R/\mathfrak{b}$ for any ideal \mathfrak{b} of R.
- (b) If depth $R/\mathfrak{a} \ge \operatorname{depth} R/\mathfrak{b}$, then $\operatorname{cd}(\mathfrak{a}, R) \le \operatorname{cd}(\mathfrak{b}, R)$ for any squarefree monomial ideal \mathfrak{b} of R.
- (c) $cd(\mathfrak{a}, R) = cd(\mathfrak{b}, R)$ if and only if depth $R/\mathfrak{a} = depth R/\mathfrak{b}$ for any squarefree monomial ideal \mathfrak{b} of R.

Proof. Assume that $cd(\mathfrak{a}, R) \leq cd(\mathfrak{b}, R)$. From Lemma 2.3, the equality (1.2) and [8, Remark 3.1] we have depth $R/\mathfrak{a} \geq n - cd(\mathfrak{b}, R) = \text{fgrade}(\mathfrak{b}, R) \geq \text{depth } R/\mathfrak{b}$. This proves (a). In order to prove (b), assume that depth $R/\mathfrak{a} \geq \text{depth } R/\mathfrak{b}$. To the contrary, suppose that $cd(\mathfrak{a}, R) > cd(\mathfrak{b}, R)$. It implies that $n - \text{depth } R/\mathfrak{a} > cd(\mathfrak{b}, R)$. This in conjunction with Lemma 2.3 yields depth $R/\mathfrak{b} > \text{depth } R/\mathfrak{a}$ which is a contradiction. Now, (c) follows from (a) and (b).

3. STABILITY OF COHOMOLOGICAL DIMENSION UNDER LINKAGE

Let a and b be ideals of a local ring (R, \mathfrak{m}) . In this section, we try to find out some conditions for the stability of cohomological dimension under linkage, i.e. $cd(\mathfrak{a}, R) = cd(\mathfrak{b}, R)$ whenever a is linked to b. It should be noted that the cohomological dimension is not preserved under linkage, in general. To be more precise, consider the next example.

Example 3.1. Let $X = (x_{ij})$ be the generic 4×3 matrix and X' be the result of dropping the first two rows from X. Let $R = k[X]_{(X)}$ be a polynomial ring over a field k localized at the maximal ideal (X). Put $\mathfrak{a} := I_3(X)$ (the ideal generated by the 3-minors of X in R) and $\mathfrak{b} := I_2(X')$ (the ideal generated by the 2-minors of X' in R). By virtue of [9, Proposition 21.24] we see that \mathfrak{a} and \mathfrak{b} are Cohen-Macaulay linked ideals, but $cd(\mathfrak{a}, R) = 4$ while $cd(\mathfrak{b}, R) = 3$ (cf. [4, Corollary, pp. 440]).

According to Corollary 2.4 and Corollary 4.1 of [8], under certain conditions, we see some relations between $cd(\mathfrak{a}, R)$, depth R/\mathfrak{a} and $fgrade(\mathfrak{a}, R)$.

One of the technical tools we use to show the stability of cohomological dimension under linkage is the depth, i.e. we investigate the equality depth $R/\mathfrak{a} = \operatorname{depth} R/\mathfrak{b}$. Take into account that from the stability of depth one can not deduce the stability of cohomological dimension under linkage, in general, as the following concrete example shows it.

Example 3.2. Let $R = k[x_0, x_1, x_2, x_3]$ be a polynomial ring over a field k. Let $\mathfrak{a} = (x_0, x_1) \cap (x_2, x_3)$ be the defining ideal of the union of the two skew lines in \mathbb{P}^3 and $\mathfrak{b} = (x_0x_3 - x_1x_2, x_1^3 - x_0^2x_3, x_2^3 - x_1x_3^2)$ be the defining ideal of the twisted quartic curve in \mathbb{P}^3 . Then it is not hard to show that

$$\mathfrak{a} \cap \mathfrak{b} = (x_0 x_3 - x_1 x_2, x_0 x_2^2 - x_1^2 x_3)$$

is a complete intersection. Therefore b is linked to a by $\mathfrak{a} \cap \mathfrak{b}$. Using CoCoA [5], we see that depth(R/\mathfrak{a}) = 1 = depth(R/\mathfrak{b}). But cd(\mathfrak{a} , R) = 3 and cd(\mathfrak{b} , R) = 2.

According to the above examples and what we will see in the sequel, the cohomological dimension, formal grade and depth are not stable, in general, under linkage.

Proposition 3.3. Suppose that (R, \mathfrak{m}) is a Gorenstein local ring, and \mathfrak{a} and \mathfrak{b} are two linked ideals of R with $t := \dim(R/\mathfrak{a})$. Then $\operatorname{depth}(R/\mathfrak{b}) = \operatorname{depth}(K_{R/\mathfrak{a}}) - 1$ provided that R/\mathfrak{a} is not a Cohen-Macaulay ring.

Proof. By the assumption, we have the following exact sequence

$$0 \longrightarrow K_{R/\mathfrak{a}} \longrightarrow R/\mathfrak{c} \longrightarrow R/\mathfrak{b} \longrightarrow 0.$$

The above exact sequence together with [3, Proposition 1.2.9] yield the following inequalities

(3.1)
$$\operatorname{depth}(K_{R/\mathfrak{a}}) \geq \min\{\operatorname{depth}(R/\mathfrak{c}), \operatorname{depth}(R/\mathfrak{b}) + 1\},\$$

(3.2) $\operatorname{depth}(R/\mathfrak{b}) \geq \min\{\operatorname{depth}(K_{R/\mathfrak{a}}) - 1, \operatorname{depth}(R/\mathfrak{c})\}.$

Note that, by Lemma 1.1(b), R/\mathfrak{b} is not Cohen-Macaulay. So using 3.1, one can deduce that depth($K_{R/\mathfrak{a}}$) $\geq \min\{t, \operatorname{depth}(R/\mathfrak{b}) + 1\} = \operatorname{depth}(R/\mathfrak{b}) + 1$. On the other hand, by 3.2, we have depth(R/\mathfrak{b}) $\geq \min\{\operatorname{depth}(K_{R/\mathfrak{a}}) - 1, t\} = \operatorname{depth}(K_{R/\mathfrak{a}}) - 1$. Therefore depth(R/\mathfrak{b}) $= \operatorname{depth}(K_{R/\mathfrak{a}}) - 1$.

Corollary 3.4. *Suppose that* (R, \mathfrak{m}) *is a Gorenstein local ring, and* \mathfrak{a} *and* \mathfrak{b} *are evenly linked ideals of* R. *Then* depth $(R/\mathfrak{a}) = depth(R/\mathfrak{b})$.

There exists a Buchsbaum quasi-Gorenstein domain A which is a homomorphic image of a polynomial ring over a field k with 11 indeterminates such that dim A = 3 and depth A = 2 (cf. [15]). Recall that a local ring (A, \mathfrak{n}) is quasi-Gorenstein if $H_{\mathfrak{n}}^{\dim A}(A) \cong E_A(A/\mathfrak{n})$ the injective hull of A/\mathfrak{n} . Now, we have a prime ideal \mathfrak{p} in R (a polynomial ring over a field k with 11 indeterminates) with dim $(R/\mathfrak{p}) = 3$ and depth $(R/\mathfrak{p}) = 2$ such that $K_{R/\mathfrak{p}} \cong R/\mathfrak{p}$. Put $\mathfrak{b} := \underline{x} : \mathfrak{p}$, where \underline{x} is a maximal regular sequence contained in \mathfrak{p} . Then \mathfrak{b} is linked to \mathfrak{p} . But

$$depth(R/\mathfrak{b}) = depth(K_{R/\mathfrak{p}}) - 1 = 1 \neq 2 = depth(R/\mathfrak{p}),$$

where the first equality follows by Proposition 3.3.

This example encouraged us to bring the following corollaries.

Corollary 3.5. Suppose that a is an unmixed, non Cohen-Macaulay and quasi-Gorenstein ideal in the Gorenstein ring (R, \mathfrak{m}) . Further assume b is an ideal linked to a. Then depth $(R/\mathfrak{b}) = depth(R/\mathfrak{a}) - 1$.

Corollary 3.6. Suppose that a is an unmixed, non Cohen-Macaulay and Buchsbaum ideal in the Gorenstein ring (R, \mathfrak{m}) . Suppose that $K_{R/\mathfrak{a}}$ is Cohen-Macaulay and dim $(R/\mathfrak{a}) > 2$. Further assume that b is an ideal linked to a. Then depth $(R/\mathfrak{b}) > 1 = depth(R/\mathfrak{a})$.

Proof. First, note that by [11, Theorem 5.4] we have $H^i_{\mathfrak{m}}(R/\mathfrak{a}) = 0$ for all $1 < i < \dim(R/\mathfrak{a})$. Then $\operatorname{depth}(R/\mathfrak{a}) = 1$. On the other hand, one has

$$\operatorname{depth}(R/\mathfrak{b}) = \operatorname{depth}(K_{R/\mathfrak{a}}) - 1 = \operatorname{dim}(K_{R/\mathfrak{a}}) - 1 > 1.$$

EGHBALI AND SHIRMOHAMMADI

This completes the proof.

As a consequence of Proposition 3.3 we derive the following result for the equality of depth under linkage. It shows that the equality of depth in Example 3.2 is not haphazard.

Corollary 3.7. Suppose that (R, \mathfrak{m}) is a Gorenstein local ring, and \mathfrak{a} and \mathfrak{b} are two linked ideals of R with $\dim(R/\mathfrak{a}) = 2$. Then $\operatorname{depth}(R/\mathfrak{b}) = \operatorname{depth}(R/\mathfrak{a}) = 1$ whenever R/\mathfrak{a} is not Cohen-Macaulay.

Proof. As $K_{R/a} = \text{Hom}(R/a, R/\underline{x})$, where \underline{x} is a maximal regular sequence in \mathfrak{a} , then by [3, Exercise 1.4.19] we can see that

$$2 \geq \operatorname{depth}(K_{R/\mathfrak{a}})$$
$$\geq \min\{2, \operatorname{depth}(R/\underline{x})\}$$
$$= \min\{2, \dim(R/\underline{x})\} = 2$$

So by Proposition 3.3 we have depth(R/\mathfrak{b}) = 1. With a similar argument we get depth(R/\mathfrak{a}) = 1.

Theorem 3.8. Suppose that \mathfrak{a} and \mathfrak{b} are ideals of a local ring (R, \mathfrak{m}) .

- (a) Further assume (R, \mathfrak{m}) is a regular local ring of positive characteristic and \mathfrak{a} is linked to \mathfrak{b} . If R/\mathfrak{a} is a Cohen-Macaulay ring, then $cd(\mathfrak{a}, R) = cd(\mathfrak{b}, R)$.
- (b) Assume that R = k[x₁,...,x_n]_(x₁,...,x_n) is a polynomial ring over a field k of positive characteristic localized at (x₁,...,x_n) with a is a squarefree monomial ideal. If a is evenly linked to b, then cd(b, R) ≤ cd(a, R).
- (c) Assume that $R = k[x_1, ..., x_n]_{(x_1, ..., x_n)}$ is a polynomial ring over a field k localized at $(x_1, ..., x_n)$. If a and b are two evenly linked squarefree monomial ideals, then one has cd(a, R) = cd(b, R).

Proof. (a) Our assumptions in conjunction with [8, 3.1] prove the assertion.

(b) As a is a square free monomial ideal, then [8, 4.2] implies that depth $R/\mathfrak{a} = \text{fgrade}(\mathfrak{a}, R)$. Since a is evenly linked to b then depth $R/\mathfrak{a} = \text{depth } R/\mathfrak{b}$ (cf. Corollary 3.4). Now, the claim follows by [8, 3.1] and the equality (1.2).

(c) The claim follows by Corollary 3.4 and [8, Theorem 4.2].

It is noteworthy to say that assumptions in Theorem 3.8(a) are not too much as we have seen in Example 3.1.

Remark 3.9. Suppose that a is an ideal in a regular local ring (R, \mathfrak{m}) of positive characteristic in a linkage class of a complete intersection ideal. Immediately, it follows from Theorem 3.8(a) that ht $\mathfrak{a} = \mathrm{cd}(\mathfrak{a}, R)$, that is a is a cohomologically complete intersection ideal.

Acknowledgment. The authors are deeply grateful to Professor K. Divaani-Aazar for careful reading of the first draft of the paper and useful comments.

References

 M. Asgharzadeh and K. Divaani-Aazar, Finiteness properties of formal local cohomology modules and Cohen-Macaulayness, Comm. Alg. 39, no. 3, 1082–1103 (2011).

- M. Brodmann and R. Y. Sharp, Local cohomology: an algebraic introduction with geometric applications, Cambridge Univ. Press, 60, Cambridge, (1998).
- [3] W. Bruns and J. Herzog, *Cohen–Macaulay rings*, Rev. ed. Cambridge Studies in Advanced Mathematics 39, Cambridge, Cambridge University Press (1998).
- [4] W. Bruns and R. Schwänzl, *The number of equations defining a determinantal variety*, Bull. London Math. Soc. 22, 439–445 (1990).
- [5] CoCoATeam, CoCoA: a system for doing Computations in Commutative Algebra, Available at http://cocoa.dima.unige.it.
- [6] K. Divaani-Aazar, R. Naghipour and M. Tousi, *Cohomological dimension of certain algebraic varieties*, Proc. Amer. Math. Soc. (130)12, 3537–3544 (2002).
- [7] M. Eghbali, On Artinianness of Formal Local Cohomology, Colocalization and Coassociated Primes, to appear in Math. Scand.
- [8] M. Eghbali, On set theoretically and cohomologically complete intersection ideals, Preprint, ArXiv:1211.5942[math.AC].
- [9] D. Eisenbud, Commutative Algebra (With a View Towards Algebraic Geometry), Springer-Verlag, (1995).
- [10] G. Faltings, Über lokale Kohomologiegruppen hoher Ordnung, J. Reine Angew. Math. 313, 43–51 (1980).
- [11] Sh. Goto and Y. Shimoda, On Rees algebras over Buchsbaum rings, J. Math. Kyoto Univ. (20)4, 691–708 (1980).
- [12] A. Grothendieck, Local cohomologyg, Notes by R. Hartshorne, Lect. Notes in Math., 20, Springer, (1966).
- [13] R. Hartshorne, Cohomological dimension of algebraic varieties, Annals of Math. 88, 403–450 (1968).
- [14] R. Hartshorne and R. Speiser, Local cohomological dimension in characteristic p, Annals of Math. 1052, 45–79 (1977).
- [15] M. Hermann and N. V. Trung, Examples of Buchsbaum quasi-Gorenstein rings, Proc. Amer. Math. Soc. V. 117, 619-625 (1993).
- [16] C. Huneke, G. Lyubeznik, On the vanishing of local cohomology modules, Inv. Math. 102 73–93 (1990).
- [17] A. Ogus, Local cohomological dimension, Annals of Math. 982, 327-365 (1973).
- [18] C. Peskine and L. Szpiro, Dimension projective finie et cohomologie locale, Publ. Math. I.H.E.S., 42, 323–395 (1973).
- [19] P. Schenzel, Notes on liaison and duality, J. Math. Kyoto Univ., 22(3), 485-498 (1982).
- [20] P. Schenzel, On formal local cohomology and connectedness, J. Algebra, 315(2), 894–923 (2007).
- [21] J. Stückrad and W. Vogel, Buchsbaum Rings and Applications, VEB Deutscher Verlag der Wissenschaften, Berlin, Germany, (1986).
- [22] K. I. Yoshida, Cofiniteness of local cohomology modules for ideals of dimension one, Nagoya Math. J. 147, 179-191 (1997).

School of Mathematics, Institute for Research in Fundamental Sciences (IPM), P. O. Box: 19395-5746, Tehran-Iran

E-mail address: m.eghbali@yahoo.com

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TABRIZ, TABRIZ, IRAN.

E-mail address: shirmohammadi@tabrizu.ac.ir