

# Means of algebraic numbers in the unit disk

Igor E. Pritsker<sup>1</sup>

*Department of Mathematics, Oklahoma State University, Stillwater, OK 74078, U.S.A.*

## Abstract

Schur studied limits of the arithmetic means  $s_n$  of zeros for polynomials of degree  $n$  with integer coefficients and simple zeros in the closed unit disk. If the leading coefficients are bounded, Schur proved that  $\limsup_{n \rightarrow \infty} |s_n| \leq 1 - \sqrt{e}/2$ . We show that  $s_n \rightarrow 0$ , and estimate the rate of convergence by generalizing the Erdős-Turán theorem on the distribution of zeros. *To cite this article: I. E. Pritsker, C. R. Acad. Sci. Paris, Ser. I 336 (2003).*

## Résumé

**Moyennes de nombres algébriques dans le disque unité.** Schur a étudié les limites des moyennes arithmétiques  $s_n$  des zéros pour les polynômes à coefficients entiers de degré  $n$  ayant des zéros simples dans le disque unité fermé. Lorsque les coefficients dominants restent bornés, Schur a démontré que  $\limsup_{n \rightarrow \infty} |s_n| \leq 1 - \sqrt{e}/2$ . Nous prouvons que  $s_n \rightarrow 0$ . Nous donnons une estimation du taux de convergence, grâce à une généralisation d'un théorème de Erdős-Turán sur la distribution des zéros. *Pour citer cet article : I. E. Pritsker, C. R. Acad. Sci. Paris, Ser. I 336 (2003).*

## 1. Schur's problem and equidistribution of zeros

Let  $\mathbb{Z}_n(D)$  be the set of polynomials of degree  $n$  with integer coefficients and all zeros in the closed unit disk  $D$ . We denote the subset of  $\mathbb{Z}_n(D)$  with simple zeros by  $\mathbb{Z}_n^1(D)$ . Given  $M > 0$ , we write  $P_n = a_n z^n + \dots \in \mathbb{Z}_n^1(D, M)$  if  $|a_n| \leq M$  and  $P_n \in \mathbb{Z}_n^1(D)$  (respectively  $P_n \in \mathbb{Z}_n(D, M)$  if  $|a_n| \leq M$  and  $P_n \in \mathbb{Z}_n(D)$ ). Schur [9, §8] studied the limiting behavior of the arithmetic means  $s_n$  of zeros for polynomials from  $\mathbb{Z}_n^1(D, M)$  as  $n \rightarrow \infty$ , where  $M > 0$  is an arbitrary fixed number. He showed that  $\limsup_{n \rightarrow \infty} |s_n| \leq 1 - \sqrt{e}/2$ , and remarked that this lim sup is equal to 0 for *monic* polynomials from  $\mathbb{Z}_n(D)$  by Kronecker's theorem [6]. We prove that  $\lim_{n \rightarrow \infty} s_n = 0$  for any sequence of polynomials from Schur's class  $\mathbb{Z}_n^1(D, M)$ ,  $n \in \mathbb{N}$ . This result is obtained as a consequence of the asymptotic equidistribution of zeros near the unit circle. Namely, if  $\{\alpha_k\}_{k=1}^n$  are the zeros of  $P_n$ , we define the counting measure  $\tau_n :=$

*Email address:* [igor@math.okstate.edu](mailto:igor@math.okstate.edu) (Igor E. Pritsker).

*URL:* <http://www.math.okstate.edu/~igor/> (Igor E. Pritsker).

1 . Research was partially supported by NSA, and by the Alexander von Humboldt Foundation.

$\frac{1}{n} \sum_{k=1}^n \delta_{\alpha_k}$ , where  $\delta_{\alpha_k}$  is the unit point mass at  $\alpha_k$ . Consider the normalized arclength measure  $\mu$  on the unit circumference  $\mathbb{T}$ , with  $d\mu(e^{it}) := \frac{1}{2\pi} dt$ . If the  $\tau_n$  converge weakly to  $\mu$  as  $n \rightarrow \infty$  ( $\tau_n \xrightarrow{*} \mu$ ) then  $\lim_{n \rightarrow \infty} s_n = \lim_{n \rightarrow \infty} \int z d\tau_n(z) = \int z d\mu(z) = 0$ . Thus Schur's problem is solved by the following result.

**Theorem 1.1** *If  $P_n(z) = a_n z^n + \dots \in \mathbb{Z}_n^1(D)$ ,  $n \in \mathbb{N}$ , satisfy  $\lim_{n \rightarrow \infty} |a_n|^{1/n} = 1$ , then  $\tau_n \xrightarrow{*} \mu$  as  $n \rightarrow \infty$ .*

Ideas on the equidistribution of zeros date back to Jentzsch and Szegő, cf. [1, Ch. 2]. They were developed further by Erdős and Turán [4], and many others; see [1] for history and additional references. More recently, this topic received renewed attention in number theory, e.g. in the work of Bilu [2]. If the leading coefficients of polynomials are bounded, then we can allow even certain multiple zeros. Define the multiplicity of an irreducible factor  $Q$  of  $P_n$  as the integer  $m_n \geq 0$  such that  $Q^{m_n}$  divides  $P_n$ , but  $Q^{m_n+1}$  does not divide  $P_n$ . If a factor  $Q$  occurs infinitely often in a sequence  $P_n$ ,  $n \in \mathbb{N}$ , then  $m_n = o(n)$  means  $\lim_{n \rightarrow \infty} m_n/n = 0$ . If  $Q$  is present only in finitely many  $P_n$ , then  $m_n = o(n)$  by definition.

**Theorem 1.2** *Assume that  $P_n \in \mathbb{Z}_n(D, M)$ ,  $n \in \mathbb{N}$ . If every irreducible factor in the sequence of polynomials  $P_n$  has multiplicity  $o(n)$ , then  $\tau_n \xrightarrow{*} \mu$  as  $n \rightarrow \infty$ .*

**Corollary 1.3** *If  $P_n(z) = a_n \prod_{k=1}^n (z - \alpha_k)$ ,  $n \in \mathbb{N}$ , satisfy the assumptions of Theorem 1.1 or 1.2, then*

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n \alpha_k^m = 0, \quad m \in \mathbb{N}.$$

We also show that the norms  $\|P_n\|_\infty := \max_{|z|=1} |P_n(z)|$  have at most subexponential growth.

**Corollary 1.4** *If  $P_n$ ,  $n \in \mathbb{N}$ , satisfy the assumptions of Theorem 1.1 or Theorem 1.2, then*

$$\lim_{n \rightarrow \infty} \|P_n\|_\infty^{1/n} = 1.$$

This result is somewhat unexpected, as we have no direct control of the norm or coefficients (except for the leading one). For example,  $P_n(z) = (z - 1)^n$  has norm  $\|P_n\|_\infty = 2^n$ .

We now consider quantitative aspects of the convergence  $\tau_n \xrightarrow{*} \mu$ . As an application, we obtain estimates of the convergence rate of  $s_n$  to 0 in Schur's problem. A classical result on the distribution of zeros is due to Erdős and Turán [4]. For  $P_n(z) = \sum_{k=0}^n a_k z^k$  with  $a_k \in \mathbb{C}$ , let  $N(\phi_1, \phi_2)$  be the number of zeros in the sector  $\{z \in \mathbb{C} : 0 \leq \phi_1 \leq \arg(z) \leq \phi_2 < 2\pi\}$ , where  $\phi_1 < \phi_2$ . Erdős and Turán [4] proved that

$$\left| \frac{N(\phi_1, \phi_2)}{n} - \frac{\phi_2 - \phi_1}{2\pi} \right| \leq 16 \sqrt{\frac{1}{n} \log \frac{\|P_n\|_\infty}{\sqrt{|a_0 a_n|}}}. \quad (1)$$

The constant 16 was improved by Ganelius, and  $\|P_n\|_\infty$  was replaced by weaker integral norms by Amoroso and Mignotte; see [1] for more history and references. Our main difficulty in applying (1) to Schur's problem is the absence of an effective estimate for  $\|P_n\|_\infty$ ,  $P_n \in \mathbb{Z}_n^1(D, M)$ . We prove a new "discrepancy" estimate via energy considerations from potential theory. These ideas originated in part in the work of Kleiner, and were developed by Sjögren and Hüsing, see [1, Ch. 5]. We also use the Mahler measure of a polynomial  $P_n(z) = a_n \prod_{k=1}^n (z - \alpha_k)$ , defined by  $M(P_n) := \exp\left(\frac{1}{2\pi} \int_0^{2\pi} \log |P_n(e^{it})| dt\right)$ . Note that

$M(P_n) = \lim_{p \rightarrow 0} \|P_n\|_p$ , where  $\|P_n\|_p := \left(\frac{1}{2\pi} \int_0^{2\pi} |P_n(e^{it})|^p dt\right)^{1/p}$ ,  $p > 0$ . Jensen's formula readily gives  $M(P_n) = |a_n| \prod_{k=1}^n \max(1, |\alpha_k|)$  [3, p. 3]. Hence  $M(P_n) = |a_n| \leq M$  for any  $P_n \in \mathbb{Z}_n(D, M)$ .

**Theorem 1.5** *Let  $\phi : \mathbb{C} \rightarrow \mathbb{R}$  satisfy  $|\phi(z) - \phi(t)| \leq A|z - t|$ ,  $z, t \in \mathbb{C}$ , and  $\text{supp}(\phi) \subset \{z : |z| \leq R\}$ . If  $P_n(z) = a_n \prod_{k=1}^n (z - \alpha_k)$  is a polynomial with integer coefficients and simple zeros, then*

$$\left| \frac{1}{n} \sum_{k=1}^n \phi(\alpha_k) - \int \phi d\mu \right| \leq A(2R + 1) \sqrt{\frac{\log \max(n, M(P_n))}{n}}, \quad n \geq 55. \quad (2)$$

This theorem is related to recent results of Favre and Rivera-Letelier [5], obtained in a different setting. Choosing  $\phi$  appropriately, we obtain an estimate of the means  $s_n$  in Schur's problem.

**Corollary 1.6** *If  $P_n \in \mathbb{Z}_n^1(D, M)$  then*

$$\left| \frac{1}{n} \sum_{k=1}^n \alpha_k \right| \leq 8 \sqrt{\frac{\log n}{n}}, \quad n \geq \max(M, 55).$$

We also have an improvement of Corollary 1.4 for Schur's class  $\mathbb{Z}_n^1(D, M)$ .

**Corollary 1.7** *If  $\{P_n\}_{n=1}^\infty \in \mathbb{Z}_n^1(D, M)$  then there is some  $c > 0$  such that  $\|P_n\|_\infty \leq e^{c\sqrt{n} \log n}$  as  $n \rightarrow \infty$ .*

The proof of Theorem 1.5 gives a result for arbitrary polynomials with simple zeros, and for any continuous  $\phi$  with finite Dirichlet integral  $D[\phi] = \iint (\phi_x^2 + \phi_y^2) dA$ . Moreover, all arguments may be extended to general sets of logarithmic capacity 1, e.g. to  $[-2, 2]$ . Using the characteristic function  $\phi = \chi_E$ , we can prove general discrepancy estimates on arbitrary sets, and obtain an Erdős-Turán-type theorem. Our results have a number of applications to the problems on integer polynomials considered in [3].

## 2. Proofs

*Proof of Theorem 1.1.* Observe that the discriminant  $\Delta(P_n) := a_n^{2n-2} \prod_{1 \leq j < k \leq n} (\alpha_j - \alpha_k)^2$  is an integer, as a symmetric form in the zeros of  $P_n$ . Since  $P_n$  has simple roots, we have  $\Delta(P_n) \neq 0$  and  $|\Delta(P_n)| \geq 1$ . Using weak compactness, we assume that  $\tau_n \xrightarrow{*} \tau$ , where  $\tau$  is a probability measure on  $D$ . Let  $K_M(x, t) := \min(-\log|x-t|, M)$ . Since  $\tau_n \times \tau_n \xrightarrow{*} \tau \times \tau$ , we obtain for the energy of  $\tau$  that

$$\begin{aligned} I[\tau] &:= - \iint \log|x-t| d\tau(x) d\tau(t) = \lim_{M \rightarrow \infty} \left( \lim_{n \rightarrow \infty} \iint K_M(x, t) d\tau_n(x) d\tau_n(t) \right) \\ &= \lim_{M \rightarrow \infty} \left( \lim_{n \rightarrow \infty} \left( \frac{1}{n^2} \sum_{j \neq k} K_M(\alpha_j, \alpha_k) + \frac{M}{n} \right) \right) \leq \lim_{M \rightarrow \infty} \left( \liminf_{n \rightarrow \infty} \frac{1}{n^2} \sum_{j \neq k} \log \frac{1}{|\alpha_j - \alpha_k|} \right) \\ &= \liminf_{n \rightarrow \infty} \frac{1}{n^2} \log \frac{|a_n|^{2n-2}}{\Delta(P_n)} \leq \liminf_{n \rightarrow \infty} \frac{1}{n^2} \log |a_n|^{2n-2} = 0. \end{aligned}$$

Thus  $I[\tau] \leq 0$ . But  $I[\nu] > 0$  for any probability measure  $\nu$  on  $D$ , except for  $\mu$  [7]. Hence  $\tau = \mu$ .

*Proof of Theorem 1.2.* Let  $\phi \in C(\mathbb{C})$ . Note that for any  $\epsilon > 0$  there are finitely many irreducible factors  $Q$  in the sequence  $P_n$  such that  $|\int \phi d\tau(Q) - \int \phi d\mu| \geq \epsilon$ , where  $\tau(Q)$  is the zero counting measure for  $Q$ . Indeed, if we have an infinite sequence of such  $Q_m$ , then  $\deg(Q_m) \rightarrow \infty$ , as there are only finitely many  $Q_m \in \mathbb{Z}_n(D, M)$  of bounded degree. Hence  $\int \phi d\tau(Q_m) \rightarrow \int \phi d\mu$  by Theorem 1.1. Let the number of such exceptional factors  $Q_m$  be  $N$ . Then we have  $|n \int \phi d\tau_n - n \int \phi d\mu| \leq N o(n) \max_D |\phi - \int \phi d\mu| + (n - N)\epsilon$ ,  $n \in \mathbb{N}$ . Hence  $\limsup_{n \rightarrow \infty} |\int \phi d\tau_n - \int \phi d\mu| \leq \epsilon$ , and  $\lim_{n \rightarrow \infty} \int \phi d\tau_n = \int \phi d\mu$  after letting  $\epsilon \rightarrow 0$ .

*Proof of Corollary 1.3.* Let  $\phi(z) = z^m$  and write  $\lim_{n \rightarrow \infty} \int z^m d\tau_n(z) = \int z^m d\mu(z) = 0$ .

*Proof of Corollary 1.4.* Let  $\|P_n\|_\infty = |P_n(z_n)|$ ,  $z_n \in D$ , and assume  $\lim_{n \rightarrow \infty} z_n = z_0 \in D$  by compactness. Then  $\|P_n\|_\infty = \exp(\log |P_n(z_n)|) = |a_n| \exp(n \int \log|z_n - t| d\tau_n(t))$ . Since  $\tau_n \xrightarrow{*} \mu$ , Theorem I.6.8 of [8] gives  $\limsup_{n \rightarrow \infty} \|P_n\|_\infty^{1/n} \leq \exp(\int \log|z_0 - t| d\mu(t)) = 1$  [8, p. 22]. But  $\|P_n\|_\infty \geq |a_n| \geq 1$ , see [1, p. 16].

*Proof of Theorem 1.5.* Given  $r > 0$ , define the measures  $\nu_k^r$  with  $d\nu_k^r(\alpha_k + re^{it}) = dt/(2\pi)$ ,  $t \in [0, 2\pi]$ . Let  $\tau_n^r := \frac{1}{n} \sum_{k=1}^n \nu_k^r$ , and estimate  $|\int \phi d\tau_n - \int \phi d\tau_n^r| \leq \frac{1}{n} \sum_{k=1}^n \frac{1}{2\pi} \int_0^{2\pi} |\phi(\alpha_k) - \phi(\alpha_k + re^{it})| dt \leq \omega_\phi(r)$ , where  $\omega_\phi(r) := \sup_{|z-\zeta| \leq r} |\phi(z) - \phi(\zeta)|$  is the modulus of continuity of  $\phi$ .

Let  $p_\nu(z) := -\int \log|z-t| d\nu(t)$  be the potential of a measure  $\nu$ . A direct evaluation gives that  $p_{\nu_k^r}(z) = -\log \max(r, |z - \alpha_k|)$  and  $p_\mu(z) = -\log \max(1, |z|)$  [8, p. 22]. Consider  $\sigma := \tau_n^r - \mu$ ,  $\sigma(\mathbb{C}) = 0$ . One computes (or see [8, p. 92]) that  $d\sigma = -\frac{1}{2\pi} (\partial p_\sigma / \partial n_+ + \partial p_\sigma / \partial n_-) ds$ , where  $ds$  is the arclength on  $\text{supp}(\sigma) = \{|z| = 1\} \cup (\cup_{k=1}^n \{|z - \alpha_k| = r\})$ , and  $n_\pm$  are the inner and the outer normals. We now use

Green's identity  $\iint_G u \Delta v \, dA = \int_{\partial G} u \frac{\partial v}{\partial n} \, ds - \iint_G \nabla u \cdot \nabla v \, dA$  with  $u = \phi$  and  $v = p_\sigma$  in each component  $G$  of  $\{|z| < R\} \setminus \text{supp}(\sigma)$ . Since  $\Delta p_\sigma = 0$  in  $G$ , adding the identities for all  $G$ , we obtain that

$$\left| \int \phi \, d\sigma \right| = \frac{1}{2\pi} \left| \iint_{|z| \leq R} \nabla \phi \cdot \nabla p_\sigma \, dA \right| \leq \frac{1}{2\pi} \sqrt{D[\phi]} \sqrt{D[p_\sigma]},$$

where  $D[\phi] = \iint (\phi_x^2 + \phi_y^2) \, dA$  is the Dirichlet integral of  $\phi$ . It is known that  $D[p_\sigma] = 2\pi I[\sigma]$  [7, Thm 1.20], where  $I[\sigma] = -\iint \log |z - t| \, d\sigma(z) \, d\sigma(t) = \int p_\sigma \, d\sigma$ . Since  $p_\mu(z) = -\log \max(1, |z|)$ , we observe that  $\int p_\mu \, d\mu = 0$ , so that  $I[\sigma] = \int p_{\tau_n^r} \, d\tau_n^r - 2 \int p_\mu \, d\tau_n^r$ . Further,  $-\int p_\mu \, d\tau_n^r = \int \log \max(1, |z|) \, d\tau_n^r(z) \leq \left( \sum_{|\alpha_k| \leq 1+r} \log(1+2r) + \sum_{|\alpha_k| > 1+r} \log |\alpha_k| \right) / n \leq \log(1+2r) + \frac{1}{n} \log M(P_n) - \frac{1}{n} \log |a_n|$ . We also have that  $\int p_{\tau_n^r} \, d\tau_n^r \leq \left( -\sum_{j \neq k} \log |\alpha_j - \alpha_k| - n \log r \right) / n^2$ . We next combine the energy estimates to obtain

$$I[\sigma] \leq \frac{2}{n} \log M(P_n) - \frac{1}{n^2} \log |a_n^2 \Delta(P_n)| - \frac{1}{n} \log r + 4r.$$

Collecting all estimates, we proceed with  $\left| \int \phi \, d\tau_n - \int \phi \, d\mu \right| \leq \left| \int \phi \, d\tau_n - \int \phi \, d\tau_n^r \right| + \left| \int \phi \, d\tau_n^r - \int \phi \, d\mu \right| \leq \omega_\phi(r) + \sqrt{D[\phi]} \sqrt{D[p_\sigma]} / (2\pi) = \omega_\phi(r) + \sqrt{D[\phi]} \sqrt{I[\sigma]} / (2\pi)$ . Thus we arrive at the main inequality:

$$\left| \int \phi \, d\tau_n - \int \phi \, d\mu \right| \leq \omega_\phi(r) + \sqrt{\frac{D[\phi]}{2\pi}} \left( \frac{2}{n} \log M(P_n) - \frac{1}{n^2} \log |a_n^2 \Delta(P_n)| - \frac{1}{n} \log r + 4r \right)^{1/2}. \quad (3)$$

Note that  $D[\phi] \leq 2\pi R^2 A^2$ , as  $|\phi_x| \leq A$  and  $|\phi_y| \leq A$  a.e. in  $\mathbb{C}$ . Also,  $\omega_\phi(r) \leq Ar$ . Since  $|\Delta(P_n)| \geq 1$  and  $|a_n| \geq 1$ , we have  $|a_n^2 \Delta(P_n)| \geq 1$ . Hence (2) follows from (3) by letting  $r = 1/\max(n, M(P_n))$ .

*Proof of Corollary 1.6.* Since  $P_n$  has real coefficients, we have that  $s_n = \int z \, d\tau_n(z) = \int \Re(z) \, d\tau_n(z)$ . We let  $\phi(z) = \Re(z)$ ,  $|z| \leq 1$ ;  $\phi(z) = \Re(z)(1 - \log |z|)$ ,  $1 \leq |z| \leq e$ ; and  $\phi(z) = 0$ ,  $|z| \geq e$ . An elementary computation shows that  $|\phi_x(z)| \leq 1$  and  $|\phi_y(z)| \leq 1/2$  for all  $z = x + iy \in \mathbb{C}$ . The Mean Value Theorem gives  $|\phi(z) - \phi(t)| \leq |z - t| \max_{\mathbb{C}} \sqrt{\phi_x^2 + \phi_y^2}$ . Hence we can use Theorem 1.5 with  $A = \sqrt{5}/2$  and  $R = e$ .

*Proof of Corollary 1.7.* Note that  $\log |P_n(z)| = n \int \log |z - w| \, d\tau_n(w)$ . For  $|z| = 1 + 1/n$ , we let  $\phi(w) = \log |z - w|$ ,  $|w| \leq 1$ ;  $\phi(w) = (1 - \log |w|) \log |1 - \bar{z}w|$ ,  $1 \leq |w| \leq e$ ; and  $\phi(w) = 0$ ,  $|w| \geq e$ . Then  $|\phi_x(w)| = O(|z - w|^{-1})$ ,  $|w| \leq 1$ ;  $|\phi_x(w)| = O(|1 - \bar{z}w|^{-1})$ ,  $1 \leq |w| \leq e$ ; and the same estimates hold for  $|\phi_y|$ . Hence  $D[\phi] = O\left(\iint_{|w| \leq 1} |z - w|^{-2} \, dA(w)\right) = O\left(\int_{1/n}^1 r^{-1} \, dr\right) = O(\log n)$ , and  $\omega_\phi(r) \leq r \max_{\mathbb{C}} \sqrt{\phi_x^2 + \phi_y^2} = rO(n)$ . Let  $r = 1/n^2$  and use (3) to obtain  $|\log |P_n(z)|| - n \log |z| = O(\sqrt{n} \log n)$ .

## References

- [1] V. V. Andrievskii and H.-P. Blatt, *Discrepancy of Signed Measures and Polynomial Approximation*, Springer-Verlag, New York, 2002.
- [2] Y. Bilu, Limit distribution of small points on algebraic tori, *Duke Math. J.* 89 (1997), 465-476.
- [3] P. Borwein, *Computational Excursions in Analysis and Number Theory*, Springer-Verlag, New York, 2002.
- [4] P. Erdős and P. Turán, On the distribution of roots of polynomials, *Ann. Math.* 51 (1950), 105-119.
- [5] C. Favre and J. Rivera-Letelier, Equidistribution quantitative des points de petite hauteur sur la droite projective, *Math. Ann.* 335 (2006), 311-361; Corrigendum in *Math. Ann.* 339 (2007), 799-801.
- [6] L. Kronecker, Zwei Sätze über Gleichungen mit ganzzahligen Coëfficienten, *J. reine ang. Math.* 53 (1857), 173-175.
- [7] N. S. Landkof, *Foundations of Modern Potential Theory*, Springer-Verlag, New York, 1972.
- [8] E. B. Saff and V. Totik, *Logarithmic Potentials with External Fields*, Springer-Verlag, Berlin, 1997.
- [9] I. Schur, Über die Verteilung der Wurzeln bei gewissen algebraischen Gleichungen mit ganzzahligen Koeffizienten, *Math. Zeit.* 1 (1918), 377-402.