

# On the Diagonal Susceptibility of the 2D Ising Model

**Craig A. Tracy**

*Department of Mathematics  
University of California  
Davis, CA 95616, USA*

**Harold Widom**

*Department of Mathematics  
University of California  
Santa Cruz, CA 95064, USA*

## Abstract

We consider the diagonal susceptibility of the isotropic 2D Ising model for temperatures below the critical temperature. For a parameter  $k$  related to temperature and the interaction constant, we extend the diagonal susceptibility to complex  $k$  inside the unit disc, and prove the conjecture that the unit circle is a natural boundary.

## I. Introduction

For the 2D Ising model [15, 16, 23], after the zero-field free energy [20] and the spontaneous magnetization [21, 28], the most important zero-field thermodynamic quantity is the magnetic susceptibility  $\chi$ . Since the free energy is known only in zero magnetic field, the susceptibility is usually studied through its relation with the zero-field spin-spin correlation function:

$$\beta^{-1}\chi = \sum_{M,N \in \mathbb{Z}} \{ \langle \sigma_{0,0} \sigma_{M,N} \rangle - \mathcal{M}^2 \} \quad (1)$$

where  $\beta = (k_B T)^{-1}$ ,  $T$  is temperature,  $k_B$  is Boltzmann's constant and  $\mathcal{M}$  is the spontaneous magnetization. If  $T_c$  denotes the critical temperature, we recall that for the isotropic 2D Ising model, i.e. horizontal and vertical interaction constants have the same value  $J$ , the spontaneous magnetization is given for  $T < T_c$  by

$$\mathcal{M} = (1 - k^2)^{1/8} \quad (2)$$

where  $k := (\sinh 2\beta J)^{-2}$  and  $\mathcal{M}$  is zero for  $T > T_c$ . (For  $0 < T < T_c$  we have  $0 < k < 1$ .)

The analysis of  $\chi = \chi(T)$  in the neighborhood of the critical temperature  $T_c$  has a long history. We refer the reader to McCoy *et al.* [17] for a review of these developments. The analysis of  $\chi$  for *complex temperatures* was initiated by Guttman and Enting [13] and by Nickel [18, 19]. (For further developments see [10, 22].) Nickel's analysis takes as its beginning the (commonly called) form-factor or particle expansion of the spin-spin correlation function [27]. For  $T < T_c$  this expansion is an infinite sum whose  $n$ th summand is a  $2n$ -dimensional integral. From an asymptotic analysis of these integrals, Nickel was led to conjecture that  $|k| = 1$  is a *natural boundary* for  $\chi$ . As Nickel himself noted, the analysis is nonrigorous since one must show that there are no cancellations of singularities in the sum. This has turned out to be a difficult problem to resolve rigorously.

In Boukraa *et al.* [7], these authors, building on results of [14], introduce a simplified model for  $\chi$ , called the *diagonal susceptibility*  $\chi_d$ , which is defined by having “a magnetic field which acts only on one diagonal of the lattice.” (See [2] for further developments.) Thus the analogue of (1) is

$$\beta^{-1}\chi_d = \sum_{N \in \mathbb{Z}} \{ \langle \sigma_{0,0} \sigma_{N,N} \rangle - \mathcal{M}^2 \}. \quad (3)$$

In this paper we consider  $\chi_d$  only for  $T < T_c$ , in which case  $k < 1$ . Then we extend  $\chi_d$  to  $k$  complex with  $|k| < 1$ . Using the Toeplitz determinant representation of the diagonal correlations [15, 24], we first derive the known representation of  $\chi_d$  in terms of a sum of multiple integrals  $\mathcal{S}_n$ . The derivation is different from those in [8, 9], [14] and [26]. As in [26] we use the identity of Geronimo-Case [12] and Borodin-Okounkov [5] relating a Toeplitz determinant to the Fredholm determinant of a product of Hankel operators. (For simplified proofs of the GCBO formula, see [3, 6].) But here we go from there to the multiple integral representation directly using a general identity for the integral of a product of determinants [1] (see eqn. (1.3) in [25]). For further background on the relationship between Toeplitz determinants and Ising correlations, see [4, 11].

In Section 4 we show that for each root of unity  $\epsilon \neq \pm 1$  a certain derivative of a certain  $\mathcal{S}_n$  is unbounded as  $k \rightarrow \epsilon$  radially, while the same derivative of the sum of the other  $\mathcal{S}_n$  remains bounded (Lemma 4). Thus, the unit circle  $|k| = 1$  is a natural boundary for  $\chi_d$ . This proves the conjecture by Boukraa *et al.* [7]. We note that in [7] the authors present an argument that the singularity of  $\mathcal{S}_n$  at an  $n$ th root of unity  $\epsilon$  is of the form  $(k - \epsilon)^{2n^2-1} \log(k - \epsilon)$ . Lemma 2 in Section IV formalizes this statement and fills in details of the proof.

## II. Toeplitz determinant representation

It was shown in [15, 24] that for  $N > 1$  the diagonal correlation has a representation as an  $N \times N$  Toeplitz determinant:

$$\langle \sigma_{0,0} \sigma_{N,N} \rangle = \det (\varphi_{m-n})_{1 \leq m, n \leq N}.$$

Here

$$\varphi(\xi) = \left[ \frac{1 - k\xi^{-1}}{1 - k\xi} \right]^{1/2},$$

and

$$\varphi_m = \frac{1}{2\pi i} \int \varphi(\xi) \xi^{-m-1} d\xi, \quad (4)$$

with integration over the unit circle. (We have  $\langle \sigma_{0,0}^2 \rangle = 1$ .)

As in [26] we invoke the formula of Geronimo-Case [12] and Borodin-Okounkov [5] to write the Toeplitz determinant in terms of the Fredholm determinant of a product of Hankel operators. We have  $\varphi(\xi) = \varphi_+(\xi) \varphi_-(\xi)$ , where

$$\varphi_+(\xi) = (1 - k\xi)^{-1/2} \quad \text{and} \quad \varphi_-(\xi) = (1 - k\xi^{-1})^{1/2}.$$

Since  $|k| < 1$  these extend analytically inside and outside the unit circle, respectively. The square roots are determined by  $\varphi_+(0) = \varphi_-(\infty) = 1$ .

The Hankel operator  $H(\psi)$  is the operator on  $\ell^2(\mathbb{Z}^+)$  with kernel  $(\psi_{i+j+1})_{i,j \geq 0}$ , where  $\psi_m$  given in analogy with (4). The operator  $H_N(\psi)$  has kernel  $(\psi_{N+i+j+1})$ .

Using  $\varphi_{\pm}(\xi) = 1/\varphi_{\mp}(\xi^{-1})$ , we find that the formula of GCBO gives

$$\det (\varphi_{m-n})_{1 \leq m, n \leq N} = \mathcal{M}^2 \det \left( I - H_N \left( \frac{\varphi_-}{\varphi_+} \right) H_N \left( \frac{\varphi_+}{\varphi_-} \right) \right).$$

Thus, if we define

$$\Lambda(\xi) = \frac{\varphi_-(\xi)}{\varphi_+(\xi)} = \sqrt{(1 - k\xi)(1 - k/\xi)}, \quad K_N = H_N(\Lambda) H_N(\Lambda^{-1}), \quad (5)$$

then

$$\beta^{-1} \chi_d = 1 - \mathcal{M}^2 + 2\mathcal{M}^2 \sum_{N=1}^{\infty} [\det(I - K_N) - 1] = 1 + \mathcal{M}^2(2\mathcal{S} - 1),$$

where

$$\mathcal{S} = \sum_{N=1}^{\infty} [\det(I - K_N) - 1]. \quad (6)$$

In what follows we extend  $\Lambda$  to be holomorphic in the complex plane cut along  $[0, k] \cup [k^{-1}, \infty]$ .

### III. Formula for $\mathcal{S}$

We use a slightly different notation for Hankel operators here.

**Proposition.** Let  $H_N(du)$  and  $H_N(dv)$  be two Hankel matrices acting on  $\ell^2(\mathbb{Z}^+)$  with  $i, j$  entries

$$\int x^{N+i+j} du(x), \quad \int y^{N+i+j} dv(y), \quad (7)$$

respectively, where  $u$  and  $v$  are measures supported inside the unit circle. Set  $K_N = H_N(du)H_N(dv)$ . Then

$$\begin{aligned} & \sum_{N=1}^{\infty} [\det(I - K_N) - 1] \\ &= \sum_{n=1}^{\infty} \frac{(-1)^n}{(n!)^2} \int \cdots \int \frac{\prod_i x_i y_i}{1 - \prod_i x_i y_i} \left( \det \left( \frac{1}{1 - x_i y_j} \right) \right)^2 \prod_i du(x_i) dv(y_i), \end{aligned}$$

where indices in the integrand run from 1 to  $n$ .

**Proof.** The Fredholm expansion is

$$\det(I - K_N) = 1 + \sum_{n=1}^{\infty} \frac{(-1)^n}{n!} \sum_{p_1, \dots, p_n \geq 0} \det(K_N(p_i, p_j)).$$

Therefore it suffices to show that

$$\begin{aligned} & \sum_{N=1}^{\infty} \sum_{p_1, \dots, p_n \geq 0} \det(K_N(p_i, p_j)) \\ &= \frac{1}{n!} \int \cdots \int \frac{\prod_i x_i y_i}{1 - \prod_i x_i y_i} \left( \det \left( \frac{1}{1 - x_i y_j} \right) \right)^2 du(x_1) \cdots du(x_n) dv(y_1) \cdots dv(y_n). \end{aligned}$$

We have

$$K_N(p_i, p_j) = \int \int \frac{x^{N+p_i} y^{N+p_j}}{1 - xy} du(x) dv(y).$$

It follows by a general identity [1] (eqn. (1.3) in [25]) that

$$\begin{aligned} \det(K_N(p_i, p_j)) &= \frac{1}{n!} \int \cdots \int \det(x_i^{N+p_j}) \det(y_i^{N+p_j}) \prod_i \frac{1}{1 - x_i y_i} \prod_i du(x_i) dv(y_i) \\ &= \frac{1}{n!} \int \cdots \int \left( \prod_i x_i y_i \right)^N \det(x_i^{p_j}) \det(y_i^{p_j}) \prod_i \frac{1}{1 - x_i y_i} \prod_i du(x_i) dv(y_i). \end{aligned}$$

Summing over  $N$  gives

$$\sum_{N=1}^{\infty} \det(K_N(p_i, p_j)) = \frac{1}{n!} \int \cdots \int \frac{\prod_i x_i y_i}{1 - \prod_i x_i y_i} \det(x_i^{p_j}) \det(y_i^{p_j}) \prod_i \frac{1}{1 - x_i y_i} \prod_i du(x_i) dv(y_i).$$

(Interchanging the sum with the integral is justified since the supports of  $u$  and  $v$  are inside the unit circle.)

Now we sum over  $p_1, \dots, p_n \geq 0$ . Using the general identity again (but in the other direction) gives

$$\sum_{p_1, \dots, p_n \geq 0} \det(x_i^{p_j}) \det(y_i^{p_j}) = n! \det \left( \sum_{p \geq 0} x_i^p y_j^p \right) = n! \det \left( \frac{1}{1 - x_i y_j} \right).$$

We almost obtained the desired result. It remain to show that

$$\det \left( \frac{1}{1 - x_i y_j} \right) \prod_i \frac{1}{1 - x_i y_i}, \quad (8)$$

which we obtain in the integrand, may be replaced by

$$\frac{1}{n!} \left( \det \left( \frac{1}{1 - x_i y_j} \right) \right)^2. \quad (9)$$

This follows by symmetrization over the  $x_i$ . (The rest of the integrand is symmetric.) For a permutation  $\pi$ , replacing the  $x_i$  by  $x_{\pi(i)}$  multiplies the determinant in (8) by  $\text{sgn } \pi$ , so to symmetrize we replace the other factor by

$$\frac{1}{n!} \sum_{\pi} \text{sgn } \pi \frac{1}{1 - x_{\pi(i)} y_i} = \frac{1}{n!} \det \left( \frac{1}{1 - x_i y_j} \right).$$

Thus, symmetrizing (8) gives (9).  $\square$

We apply this to the operator  $K_N = H_N(\Lambda) H_N(\Lambda^{-1})$  given by (5). The matrix for  $H_N(\Lambda)$  has  $i, j$  entry

$$\frac{1}{2\pi i} \int \Lambda(\xi) \xi^{-N-i-j-2} d\xi,$$

where the integration may be taken over a circle with radius in  $(1, |k|^{-1})$ . Setting  $\xi = 1/x$  and using  $\Lambda(1/x) = \Lambda(x)$  we see that the entries of  $H_N(\Lambda)$  are given as in (7) with

$$du(x) = \frac{1}{2\pi i} \Lambda(x) dx,$$

and integration is over a circle  $\mathcal{C}$  with radius in  $(|k|, 1)$ . Similarly  $H_N(\Lambda^{-1}) = H_N(v)$  where in (7)

$$d\psi(y) = \frac{1}{2\pi i} \Lambda(y)^{-1} dy,$$

with integration over the same circle  $\mathcal{C}$ .

Hence the Proposition gives

$$\mathcal{S} = \sum_{n=1}^{\infty} \mathcal{S}_n, \quad (10)$$

where

$$\mathcal{S}_n = \frac{(-1)^n}{(n!)^2} \frac{1}{(2\pi i)^{2n}} \int \cdots \int \frac{\prod_i x_i y_i}{1 - \prod_i x_i y_i} \left( \det \left( \frac{1}{1 - x_i y_j} \right) \right)^2 \prod_i \frac{\Lambda(x_i)}{\Lambda(y_i)} \prod_i dx_i dy_i,$$

with all integrations over  $\mathcal{C}$ .

We deform  $\mathcal{C}$  to the contour back and forth along the interval  $[0, k]$ , and then make the substitutions  $x_i \rightarrow kx_i$ ,  $y_i \rightarrow ky_i$ . We obtain

$$\mathcal{S}_n = \frac{1}{(n!)^2} \frac{\kappa^{2n}}{\pi^{2n}} \int_0^1 \cdots \int_0^1 \frac{\prod_i x_i y_i}{1 - \kappa^n \prod_i x_i y_i} \left( \det \left( \frac{1}{1 - \kappa x_i y_j} \right) \right)^2 \prod_i \frac{\Lambda_1(x_i)}{\Lambda_1(y_i)} \prod_i dx_i dy_i, \quad (11)$$

where we have set

$$\kappa = k^2, \quad \Lambda_1(x) = \sqrt{\frac{(1-x)(1-\kappa x)}{x}}.$$

Using the fact that the determinant in the integrand is a Cauchy determinant we obtain the alternative expression

$$\mathcal{S}_n = \frac{1}{(n!)^2} \frac{\kappa^{n(n+1)}}{\pi^{2n}} \int_0^1 \cdots \int_0^1 \frac{\prod_i x_i y_i}{1 - \kappa^n \prod_i x_i y_i} \frac{\Delta(x)^2 \Delta(y)^2}{\prod_{i,j} (1 - \kappa x_i y_j)^2} \prod_i \frac{\Lambda_1(x_i)}{\Lambda_1(y_i)} \prod_i dx_i dy_i, \quad (12)$$

where  $\Delta(x)$  and  $\Delta(y)$  are Vandermonde determinants. Clearly,  $\mathcal{S}_n$  is holomorphic in  $\kappa$  for  $|\kappa| < 1$ . It is straightforward to prove that the sum (10) converges uniformly in  $\kappa$  for  $|\kappa| \leq r$  for all  $0 < r < 1$ ; and hence,  $\mathcal{S}$  is holomorphic in the  $\kappa$  unit disc.

#### IV. Natural boundary

**Theorem.** The unit circle  $|\kappa| = 1$  is a natural boundary for  $\mathcal{S}$ .

There will four lemmas. In these,  $\epsilon \neq 1$  will be an  $n$ th root of unity and we consider the behavior of  $\mathcal{S}$  as  $\kappa \rightarrow \epsilon$  radially.

For  $\ell \geq 0$  we use the representation (12) and look at

$$\int_0^1 \cdots \int_0^1 \frac{\prod_i x_i y_i}{(1 - \kappa^n \prod_i x_i y_i)^{\ell+1}} \frac{\Delta(x)^2 \Delta(y)^2}{\prod_{i,j} (1 - \kappa x_i y_j)^2} \prod_i \frac{\Lambda_1(x_i)}{\Lambda_1(y_i)} \prod_i dx_i dy_i, \quad (13)$$

where all indices run from 1 to  $n$ . This will be the main contribution to  $d^\ell \mathcal{S}_n / d\kappa^\ell$ .

**Lemma 1.** The integral (13) is bounded when  $\ell < 2n^2 - 1$  and it is of the order  $\log(1 - |\kappa|)^{-1}$  when  $\ell = 2n^2 - 1$ .

**Proof.** First we establish the first part of the statement. The numerator in the first factor is bounded and the denominator in the second factor is bounded away from zero as  $\kappa \rightarrow \epsilon$  since  $\epsilon \neq 1$ .

If  $\prod_i x_i y_i < 1 - \delta$  then the rest of the integrand is bounded except for the last quotient, and the integral of that is  $O(1)$ .

If  $\prod_i x_i y_i > 1 - \delta$  then each  $x_i, y_i > 1 - \delta$  and the integrand has absolute value at most a constant times

$$\frac{\Delta(x)^2 \Delta(y)^2}{|\kappa^{-n} - \prod_i x_i y_i|^{\ell+1}} \prod_i \sqrt{\frac{1 - x_i}{1 - y_i}}.$$

We assumed that  $\kappa \rightarrow \epsilon$  along a radius, so  $\kappa^{-n} > 1$ . Therefore we get an upper bound if we replace  $\kappa^{-n}$  by 1. Then in the integral we make the substitutions  $x_i = 1 - \xi_i$ ,  $y_i = 1 - \eta_i$  (so  $\xi_i, \eta_i < \delta$ ) and we obtain

$$\frac{\Delta(\xi)^2 \Delta(\eta)^2}{(1 - \prod_i (1 - \xi_i)(1 - \eta_i))^{\ell+1}} \prod_i \sqrt{\frac{\xi_i}{\eta_i}}.$$

Whenever  $z_i \in [0, 1]$  ( $i = 1, \dots, m$ ) we have  $z_1 \cdots z_m \leq z_i$  for each  $i$ , and so averaging gives

$$z_1 \cdots z_m \leq \left( \sum_j z_j \right) / m,$$

and therefore

$$1 - z_1 \cdots z_m \geq \sum_j (1 - z_j) / m.$$

It follows that

$$1 - \prod_i (1 - \xi_i)(1 - \eta_i) \geq \sum_i (\xi_i + \eta_i) / 2n. \quad (14)$$

Therefore the integrand above is at most  $(2n)^{\ell+1}$  times

$$\frac{\Delta(\xi)^2 \Delta(\eta)^2}{(\sum_i (\xi_i + \eta_i))^{\ell+1}} \prod_i \sqrt{\frac{\xi_i}{\eta_i}}.$$

This is homogeneous of degree  $2n(n-1) - \ell - 1$ . We first integrate over the region  $\sum_i(\xi_i + \eta_i) = r$  and then over  $r$ . The resulting integral is at most a constant times

$$\int_0^{2n\delta} r^{2n^2 - \ell - 2} dr.$$

This is finite when  $\ell < 2n^2 - 1$ , and so the first statement of the lemma is established. We note that the  $(2n-1)$ -dimensional volume of the region  $\sum_i(\xi_i + \eta_i) = 1$  is  $1/\Gamma(2n)$ , another nice factor which we can use if needed. But it won't be.

We now consider the integral when  $\ell = 2n^2 - 1$ . As before, the integral over the region  $\prod_i x_i y_i < 1 - \delta$  is  $O(1)$ , so we assume  $\prod_i x_i y_i > 1 - \delta$ . In particular each  $x_i, y_i > 1 - \delta$ . The factors  $1 - \kappa x_i y_j$  in the second denominator equal  $1 - \kappa(1 + O(\delta)) = (1 - \kappa)(1 + O(\delta))$  since  $\kappa$  is bounded away from 1. From this we see that if we factor out  $\kappa^{2n^3}$  from the first denominator and  $(1 - \kappa)^{n^2}$  from the second, the integrand becomes

$$\frac{\Delta(x)^2 \Delta(y)^2}{(\kappa^{-n} - \prod_i x_i y_i)^{2n^2}} \prod_i \sqrt{\frac{1 - x_i}{1 - y_i}} (1 + O(\delta)).$$

We again make the substitutions  $x_i = 1 - \xi_i, y_i = 1 - \eta_i$  and set  $r = \sum_i(\xi_i + \eta_i)$ . Then since  $\prod_i(1 - \xi_i)(1 - \eta_i) = 1 - r + O(r^2)$  the integrand becomes

$$\frac{\Delta(\xi)^2 \Delta(\eta)^2}{(\kappa^{-n} - 1 + r + O(r^2))^{2n^2}} \prod_i \sqrt{\frac{\xi_i}{\eta_i}} (1 + O(\delta)).$$

The integration domain  $\prod_i x_i y_i > 1 - \delta$  becomes  $r + O(r^2) < \delta$ , which is contained in  $r < 2\delta$  and contains  $r < \delta/2$ . The integral without the  $O(\delta)$  term is at least a constant times

$$\int_0^{\delta/2} \frac{r^{2n^2 - 1}}{(\kappa^{-n} - 1 + 2r)^{2n^2}} dr,$$

which is asymptotically a constant independent of  $\delta$  times  $\log(\kappa^{-n} - 1)^{-1}$  as  $\kappa \rightarrow \epsilon$ . Similarly the integral of the  $O(\delta)$  term is at most a constant independent of  $\delta$  times  $\delta \log(\kappa^{-n} - 1)^{-1}$ . Since  $\delta$  is arbitrarily small, this proves the lemma.  $\square$

**Lemma 2.** We have

$$\left(\frac{d}{d\kappa}\right)^{2n^2 - 1} \mathcal{S}_n \approx \log(1 - |\kappa|)^{-1}.$$

**Proof.** To compute the derivative of the integral in (12) one integral we get is a constant depending on  $n$  times (13) with  $\ell = 2n^2 - 1$ . The other integrals are similar but in each the power in the denominator is less than  $2n^2 - 1$  while we get extra factors obtained by differentiating the rest of the integrand for  $\mathcal{S}_n$ . These factors are of the form  $(1 - \kappa x_i y_i)^{-1}$ ,  $(1 - \kappa x_i)^{-1}$ , or  $(1 - \kappa y_i)^{-1}$ . By an obvious modification

of the first statement of Lemma 1 we see that these other integrals are all bounded. The lemma follows.  $\square$

**Lemma 3.** If  $\epsilon^m \neq 1$  then

$$\left(\frac{d}{d\kappa}\right)^{2n^2-1} \mathcal{S}_m = O(1).$$

**Proof.** If  $\epsilon^m \neq 1$  all integrands obtained by differentiating the integral in (12) are bounded as  $\kappa \rightarrow \epsilon$ .  $\square$

**Lemma 4.** We have

$$\sum_{m>n} \left(\frac{d}{d\kappa}\right)^{2n^2-1} \mathcal{S}_m = O(1).$$

**Proof.** We shall show that for  $\kappa$  sufficiently close to  $\epsilon$  all integrals we get by differentiating the integral for  $\mathcal{S}_m$  are at most  $A^m m^m$ , where  $A$  is some constant. Note that the value of  $A$  will change with each of its appearances. It may depend on  $n$ , but not on  $m$ . Because of the  $1/(m!)^2$  appearing in front of the integrals this will show that the sum is bounded.

As before, we first use (12) (with  $n$  replaced by  $m$ ) and consider the integral we get when the first factor in the integrand is differentiated  $2n^2 - 1$  times. All indices in the integrands now run from 1 to  $m$ .

First,

$$|1 - \kappa^m \prod_i x_i y_i| = |\kappa^m| |\kappa^{-m} - \prod_i x_i y_i| \geq |\kappa|^m (1 - \prod_i x_i y_i).$$

Next,  $|1 - \kappa x_i| \leq 2$ . Since  $y_i \in [0, 1]$  and  $\kappa \in [0, \epsilon]$  we also have  $\kappa y_i \in [0, \epsilon]$ . Therefore  $|1 - \kappa y_i| \geq a$ , where  $a = \text{dist}(1, [0, \epsilon])$ . Hence the integrand in (12) after differentiating the first factor has absolute value at most  $A^m$  times

$$\frac{1}{(1 - \prod x_i y_i)^{2n^2}} \frac{\Delta(x)^2 \Delta(y)^2}{\prod_{i,j} |1 - \kappa x_i y_j|^2} \prod_i \sqrt{\frac{1 - x_i}{1 - y_i} \frac{y_i}{x_i}}. \quad (15)$$

Since we also have

$$|1 - \kappa x_i y_j| \geq a, \quad (16)$$

(15) is at most

$$a^{-m^2} \frac{\Delta(x)^2 \Delta(y)^2}{(1 - \prod x_i y_i)^{2n^2}} \prod_i \sqrt{\frac{1 - x_i}{1 - y_i} \frac{y_i}{x_i}}.$$

With  $x_i = 1 - \xi_i$ ,  $y_i = 1 - \eta_i$  and  $r = \sum_i(\xi_i + \eta_i)$  again, we first integrate over  $r < \delta$ , where the small  $\delta$  will be chosen below. Using (14) again, we see that the integrand is at most  $A^m$  times

$$a^{-m^2} \frac{\Delta(\xi)^2 \Delta(\eta)^2}{(\sum_i(\xi_i + \eta_i))^{2n^2}} \prod_i \sqrt{\xi_i/\eta_i}.$$

(The factor  $m^{2n^2}$  coming from using (14) and a bound for  $\prod \sqrt{y_i/x_i}$  appearing in (15) were absorbed into  $A^m$ .) When  $\xi_i, \eta_i < 1$  we have  $\Delta(\xi)^2, \Delta(\eta)^2 < 1$ , so integrating with respect to  $r$  over  $r < \delta$ , using homogeneity, gives at most a constant times

$$a^{-m^2} \int_0^\delta r^{2m(m-1)-2n^2+2m-1} dr = a^{-m^2} \int_0^\delta r^{2m^2-2n^2-1} dr.$$

(The integral of the last factor over  $r = 1$  equals  $(\pi/2)^m/\Gamma(2m)$ .) The integral is  $O(\delta^{2m^2})$  since  $m > n$ , and so the above is exponentially small in  $m$  if we choose  $\delta^2 < a$ .

There remains the integral over the region  $r > \delta$ , and for this we use the representation (11). We are led to (15) with the second factor replaced by the absolute value of

$$\left( \det \left( \frac{1}{1 - \kappa x_i y_j} \right) \right)^2.$$

From (14) we see that in this region the first factor in (15) is at most  $(2m/\delta)^{2n^2}$ . By (16) and the Hadamard inequality the square of the determinant has absolute value at most  $a^{-2m} m^m$ . Therefore the integral over this region has absolute value at most

$$\left( \frac{2m}{\delta} \right)^{2n^2} a^{-2m} m^m \int_0^1 \cdots \int_0^1 \prod_i \sqrt{\frac{1-x_i y_i}{1-y_i x_i}} \prod_i dx_i dy_i.$$

The integral here is  $A^m$ , and so we have shown that the integral in the region  $r > \delta$  is at most  $A^m m^m$ .

This is a bound for only one term we get when we differentiate  $2n^2 - 1$  times the integrand for  $\mathcal{S}_m$ . The number of factors in the integrand involving  $\kappa$  is  $O(m^2)$  so if we differentiate  $2n^2 - 1$  times we get a sum of  $O(m^{4n^2})$  terms. In each of the other terms the denominator in the first factor has a power no larger than  $2n^2$  and at most  $2n^2$  extra factors appear which are of the form  $(1 - \kappa x_i y_i)^{-1}$ ,  $(1 - \kappa x_i)^{-1}$ , or  $(1 - \kappa y_i)^{-1}$ . Each has absolute value at most  $a^{-1}$ , so their product is  $O(1)$ . It follows that we have the bound  $A^m m^m$  for the sum of these integrals. Lemma 4 is established.  $\square$

**Proof of the Theorem.** Suppose  $\kappa \rightarrow \epsilon$  radially, where  $\epsilon \neq 1$  is a root of unity. It is a primitive  $n$ th root of unity for some  $n$ . Then  $\epsilon^m \neq 1$  when  $m < n$  so Lemma 3

applies for these  $m$ . Combining this with Lemma 4 and Lemma 2 gives

$$\left(\frac{d}{d\kappa}\right)^{2n^2-1} \mathcal{S} \approx \log(1 - |\kappa|)^{-1}.$$

Therefore  $\mathcal{S}$  cannot be analytically continued beyond any such  $\epsilon$ , and these are dense in the unit circle.  $\square$

**Remark.** From the proofs of Lemma 3 and 4, with  $2n^2 - 1$  replaced by  $\ell$  in both, one can see that  $\mathcal{S}$  extends to a  $C^\ell$  function of  $\kappa$  up to the boundary except for the  $m$ th roots of unity with  $m \leq \sqrt{(\ell + 1)/2}$ . In particular,  $\mathcal{S}$  extends to a function of class  $C^6$  up to the boundary except for  $\kappa = 1$ .

### Acknowledgments

The authors wish to thank Bernie Nickel and Barry McCoy for their helpful comments. This work was supported by the National Science Foundation through grants DMS-1207995 (first author) and DMS-0854934 (second author).

## References

- [1] C. Andréief, Note sur une relation les intégrales définies des produits des fonctions, *Mém. de la Soc. Sci., Bordeaux* **2** (1883), 1–14.
- [2] M. Assis, S. Boukraa, S. Hassani, M. Van Hoesij, J.-M. Maillard, and B. M. McCoy, Diagonal Ising susceptibility: elliptic integrals, modular forms and Calabi-Yau equations. *J. Phys. A: Math. Theor.* **45** (2012), 075205 (32 pp).
- [3] E. Basor and H. Widom, On a Toeplitz determinant identity of Borodin and Okounkov, *Integral Equations and Operator Theory* **37** (2000), 397–401.
- [4] R. J. Baxter, Onsager and Kaufman’s calculation of the spontaneous magnetization of the Ising model, *J. Stat. Phys.* **145** (2011), 518–548.
- [5] A. Borodin and A. Okounkov, A Fredholm determinant formula for Toeplitz determinants, *Integral Equations and Operator Theory* **37** (2000), 386–396.
- [6] A. Böttcher, One more proof of the Borodin-Okounkov formula for Toeplitz determinants, *Integral Equations and Operator Theory* **41** (2001), 123–125.
- [7] S. Boukraa, S. Hassani, J.-M. Maillard, B. M. McCoy and N. Zenine, The diagonal Ising susceptibility, *J. Phys. A: Math. Theor.* **40** (2007), 8219–8236.

- [8] A. I. Bugrii, Correlation function of the two-dimensional Ising model on a finite lattice: I, *Theor. and Math. Phys.* **127** (2001), 528–548.
- [9] A. I. Bugrii and O. O. Lisovyy, Correlation function of the two-dimensional Ising model on a finite lattice: II, *Theor. and Math. Phys.* **140** (2004), 987–1000.
- [10] Y. Chan, A. J. Guttmann, B. G. Nickel and J. H. H. Perk, The Ising susceptibility scaling function, *J. Stat. Phys.* **145** (2011), 549–590.
- [11] P. Deift, A. Its and I. Krasovsky, Toeplitz matrices and Toeplitz determinants under the impetus of the Ising model: Some history and some recent results, *Commun. on Pure and Appl. Math.* **66** (2013), 1360–1438.
- [12] J. S. Geronimo and K. M. Case, Scattering theory and polynomials orthogonal on the unit circle, *J. Math. Phys.* **20** (1979), 299–310.
- [13] A. J. Guttmann and I. G Enting, Solvability of some statistical mechanical systems, *Phys. Rev. Letts.* **76** (1996), 344–347.
- [14] I. Lyberg and B. M. McCoy, Form factor expansion of the row and diagonal correlation functions of the two-dimensional Ising model, *J. Phys. A: Math. Theor.* **40** (2007), 3329–3346.
- [15] B. M. McCoy and T. T. Wu, *The Two-Dimensional Ising Model*, Harvard University Press, 1973.
- [16] B. M. McCoy, *Advanced Statistical Mechanics*, Oxford University Press, Great Clarendon Street, Oxford, 2010.
- [17] B. M. McCoy, M. Assis, S. Boukraa, S. Hassani, J.-M. Maillard, W. P. Orrick and N. Zenine, The saga of the Ising susceptibility, in *New Trends in Quantum Integrable Systems: Proceedings of the Infinite Analysis 09*, eds. B. L. Feigin, M. Jimbo and M. Okado, World Scientific, 2010, pp. 287–306. [preprint available at arXiv:1003.0751.].
- [18] B. Nickel, On the singularity structure of the 2D Ising model, *J. Phys. A: Math. Gen.* **32** (1999), 3889–3906.
- [19] B. Nickel, Addendum to ‘On the singularity structure of the 2D Ising model’, *J. Phys. A: Math. Gen.* **33** (2000), 1693–1711.
- [20] L. Onsager, Crystal statistics, I. A two-dimensional model with an order-disorder transition, *Phys. Rev.* **65** (1944), 117–149.
- [21] L. Onsager, discussion, *Nuovo Cimento* **6** Suppl. (1949), 261.

- [22] W. P. Orrick, B. Nickel, A. J. Guttmann and J. H. H. Perk, The susceptibility of the square lattice Ising model: New developments, *J. Stat. Phys.* **102** (2001), 795–841.
- [23] J. Palmer, *Planar Ising Correlations*, Progress in Mathematical Physics, Vol. 49, Birkhäuser, Boston, 2007.
- [24] J. Stephenson, Ising-model spin correlations on the triangular lattice, *J. Math. Phys.* **5** (1964), 1009–1024.
- [25] C. A. Tracy and H. Widom, Correlation functions, cluster functions, and spacing distributions for random matrices, *J. Stat. Phys.* **92** (1998), 809–835.
- [26] N. S. Witte and P. J. Forrester, Fredholm determinant evaluations of the Ising model diagonal correlations and their  $\lambda$  generalization, *Studies in Appl. Math.* **128** (2011), 183–223.
- [27] T. T. Wu, B. M. McCoy, C. A. Tracy and E. Barouch, Spin-spin correlation functions for the two-dimensional Ising model: Exact theory in the scaling region, *Phys. Rev.* **B13** (1976), 315–374.
- [28] C. N. Yang, The spontaneous magnetization of the two-dimensional Ising model, *Phys. Rev.* **85** (1952), 808–816.