

Adsorbed self-avoiding walks subject to a force

E. J. Janse van Rensburg¹ and S. G. Whittington²

¹Department of Mathematics and Statistics, York University, Toronto, Canada

²Department of Chemistry, University of Toronto, Toronto, Canada

Abstract. We consider a self-avoiding walk model of polymer adsorption where the adsorbed polymer can be desorbed by the application of a force. In this paper the force is applied normal to the surface at the last vertex of the walk. We prove that the appropriate limiting free energy exists where there is an applied force and a surface potential term, and prove that this free energy is convex in appropriate variables. We then derive an expression for the limiting free energy in terms of the free energy without a force and the free energy with no surface interaction. Finally we show that there is a phase boundary between the adsorbed phase and the desorbed phase in the presence of a force, prove some qualitative properties of this boundary and derive bounds on the location of the boundary.

1. Introduction

Polymer adsorption at an impenetrable surface [1] is a topic that has been attracting interest from theorists for at least fifty years [24, 26]. Various polymer models have been examined, including random walks [24, 27], directed and partially directed walks [2, 14, 29] and self-avoiding walks [4, 7, 12, 18]. Techniques like atomic force microscopy [6] allow adsorbed polymers to be pulled off a surface and this problem has also attracted attention [19, 20, 23, 27]. Variants of the problem such as pulling at an angle [21, 22, 25, 28] and from inhomogeneous surfaces [9, 10] have also been studied.

The standard lattice model of polymer adsorption is self-avoiding walks. In the absence of a force this problem is quite well understood [1, 4, 12], though many details remain as open questions. For instance, although there is a rigorous proof of the existence of a phase transition, the order of the transition is not known rigorously. When the adsorbed self-avoiding walk is subject to a force, numerical evidence shows that self-avoiding walks are well approximated by partially directed walks [9], at least for large forces. Nevertheless, at the rigorous level we know very little about the behaviour of self-avoiding walks in these circumstances. The phase transition (*i.e.* desorption) under the influence of a force is expected to be first order but this is not known rigorously.




Figure 1. A pulled and adsorbing self-avoiding walk in the positive half-lattice. Visits of the walk to the adsorbing line are weighted by a , while a force f is pulling the path at its endpoint in the vertical direction.

In this paper we consider the simplest case, that of a self-avoiding walk terminally attached to an impenetrable surface at which the walk adsorbs, and pulled from the other unit degree vertex in a direction *normal* to the surface. We prove the existence of the appropriate thermodynamic limit and some qualitative properties of the limiting free energy. In addition we derive some information about the form of the phase diagram for the model.

2. Thermodynamic limits and convexity

Consider the d -dimensional hypercubic lattice Z^d where the vertices have integer coordinates (x, y, \dots, z) . We write (x_i, y_i, \dots, z_i) , $i = 0, 1, 2, \dots, n$ for the coordinates of the i -th vertex of an n -

step self-avoiding walk on Z^d . The number of n -step self-avoiding walks from the origin is denoted by c_n . It is known that $\lim_{n \rightarrow \infty} \frac{1}{n} \log c_n = \log \mu_d$ exists [3], where μ_d is the *growth constant* of self-avoiding walks.

A *positive walk* is a self-avoiding walk on Z^d that starts at the origin and is constrained to have $z_i \geq 0$ for all $0 \leq i \leq n$. The number of n -step positive walks from the origin is denoted by c_n^+ . It is known that $\lim_{n \rightarrow \infty} \frac{1}{n} \log c_n^+ = \log \mu_d$ [4]. Vertices of a positive walk in the hyperplane $z = 0$ are *visits* although, by convention, the vertex at the origin is not counted as a visit. The number of positive walks of n -steps from the origin with v visits is denoted by $c_n^+(v)$.

An n -step positive walk is a *loop* if $z_n = 0$. A positive walk (or a loop) is *unfolded in the x -direction* if $0 \leq x_i < x_n$, $i = 1, 2, \dots, n-1$, with similar definitions for the y -direction, up to and including the z -direction. See [5].




Figure 2. A loop and an unfolded loop on the square lattice.

The number of n -step loops with v visits is denoted $l_n(v)$. Similarly we write $c_n^\ddagger(v)$ and $l_n^\ddagger(v)$ for the corresponding numbers of positive walks and loops that are unfolded in the x -direction. We define the corresponding partition functions

$$C_n(a) = \sum_v c_n^+(v) a^v, \quad L_n(a) = \sum_v l_n(v) a^v \quad (2.1)$$

and

$$C_n^\ddagger(a) = \sum_v c_n^\ddagger(v) a^v, \quad L_n^\ddagger(a) = \sum_v l_n^\ddagger(v) a^v. \quad (2.2)$$

A *tail* is a positive walk that leaves the surface on its first step and never returns. Clearly the number of n -step tails is $c_n^+(0)$.

Next we recall a useful lemma.

Lemma 1 *For unfolding in the x -direction and $a > 0$ the following limits exist and are equal:*

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log C_n(a) = \lim_{n \rightarrow \infty} \frac{1}{n} \log L_n(a) = \lim_{n \rightarrow \infty} \frac{1}{n} \log C_n^\ddagger(a) = \lim_{n \rightarrow \infty} \frac{1}{n} \log L_n^\ddagger(a) \equiv \kappa(a).$$

Also, $\kappa(a) = \log \mu_d$ for $a \leq 1$. \square

See [4] for the proof of this lemma.

We generalize the model above to a two-parameter model of *pulled adsorbing positive walks*.

Let $c_n^+(v, h)$ be the number of n -step positive walks with v visits and with $z_n = h$. Then $c_n^+(0, h)$ is the number of n -step tails with $z_n = h$. We call h the *height* of the last vertex.

We write $c_n^\dagger(v, h)$ for the numbers of the corresponding unfolded walks. The corresponding partition functions are

$$C_n(a, y) = \sum_{v, h} c_n^\dagger(v, h) a^v y^h, \quad T_n(y) = \sum_h c_n^\dagger(0, h) y^h \quad (2.3)$$

and

$$C_n^\ddagger(a, y) = \sum_{v, h} c_n^\ddagger(v, h) a^v y^h, \quad T_n^\ddagger(y) = \sum_h c_n^\ddagger(0, h) y^h. \quad (2.4)$$

The following lemma was proved in [17].

Lemma 2 *For unfolding in the x-direction the following limits exist and are equal:*

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log T_n(y) = \lim_{n \rightarrow \infty} \frac{1}{n} \log T_n^\ddagger(y) = \lambda(y).$$




Figure 3. The circled vertex is the last visit of the walk. At this vertex the walk can be decomposed into a loop and a tail.

We extend these two lemmas by proving the existence of the thermodynamic limit for the two parameter adsorbing and pulled positive walk model.

Theorem 1 *The limit $\lim_{n \rightarrow \infty} \frac{1}{n} \log C_n(a, y)$ exists and is equal to $\max[\kappa(a), \lambda(y)]$.*

Proof: Every positive walk is either a loop or it eventually leaves the surface for the last time. This decomposes positive walks into a loop, followed by a tail (either the loop, or the tail, may be empty) – see Figure 3.

If the loop and the tail in the decomposition are both unfolded, then a lower bound is obtained:

$$C_n(a, y) \geq \sum_{m=0}^n L_m^\ddagger(a) T_{n-m}^\ddagger(y)$$

since the two unfolded parts must be distinct except for the vertex where the loop ends and the tail begins. In particular, by only considering the $m = 0$ or $m = n$ terms on the right hand side, the following lower bound on $C_n(a, y)$ is obtained:

$$C_n(a, y) \geq \max[L_n^\ddagger(a), T_n^\ddagger(y)]$$

and therefore

$$\liminf_{n \rightarrow \infty} \frac{1}{n} \log C_n(a, y) \geq \max[\kappa(a), \lambda(y)]$$

where we have made use of Lemma 1 and Lemma 2.

To obtain an upper bound on $C_n(a, y)$ we again consider the loop plus tail decomposition in Figure 3. If all loops and tails are considered, then an upper bound on $C_n(a, y)$ follows:

$$C_n(a, y) \leq \sum_m L_m(a) T_{n-m}(y).$$

Define the generating functions

$$\widehat{L}(a, t) = \sum_n L_n(a) t^n, \quad \widehat{T}(y, t) = \sum_n T_n(y) t^n.$$

Using the convolution theorem

$$\widehat{L}(a, t) \widehat{T}(y, t) = \sum_n \sum_{m=0}^n L_m(a) T_{n-m}(y) t^n = \sum_n B_n(a, y) t^n = \widehat{B}(a, y, t)$$

and so $C_n(a, y) \leq B_n(a, y)$.

The radius of convergence of $\widehat{L}(a, t)$ is $t = t_1 = \exp(-\kappa(a))$ and of $\widehat{T}(y, t)$ is $t = t_2 = \exp(-\lambda(y))$. Thus, the product $\widehat{B}(a, y, t)$ has radius of convergence $\min\{t_1, t_2\}$ and is singular at $t = t_1$ and at $t = t_2$.

Hence $\limsup_{n \rightarrow \infty} \frac{1}{n} \log B_n(a, y) = \max[\kappa(a), \lambda(y)]$ and therefore

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \log C_n(a, y) \leq \max[\kappa(a), \lambda(y)].$$

Hence the thermodynamic limit exists and $\lim_{n \rightarrow \infty} \frac{1}{n} \log C_n(a, y) = \max[\kappa(a), \lambda(y)]$. \square

We shall write

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log C_n(a, y) = \kappa(a, y) \tag{2.5}$$

so $\kappa(a, y) = \max[\kappa(a), \lambda(y)]$.

Theorem 2 *The limiting free energy $\kappa(a, y) = \lim_{n \rightarrow \infty} \frac{1}{n} \log C_n(a, y)$ is a convex function of $\log a$ and $\log y$.*

Proof: The theorem is a consequence of Hölder's inequality. For any p such that $0 \leq p \leq 1$,

$$\sum_{v,h} c_n^+(v, h) (a_1^p a_2^{1-p})^v (y_1^p y_2^{1-p})^h \leq \left(\sum_{v,h} c_n^+(v, h) a_1^v y_1^h \right)^p \left(\sum_{v,h} c_n^+(v, h) a_2^v y_2^h \right)^{1-p} \tag{2.6}$$

Taking logarithms and dividing by n gives

$$\frac{1}{n} \log C_n(a_1^p a_2^{1-p}, y_1^p y_2^{1-p}) \leq p \frac{1}{n} \log C_n(a_1, y_1) + (1-p) \frac{1}{n} \log C_n(a_2, y_2) \tag{2.7}$$

so $\frac{1}{n} \log C_n(a, y)$ is a convex function of $\log a$ and $\log y$. Since the limit of a sequence of convex functions, when it exists, is also convex, this proves the theorem. \square

It follows that the free energy is convex as a surface and not only in the coordinate directions.

3. The free energy

In this section we turn our attention to the free energy $\kappa(a, y)$ given by equation (2.5) in terms of $\kappa(a)$ (the free energy of an adsorbing walk) and $\lambda(y)$ (the free energy of a pulled walk). We first review the properties of $\kappa(a)$ and $\lambda(y)$, and prove that $\kappa(a)$ is asymptotic to $\log \mu_{d-1} + \log a$ and $\lambda(y)$ is asymptotic to $\log y$.

3.1. Bounds on $\kappa(a)$

We first recall some earlier results for the adsorption problem without a force [4].

It is known that $\kappa(a)$ is singular at a critical point at $a = a_c^o$ corresponding to the adsorption transition in this model. It is known that $1 < a_c^o < \frac{\mu_d}{\mu_{d-1}}$, see for example [4], or [13] for a review. The order of the adsorption transition is not known rigorously but is thought to be continuous [4, 7, 18]. For $a < a_c^o$, $\kappa(a) = \kappa(1) = \log \mu_d$, independent of a , and for $a > a_c^o$

$$\max \{ \log \mu_d, \log \mu_{d-1} + \log a \} \leq \kappa(a) \leq \log \mu_d + \log a. \quad (3.1)$$

The average number of visits in an adsorbing loop is given by

$$\langle v \rangle_n = a \frac{d}{da} \log \sum_v l_n(v) a^v = \frac{\sum_v v l_n(v) a^v}{\sum_v l_n(v) a^v}; \quad (3.2)$$

see equation (2.1). Since $\frac{1}{n} \log L_n(a)$ is a sequence of convex functions converging to a convex limit $\kappa(a)$, one may interchange the limit and the derivative almost everywhere in

$$\mathcal{E}(a) = \lim_{n \rightarrow \infty} \frac{1}{n} \langle v \rangle_n = \lim_{n \rightarrow \infty} \frac{a}{n} \frac{d}{da} \log L_n(a) = a \frac{d}{da} \kappa(a) \quad (3.3)$$

where $\mathcal{E}(a)$ is equal to the density of visits almost everywhere.

The left- and right-densities of visits may be defined as follows

$$\mathcal{E}_-(a) = a \frac{d^-}{da} \kappa(a), \quad \mathcal{E}_+(a) = a \frac{d^+}{da} \kappa(a). \quad (3.4)$$

These exist for every finite $a > 0$ since $\kappa(a)$ is a convex function of $\log a$. Moreover $\mathcal{E}_-(a)$ is lower semi-continuous and $\mathcal{E}_+(a)$ is upper semi-continuous. It follows that $\mathcal{E}_-(a) = \mathcal{E}_+(a) = \mathcal{E}(a)$ for almost every a , and $\mathcal{E}_-(a) \leq \mathcal{E}_+(a)$ for every $a > 0$.

The density of visits, and $\mathcal{E}_-(a)$ and $\mathcal{E}_+(a)$ are monotone functions, and they are differentiable almost everywhere. Clearly $\mathcal{E}(a) = 0$ if $a < a_c^o$ and $\mathcal{E}(a) > 0$ for almost all $a > a_c^o$.

The Legendre transform of $\kappa(a)$ is defined by

$$\log \mathcal{P}(\alpha) = \inf_{a > 0} \{ \kappa(a) - \alpha \log a \}. \quad (3.5)$$

The function $\log \mathcal{P}(\alpha)$ is a concave function of α (see for example [13]). The inverse transform is

$$\kappa(a) = \sup_{\alpha \in [0,1]} \{ \log \mathcal{P}(\alpha) + \alpha \log a \}, \quad (3.6)$$

see reference [13].

The supremum in equation (3.6) may be realised when $\alpha = \alpha^* = 0$ or $\alpha = \alpha^* = 1$, in which case $\kappa(a) = \log \mathcal{P}(0)$ or $\kappa(a) = \log \mathcal{P}(1) + \log a$. (This occurs at boundaries of the range of α in equation (3.6)). In these cases we note by equation (3.3) that $\mathcal{E}(a) = 0 = \alpha^*$ or $\mathcal{E}(a) = 1 = \alpha^*$, respectively.

Otherwise we notice that $\log \mathcal{P}(\alpha) + \alpha \log a$ is a concave function of $\log a$. Hence, it has left-derivatives and right-derivatives everywhere. Since $\kappa(a)$ is defined for every $a \geq 0$, there exists an

$\alpha^* \in [0, 1]$ solving equation (3.6). If $\alpha^* \neq 0$ and $\alpha^* \neq 1$, then it must be in $(0, 1)$, in which case one may instead consider the solutions of

$$\frac{d^-}{d\alpha} (\log \mathcal{P}(\alpha) + \alpha \log a) = 0, \quad \text{or} \quad \frac{d^+}{d\alpha} (\log \mathcal{P}(\alpha) + \alpha \log a) = 0. \quad (3.7)$$

The solutions of these (α_- and α_+ , respectively) will be equal for almost every a (because of concavity). Notice that the solutions will be continuous almost everywhere, except where $\log \mathcal{P}(\alpha) + \alpha \log a$ is not differentiable. In that case $\alpha_- < \alpha_+$ and since $\log \mathcal{P}(\alpha)$ is a concave function of α , both α_- and α_+ are monotone and so differentiable almost everywhere, while $\alpha_- \leq \alpha^* \leq \alpha_+$.

In addition, $\kappa(a) = \log \mathcal{P}(\alpha_-) + \alpha_- \log a$ for almost all $a > 0$, and similarly, $\kappa(a) = \log \mathcal{P}(\alpha_+) + \alpha_+ \log a$ for almost all $a > 0$.

Observe that $\frac{d^-}{d\alpha} \log \mathcal{P}(\alpha) = -\log a$ if $\alpha = \alpha_-$ and $\frac{d^+}{d\alpha} \log \mathcal{P}(\alpha) = -\log a$ if $\alpha = \alpha_+$.

Taking the left derivative of $\kappa(a)$ with respect to $\log a$ gives the left-density of visits $\mathcal{E}_-(a)$ which exists everywhere. Using the chain rule for differentiation it follows that for almost every a ,

$$\begin{aligned} \mathcal{E}_-(a) &= a \frac{d^-}{da} \kappa(a) = a \frac{d^-}{da} (\log \mathcal{P}(\alpha_-) + \alpha_- \log a) \\ &= \left[a \frac{d^-}{da} \alpha_- \right] \frac{d^-}{d\alpha} \log \mathcal{P}(\alpha)|_{\alpha=\alpha_-} + a \log a \left[\frac{d^-}{da} \alpha_- \right] + \alpha_- \\ &= \left[a \frac{d^-}{da} \alpha_- \right] (-\log a) + a \log a \left[\frac{d^-}{da} \alpha_- \right] + \alpha_- = \alpha_-. \end{aligned}$$

This is true for almost all a since $\log \mathcal{P}(\alpha)$ is a concave function of α , and has left- and right-derivatives everywhere while α_- is monotone and has left-derivatives almost everywhere.

Similarly, one may show that $\mathcal{E}_+(a) = a \frac{d^+}{da} \kappa(a) = \alpha_+$. Since $\alpha_- = \alpha_+$ almost everywhere, it follows that $\mathcal{E}_+(a) = \mathcal{E}_-(a)$ almost everywhere and is equal to $a \frac{d}{da} \kappa(a)$ almost everywhere.

Thus, $\mathcal{E}_+(a) = \mathcal{E}_-(a) = \mathcal{E}(a)$ almost everywhere, and we may redefine $\mathcal{E}(a) = \mathcal{E}_-(a) = \alpha_- = \alpha^*$. This shows that the supremum in equation (3.6) is realized by α^* , or by the density of visits, for almost every a . Hence

$$\lim_{n \rightarrow \infty} \frac{1}{n} \langle v \rangle_n = \mathcal{E}(a) = \alpha^*, \quad \text{for almost every } a > 0. \quad (3.8)$$

Define the function $v_n^* = \lfloor \langle v \rangle_n \rfloor$. Then we have proven the following theorem.

Theorem 3 *For almost every value of $a > 0$,*

$$\lim_{n \rightarrow \infty} \frac{1}{n} v_n^* = \mathcal{E}(a) = \alpha^*$$

where α^* realises the supremum in equation (3.6). \square

Observe that $v_n^* \rightarrow \infty$ as $n \rightarrow \infty$ if $a > a_c^o$. We proceed by finding a lower bound on the partition function $L_n(a)$ in terms of unfolded loops.

First consider walks from the origin and confined to the adsorbing plane. Suppose the walks are unfolded in the x -direction and are of length v_n^* . The number of such walks is denoted by $c_{v_n^*}^{(d-1)}$, since these are unfolded self-avoiding walks in the $(d-1)$ -dimensional lattice which is the adsorbing plane. The number of visits of these walks is v_n^* .

Next, consider unfolded loops of length $n - v_n^*$ with exactly 1 visit (the endpoint). Such loops start in the origin, step into the positive half-lattice above the adsorbing plane, and only return to the adsorbing plane at their final step. The number of such loops is $l_{n-v_n^*}^\ddagger(1)$.

Corollary 1 For every $\delta > 0$ there exists an a_δ such that for all $a > a_\delta$,

$$\log \mu_{d-1} + \log a \leq \kappa(a) \leq \log \mu_{d-1} + (1 + \delta) \log a.$$

Proof: Substitute the upper bound $\kappa(z) \leq \log \mu_d + \log z$ in the right hand side of Theorem 4 to find that for $a > z > \frac{\mu_d}{\mu_{d-1}}$,

$$\kappa(a) - \log \mu_d \leq \left(\log a - \log \frac{\mu_d}{\mu_{d-1}} \right) \left(\frac{\log z}{\log z - \log \frac{\mu_d}{\mu_{d-1}}} \right). \quad (3.11)$$

Let $\delta > 0$ and choose z so large that

$$1 \leq \frac{\log z}{\log z - \log \frac{\mu_d}{\mu_{d-1}}} \leq 1 + \delta.$$

Put $a_\delta = z$ so that $a > a_\delta$. Then for $a > a_\delta$ equation (3.11) becomes

$$\log a + \log \mu_{d-1} \leq \kappa(a) \leq \log \mu_d + (1 + \delta) \left(\log a - \log \frac{\mu_d}{\mu_{d-1}} \right),$$

where we recall the lower bound from equation (3.1). Expand and simplify the right hand side to obtain

$$\log a + \log \mu_{d-1} \leq \kappa(a) \leq (1 + \delta) \log a + (1 + \delta) \log \mu_{d-1} - \delta \log \mu_d.$$

This may be put in the form

$$\log a + \log \mu_{d-1} \leq \kappa(a) \leq \log \mu_{d-1} + (1 + \delta) \log a + \delta \log \frac{\mu_{d-1}}{\mu_d}.$$

Since $\frac{\mu_d}{\mu_{d-1}} > 1$ the last term may be discarded. \square

That is, $\kappa(a)$ is asymptotic to $\log \mu_{d-1} + \log a$ in the sense of Corollary 1.

Since $\kappa(a) = \log \mu_d$ if $a \leq 1$ and $\kappa(a) > \log \mu_d$ if $a > a_c^o$, there is an *adsorption transition* in the model where $\mathcal{E}(a)$ becomes strictly positive. These results are taken together in the following theorem.

Theorem 5 There is a critical point a_c^o separating a phase of free positive walks from a phase of adsorbed walks.

For $a < a_c^o$, $\kappa(a) = \log \mu_d$.

If $a > a_c^o$, then

$$\max\{\log \mu_d, \log \mu_{d-1} + \log a\} \leq \kappa(a) \leq \max\{\log \mu_d, \log \mu_d + \log a\}.$$

Moreover, $\kappa(a)$ is asymptotic to $\log \mu_{d-1} + \log a$ for large a , and $a_c^o \in \left[1, \frac{\mu_d}{\mu_{d-1}}\right]$. \square

3.2. Bounds on $\lambda(y)$

We first recall some earlier results for pulled self-avoiding walks (see section 2.3 in reference [17]).

It is known that $\lambda(y)$ is a non-analytic function of y with a singularity at $y = y_c^o$ where $1 \leq y_c^o \leq \mu_d$. It is also known that $\lambda(y) = \log \mu_d$ if $y \leq 1$. Bounds for $y > 1$ are also known, and are given by

$$\max\{\log \mu_d, \log y\} \leq \lambda(y) \leq \log \mu_d + \log y. \quad (3.12)$$

The number of tails from the origin of length n and height of last vertex h is $c_n^+(0, h)$. The number of unfolded tails of length n is $c_n^\dagger(0, h)$.

Then the partition function of unfolded tails is given by

$$T_n^\dagger(y) = \sum_h c_n^\dagger(0, h) y^h. \quad (3.13)$$

The average height of a pulled tail is given by

$$\langle h \rangle_n = y \frac{d}{dy} \log \sum_h c_n^\dagger(0, h) y^h. \quad (3.14)$$

Since $\frac{1}{n} \log T_n^\dagger(y)$ is a sequence of convex functions converging to a convex limit $\lambda(y)$, one may interchange the limit and the derivative almost everywhere in

$$\mathcal{E}^\lambda(y) = \lim_{n \rightarrow \infty} \frac{1}{n} \langle h \rangle_n = \lim_{n \rightarrow \infty} \frac{y}{n} \frac{d}{dy} \log T_n^\dagger(y) = y \frac{d}{dy} \lambda(y). \quad (3.15)$$

Similarly to equation (3.3), one may define left- and right-densities of the height of the endpoint:

$$\mathcal{E}_-^\lambda(y) = y \frac{d^-}{dy} \lambda(y), \quad \mathcal{E}_+^\lambda(y) = y \frac{d^+}{dy} \lambda(y). \quad (3.16)$$

These exist for every $y > 0$, while $\mathcal{E}_-^\lambda(y)$ is lower semi-continuous and $\mathcal{E}_+^\lambda(y)$ is upper semi-continuous. Notice that $\mathcal{E}_-^\lambda(y) \leq \mathcal{E}_+^\lambda(y)$ generally and $\mathcal{E}_-^\lambda(y) = \mathcal{E}_+^\lambda(y) = \mathcal{E}^\lambda(y)$ for almost all $y > 0$.

The Legendre transform of $\lambda(y)$ is defined by

$$\log \mathcal{P}^\lambda(\beta) = \inf_{\beta > 0} \{ \lambda(y) - \beta \log y \}. \quad (3.17)$$

The function $\log \mathcal{P}^\lambda(\beta)$ is a concave function of β on $[0, 1]$ (see for example reference [13]). The inverse transform is given by

$$\lambda(y) = \sup_{\beta \in [0, 1]} \{ \log \mathcal{P}^\lambda(\beta) + \beta \log y \}. \quad (3.18)$$

Since $\lambda(y)$ is defined for every $y \geq 0$, there exists a $\beta^* \in [0, 1]$ solving equation (3.18).

Next, define the function $h_n^* = \lfloor \langle h \rangle_n \rfloor$. Arguments similar to those leading to Theorem 3 then give the following theorem:

Theorem 6 *For almost every value of $y > 0$,*

$$\lim_{n \rightarrow \infty} \frac{1}{n} h_n^* = \mathcal{E}^\lambda(y) = \beta^*$$

where β^* realizes the supremum in equation (3.18). \square

A lower bound on $T_n(y)$ may be obtained by only considering unfolded tails of length $n - h^* + 1$ which end in a vertex at height 1, followed by a sequence of $h^* - 1$ vertical edges to end at height h^* . This shows that $c_{n-h_n^*+1}^\dagger(1) y^{h_n^*} \leq T_n(y)$. Take logarithms, divide by n and let $n \rightarrow \infty$ to obtain

$$(1 - \mathcal{E}^\lambda(y)) \log \mu_d + \mathcal{E}^\lambda(y) \log y \leq \lambda(y), \quad \text{for almost every } y \geq y_c^o, \quad (3.19)$$

and where $\mathcal{E}^\lambda(y) = y \frac{d}{dy} \lambda(y)$.

This gives the following lemma:

Lemma 4 *Suppose that $f(x) = \lambda(e^x)$. Then for almost every $x > \log y_c^o$,*

$$f(x) - \log \mu_d \geq \frac{d}{dx} f(x) (x - \log \mu_d).$$

Integrate the inequality in lemma 4 from $\log z$ to $\log y$ where $z > \mu_d$. This gives the following theorem.

Theorem 7 *The free energy of pulled walks satisfies the following inequality:*

$$\frac{\lambda(y) - \log \mu_d}{\log y - \log \mu_d} \leq \frac{\lambda(z) - \log \mu_d}{\log z - \log \mu_d}$$

whenever $y > z > \mu_d$.

Proof: Separate variables in Lemma 4 and integrate over $(\log z, \log y)$ where $z > \mu_d$. This gives

$$\int_{\log z}^{\log y} \frac{f'(x)}{f(x) - \log \mu_d} dx \leq \int_{\log z}^{\log y} \frac{dx}{x - \log \mu_d}.$$

Integrate both sides and collect terms. \square

Use the upper bound $\lambda(y) \leq \log \mu_d + \log y$ for $y \geq 1$ in Theorem 7. This shows that

$$\lambda(y) - \log \mu_d \leq (\log y - \log \mu_d) \left(\frac{\log z}{\log z - \log \mu_d} \right). \quad (3.20)$$

Let $\delta > 0$ and choose $z \geq 1$ so large that

$$1 \leq \frac{\log z}{\log z - \log \mu_d} \leq 1 + \delta.$$

Put $y_\delta = z$. Since $\lambda(y) \geq \log y$, the above bound in equation (3.20) becomes

$$\log y \leq \lambda(y) \leq \log \mu_d + (1 + \delta)(\log y - \log \mu_d)$$

whenever $y > y_\delta$. Expand and simplify the right hand side to obtain

$$\log y \leq \lambda(y) \leq (1 + \delta) \log y - \delta \log \mu_d.$$

This is true for all $y > z$. Hence, given $\delta > 0$ there is an y_δ such that these bounds are true for all $y > y_\delta$. Since $\mu_d > 1$ the last term in the upper bound is negative and may be discarded. This gives

Corollary 2 *For every $\delta > 0$ there exists an y_δ such that for all $y > y_\delta$,*

$$\log y \leq \lambda(y) \leq (1 + \delta) \log y.$$

\square

Thus, there is a phase transition where walks become ballistic. This follows in particular because $\lambda(y) = \log \mu_d$ if $y \leq 1$ and from the bounds in equation (3.12). These results may be taken together in the following theorem.

Theorem 8 *There is a critical point y_c^o separating a free phase from a ballistic phase in the limiting free energy $\lambda(y)$ of (vertically) pulled positive walks.*

For $y < y_c^o$, $\lambda(y) = \log \mu_d$.

If $y > y_c^o$, then

$$\max\{\log \mu_d, \log y\} \leq \lambda(y) \leq \log \mu_d + \log y.$$

Moreover, $\lambda(y)$ is asymptotic to $\log y$ for large y , and $y_c^o \in [1, \mu_d]$. \square

By introducing the reduced pulling force \hat{f} in the model by setting $y = \exp \hat{f}$, the last theorem states that there is a critical reduced force $\hat{f}_c \in [0, \log \mu_d]$ such that for $\hat{f} > \hat{f}_c$ the model is ballistic. Numerical simulations suggest that $\hat{f}_c = 0$ [17]. See also [11].

4. Phase transitions and the phase diagram of pulled adsorbing walks

In this section the phase diagram of a pulled adsorbing walk is examined. The free energy of the model is $\kappa(a, y)$ (see equation (2.5)) and $\kappa(a, y) = \max\{\kappa(a), \lambda(y)\}$. By Theorem 2 $\kappa(a, y)$ is a convex function of $\log a$ and $\log y$.

We next examine the properties of $\kappa(a, y)$ by using the results of section 3 on $\kappa(a)$ and $\lambda(y)$.

If $a < a_c^o$ and $y < y_c^o$ the $\kappa(a) = \lambda(y) = \log \mu_d$. Thus, $\kappa(a, y) = \log \mu_d$ in this regime, which is a *free* phase.

If $a < a_c^o$ is fixed, and $y < y_c^o$ is increased then there is a critical point at y_c^o where $\lambda(y) > \log \mu_d$ so that the model transitions into a *ballistic* phase (where $\kappa(a, y)$ is only dependent on y). This shows that the line segment $0 \leq a < a_c^o$ and $y = y_c^o$ is a critical line in the phase diagram, separating a free phase from a ballistic phase.

Similarly, if $y < y_c^o$ and $a < a_c^o$ is increased then there is a critical point at a_c^o where $\kappa(a) > \log \mu_d$ so that the model goes through a transition to an *adsorbed* phase. This shows that the line segment $a \leq y < y_c^o$ and $a = a_c^o$ is a critical line in the phase diagram, separating a free phase from an adsorbed phase.

Next, consider the cases that either $a > a_c^o$ or $y > y_c^o$.

Lemma 5 *For $a > a_c^o$ there exists a real number $y_c(a)$, $1 \leq y_c(a) \leq e^{\kappa(a,1)}$, such that the free energy is equal to $\kappa(a, 1)$ for $y < y_c(a)$ and, for $y > y_c(a)$,*

$$\kappa(a, y) \geq \max[\kappa(a, 1), \log y].$$

Proof: Fix $a > a_c^o$ and note that $\kappa(a, 1) = \kappa(a)$. For $y \leq 1$, $\kappa(a, y) \leq \kappa(a, 1)$ by monotonicity. Also

$$\kappa(a, y) \geq \lim_{n \rightarrow \infty} \frac{1}{n} \log L_n(a) = \kappa(a) = \kappa(a, 1)$$

and hence $\kappa(a, y) = \kappa(a, 1)$ for $y \leq 1$.

For $y \geq 1$, $\kappa(a, y) \geq \kappa(a, 1)$ by monotonicity. By extracting the single term in the partition function, where the walk is a straight line normal to the surface, $\kappa(a, y) \geq \log y$. Hence $\kappa(a, y)$ is singular at a point $y = y_c(a)$ where $1 \leq y_c(a) \leq e^{\kappa(a,1)} = e^{\kappa(a)}$. \square

The scaled average height of the last vertex, $\frac{1}{n} \langle h \rangle_n = \langle z_n \rangle$, is given by

$$\frac{\langle h \rangle}{n} = \frac{\sum_{v,h} h c_n(v, h) a^v y^h}{\sum_{v,h} n c_n(v, h) a^v y^h} = \frac{1}{n} \frac{\partial}{\partial \log y} \log \sum_{v,h} c_n(v, h) a^v y^h. \quad (4.1)$$

In the $n \rightarrow \infty$ limit this becomes

$$\widehat{h} = \lim_{n \rightarrow \infty} \frac{\langle h \rangle}{n} = \frac{\partial \kappa(a, y)}{\partial \log y}, \quad \text{for almost every } (a, y), \quad (4.2)$$

where we have used the convexity of $\kappa(a, y)$ when we interchanged the order of the limit and the derivative.

Since $\kappa(a, y) = \max\{\kappa(a), \lambda(y)\}$ it follows from Lemma 5 that if $a > a_c^o$ and $y < y_c(a)$ then the free energy is independent of y and is equal to $\kappa(a)$. The limiting scaled average height of the last vertex, $\widehat{h} = \lim_{n \rightarrow \infty} \frac{1}{n} \langle h \rangle_n = 0$ and $\langle h \rangle_n = o(n)$ in this regime.

For $a > a_c^o$ and $y > y_c(a)$ the free energy is given by $\kappa(a, y) = \lambda(y)$ and this is strictly increasing with y so that $\widehat{h} > 0$. That is, the walk is in a ballistic phase.

These results, together with Theorem 1, show that we have phases where the walk is adsorbed and where the walk is ballistic, but there is no phase in which the walk is partly adsorbed and partly ballistic (such phases are present in a directed model of an adsorbing copolymer pulled in the middle [8]).

We can obtain useful upper bounds on $\kappa(a, y)$ by noting that in the adsorbed phase the free energy is independent of y and in the desorbed phase it is independent of a . Therefore, in the adsorbed phase

$$\kappa(a, y) \leq \log \mu_d + \log a \tag{4.3}$$

and in the desorbed phase

$$\kappa(a, y) \leq \log \mu_d + \log y. \tag{4.4}$$

Lemma 6 For $a > a_c^0$ the phase boundary $y = y_c(a)$ between the adsorbed phase and the desorbed ballistic phase satisfies the bounds

$$\max \left[1, a \frac{\mu_{d-1}}{\mu_d} \right] \leq y_c(a) \leq a \mu_d.$$

Proof: Fix $a > a_c^0$ and suppose that $y \geq 1$. If $y \leq y_c(a)$ we are in the adsorbed phase and $\kappa(a, y) \leq \log \mu_d + \log a$. But $\kappa(a, y) \geq \log y$. Hence if $\log y > \log \mu_d + \log a$ we have a contradiction and we are in the ballistic phase. Therefore $y_c(a) \leq a \mu_d$.

Similarly if $y \geq y_c(a)$ then $\kappa(a, y) \leq \log \mu_d + \log y$. But $\kappa(a, y) \geq \max[\log \mu_d, \log \mu_{d-1} + \log a]$. Hence $y_c(a) \geq \max \left[1, a \frac{\mu_{d-1}}{\mu_d} \right]$. \square

Taking the results above together gives the phase diagram in Figure 5.




Figure 5. The phase diagram of pulled adsorbing walks. The critical curve $y_c(a)$ separating the ballistic and adsorbed phases is asymptotic to the dashed line in the sense that $\log y_c(a) = \log(a\mu_{d-1}) + o(\log(a))$. There are numerical results indicating that $y_c^0 = 1$ (see also [11]). It is known that $a_c^0 > 1$ [4].

The phase boundary $y_c(a)$ is monotone strictly increasing, as shown below.

Lemma 7 *The phase boundary $y = y_c(a)$ is monotone strictly increasing.*

Proof: Take two points, (a_1, y_1) and (a_2, y_2) on the phase boundary. Suppose that $a_2 > a_1$. Then $\kappa(a_1) = \lambda(y_1)$ and $\kappa(a_2) = \lambda(y_2)$. But $\kappa(a)$ is strictly increasing so $\kappa(a_2) > \kappa(a_1)$ and therefore $\lambda(y_2) > \lambda(y_1)$. But $\lambda(y)$ is also strictly increasing so $y_2 > y_1$. \square

Finally, it is a corollary of Corollaries 1 and 2 that $y_c(a)$ is asymptotic to $a\mu_{d-1}$ as $a \rightarrow \infty$.

To see this, notice that the critical curve $y_c(a)$ is given by the solution of $\lambda(y) = \kappa(a)$, by definition of $\kappa(a, y)$. That is $\lambda(y_c(a)) = \kappa(a)$ for almost all $a > a_c^o$ by monotonicity of $y_c(a)$.

Choose $\delta > 0$. Since $y_c(a)$ strictly increases with a , there is an a_1 such that if $a > a_1$, then $y_c(a) > y_\delta$ in Corollary 2. Increase a , if necessary, until $a > \max\{a_1, a_\delta\}$ in Corollary 1. Then both the conditions of Corollaries 1 and 2 are valid, and so

$$\begin{aligned} \log \mu_{d-1} + \log a &\leq \kappa(a) \leq \log \mu_{d-1} + (1 + \delta) \log a, \\ \log y_c(a) &\leq \lambda(y_c(a)) \leq (1 + \delta) \log y_c(a). \end{aligned} \quad (4.5)$$

Noting that $\kappa(a) = \lambda(y_c(a))$ for almost all $a > a_c^o$, it follows that

$$\log \mu_{d-1} + \log a \leq (1 + \delta) \log y_c(a). \quad (4.6)$$

Rearrange this into $\frac{\log y_c(a)}{\log(a\mu_{d-1})} \geq \frac{1}{1+\delta}$ and take $a \rightarrow \infty$. Then $y_c(a) \rightarrow \infty$ by Lemma 7 so that for any small $\delta > 0$

$$\liminf_{a \rightarrow \infty} \frac{\log y_c(a)}{\log(a\mu_{d-1})} \geq \frac{1}{1 + \delta}. \quad (4.7)$$

On the other hand, it similarly follows from equation (4.5) that

$$\log y_c(a) \leq \log \mu_{d-1} + (1 + \delta) \log a. \quad (4.8)$$

Rearrange this into $\frac{\log y_c(a)}{\log(a\mu_{d-1})} \leq 1 + \frac{\delta \log a}{\log a + \log \mu_{d-1}}$ and take $a \rightarrow \infty$. Then $y_c(a) \rightarrow \infty$ by Lemma 7 so that for any small $\delta > 0$

$$\limsup_{a \rightarrow \infty} \frac{\log y_c(a)}{\log(a\mu_{d-1})} \leq 1 + \delta. \quad (4.9)$$

Taking $\delta \rightarrow 0^+$ in equations (4.7) and (4.9) gives the following theorem.

Theorem 9 *The critical curve $\log y_c(a)$ is asymptotic to $\log(a\mu_{d-1})$. That is,*

$$\lim_{a \rightarrow \infty} \frac{\log y_c(a)}{\log(a\mu_{d-1})} = 1.$$

Hence $\log y_c(a) = \log(a\mu_{d-1}) + o(\log a)$ as $a \rightarrow \infty$. \square

5. Discussion

We have considered a self-avoiding walk model of polymer adsorption where the adsorbed walk is subject to a force, applied normal to the surface, that tends to desorb the walk. We have proved several rigorous results about this model, including the existence of the limiting free energy, the convexity of the free energy and connections between this two variable model and the two one variable models where (a) the walk adsorbs but without an applied force, and (b) the walk is subject to a force but does not adsorb.

We have shown that there is a phase boundary in a relevant two variable space, proved some qualitative results about the boundary and derived bounds on its location. Our results show that

there are only three phases in the model, namely a free phase, a ballistic phase and an adsorbed phase. There is no phase which exhibits both the characteristic of being adsorbed and being ballistic in this model, in contrast with results on a directed path model (with the pulling force in the middle) [15, 16].

There are interesting questions about reentrant phase boundaries in models such as the one considered here [19]. The asymptotic result about the shape of the phase boundary in the large a limit, derived in the last section, in fact establishes reentrance in this model in three and more dimensions. By corollary 1 and lemma 7 for any fixed $\delta > 0$ we can take a large enough that

$$\log \mu_{d-1} + \log a \leq (1 + \delta) \log y_c(a) \quad \text{and} \quad \log y_c(a) \leq \log \mu_{d-1} + (1 + \delta) \log a. \quad (5.1)$$

If we connect to physical variables by writing $a = \exp[-\epsilon/kT]$ and $y = \exp[f/kT]$ where $\epsilon < 0$ is the energy associated with a visit, f is the applied force, k is Boltzmann's constant and T is the temperature, this relation becomes

$$(1 - \delta) (kT \log \mu_{d-1} - \epsilon) \leq f \leq kT \log \mu_{d-1} - (1 + \delta) \epsilon \quad (5.2)$$

for sufficiently small values of T . If $d \geq 3$ then $\mu_{d-1} > 1$. This shows that the critical force is an increasing function of T at sufficiently low T provided that $d \geq 3$. This establishes reentrance. Moreover,

$$\mathcal{S}_d = \lim_{T \rightarrow 0} \frac{df}{dT} = k \log \mu_{d-1} \quad (5.3)$$

which is the ground state entropy, and this is strictly positive if $d \geq 3$ (but $\mathcal{S}_2 = 0$).

This model has several interesting extensions. What happens if the force is applied at an angle to the surface? What happens if the walk or the surface has (regular) heterogeneity? We intend to explore these questions in subsequent papers.

Acknowledgements

This research was partially supported by NSERC of Canada.

References

- [1] De'Bell K and Lookman T 1993 *Rev. Mod. Phys.* **65** 87-113
- [2] Forgacs G, Privman V and Frisch H L 1989 *J. Chem. Phys.* **90** 3339-3345
- [3] Hammersley J M and Morton K W 1954 *J. Roy. Stat. Soc. B* **16** 23-38
- [4] Hammersley J M, Torrie G M and Whittington S G 1982 *J. Phys. A: Math. Gen.* **15** 539-571
- [5] Hammersley J M and Welsh D J A 1962 *Oxford Quart. J. Math.* **13** 108
- [6] Haupt B J, Ennis J and Sevick E M 1999 *Langmuir* **15** 3886-3892
- [7] Hegger R and Grassberger P 1994 *J. Phys. A: Math. Gen.* **27** 4069-4081
- [8] Iliev G K and Janse van Rensburg E J 2012 *J. Stat. Mech.* P01019
- [9] Iliev G K, Orlandini E and Whittington S G 2013 *J. Phys. A: Math. Theor.* **46** 055001
- [10] Iliev G K and Whittington S G 2012 *J. Phys. A: Math. Theor.* **45** 185003
- [11] Ioffe D and Velenik Y, *Ballistic phase of self-interacting random walks*, 2008, *Analysis and Stochastics of Growth Processes and Interface Models* ed P Morters, R Moser, M Penrose, H Schwetlick and J Zimmer Oxford: Oxford University Press) pp 55-79
- [12] Janse van Rensburg E J 1998 *J. Phys. A: Math. Gen.* **31** 8295-8306
- [13] Janse van Rensburg E J 2000 *The Statistical Mechanics of Interacting Walks, Polygons, Animals and Vesicles.* Oxford University Press.
- [14] Janse van Rensburg E J 2003 *J. Phys. A: Math. Gen.* **36** R11-R61

- [15] Janse van Rensburg E J 2010 *J. Phys. A: Math. Theo.* **43** 215001
- [16] Janse van Rensburg E J 2010 *J. Phys. A: Math. Theo.* **43** 335001
- [17] Janse van Rensburg E J, Orlandini E, Tesi M C and Whittington S G 2009 *J. Stat. Mech.* P07014
- [18] Janse van Rensburg E J and Rechnitzer A 2004 *J. Phys. A: Math. Gen.* **37** 6875-6898
- [19] Krawczyk J, Owczarek A L, Prellberg T and Rechnitzer A 2005 *J. Stat. Mech.* P05008
- [20] Orlandini E, Tesi M C and Whittington S G 1999 *J. Phys. A: Math. Gen.* **32** 469-477
- [21] Orlandini E and Whittington S G 2010 *J. Phys. A: Math. Theor.* **43** 485005
- [22] Osborn J and Prellberg T 2010 *J. Stat. Mech.* P09018
- [23] Owczarek A L 2010 *J. Phys. A: Math. Theor.* **43** 225002
- [24] Rubin R J 1965 *J. Chem. Phys.* **43** 2392-2407
- [25] Serr A and Netz R R 2007 *Europhys. Lett.* **78** 68006
- [26] Silberberg A 1962 *J. Phys. Chem.* **66** 1872-1883
- [27] Skvortsov A M, Klushin L I, Polotsky A A and Binder K 2012 *Phys. Rev. E* **85** 031803
- [28] Tabbara R and Owczarek A L 2012 *J. Phys. A: Math. Theor.* **45** 435002
- [29] Whittington S G 1998 *J. Phys. A: Math. Gen.* **31** 8797-8803