EXPANSION OF CONJUGACY CLASSES IN $PSL_2(q)$

SHELLY GARION

ABSTRACT. For any conjugacy class C in $G = PSL_2(q)$ we compute C^2 and discuss whether C contains a triple of elements whose product is 1 which generate G. Moreover, we determine which elements in G can be written as a product of two conjugate elements that generate G.

1. INTRODUCTION

There is a great interest in expansion properties of conjugacy classes in finite simple (nonabelian) groups. Thompson conjectured that every finite simple group G has a conjugacy class C such that $C^2 = G$. This conjecture was verified for many families of finite simple groups, including the alternating groups and the sporadic groups, and Ellers and Gordeev [4] proved this conjecture for all finite simple groups of Lie type defined over fields with more than 8 elements, but nevertheless it is still very much open today. On the other hand, the expansion of small enough conjugacy classes C in sufficiently large finite simple groups, has been investigated by Schul [13]. Moreover, Guralnick and Malle [8] showed that every finite simple group G has a conjugacy class C that contains a triple of elements which have product 1 and generate G.

In this paper we consider *all* the conjugacy classes in the group $G = \text{PSL}_2(q)$, and extend previous results of Guralnick and Malle [8, Lemma 3.14, Lemma 3.15 and Theorem 7.1].

Theorem 1. Let $G = PSL_2(q)$ where q > 3 and let C be a non-trivial G-conjugacy class. Then C contains elements x, y such that $\langle x, y \rangle = G$, with only the following exceptions:

- C is a conjugacy class of an element of order 2;
- q = 9 and C is a conjugacy class of a unipotent element (of order 3).

In addition, C contains three elements x, y, z such that xyz = 1 and $\langle x, y \rangle = G$ if and only if one of the following holds:

- C is a conjugacy class of a semisimple element of a q-minimal order greater than 3;
- q > 3 is prime and C is a conjugacy class of a unipotent element.

Definition 1. Let $q = p^e$ be a prime power and let n > 1 be an integer. Then n is called a *q*-minimal order if

$$= \min\{f > 0 : p^f \equiv \pm 1 \mod (\gcd(2, n) \cdot n)\}.$$

(Namely, $PSL_2(p^e)$ contains an element of order n, but no $PSL_2(p^f)$, f < e contains such an element.)

e

²⁰⁰⁰ Mathematics Subject Classification. 20D06.

The author was supported by the SFB 878 "Groups, Geometry and Actions".

Note that similar results appear in [9, 10, 11] regarding generation properties of $PSL_2(q)$ by a triple of elements (x, y, z) with product 1 of prescribed orders (see also [5, 12]).

Theorem 2. Let $G = PSL_2(q)$ where q > 2 is even.

Let C be the G-conjugacy class of an element x.

- (1) If x is a semisimple element whose order divides q-1 then $C^2 = G$.
- (2) If x is a semisimple element whose order divides q + 1 then $C^2 = G \setminus \{unipotents\}$.
- (3) If x is a unipotent element then $C^2 = G$.

In addition, only a semisimple element in G can be written as a product of two G-conjugate (semisimple) elements that generate G.

Theorem 3. Let $G = PSL_2(q)$ where q > 3 is odd.

Let C be the G-conjugacy class of an element x.

(1) If x is a semisimple element whose order is greater than 2 then $C^2 = G$.

(2) If x is an element of order 2 then
$$C^2 = \begin{cases} G & \text{if } q \equiv 1 \mod 4 \\ G \setminus \{\text{unipotents}\} & \text{if } q \equiv 3 \mod 4 \end{cases}$$

(3) If x is a unipotent element then

$$\mathcal{C}^{2} = \begin{cases} \{unipotents\} \cup \{semisimples \ of \ q-good \ orders\} \cup \{1\} & if \ q \equiv 1 \ \text{mod} \ 4 \\ \{unipotents\} \cup \{semisimples \ of \ q-good \ orders\} & if \ q \equiv 3 \ \text{mod} \ 4 \end{cases}$$

In addition, any non-trivial element in G can be written as a product of two G-conjugate semisimple elements that generate G. And moreover, a semisimple element can be written as a product of two G-conjugate unipotents that generate G if and only if its order is q-minimal and q-good, where $q \neq 9$.

Definition 2. Let $q = p^e$ be a prime power and let n > 1 be an integer. Then n is called a *q*-good order if one of the following holds:

- *n* is odd and divides either q 1 or q + 1;
- n is even and 4n divides either q 1 or q + 1.

Corollary 4. Let $G = PSL_2(q)$ and let C be the G-conjugacy class of an element x.

(1) If q is even then

$$\frac{|\mathcal{C}|}{|G|^{2/3}} \to 1 \text{ and } \frac{|\mathcal{C}^2|}{|G|} \to 1 \text{ as } q \to \infty.$$

(2) If q is odd and x is a semisimple element whose order is greater than 2 then

$$\frac{|\mathcal{C}|}{|G|^{2/3}} \to \sqrt[3]{4} and \ \frac{|\mathcal{C}^2|}{|G|} \to 1 \ as \ q \to \infty.$$

(3) If q is odd and x is an element of order 2 then

$$\frac{|\mathcal{C}|}{|G|^{2/3}} \to \frac{1}{\sqrt[3]{2}} \text{ and } \frac{|\mathcal{C}^2|}{|G|} \to 1 \text{ as } q \to \infty.$$

(4) If q is odd and x is a unipotent element then

$$\frac{|\mathcal{C}|}{|G|^{2/3}} \to \frac{1}{\sqrt[3]{2}} \text{ and } \frac{|\mathcal{C}^2|}{|G|} \to \frac{3}{4} \text{ as } q \to \infty.$$

 $\mathbf{2}$

q	order of a	q-minimal orders	q-minimal orders
	unipotent	which are q -good	which are not q -good
2	2	3	_
3	3	_	2
4	2	5	—
5	5	3	2
7	7	2,3	4
8	2	7, 9	—
9	3	5	4
11	11	3, 5	2, 6
13	13	3,7	2, 6
16	2	15, 17	—
17	17	2, 3, 4, 9	8
19	19	3, 5, 9	2,10
23	23	2, 3, 6, 11	4, 12
25	5	6, 13	4, 12
27	3	7, 13	14
29	29	3,5,7,15	2,14

Example 5. For any prime power q < 30 the following table presents all the *q*-minimal orders, divided according to whether they are *q*-good or not.

TABLE 1. Orders of elements in $PSL_2(q)$

2. Preliminaries – basic properties of $PSL_2(q)$

2.1. Elements and conjugacy classes. (see [2, 3, 6, 14]). Let $q = p^e$, where p is a prime number and $e \ge 1$. Recall that the order of $G = \text{PSL}_2(q)$ is $q(q^2 - 1)/d$, where

$$d = \gcd(2, q - 1) = \begin{cases} 1 & \text{if } q \text{ is even} \\ 2 & \text{if } q \text{ is odd} \end{cases}$$

One can classify the elements of $\text{PSL}_2(q)$ according to the Jordan form of their pre-image in $\text{SL}_2(q)$. The following Table 2 lists the three types of elements, according to whether the characteristic polynomial $P(\lambda) := \lambda^2 - \alpha \lambda + 1$ of the matrix $A \in \text{SL}_2(q)$ (where $\alpha = \text{tr}(A)$) has 0, 1 or 2 distinct roots in \mathbb{F}_q .

has 0, 1 or 2 distinct roots in \mathbb{F}_q . We denote $\kappa(\alpha) = \begin{cases} 1 & \text{if } \alpha \neq 0 \\ 2 & \text{if } \alpha = 0 \end{cases}$.

2.2. Subgroups. Table 3 specifies all the subgroups of $G = PSL_2(q)$ up to isomorphism following [14, Theorems 6.25 and 6.26].

These subgroups can be divided into the following three classes, following Macbeath [11]. The subgroups isomorphic to $PSL_2(q_1)$ or $PGL_2(q_1)$ are usually called *subfield* subgroups (since \mathbb{F}_{q_1} is a subfield of \mathbb{F}_q). Since A_4 , S_4 , A_5 and dihedral groups correspond to the finite triangle groups, that is, triangle groups $T_{r,s,t}$ such that 1/r + 1/s + 1/t > 1, we will call them

SHELLY GARION

type	roots of $P(\lambda)$	Jordan form in $\mathrm{SL}_2(\overline{\mathbb{F}}_p)$	order	G-conjugacy classes
unipotent	1 root	$\begin{pmatrix} \pm 1 & 1 \\ 0 & \pm 1 \end{pmatrix}$	p	d classes in G
		$\alpha = \pm 2$		each of size $\frac{q^2-1}{d}$
semisimple	2 roots	$\begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix}$	divides	for each α :
split		where $a \in \mathbb{F}_q^*$	$\frac{q-1}{d}$	one conjugacy class
		and $a + a^{-1} = \alpha$		of size $\frac{q(q+1)}{\kappa(\alpha)}$
semisimple	no roots	$\begin{pmatrix} a & 0 \\ 0 & a^q \end{pmatrix}$	divides	for each α :
non-split		where $a \in \mathbb{F}_{q^2}^* \setminus \mathbb{F}_q^*$	$\frac{q+1}{d}$	one conjugacy class
		$a^{q+1} = 1$ and $a^{-1} + a^q = \alpha$		of size $\frac{q(q-1)}{\kappa(\alpha)}$

TABLE 2. Elements in $PSL_2(q)$

· · ·	· 1 1	1.4.	
type	maximal order	conditions	
<i>p</i> -group	q	_	
Frobenius (Borel)	q(q-1)/d	-	
cyclic (split)	(q-1)/d	-	
dihedral (split)	2(q-1)/d		
cyclic (non-split)	(q+1)/d		
dihedral (non-split)	2(q+1)/d	_	
$\mathrm{PSL}_2(q_1)$		$q = q_1^m \ (m \in \mathbb{N})$	
$\mathrm{PGL}_2(q_1)$		$q \text{ is odd}, q = q_1^{2m} \ (m \in \mathbb{N})$	
A_4	12	q is odd; or $q = 2^e$, e even	
S_4	24	$q^2 \equiv 1 \mod 16$	
A_5	60	$p = 5 \text{ or } q^2 \equiv 1 \mod 5$	

TABLE 3. Subgroups of $PSL_2(q)$

small subgroups. For convenience we will refer to the other subgroups, namely subgroups of the Borel and cyclic subgroups, as *structural* subgroups.

Macbeath [11] classified the pairs of elements in $PSL_2(q)$ in a way which makes it easy to decide what kind of subgroup they generate.

Theorem 6. [11, Theorem 1]. For every $\alpha, \beta, \gamma \in \mathbb{F}_q$ there exist three matrices $A, B, C \in SL_2(q)$ satisfying $tr(A) = \alpha$, $tr(B) = \beta$, $tr(C) = \gamma$ and ABC = I.

Definition 3. A triple $(\alpha, \beta, \gamma) \in \mathbb{F}_q^3$ is called *singular* if $\alpha^2 + \beta^2 + \gamma^2 - \alpha\beta\gamma - 4 = 0$.

Theorem 7. [11, Theorem 2]. Let $(A, B, C) \in SL_2(q)^3$ be a triple of matrices satisfying $tr(A) = \alpha$, $tr(B) = \beta$, $tr(C) = \gamma$ and ABC = I. Then $(\alpha, \beta, \gamma) \in \mathbb{F}_q^3$ is singular if and only if the group generated by the images of A and B is a structural subgroup of $PSL_2(q)$.

2.3. Orders and traces. For an integer n > 1 we denote

(1)
$$\mathcal{T}_q(n) = \{ \alpha \in \mathbb{F}_q : \alpha = \operatorname{tr}(A), A \in \operatorname{SL}_2(q), |\bar{A}| = n \}.$$

It is easy to see from Table 2 that for any prime power q, $\mathcal{T}_q(2) = \{0\}$, $\mathcal{T}_q(3) = \{\pm 1\}$, and for any odd $q = p^e$, $\mathcal{T}_q(p) = \{\pm 2\}$. Moreover, when q is odd, then $\alpha \in \mathcal{T}_q(n)$ if and only if $-\alpha \in \mathcal{T}_q(n)$. In fact, for any prime power q and integer n > 1, $\mathcal{T}_q(n)$ can be effectively computed as follows.

Proposition 8. Denote by $\mathcal{P}_q(n)$ the set of primitive roots of unity of order n in \mathbb{F}_q .

• Let $q = 2^e$ for some positive integer e and let n > 1 be an integer, then

(2)
$$\mathcal{T}_q(n) = \begin{cases} \{0\} & \text{if } n = 2\\ \{a + a^{-1} : a \in \mathcal{P}_q(n)\} & \text{if } n \text{ divides } q - 1\\ \{b + b^q : b \in \mathcal{P}_{q^2}(n)\} & \text{if } n \text{ divides } q + 1\\ \emptyset & \text{otherwise} \end{cases}$$

• Let $q = p^e$ for some odd prime p and some positive integer e and let n > 1 be an integer, then

(3)
$$\mathcal{T}_{q}(n) = \begin{cases} \{\pm 2\} & \text{if } n = p \\ \{\pm (a + a^{-1}) : a \in \mathcal{P}_{q}(2n)\} & \text{if } n \text{ divides } \frac{q-1}{2} \\ \{\pm (b + b^{q}) : b \in \mathcal{P}_{q^{2}}(2n)\} & \text{if } n \text{ divides } \frac{q+1}{2} \\ \emptyset & \text{otherwise} \end{cases}$$

Proof. We prove the case where q is odd and n divides (q-1)/2. The other cases are similar.

Assume first that n is even. Let a be a primitive root of unity of order 2n. Then -a is also a primitive root of unity of order 2n, and $(-a)^n = a^n = -1$. Thus the matrices

$$A = \begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix} \quad \text{and} \quad -A = \begin{pmatrix} -a & 0 \\ 0 & -a^{-1} \end{pmatrix}$$

both reduce to the same element $\overline{A} \in G$ of order n. Hence, $a + a^{-1}$ and $-(a + a^{-1})$ both belong to $\mathcal{T}_q(n)$. $\mathcal{T}_q(n)$ contains only the elements of the claimed form, since it suffices to consider only the conjugacy classes of elements of G, described in Table 2.

Now assume that n is odd. Then a is a primitive root of unity of order n if and only if -a is a primitive root of unity of order 2n. In this case, $(-a)^n = -a^n = -1$. Thus the matrices

$$A = \begin{pmatrix} a & 0\\ 0 & a^{-1} \end{pmatrix} \quad \text{and} \quad -A = \begin{pmatrix} -a & 0\\ 0 & -a^{-1} \end{pmatrix}$$

both reduce to the same element $\overline{A} \in G$ of order n. Hence, $a + a^{-1}$ and $-(a + a^{-1})$ both belong to $\mathcal{T}_q(n)$. Again, considering the conjugacy classes appearing in Table 2 shows that $\mathcal{T}_q(n)$ contains only the elements of the claimed form. \Box

Proposition 9. Assume that $q = p^e$ is odd. Let $C \in SL_2(q)$, and denote $\gamma = tr(C)$ and $t = |\overline{C}|$. Assume that $\gamma \neq \pm 2$ (or equivalently, $t \neq p$). Then t is a q-good order if and only if one of $2 + \gamma$, $2 - \gamma$ is a square in \mathbb{F}_q .

Proof. As $t \neq p$ then t divides either (q-1)/2 or (q+1)/2.

If t divides (q-1)/2 then $\gamma = a + a^{-1}$ or $\gamma = -(a + a^{-1})$, for some primitive root of unity a of order 2t in \mathbb{F}_q (see Proposition 8). Hence,

$$\{2+\gamma, 2-\gamma\} = \{a+2+a^{-1}, (-a)+2+(-a)^{-1}\}.$$

Therefore, $2 + \gamma$ or $2 - \gamma$ is a square in \mathbb{F}_q if and only if $a = c^2$ or $-a = c^2$ for some $c \in \mathbb{F}_q$. Indeed, $a + 2 + a^{-1}$ is a square if and only if $(a + 1)^2/a$ is a square if and only if a is a square in \mathbb{F}_q , and similarly for $(-a) + 2 + (-a)^{-1}$.

Now, one the following necessarily holds:

- If t is even then -a is also a primitive root of unity of order 2t. Hence, a is a square in \mathbb{F}_q if and only if -a is a square is \mathbb{F}_q . In addition, a is a square in \mathbb{F}_q if and only if \mathbb{F}_q contains a primitive root of unity of order 4t, namely, if and only if 4t divides q-1.
- If t is odd and $q \equiv 1 \mod 4$ then 4t divides q 1, and so \mathbb{F}_q contains a primitive root of unity of order 4t. Thus a, which is a primitive root of unity of order 2t, is a square in \mathbb{F}_q , as required.
- If t is odd and $q \equiv 3 \mod 4$ then \mathbb{F}_q contains a primitive root of unity of order 2t but does not contain a primitive root of unity of order 4t, and so, a is a non-square in \mathbb{F}_q . However, -a is a primitive root of unity of order t, and so, it is necessarily a square in \mathbb{F}_q , as required.

In conclusion, $a = c^2$ or $-a = c^2$ for some $c \in \mathbb{F}_q$ if and only if either t is odd and divides q - 1 or t is even and 4t divides q - 1.

If t divides (q+1)/2 then $\gamma = a + a^q$ or $\gamma = -(a + a^q)$, for some primitive root of unity a of order 2t in \mathbb{F}_{q^2} (see Proposition 8). Hence,

$$\{2+\gamma, 2-\gamma\} = \{a+2+a^q, (-a)+2+(-a)^q\}.$$

Therefore, $2 + \gamma$ or $2 - \gamma$ is a square in \mathbb{F}_q if and only if $a = c^2$ or $-a = c^2$ for some $c \in \mathbb{F}_{q^2}$ satisfying $c^{q+1} = 1$.

Now, one the following necessarily holds:

- If t is even then -a is also a primitive root of unity of order 2t. Hence, $a = c^2$ for some $c \in \mathbb{F}_{q^2}$ satisfying $c^{q+1} = 1$ if and only if $-a = b^2$ for some $b \in \mathbb{F}_{q^2}$ satisfying $b^{q+1} = 1$. This is equivalent to the condition that 4t divides q + 1.
- If t is odd and $q \equiv 3 \mod 4$ then \mathbb{F}_{q^2} contains a primitive root of unity b of order 4t satisfying $b^{q+1} = 1$. Hence, $a = c^2$ for some $c \in \mathbb{F}_{q^2}$ satisfying $c^{q+1} = 1$, as required.
- If t is odd and $q \equiv 1 \mod 4$ then \mathbb{F}_{q^2} does not contain a primitive root of unity c of order 4t satisfying $c^{q+1} = 1$. However, in this case, $-a = b^2$ for some $b \in \mathbb{F}_{q^2}$ satisfying $b^{q+1} = 1$, as required.

In conclusion, $a = c^2$ or $-a = c^2$ for some $c \in \mathbb{F}_{q^2}$ satisfying $c^{q+1} = 1$ if and only if either t is odd and divides q + 1 or t is even and 4t divides q + 1.

2.4. Number of elements.

Definition 4. Let q be a prime power and let $\alpha \in \mathbb{F}_q$.

If $\alpha = \pm 2$ we say that α is a *unipotent* trace. If α is a trace of a semisimple split (respectively, non-split) matrix in $SL_2(q)$ we say that α is *split* (respectively, *non-split*).

Assume that q is odd and $\alpha \neq \pm 2$. If at least one of $2 + \alpha$, $2 - \alpha$ is a square in \mathbb{F}_q we say that α is a good trace. If none of $2 + \alpha$, $2 - \alpha$ is a square in \mathbb{F}_q we say that α is a bad trace.

Lemma 10.

- (1) If q is even then \mathbb{F}_q contains one unipotent, (q-2)/2 split and q/2 non-split traces.
- (2) If q is odd then \mathbb{F}_q contains 2 unipotent, (q-3)/2 split and (q-1)/2 non-split traces.
- (3) If $q \equiv 1 \mod 4$ then \mathbb{F}_q contains (q-1)/4 bad traces.
- (4) If $q \equiv 3 \mod 4$ then \mathbb{F}_q contains (q+1)/4 bad traces.

Proof. (1) and (2) are due to [11, Lemma 2].

(3) If $q \equiv 1 \mod 4$ then (q-1)/2 is even, and so all the bad traces are necessarily split. Hence, there are $\frac{1}{2}(\frac{1}{2}(q-3)+1)$ bad traces.

(4) If $q \equiv 3 \mod 4$ then (q+1)/2 is even, and so all the bad traces are non-split. Hence, there are $\frac{1}{2}(\frac{1}{2}(q-1)+1)$ bad traces.

Now, the following corollaries easily follow from Table 2, Proposition 9 and Lemma 10.

Corollary 11. If q is even then $G = PSL_2(q)$ contains:

- $q^2 1$ unipotent elements,
- $\frac{1}{2}q(q+1)(q-2)$ semisimple split elements,
- $\frac{1}{2}q^2(q-1)$ semisimple non-split elements.

Corollary 12. If q is even then $G = PSL_2(q)$ contains:

- $q^2 1$ unipotent elements,
- ¹/₄q(q + 1)(q 3) semisimple split elements,
 ¹/₄q(q 1)² semisimple non-split elements.
- $\frac{1}{8}q(q^2-1) = \frac{|G|}{4}$ semisimple elements whose order is not q-good.

2.5. Unipotent elements when q is odd. Consider the following matrices in $G_0 = SL_2(q)$ (where q is odd):

$$U_{1} = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \quad U_{-1} = \begin{pmatrix} -1 & 1 \\ 0 & -1 \end{pmatrix} = -U_{1}^{-1},$$
$$U_{1}' = XU_{1}X^{-1} = \begin{pmatrix} 1 & x^{2} \\ 0 & 1 \end{pmatrix} \in G_{0}, \quad U_{-1}' = XU_{-1}X^{-1} = \begin{pmatrix} -1 & x^{2} \\ 0 & -1 \end{pmatrix} \in G_{0}$$

where $x \in \mathbb{F}_{q^2} \setminus \mathbb{F}_q$ satisfies that $x^2 \in \mathbb{F}_q$ and $X = \begin{pmatrix} x & 0 \\ 0 & x^{-1} \end{pmatrix} \in \mathrm{SL}_2(q^2).$

Proposition 13. Let $G_0 = SL_2(q)$ when q is odd. Then, for any $A \in G_0$, $XAX^{-1} \in G_0$. Moreover,

- If A ≠ I and tr(A) = 2 then A is G₀-conjugate to either U₁ or U'₁.
 If A ≠ -I and tr(A) = -2 then A is G₀-conjugate to either U₋₁ or U'₋₁.

Proof. Indeed, if $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in G_0$ then $XAX^{-1} = \begin{pmatrix} a & bx^2 \\ cx^{-2} & d \end{pmatrix} \in G_0$. Moreover, U_1 is not G_0 -conjugate to $U'_1 = X U_1 X^{-1}$, since x^2 is not a square of some element in \mathbb{F}_q . Hence, any $I \neq A \in G_0$ with tr(A) = 2 is G_0 -conjugate to either U_1 or U'_1 (see Table 2).

Corollary 14. Let $G = PSL_2(q)$ when q is odd. Then, for any $\overline{A} \in G$, $\overline{X}\overline{A}\overline{X}^{-1} \in G$. If moreover, \overline{A} is unipotent then it is G-conjugate to either \overline{U}_1 or $\overline{U'}_1 = \overline{X}\overline{U}_1\overline{X}^{-1}$. In addition,

- If $q \equiv 1 \mod 4$ then \overline{U}_1 and \overline{U}_{-1} are G-conjugate.
- If $q \equiv 3 \mod 4$ then \overline{U}_1 and $\overline{U'}_{-1}$ are G-conjugate.

3. Proof of the main results

Proof of Theorem 1. The proof follows from the following Proposition 15, Proposition 17, Proposition 18, Remark 19, Proposition 32 and Remark 33.

Proof of Theorem 2. The proof follows from the following Proposition 16, Proposition 22, Proposition 23, Proposition 24 and Proposition 28.

Proof of Theorem 3. The proof follows from the following Proposition 16, Proposition 21, Proposition 22, Proposition 24, Proposition 25, Corollary 27, Corollary 31 and Proposition 32.

Proof of Corollary 4. The proof follows from Theorem 2, Theorem 3, Table 2 and §2.4.

- (1) Let $G = PSL_2(q)$ when q is even, then $|G| = q(q^2 1)$.
 - If \mathcal{C} is a conjugacy class of a semisimple split element then

$$C| = q(q+1) \text{ and } |C^2| = q(q^2-1).$$

• If \mathcal{C} is a conjugacy class of a semisimple non-split element then

$$\mathcal{C}| = q(q-1) \text{ and } |\mathcal{C}^2| = q(q^2-1) - (q^2-1) = (q-1)(q^2-1).$$

• If ${\mathcal C}$ is a conjugacy class of a unipotent element then

$$|\mathcal{C}| = q^2 - 1 \text{ and } |\mathcal{C}^2| = q(q^2 - 1).$$

Now let $G = PSL_2(q)$ when q is odd, then $|G| = q(q^2 - 1)/2$.

(2) If \mathcal{C} is a conjugacy class of semisimple element whose order is greater than 2 then

$$|\mathcal{C}| = q(q \pm 1) \text{ and } |\mathcal{C}^2| = q(q^2 - 1)/2$$

- (3) If \mathcal{C} is a conjugacy class of semisimple element x of order 2 then
 - If $q \equiv 1 \mod 4$ then x is split and so

$$C| = q(q+1)/2 \text{ and } |C^2| = q(q^2-1)/2.$$

• If $q \equiv 3 \mod 4$ then x is non-split and so

$$\mathcal{C}| = q(q-1)/2 \ and \ |\mathcal{C}^2| = q(q^2-1)/2 - (q^2-1) = (q-2)(q^2-1)/2.$$

(4) If C is a conjugacy class of a unipotent element, then by Corollary 12,

$$\begin{aligned} |\mathcal{C}| &= (q^2 - 1)/2 \ and \ |\mathcal{C}^2| = q(q^2 - 1)/2 - q(q^2 - 1)/8 - \epsilon = 3q(q^2 - 1)/8 - \epsilon, \\ \text{where } \epsilon &= \begin{cases} 1 & if \ q \equiv 1 \mod 4 \\ 0 & if \ q \equiv 3 \mod 4. \end{cases} \end{aligned}$$

3.1. Generation properties of C.

Proposition 15. Let $G = PSL_2(q)$ when q > 3 and let C be a conjugacy class of a semisimple element s of order greater than 2. Then C contains two elements x, y that generate G.

Proof. Let $s \in G$ be a semisimple element and let C be the conjugacy class of s. Let $S \in G_0 = \operatorname{SL}_2(q)$ be the preimage of s. Denote the order of s by n and $\alpha = \operatorname{tr}(S)$, thus $\alpha \notin \{0, \pm 2\}$. Recall that $d = \operatorname{gcd}(2, q - 1)$.

If $q \neq 5,7$ take some $\gamma \in \mathcal{T}_q((q+1)/d)$. If (α, α, γ) is singular then $(2-\gamma)(\alpha^2 - \gamma - 2) = 0$ and so $\gamma = \alpha^2 - 2$. Thus, we may replace γ by $-\gamma$ (or by another $\gamma \in \mathcal{T}_q((q+1)/d))$ to obtain a non-singular triple (α, α, γ) . When q = 9 one can moreover make sure that (α, α, γ) does not correspond to a triple of matrices (A, B, C) with ABC = I satisfying $\langle \bar{A}, \bar{B} \rangle \cong A_5$ (see [10]). When q = 5 or 7 take $\gamma = -2$, and then the triple $(\alpha, \alpha, -2)$ is also not singular.

By Theorem 6, there exists a triple of matrices (A, B, C) such that ABC = I, $tr(A) = tr(B) = \alpha$ and $tr(C) = \gamma$. Moreover, by Theorem 7, $\langle \bar{A}, \bar{B} \rangle$ is not a structural subgroup of G. By considering the possible subgroups of G detailed in Table 3 we see that the only non-structural subgroup containing \bar{C} is G itself, and hence $\langle \bar{A}, \bar{B} \rangle = G$.

Proposition 16. Let $G = PSL_2(q)$ when q > 3.

- If q is odd then any non-trivial element in G can be written as a product of two G-conjugate semisimple elements that generate G;
- If q is even then only a semisimple element in G can be written as a product of two G-conjugate (semisimple) elements that generate G.

Proof. Assume that q is even. Let $Z \in G$ and denote $\gamma = \operatorname{tr}(Z)$. Assume that Z = XY where X is G-conjugate to Y, and let $\alpha = \operatorname{tr}(X) = \operatorname{tr}(Y)$. If Z is unipotent then $\gamma = 0$. Since $(\alpha, \alpha, 0)$ is singular then by Theorem 7 the subgroup $\langle X, Y \rangle$ is a structural subgroup of G, and so X and Y cannot generate G. If Z is semisimple then $\gamma \neq 0$, and we can choose some $\alpha \in \mathcal{T}_q(q+1)$ such that $\gamma \neq \alpha^2$ and then (α, α, γ) is not singular.

Assume that q is odd. Let $1 \neq z \in G$ and let $Z \in G_0 = \operatorname{SL}_2(q)$ be the preimage of z. Denote $\gamma = \operatorname{tr}(Z)$. If $q \neq 5,7$ take some $\alpha \in \mathcal{T}_q((q+1)/2)$. If (α, α, γ) is singular then $(2-\gamma)(\alpha^2 - \gamma - 2) = 0$ and so $\gamma = -2$ or $\gamma = \alpha^2 - 2$. Thus, we may replace γ by $-\gamma$ to obtain a non-singular triple (α, α, γ) . When q = 9 one can moreover make sure that (α, α, γ) does not correspond to a triple of matrices (A, B, C) with ABC = I satisfying $\langle \overline{A}, \overline{B} \rangle \cong A_5$ (see [10]). When q = 5 or 7 take $\alpha = -2$ (and $\gamma \neq 2$) and then the triple $(-2, -2, \gamma)$ is also not singular.

By Theorem 6, there exists a triple of matrices (A, B, C) such that ABC = I, $tr(A) = tr(B) = \alpha$ and $tr(C) = \gamma$. Moreover, by Theorem 7, $\langle \bar{A}, \bar{B} \rangle$ is not a structural subgroup of G. By considering the possible subgroups of G detailed in Table 3 we see that the only non-structural subgroup containing \bar{A} is G itself, and hence $\langle \bar{A}, \bar{B} \rangle = G$.

Proposition 17. Let $G = PSL_2(q)$ when q > 3 and let C be a conjugacy class of a semisimple element s. Then C contains three elements x, y, z such that xyz = 1 and $\langle x, y \rangle = G$ if and only if the order of s is q-minimal and greater than 3.

Proof. Let $s \in G$ be a semisimple element and let C be the conjugacy class of s. Let $S \in G_0 = \operatorname{SL}_2(q)$ be the preimage of s. Denote the order of s by n and $\alpha = \operatorname{tr}(S)$.

If x, y, z are three elements of order 2 (respectively, 3) such that xyz = 1, then any finite group generated by x and y is necessarily abelian (respectively, solvable) and so x and y cannot generate G (see Remark 19).

If n is not q-minimal then α belongs to some proper subfield \mathbb{F}_{q_1} of \mathbb{F}_q . In this case, for any three elements $x, y, z \in \mathcal{C}$ satisfying xyz = 1, the subgroup $\langle x, y \rangle$ is necessarily contained in a proper subgroup of G isomorphic to $\mathrm{PSL}_2(q_1)$ (see [11] and [9]).

If n > 3 is q-minimal then $\alpha \notin \{0, \pm 1, \pm 2\}$ and hence $(\alpha - 2)^2(\alpha + 1) \neq 0$ implying that the triple (α, α, α) is non-singular. If q > 5 and n = 5 one can moreover make sure that the triple (α, α, α) does not correspond to a triple of matrices (A, B, C) with ABC = Isatisfying $\langle \overline{A}, \overline{B} \rangle \cong A_5$ (see [10]).

Therefore, by Theorems 6 and 7, there exists a triple of matrices (A, B, C) such that ABC = I, $tr(A) = tr(B) = tr(C) = \alpha$ and $\langle \bar{A}, \bar{B} \rangle$ is a subfield subgroup of G. By the q-minimality of n, s does not belong to any subgroup isomorphic to $PSL_2(q_1)$ for some proper subfield $\mathbb{F}_{q_1} \subset \mathbb{F}_q$. In addition, if $\langle x, y \rangle = PGL_2(q_1)$ and xyz = 1 then exactly two of x, y, z belong to $PGL_2(q_1) \setminus PSL_2(q_1)$ and the third one belongs to $PSL_2(q_1)$, so x, y, z cannot all have the same order n. Therefore, $\langle \bar{A}, \bar{B} \rangle$ cannot generate a subgroup isomorphic to $PGL_2(q_1)$, hence $\langle \bar{A}, \bar{B} \rangle = G$ as needed.

Proposition 18. Let $G = PSL_2(q)$ where q is a prime power. Then there exist three Gconjugate unipotent elements A, B and C such that ABC = 1 and $\langle A, B \rangle = G$ if and only q > 3 is prime.

Proof. If $A, B, C \in G$ are three elements of prime order p such that ABC = 1 then $\langle A, B \rangle$ is contained in a subgroup isomorphic to $PSL_2(p)$, implying that necessarily q = p > 3 is prime (see [11] and [9]).

Now, choose some $a \in \mathbb{F}_p \setminus \{0, \pm 2\}$ and consider the following matrices in $SL_2(p)$:

$$M = \begin{pmatrix} a+1 & -a/2 - 1 \\ 2 & -1 \end{pmatrix}, \quad K = \begin{pmatrix} a & -1/2 \\ 2 & 0 \end{pmatrix}.$$

Let $A = U_{-1} \in G_0$ be as in §2.5, $B = MU_{-1}M^{-1}$ and $C = KU_{-1}K^{-1}$, then ABC = I.

As $\operatorname{tr}(U_{-1}) = -2$ and (-2, -2, -2) is a *non-singular* triple, then by Theorem 7, the subgroup generated by the images of A and B is not a structural subgroup of G. By considering the possible subgroups of G detailed in Table 3 we conclude that $\langle \overline{A}, \overline{B} \rangle = G$, as needed

Remark 19. Recall that any finite group generated by two non-commuting involutions is dihedral. In addition, the group presented by $\langle x, y | x^3 = y^3 = (xy)^3 \rangle$ is an infinite solvable group (see e.g. [1]).

3.2. Basic properties of C^2 .

Lemma 20. Let $A, B \in G_0 = \operatorname{SL}_2(q)$ be two non-central matrices such that $\operatorname{tr}(A) = \operatorname{tr}(B)$ or $\operatorname{tr}(A) = -\operatorname{tr}(B)$. Let \mathcal{C} be some conjugacy class in $G = \operatorname{PSL}_2(q)$. If $\overline{A} \in \mathcal{C}^2$ then also $\overline{B} \in \mathcal{C}^2$.

Proof. If $\overline{A} \in \mathcal{C}^2$ then $\overline{A} = z(gzg^{-1})$ for some $z \in \mathcal{C}, g \in G$.

If q is even, or if q is odd and $tr(A) \neq \pm 2$, then necessarily $\overline{B} = h\overline{A}h^{-1}$ for some $h \in G$ (see Table 2), and so $\overline{B} = (hzh^{-1})(hgzg^{-1}h^{-1}) \in \mathcal{C}^2$.

Assume that q is odd, z is semisimple and \overline{A} is unipotent. Let $Z \in G_0$ be the preimage of z. Then by Corollary 14, either $\overline{B} = h\overline{A}h^{-1}$ or $\overline{B} = h\overline{X}A\overline{X}^{-1}h^{-1}$ for some $h \in G$ and a matrix $X \in SL_2(q^2)$ as in §2.5. Moreover by Proposition 13, $XZX^{-1} \in G_0$, and since $\operatorname{tr}(XZX^{-1}) = \operatorname{tr}(Z)$ then $\overline{X}z\overline{X}^{-1} \in \mathcal{C}$ (see Table 2). Similarly, $\overline{X}gzg^{-1}\overline{X}^{-1} \in \mathcal{C}$.

Hence, either

$$\bar{B} = h\bar{A}h^{-1} = (hzh^{-1})(hgzg^{-1}h^{-1}) \in \mathcal{C}^2$$

or

$$\bar{B} = h\bar{X}\bar{A}\bar{X}^{-1}h^{-1} = (h\bar{X}z\bar{X}^{-1}h^{-1})(h\bar{X}gzg^{-1}\bar{X}^{-1}h^{-1}) \in \mathcal{C}^2$$

If q is odd and z is unipotent, then the following Proposition 21 shows that C^2 contains all the unipotent elements in G, as needed.

Proposition 21. Let $G = PSL_2(q)$ when q is odd and let C be the G-conjugacy class of a unipotent element x. Then C^2 contains all the unipotent elements in G. Moreover, $1 \in C^2$ if and only if $q \equiv 1 \mod 4$.

Proof. For q = 5 this claim can be easily verified. When q > 5 there exist elements $a, b \in \mathbb{F}_q$ such that a, b and a + 1 are squares in \mathbb{F}_q but b + 1 is a non-square in \mathbb{F}_q . Let U_1 be as in §2.5, let $A = \begin{pmatrix} 1 & a \\ 0 & 1 \end{pmatrix}$, $B = \begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix}$, then $U_1A = \begin{pmatrix} 1 & a+1 \\ 0 & 1 \end{pmatrix}$ and $U_1B = \begin{pmatrix} 1 & b+1 \\ 0 & 1 \end{pmatrix}$. Moreover, A, B and U_1A are conjugate to U_1 in $\mathrm{SL}_2(q)$, while U_1B is conjugate to U'_1 in $\mathrm{SL}_2(q)$ (see Proposition 13). By Corollary 14, any unipotent element in G is G-conjugate to either \overline{U}_1 or to \overline{U}'_1 , as needed.

In addition, by Corollary 14, \overline{U}_1 is *G*-conjugate to \overline{U}_1^{-1} if and if $q \equiv 1 \mod 4$.

Proposition 22. Let $G = PSL_2(q)$ and let C be the G-conjugacy class of a semisimple element x. Then C^2 contains all the semisimple elements in G. Moreover $1 \in C^2$.

Proof. Let $z \in G$ be some semisimple element. Let $X, Z \in SL_2(q)$ be the pre-images of x, z respectively. Denote $\alpha = tr(X), \gamma = tr(Z)$. By Theorem 6, there exist matrices $X', Y', Z' \in SL_2(q)$ such that $tr(X') = tr(Y') = \alpha$ and $tr(Z') = \gamma$. By Table 2, $\bar{X}', \bar{Y}' \in C$ and so $\bar{Z}' \in C^2$. Hence by Lemma 20, $z \in C^2$ as needed.

Moreover, since $\operatorname{tr}(X) = \operatorname{tr}(X^{-1})$ then according to Table 2, $x^{-1} \in \mathcal{C}$ as well, and so $xx^{-1} = 1 \in \mathcal{C}^2$.

Note that the previous proposition is a specific case of a more general result of Gow [7].

Proposition 23. Let $G = PSL_2(q)$ when q is even and let C be the G-conjugacy class of a unipotent element x. Then $C^2 = G$.

Proof. Let $z \in G$ be any semisimple element and denote $\gamma = \operatorname{tr}(z)$. By Theorem 6, there exist matrices $x', y', z' \in G$ such that $\operatorname{tr}(x') = \operatorname{tr}(y') = 0$ and $\operatorname{tr}(z') = \gamma$. Thus $x', y' \in \mathcal{C}$ and so $z' \in \mathcal{C}^2$. Hence by Lemma 20, $z \in \mathcal{C}^2$ as needed.

Moreover, since x has order 2 then $x^2 = 1 \in \mathcal{C}^2$. Without loss of generality we may assume that $x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ and take $y = \begin{pmatrix} a+1 & a \\ a & a+1 \end{pmatrix} \in \mathrm{SL}_2(q)$ (where $a \neq 0, 1$). Then $xy = yx = \begin{pmatrix} a & a+1 \\ a+1 & a \end{pmatrix}$. Since $\mathrm{tr}(x) = \mathrm{tr}(y) = \mathrm{tr}(xy) = 0$ then x, y, xy are unipotents. Hence by Lemma 20, \mathcal{C}^2 contains all the unipotent elements.

3.3. Unipotent elements contained in C^2 .

Proposition 24. Let $G = PSL_2(q)$ and let C be the G-conjugacy class of a semisimple split element x. Then C^2 contains all the unipotent elements in G.

Proof. By Table 2, we may assume x is an image of a matrix $X = \begin{pmatrix} a & 1 \\ 0 & a^{-1} \end{pmatrix} \in SL_2(q)$ (where $a \neq 0, \pm 1$). Let $Y = \begin{pmatrix} a^{-1} & a^{-1}(1-a) \\ 0 & a \end{pmatrix}$, then its image in $G = PSL_2(q)$ is G-conjugate to x, and $XY = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$. Hence C^2 contains a unipotent element, and the result now follows from Lemma 20.

Proposition 25. Let $G = PSL_2(q)$ when q is odd and let C be the G-conjugacy class of a semisimple non-split element x of order greater than 2. Then C^2 contains all the unipotent elements in G.

Proof. By Table 2, we may assume that x is an image of a matrix $X = \begin{pmatrix} a & -1 \\ 1 & 0 \end{pmatrix} \in SL_2(q)$

(where $a \neq 0$). Let $Y = \begin{pmatrix} 0 & 1 \\ -1 & -a \end{pmatrix}$, then its image in $G = \text{PSL}_2(q)$ is G-conjugate to x, and $XY = \begin{pmatrix} 1 & 2a \\ 0 & 1 \end{pmatrix}$. Hence \mathcal{C}^2 contains a unipotent element, and the result now follows from Lemma 20.

Proposition 26. Let $G = PSL_2(q)$ when q is odd and let C be the G-conjugacy class of a semisimple non-split element x of order 2. Then C^2 does not contain any unipotent element of G.

Proof. Assume that C^2 contains a unipotent element $z \in G$. Then z = xy for some non-split elements x, y of order 2. Let $X, Y, Z \in SL_2(q)$ be the pre-images of x, y, z respectively. Then tr(X) = tr(Y) = 0 and $tr(Z) = \pm 2$. Observe that $(0, 0, \pm 2)$ is a *singular* triple. Therefore, by Theorem 7, the subgroup $H = \langle x, y \rangle$ is a *structural* subgroup of G. By observing the possible subgroups of G detailed in Table 3, H cannot be a subgroup of the Borel subgroup since it contains a non-split element, and hence cannot contain the unipotent element z, yielding a contradiction.

Corollary 27. Let $G = PSL_2(q)$ when q is odd and let C be the G-conjugacy class of a semisimple element x of order 2. Then

- If $q \equiv 1 \mod 4$ then C^2 contains all the unipotent elements in G.
- If $q \equiv 3 \mod 4$ then C^2 does not contain any unipotent element of G.

Proof. If $q \equiv 1 \mod 4$ then an element of order 2 is split and the result follows from Proposition 24. If $q \equiv 3 \mod 4$ then an element of order 2 is non-split and the result follows from Proposition 26.

Proposition 28. Let $G = PSL_2(q)$ when q is even and let C be the G-conjugacy class of a semisimple non-split element x. Then C^2 does not contain any unipotent element of G.

Proof. Assume that C^2 contains a unipotent element $z \in G$. Then z = xy for some non-split matrices x, y. Then $tr(x) = tr(y) = \alpha \neq 0$ and tr(z) = 0. Observe that $(\alpha, \alpha, 0)$ is a singular

triple. Therefore, by Theorem 7, the subgroup $H = \langle x, y \rangle$ is a *structural* subgroup of G. By observing the possible subgroups of G detailed in Table 3, H cannot be a subgroup of the Borel subgroup since it contains a non-split element, and hence cannot contain the unipotent element z, yielding a contradiction.

3.4. Unipotent conjugacy classes C when q is odd.

Proposition 29. Let $G_0 = SL_2(q)$ when $q = p^e$ is odd, and let $A, B \in G_0$, $A, B \neq \pm I$ satisfy $tr(A), tr(B) \in \{\pm 2\}$. Denote C = AB and $\gamma = tr(C)$.

- (1) If tr(A) = tr(B), then A is G_0 -conjugate to B if and only if 2γ is a square in \mathbb{F}_q .
- (2) If tr(A) = -tr(B), then A is G₀-conjugate to -B if and only if $2 + \gamma$ is a square in \mathbb{F}_q .

Proof. Without loss of generality we may assume that $A = U_1$.

(1) If
$$B = MU_1M^{-1}$$
 for some matrix $M = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in G_0$, then
 $\gamma = \operatorname{tr}(C) = \operatorname{tr}(AB) = \operatorname{tr}(U_1MU_1M^{-1}) = \operatorname{tr}\begin{pmatrix} 1 - ac - c^2 & 1 + ac + a^2 \\ -c^2 & 1 + ac \end{pmatrix} = 2 - c^2$,

and so $2 - \gamma$ is a square in \mathbb{F}_q .

If $B = MU'_1 M^{-1}$ for some matrix $M \in G_0$, then

$$\gamma = \operatorname{tr}(C) = \operatorname{tr}(AB) = \operatorname{tr}(U_1 M U_1' M^{-1}) = 2 - x^2 c^2,$$

and since $x \in \mathbb{F}_{q^2} \setminus \mathbb{F}_q$ then $2 - \gamma$ is a non-square in \mathbb{F}_q .

(2) Similarly, if $B = M(-U_1)M^{-1}$ for some matrix $M \in G_0$, then

$$\gamma = \operatorname{tr}(C) = \operatorname{tr}(AB) = \operatorname{tr}(U_1M(-U_1)M^{-1}) = -2 + c^2,$$

and so $2 + \gamma$ is a square in \mathbb{F}_q .

If $B = M(-U'_1)M^{-1}$ for some matrix $M \in G_0$, then

$$\gamma = \operatorname{tr}(C) = \operatorname{tr}(AB) = \operatorname{tr}(U_1M(-U_1')M^{-1}) = -2 + x^2c^2,$$

and so $2 + \gamma$ is a non-square in \mathbb{F}_q .

Therefore, in order to decide whether C = AB, where $A, B \in G$ are G-conjugate unipotent elements and C is of order $t \neq p$, one needs to determine whether for $\gamma \in \mathcal{T}_q(t)$, $2 - \gamma$ or $2 + \gamma$ is a square in \mathbb{F}_q . In Proposition 9 we showed that this is equivalent to decide whether t is a q-good order or not (see Definition 2).

Corollary 30. Let $G = PSL_2(q)$ when $q = p^e$ is odd. Let $A, B \in G$ be two elements of order p and assume that the order of C = AB is $t \neq p$. Then A is G-conjugate to B if and only if t is a q-good order.

Proof. Let $A, B \in G_0 = \operatorname{SL}_2(q)$ and C = AB and assume that their images $\overline{A}, \overline{B}, \overline{C} \in G = \operatorname{PSL}_2(q)$ have respective orders $(p, p, t), t \neq p$. Denote $\gamma = \operatorname{tr}(C)$. Then \overline{A} and \overline{B} are unipotent if and only if $A, B \neq \pm I$ and $\operatorname{tr}(A), \operatorname{tr}(B) \in \{\pm 2\}$. Moreover, \overline{A} and \overline{B} are G-conjugate if and only if either $\operatorname{tr}(A) = \operatorname{tr}(B)$ and A and B are G_0 -conjugate or $\operatorname{tr}(A) = -\operatorname{tr}(B)$ and A and -B are G_0 -conjugate. From Proposition 29 we deduce that \overline{A}

and \overline{B} are *G*-conjugate if and only if $2 - \gamma$ or $2 + \gamma$ is a square in \mathbb{F}_q . By Proposition 9, the latter is equivalent to *t* being a *q*-good order.

Corollary 31. Let $G = PSL_2(q)$ when q is odd and let C be the G-conjugacy class of a unipotent element x. Let z be a semisimple element in G. Then C^2 contains z if and only if the order of z is q-good.

Proof. Follows from Corollary 30 and Lemma 20.

Proposition 32. Assume that $q = p^e$ where p is odd and $5 \le q \ne 9$ and let $G = PSL_2(q)$. Then there exist two G-conjugate unipotent elements A and B such that $\langle A, B \rangle = G$.

Moreover, if C = AB is semisimple then the order of C is q-minimal and q-good.

Proof. Let $(A, B, C) \in G^3$ be a triple of respective orders (p, p, t) such that $t \neq p$. If t is not q-good then by Corollary 30, A and B are not G-conjugate. If t is not q-minimal then the subgroup $\langle A, B \rangle$ is contained in a subfield subgroup isomorphic to $PSL_2(q_1)$ for some proper subfield $\mathbb{F}_{q_1} \subset \mathbb{F}_q$ (see [11] and [9]).

Now, let t be a q-minimal order which is also a q-good order. By Theorem 6, there exist a triple $(A, B, C) \in G^3$ of respective orders (p, p, t) such that ABC = 1. By Corollary 30, A is G-conjugate to B.

Let $\gamma \in \mathcal{T}_q(t)$, as $t \neq p$ then $\gamma \neq \pm 2$ and so $(\gamma \pm 2)^2 \neq 0$ implying that $(2, 2, \gamma)$ is non-singular, hence by Theorem 7, $\langle A, B \rangle$ is not a structural subgroup of G. Since t is q-minimal, then $\langle A, B \rangle$ is not isomorphic neither to $\mathrm{PSL}_2(q_1)$ nor to $\mathrm{PGL}_2(q_1)$, for some proper subfield \mathbb{F}_{q_1} of \mathbb{F}_q . Moreover, if $5 < q \neq 9$ then either p > 5; or p = 5 and e > 1implying that $t \neq 2, 3, 5$; or p = 3 and e > 2 implying that t > 5 (see Table 1). Therefore, $\langle A, B \rangle$ cannot be a small subgroup, hence $\langle A, B \rangle = G$.

Remark 33. Let $G = PSL_2(9) \cong A_6$, let A, B be two unipotent elements (of order 3) and $C = (AB)^{-1}$. Denote the order of C by t, then $t \in \{2, 3, 4, 5\}$.

- Clearly if t = 2 or t = 3 then $\langle A, B \rangle \neq G$ (see Table 1).
- t = 4 is not 9-good and so A and B are not G-conjugate, by Corollary 30.
- t = 5 is 9-good, however if A is G-conjugate to B, then one can verify that $\langle A, B \rangle \cong A_5$ is a small subgroup of G (see e.g. [6, §2, Theorem 8.4]).

Acknowledgement. The author would like to thank Gili Schul for useful discussions.

References

- H.S.M. Coxeter, W.O.J. Moser, Generators and relations for discrete groups, Ergebnisse der Mathematik und Ihrer Grenzgebrete, New Series, no. 14. Springer, Berlin-Gottingen-Heidelberg, 1957.
- [2] L.E. Dickson, *Linear Groups with an Exposition of the Galois Field Theory*, Teubner (1901).
- [3] L. Dornhoff, Group Representation Theory, Part A, Marcel Dekker, 1971.
- [4] E.W. Ellers, N. Gordeev, On the conjectures of J. Thompson and O. Ore, Trans. Amer. Math. Soc. 350 (1998), 3657-3671
- [5] S. Garion, On Beauville Structures for PSL(2, q), arXiv:1003.2792.
- [6] D. Gorenstein, *Finite groups*, Chelsea Publishing Co., New York, 1980.
- [7] R. Gow, Commutators in finite simple groups of Lie type, Bull. London Math. Soc. 32 (2000), no. 3, 311–315.
- [8] R. Guralnick, G. Malle, Products of conjugacy classes and fixed point spaces, J. Amer. Math. Soc. 25 (2012), 77–121.
- U. Langer, G. Rosenberger, Erzeugende endlicher projektiver linearer Gruppen, Results Math. 15 (1989), no. 1-2, 119–148.

- [10] F. Levin, G. Rosenberger, Generators of finite projective linear groups. II., Results Math. 17 (1990), no. 1-2, 120–127.
- [11] A.M. Macbeath, Generators of the linear fractional groups, Number Theory (Proc. Sympos. Pure Math., Vol. XII, Houston, Tex., 1967), Amer. Math. Soc., Providence, R.I. (1969), 14–32.
- [12] C. Marion, Triangle groups and $PSL_2(q)$, J. Group Theory 12 (2009), 689–708.
- [13] G. Schul, Expansion in finite simple groups.
- [14] M. Suzuki, Group Theory I, Springer, Berlin (1982).

FACHBEREICH MATHEMATIK UND INFORMATIK, UNIVERSITÄT MÜNSTER, EINSTEINSTRASSE 62, D-48149 MÜNSTER, GERMANY

 $E\text{-}mail\ address:\ \texttt{shelly.garion} @uni-muenster.de$