ON THE q-ANALOGUE OF LAPLACE TRANSFORM

WON SANG CHUNG¹ AND TAEKYUN KIM²

ABSTRACT. In this paper, we consider a q-analogue of Laplace transform and we investigate some properties of q-Laplace transform. From our investigation, we derive some interesting formulae related to q-Laplace transform.

1. Introduction

For $q \in [0,1]$, we define the q-Shifted factorials by $(a:q)_0 = 1$, $(a:q)_n = \prod_{i=0}^{n-1} (1-aq^i)$, and $(a:q)_\infty = \lim_{n\to\infty} (a:q)_n = \prod_{i=0}^\infty (1-aq^i)$. If x is classical object, such as a complex number, its q-version is defined by $[x]_q = \frac{1-q^x}{1-q}$. As is well-known, the q-exponential functions are given by

$$E_q(-z) = ((1-q)z:q))_{\infty} = \sum_{n=0}^{\infty} \frac{(-1)^n q^{\binom{n}{2}}}{[n]_q!} z^n,$$
 (1.1)

and

$$e_q(z) = \frac{1}{((1-q)z:q)_{\infty}} = \sum_{n=0}^{\infty} \frac{1}{[n]_q!} z^n$$
, (see [1-8]), (1.2)

where $[n]_q! = [n]_q[n-1]_q \cdots [2]_q[1]_q$.

For $t, x, y \in \mathbb{R}$ and $n \in \mathbb{Z} \geq 0$, the *q-binomial theorem* is also given by

$$(x+y)_q^n = \prod_{j=0}^{n-1} (x+q^j y) = \sum_{k=0}^n \binom{n}{k}_q x^{n-k} q^{\binom{2}{k}} y^k, \tag{1.3}$$

$$(1+x)_q^t = \frac{(x:q)_\infty}{(q^t x:q)_\infty}, \text{ and } \frac{1}{(x-y)_q^n} = \sum_{k=0}^\infty \binom{n+k-1}{k}_q x^{n-k} y^k,$$
 (1.4)

where

$$\binom{n}{k}_q = \frac{[n]_q!}{[n-k]_q![k]_q!} = \frac{[n]_q[n-1]_q \cdots [n-k+1]_q}{[k]_q!}, \text{ (see [2,6])}.$$

From (1.4), we note that

$$(1+x)_a^{s+t} = (1+x)_a^s (1+q^s x)_a^t. (1.5)$$

For $s \in \mathbb{C}$ with s > 0, the gamma function is defined by

$$\Gamma(s) = \int_0^\infty e^{-t} t^{s-1} dt. \tag{1.6}$$

¹⁹⁹¹ Mathematics Subject Classification. 05A40, 05A19.

Key words and phrases. Frobenius-type Eulerian polynomials, Frobenius-type Eulerian numbers.umbral calculus.

By (1.6), we get

$$\Gamma(s+1) = s\Gamma(s), \ \Gamma(n+1) = n!, \ (n \in \mathbb{N}).$$

The Jackson q-derivative is defined by

$$D_q f(x) = \frac{\partial_q f(x)}{\partial_q x} = \frac{f(x) - f(qx)}{(1 - q)x}, \text{ (see [1-8])}.$$
 (1.7)

Note that $\lim_{q\to 1} D_q f(x) = f'(x)$. The definite Jackson q-integral is given by

$$\int_0^x f(t)d_q t = (1 - q) \sum_{a=0}^\infty f(q^a x) x q^a, \text{ (see [3-4])}.$$
 (1.8)

Thus, by (1.7) and (1.8), we get

$$\int_0^x \frac{\partial_q}{\partial_q t} f(t) d_q t = f(x) - f(0). \tag{1.9}$$

The improper q-integral of f is given by

$$\int_0^{\frac{\infty}{a}} f(x)d_q x = (1-q) \sum_{n \in \mathbb{Z}} \frac{q^n}{a} f\left(\frac{q^n}{a}\right). \tag{1.10}$$

Thus, by (1.10), we get

$$\int_{x}^{\infty} f(y)d_{q}y = \int_{0}^{x \cdot \infty} f(y)d_{1}y - \int_{0}^{x} f(y)d_{q}y$$

$$= x(1-q)\sum_{n=0}^{\infty} q^{-n}f(q^{-n}x).$$
(1.11)

Let f be a function defined for $t \geq 0$. Then the integral

$$\mathcal{L}(f(t)) = \int_0^\infty e^{-st} f(t) dt$$
, (see [9])

is said to be the $Laplace\ transform$ of f, provided the integral converges.

In this paper, we consider a q-analogue of Laplace transform, which is called by q-Laplace transform, and we investigate some properties of q-Laplace transform. From our investigation, we derive some interesting formulae related to q-Laplace transform.

2. q-Laplace transforms

Now we consider two types of the q-Laplace transform. The q-Laplace transform of the first kind is defined by

$$F(s) = L_q(f(t)) = \int_0^\infty E_q(-qst)f(t)d_q(t), \ (s > 0).$$
 (2.1)

Then, by (2.1), we get

$$L_q(\alpha f(t) + \beta g(t)) = \alpha L_q(f(t)) + \beta_q L_g(t), \qquad (2.2)$$

where α, β are constants.

The q-extension of gamma function is defined by

$$\Gamma_q(t) = \int_0^\infty x^{t-1} E_q(-qx) d_q x, \ (t > 0), \ (\text{see [2,6]}).$$
 (2.3)

Thus, by (2.3), we get

$$\Gamma_q(t+1) = [t]_q \Gamma_q(t), \ \Gamma_q(n+1) = [n]_q!, \ (\text{see } [2,6]).$$
 (2.4)

From (1.2), we can derive

$$e_q(iz) = \sum_{n=0}^{\infty} \frac{i^n z^n}{[n]_q!} = \sum_{n=0}^{\infty} \frac{(-1)^n}{[2n]_q!} z^{2n} + i \sum_{n=0}^{\infty} \frac{(-1)^n z^{2n+1}}{[2n+1]_q!}. \tag{2.5}$$

By (2.5), we define *q-cosine* and *q-sine function* as follows:

$$\cos_q(z) = \sum_{n=0}^{\infty} \frac{(-1)^n z^{2n}}{[2n]_q!} = \frac{1}{2} \left(e_q(iz) + e_q(-iz) \right), \tag{2.6}$$

and

$$\sin_q(z) = \sum_{n=0}^{\infty} \frac{(-1)^n}{[2n+1]_q!} = \frac{1}{2i} \left(e_q(iz) - e_q(-iz) \right), \tag{2.7}$$

where $i = \sqrt{-1}$. From (2.1) and (2.4), we note that

$$L_q(1) = \frac{1}{s} (s > 0), \ L_q(t) = \frac{1}{s^2} (s > 0), \ L_q(e_q(-3t)) = \frac{1}{s+3} (s > -3),$$

 $L_q(\sin_q 2t) = \frac{2}{s^2+4}, \ L_q(1+5t) = \frac{1}{s} + \frac{5}{s^2} (s > 0), \cdots.$

We state the generalization of some of the preceding examples by mean of the next theorem. From this point on we shall also refrain from stating any restrictions on s; it is understood that s is sufficiently restricted to guarantee the convergence of the appropriate q-Laplace transform.

For $\alpha \in \mathbb{R}$ with $\alpha > -1$, we have

$$L_{q}(t^{\alpha}) = \int_{0}^{\infty} E_{q}(-qst)t^{\alpha}d_{q}t = \frac{1}{s^{\alpha+1}} \int_{0}^{\infty} E_{q}(-qt)t^{\alpha}d_{q}t$$

$$= \frac{1}{s^{\alpha+1}} \Gamma_{q}(\alpha+1).$$
(2.8)

In particular, $\alpha = n \in \mathbb{N}$, by (2.8), we get

$$L_q(t^n) = \frac{1}{s^{n+1}} \Gamma_q(n+1) = \frac{[n]_q!}{s^{n+1}}.$$
 (2.9)

Let us take $\alpha = -\frac{1}{2}$ and $\alpha = \frac{1}{2}$. Then we see that

$$L_q\left(t^{-\frac{1}{2}}\right) = \frac{1}{s^{\frac{1}{2}}}\Gamma_q\left(\frac{1}{2}\right) = \frac{1}{\sqrt{s}}\Gamma_q\left(\frac{1}{2}\right),\tag{2.10}$$

and

$$L_q\left(t^{\frac{1}{2}}\right) = \frac{1}{s^{\frac{3}{2}}}\Gamma_q\left(\frac{3}{2}\right) = \frac{\Gamma_q\left(\frac{1}{2}\right)}{\left[2\right]_{\frac{1}{2}}\sqrt{s^3}}.$$
 (2.11)

By (1.1) and (1.2), we get

$$L_{s}(e_{q}(at)) = \int_{0}^{\infty} E_{q}(-qst)e_{q}(at)d_{q}t$$

$$= \sum_{n=0}^{\infty} \frac{a^{n}}{[n]_{q}!} \int_{0}^{\infty} E_{q}(-qst)t^{n}d_{q}t$$

$$= \sum_{n=0}^{\infty} \frac{a^{n}}{[n]_{q}!} \frac{\Gamma_{q}(n+1)}{s^{n+1}} = \frac{1}{s} \sum_{n=0}^{\infty} \left(\frac{a}{s}\right)^{n}$$

$$= \frac{1}{s} \left(\frac{1}{1 - \frac{a}{s}}\right) = \frac{1}{s - a},$$
(2.12)

and

$$L_{q}(E_{q}(at)) = \int_{0}^{\infty} E_{q}(-qst)E_{q}(at)d_{q}t$$

$$= \sum_{n=0}^{\infty} \frac{(-1)^{n}q^{\binom{n}{2}}a^{n}}{[n]_{q}!} \int_{0}^{\infty} E_{q}(-qst)t^{n}d_{q}t$$

$$= \sum_{n=0}^{\infty} \frac{(-1)^{n}}{[n]_{q}!}q^{\binom{n}{2}}a^{n}\frac{\Gamma_{q}(n+1)}{s^{n+1}} = \sum_{n=0}^{\infty} (-1)^{n}q^{\binom{n}{2}}\frac{a^{n}}{s^{n+1}}.$$
(2.13)

Therefore, by (2.8)-(2.13), we obtain the following theorem.

Theorem 2.1. For $\alpha \in \mathbb{R}$ with $\alpha > -1$, we have

$$L_q(t^{\alpha}) = \frac{1}{s^{n+1}} \Gamma_q(\alpha + 1).$$

In particular, $\alpha = n \in \mathbb{N}$, we get

$$L_q(t^n) = \frac{[n]_q!}{s^{n+1}}.$$

Moreover,

$$L_q\left(t^{-\frac{1}{2}}\right) = \frac{1}{\sqrt{s}}\Gamma_q\left(\frac{1}{2}\right), \ L_q\left(t^{\frac{1}{2}}\right) = \frac{\Gamma_q\left(\frac{1}{2}\right)}{\left[2\right]_{q^{\frac{1}{2}}}\sqrt{s^3}}.$$

and

$$L_s(e_q(at)) = \frac{1}{s-a}, \ L_q(E_q(at)) = \sum_{n=0}^{\infty} (-1)^n q^{\binom{n}{2}} \frac{a^n}{s^{n+1}}.$$

From (2.6) and (2.7), we have

$$L_{q}(\cos_{q} at) = \sum_{n=0}^{\infty} \frac{(-1)^{n} a^{2n}}{[2]_{q}!} \int_{0}^{\infty} E_{q}(-qst) t^{2n} d_{q} t$$

$$= \sum_{n=0}^{\infty} \frac{(-1)^{n} a^{2n}}{[2n]_{q}!} \frac{1}{s^{2n+1}} \Gamma_{q}(2n+1)$$

$$= \frac{1}{s} \sum_{n=0}^{\infty} (-1)^{n} \left(\frac{a}{s}\right)^{2n} = \frac{1}{s} \frac{1}{1 + \left(\frac{a}{s}\right)^{2}} = \frac{s}{s^{2} + a^{2}},$$
(2.14)

and

$$L_{q}(\sin_{q} at) = \sum_{n=0}^{\infty} \frac{(-1)^{n} a^{2n+1}}{[2n+1]_{q}!} \int_{0}^{\infty} E_{q}(-qst) t^{2n+1} d_{q}t$$

$$= \sum_{n=0}^{\infty} \frac{(-1)^{n} a^{2n+1}}{[2n+1]_{q}!} \frac{\Gamma_{q}(2n+2)}{s^{2n+2}}$$

$$= \frac{1}{s} \sum_{n=0}^{\infty} (-1)^{n} \left(\frac{a}{s}\right)^{2n+1} = \frac{1}{s} \frac{\frac{a}{s}}{1 + \left(\frac{a}{s}\right)^{2}} = \frac{a}{s^{2} + a^{2}}.$$
(2.15)

Let us define hyperbolic q-cosine and hyperbolic q-sine functions as follows:

$$\cosh_q t = \frac{e_q(t) + e_q(-t)}{e}, \ \sinh_q t = \frac{e_q(t) - e_q(-t)}{2}.$$
 (2.16)

From (2.1), (2.12) and (2.16), we note that

$$L_q(\cosh_q at) = \frac{1}{2} \left\{ L_q(e_q(at)) + L_q(e_q(-at)) \right\}$$

$$= \frac{1}{2} \left\{ \frac{1}{s-a} + \frac{1}{s+a} \right\} = \frac{s}{s^2 - a^2},$$
(2.17)

and

$$L_q(\sinh_q at) = \frac{1}{2} \left\{ L_q(e_q(at)) - L_q(e_q(-at)) \right\}$$

$$= \frac{1}{2} \left\{ \frac{1}{s-a} - \frac{1}{s+a} \right\} = \frac{a}{s^2 - a^2}.$$
(2.18)

Therefore, by (2.14)-(2.18), we obtain the following theorem.

Theorem 2.2. (Transforms of q-trigonometric function)

$$L_q(\cos_q at) = \frac{s}{s^2 + a^2}, \ L_q(\sin_q at) = \frac{a}{s^2 + a^2}$$

 $L_q(\cosh_q at) = \frac{s}{s^2 - a^2}, \ L_q(\sinh_q at) = \frac{a}{s^2 - a^2}$

Now, we observe that

$$E_{q}(-qst) = \sum_{n=0}^{\infty} \frac{(-1)^{n} q^{\binom{n}{2}}}{[n]_{q}!} q^{n} s^{n} t^{n} = \sum_{n=0}^{\infty} \frac{(-1)^{n} q^{\binom{n}{2}}}{[n]_{q}!} ([n]_{q} (q-1) + 1) s^{n} t^{n}$$

$$= (q-1) \sum_{n=1}^{\infty} \frac{(-1)^{n} q^{\binom{n}{2}}}{[n-1]_{q}!} s^{n} t^{n} + \sum_{n=0}^{\infty} \frac{(-1)^{n} q^{\binom{n}{2}}}{[n]_{q}!} s^{n} t^{n}$$

$$= - (q-1) st \sum_{n=0}^{\infty} \frac{(-1)^{n} q^{\binom{n}{2}}}{[n]_{q}!} q^{n} s^{n} t^{n} + \sum_{n=0}^{\infty} \frac{(-1)^{n} q^{\binom{n}{2}}}{[n]_{q}!} s^{n} t^{n}$$

$$= - (q-1) st E_{q}(-qst) + E_{q}(-st).$$

$$(2.19)$$

Thus, by (2.19), we get

$$E_{q}(-qst) = \frac{1}{1 + (q-1)st} E_{q}(-st) = \frac{1}{(1 + (q-1)st)(1 + (q-1)q^{-1}st)} E_{q}(-q^{-1}st)$$

$$= \dots = \frac{1}{(1 + (q-1)st)(1 + (q-1)q^{-1}st) \dots (1 + (q-1)q^{-k}st)} E_{q}(-q^{-k}st).$$
(2.20)

Now, we discuss the q-differential equation. The main purpose of q-Laplace transform is in converting q-differential equation into simpler form which may be solved more easily. Like the ordinary Laplace transform, we can compute the q-Laplace transform of derivative by using the definition of q-Laplace transform.

Now, we oberve that

$$D_{q}(f(t)g(t)) = \frac{\partial}{\partial_{q}t}(f(t)g(t)) = \frac{\partial f(t)}{\partial_{q}t}g(t) + f(qt)\frac{\partial g(t)}{\partial_{q}t}$$

$$= (D_{q}f(t))g(t) + f(qt)(D_{q}g(t)).$$
(2.21)

Thus, by (2.21), we get

$$\int_0^\infty (D_q f(t)) g(t) d_q t = \int_0^\infty D_q(f(t)g(t)) d_q t - \int_0^\infty f(qt) (D_q g(t)) d_q t.$$
 (2.22)

It is easy to show that

$$D_{q}E_{q}(-qst) = \frac{\partial}{\partial_{q}t} \sum_{n=1}^{\infty} \frac{q^{\binom{n}{2}}(-1)^{n}}{[n]_{q}!} q^{n} s^{n} t^{n} = \sum_{n=1}^{\infty} \frac{q^{\binom{n}{2}}(-1)^{n}}{[n-1]_{q}!} q^{n} s^{n} t^{n-1}$$

$$= -s \sum_{n=0}^{\infty} \frac{q^{\binom{n}{2}}(-1)^{n}}{[n]_{q}!} q^{n+1} s^{n} t^{n}.$$
(2.23)

From (2.20), (2.22) and (2.23), we have

$$L_{q}(D_{q}f(t)) = \int_{0}^{\infty} (D_{q}f(t)) E_{q}(-qst) d_{q}t = -f(0) - \int_{0}^{\infty} f(qt) \frac{\partial}{\partial_{q}t} E_{q}(-qst) d_{q}t$$

$$= -f(0) + s \sum_{n=0}^{\infty} \frac{q^{\binom{n+1}{2}}}{[n]_{q}!} (-1)^{n} \int_{0}^{\infty} f(qt) q^{n+1} s^{n} t^{n} d_{q}t$$

$$= -f(0) + s \int_{0}^{\infty} f(t) \sum_{n=0}^{\infty} \frac{(-1)^{n} q^{\binom{n}{2}}}{[n]_{q}!} (qst)^{n} d_{q}t$$

$$= -f(0) + s \int_{0}^{\infty} f(t) E_{q}(-qst) d_{q}t = -f(0) + s L_{q}(f(t)).$$
(2.24)

If we replace f(t) by $D_q f(t)$, we see that

$$L_{q}(D_{q}^{2}f(t)) = -(D_{q}f)(0) + s \int_{0}^{\infty} (D_{q}f(t)) E_{q}(-q(t)) d_{q}t$$

$$= -(D_{q}f)(0) + sL_{q}(D_{q}f(t))$$

$$= -(D_{q}f)(0) + s(-f(0)) + sL_{q}(f(t))$$

$$= -(D_{q}f)(0) - sf(0) + s^{2}L_{q}(f(t)).$$
(2.25)

Continuing this process, we get

$$L_q\left(f^{(n)}(t)\right) = s^n L_q\left(f(t)\right) - \sum_{i=0}^{n-1} s^{n-1-i} f^{(i)}(0), \tag{2.26}$$

where
$$f^{(n)}(t) = \left(\frac{\partial_q}{\partial_q t}\right)^n f(t) = D_q^n f(t), \ f^{(n)}(0) = \left. f^{(n)}(t) \right|_{t=0}$$
.

where $f^{(n)}(t) = \left(\frac{\partial_q}{\partial_q t}\right)^n f(t) = D_q^n f(t), f^{(n)}(0) = f^{(n)}(t)\big|_{t=0}$. A function f is said to be of exponential order c if there exists c, M>0 and T > 0 such that

$$|f(t)| < Me^{ct}$$
 for all $t > T$.

If f(t) is piecewise continuous on the interval $(0, \infty)$ and of exponential order c, then $L_q(f(t))$ exists for s > c. Therefore, by (2.26), we obtain the following theorem.

Theorem 2.3. If $f, f', \dots, f^{(n-1)}$ are continuous on $(0, \infty)$ and are of exponential order and if $f^{(n)}(t)$ is piecewise continuous on $(0, \infty)$, then we have

$$L_q\left(f^{(n)}(t)\right) = s^n L_q\left(f(t)\right) - \sum_{i=0}^{n-1} s^{n-1-i} f^{(i)}(0),$$

where $f^{(n)}(t) = \left(\frac{\partial_q}{\partial_q t}\right)^n f(t)$

Let us consider the following q-derivative in s:

$$\frac{\partial F(q^{-1}s)}{\partial_q s} = \int_0^\infty \left(\frac{\partial_q}{\partial_q s} E_q(-st)\right) f(t) d_q t$$

$$= -\int_0^\infty t E_q(-qst) f(t) d_q t$$

$$= -L_q(tf(t)),$$
(2.27)

and

$$q\left(\frac{\partial_{q}}{\partial_{q}s}\right)^{2} F(q^{-2}s) = q\left(\frac{\partial_{q}}{\partial_{q}s}\right)^{2} \int_{0}^{\infty} E_{q}\left(-q^{-1}st\right) f(t) d_{q}t$$

$$= q \int_{0}^{\infty} \left(\left(\frac{\partial_{q}}{\partial_{q}s}\right)^{2} E_{q}\left(-q^{-1}st\right)\right) f(t) d_{q}t$$

$$= q \sum_{n=0}^{\infty} \frac{(-1)^{n+2}}{[n]_{q}!} q^{\binom{n}{2}} q^{n-1} s^{n} \int_{0}^{\infty} t^{n+2} f(t) d_{q}t \qquad (2.28)$$

$$= \int_{0}^{\infty} (-1)^{2} \left(\sum_{n=0}^{\infty} \frac{(-1)^{n} q^{\binom{n}{2}}}{[n]_{q}!} q^{n} s^{n} t^{n}\right) t^{2} f(t) d_{q}t$$

$$= (-1)^{2} \int_{0}^{\infty} E_{q}(-qst) t^{2} f(t) d_{q}t = (-1)^{2} L_{q}\left(t^{2} f(t)\right).$$

Continuing this process, we get

$$L_q(t^n f(t)) = (-1)^n q^{\binom{n}{2}} \left(\frac{\partial_q}{\partial_q s}\right)^n F(q^{-ns}). \tag{2.29}$$

Therefore, by (2.29), we obtain the following theorem.

Theorem 2.4 (q-Derivative of Transforms). For $n \in \mathbb{N}$, we have

$$L_q(t^n f(t)) = (-1)^n q^{\binom{n}{2}} \left(\frac{\partial_q}{\partial_q s}\right)^n F(q^{-n} s).$$

From Theorem 2.4, we note that

$$L_{q}(t^{n}e_{q}(at)) = (-1)^{n}q^{\binom{n}{2}} \left(\frac{\partial_{q}}{\partial_{q}s}\right)^{n} \int_{0}^{\infty} E_{q}(-q^{-n+1}st)e_{q}(at)d_{q}t$$

$$= (-1)^{n}q^{\binom{n}{2}} \left(\frac{\partial_{q}}{\partial_{q}s}\right)^{n} \left(\frac{1}{q^{-n}s-a}\right)$$

$$= \frac{(-1)^{n}q^{\binom{n}{2}}[n]_{q}!(-1)^{n}q^{-n^{2}}}{(s-a)(q^{-1}s-a)\cdots(q^{-n}s-a)}$$

$$= \frac{q^{-\binom{n+1}{2}}[n]_{q}!}{(s-a)(q^{-1}s-a)\cdots(q^{-n}s-a)},$$
(2.30)

and

$$L_{q}(e_{q}(at)f(t)) = \sum_{n=0}^{\infty} \frac{a^{n}}{[n]_{q}!} L_{q}(t^{n}f(t))$$

$$= \sum_{n=0}^{\infty} \frac{a^{n}}{[n]_{q}!} (-1)^{n} q^{\binom{n}{2}} \left(\frac{\partial_{q}}{\partial_{q}s}\right)^{n} F(q^{-n}s)$$

$$= \sum_{n=0}^{\infty} \frac{(-a)^{n}}{[n]_{q}!} q^{\binom{n}{2}} \left(\frac{\partial_{q}}{\partial_{q}s}\right)^{n} F(q^{-n}s).$$
(2.31)

Therefore, by (2.30) and (2.31), we obtain the following corollary.

Corollary 2.5. For $n \in \mathbb{N}$, we have

$$L_q\left(e_q(at)f(t)\right) = \sum_{n=0}^{\infty} \frac{(-a)^n}{[n]_q!} q^{\binom{n}{2}} \left(\frac{\partial_q}{\partial_q s}\right)^n F(q^{-n}s).$$

In particular, for $f(t) = t^n$ $(n \in \mathbb{N})$, we have

$$L_q(t^n e_q(at)) = \frac{q^{-\binom{n+1}{2}}[n]_q!}{(s-a)(q^{-1}s-a)\cdots(q^{-n}s-a)}.$$

For t, s > 0, the *q-beta function* is defined by

$$B_q(t,s) = \int_0^1 x^{t-1} (1 - qx)_q^{s-1} d_q x, \text{ (see [2,6])}.$$
 (2.32)

From (2.32), we note that

$$B_q(t,s) = \frac{\Gamma_q(t)\Gamma_q(s)}{\Gamma_q(t+s)}, \text{ (see [2,6])}.$$
(2.33)

It is easy to show that

$$T_q^{\alpha}(f) = \int_0^t (t - qs)_q^{\alpha - 1} f(s) d_q s = t \int_0^1 (t - qrt)_q^{\alpha - 1} f(rt) d_q r, \ (\alpha > 0).$$
 (2.34)

In particular, if we take $f(t) = t^{\beta}$ ($\beta > 0$), then

$$\int_{0}^{t} (t - qs)_{q}^{\alpha - 1} t^{\beta} d_{q} s = t^{\alpha + \beta} \int_{0}^{1} (1 - qr)_{q}^{\alpha - 1} r^{\beta} d_{q} r$$

$$= B_{q}(\alpha, \beta + 1) t^{\alpha + \beta}.$$
(2.35)

If functions f and g are piecewise continuous on $(0, \infty)$, then a special product, denote by f*g is defined by the integral

$$(f*g)(t) = \int_0^t f(\tau)g(t-\tau)d\tau, \qquad (2.36)$$

and is called *convolution* of f and g.

Now, we consider the q-analogue of convolution of f and g.

Let $f_1(t) = t^{\alpha}$, $g(t) = t^{\beta-1}$ $(\alpha, \beta > 0)$. Then we define the *q*-convolution of f and g as follows:

$$(f_1 * g)(t) = \int_0^t f_1(\tau)g(t - q\tau)d_q\tau, \tag{2.37}$$

where $g(t - q\tau) = (t - q\tau)_q^{\beta - 1}$. From (2.37), we have

$$(f_1 * g)(t) = \int_0^t \tau^{\alpha} (t - \tau q)_q^{\beta - 1} d_q \tau = t \int_0^1 (rt)^{\alpha} (t - qrt)_q^{\beta - 1} d_q r$$

$$= t^{\alpha + \beta} \int_0^1 r^{\alpha} (1 - qr)^{\beta - 1} d_q r = t^{\alpha + \beta} B_q(\alpha + 1, \beta)$$

$$= \frac{\Gamma_q(\alpha + 1) \Gamma_q(\beta)}{\Gamma_q(\alpha + \beta + 1)} t^{\alpha + \beta}.$$
(2.38)

Thus, by (2.38), we get

$$L_{q}(f_{1}*g) = B_{q}(\alpha + 1, \beta) \int_{0}^{\infty} E_{q}(-qst)t^{\alpha+\beta}d_{q}t$$

$$= B_{q}(\alpha + 1, \beta) \frac{1}{s^{\alpha+\beta+1}} \Gamma_{q}(\alpha + \beta + 1)$$

$$= \left(\frac{\Gamma_{q}(\alpha + 1)}{s^{\alpha+1}}\right) \left(\frac{\Gamma_{q}(\beta)}{s^{\beta}}\right).$$
(2.39)

By (2.8), we see that

$$L_q(f_1) = \frac{\Gamma_q(\alpha + 1)}{s^{\alpha + 1}}, \ L_q(g) = \frac{1}{s^{\beta}} \Gamma_q(\beta). \tag{2.40}$$

Hence, by (2.39) and (2.40), we get

$$L_q(f_1 * g) = L_q(f_1)L_q(g). (2.41)$$

Assume that f(t) is of the type such that equation (2.41). Then we have

$$f*g = \int_0^t (t - \tau q)_q^{\beta - 1} f(\tau) d_q \tau = t \int_0^1 (t - qrt)_q^{\beta - 1} f(rt) d_q r$$

$$= T_q^{\beta} (f(t)), \qquad (2.42)$$

where $g(t) = t^{\beta - 1}$, and

$$L_q\left(T_q^{\beta}f(t)\right) = L_q(f*g) = L_q(f)L_q(g)$$

$$= \frac{\Gamma_q(\beta)}{c^{\beta}}L_q(f). \tag{2.43}$$

If $f(t) = \sum_{i} a_i t^{\alpha_i}$, then we have

$$L_{q}(f*g) = \sum_{i} a_{i}L_{q}(t^{\alpha_{i}}*g)$$

$$= \sum_{i} a_{i}L_{q}(t^{\alpha_{i}})L_{q}(g)$$

$$= L_{q}\left(\sum_{i} a_{i}t^{\alpha_{i}}\right)L_{q}(g)$$

$$= L_{q}(f)L_{q}(g).$$
(2.44)

Therefore, by (2.44), we obtain the following theorem.

Theorem 2.6. For $f(t) = \sum_i a_i t^{\alpha_i}$ and $g(t) = t^{\beta-1}$, we have

$$L_q(f*g) = L_q(f)L_q(g).$$

For $\beta = 1$, let $T_q f(t) = T_q^1 f(t)$. Then we see that

$$L_q(T_q \sin_q t) = L_q(\sin_q t * 1) = L_q(\sin_q t) L_q(1)$$

$$= \left(\frac{1}{s^2 + 1}\right) \left(\frac{1}{s}\right) = \frac{1}{s(s^2 + 1)},$$
(2.45)

and

$$L_q(T_q \sin_q t) = L_q\left(\int_0^t \sin_q \tau d_q \tau\right). \tag{2.46}$$

By (2.45) and (2.46), we get

$$L_q \left(\int_0^t \sin_q \tau d_q \tau \right) = \frac{1}{s(s^2 + 1)}.$$
 (2.47)

Note that

$$L_{q}\left(\int_{0}^{t} (1 - \cos_{q} \tau) d_{q} \tau\right) = L_{q} \left(T_{q} \left(1 - \cos_{q} t\right)\right)$$

$$= L_{q} \left((1 - \cos_{q} t) * 1\right) = L_{q} \left(1 - \cos_{q} t\right) L_{q}(1)$$

$$= \left(\frac{1}{s} - \frac{s}{s^{2} + 1}\right) \frac{1}{s} = \frac{1}{s^{2}(s^{2} + 1)},$$
(2.48)

and

$$L_{q}\left(\int_{0}^{t} (\tau - \sin_{q} t) d_{q} \tau\right)$$

$$= L_{q}\left(T_{q}\left(t - \sin_{q} t\right)\right) = L_{q}\left((t - \sin_{q} t) * 1\right) = L_{q}\left(t - \sin_{q} t\right) L_{q}(1)$$

$$= \left(\frac{1}{s^{2}} - \frac{1}{s^{2} + 1}\right) \frac{1}{s} = \frac{1}{s^{2}(s^{2} + 1)} \frac{1}{s} = \frac{1}{s^{3}(s^{2} + 1)}.$$
(2.49)

Therefore, by (2.48) and (2.49), we obtain the following corollary.

Corollary 2.7. For $t \geq 0$, we have

$$L_q\left(\int_0^t (1-\cos_q \tau) d_q \tau\right) = \frac{1}{s^2(s^2+1)},$$

and

$$L_q\left(\int_0^t (\tau - \sin_q \tau) d_q \tau\right) = \frac{1}{s^3(s^2 + 1)},$$

Let u(t-a) be Heaviside function which defined by

$$u(t-a) = \begin{cases} 1 & \text{if } t \ge a, \\ 0 & \text{if } 0 \le t < a. \end{cases}$$
 (2.50)

Then, we have

$$L_{q}(u(t-a)) = \int_{0}^{\infty} E_{q}(-qst)u(t-a)d_{q}t = \int_{a}^{\infty} E_{q}(-qst)d_{q}t$$

$$= \int_{0}^{\infty} E_{q}(-qst)d_{q}t - \int_{0}^{a} E_{q}(-qst)d_{q}t$$

$$= \frac{1}{s} - \int_{0}^{a} E_{q}(-qst)d_{q}t = \frac{1}{s} - \sum_{n=0}^{\infty} \frac{(-1)^{n}q^{\binom{n}{2}}}{[n]_{q}!}(qs)^{n} \int_{0}^{a} t^{n}d_{q}t \quad (2.51)$$

$$= \frac{1}{s} + \frac{1}{s} \sum_{n=1}^{\infty} \frac{(-1)^{n}}{[n]_{q}!} q^{\binom{n}{2}} s^{n} a^{n} = \frac{1}{s} \sum_{n=0}^{\infty} \frac{(-1)^{n}q^{\binom{n}{2}}}{[n]_{q}!}(as)^{n}$$

$$= \frac{1}{s} E_{q}(-qas).$$

Therefore, by (2.51), we obtain the following theorem.

Theorem 2.8. Let u(t-a) be Heaviside function. Then we have

$$L_q(u(t-a)) = \frac{1}{s}E_q(-qas).$$

3. q-Laplace transform of the second kind

In this section, we consider the q-Laplace transform with $e_q(-st)$ and is called q-Laplace transform of the second kind. The q-Laplace transform of the second kind is defined by

$$\tilde{F}(s) = \tilde{L}_q(f(t)) = \int_0^\infty e_q(-st)f(t)d_qt, \ (s>0).$$
 (3.1)

As is well known, the q-gamma function of the second kind is defined by

$$\gamma_q(t) = \int_0^\infty x^{t-1} e_q(-x) d_q x, \ (t > 0), \ (\text{see [2,6]}).$$
 (3.2)

Thus, by (3.2), we get

$$\gamma_q(1) = 1, \ \gamma_q(t+1) = q^{-t}[t]_q \gamma_q(t), \ \gamma_q(n) = q^{-\binom{n}{2}} \Gamma_q(n), \ (n \in \mathbb{N}).$$
 (3.3)

For $\alpha \in \mathbb{R}$ with $\alpha > -1$, we have

$$\tilde{L}_{q}(t^{\alpha}) = \int_{0}^{\infty} e_{q}(-st)t^{\alpha}d_{q}t = \frac{1}{s^{\alpha+1}} \int_{0}^{\infty} e_{q}(-t)t^{\alpha}d_{q}t$$

$$= \frac{1}{s^{\alpha+1}} \gamma_{q}(\alpha+1).$$
(3.4)

In particular, for $\alpha = n \in \mathbb{N}$, by (3.3) and (3.4), we get

$$\begin{split} \tilde{L}_{q}\left(t^{n}\right) &= \frac{1}{s^{n+1}} \gamma_{q}(n+1) = \frac{1}{s^{n+1}} q^{-\binom{n+1}{2}} \Gamma_{q}(n+1) \\ &= \frac{1}{s^{n+1}} q^{-\frac{(n+1)n}{2}} [n]_{q}!. \end{split} \tag{3.5}$$

Let us take $\alpha = -\frac{1}{2}$. Then we have

$$\tilde{L}_q\left(t^{-\frac{1}{2}}\right) = \frac{1}{\sqrt{s}}\gamma_q\left(\frac{1}{2}\right), \ \tilde{L}_q\left(t^{\frac{1}{2}}\right) = \frac{1}{\sqrt{s^3}q[2]_{q^{\frac{1}{2}}}}\gamma_q\left(\frac{1}{2}\right).$$
 (3.6)

In a similar way, we can give the q-Laplace transform of the second kind for $e_q(at)$ and $E_q(at)$.

$$\tilde{L}_{q}\left(e_{q}(at)\right) = \sum_{n=0}^{\infty} \frac{a^{n}}{[n]_{q}!} \int_{0}^{\infty} e_{q}(-st)t^{n}d_{q}t = \sum_{n=0}^{\infty} \frac{a^{n}}{[n]_{q}!} \tilde{L}_{q}(t^{n})$$

$$= \sum_{n=0}^{\infty} \frac{a^{n}}{[n]_{q}!} \frac{q^{-\binom{n+1}{2}}}{s^{n+1}} \Gamma_{q}(n+1) = \sum_{n=0}^{\infty} \frac{a^{n}}{s^{n+1}} q^{-\binom{n+1}{2}},$$
(3.7)

and

$$\tilde{L}_{q}(E_{q}(at)) = \sum_{n=0}^{\infty} \frac{q^{\binom{n}{2}} a^{n}}{[n]_{q}!} \int_{0}^{\infty} t^{n} e_{q}(-st) d_{q}t = \sum_{n=0}^{\infty} \frac{q^{\binom{n}{2}} a^{n}}{[n]_{q}!} \tilde{L}_{q}(t^{n})$$

$$= \sum_{n=0}^{\infty} \frac{q^{\binom{n}{2}} a^{n}}{[n]_{q}!} \frac{q^{-\binom{n+1}{2}}}{s^{n+1}} \Gamma_{q}(n+1) = \frac{1}{s} \sum_{n=0}^{\infty} \left(\frac{a}{qs}\right)^{n}$$

$$= \frac{1}{s} \left(\frac{qs}{qs-a}\right) = \frac{q}{qs-a}.$$
(3.8)

Therefore, by (3.4)-(3.8), we obtain the following theorem.

Theorem 3.1. For $\alpha \in \mathbb{R}$ with $\alpha > -1$, we have

$$\tilde{L}_q(t^{\alpha}) = \frac{1}{s^{\alpha+1}} \gamma_q(\alpha+1), (s>0).$$

In particular, if $\alpha = n \in \mathbb{N}$, then we see that

$$\tilde{L}_q(t^n) = \frac{1}{s^{n+1}} q^{-\binom{n+1}{2}} [n]_q!.$$

Moreover,

$$\tilde{L}_q(e_q(at)) = \sum_{n=0}^{\infty} \frac{a^n}{s^{n+1}} q^{-\binom{n+1}{2}}, \ \tilde{L}_q(E_q(at)) = \frac{q}{qs-a},$$

and

$$\tilde{L}_q\left(t^{-\frac{1}{2}}\right) = \frac{1}{\sqrt{s}}\gamma_q\left(\frac{1}{2}\right), \ \ \tilde{L}_q\left(t^{\frac{1}{2}}\right) = \frac{1}{\sqrt{s^3}q[2]_{q^{\frac{1}{2}}}}\gamma_q\left(\frac{1}{2}\right).$$

From (1.1), we note that

$$E_q(it) = \sum_{n=0}^{\infty} \frac{q^{\binom{n}{2}}}{[n]_q!} i^n t^n.$$
 (3.9)

Thus, by (3.9), we get

$$E_q(it) = \sum_{n=0}^{\infty} \frac{(-1)^n q^{\binom{2n}{2}}}{[2n]_q!} t^{2n} + i \sum_{n=0}^{\infty} \frac{(-1)^n q^{\binom{2n+1}{2}}}{[2n+1]_q!} t^{2n+1}.$$
 (3.10)

By (3.10), we can define new *q-Sine* and *q-Cosine* function as follows:

$$\operatorname{Cos}_{q}(at) = \sum_{n=0}^{\infty} \frac{q^{\binom{2n}{2}} (-1)^{n} a^{2n}}{[2n]_{q}!} t^{2n} = \frac{1}{2} \left[E_{q}(iat) + E_{q}(-iat) \right], \tag{3.11}$$

and

$$\operatorname{Sin}_{q}(at) = \sum_{n=0}^{\infty} \frac{q^{\binom{2n+1}{2}}(-1)^{n}a^{2n+1}}{[2n+1]_{q}!}t^{2n+1} = \frac{1}{2}\left[E_{q}(iat) - E_{q}(-iat)\right].$$
(3.12)

Now, we give the q-Laplace transform of the second kind for q-Sine and q-Cosine functions:

From (2.16) and (3.7), we note that

$$\tilde{L}_{q}(\cos_{q} at) = \frac{1}{2} \left[\tilde{L}_{q}(e_{q}(iat)) + \tilde{L}_{q}(-iat) \right]
= \frac{1}{2} \left[\sum_{l=0}^{\infty} \frac{q^{-\binom{n+1}{2}}}{s^{n+1}} (ia)^{n} + \sum_{n=0}^{\infty} \frac{q^{-\binom{n+1}{2}}}{s^{n+1}} (-ia)^{n} \right]
= \frac{1}{2} \times 2 \sum_{n=0}^{\infty} \frac{q^{-n(2n+1)}}{s^{2n+1}} \left(\frac{a}{s} \right)^{2n} = \frac{1}{s} \sum_{n=0}^{\infty} (-1)^{n} q^{-n(2n+1)} \left(\frac{a}{s} \right)^{2n},$$
(3.13)

and

$$\tilde{L}_{q}\left(\sin_{q} at\right) = \sum_{n=0}^{\infty} \frac{(-1)^{n} a^{2n+1}}{[2n+1]_{q}!} \int_{0}^{\infty} e_{q}(-st) t^{2n+1} d_{q} t = \sum_{n=0}^{\infty} \frac{(-1)^{n} a^{2n+1}}{[2n+1]_{q}!} \tilde{L}_{q}(t^{2n+1})$$

$$= \sum_{n=0}^{\infty} \frac{(-1)^{n} a^{2n+1}}{[2n+1]_{q}!} \times \frac{[2n+1]_{q}!}{s^{2n+1}} q^{-\frac{(2n+1)(2n+2)}{2}}$$

$$= \frac{1}{s} \sum_{n=0}^{\infty} (-1)^{n} q^{-(n+1)(2n+1)} \left(\frac{a}{s}\right)^{2n+1}.$$
(3.14)

Therefore, by (3.13) and (3.14), we obtain the following theorem.

Theorem 3.2 (Transforms of q-cosine and q-sine).

$$\tilde{L}_q(\cos_q at) = \frac{1}{s} \sum_{n=0}^{\infty} (-1)^n q^{-n(2n+1)} \left(\frac{a}{s}\right)^{2n},$$

$$\tilde{L}_q(\sin_q at) = \frac{1}{s} \sum_{n=0}^{\infty} (-1)^n q^{-(n+1)(2n+1)} \left(\frac{a}{s}\right)^{2n+1}.$$
(3.15)

We consider q-Laplace transforms of the second kind for the q-Cosine and q-Sine functions of the second kind. By (3.11) and (3.12), we get

$$\tilde{L}_{q}\left(\operatorname{Cos}_{q}(at)\right) = \sum_{n=0}^{\infty} \frac{q^{\binom{2n}{2}}(-1)^{n}a^{2n}}{[2n]_{q}!} \int_{0}^{\infty} t^{2n}e_{q}(-st)d_{q}t$$

$$= \sum_{n=0}^{\infty} \frac{q^{\binom{2n}{2}}(-1)^{n}a^{2n}}{[2n]_{q}!} \times \frac{[2n]_{q}!}{s^{2n+1}}q^{-\binom{2n+1}{2}}$$

$$= \frac{1}{s} \sum_{n=0}^{\infty} \left(\frac{a}{qs}\right)^{2n} (-1)^{n} = \frac{q^{2}s}{(qs)^{2} + a^{2}},$$
(3.16)

and

$$\tilde{L}_{q}(\operatorname{Sin}_{q}(at)) = \frac{1}{2i} \left\{ \tilde{L}_{q}(E_{q}(iat)) - \tilde{L}_{q}(E_{q}(-iat)) \right\}
= \frac{1}{2i} \left\{ \frac{q}{qs - ia} - \frac{q}{qs + ia} \right\} = \frac{1}{2i} \times \frac{2qia}{(qs)^{2} + a^{2}} = \frac{qa}{(qs)^{2} + a^{2}}.$$
(3.17)

Therefore, by (3.16) and (3.17), we obtain the following theorem.

Theorem 3.3 (Transform of q-Sine and q-Cosine function).

$$\tilde{L}_q(\text{Cos}_q(at)) = \frac{q^2s}{(qs)^2 + a^2}, \ \tilde{L}_q(\text{Sin}_q(at)) = \frac{qa}{(qs)^2 + a^2}.$$

Here, we compute the q-Laplace transform of derivative by using the definition of the q-Laplace transform of the second kind. It is easy to show that

$$\int_{0}^{x} f'(t)g(t)d_{q}t = (f(x)g(t) - f(0)g(0)) - \int_{0}^{x} f(qt)g'(t)d_{q}t,$$
 (3.18)

where $f'(t) = D_q f(t) = \frac{\partial_q f(t)}{\partial_q t}$. By (1.2), (3.1) and (3.18), we get

$$\tilde{L}_{q}\left(f'(t)\right) = \int_{0}^{\infty} f'(t)e_{q}(-st)d_{q}t = -f(0) - \int_{0}^{\infty} f(qt)e_{q}'(-st)d_{q}t. \tag{3.19}$$

From (1.2), we note that

$$e_{q}'(-st) = \frac{\partial_{q}}{\partial_{q}t}e_{q}(-st) = \sum_{n=0}^{\infty} \frac{(-1)^{n}}{[n]_{q}!}s^{n}\frac{\partial_{q}t^{n}}{\partial_{q}t} = \sum_{n=1}^{\infty} \frac{(-1)^{n}s^{n}}{[n-1]_{q}!}t^{n-1}$$

$$= -s\sum_{n=0}^{\infty} \frac{(-1)^{n}}{[n]_{q}!}s^{n}t^{n} = -se_{q}(-st).$$
(3.20)

By (3.19) and (3.20), we get

$$\tilde{L}_{q}\left(f'(t)\right) = -f(0) + s \int_{0}^{\infty} f(qt)e_{q}(-st)d_{q}t
= -f(0) + sq^{-1} \int_{0}^{\infty} f(t)e_{q}\left(-q^{-1}st\right)d_{q}t
= -f(0) + sq^{-1}\tilde{F}\left(q^{-1}s\right),$$
(3.21)

where $f'(t) = D_q f(t) = \frac{\partial_q f(t)}{\partial_q t}$. If we replace f(t) by $D_q f(t) = \frac{\partial_q f(t)}{\partial_q t}$, then we have

$$\tilde{L}_{q}\left(f^{(2)}(t)\right) = s^{2}q^{-3}\tilde{F}\left(q^{-2}s\right) - q^{-1}sf(0) - f'(0), \tag{3.22}$$

and

$$\tilde{L}_{q}\left(f^{(3)}(t)\right) = s^{3}q^{-6}\tilde{F}\left(q^{-3}s\right) - s^{2}q^{-3}f(0) - sq^{-1}f'(0) - f^{(2)}(0),$$

where $f^{(3)}(t) = D_q^3 f(t) = \left(\frac{\partial_q}{\partial_q t}\right)^3 f(t)$, $f^{(2)}(0) = \left(\frac{\partial_q}{\partial_q t}\right)^2 f(t)\Big|_{t=0}$. Continuing this process, we get

$$\tilde{L}_q\left(f^{(n)}(t)\right) = s^n q^{-\binom{n+1}{2}} \tilde{F}\left(q^{-n}s\right) - \sum_{i=0}^{n-1} s^{n-1-i} q^{-\binom{n-i}{2}} f^{(i)}(0), \tag{3.23}$$

where $f^{(n)}(t) = D_q^n f(t) = \left(\frac{\partial_q}{\partial_q t}\right)^n f(t)$, $f^{(n)}(0) = \left(\frac{\partial_q}{\partial_q t}\right)^n f(t)\Big|_{t=0}$. Therefore, by (3.23), we obtain the following theorem.

Theorem 3.4. For $n \in \mathbb{N}$, we have

$$\tilde{L}_q\left(f^{(n)}(t)\right) = s^n q^{-\binom{n+1}{2}} \tilde{F}\left(q^{-n}s\right) - \sum_{i=0}^{n-1} s^{n-1-i} q^{-\binom{n-i}{2}} f^{(i)}(0),$$

where
$$\tilde{F}(s) = \tilde{L}_q(f(t)) = \int_0^\infty e_q(-st)f(t)d_qt$$
, $f^{(n)}(t) = \left(\frac{\partial_q}{\partial_q t}\right)^n f(t)$.

Let us consider the following s-derivative for the q-Laplace transform of the second kind.

$$\frac{\partial_{q}\tilde{F}(s)}{\partial_{q}s} = \frac{\partial_{q}\tilde{L}_{q}(f(t))}{\partial_{q}s} = \frac{\partial_{q}}{\partial_{q}s} \int_{0}^{\infty} e_{q}(-st)f(t)d_{q}t
= \int_{0}^{\infty} (-te_{q}(-st))f(t)d_{q}t = -\tilde{L}_{q}(tf(t)).$$
(3.24)

Thus, by (3.24), we get

$$\tilde{L}_q(tf(t)) = -\frac{\partial_q}{\partial_q s} \tilde{F}(s). \tag{3.25}$$

Continuing this process, we get

$$\tilde{L}_q(t^n f(t)) = (-1)^n \left(\frac{\partial_q}{\partial_q s}\right)^n \tilde{F}(s). \tag{3.26}$$

Therefore, by (3.26), we obtain the following theorem.

Theorem 3.5. For $n \in \mathbb{N}$, we have

$$\tilde{L}_q(t^n f(t)) = (-1)^n \left(\frac{\partial_q}{\partial_q s}\right)^n \tilde{F}(s).$$

For example, if we take $f(t) = e_q(at)$, then

$$\tilde{L}_q(t^n e_q(at)) = (-1)^n \left(\frac{\partial_q}{\partial_q s}\right)^n \tilde{L}_q(e_q(at)) = (-1)^n \left(\frac{\partial_q}{\partial_q s}\right)^n \sum_{k=0}^{\infty} \frac{q^{-\frac{k(k+1)}{2}}}{s^{k+1}} a^k$$

$$= (-1)^n \sum_{k=0}^{\infty} a^k (-1)^k q^{-nk - \binom{k+1}{2} - \binom{n+1}{2}} \times \frac{[k+1]_q [k+2]_q \cdots [k+n]_q}{s^{k+n+1}},$$

and

$$\tilde{L}_{q}(t^{n}E_{q}(at)) = (-1)^{n} \left(\frac{\partial_{q}}{\partial_{q}s}\right)^{n} \tilde{L}_{q}(E_{q}(at)) = (-1)^{n} \left(\frac{\partial_{q}}{\partial_{q}s}\right)^{n} \frac{q}{qs-a}$$

$$= (-1)^{n} (-1)^{n} q^{n+1} [n]_{q}! \frac{1}{(qs-a)(q^{2}s-a)\cdots(q^{n+1}s-a)}$$

$$= q^{n+1} \frac{[n]_{q}!}{(qs-a)(q^{2}s-a)\cdots(q^{n+1}s-a)}.$$

Finally, let us consider

$$L_{q}(e_{q}(at)f(t)) = \int_{0}^{\infty} e_{q}(-st)e_{q}(at)f(t)d_{q}t$$

$$= \sum_{n=0}^{\infty} \frac{a^{n}}{[n]_{q}!} \int_{0}^{\infty} e_{q}(-st)t^{n}f(t)d_{q}t$$

$$= \sum_{n=0}^{\infty} \frac{a^{n}}{[n]_{q}!} (-1)^{n} \tilde{L}_{q}(t^{n}f(t))$$

$$= \sum_{n=0}^{\infty} \frac{a^{n}}{[n]_{q}!} (-1)^{n} \left(\frac{\partial_{q}}{\partial_{q}s}\right)^{n} \tilde{L}_{q}(f(t))$$

$$= \sum_{n=0}^{\infty} \frac{a^{n}}{[n]_{q}!} (-1)^{n} \left(\frac{\partial_{q}}{\partial_{q}s}\right)^{n} \tilde{F}(s).$$

References

- G. Andrews, R. Askey and R. Roy, Special functions, Cambridge University Press, Cambridge, 1999.
- [2] R. Diaz and C. Teruel, q, k-generalized gamma and beta functions, J. Nonlinear Math. Phys. 12 (2005), no. 1, 118-134.
- [3] T. Mansour and A. S. Shabani, Generalization of some inequalities for the (q_1, \dots, q_s) -gamma function, Matematiche (Catania), **67** (2012), o. 2, 119-130.
- [4] M. Mansour, Determining the k-generalized gamma function $\Gamma_k(x)$ by functional equations, Int. J. Contemp. Math. Sci., 4 (2009), no. 21-24, 1037-1042.
- [5] T. Mansour, Some inequalities for the q-gamma function, J. Inequal. Pure Appl. Math., 9 (2008) no. 1, Article 18, 4 pp.
- [6] A. de Sole and V. Kac, On integral representations of q-gamma and q-beta functions, arXiv: math. QA/0302032, 2003.
- [7] F. Ucar and D. Albayrak, On q-Laplace type integral operators and their applications, J. Difference Equ. Appl., 18 (2012), no. 6, 1001-1014.
- [8] R. K. Yadav, S. O. Purohit and P. Nirwan, On q-Laplace transforms of a general class of q-polynomials and q-hypergeometric functions, Math. Maced., 7 (2009), 81-88.
- [9] D. G. Zill and M. R. Cullen, Advanced Engineering Mathematics, ones and Bartlett, 2005.
- 1 Department of Physics, Gyeongsang National University, Jinju $660,\!701,$ Republic of Korea.

 $E ext{-}mail\ address: mimip4444@hanmail.net}$

 2 Department of Mathematics, Kwangwoon University, Seoul 139-701, Republic of Korea.

E-mail address: tkkim@kw.ac.kr