

Affine Connected Generalized Dynkin Diagram with Rank 4

Zhengtang Tan ^a, Weicai Wu ^b, Shouchuan Zhang ^c

a. School of Engineering and Design, Hunan Normal University

Changsha 410081, P.R. China

b. School of Mathematics, Hunan Institute of Science and Technology

Yueyang 414006, P.R. China

c. Department of Mathematics, Hunan University

Changsha 410082, P.R. China

Emails: 1843186255@qq.com (ZTT); weicaiwu@hnu.edu.cn; z9491@sina.cn (SZ)(WCW)

Abstract

All affine connected Generalized Dynkin Diagram with rank 4 are found.

2010 Mathematics Subject Classification: 16W30, 22E60, 05L25

Keywords: Affine, Nichols algebra, Generalized Dynkin Diagram, Arithmetic root system.

Contents

1	Introduction	5
2	Affine over connected Generalized Dynkin Diagram with Rank 3	6
2.1	GDD 1 of Row 1	6
2.2	GDD 1 of Row 2	8
2.3	GDD 1 of Row 3	9
2.4	GDD 1 of Row 4	11
2.5	GDD 2 of Row 4	14
2.6	GDD 1 of Row 5	18
2.7	GDD 2 of Row 5	20
2.8	GDD 3 of Row 5	21

2.9	GDD 1 of Row 6	23
2.10	GDD 2 of Row 6	26
2.11	GDD 3 of Row 6	30
2.12	GDD 1 of Row 7	37
2.13	GDD 2 of Row 7	38
2.14	GDD 3 of Row 7	41
2.15	GDD 4 of Row 7	45
2.16	GDD 1 of Row 8	46
2.17	GDD 2 of Row 8	48
2.18	GDD 3 of Row 8	52
2.19	GDD 4 of Row 8	59
2.20	GDD 1 of Row 9	59
2.21	GDD 2 of Row 9	64
2.22	GDD 3 of Row 9	64
2.23	GDD 4 of Row 9	69
2.24	GDD 1 of Row 10	69
2.25	GDD 2 of Row 10	72
2.26	GDD 3 of Row 10	75
2.27	GDD 1 of Row 11	80
2.28	GDD 2 of Row 11	82
2.29	GDD 1 of Row 12	84
2.30	GDD 1 of Row 13	85
2.31	GDD 2 of Row 13	87
2.32	GDD 1 of Row 14	88
2.33	GDD 2 of Row 14	89
2.34	GDD 3 of Row 14	91
2.35	GDD 1 of Row 15	92
2.36	GDD 2 of Row 15	94
2.37	GDD 3 of Row 15	97
2.38	GDD 4 of Row 15	100
2.39	GDD 1 of Row 16	100
2.40	GDD 2 of Row 16	102
2.41	GDD 3 of Row 16	104
2.42	GDD 4 of Row 16	105
2.43	GDD 5 of Row 16	106
2.44	GDD 1 of Row 17	106
2.45	GDD 2 of Row 17	109
2.46	GDD 3 of Row 17	110

2.47	GDD 4 of Row 17	111
2.48	GDD 5 of Row 17	114
2.49	GDD 6 of Row 17	115
2.50	GDD 7 of Row 17	116
2.51	GDD 8 of Row 17	118
2.52	GDD 9 of Row 17	119
2.53	GDD 1 of Row 18	120
2.54	GDD 2 of Row 18	121
3	Properties about arithmetic root system with rank 3 and 4	121
3.1	Classical types	121
3.2	Properties about arithmetic root system with rank 3	123
3.3	Properties about arithmetic root system with rank 4	137
3.4	Affine with circle	157
4	The proof of Theorem 2.1 from subsection 4.1 to subsection 4.55	158
4.1	GDD 1 of Row 1	158
4.2	GDD 1 of Row 2	162
4.3	GDD 1 of Row 3	166
4.4	GDD 1 of Row 4	169
4.5	GDD 2 of Row 4	178
4.6	GDD 1 of Row 5	189
4.7	GDD 2 of Row 5	194
4.8	GDD 3 of Row 5	197
4.9	GDD 1 of Row 6	201
4.10	GDD 2 of Row 6	206
4.11	GDD 3 of Row 6	217
4.12	GDD 1 of Row 7	227
4.13	GDD 2 of Row 7	229
4.14	GDD 3 of Row 7	237
4.15	GDD 4 of Row 7	250
4.16	GDD 1 of Row 8	252
4.17	GDD 2 of Row 8	259
4.18	GDD 3 of Row 8	267
4.19	GDD 4 of Row 8	278
4.20	GDD 1 of Row 9	278
4.21	GDD 2 of Row 9	287
4.22	GDD 3 of Row 9	287
4.23	GDD 4 of Row 9	295

4.24	GDD 1 of Row 10	295
4.25	GDD 2 of Row 10	300
4.26	GDD 3 of Row 10	307
4.27	GDD 1 of Row 11	318
4.28	GDD 2 of Row 11	324
4.29	GDD 1 of Row 12	329
4.30	GDD 1 of Row 13	331
4.31	GDD 2 of Row 13	335
4.32	GDD 1 of Row 14	336
4.33	GDD 2 of Row 14	339
4.34	GDD 3 of Row 14	344
4.35	GDD 1 of Row 15	347
4.36	GDD 2 of Row 15	352
4.37	GDD 3 of Row 15	358
4.38	GDD 4 of Row 15	367
4.39	GDD 1 of Row 16	369
4.40	GDD 2 of Row 16	372
4.41	GDD 3 of Row 16	380
4.42	GDD 4 of Row 16	389
4.43	GDD 5 of Row 16	392
4.44	GDD 1 of Row 17	393
4.45	GDD 2 of Row 17	400
4.46	GDD 3 of Row 17	402
4.47	GDD 4 of Row 17	407
4.48	GDD 5 of Row 17	414
4.49	GDD 6 of Row 17	419
4.50	GDD 7 of Row 17	420
4.51	GDD 8 of Row 17	426
4.52	GDD 9 of Row 17	430
4.53	GDD 1 of Row 18	432
4.54	GDD 2 of Row 18	434
4.55	More about GDD 3 of Row 15	435

5 Appendix 436

5.1	Affine over connected Generalized Dynkin Diagram with Rank 1 or 2	436
-----	-------------------------------------------------------------------	-----

1 Introduction

Nichols algebras play a fundamental role in the classification of finite-dimensional complex pointed Hopf algebras by means of the lifting method developed by Andruskiewitsch and Schneider [AS02, AS10, AHS08]. Heckenberger [He06a, He05] classified arithmetic root systems. Heckenberger [He06b] proved a GDD is an arithmetic root system if and only if corresponding matrix is a finite Cartan matrix for GDDs of Cartan types. W. Wu, S. Zhang and Y.-Z. Zhang [WZZ15b] proved a Nichols Lie algebra is a finite dimensional if and only if its GDD is an arithmetic root system.

In this paper, using Table *A1*, *A2*, *B* and *C* in [He06a, He05], we find all affine connected Generalized Dynkin Diagram with rank 4.

Preliminaries

We now recall some basic concepts of the graph theory (see [Ha]). Let Γ_1 be a non-empty set and $\Gamma_2 \subseteq \{\{u, v\} \mid u, v \in \Gamma_1, \text{ with } u \neq v\} \subseteq 2^{\Gamma_1}$. Then $\Gamma = (\Gamma_1, \Gamma_2)$ is called a graph; Γ_1 is called the vertex set of Γ ; Γ_2 is called the edge set of Γ ; Element $\{u, v\} \in \Gamma_2$ is called an edge, written $a_{u,v}$. Let F be an algebraically closed field of characteristic zero. If $\{x_1, \dots, x_n\}$ is a basis of vector space V and $C(x_i \otimes x_j) = q_{ij}x_j \otimes x_i$ with $q_{ij} \in F$, then V is called a braided vector space of diagonal type, $\{x_1, \dots, x_n\}$ is called canonical basis and $(q_{ij})_{n \times n}$ is called braided matrix. Let $\tilde{q}_{i,j} := q_{ij}q_{ji}$ for $i, j \in \{1, 2, \dots, n\}$ with $i \neq j$. If let $\Gamma_1 = \{1, 2, \dots, n\}$ and $\Gamma_2 = \{\{u, v\} \mid \tilde{q}_{u,v} \neq 1, u, v \in \Gamma_1, \text{ with } u \neq v\}$, then $\Gamma = (\Gamma_1, \Gamma_2)$ is a graph. Set q_{ii} over vertex i and $\tilde{q}_{i,j}$ over edge a_{ij} for $i, j \in \Gamma_1$ with $i \neq j$. Then $\Gamma = (\Gamma_1, \Gamma_2)$ is called a Generalized Dynkin Diagram of braided vector space V , written a GDD in short. If $\Delta(\mathfrak{B}(V))$ is an arithmetic root system. then we call its GDD an arithmetic root system for convenience.

Let $(q_{ij})_{n \times n}$ be a matrix. If $q_{ij}q_{ji} \begin{cases} \neq 1, & \text{when } |j - i| = 1 \\ = 1, & \text{when } |j - i| \neq 1 \end{cases}$ for any $1 \leq i \neq j \leq n$, then $(q_{ij})_{n \times n}$ is called a chain or labelled chain. If $(q_{ij})_{n \times n}$ is a chain and

$$(q_{11}q_{1,2}q_{2,1} - 1)(q_{11} + 1) = 0; (q_{n,n}q_{n,n-1}q_{n-1,n} - 1)(q_{n,n} + 1) = 0; \quad (1)$$

i.e.

$$q_{ii} + 1 = q_{i,i-1}q_{i-1,i}q_{i,i+1}q_{i+1,i} - 1 = 0 \quad (2)$$

$$\text{or } q_{ii}q_{i,i-1}q_{i-1,i} = q_{ii}q_{i,i+1}q_{i+1,i} = 1, \quad (3)$$

$1 < i < n$, then $(q_{ij})_{n \times n}$ is called a simple chain (see [He06a, Def.1]). Conditions (1), (2) and (3) are called simple chain conditions. Let $q := q_{n,n}^2 q_{n-1,n} q_{n,n-1}$. For $0 \leq j \leq n$

and $0 < i_1 < i_2 < \cdots < i_j \leq n$, let $C_{n,q,i_1,i_2,\dots,i_j}$ denote a simple chain which satisfies the condition: $\tilde{q}_{i,i-1} = q$ if and only if $i \in \{i_1, i_2, \dots, i_j\}$, where $\tilde{q}_{1,0} := \frac{1}{q_{11}^2 \tilde{q}_{12}}$. For example,

$$n \geq 2, \quad \begin{array}{ccccccc} q & q^{-1}q & q^{-1}q & \cdots & q & q^{-1}q & \\ \bullet & \bullet & \bullet & \cdots & \bullet & \bullet & \\ 1 & 2 & 3 & & n-1 & n & \end{array}, q \in F^*/\{1, -1\}.$$

is $C_{n,q,i_1,i_2,\dots,i_j}$ with $j = 0$.

$$\begin{array}{ccccccc} q^{-1}q & q^{-1}q & q^{-1} & \cdots & q^{-1}q & -1 & \\ \bullet & \bullet & \bullet & \cdots & \bullet & \bullet & \\ 1 & 2 & 3 & & n-1 & n & \end{array}, q \in F^*/\{1\}.$$

is $C_{n,q,i_1,i_2,\dots,i_j}$ with $j = n$.

For the convenience, we let $C_{1,q,i_1,i_2,\dots,i_j}$ denote the GDD with length 1 satisfied the following conditions: $q = q_{11}$ when $q_{11} \neq -1$; q can be any number when $q_{11} = -1$.

Every connected subGDD of every arithmetic root system in Row 1-10 in Table C is called a classical GDD. Every connected GDD which is not classical is called an exception.

If omitting every vertex in a connected GDD with rank n is an arithmetic root system and this GDD is not an arithmetic root system, then this GDD is called an affine GDD over connected subGDD of this GDD with rank $n-1$, affine GDD in short. In other word, If a GDD is affine of a braided vector space V , then Nichols algebra of every proper subGDD is finite dimensional and $\dim \mathfrak{B}(V) = \infty$. If a GDD is obtained by relabelling other GDD, then we view the two GDD are the same since their YD modules are isomorphic. For example,

$$\begin{array}{ccc} q & q^{-1}q & q^{-1}q^{-3} \\ \bullet & \bullet & \bullet \end{array}, q \in R_9.$$

and

$$\begin{array}{ccc} q^{-3} & q^{-1}q & q^{-1}q \\ \bullet & \bullet & \bullet \end{array}, q \in R_9.$$

are the same.

2 Affine over connected Generalized Dynkin Diagram with Rank 3

Theorem 2.1. *A connected GDD with rank 4 is affine if and only if it is one of following lists from subsection 2.1 to subsection 2.54.*

2.1 GDD 1 of Row 1

(1).

$$\begin{array}{c} -1 \quad q^{-2} \quad q \quad q^{-1}q \quad q^{-1} \quad q \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ , q \in R_3 \cup R_6 \end{array}$$

$$\begin{array}{c} q^{-3} \quad (2) \cdot \\ q^{-1} \quad q \quad q^{-1}q \quad q^{-1} \quad q \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ , q \in R_9 \end{array}$$

$$(3). \quad \begin{array}{c} \bullet \quad q^2 \\ | \\ q^{-2} \\ \bullet \text{---} \bullet \text{---} \bullet \\ q \quad q^{-2} \quad q^2 \quad q^{-2}q^2 \end{array}$$

$$, q \in F^* \setminus \{1, -1\}.$$

$$(4). \quad \begin{array}{c} \bullet \quad q \\ | \\ q^{-1} \\ \bullet \text{---} \bullet \text{---} \bullet \\ q^2 \quad q^{-2} \quad q \quad q^{-1}q \end{array}$$

$$, q \in F^* \setminus \{1, -1\}.$$

$$(5). \quad \begin{array}{c} \bullet \quad -q^{-1} \\ | \\ -q \\ \bullet \text{---} \bullet \text{---} \bullet \\ q \quad -q \quad -q^{-1}q \quad -q^{-1} \end{array}$$

$$, q \in R_3.$$

$$(6). \quad \begin{array}{c} \bullet \quad q \\ | \\ q^{-1} \\ \bullet \text{---} \bullet \text{---} \bullet \\ -1 \quad q^{-2} \quad q \quad q^{-1}q \end{array}$$

$$, q \in R_3 \cup R_6.$$


$$(7). \quad \begin{array}{c} \bullet \quad -q \\ | \\ -q^{-1} \\ \bullet \text{---} \bullet \text{---} \bullet \\ q \quad -q^{-1}q \quad -q^{-1}q \quad -q^{-1}q \end{array}$$

$$, q \in R_3.$$

$$(8). \quad \begin{array}{c} \bullet \quad q \\ | \\ q^{-1} \\ \bullet \text{---} \bullet \text{---} \bullet \\ q^{-3}q^{-1} \quad q \quad q^{-1}q \end{array}$$


$$, q \in R_9.$$

$$(9)$$




$$q \neq 1$$

$$(10)$$



$$q \neq 1$$

$$(11)$$



$$q \in R_3$$

2.2 GDD 1 of Row 2

$$(1).$$

$$q \quad q^{-2} \quad q^2 \quad q^{-2}q^2 \quad q^{-2} \quad q$$

$$, q \in F^* \setminus \{1, -1\}$$

$$(2).$$

$$q^4 \quad q^{-4} \quad q^2 \quad q^{-2}q^2 \quad q^{-2} \quad q$$

$$q^4 \neq 1,$$

$$(3).$$

$$q^2 \quad -q^{-2}q^{-2}q^{-2}q^{-2}q^{-2}q^2 \quad iq$$

$$q \in R_3$$

$$(4).$$

$$-1 \quad q^{-4} \quad q^2 \quad q^{-2}q^2 \quad q^{-2} \quad q$$

$$q \in R_3 \cup R_6.$$

$$(5).$$


$$q^2 \quad -q^{-2}q^2 \quad -q^{-2}q^2 \quad -q^{-2}q^2$$

$$q \in R_3$$

$$(6).$$


$$q^{-6} \quad q^{-2} \quad q^2 \quad q^{-2}q^2 \quad q^{-2} \quad q$$

$$q \in R_9$$




$q^2 \neq 1.$

(8)




$q^2 \neq 1.$

(9)




$q^2 \neq 1.$

(11)




$q \in R_3$

(12)




$q^2 \neq 1.$


2.3 GDD 1 of Row 3



$q^4 \neq 1.$



$q^2 \neq 1.$



$q \in R_3$

(4).


$$\begin{array}{c} -1 \quad q^{-2} \quad q \quad q^{-1}q \quad q^{-2} \quad q^2 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ q \in R_3 \cup R_6. \end{array}$$

$$\begin{array}{c} (5). \\ q \quad -q^{-1}q \quad -q^{-1}q \quad q^{-2} \quad q^2 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ q \in R_3 \end{array}$$

$$\begin{array}{c} (6). \\ q^{-3} \quad q^{-1} \quad q \quad q^{-1}q \quad q^{-2} \quad q^2 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array}$$

$$q \in R_9$$


(7)




$$q^2 \neq 1.$$

(8)

$$q \notin R_3 .$$




(9)




$$\text{and } q = q^4$$

(10)




$$q^2 = -1.$$

(11)



$$q^2 \neq 1.$$


(12)



$$q^2 \neq 1.$$

(13)

$q^2 \neq 1$.



2.4 GDD 1 of Row 4

$$\begin{array}{cccc} \text{(1)} & & & \\ -1 & q & -1 & q^{-2}q^2 & q^{-2} & q^2 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

when $q \in R_3$.

$$\begin{array}{cccc} \text{(2)} & & & \\ -1 & q & -1 & q^{-3}q^3 & q^{-3} & q^3 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

$q^6 \neq 1$.

$$\begin{array}{cccc} \text{(3)} & & & \\ -q & q^2 & -1 & q^{-3}q^3 & q^{-3} & q^3 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

$q^6 \neq 1$.

$$\begin{array}{cccc} \text{(4)} & & & \\ r & r^{-1} & -1 & q^{-1}q & q^{-1} & q \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

except $r = -1, q \in R_3; r = -q^{-1}, q^2 \neq 1; q^{-2} = r, q^2 \neq 1; q^3 = q^{-2}, q^6 \neq 1$.

$$\begin{array}{cccc} \text{(5)} & & & \\ q & q^{-1} & -1 & q^2 & q^{-2}q^2 & q^{-2} \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

when $q \notin R_4$

$$\begin{array}{cccc} \text{(6)} & & & \\ q & q^{-1} & -1 & q^{-1}q & q^{-1} & q \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

$q^2 \neq 1$.

$$\overset{(7)}{-1 \quad -1 \quad -q^{-1}-q \quad -q^{-1}-q}$$

$$q^2 \neq 1.$$

$$(8)$$

$$q^2 \neq 1.$$

$$(9)$$

$$q^3 \neq 1.$$

$$(10)$$

$$q^2 \neq -1.$$

$$(11)$$

$$q \in R_3.$$

$$(12)$$

$$q \in R_3$$

$$(13)$$









$$q \in R_6$$

$$(14)$$


$$q \notin R_2 \cup R_3$$

$$(15)$$

$$q \in R_3$$




(16)



$$q^{-1} = q^2.$$


(17)

$$q = r^2, q^2 \neq 1.$$




(18)

$$\text{when } r = q^2, q^2 \neq 1.$$




(19)

$$\text{when } -1 = r, q^2 \neq 1.$$




(20)

$$\text{when } r = -q^{-1}, q \in R_3$$




(21)

$$\text{when } q \in R_3$$




(22)

$$\text{when } q^{-1} = q^4$$




(23)

$$q \notin R_2 \cup R_3$$




(24)

$$\text{when } q = q^{-4}.$$




(25)

$q \notin R_2 \cup R_3$




(26)

when $-1 = q^{-2}$




(27)

$q \in R_3$




(28)


$q \in R_6$




2.5 GDD 2 of Row 4




$q \notin R_2 \cup R_3$




$q \in R_3$











$q \in R_3$



$q \in R_3$









$$q \in R_3$$

(14)




$$q^2 \neq 1.$$

(15)




$$q^2 \neq 1.$$

(16)




$$\text{when } q^2 = -1.$$

(17)




$$\text{when } q^4 = q.$$

(18)




$$\text{when } -1 = q^3$$

(19)




$$q^6 = q$$

(20)




when $q = q^{-2}$ or $q = q^{-3}$.

(21)




$q \in R_3$

(22)




when $-q^{-2} = q$.

(23).




$q \in R_3$.

(24)




$q \in R_3$.

(25)




$q \in R_3$.

(26)



$q \in R_3$.

(27)



$q \in R_3$.

2.6 GDD 1 of Row 5

$$\begin{array}{cccc} \textcircled{1} & & & \\ -q^{-1}q^2 & -1 & q^{-2}q^2 & q^{-2} \quad q \\ \bullet & \bullet & \bullet & \bullet \end{array}$$

$$q^4 \neq 1.$$

$$\begin{array}{cccc} \textcircled{2} & & & \\ -1 & q & -1 & q^{-2}q^2 \quad q^{-2} \quad q \\ \bullet & \bullet & \bullet & \bullet \end{array}$$

$$q^4 \neq 1.$$

$$\begin{array}{cccc} \textcircled{3} & & & \\ -1 & q^2 & -1 & q^{-6}q^6 \quad q^{-6} \quad q^3 \\ \bullet & \bullet & \bullet & \bullet \end{array}$$

$$q^2 \notin R_2 \cup R_3.$$

$$\begin{array}{cccc} \textcircled{4} & & & \\ -q^{-2}q^4 & -1 & q^{-6}q^6 & q^{-6} \quad q^3 \\ \bullet & \bullet & \bullet & \bullet \end{array}$$

$$q^{12} \neq 1.$$

$$\begin{array}{cccc} \textcircled{5} & & & \\ r^2 & r^{-2} & -1 & q^{-2}q^2 \quad q^{-2} \quad q \\ \bullet & \bullet & \bullet & \bullet \end{array}$$

$$r^{-2}, q^4 \neq 1. \quad q, q^{-1} \neq r.$$

$$\begin{array}{cccc} \textcircled{6} & & & \\ q & q^{-1} & -1 & q^2 \quad q^{-2} \quad q^{-2} \quad q \\ \bullet & \bullet & \bullet & \bullet \end{array}$$

$$q \notin R_2 \cup R_3, \quad q^4 \neq 1.$$

$$\begin{array}{cccc} \textcircled{7} & & & \\ q^2 & q^{-2} & -1 & q^{-2}q^2 \quad q^{-2} \quad q \\ \bullet & \bullet & \bullet & \bullet \end{array}$$

$$q^2 \notin R_2 \cup R_3, \quad q^4 \neq 1.$$

$$\begin{array}{cccc} \textcircled{8} & & & \\ q^{-1} & q & -1 & q^{-2}q^2 \quad q^{-2} \quad q \\ \bullet & \bullet & \bullet & \bullet \end{array}$$

$$q \notin R_2 \cup R_3. \quad q^4 \neq 1.$$

$$\begin{array}{cccc} \textcircled{9} & & & \\ q^2 & q^{-2} & -1 & q^{-2}q^2 \quad q^{-2} \quad q \\ \bullet & \bullet & \bullet & \bullet \end{array}$$

$$, q^2 \in R_3. \quad q^4 \neq 1.$$

$$q^{-2} \begin{matrix} (10) \\ -q^{-2} - 1 - q^{-2} - q^{-2} \end{matrix} \begin{matrix} i q \\ \bullet \end{matrix}$$

, $q \in R_3$


$$-1 \begin{matrix} (11) \\ q^2 - 1 - q^{-2} - q^2 - q^{-2} \end{matrix} \begin{matrix} i q \\ \bullet \end{matrix}$$

, $q \in R_3$

$$q^2 \begin{matrix} (12) \\ -1 - q^{-2} - q^2 - q^{-2} \end{matrix} \begin{matrix} i q \\ \bullet \end{matrix}$$


, $q \in R_3$

(13)



$q^4 \neq 1$.

(14)




$q^4 \neq 1$.

(15)


when $q = q^4$ or $q = q^6$.

(16)





$q^4 \neq 1$.

(17)




when $q = q^{-2}$.

(18)




when $q = r^2$. $q^4 \neq 1$. $r^2 \neq 1$; $r^2 \neq q^2, q^{-2}$.

(19)




when $-1 = r^2$. $q^4 \neq 1$. $r^2 \neq 1$; $r^2 \neq q^2, q^{-2}$.

(20)




$q^4 \neq 1$.


2.7 GDD 2 of Row 5




$q^4 \neq 1$.




$q^4 \neq 1$.




$q^4 \neq 1$. $q^2 \notin R_3$.




, $q^2 \in R_3$.



, $q^2 \in R_3$.




$q^2 \in R_3$.




, $q \in R_3$.

(8).




$q \in R_3$




, $q \in R_3$

(10)




when $q^6 = 1$.

(11)



$q^4 \neq 1$.


(12)




when $q \in R_4 \cup R_5 \cup R_7 \cup R_9$.

(13)

$q \in R_6$.




(14)




$q \in R_3 \cup R_6$.

(15)



when $q = q^4$,

2.8 GDD 3 of Row 5



$$, q^2 \in F^* \setminus \{1, -1\}$$

$$(2). \quad q^6 \quad q^{-6} \quad q^2 \quad q^{-2}-1 \quad q^2 \quad -q^{-1}$$

$$, q^2 \in F^* \setminus \{1, -1\}, q^2 \notin R_3.$$

$$(3). \quad q^2 \quad -q^2 \quad -q^{-2} \quad -q^2-1 \quad -q^{-2}-iq$$

$$, q \in R_3.$$

$$(4). \quad -1 \quad q^2 \quad q^{-2}-1 \quad q^2 \quad -q^{-1}$$

$$, q \in R_3.$$

$$(5). \quad -q^2 \quad -q^{-2}q^2 \quad q^{-2}-1 \quad q^2 \quad -q^{-1}$$

$$, q \in R_3.$$

$$(6). \quad -1 \quad -1 \quad q^2 \quad q^{-2}-1 \quad q^2 \quad -q^{-1}$$

$$, q^2 \in R_3.$$

$$(7). \quad -1 \quad q^2 \quad -q^2 \quad -q^{-2}-1 \quad -q^2 \quad -(iq)^{-1}$$

$$, q^2 \in R_3$$

$$(8). \quad -1 \quad -q^2 \quad q^2 \quad q^{-2}-1 \quad q^2 \quad -q^{-1}$$

$$, q^2 \in R_3.$$



$$(9). \quad q \quad q^{-2} \quad q^2 \quad q^{-2}-1 \quad q^2 \quad -q^{-1}$$

$$q^4 \neq 1.$$

$$(10)$$


$$q^4 \neq 1,$$

$$(11)$$




$$\text{when } q \in R_6 \cup R_8$$

$$(12)$$




$$q^4 \neq 1, q \in R_3 \cup R_2$$

(13)




$$q^4 \neq 1.$$

(14)




$$q^4 \neq 1.$$

(15)



$$q^2 \in R_3$$


(16)




$$q^2 \in R_3.$$

2.9 GDD 1 of Row 6


(1)



$$q^4 \neq 1,$$




$$q^4 \neq 1,$$




$$q \notin R_3 \cup R_2, q^6 \neq 1,$$

$q \notin R_2 \cup R_3.$
(15)




$q^2 \neq 1.$

(16)




when $q^2 = -1$

(17)




$q \in R_3.$

(18)




when $q \in R_5 \cup R_4$

(19)




when $q^2 = q^{-1}$

(20)




when $r = -1. q^2 \neq 1.$

(21)




when $r = q^2. q^2 \neq 1.$

(22)




when $-1 = q^{-2}$.

(23)



when $q^2 = q^{-2}$.

(24)



$q \in R_3$

2.10 GDD 2 of Row 6

(1).
 $\bullet \xrightarrow{q} \bullet \xrightarrow{q^{-2}} \bullet \xrightarrow{-1} \bullet \xrightarrow{q^2} \bullet \xrightarrow{-1} \bullet \xrightarrow{q^{-4}} \bullet \xrightarrow{q^4}$
 $, q \in F^* \setminus \{1, -1\}.$


(2).
 $\bullet \xrightarrow{q^2} \bullet \xrightarrow{q^{-2}} \bullet \xrightarrow{-1} \bullet \xrightarrow{q} \bullet \xrightarrow{-1} \bullet \xrightarrow{q^{-2}} \bullet \xrightarrow{q^2}$
 $, q \in F^* \setminus \{1, -1\}.$

(3).
 $\bullet \xrightarrow{q^3} \bullet \xrightarrow{q^{-3}} \bullet \xrightarrow{-1} \bullet \xrightarrow{q} \bullet \xrightarrow{-1} \bullet \xrightarrow{q^{-2}} \bullet \xrightarrow{q^2}$
 $, q \in F^* \setminus \{1, -1\}. q \notin R_3,$


(4).
 $\bullet \xrightarrow{q} \bullet \xrightarrow{-q} \bullet \xrightarrow{-q^{-1}} \bullet \xrightarrow{1} \bullet \xrightarrow{q^2} \bullet \xrightarrow{q^{-2}}$
 $, q \in R_3.$

(5).
 $\bullet \xrightarrow{-1} \bullet \xrightarrow{q} \bullet \xrightarrow{-1} \bullet \xrightarrow{q} \bullet \xrightarrow{-1} \bullet \xrightarrow{q^{-2}} \bullet \xrightarrow{q^2}$
 $, q \in R_3.$


(6).
 $\bullet \xrightarrow{-q} \bullet \xrightarrow{-q^{-1}} \bullet \xrightarrow{1} \bullet \xrightarrow{q} \bullet \xrightarrow{-1} \bullet \xrightarrow{q^{-2}} \bullet \xrightarrow{q^2}$
 $q \in R_3.$




, $q \in R_3$



$q \in R_3$




, $q \in R_3$.




, $q \in R_3$.

(19)




when $q = q^4$.

(20)




$q^2 \neq 1$.

(21)




$q^4 = q$

(22)




and $q^6 = q^2$

(23)




and $q^2 = -1$

(24)




$q^2 \neq 1$.

(25)




and $q^3 = -1$

(26)




$q^2 \neq 1$.

(27)




when $q \in R_4 \cup R_5 \cup R_3$

(28)




and $-1 = q^2$

(29)




$q \in R_3$

(30)




$q \in R_3$

(31)




$q \in R_3$

(32)




$q \in R_3$

(33)




$q \in R_3$

(34)




$q \in R_3$

(35)




$q \in R_3$

(36)




$q \in R_6$

2.11 GDD 3 of Row 6






, $q \in F^* \setminus \{1, -1\}$.


(2).




when $q \in R_3, q \in F^* \setminus \{1, -1\}$




(6).



, $q \in R_3.$




(8).




, $q \in R_3$.


(9).



, $q \in R_3$.




, $q \in R_3$.




, $q \in R_3$.

(12).



, $q \in R_3$.



$q^4 \neq 1$,


$$(14) \begin{array}{c} q^4 \quad -1 \\ \bullet \quad \bullet \\ \bullet \quad \bullet \\ -q^{-1}q^2 \quad -1 \quad q^{-2}q^2 \end{array}$$

when $q \in R_6$

$$(15) \begin{array}{c} q^4 \quad -1 \\ \bullet \quad \bullet \\ \bullet \quad \bullet \\ -1 \quad q \quad -1 \quad q^{-2}q^2 \end{array}$$

when $q \in R_3$

(16)




when $q \in R_3 \cup R_5$

$$(17) \begin{array}{c} q^6 \quad -1 \\ \bullet \quad \bullet \\ \bullet \quad \bullet \\ -1 \quad q \quad -1 \quad q^{-3}q^3 \end{array}$$


when $q \in R_7$

(18)




when $q \in R_4 \cup R_6$.


(19)




when $q \in R_8$.




when $q \in R_4 \cup R_5 \cup R_7 \cup R_6$




when $r \neq q^2, q^2 \neq 1, r \neq 1, q, q^{-1} \neq r$.




when $q \in R_9 \cup R_8 \cup R_{13}$.



when $q \in R_5 \cup R_4$




when $q \in R_6 \cup R_8$.




, $q \in R_3$.

(26)




, $q \in R_3$.




, $q \in R_3$.

(28)




, $q \in R_3$.

(29)




when $\tilde{q}_{34} = q^2$ and $q^2 \in R_3$.

(30)




when $\tilde{q}_{34} = q^3$ and $q^8 = 1$.

(31)




when $\tilde{q}_{34} = q^{-1}$.

(32)




when $\tilde{q}_{34} = -q^{-1}$ and $q \in R_3$.

(33)




$q \in R_6$.

(34)




when $\tilde{q}_{34} = q^3$, $q^2 = q^{-6}$.

(35)



with $q \in R_3$. when $\tilde{q}_{34} = q^{-1}$ and $q^2 \neq 1$.

(36)



with $q \in R_3$. $\tilde{q}_{34} = -q^{-1}$.

2.12 GDD 1 of Row 7

$$\begin{array}{c} (1) \\ -1 \quad q \quad -1 \quad q^{-1}q \quad q^{-3} \quad q^3 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ q \notin R_2 \cup R_3. \end{array}$$

$$\begin{array}{c} (2) \\ -q \quad \frac{1}{q^2} \quad -1 \quad q^{-2}q^2 \quad q^{-6} \quad q^6 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ q^4 \neq 1, q \notin R_2 \cup R_3. \end{array}$$

$$\begin{array}{c} (3) \\ -1 \quad q \quad -1 \quad q^{-2}q^2 \quad q^{-6} \quad q^6 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ q \notin R_2 \cup R_3. \end{array}$$

$$\begin{array}{c} (4) \\ -1 \quad q \quad -1 \quad q^{-3}q^3 \quad q^{-9} \quad q^9 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ q^3, q \notin R_2 \cup R_3, \end{array}$$

(5)

$$\begin{array}{c} -q \quad \frac{1}{q^2} \quad -1 \quad q^{-3}q^3 \quad q^{-9} \quad q^9 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ q^2, q \notin R_2 \cup R_3, \end{array}$$

$$\begin{array}{c} (6) \\ q^{-1} \quad q \quad -1 \quad q^{-1}q \quad q^{-3} \quad q^3 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ q \notin R_2 \cup R_3, q \notin R_4 \end{array}$$

$$\begin{array}{c} (7) \\ r \quad r^{-1} \quad -1 \quad q^{-1}q \quad q^{-3} \quad q^3 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ q \notin R_2 \cup R_3. \end{array}$$

(8)

$$\begin{array}{c}
 q \quad q^{-1} \quad -1 \quad q^2 \quad q^{-2} \quad q^{-3} \quad q^3 \\
 \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \\
 \hline
 q \notin R_2 \cup R_3.
 \end{array}$$

$$\begin{array}{c}
 (9) \\
 q \quad q^{-1} \quad -1 \quad q^{-1}q \quad q^{-3} \quad q^3 \\
 \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \\
 \hline
 q \notin R_2 \cup R_3,
 \end{array}$$

$$\begin{array}{c}
 (10) \\
 q^{-2} \quad q^2 \quad -1 \quad q^{-1}q \quad q^{-3} \quad q^3 \\
 \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \\
 \hline
 q \notin R_2 \cup R_3.
 \end{array}$$

$$\begin{array}{c}
 (11) \\
 q^{-1} \quad -q^{-1} \quad -1 \quad -q \quad -q^{-1} \quad -q^{-3} \quad q^3 \\
 \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \\
 \hline
 , q \in R_3.
 \end{array}$$

$$\begin{array}{c}
 (12) \\
 -1 \quad q \quad -1 \quad -q^{-1}q \quad -q^{-3} \quad q^3 \\
 \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \\
 \hline
 , q \in R_3.
 \end{array}$$

$$\begin{array}{c}
 (13) \\
 q \quad -1 \quad -1 \quad -q^{-1}q \quad -q^{-3} \quad q^3 \\
 \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \\
 \hline
 , q \in R_3.
 \end{array}$$

$$\begin{array}{c}
 (14) \\
 \begin{array}{c}
 -1 \quad q^{-1} \quad q \quad q^{-1} \quad -1 \\
 \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \\
 \hline
 \begin{array}{c}
 \bullet \quad \bullet \quad \bullet \\
 \hline
 q^{-3} \quad q^3
 \end{array}
 \end{array} \\
 q \notin R_2 \cup R_3.
 \end{array}$$

2.13 GDD 2 of Row 7

$$\begin{array}{c}
 (1). \\
 q \quad q^{-1} \quad -1 \quad q \quad -1 \quad q^{-3} \quad q^3 \\
 \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \\
 \hline
 , q \in F^* \setminus \{1, -1\} \quad q \notin R_2 \cup R_3.
 \end{array}$$

$$\begin{array}{c}
 (2). \\
 q \quad q^{-2} \quad -1 \quad q^2 \quad -1 \quad q^{-6} \quad q^6 \\
 \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \\
 \hline
 q^2 \notin R_2 \cup R_3, \quad q^4 \neq 1,
 \end{array}$$

$$\begin{array}{c}
 (3). \\
 q^2 \quad q^{-2} \quad -1 \quad q \quad -1 \quad q^{-3} \quad q^3 \\
 \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \\
 \hline
 \end{array}$$

$$q \notin R_2 \cup R_3,$$

$$\begin{array}{c}
 (4). \\
 q^3 \quad q^{-3} \quad -1 \quad q \quad -1 \quad q^{-3} \quad q^3 \\
 \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \\
 \hline
 \end{array}$$

$$q \notin R_2 \cup R_3,$$

$$(5). \quad \begin{array}{ccccccc} -1 & q^{-1} & -1 & q & -1 & q^{-3} & q^3 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \\ \hline q & \notin R_3, & q \in F^* \setminus \{1, -1\}. \end{array}$$

$$(6). \quad \begin{array}{ccccccc} q & -q & -1 & -q^{-1} & -1 & -q^3 & -q^{-3} \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

, $q \in R_3$.

$$(7). \quad \begin{array}{ccccccc} & & q^3 & & & & \\ & & \bullet & & & & \\ & & | & & & & \\ & & q^{-3} & & & & \\ \bullet & & \bullet & & \bullet & & \\ \hline q & q^{-1} & -1 & q & -1 \end{array}$$

$q \notin R_2 \cup R_3$,

$$(8). \quad \begin{array}{ccccccc} & & q^6 & & & & \\ & & \bullet & & & & \\ & & | & & & & \\ & & q^{-6} & & & & \\ \bullet & & \bullet & & \bullet & & \\ \hline q & q^{-2} & -1 & q^2 & -1 \end{array}$$

when $q \in R_8 \cup R_{18}$

$$(9). \quad \begin{array}{ccccccc} & & q^3 & & & & \\ & & \bullet & & & & \\ & & | & & & & \\ & & q^{-3} & & & & \\ \bullet & & \bullet & & \bullet & & \\ \hline q^2 & q^{-2} & -1 & q & -1 \end{array}$$

$q \notin R_2 \cup R_3$,

$$(10). \quad \begin{array}{ccccccc} & & q^3 & & & & \\ & & \bullet & & & & \\ & & | & & & & \\ & & q^{-3} & & & & \\ \bullet & & \bullet & & \bullet & & \\ \hline q^3 & q^{-3} & -1 & q & -1 \end{array}$$


$q \notin R_2 \cup R_3$,

$$(11). \quad \begin{array}{ccccccc} & & q^3 & & & & \\ & & \bullet & & & & \\ & & | & & & & \\ & & q^{-3} & & & & \\ \bullet & & \bullet & & \bullet & & \\ \hline -1 & q^{-1} & -1 & q & -1 \end{array}$$

when $q \in R_4 \cup R_5 \cup R_6$.

$$(12). \quad \begin{array}{ccccccc} & & -q^{-3} & & & & \\ & & \bullet & & & & \\ & & | & & & & \\ & & -q^3 & & & & \\ \bullet & & \bullet & & \bullet & & \\ \hline q & -q & -1 & -q^{-1} & -1 \end{array}$$


, $q \in R_3$.



, $q \in R_3$.


(14)

and $q^2 = q^6$.




(15)

and $-1 = q^2$.




(16)

$q^3 = q^9$.



(17)


and $-1 = q^3$.



(18)



$q^6 \neq 1..$

(19)



$q^{12} \neq 1.$


(20)



when $q \in R_4$ or $q^3 = q^{-2}$.


2.14 GDD 3 of Row 7

(1).




when $q \in R_6$.


(2).




$q \in R_6$




$q \notin R_2 \cup R_3$.




, $q \in R_3$.




when $q \in R_6$.



when $q \in R_4 \cup R_6$.




when $q \in R_8$




when $q \in R_7$.


(9)




when $q \in R_6$.



$q \in R_{10}$



when $q \in R_{11}$.



$$q \in R_4 \cup R_7 \cup R_{10} \cup R_6.$$

$$(13) \begin{array}{c} q^3 \quad -1 \quad q^{-2} \\ \bullet \quad \bullet \quad \bullet \\ \text{---} \quad \diagup \quad \diagdown \\ r \quad r^{-1} \quad -1 \quad q^{-1} \quad q \end{array}$$

when $r = q^3$, $q \notin R_6$ or $r = q^6$, $q \notin R_{12}$ or $r = -q^{-3}$, $q \in R_9$ or $q^3 = -1$, or $r = -1$; $r \neq 1$, $r \neq q, q^{-1}$.

$$(14) \begin{array}{c} q^{-6} \quad -1 \quad q^4 \\ \bullet \quad \bullet \quad \bullet \\ \text{---} \quad \diagup \quad \diagdown \\ q \quad q^{-1} \quad -1 \quad q^2 \quad q^{-2} \end{array}$$

$$\text{when } q \in R_7 \cup R_{13} \cup R_{19} \cup R_{12}$$

$$(15) \begin{array}{c} q^3 \quad -1 \quad q^{-2} \\ \bullet \quad \bullet \quad \bullet \\ \text{---} \quad \diagup \quad \diagdown \\ q \quad q^{-1} \quad -1 \quad q^{-1} \quad q \end{array}$$

$$\text{when } q \in R_5 \cup R_8 \cup R_6 .$$

$$(16) \begin{array}{c} q^3 \quad -1 \quad q^{-2} \\ \bullet \quad \bullet \quad \bullet \\ \text{---} \quad \diagup \quad \diagdown \\ q^{-2} \quad q^2 \quad -1 \quad q^{-1} \quad q \end{array}$$

$$\text{when } q \in R_4 \cup R_5 \cup R_8 \cup R_{11} \cup R_6 .$$

$$(17) \begin{array}{c} -q^{-3} \quad -1 \quad q^2 \\ \bullet \quad \bullet \quad \bullet \\ \text{---} \quad \diagup \quad \diagdown \\ q^{-1} \quad -q^{-1} \quad -1 \quad -q \quad -q^{-1} \end{array}$$


$$\text{when } q \in R_3.$$

$$(18) \begin{array}{c} -q^3 \quad -1 \quad q^{-2} \\ \bullet \quad \bullet \quad \bullet \\ \text{---} \quad \diagup \quad \diagdown \\ -1 \quad q \quad -1 \quad -q^{-1} \quad -q \end{array}$$

$$\text{when } q \in R_3.$$


$$(19) \begin{array}{c} q^{-2} \quad q \quad q^{-1} \\ \bullet \quad \bullet \quad \bullet \\ \text{---} \quad \diagup \quad \diagdown \\ q^3 \quad q^{-3} \quad -1 \quad q^3 \quad -1 \end{array}$$

$$, q^3 \in F^* \setminus \{1, -1\}. \quad q \in R_5 \cup q \in R_4.$$




when $q^3 \in F^* \setminus \{1, -1\}$. $q^8 = 1$, $q \in R_{18}$.

(21).




when $q \in R_5$, $q \in R_{12}$.




when $q^{11} = 1$ or $q^{12} = -1$

(23).




$q \notin R_6 \cup R_3$



when $q \in R_5$.


(25).



when CON 1, CON 2, CON 3 and CON 4 are satisfied, where

CON 1: $, q^3, s^3, r^3 \in F^* \setminus \{1\}, q^3 s^3 r^3 = 1, q^3 \neq s^3, q^3 \neq r^3, r^3 \neq s^3$. and CON 2: $q = r^3$, or $r^6 = q$, or $q \neq r^9$, or $q = -r^3$, or $q^3 \in R_3$, or $r^3 = -1$. CON 3: $s^3 = q^2$. $q \notin R_5$. CON 4: $q \notin R_2 \cup R_3$.

(26)



$\tilde{q}_{14} = q^{-2}$ and $q \in R_5$.

2.15 GDD 4 of Row 7

(1).
 $q^6 \quad q^{-6} \quad q^3 \quad q^{-3} \quad -1 \quad q^2 \quad -q^{-1}$
 $\bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet$
 $q \notin R_3 \cup R_2, q^3 \in F^* \setminus \{1, -1\}$.

(2).
 $q^9 \quad q^{-9} \quad q^3 \quad q^{-3} \quad -1 \quad q^2 \quad -q^{-1}$
 $\bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet$
 $q^3 \notin R_3 \cup R_2, q \notin R_3 \cup R_2$,

(3).
 $-1 \quad q^{-3} \quad q^3 \quad q^{-3} \quad -1 \quad q^2 \quad -q^{-1}$
 $\bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet$
 $q \notin R_3 \cup R_2, q^3 \in F^* \setminus \{1, -1\}$.

(4).
 $-q^{-3} \quad q^3 \quad q^{-3} \quad -1 \quad q^2 \quad -q^{-1}$
 $\bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet$
 $q \notin R_3 \cup R_2, -q^3 \in R_3$.

(5).
 $-1 \quad q^3 \quad q^3 \quad q^{-3} \quad -1 \quad q^2 \quad -q^{-1}$
 $\bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet$
 $q \notin R_3 \cup R_2, q^3 \in R_3$.

(6).

$$\begin{array}{c} -q^3 - q^{-3}q^3 \quad q^{-3}-1 \quad q^2 \quad -q^{-1} \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ q \notin R_3 \cup R_2, q^3 \in R_3. \end{array}$$

$$\begin{array}{c} (7). \\ -1 \quad -1 \quad q^3 \quad q^{-3}-1 \quad q^2 \quad -q^{-1} \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ q \notin R_3 \cup R_2, q^3 \in R_3. \end{array}$$

$$\begin{array}{c} (8). \\ -1 \quad q^3 \quad -q^3 - q^{-3}-1 \quad q^2 \quad q^{-1} \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ q \notin R_3 \cup R_2, q^3 \in R_3. \end{array}$$


$$\begin{array}{c} (9). \\ -1 \quad -q^3 \quad q^3 \quad q^{-3}-1 \quad q^2 \quad -q^{-1} \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ q \notin R_3 \cup R_2, q^3 \in R_3. \end{array}$$

$$\begin{array}{c} (10). \\ q^3 \quad q^{-3} \quad q^3 \quad q^{-3}-1 \quad q^2 \quad -q^{-1} \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ q \notin R_3 \cup R_2, q^3 \notin R_2. \end{array}$$


$$\begin{array}{c} (11). \\ q^3 \quad q^{-6} \quad q^6 \quad q^{-6}-1 \quad q^4 \quad -q^{-2} \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ q^2 \notin R_3 \cup R_2, q^3 \notin R_2 \cup R_4, \end{array}$$

(12)

$$q \notin R_3 \cup R_2.$$




(13)



when $q \in R_5 \cup R_7 \cup R_9 \cup R_6 \cup R_4$.

(14)

$$q \notin R_3 \cup R_2,$$



2.16 GDD 1 of Row 8

$$\begin{array}{c} (1). \\ q^3 \quad q^{-3} \quad q \quad q^{-1}-1 \quad q \quad q^{-1} \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ \text{when } q \notin R_4 \end{array}$$

$$\begin{array}{c} (2). \\ -q \quad -q^{-1}q \quad q^{-1}-1 \quad q \quad q^{-1} \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ , q \in R_3. \end{array}$$

$$(3). \quad \begin{array}{ccccccc} -1 & & q & & q^{-1} & -1 & q & & q^{-1} \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$

, $q \in R_3$.

$$(4). \quad \begin{array}{ccccccc} -1 & & q & & -q & -q^{-1} & -1 & -q & & -q^{-1} \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$

, $q \in R_3$.

$$(5). \quad \begin{array}{ccccccc} -1 & & q & & q^{-1} & -1 & q & & q^{-1} \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$

, $q \in R_3$.

$$(6) \quad \begin{array}{ccccccc} & & & & q^{-1} & & & & \\ & & & & q & & & & \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \\ -1 & q & -1 & q^{-1} & q & & & & \end{array}$$

when $q \in R_3 \cup R_4$.

$$(7) \quad \begin{array}{ccccccc} & & & & q^{-2} & & & & \\ & & & & q^2 & & & & \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \\ -1 & q & -1 & q^{-2} & q^2 & & & & \end{array}$$

when $q \notin R_3$

$$(8) \quad \begin{array}{ccccccc} & & & & q^{-3} & & & & \\ & & & & q^3 & & & & \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \\ -1 & q & -1 & q^{-3} & q^3 & & & & \end{array}$$

when $q \in R_4 \cup R_6 \cup R_5$.


$$(9) \quad \begin{array}{ccccccc} & & & & q^{-3} & & & & \\ & & & & q^3 & & & & \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \\ -q & q^2 & -1 & q^{-3} & q^3 & & & & \end{array}$$

when $q \in R_5 \cup R_6$.


$$(10) \quad \begin{array}{ccccccc} & & & & q^{-1} & & & & \\ & & & & q & & & & \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \\ r & r^{-1} & -1 & q^{-1} & q & & & & \end{array}$$

$q^2 \neq 1, r \neq 1, r \neq q, q^{-1}$.

(11)




$q \notin R_2 \cup R_3.$




when $q \notin R_2 \cup R_3.$

(13)




and $q^{-1} = q^2.$

(14)



$q^2 \neq 1.$

(15)



and $q^2 = q^{-1}$

2.17 GDD 2 of Row 8

(1).
 q^3
 $q^{-3} \quad -1 \quad q \quad -1 \quad q^{-1} \quad -1$

 when $q \notin R_4, q^4, q^3 \neq 1.$

(2).
 $-q \quad -q^{-1} \quad -1 \quad q \quad -1 \quad q^{-1} \quad -1$

 $q \in R_3.$

(3).
 $-1 \quad -1 \quad -1 \quad q \quad -1 \quad q^{-1} \quad -1$

 $q \in R_3.$

(4).

$$\begin{array}{ccccccc} -1 & -q & -1 & q & -1 & q^{-1} & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

$q \in R_3.$

$$(5). \begin{array}{ccccccc} -1 & q & -1 & -q & -1 & -q^{-1}-1 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

$q \in R_3.$

$$(6). \begin{array}{ccccccc} & & -1 & & & & \\ & & \bullet & & & & \\ & & q^{-1} & & & & \\ \bullet & & \bullet & & \bullet & & \\ q & q^{-1} & -1 & q & -1 & & \end{array}$$

when $q \in R_4.$

$$(7). \begin{array}{ccccccc} & & -1 & & & & \\ & & \bullet & & & & \\ & & q^{-1} & & & & \\ \bullet & & \bullet & & \bullet & & \\ q^2 & q^{-2} & -1 & q & -1 & & \end{array}$$

when $q \in R_4 \cup R_5 .$

$$(8). \begin{array}{ccccccc} & & -1 & & & & \\ & & \bullet & & & & \\ & & q^{-1} & & & & \\ \bullet & & \bullet & & \bullet & & \\ q^3 & q^{-3} & -1 & q & -1 & & \end{array}$$

when $q \in R_4 \cup R_5 \cup R_6.$

$$(9). \begin{array}{ccccccc} & & -1 & & & & \\ & & \bullet & & & & \\ & & q^{-1} & & & & \\ \bullet & & \bullet & & \bullet & & \\ -1 & q^{-1} & -1 & q & -1 & & \end{array}$$

, $q \in F^* \setminus \{1, -1\}.$

$$(10). \begin{array}{ccccccc} & & -1 & & & & \\ & & \bullet & & & & \\ & & q^{-1} & & & & \\ \bullet & & \bullet & & \bullet & & \\ -1 & q & -1 & q & -1 & & \end{array}$$

, $q \in R_3.$


$$(11). \begin{array}{ccccccc} & & -1 & & & & \\ & & \bullet & & & & \\ & & q^{-1} & & & & \\ \bullet & & \bullet & & \bullet & & \\ -1 & -1 & -1 & q & -1 & & \end{array}$$

, $q \in R_3.$

(12)


and $q^{-1} = q^2$

(13)




when $q \in R_3$.

(14)




and $q = q^{-3}$

(15)




$q \in R_3$.

(16)




and $q^2 = q^{-3}$

(17)





and $q^2 = -1$.

(18)




and $q^3 = q^{-3}$

(19)




and $q^3 = -1$.

(20)




$q^3 = -1$,

(21)




$q^2 \neq 1$,

(22)




when $q \in R_3$

(23)




when $q \in R_3$

(24)




when $q \in R_3$

(25)




$q \in R_6$.

(26)




when $q \in R_3$

(27)




when $q \in R_3$

(28)




when $q \in R_3$

(29)




when $q \in R_3$

(30)



when $q \in R_3$

(31)



$q \in R_6$.

2.18 GDD 3 of Row 8

$$\overset{(1)}{\begin{array}{ccccccc} -1 & q & -1 & q^{-2}q^2 & q^{-2} & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}}$$

when $q \notin R_2 \cup R_3$

$$\overset{(2)}{\begin{array}{ccccccc} -1 & q & -1 & q^{-3}q^3 & q^{-3} & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}}$$

$q \notin R_2 \cup R_3$

$$\overset{(3)}{\begin{array}{ccccccc} -q & q^2 & -1 & q^{-3}q^3 & q^{-3} & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}}$$

$$q \notin R_2 \cup R_3$$

$$(4) \quad r \quad \begin{array}{cccc} r^{-1} & -1 & q^{-1}q & q^{-1} & -1 \end{array}$$

$$\text{when } r \neq q^2, r \neq 1, r \neq q, q^{-1}.$$

$$(5) \quad q^{-1} \quad \begin{array}{cccc} q & -1 & q^{-2}q^2 & q^{-2} & -1 \end{array}$$

$$, q \notin R_3 \cup R_2.$$

$$(6) \quad q \quad \begin{array}{cccc} q^{-1} & -1 & q^{-1}q & q^{-1} & -1 \end{array}$$

$$q \notin R_4 \cup R_3.$$

$$(7) \quad -1 \quad \begin{array}{cccc} q & -1 & -q^{-1}-q & -q^{-1}-1 \end{array}$$

$$, q \in R_3.$$

$$(8) \quad q \quad \begin{array}{cccc} -1 & -1 & -q^{-1}-q & -q^{-1}-1 \end{array}$$

$$, q \in R_3.$$


$$(9) \quad \begin{array}{c} -1 \\ q^{-1} \\ \bullet \\ \bullet \end{array} \quad \begin{array}{cccc} q^2 & q^{-2} & q & q^{-1}-1 \end{array}$$

$$\text{when } q \notin R_2 \cup R_3$$


$$(10) \quad \begin{array}{c} -1 \\ q^{-1} \\ \bullet \\ \bullet \end{array} \quad \begin{array}{cccc} q^3 & q^{-3} & q & q^{-1}-1 \end{array}$$

$$, q \in F^* \setminus \{1, -1\}, q \notin R_3.$$


$$(11).$$




, $q \in F^* \setminus \{1, -1\}$.




, $q \in R_3$.




, $q \in R_3$.




, $q \in R_3$.




, $q \in R_3$.




, $q \in R_3$.



, $q \in R_3$.




when $q \notin R_2 \cup R_3$




$q^4 \neq 1$.

(20)




, $q \in R_3$.

(21)




, $q \in R_3 \cup R_4$.

(22)




when $q^3 = q^{-1}$, or $q^3 = -1$

(23)




$r = -1$, $q \in R_3 \cup R_4 \cup R_6$

(24)




and $-1 = q^{-2}$

(25)




when $q \in R_6$

(26)




and $r^2 = -q^{-1}$, $q \in R_6$

(27)




and $r^2 = -q^{-1}$, $q \in R_6$

(28)




$q^2 \neq 1$.

(29)




when $q^3 = q^{-2}$.

(30)




and $-1 = q^{-2}$

(31)




when $q \in R_3 \cup R_4 \cup R_6$, $-1 = r^2$.

(32)




and $-1 = q^2$

(33)




$$q \in R_3]]$$

$$(34)$$




$$q \in R_8.$$

$$(35)$$




$$\text{and } -1 = r^3, q \in R_4 \cup R_6, r \neq 1, r \neq q, q^{-1}.$$

$$(36)$$




$$\text{and } -1 = q^3$$

$$(37)$$




$$\text{and } -1 = q^{-6}$$

$$(38)$$




$$\text{and } q^{-3} = -q^{-1}$$

$$(39)$$




$$\text{and } r^3 = -q^{-1}, q \in R_{18} \cup R_6. r \neq 1, r \neq q, q^{-1}.$$

$$(40)$$




$$q^2 \neq 1.$$

(41)




and $r' = r$, $q^2 \neq 1$, $r \neq 1$, $r \neq q, q^{-1}$.

(42)




and $q^{-1} = r^2$, $q^2 \neq 1$, $r \neq 1$, $r \neq q, q^{-1}$.

(43)




and $r = q^{-2}$, $q \notin R_2 \cup R_3$

(44)




and $-q = r$ with $q \in R_6$

(45)




and $r = -1$, $q \in R_6$

(46)




when $q \in R_6$, $-q = r$

(47)




$q \notin R_2 \cup R_3$.

(48)




and $-q^2 = q^{-1}$

(49)




$q \notin R_2 \cup R_3$

(50)




$q \notin R_2 \cup R_3$

(51)




when $q \in R_6$

(52)



$q \notin R_3$


(53)



$q \notin R_6$

2.19 GDD 4 of Row 8

2.20 GDD 1 of Row 9



when $r \neq q^{-3}$, $q^2 \neq 1$. $r \neq 1$, $r \neq q, q^{-1}$.

$$\begin{array}{c} (2). \\ q^3 \quad q^{-3} \quad q \quad q^{-1} \quad -1 \quad r^{-1} \quad r \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ , q \in F^* \setminus \{1, -1\}, q \notin R_3. r \neq 1, r \neq q, q^{-1}. \end{array}$$

$$\begin{array}{c} (3). \\ -1 \quad q^{-1} \quad q \quad q^{-1} \quad -1 \quad r^{-1} \quad r \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ \text{when } r \neq q^{-2}, q \in F^* \setminus \{1, -1\} r \neq 1, r \neq q, q^{-1}. \end{array}$$

$$\begin{array}{c} (4). \\ q \quad -q \quad -q^{-1} \quad -q \quad -1 \quad -r \quad -r^{-1} \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ , q \in R_3. r \neq 1, r \neq q, q^{-1}. \end{array}$$

$$\begin{array}{c} (5). \\ -1 \quad q \quad q \quad q^{-1} \quad -1 \quad r^{-1} \quad r \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ , q \in R_3. r \neq 1, r \neq q, q^{-1}. \end{array}$$

$$\begin{array}{c} (6). \\ -q \quad -q^{-1} \quad q \quad q^{-1} \quad -1 \quad r^{-1} \quad r \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ \text{when } r \neq -q, q \in R_3. r \neq 1, r \neq q, q^{-1}. \end{array}$$

$$\begin{array}{c} (7). \\ -1 \quad -1 \quad q \quad q^{-1} \quad -1 \quad r^{-1} \quad r \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ , q \in R_3. r \neq 1, r \neq q, q^{-1}. \end{array}$$

$$\begin{array}{c} (8). \\ -1 \quad q \quad -q \quad -q^{-1} \quad -1 \quad -r^{-1} \quad -r \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ , q \in R_3. r \neq 1, r \neq q, q^{-1}. \end{array}$$

$$\begin{array}{c} (9). \\ -1 \quad -q \quad q \quad q^{-1} \quad -1 \quad r^{-1} \quad r \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ , q \in R_3. r \neq 1, r \neq q, q^{-1}. \end{array}$$

$$\begin{array}{c} (10). \\ q^{-1} \quad q^{-1} \quad q \quad -1 \quad r \quad r^{-1} \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array}$$

when $q \neq -r^{-1}$ and $q^2 \neq 1$; $q^3 \neq r^2$ or $q^6 = 1$; $q^2 \neq r^{-1}$; $r \neq 1$, $r \neq q, q^{-1}$; $q^2 \neq 1$.

$$\begin{array}{c} (11) \\ q \quad q^{-2} \quad q^2 \quad q^{-2} \quad -1 \quad r^{-2} \quad r^2 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ q^4 \neq 1, r \neq 1, r \neq q, q^{-1}. \end{array}$$

$$\begin{array}{c} (12) \\ -1 \quad q \quad -1 \quad -1 \quad -1 \quad r^{-1} \quad r \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ \text{when } r \neq -1, q \in R_3. r \neq 1, r \neq q, q^{-1}. \end{array}$$

(13)

$$q^{-1} \quad -q^{-1} \quad 1 \quad -1 \quad -1 \quad r^{-1} \quad r$$

when $q \in R_3$, $r \neq -1$. $r \neq 1$, $r \neq q$, q^{-1} .

$$(14) \quad \begin{array}{c} r \\ \bullet \\ r^{-1} \\ | \\ \bullet \\ -1 \quad q \quad -1 \quad q^{-1} \quad q \\ \bullet \end{array}$$

when $r = -q$, and $q \in R_3$ or $r = q^3$, and $q^4 \neq 1$ or $r = q^2$ and $r \notin R_3$. $r \neq 1$, $r \neq q$, q^{-1} .

$$(15) \quad \begin{array}{c} r^2 \\ \bullet \\ r^{-2} \\ | \\ \bullet \\ -q^{-1}q^2 \quad -1 \quad q^{-2}q^2 \\ \bullet \end{array}$$

when $r^{-2} = q^{-3}$ or $r^{-2} = q^2$, $q \notin R_3$; $r \neq 1$, $r \neq q$, q^{-1} ; $q^2 \neq 1$.

$$(16) \quad \begin{array}{c} r^2 \\ \bullet \\ r^{-2} \\ | \\ \bullet \\ -1 \quad q \quad -1 \quad q^{-2}q^2 \\ \bullet \end{array}$$

when $r^2 = q$, with $q^2, q^3 \neq 1$; or $r^2 = q^3$, $q^3, q^5 \neq 1$; or $r^2 = -q, q \in R_3$; or $r^2 = -1$, $q \in R_3 \cup R_6$. $r^2 \neq 1$, $r^2 \neq q^2, q^{-2}$; $q^4 \neq 1$.

$$(17) \quad \begin{array}{c} r^3 \\ \bullet \\ r^{-3} \\ | \\ \bullet \\ -1 \quad q \quad -1 \quad q^{-3}q^3 \\ \bullet \end{array}$$

when $r^3 = q$ or $r^3 = q^2$, $q^5 \neq 1$ or $q^5 = -1$, $q \in R_4$; $r^3 = q$ or $r^3 = q^2$, $q^5 \neq 1$ or $q^5 = -1$, $q \in R_4$; $r \neq 1$; $r^3 \neq q^3, q^{-3}$. $r^3 \neq 1$; $q \notin R_2 \cup R_3$.

$$(18) \quad \begin{array}{c} r^3 \\ \bullet \\ r^{-3} \\ | \\ \bullet \\ -q^{-1}q^2 \quad -1 \quad q^{-3}q^3 \\ \bullet \end{array}$$


when $r^3 = q^2$, $q \notin R_5$, $q \notin R_2 \cup R_3$; $r^3 \neq q^3, q^{-3}$. $r^3 \neq 1$.

$$(19) \quad \begin{array}{c} r \\ \bullet \\ r^{-1} \\ | \\ \bullet \\ q^{-1} \quad q \quad -1 \quad q^{-1} \quad q \\ \bullet \end{array}$$


$r \neq 1$, $r \neq q, q^{-1}$; $q^2 \neq 1$.

$$(20) \quad \begin{array}{c} r \\ \bullet \\ r^{-1} \\ | \\ \bullet \\ s \quad s^{-1} \quad -1 \quad q^{-1} \quad q \\ \bullet \end{array}$$


$r \neq 1, r \neq q, q^{-1}; q^2 \neq 1.$




$q \notin R_2 \cup R_3, r \neq 1, r \neq q, q^{-1}.$




$q \notin R_2 \cup R_3, r \neq 1, r \neq q, q^{-1}.$




$, q \notin R_3, r \neq 1, r \neq q, q^{-1}.$




$q \notin R_2 \cup R_3, r \neq 1, r \neq q, q^{-1}.$



when $-r = -1, q \in R_3, r \neq 1, r \neq q, q^{-1}.$




when $r = -q; \text{ or } r = -q^2; \text{ or } r = 1, q \in R_3; r \neq 1, r \neq q, q^{-1}.$




when $r = -q, q \in R_3 \text{ or } r = -1, q \in R_3; r \neq 1, r \neq q, q^{-1}.$


(28)

and $r^2 = q^2$, $r \neq 1$, $r \neq q, q^{-1}$. $q^2 \neq 1$.


(29)

and $r = q^2$, $r \neq 1$, $r \neq q, q^{-1}$. $q^2 \neq 1$.


(30)

and $-1 = q^3$, $r \neq 1$, $r \neq q, q^{-1}$.


(31)

 $r \neq 1$, $r \neq q, q^{-1}$. $q^2 \neq 1$.


(32)

when $q = -r$ and $r \in R_3$; $r \neq 1$, $r \neq q, q^{-1}$. $q^2 \neq 1$.


(33)

when $-r^{-1} = q^{-1}$, $r \in R_6$; $r \neq 1$, $r \neq q, q^{-1}$. $q^2 \neq 1$.

(34)


when $r = -q$; $r \in R_3$. $r \neq 1$, $r \neq q, q^{-1}$.

(35)




and $s \in R_3$, $q = s^2$ or $q = s$ or $q = s^3$ or $q = -s$; $r \neq 1$, $r \neq q, q^{-1}$.

(36)




when $s = q^{-1}$ with $q, s \in R_3$; $r \neq 1$, $r \neq q, q^{-1}$.

(37)




and $-q = r$: $r \neq 1$, $r \neq q, q^{-1}$. $q \in R_3$.

(38)




when $r = q^2$; $r \neq 1$, $r \neq q, q^{-1}$; $q^2 \neq 1$. $r^4 \neq 1$.

(39)



and $s^{-1} = q$; $s \in R_3$; $q^2 \neq 1$.


(40)



when $s = t$ or $s^{-1} = t$, $s, t \in R_3$.


2.21 GDD 2 of Row 9

2.22 GDD 3 of Row 9




when $q = s^2$, $s^3 \neq 1$, or $q = s^3$, with $s^2, s^3, s^4 \neq 1$, or $q = -s$, $s \in R_3$ or $s = -1$, $q^2, r^2 \neq 1$; $q, r, s \in F^* \setminus \{1\}$, $qrs = 1$, $q \neq r$, $q \neq s$, $s \neq r$.

(2).




, $q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 5.

when $s = -q$. $q, r, s \in F^* \setminus \{1\}$, $qrs = 1$, $q \neq r$, $q \neq s$, $s \neq r$.




when $q^2 = s$, or $s^2 = q^2$, or $s^3 = q^2$, or $q^2 = -s$, $s \in R_3$ or $s = -1$ or $q^2 = -1$, $s \in R_2 \cup R_3 \cup R_4 \cup R_6$, $q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 6; , $q, r, s \in F^* \setminus \{1\}$, $qrs = 1$, $q \neq r$, $q \neq s$, $s \neq r$.

(4).




when $q = s^2$, $1 \neq s^3$, or $q = s^3$ with $s^2, s^3, s^4 \neq 1$, or $s = -1$ or $s = -q$, $s \in R_3$; $q = r^2$, $1 \neq r^3$, or $q = r^3$ with $r^2, r^3, r^4 \neq 1$, or $r = -1$ or $r = -q$, $r \in R_3$; , $q, r, s \in F^* \setminus \{1\}$, $qrs = 1$, $q \neq r$, $q \neq s$, $s \neq r$.




when $q^3 = s$ or $s^2 = q^3$, or $q^3 = s^3$ or $-s = q^3$, $s \in R_3$ or $q^3 = -1$, $s \in R_2 \cup R_3 \cup R_4 \cup R_6$; $q \in F^* \setminus \{1, -1\}$. $q \notin R_3$; , $q, r, s \in F^* \setminus \{1\}$, $qrs = 1$, $q \neq r$, $q \neq s$, $s \neq r$.

(6).




when $s^3 = q$ or $s^6 = q$ or $s^9 = q$ or $-s^3 = q$, $s \in R_9$ or $s^3 = -1$; $r^3 = q^2$; , $q, r, s \in F^* \setminus \{1\}$, $qrs = 1$, $q \neq r$, $q \neq s$, $s \neq r$.

(7).



when $s = -1$, $q \in R_3 \cup R_4 \cup R_6$ or $s = -q$ with $s \in R_3$ or $q \in R_3$; $, q, r, s \in F^* \setminus \{1\}$, $qrs = 1$, $q \neq r$, $q \neq s$, $s \neq r$.

(8).




$, q, r, s \in F^* \setminus \{1\}$, $qrs = 1$, $q \neq r$, $q \neq s$, $s \neq r$. GDD 3 of Row 9.

$q', r', s' \in F^* \setminus \{1\}$, $q'r's' = 1$, $q' \neq r'$, $q' \neq s'$, $s' \neq r'$. $q = q'$

when $s' = s$, $s \in R_3$ or $s' = s^{-1}$, or $s' = -s$ with $s \in R_3 \cup R_6$, or $s' = -1$, $s \in R_2 \cup R_3 \cup R_4 \cup R_6$ or $s = -1$, $s' \in R_2 \cup R_3 \cup R_4 \cup R_6$; $r' = r$, $r \in R_3$ or $r' = r^{-1}$ or $r' = -r$ with $r \in R_3 \cup R_6$, or $r' = -1$, $r \in R_2 \cup R_3 \cup R_4 \cup R_6$ or $r = -1$, $r' \in R_2 \cup R_3 \cup R_4 \cup R_6$; $, q, r, s \in F^* \setminus \{1\}$, $qrs = 1$, $q \neq r$, $q \neq s$, $s \neq r$.


(9).



$, q \in F^* \setminus \{1, -1\}$, $q^3 \neq 1$. GDD 3 of Row 10.


when $q = -s$ with $q \in R_3 \cup R_6$, or $s = -1$, $q \in R_3 \cup R_4 \cup R_6$; $r = q^{-2}$ with $r \in R_3$, or $r^{-1} = q^{-2}$ or $-r = q^{-2}$ with $r \in R_3 \cup R_6$, or $-1 = q^{-2}$, $r \in R_2 \cup R_3 \cup R_4 \cup R_6$. or $-1 = r$, $q^{-2} \in R_2 \cup R_3 \cup R_4 \cup R_6$; $, q \in F^* \setminus \{1, -1\}$, $q^3 \neq 1$; $q, r, s \in F^* \setminus \{1\}$, $qrs = 1$, $q \neq r$, $q \neq s$, $s \neq r$.

(10).




when $q = s^2$ or $q = -s^{-2}$, $q \in R_6$ or $-1 = s^{-2}$, $q \in \cup R_4 \cup R_6$; $q = r^2$ or $q = -r^{-2}$, $q \in R_6$ or $-1 = r^{-2}$, $q \in R_4 \cup R_6$; , $q \in F^* \setminus \{1, -1\}$, $q^3 \neq 1$; $q, r, s \in F^* \setminus \{1\}$, $qrs = 1$, $q \neq r, q \neq s, s \neq r$.


(11).




when $q = -s$ or $s = -1$; $-r = q$ or $r = -1$; $q \in R_3$; $q, r, s \in F^* \setminus \{1\}$, $qrs = 1$, $q \neq r, q \neq s, s \neq r$.




when $q = -s^{-1}$, $q = s^{-2}$, $-s^{-3} = q$; $q \in R_3$; $q, r, s \in F^* \setminus \{1\}$, $qrs = 1$, $q \neq r, q \neq s, s \neq r$.



when $q = s$ (omitting) or $s = q^{-1}$ (omitting), or $s = -q$ ($q \in R_3$) or $s = -1$ ($q \in R_3$); $q \in R_3$; , $q, r, s \in F^* \setminus \{1\}$, $qrs = 1$, $q \neq r, q \neq s, s \neq r$.




when $s = -q$ or $-q = s^2$ or $-q = s^3$, $s^3 \neq 1$, or $s = -1$; $q \in R_3$; $q, r, s \in F^* \setminus \{1\}$, $qrs = 1$, $q \neq r, q \neq s, s \neq r$.




when $t = s$, or $t = s^{-1}$, or $t = -s$, or $s = -1$; , $t \in R_3$; $q, r, s \in F^* \setminus \{1\}$, $qrs = 1$, $q \neq r, q \neq s, s \neq r$.

(16).




when $s \in R_2 \cup R_3 \cup R_4 \cup R_6$; $q \in R_3$; $q, r, s \in F^* \setminus \{1\}$, $qrs = 1$, $q \neq r$, $q \neq s$, $s \neq r$.

(17).




when $q = -s^{-1}$ or $s = -1$; $q \in R_3$; $q, r, s \in F^* \setminus \{1\}$, $qrs = 1$, $q \neq r$, $q \neq s$, $s \neq r$.

(18).




when $q = -s$ or $q = -s^{-1}$ or $s = -1$; $q \in R_3$; $q, r, s \in F^* \setminus \{1\}$, $qrs = 1$, $q \neq r$, $q \neq s$, $s \neq r$.

(19).




when $q = s^2$ or $q = s^3$; $r = -q^{-1}$; $q \in R_3$; $q, r, s \in F^* \setminus \{1\}$, $qrs = 1$, $q \neq r$, $q \neq s$, $s \neq r$.

(20).




when $s = -q$; $q \in R_3$; $q, r, s \in F^* \setminus \{1\}$, $qrs = 1$, $q \neq r$, $q \neq s$, $s \neq r$.

(21)




when $\tilde{q}_{34} = r^{-1}s^{-1}$. $s = r'$. $s' = r$; or $\tilde{q}_{34} = -r^{-1}s^{-1}$. $r' = -s$, $s' = -r$ or $s' = s$, $r' = r$. $\tilde{q}_{34} = s^{-2}$. $s = -r$; $q, r, s \in F^* \setminus \{1\}$, $qrs = 1$, $q \neq r$, $q \neq s$, $s \neq r$; $q, r', s' \in F^* \setminus \{1\}$, $qr's' = 1$, $q \neq r'$, $q \neq s'$, $s' \neq r'$.

(22)




when $\tilde{q}_{34} = r$. $q = -r$. $q^3, q^2 \neq 1$; $q, r, s \in F^* \setminus \{1\}$, $qrs = 1$, $q \neq r$, $q \neq s$, $s \neq r$.
(23)




with $q^3, q^2 \neq 1$, $q^{-2} = \tilde{q}_{34}$ and $s^{-2} = q$. $q^2, r^2, s^2 \in F^* \setminus \{1\}$, $q^2 r^2 s^2 = 1$, $q^2 \neq r^2$, $q^2 \neq s^2$, $s^2 \neq r^2$.

2.23 GDD 4 of Row 9


2.24 GDD 1 of Row 10




when $q \notin R_4 \cup R_2 \cup R_3$;



when $q \notin R_2 \cup R_3$;



when $q \notin R_4 \cup R_3 \cup R_2$.



, $q \in R_3$.

(5).

$$\begin{array}{ccccccc} -1 & & -q & -q^{-1} & -1 & & -q^{-1} & -q \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \end{array}$$

, $q \in R_3$

(6).

$$\begin{array}{ccccccc} q & & q^{-1} & & q & & q^{-1} & -1 & q^{-1} & & q \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \end{array}$$

when $q \notin R_{10} \cup R_4 \cup R_3 \cup R_6 \cup R_2$

(7)

$$\begin{array}{ccccccc} q & & q^{-2} & & q^2 & & q^{-2} & -1 & q^{-2} & & q^2 \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \end{array}$$

$q \notin R_4 \cup R_2 \cup R_3$.

(8)

$$\begin{array}{ccccccc} & & & & q^2 & & & & & & & \\ & & & & \bullet & & & & & & & \\ & & & & | & & & & & & & \\ & & & & q^{-2} & & & & & & & \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \\ -q^{-1}q^2 & & -1 & & q^{-2}q^2 & & & & & & & \end{array}$$

$q^4 \neq 1$.

(9)

$$\begin{array}{ccccccc} & & & & q^2 & & & & & & & \\ & & & & \bullet & & & & & & & \\ & & & & | & & & & & & & \\ & & & & q^{-2} & & & & & & & \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \\ -1 & & q & & -1 & & q^{-2}q^2 & & & & & \end{array}$$

$q \notin R_2 \cup R_3$.

(10)


$$\begin{array}{ccccccc} & & & & q^3 & & & & & & & \\ & & & & \bullet & & & & & & & \\ & & & & | & & & & & & & \\ & & & & q^{-3} & & & & & & & \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \\ -1 & & q & & -1 & & q^{-3}q^3 & & & & & \end{array}$$

$q \notin R_2 \cup R_3$


(11)

$$\begin{array}{ccccccc} & & & & q^3 & & & & & & & \\ & & & & \bullet & & & & & & & \\ & & & & | & & & & & & & \\ & & & & q^{-3} & & & & & & & \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \\ -q^{-1}q^2 & & -1 & & q^{-3}q^3 & & & & & & & \end{array}$$


$q \notin R_2 \cup R_3, q^3 \notin R_2 \cup R_3$




$$q \notin R_2 \cup R_3$$




$$q \notin R_2 \cup R_3, q^2 \notin R_2 \cup R_3$$




$$q \notin R_2 \cup R_3$$




$$q \notin R_2 \cup R_3$$



$$-q \notin R_2 \cup R_3$$




$$q \in R_3.$$




$$q \in R_3.$$

(19)




$q \notin R_2 \cup R_3$

(20)




$q \notin R_2 \cup R_3$


2.25 GDD 2 of Row 10




$q \notin R_2 \cup R_3$




$q \notin R_2 \cup R_3$.




, $q \in R_3$,




, $q \in R_3$




$q^2 \notin R_2 \cup R_3$




when $q \in R_4 \cup R_5$




when $q \in R_7$.




when $q \in R_5 \cup R_7$.




when $q \in R_7 \cup R_8 \cup R_9 \cup R_{14} \cup R_{10}$.




when $q \in R_8$, $q^{-6} = q^3$.




$q \notin R_2 \cup R_3$.




$q \notin R_2 \cup R_3$. $r \neq 1$, $r \neq q, q^{-1}$.




when $q^4 = -1$, $q^2 \notin R_3 \cup R_2$.



$q \notin R_2 \cup R_3$.




$q \notin R_2 \cup R_3.$




$, q \in R_3.$

(17)




$-1 = q^{-2}$

(18)




and $q^{-2} = q^2.$

(19)




$q \notin R_2 \cup R_3$

(20)




and $-1 = q^2$

(21)



when $q \in R_6$


(22)




and $q^4 = q^{-1}$.

(23)


and $q^{-1} = q^2$.




2.26 GDD 3 of Row 10



when $q \in R_5 \cup R_7 \cup R_6 \cup R_4$




when $q \in R_6$.




when $q \in R_4 \cup R_6 \cup R_8$.


(4).



when $q \in R_5 \cup R_6 \cup R_7 \cup R_4$.




when $q \in R_5 \cup R_7 \cup R_9 \cup R_6 \cup R_4$




when $q \in R_4$

(7).




when $s = q^2$ or $s = q^{-2}$, $q \in R_3 \cup R_6$ or $s = -q^{-2}$, $s \in R_3 \cup R_6$ or $s = -1$, $q^{-2} \in R_2 \cup R_3 \cup R_4 \cup R_6$ or $q^{-2} = -1$, $s \in R_2 \cup R_3 \cup R_4 \cup R_6$; $r = -q, q \in R_6$, or $r = -1$, $q \in R_4 \cup R_6$; $q, r, s \in F^* \setminus \{1\}$, $qrs = 1$, $q \neq r, q \neq s, s \neq r$.

(8).




when $q \in R_6 \cup R_4$.

(9).




when $q \in R_6$.


(10).



, $q \in R_3$.




$q \notin R_2 \cup R_3 \cup R_4$.




$q \notin R_2 \cup R_3$

(13).




when $q \in R_5 \cup R_7 \cup R_6 \cup R_4$.




$q \notin R_2 \cup R_3$

(15).




when $q \in R_8$.

(16).





when $s = -q$, $q \in R_6$ or $-1 = s$, $q \in R_4 \cup R_6$; $r = q^2$ or $r = -q^{-2} q \in R_6$ or $-1 = q^{-2}$, $r \in R_2 \cup R_3 \cup R_4 \cup R_6$ or $-1 = r$, $q^{-2} \in R_2 \cup R_3 \cup R_4 \cup R_6$; $q, r, s \in F^* \setminus \{1\}$, $qrs = 1$, $q \neq r, q \neq s, s \neq r$.


(17).



when $q \in R_6$.



(20)



when $\tilde{q}_{34} = q^{-1}$ and $q^8 = 1$. $q^2, q \notin R_2 \cup R_3$

(21)

when $\tilde{q}_{34} = s$. $q = -s$. or $\tilde{q}_{34} = q^2 s^{-1}$. $q^3 = sr^{-1}$. or $\tilde{q}_{34} = q^2 s^{-1}$. $q^2 s^{-1} = r$. $q = r^{-2}$
 with $, q, r, s \in F^* \setminus \{1\}$, $qrs = 1$, $q \neq r$, $q \neq s$, $s \neq r$..

(22)

when $\tilde{q}_{34} = q$ with $q^2, q^3 \neq 1$.

(23)

when $\tilde{q}_{34} = q$ and $q \in R_6$.

(24)

when $\tilde{q}_{34} = r$ and $q = -r$, or $\tilde{q}_{34} = s$ and $q = s^{-2}$. $rsq^{-2} = 1$.

(25)

$$q \notin R_2 \cup R_3,$$

when $\tilde{q}_{34} = r$ and $q = -r$, or $\tilde{q}_{34} = s$ and $q = s^{-2}$. $rsq^{-2} = 1$; $q \notin R_2 \cup R_3$.

$$(26) \quad \begin{array}{ccccc} & & q & & \\ -1 & \bullet & \square & \bullet & -1 \\ & q & q^{-2} & q^4 & \\ -1 & \bullet & & \bullet & -1 \\ & & q^{-2} & & \end{array}$$

when $\tilde{q}_{34} = q$, $q \in R_6$.

2.27 GDD 1 of Row 11

$$(1). \quad \begin{array}{ccccccc} q^2 & q^{-2} & q & q^{-1} & -1 & q^{-1} & q \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

$q \notin R_4$, $q \in F^* \setminus \{1, -1\}$.

$$(2). \quad \begin{array}{ccccccc} -1 & q & q & q^{-1} & -1 & q^{-1} & q \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

when $q \in R_3$

$$(3) \quad \begin{array}{ccccccc} -q & -q^{-1} & q & q^{-1} & -1 & q^{-1} & q \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

$, q \in R_3$

$$(4). \quad \begin{array}{ccccccc} -1 & -1 & q & q^{-1} & -1 & q^{-1} & q \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

$, q \in R_3$


$$(5). \quad \begin{array}{ccccccc} -1 & -q & q & q^{-1} & -1 & q^{-1} & q \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

$, q \in R_3$.

$$(6) \quad \begin{array}{ccccccc} q & q^{-2} & q^2 & q^{-2} & -1 & q^{-2} & q^2 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$


$$q^4 \neq 1,$$

(7)




$$q^4 \neq 1,$$

(8)




$$q^3 \in R_3.$$

(9)




$$r \neq 1, r \neq q, q^{-1}; r \neq 1, r \neq q, q^{-1}.$$

(10)




$$, q \notin R_6.$$

(11)




$$, q \in R_3.$$

(12)




$$\text{when } q \in R_3$$

(13)



$$, q \in R_3.$$



, $q \in R_3$.

(15)

$q^2 \neq 1$,

(16)





$q \in R_3$.

(17)


$q \in R_3$.

(18)


is affine.



2.28 GDD 2 of Row 11




when $q \in R_3$.




when $q \in R_3$.

(3).




when $q \in R_3$.

(4).




when $-q = r$ or $r = -1$ and when $-q = s$ or $s = -1$; $q, r, s \in F^* \setminus \{1\}, qrs = 1, q \neq r, q \neq s, s \neq r; q \in R_3$.


(5).




, $q \in R_3$




, $q \in R_3$



$q \in R_3$.




$q \in R_3$.




$q \in R_3$.

(10)



when $\tilde{q}_{34} = q^{-1}$. $q \in R_3$.

(11)



when $\tilde{q}_{34} = q^{-2}$; $q \in R_3$.

2.29 GDD 1 of Row 12

(1).

A path graph with four vertices and three edges. Edges are labeled with q , q^{-2} , q^2 , $q^{-2}q^2$, q^{-2} , and $-q^{-2}$.

$q = -\xi, \xi \in R_3, q \in F^* \setminus \{1, -1\}$.

(2).

A path graph with four vertices and three edges. Edges are labeled with q^2 , q^{-2} , q , $q^{-1}q$, q^{-1} , and $-q^{-1}$.

$q = -\xi, \xi \in R_3, q \in F^* \setminus \{1, -1\}$.

$$(3). \quad \begin{array}{ccccccc} q & & -q^{-1} & & -q^{-1} & & -q^{-1} & & q \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$

$, q \in R_3.$

$$(4). \quad \begin{array}{ccccccc} -1 & & q^{-2} & & q & & q^{-1} & & q & & -q^{-1} \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$

$q = -\xi, \xi \in R_3.$

$$(5). \quad \begin{array}{ccccccc} q & & -q^{-1} & & q & & -q^{-1} & & q^{-1} & & q^{-1} \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$

$, q \in R_3.$

$$(6). \quad \begin{array}{ccccccc} & & & & -q^{-1} & & \\ & & & & \bullet & & \\ & & & & | & & \\ & & & & q^{-1} & & \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \\ q & & q^{-1} & & q & & q^{-1} & & q \end{array}$$

$$q = -\xi, \xi \in R_3, q \in F^* \setminus \{1\}.$$

$$(7). \quad \begin{array}{ccccccc} & & & & -q^{-1} & & \\ & & & & \bullet & & \\ & & & & | & & \\ & & & & q^{-1} & & \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \\ -1 & & q^{-1} & & q & & q^{-1} & & q \end{array}$$

$q \in R_6.$

2.30 GDD 1 of Row 13

$$(1). \quad \begin{array}{ccccccc} q & & q^{-1} & & q & & q^{-1} & & q & & q^{-2} & & -1 \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$

$q \in R_6 \cup R_3,$

$$(2). \quad \begin{array}{ccccccc} q & & q^{-2} & & q^2 & & q^{-2} & & q^2 & & q^{-4} & & -1 \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$

$q \notin R_6 \cup R_3,$

$$(3). \quad \begin{array}{ccccccc} q^2 & & q^{-2} & & q & & q^{-1} & & q & & q^{-2} & & -1 \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$

when $q \in R_3 \cup R_6$.

$$(4). \quad \begin{array}{ccccccc} -1 & & & & & & \\ q^{-1} & q & q^{-1}q & q^{-2} & -1 & & \\ \bullet & \bullet & \bullet & \bullet & \bullet & & \end{array}$$

with $q \in R_6$.

$$(5). \quad \begin{array}{ccccccc} q & & & & & & \\ -q & -q^{-1} & -q^{-1}q & -q^{-1}q^2 & -1 & & \\ \bullet & \bullet & \bullet & \bullet & \bullet & & \end{array}$$

, $q \in R_3$.

$$(6). \quad \begin{array}{ccccccc} -1 & & & & & & \\ q^{-2} & q & q^{-1}q & q^{-2} & -1 & & \\ \bullet & \bullet & \bullet & \bullet & \bullet & & \end{array}$$

when $q \in R_3 \cup R_6$

$$(7). \quad \begin{array}{ccccccc} q & & & & & & \\ -q^{-1}q & -q^{-1}q & -q^{-1}q & q^{-2} & -1 & & \\ \bullet & \bullet & \bullet & \bullet & \bullet & & \end{array}$$

when $q \in R_3$

$$(8). \quad \begin{array}{cccc} & & -1 & \\ & & q^{-2} & \\ \bullet & & \bullet & \\ q & q^{-1} & q & q^{-1}q \end{array}$$

when $q \in R_3 \cup R_6$.


$$(9). \quad \begin{array}{cccc} & & -1 & \\ & & q^{-2} & \\ \bullet & & \bullet & \\ -1 & q^{-1} & q & q^{-1}q \end{array}$$

when $q \in R_3$

$$(10). \quad \begin{array}{cccc} & & -1 & \\ & & q^{-2} & \\ \bullet & & \bullet & \\ -1 & q^{-2} & q & q^{-1}q \end{array}$$


$q \in R_6$.

(10)




$$q = q^{-2}$$

(11)




with $q \in R_3$

(12)



when $q \in R_3 \cup R_6$

(13)



with $-q = q^{-2}$

2.31 GDD 2 of Row 13

(1)
$$\begin{array}{cccccc} q & q^{-1}q & q^{-1} & -q^{-1}q^2 & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

when $q \in R_3 \cup R_6$.

(2)
$$\begin{array}{cccccc} -1 & q^{-1}q & q^{-1} & -q^{-1}q^2 & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$


when $q \in R_6$.

(3)
$$\begin{array}{cccc} -1 & q^2 & q^2 & -1 \\ \bullet & \bullet & \bullet & \bullet \\ & -q^{-1} & q^{-1} & q \end{array}$$

when $q \in R_3$.


(4)

when $q \in R_6$.



(5)

when $q \in R_3 \cup R_6$.



2.32 GDD 1 of Row 14

$$\begin{array}{cccc} (1) & -q^{-1/2} & -1 & q^{-2}q^2 & q^{-2} & -q^{-2} \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \\ q^2 = -\xi, \xi \in R_3. & & & & & \end{array}$$

$$\begin{array}{cccc} (2) & -1 & -1 & q^{-2}q^2 & q^{-2} & -q^{-2} \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \\ q^2 = -\xi, \xi \in R_3.. & & & & & \end{array}$$

$$\begin{array}{cccc} (3) & -1 & -1 & q^{-3}q^3 & q^{-3} & -q^{-3} \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \\ q^3 \in R_6. & & & & & \end{array}$$

$$\begin{array}{cccc} (4) & -q^{-1/2} & -1 & q^{-3}q^3 & q^{-3} & -q^{-3} \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \\ \text{when } q \in R_{18}. & & & & & \end{array}$$

$$\begin{array}{cccc} (5) & r & -1 & q^{-1}q & q^{-1} & -q^{-1} \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \\ \text{when } q \in R_6; r \neq 1, r \neq q, q^{-1}. & & & & & \end{array}$$

$$\begin{array}{cccc} (6) & q & -1 & q^2 & q^{-2}q^2 & -q^2 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \\ q \in R_6. & & & & & \end{array}$$

$$\begin{array}{cccc} (7) & q & -1 & q^{-1}q & q^{-1} & -q^{-1} \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \\ q \in R_6; r \neq 1, r \neq q, q^{-1}. & & & & & \end{array}$$


$$\begin{array}{cccc} (8) & q^{-2} & -1 & q^{-1}q & q^{-1} & -q^{-1} \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \\ q \in R_6. & & & & & \end{array}$$

$$\begin{array}{cccc} (9) & q^{-1} & -q^{-1} & -q & -q^{-1} & q \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \\ , q \in R_3. & & & & & \end{array}$$

$$\begin{array}{cccc} (10) & -1 & -1 & -q^{-1}q & -q^{-1}q^{-1} \\ \bullet & \bullet & \bullet & \bullet & \bullet \\ , q \in R_3. & & & & \end{array}$$


$$\begin{array}{cccc} (11) & q & -1 & -q^{-1}q & -q^{-1}q^{-1} \\ \bullet & \bullet & \bullet & \bullet & \bullet \\ , q \in R_3. & & & & \end{array}$$

(12)




when $q \in R_3 \cup R_6$.

(13)




when $q \in R_3 \cup R_6$.

(14)




when $r = -1, q \in R_6$.

(15)




with $q \in R_6$.


2.33 GDD 2 of Row 14



$q = -\xi, \xi \in R_3$.




$q = -\xi, \xi \in R_3$.




$q = -\xi, \xi \in R_3$.

(4).





$q = -\xi, \xi \in R_3$.




$q = -\xi, \xi \in R_3$.

(6).




(8)




and $q \in R_6$

(9)




with $q \in R_6.$

(10)




with $q \in R_6.$

(11)




when $q \in R_6.$

(12)



when $q \in R_6.$


(13)



with $q \in R_6$.

(14)

when $q \in R_6$.



2.34 GDD 3 of Row 14

(1).
 $q^2 \quad q^{-2} \quad q \quad q^{-1} - 1 \quad q \quad -q$

 $q = -\xi, \xi \in R_3, q \in R_6$.

(2).
 $q^3 \quad q^{-3} \quad q \quad q^{-1} - 1 \quad q \quad -q$

 $, q \notin R_3$.

(3).
 $q \quad -q \quad -q^{-1} - q \quad -1 \quad -q^{-1} q^{-1}$

 $, q \in R_3$.


(4).
 $-1 \quad q \quad -q - q^{-1} - 1 \quad -q \quad q$

 $, q \in R_3$.

(5).
 $q \quad q^{-2} \quad q^2 \quad q^{-2} - 1 \quad q \quad -q$


 $q = -\xi, \xi \in R_3$

(6)




$q \in R_3$.

(7)




$q \in R_3$

(8)



with $q \in R_6$.

(9)



with $q \in R_6$.

2.35 GDD 1 of Row 15

$$\begin{array}{cccc} (1) & & & \\ -q & -1 & q^2 & -1 \\ q^2 & & & \\ \bullet & \bullet & \bullet & \bullet \end{array}$$

when $q^2 \in R_3$.

$$\begin{array}{cccc} (2) & & & \\ -1 & -1 & q^2 & -1 \\ q & q^{-2} & q^2 & \\ \bullet & \bullet & \bullet & \bullet \end{array}$$

$q^2 \in R_3$.

$$\begin{array}{cccc} (3) & & & \\ -1 & -1 & q^3 & -1 \\ q & q^{-3} & q^3 & \\ \bullet & \bullet & \bullet & \bullet \end{array}$$

when $q^3 \in R_3$.

(4)

$$\begin{array}{cccc} -q & -1 & q^3 & -1 \\ q^2 & q^{-3} & q^3 & \\ \bullet & \bullet & \bullet & \bullet \end{array}$$

when $q^3 \in R_3$.

$$\begin{array}{cccc} (5) & & & \\ r & -1 & q & -1 \\ r^{-1} & q^{-1} & q & \\ \bullet & \bullet & \bullet & \bullet \end{array}$$


when $q \in R_3$; $r \neq 1, r \neq q, q^{-1}$.

$$\begin{array}{cccc} (6) & & & \\ q & -1 & q^{-2} & -1 \\ q^{-1} & q^2 & q^{-2} & \\ \bullet & \bullet & \bullet & \bullet \end{array}$$


with $q \in R_6$.

$$\begin{array}{cccc} (7) & & & \\ q & -1 & q & -1 \\ q^{-1} & q^{-1} & q & \\ \bullet & \bullet & \bullet & \bullet \end{array}$$


, $q \in R_3$.



when $q \in R_3$.




, $q \in R_3$.




when $q \in R_3$

(11)




when $q \in R_3$

(12)




when $q \in R_3$

(13)




with $r = -1$. $q \in R_3$

(14)




and $r = q^{-2}$, $q \in R_3$

(15)




and $q = q^4$

(16)




and $q = q^{-2}$


2.36 GDD 2 of Row 15




$q \in R_3.$




$q \in R_3.$




$, q \in R_3.$




$, q \in R_3.$




$, q \in R_3.$




$, q \in R_3.$




$q \in R_3.$




$q \in R_3.$




$q \in R_3.$




$q \in R_3.$



$, q \in R_3.$




$, q \in R_3.$



$, q \in R_3.$

(13).



, $q \in R_3$.

(14)

when $q \in R_3$

(15)

when $q \in R_3$.

(16).

when $q \in R_3$.

(17)

when $q \in R_3$.

(18)

when $q \in R_3$.








(19)


when $q \in R_3$.

(20)

when $q \in R_3$.


(21)






when $q \in R_3$.

(22)




when $q \in R_3$.

(23)




when $q \in R_3$.

(24)




when $q \in R_3$.

(25)




when $q \in R_3$.

2.37 GDD 3 of Row 15




when $q \in R_3$.

(2).




when $q \in R_3$.




when $q \in R_3$.


(4).




when $q \in R_3$.




when $q \in R_3$.




when $q \in R_3$.




when $q \in R_3$.




when $q \in R_3$.



when $q \in R_3$.




when $q \in R_3$.




when $q \in R_3$.

(12)




when $q \in R_3$.




when $q \in R_3$.

(14)





with $q \in R_3$. when $\tilde{q}_{43} = q^{-1}$.



$q \in R_3,$

2.38 GDD 4 of Row 15




when $q \in R_3$



when $q \in R_3.$

2.39 GDD 1 of Row 16



$$\begin{array}{cccc} -q^{-1} & -1 & q^{-2}q^2 & -q^{-2}-q^2 \\ \bullet & \bullet & \bullet & \bullet \\ \hline \end{array}$$

when $q \in R_3$.

$$\begin{array}{cccc} (3) & & & \\ -1 & -1 & q^{-2}q^2 & -q^{-2}-q^2 \\ \bullet & \bullet & \bullet & \bullet \\ \hline \end{array}$$

when $q^2 \in R_3$.

$$\begin{array}{cccc} (4) & & & \\ -1 & -1 & q^{-3}q^3 & -q^{-1}-q \\ \bullet & \bullet & \bullet & \bullet \\ \hline \end{array}$$

when $q^3 \in R_3$.

(5)

$$\begin{array}{cccc} -q^{-1} & -1 & q^{-3}q^3 & -q^{-3}-q^3 \\ \bullet & \bullet & \bullet & \bullet \\ \hline \end{array}$$

when $q^3 \in R_3$.

$$\begin{array}{cccc} (6) & & & \\ q^{-1} & -1 & q^{-1}q & -q^{-1}-q \\ \bullet & \bullet & \bullet & \bullet \\ \hline \end{array}$$

when $q \in R_3$.

$$\begin{array}{cccc} (7) & & & \\ r & r^{-1} & -1 & q^{-1}q & -q^{-1}-q \\ \bullet & \bullet & \bullet & \bullet & \bullet \\ \hline \end{array}$$

when $r \neq -q$; $q \in R_3$.

$$\begin{array}{cccc} (8) & & & \\ q & q^{-1} & -1 & q^2 & q^{-2} & -q^{-2}-q^2 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \\ \hline \end{array}$$

with $q^2 \in R_3$.

$$\begin{array}{cccc} (9) & & & \\ q & q^{-1} & -1 & q^{-1}q & -q^{-1}-q \\ \bullet & \bullet & \bullet & \bullet & \bullet \\ \hline \end{array}$$


when $q \in R_3$.

$$\begin{array}{cccc} (10) & & & \\ & -q & & \\ & | & & \\ & -q^{-1} & & \\ & \bullet & & \\ & | & & \\ -1 & q^{-1} & q & q^{-1} & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet \\ \hline \end{array}$$


$q \in R_3$.

(11)

and $-q^{-1} = -q^2$.




(12)




and $-q = r; q \in R_3$.


2.40 GDD 2 of Row 16




$q \in R_3$




$q^2 \in R_3$.




$q \in R_3$.




$q \in R_3$.




$, q \in R_3$.




$, q \in R_3$.




, $q \in R_3$.




, $q \in R_3$.




$q \in R_3$.




$q \in R_3$




, $q \in R_3$.



$q \in R_3$




$q \in R_3$.




$q \in R_3$

(15)




when $q \in R_3$

(16)




when $q \in R_3$

(17)




when $q \in R_3$

(18)




when $q \in R_3$

(19)




when $q \in R_3$

(20)




when $q \in R_3$


2.41 GDD 3 of Row 16




$q \in R_3.$




when $q \in R_3$



when $q \in R_3$




when $(-q)^3 = r$ or $q^2 = r$; $q \in R_3$; $r \neq 1, r \neq q, q^{-1}$.



, $q \in R_3$.


(6)



when $q \in R_3$


2.42 GDD 4 of Row 16

(1)




when $q \in R_3$.

(2)




$q = -\xi, \xi \in R_3$,

(3)



when $q \in R_3$


(4)



when $q \in R_3$


2.43 GDD 5 of Row 16

(1)




$q \in R_3,$

(2)




$q \in R_3.$

(3)



when $q \in R_3$


(4)



$q \in R_3,$


2.44 GDD 1 of Row 17

(1)




$r \in R_4 \cup R_3 \cup R_6 \cup R_2, q \in R_3.$

(2)




$q, r \in R_3,$


(3)




$r \in R_6, q \in R_3$




$r \neq -1, q \in R_3.$




$q \in R_3, r \in R_4,$




when $r = q$ and $q \in R_3$, or $r = q^{-1}$, or $r = -q$ and $r \in R_6$ or $r = -1$, with $r \in R_4 \cup R_3 \cup R_6 \cup R_2$



when $q = -r$ with $q, r \in R_3.$




when $q = r^{-2}$, $r \in R_6$ and $q \in R_3.$




when $r = q$ or $r = q^2$ or $r = -q$ with $r^2 \neq 1$; $q \in R_3$

(10)




when $r = q^{-1}$, or $r = q$, or $r = -q^{-1}$, or $r = -1$; $q \in R_3.$

(11)




when $r = q$, or $r = q^{-1}$, or $r = -q$, or $r = -1$ with $q \in R_3$.

(12)




when $r \in R_2 \cup R_3 \cup R_4 \cup R_6$ with $q \in R_3$.

(13)




$r = -q^{-1}$, or $r = q$ with $q \in R_3$.

(14)




and $q = r^{-1}$ with $q \in R_3$.

(15)




and $q = r^{-2}$ with $q \in R_3$.

(16)




and $q^2 = r^{-1}$ with $q \in R_3$.

(17)




and $q = -r^{-1}$; $q \in R_3$.

(18)




and $q^2 = -r^{-1}$ with $q \in R_3$.

(19)




and $q = r$ with $q \in R_3$.

(20)




and $q = r^2$ with $q \in R_3$.

(21)



and $r = q^2$ with $q \in R_3$.


(22)



and $r = -q$ with $q \in R_3$.


2.45 GDD 2 of Row 17

(1)




$q \in R_3$.

(2)




$q \in R_3$.

(3)



and $-q^{-1} = -q^2$ with $q \in R_3$.

(4)



and $r = -q$ with $q \in R_3$.


2.46 GDD 3 of Row 17

(1)




$q \in R_3.$

(2)




$q \in R_3.$

(3)




$q \in R_3.$

(4)




$q = -\xi, \xi \in R_3.$

(5)




$q = -\xi, \xi \in R_3,$

(6)




$q = -\xi, \xi \in R_3,$


(7)




$$q^2 = -\xi^2, \xi \in R_3$$



$$q = -\xi, \xi \in R_3.$$




$$q = -\xi, \xi \in R_3.$$




$$, q \in R_3.$$


2.47 GDD 4 of Row 17




$$\text{when } q \in R_3.$$




$$q^2 \in R_3.$$



$$\text{when } q \in R_3.$$




$$\text{when } q \in R_3.$$




$$\text{when } q \in R_3.$$


(6).




when $q \in R_3$.




$q \in R_3$.




when $q \in R_3$.



when $q \in R_3$.




when $q \in R_3$.




when $q \in R_3$.

(12)




when $q \in R_3$

(13)




when $q \in R_3$

(14)




when $q \in R_3$

(15)




when $q \in R_3$

(16)




when $q \in R_3$

(17)




when $q \in R_3$

(18)




when $q \in R_3$

(19)




when $q \in R_3$

(20)




when $q \in R_3$

(21)




when $q \in R_3$

(22)



when $q \in R_3$

(23)



when $q \in R_3$

2.48 GDD 5 of Row 17

(1)

$$\begin{array}{ccccccc} q^2 & q^{-2} & -1 & q^{-1} & -q^{-1} & q & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

$q \in R_3$.

(2)

$$\begin{array}{ccccccc} -1 & q^{-1} & -1 & q & -q & q^{-1} & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

$q \in R_3$.

(3).

$$\begin{array}{ccccccc} & & -1 & & & & \\ & & | & & & & \\ & & -q & & & & \\ \bullet & & \bullet & & \bullet & & \\ q^2 & q^{-2} & q & q^{-1} & -1 & & \end{array}$$

$r = -\xi, \xi \in R_3$.

(4).

$$\begin{array}{ccccccc} & & -1 & & & & \\ & & | & & & & \\ & & -q & & & & \\ \bullet & & \bullet & & \bullet & & \\ -1 & q^{-1} & q & q^{-1} & -1 & & \end{array}$$


when $q \in R_6$.

(5).

$$\begin{array}{ccccccc} & & -1 & & & & \\ & & | & & & & \\ & & q & & & & \\ \bullet & & \bullet & & \bullet & & \\ -1 & q & -q & -q^{-1} & -1 & & \end{array}$$


when $q \in R_3$.

(6)




$-r = q^{-2}, q \in R_3.$

(7)




when $q \in R_3.$

(8)




when $q \in R_3.$

(9)




when $q \in R_3.$

(10)



when $q \in R_3.$


(11)



when $q \in R_3.$

2.49 GDD 6 of Row 17

(1)



when $q \in R_3.$

2.50 GDD 7 of Row 17

$$\begin{array}{cccc} (1) & & & \\ -1 & q & -1 & q^{-1}q & -q & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

when $q \in R_3$.

$$\begin{array}{cccc} (2) & & & \\ -q & \frac{(2)}{q^2} & -1 & q^{-2}q^2 & -q^2 & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

when $q^2 \in R_3$.

$$\begin{array}{cccc} (3) & & & \\ -1 & q & -1 & q^{-2}q^2 & -q^2 & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

when $q^2 \in R_3$.

$$\begin{array}{cccc} (4) & & & \\ -1 & q & -1 & q^{-3}q^3 & -q^3 & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

when $q \in R_3$.

$$\begin{array}{cccc} (5) & & & \\ -q & \frac{(5)}{q^2} & -1 & q^{-3}q^3 & -q^3 & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

$q^3 \in R_3$.

$$\begin{array}{cccc} (6) & & & \\ q^{-1} & q & -1 & q^{-1}q & -q & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

when $q \in R_3$

$$\begin{array}{cccc} (7) & & & \\ r & r^{-1} & -1 & q^{-1}q & -q & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

$q \in R_3; r \neq 1, r \neq q, q^{-1}$.


$$\begin{array}{cccc} (8) & & & \\ q & q^{-1} & -1 & q^{-1}q & -q & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

$q \in R_3$.

$$\begin{array}{cccc} (9) & & & \\ & & -1 & \\ & & | & \\ & & -q & \\ & & | & \\ -1 & q^{-1} & q & q^{-1} & -1 \end{array}$$

$q \in R_3$.


(10).



$$q \in R_3.$$


(11)

when $q \in R_3$.




(12)

$q \in R_3$.




(13)

$q^2 \in R_3$.




(14)

$q^2 \in R_3$.




(15)

$q^3 \in R_3$.




(16)

and $r = -1$ when $q \in R_3$.




(17)

when $r = -q$, $q \in R_3$.




(18)



and $r = -1$ when $q \in R_3$.


2.51 GDD 8 of Row 17

(1)




$r = -q^{-1}$, $r \in R_3$, when $q \in R_3$.

(2)




when $q \in R_3$, $q = -r$.

(3)




when $q \in R_3$, $r^2 \neq 1$, $r = -q$. $r \neq 1$.

(4)




when $q = r$ or $q = r^2$, $r \in R_3$, $q \in R_3$.

(5)




when $q = r^2$, $r \in R_6$, $q \in R_3$.

(6)




when $q \in R_3$.

(7)




when $q \in R_3$.

(8)



$q \in R_3$.


(f)



$q \in R_3$.


2.52 GDD 9 of Row 17

(1)




$q \in R_3, r \in R_2 \cup R_3 \cup R_4 \cup R_6$.


(2)



when $q \in R_3, r \in R_3$.




when $q \in R_3, r \in R_6$,




when $q \in R_3, r^2 \neq 1$,

(5)




when $q \in R_3, r \in R_4$,


(6)



when $q \in R_3$.



when $q \in R_3$.



when $q \in R_3$.









when $q \in R_3, q \neq r$.



when $r = q^{-1}$ or $q^{-1} = r^2$. $q \in R_3$.

2.53 GDD 1 of Row 18





when $q \in R_9$.

2.54 GDD 2 of Row 18

3 Properties about arithmetic root system with rank 3 and 4

3.1 Classical types

Lemma 3.1. *A GDD is a classical GDD if and only if it is one of classical Type 1-7, Here classical types are listed as follows:*

Type 1, $2 \leq n$.

$$C_{n-1,q,i_1,i_2,\dots,i_j} \xrightarrow{q^{-2}q^2} \bullet$$

$q \in F^* \setminus \{1, -1\}$, $0 \leq j \leq n - 1$. In particular, if $q \in R_3$, then Type 1 turn into the following:

$$C_{n-1,q,i_1,i_2,\dots,i_j} \xrightarrow{q \quad q^{-1}} \bullet, q^3 = 1.$$

Type 2. $2 \leq n$.

$$C_{n-1,q^2,i_1,i_2,\dots,i_j} \xrightarrow{q^{-2}q} \bullet$$

$q \in F^* \setminus \{1, -1\}$. $0 \leq j \leq n - 1$. In particular, if $q \in R_3$, then Type 3 turn into the following:

$$C_{n-1,q^{-1},i_1,i_2,\dots,i_j} \xrightarrow{q \quad q} \bullet, q^3 = 1.$$

Type 3, $2 \leq n$.

$$C_{n-1,q^{-2},i_1,i_2,\dots,i_j} \xrightarrow{q^2 \quad -q^{-1}} \bullet$$

$q \in F^* \setminus \{1, -1\}$, $0 \leq j \leq n - 1$,

In particular,

if $q \in R_3$, then Type 3 turn into the following:

$$C_{n-1, q^{-1}, i_1, i_2, \dots, i_j} \xrightarrow{q^{-q}} \bullet, q^3 = 1.$$

If $q \in R_4$, then Type 3 turn into the Type 2.

If $j = 0$, then Type 3 turn into the following:

$$\bullet \xrightarrow{q^{-2} q^2} \bullet \xrightarrow{q^{-2} q^2} \bullet \xrightarrow{q^{-2}} \dots \xrightarrow{q^{-2} q^2} \bullet \xrightarrow{-q^{-1}} \bullet, q \in F^*/\{1, -1\}.$$


which is Type 2.

Type 4. $2 \leq n$.

$$C_{n-1, -q^{-1}, i_1, i_2, \dots, i_j} \xrightarrow{-q^q} \bullet$$


$q^3 = 1$. $0 \leq j \leq n - 1$.

Type 5. $3 \leq n$.



$q \neq 1$, $0 \leq j \leq n - 2$.

Type 6. $3 \leq n$.



$q^2 \neq 1$. $0 \leq j \leq n - 2$,

Type 7. $1 \leq n$. $C_{n, q^{-1}, i_1, i_2, \dots, i_j}$. $q \neq 1$, $0 \leq j \leq n$.

Proof. We only consider the case $n > 4$ since every GDD classical type with $n < 5$ is subGDD of a GDD classical type with $n > 4$.

Necessity is obviously. Now we show the sufficiency.

Type 1. Type 1 is classical by GDD 1 of Row 10 in Table C when $2 \leq j \leq n - 1$. Type 1 is classical by GDD 1 of Row 9 in Table C when $j = 0$. Type 1 is classical by GDD 2 of Row 9 in Table C and by GDD 1 of Row 7 in Table C when $j = 1$.

Type 2. Type 2 is classical by GDD 1 of Row 4 in Table C when $q^4 \neq 1$ and $1 \leq j \leq n - 1$. Type 2 is classical by GDD 1 of Row 3 in Table C when $q^2 \neq 1$ and $j = 0$. Type 2 is classical by GDD 1 of Row 3 in Table C when $1 \leq j \leq n - 1$.

Type 3. Type 3 is classical by GDD 2 of Row 4 in Table C when $q^4 \neq 1$ and $1 \leq j \leq n - 1$.

Type 4. Type 4 is classical by GDD 1, 2 of Row 6 in Table C when $1 \leq j \leq n - 1$. Type 4 is classical by GDD 1 of Row 5 in Table C when $j = 0$.

Type 5. Type 5 is classical by GDD 3 of Row 10 in Table C when $1 \leq j \leq n - 2$ with $q \neq 1$. Type 5 is classical by GDD 1 of Row 8 in Table C when $j = 0$ with $q^2 \neq 1$.

Type 6. Type 6 is classical by GDD 2 of Row 10 in Table C when $1 \leq j \leq n - 2$. Type 1 is classical by GDD 3 of Row 9 in Table C when $j = 0$.

3.2 Properties about arithmetic root system with rank 3

Lemma 3.2. $n = 3$.

(I) *GDD is not an arithmetic root system if there exist two vertexes i and j with $q_{ii} \neq -1$ and $\tilde{q}_{ij} \neq 1$ such that $q_{ii}\tilde{q}_{ij} \neq 1$ and i is not a middle vertex when GDD is a chain except*

$$\begin{array}{c} (a) \\ -q \quad -1 \quad -q^{-1}q^{-1} \\ \bullet \quad \bullet \quad \bullet \end{array}, q \in R_3. \text{ GDD 3 of Row 14 or Type 4.}$$

$$\begin{array}{c} (b) \\ q \quad -q^{-1}q \quad -q^{-1}q \\ \bullet \quad \bullet \quad \bullet \end{array}, q \in R_3. \text{ GDD 5 of Row 16.}$$

$$\begin{array}{c} (c) \\ -1 \quad -1 \quad -1 \quad -q^{-1}q^{-1} \\ \bullet \quad \bullet \quad \bullet \end{array}, q \in R_3. \text{ GDD 9 of Row 17.}$$

$$\begin{array}{c} (d) \\ q \quad q^{-1} \quad q \quad q^{-1}q^{-3} \\ \bullet \quad \bullet \quad \bullet \end{array}, q \in R_9. \text{ GDD 1 of Row 18.}$$

$$\begin{array}{c} (e) \\ q \quad q^{-1} \quad q^{-4} \quad q^4 \quad q^{-3} \\ \bullet \quad \bullet \quad \bullet \end{array}, q \in R_9. \text{ GDD 2 of Row 18.}$$

$$\begin{array}{c} (f) \\ -1 \quad -q^{-1}q \quad -q \quad q \\ \bullet \quad \bullet \quad \bullet \end{array}, q \in R_3. \text{ GDD 2 of Row 14 or Type 4.}$$

$$(g) \quad \begin{array}{c} -1 \quad -1 \\ \quad \bullet \quad q^{-1} \\ \quad \bullet \quad \bullet \\ -1 \quad -q \quad q \end{array}, q \in R_3. \text{ GDD 8 of Row 17.}$$

$$\begin{array}{c} (h) \\ q^3 \quad -1 \quad q^2 \quad -q^{-1} \\ \bullet \quad \bullet \quad \bullet \end{array}, q \in F^* \setminus \{1, -1\}. \quad q \notin R_3, \text{ GDD 4 of Row 7.}$$

$$\begin{array}{c} (i) \\ q^2 \quad q^{-2} \quad q^2 \quad q^{-2}q \\ \bullet \quad \bullet \quad \bullet \end{array}, q \in F^* \setminus \{1, -1\}. \text{ GDD 1 of Row 2 or Type 2.}$$

(j) $\begin{array}{c} q \\ \bullet \text{---} \bullet \text{---} \bullet \\ -1 \quad -1 \quad -q^{-1}-q \end{array}, q \in R_3. \text{ GDD 4 of Row 16.}$

(k) $\begin{array}{c} \bullet \\ q^{-1} \quad -1 \quad -q \\ \bullet \text{---} \bullet \\ q \quad -1 \quad -1 \end{array}, q \in R_3. \text{ GDD 3 of Row 16.}$

(l) $\begin{array}{c} q \\ -q^{-1} \quad \bullet \quad q^{-1} \\ \bullet \text{---} \bullet \\ -q \quad -q^{-1}-1 \end{array}, q \in R_3. \text{ GDD 3 of Row 17.}$

(m) $\begin{array}{c} -1 \\ q^{-1} \quad \bullet \quad q^3 \\ \bullet \text{---} \bullet \\ q \quad q^{-2} \quad -1 \end{array} \quad q^2, q^3 \neq 1, \text{ GDD 3 of Row 7.}$

(n) $\begin{array}{c} q \quad q^{-2} \quad -1 \\ -1 \quad -1 \quad -1 \quad q \quad q \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}, q \in R_3. \text{ GDD 4 of Row 7.}$

(o) $\begin{array}{c} q^{-1} \quad -1 \quad -1 \quad q \\ q \quad \bullet \text{---} \bullet \text{---} \bullet \end{array}, q \in R_4. \text{ GDD 4 of Row 7.}$

(p) classical GDD satisfied conditions.

(II) Chain GDD is not an arithmetic root system if $\tilde{q}_{21}\tilde{q}_{23} \neq 1$, $q_{22} = -1$ and there exists another vertex another vertex $i \neq 2$ such that $q_{ii} = -1$ and $\tilde{q}_{i2} \neq -1$ except

(a) $\begin{array}{c} -1 \\ q \quad -1 \quad q^{-2}q^2 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}, q \in F^* \setminus \{1, -1\}. \text{ GDD 2 of Row 6 or Type 1.}$

(b) $\begin{array}{c} -1 \\ q \quad -1 \quad q^{-3}q^3 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}, q \in F^* \setminus \{1, -1\}, q^3 \neq 1, \text{ GDD 2 of Row 7.}$

(d) $\begin{array}{c} -1 \\ q \quad -1 \quad q \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}, q \in R_3. \text{ GDD 2 of Row 15.}$

(e) $\begin{array}{c} -1 \\ q \quad -1 \quad -q^{-1}q \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}, q \in R_3. \text{ GDD 2 of Row 16.}$

(f) $\begin{array}{c} -1 \\ -1 \quad -1 \quad q \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}, q \in R_3. \text{ GDD 1 of Row 17.}$

(g) $\begin{array}{c} -1 \\ q \quad -1 \quad -q \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}, q \in R_3. \text{ GDD 4 of Row 17.}$

(h) classical GDD satisfied conditions.

(III) Chain GDD is not an arithmetic root system if $q_{22} = -1$, $\tilde{q}_{21}\tilde{q}_{23} \neq 1$ and $\tilde{q}_{21}q_{11} = 1$ when $q_{33} = -1$ or $\tilde{q}_{23}q_{33} = 1$ when $q_{11} = -1$ except

$$\begin{array}{c} (a) \\ \bullet \xrightarrow{-1} \bullet \xrightarrow{q} \bullet \xrightarrow{-1} \bullet \xrightarrow{q^{-2}q^2} \bullet \end{array}, q \in F^* \setminus \{1, -1\}. \text{ GDD 2 of Row 6 or type 1.}$$

$$\begin{array}{c} (b) \\ \bullet \xrightarrow{-1} \bullet \xrightarrow{q} \bullet \xrightarrow{-1} \bullet \xrightarrow{q^{-3}q^3} \bullet \end{array}, q \in F^* \setminus \{1, -1\}, q \notin R_3, \text{ GDD 2 of Row 7.}$$

$$\begin{array}{c} (c) \\ \bullet \xrightarrow{-1} \bullet \xrightarrow{q} \bullet \xrightarrow{-1} \bullet \xrightarrow{-q^{-1}q} \bullet \end{array}, q \in R_3. \text{ GDD 2 of Row 16.}$$

$$\begin{array}{c} (d) \\ \bullet \xrightarrow{-1} \bullet \xrightarrow{-1} \bullet \xrightarrow{-1} \bullet \xrightarrow{q^{-1}q} \bullet \end{array}, q \in R_4. \text{ GDD 2 of Row 10 or type 1.}$$

$$\begin{array}{c} (e) \\ \bullet \xrightarrow{-1} \bullet \xrightarrow{-1} \bullet \xrightarrow{-1} \bullet \xrightarrow{q^{-1}q} \bullet \end{array}, q^2 \neq 1, \text{ GDD 1 of Row 9.}$$

(f) classical GDD satisfied conditions.

(IV) GDD is an arithmetic root system if $\tilde{q}_{21}q_{11} = 1$ and $\tilde{q}_{23}q_{33} = 1$ with $q_{22} = -1$.

(V) GDD is not an arithmetic root system if there exist different three i, j, k such that $q_{ii} \neq -1$, $q_{ii}\tilde{q}_{ij} \neq 1$ and $q_{ii}\tilde{q}_{ik} \neq 1$ with $\tilde{q}_{ik} \neq 1$ and $\tilde{q}_{ij} \neq 1$ except

$$\begin{array}{c} (a) \\ \bullet \xrightarrow{q} \bullet \xrightarrow{q^{-1}} \bullet \xrightarrow{-q^{-1}q^2} \bullet \xrightarrow{-1} \bullet \end{array}, q \in R_3 \cup R_6. \text{ GDD 2 of Row 13.}$$

$$\begin{array}{c} (b) \\ \bullet \xrightarrow{-1} \bullet \xrightarrow{q^{-1}} \bullet \xrightarrow{-q^{-1}q^{-1}} \bullet \xrightarrow{-1} \bullet \end{array}, q \in R_3. \text{ GDD 4 of Row 15.}$$

$$\begin{array}{c} (c) \\ \bullet \xrightarrow{-1} \bullet \xrightarrow{q^{-1}} \bullet \xrightarrow{q^{-1}q^{-1}} \bullet \xrightarrow{-1} \bullet \end{array}, q \in R_3. \text{ GDD 6 of Row 17.}$$

(VII) Chain GDD is not an arithmetic root system if $q_{11} = -1$, $\tilde{q}_{12} = q_{22} \neq -1$ or $q_{33} = -1$, $\tilde{q}_{32} = q_{22} \neq -1$ except

$$\begin{array}{c} (a) \\ \bullet \xrightarrow{-1} \bullet \xrightarrow{q^{-1}} \bullet \xrightarrow{q} \bullet \xrightarrow{-1} \bullet \end{array}, q \in R_3. \text{ GDD 1 of Row 15.}$$

$$\begin{array}{c} (b) \\ -1 \quad q^{-1} \quad q^{-1} \quad -q^{-1-1} \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}, q \in R_3. \text{ GDD 6 of Row 17.}$$

$$\begin{array}{c} (c) \\ q \quad q^{-1} \quad q \quad q^{-2-1} \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}, q \in R_3. \text{ GDD 1 of Row 13.}$$

$$\begin{array}{c} (d) \\ q \quad q^{-1} \quad -q^{-1} \quad q^2 \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}, q \in R_6. \text{ GDD 2 of Row 13.}$$

(VIII) GDD is not an arithmetic root system if there exist two different i, j such that $q_{ii} = -1 = \tilde{q}_{ij}$ with the fixed parameter $q \in R_3$ except

$$\begin{array}{c} (a) \\ -1 \quad -1 \quad -1 \quad q \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}, q \in R_3. \text{ GDD 1 of Row 17.}$$

$$\begin{array}{c} (b) \\ -1 \quad -1 \quad q \quad q^{-1-1} \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}, q \in R_3. \text{ GDD 2 of Row 17.}$$

$$\begin{array}{c} (c) \\ -1 \quad -1 \quad -1 \quad -q^{-1}q^{-1} \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}, q \in R_3. \text{ GDD 9 of Row 17.}$$

$$(d) \quad \begin{array}{c} \quad \quad -1 \\ \quad \quad \bullet \\ -1 \quad \quad q^{-1} \\ \quad \quad \bullet \\ -1 \quad -q \quad q \end{array}, q \in R_3. \text{ GDD 8 of Row 17.}$$

$$\begin{array}{c} (e) \\ q \quad q^{-1} \quad -1 \quad -1 \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array} q^2 \neq 1, \text{ GDD 1 of Row 9.}$$

$$(f) \quad \begin{array}{c} \quad \quad -1 \\ \quad \quad \bullet \\ q^{-1} \quad \quad -q \\ \quad \quad \bullet \\ q \quad -1 \quad -1 \end{array}, q \in R_3. \text{ GDD 3 of Row 16.}$$

$$\begin{array}{c} (g) \\ q \quad -1 \quad -1 \quad -q^{-1-q} \\ \bullet \text{---} \bullet \text{---} \bullet \end{array} q \in R_3. \text{ GDD 4 of Row 16.}$$

(IX) Chain GDD is not an arithmetic root system if $q_{22} = -\tilde{q}_{21} \notin R_4$ with $q_{11} = -1$ or $q_{22} = -\tilde{q}_{23} \notin R_4$ with $q_{33} = -1$ except

$$\begin{array}{c} (a) \\ q \quad q^{-1} \quad -q^{-1} \quad q^2 \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}, q \in R_3. \text{ GDD 2 of Row 13.}$$

$$\begin{array}{c} (b) \\ -q \quad -q^{-1}q^{-1} \quad -q^{-1}1 \\ \bullet \quad \bullet \quad \bullet \end{array}, q \in R_3. \text{ GDD 2 of Row 13.}$$

$$\begin{array}{c} (c) \\ -1 \quad q^{-1} \quad -q^{-1}1 \\ \bullet \quad \bullet \quad \bullet \end{array}, q \in R_3. \text{ GDD 4 of Row 15.}$$

$$\begin{array}{c} (d) \\ -1 \quad -q^{-1}q \quad -1 \\ \bullet \quad \bullet \quad \bullet \end{array}, q \in R_3. \text{ GDD 5 of Row 17.}$$

$$\begin{array}{c} (e) \\ -1 \quad q^{-1} \quad q^{-1} \\ \bullet \quad \bullet \quad \bullet \end{array}, q \in R_3. \text{ GDD 6 of Row 17.}$$

$$\begin{array}{c} (f) \\ -1 \quad q^{-1} \quad q \quad -q \quad -1 \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array}, q \in R_3. \text{ GDD 7 of Row 17.}$$

(X) Chain GDD is not an arithmetic root system if $q_{22} = q_{33} = -1$, $\tilde{q}_{32} \neq -1$ and $q_{11}\tilde{q}_{12} = 1$ or $q_{22} = q_{11} = -1$, $\tilde{q}_{12} \neq -1$ and $q_{33}\tilde{q}_{32} = 1$ except

$$\begin{array}{c} (a) \\ q \quad q^{-1} \quad -1 \quad q \quad -1 \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array}, q \in F^* \setminus \{1, -1\}. \text{ GDD 2 of Row 4.}$$

$$\begin{array}{c} (b) \\ q^2 \quad q^{-2} \quad -1 \quad q \quad -1 \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array}, q \in F^* \setminus \{1, -1\}. \text{ GDD 2 of Row 6.}$$

$$\begin{array}{c} (c) \\ q^3 \quad q^{-3} \quad -1 \quad q \quad -1 \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array}, q \in F^* \setminus \{1, -1\}. \quad q \notin R_3, \text{ GDD 2 of Row 7.}$$

$$\begin{array}{c} (d) \\ -q \quad -q^{-1}1 \quad q \quad -1 \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array}, q \in R_3. \text{ GDD 2 of Row 16.}$$

$$\begin{array}{c} (e) \\ -1 \quad -1 \quad -1 \quad q \quad -1 \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array}, q \in R_3. \text{ GDD 1 of Row 17.}$$

(XI) Chain GDD is not an arithmetic root system if $q_{11} = q_{22} = q_{33} = -1$ except

$$\begin{array}{c} (a) \\ -1 \quad q^{-1} \quad -1 \quad q \quad -1 \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array}, q \in F^* \setminus \{1, -1\}. \text{ GDD 2 of Row 8.}$$

(b).

$$\begin{array}{c} -1 \quad -1 \quad -1 \\ \bullet \xrightarrow{q} \bullet \xrightarrow{q} \bullet \end{array}, q \in R_3. \text{ GDD 2 of Row 15.}$$

$$\begin{array}{c} (c) \\ -1 \quad -1 \quad -1 \\ \bullet \xrightarrow{-q} \bullet \xrightarrow{q} \bullet \end{array}, q \in R_3. \text{ GDD 4 of Row 17.}$$

$$\begin{array}{c} (d) \\ -1 \quad -1 \quad -1 \\ \bullet \xrightarrow{q} \bullet \xrightarrow{-q} \bullet \end{array}, q \in R_3. \text{ GDD 4 of Row 17.}$$

$$\begin{array}{c} (e) \\ -1 \quad -1 \quad -1 \\ \bullet \xrightarrow{q} \bullet \xrightarrow{-1} \bullet \end{array}, q \in R_3. \text{ GDD 1 of Row 17;} \\ q \in R_6, \text{ GDD 2 of Row 7; } q \in R_4, \text{ GDD 2 of Row 6.}$$

$$\begin{array}{c} (f) \\ q \quad -1 \quad -1 \quad -1 \\ \bullet \xrightarrow{-1} \bullet \xrightarrow{-1} \bullet \end{array}, q \in R_4. \text{ GDD 1 of Row 2.}$$

(g) classical GDD satisfied conditions.

(XII) Chain GDD is not an arithmetic root system if $q_{22} \neq -1$ and $q_{11} = -1$, $q_{22}\tilde{q}_{12} \neq 1$ or $q_{22} \neq -1$ and $q_{33} = -1$ and $q_{22}\tilde{q}_{32} \neq 1$ except

$$\begin{array}{c} (a) \\ q \quad q \quad q^{-2} \\ \bullet \xrightarrow{q^{-1}} \bullet \xrightarrow{q} \bullet \end{array}, q \in R_3 \cup R_6. \text{ GDD 1 of Row 13.}$$

$$\begin{array}{c} (b) \\ q \quad -q^{-1} \quad -q^{-2} \\ \bullet \xrightarrow{q^{-1}} \bullet \xrightarrow{q^{-2}} \bullet \end{array}, q \in R_3. \text{ GDD 2 of Row 13.}$$

$$\begin{array}{c} (c) \\ -1 \quad -q^{-1} \quad -1 \\ \bullet \xrightarrow{q^{-1}} \bullet \xrightarrow{q^{-1}} \bullet \end{array}, q \in R_3. \text{ GDD 4 of Row 15.}$$

$$\begin{array}{c} (d) \\ -1 \quad q \quad q^{-1} \\ \bullet \xrightarrow{-1} \bullet \xrightarrow{q} \bullet \end{array}, q \in R_3. \text{ GDD 2 of Row 17.}$$

$$\begin{array}{c} (e) \\ -1 \quad -q \quad -1 \\ \bullet \xrightarrow{q} \bullet \xrightarrow{q^{-1}} \bullet \end{array}, q \in R_3. \text{ GDD 5 of Row 17.}$$

(f)

$$\begin{array}{c} -1 \\ \bullet \end{array} \xrightarrow{q^{-1}} \begin{array}{c} q^{-1} \\ \bullet \end{array} \xrightarrow{-q^{-1}} \begin{array}{c} -1 \\ \bullet \end{array}, q \in R_3. \text{ GDD 6 of Row 17.}$$

$$\begin{array}{c} -1 \\ \bullet \end{array} \xrightarrow{q^{-1}} \begin{array}{c} (g) \\ q \\ \bullet \end{array} \xrightarrow{-q} \begin{array}{c} -1 \\ \bullet \end{array}, q \in R_3. \text{ GDD 7 of Row 17.}$$

$$\begin{array}{c} -1 \\ \bullet \end{array} \xrightarrow{q^{-1}} \begin{array}{c} (h) \\ q \\ \bullet \end{array} \xrightarrow{q} \begin{array}{c} -1 \\ \bullet \end{array}, q \in R_3. \text{ GDD 1 of Row 15.}$$

(XIII) Cycle GDD is not an arithmetic root system if $q_{ii} \neq -1$ for $1 \leq i \leq 3$.

(XIV) GDD is not an arithmetic root system if there exist distinct two vertexes i and j with $q_{ii} \neq -1$ and $\tilde{q}_{ij} \neq 1$ such that $q_{ii}\tilde{q}_{ij} \neq 1$ and i is not a middle vertex when GDD is a chain, as well as there exists other place where simple chain condition (3) is not satisfied except

$$\begin{array}{c} q \\ \bullet \end{array} \xrightarrow{q^{-1}} \begin{array}{c} (a) \\ q^{-4} \\ \bullet \end{array} \xrightarrow{q^4} \begin{array}{c} q^{-1} \\ \bullet \end{array}, q \in R_9. \text{ GDD 2 of Row 18.}$$

(XV) Chain GDD is not an arithmetic root system if $q_{22}^{-2} = \tilde{q}_{12}$ or $q_{22}^{-2} = \tilde{q}_{32}$ except

$$\begin{array}{c} q \\ \bullet \end{array} \xrightarrow{q^{-1}} \begin{array}{c} (a) \\ q \\ \bullet \end{array} \xrightarrow{q^{-2}} \begin{array}{c} q^2 \\ \bullet \end{array}, q^2 \neq 1. \text{ GDD 1 of Row 3.}$$

$$\begin{array}{c} -1 \\ \bullet \end{array} \xrightarrow{q^{-1}} \begin{array}{c} (b) \\ q \\ \bullet \end{array} \xrightarrow{q^{-2}} \begin{array}{c} q^2 \\ \bullet \end{array}, q^2 \neq 1. \text{ GDD 1 of Row 6.}$$

$$\begin{array}{c} q \\ \bullet \end{array} \xrightarrow{q^{-1}} \begin{array}{c} (d) \\ q^{-4} \\ \bullet \end{array} \xrightarrow{q^4} \begin{array}{c} q^{-1} \\ \bullet \end{array}, q \in R_9. \text{ GDD 2 of Row 18.}$$

$$\begin{array}{c} q \\ \bullet \end{array} \xrightarrow{q^{-1}} \begin{array}{c} (e) \\ -q \\ \bullet \end{array} \xrightarrow{-q^{-1}} \begin{array}{c} -1 \\ \bullet \end{array}, q \in R_3 \cup R_6. \text{ GDD 2 of Row 13.}$$

$$\begin{array}{c} q \\ \bullet \end{array} \xrightarrow{q^{-1}} \begin{array}{c} (f) \\ q \\ \bullet \end{array} \xrightarrow{q^{-2}} \begin{array}{c} -1 \\ \bullet \end{array}, q \in R_3 \cup R_6. \text{ GDD 1 of Row 13.}$$

$$\begin{array}{c} -1 \\ \bullet \end{array} \xrightarrow{q^{-1}} \begin{array}{c} (g) \\ -q \\ \bullet \end{array} \xrightarrow{-q^{-1}} \begin{array}{c} -1 \\ \bullet \end{array}, q \in R_3. \text{ GDD 4 of Row 15.}$$

(h)

$$\overset{-1}{\bullet} \xrightarrow{q} \overset{-q}{\bullet} \xrightarrow{-q^{-1}-1} \overset{-1}{\bullet}, q \in R_3. \text{ GDD 5 of Row 17.}$$

$$\overset{-1}{\bullet} \xrightarrow{q^{-1}} \overset{(i)}{q^{-1}} \xrightarrow{q^{-1}-q^{-1}-1} \overset{-1}{\bullet}, q \in R_3. \text{ GDD 6 of Row 17.}$$

$$\overset{-1}{\bullet} \xrightarrow{q^{-1}} \overset{(j)}{q} \xrightarrow{q} \overset{-1}{\bullet}, q \in R_3. \text{ GDD 1 of Row 15.}$$

(XVI) Cycle GDD is not an arithmetic root system if $q_{ii} \neq -1$ for $1 \leq i \leq 3$

(XVII) Cycle GDD is not an arithmetic root system if only one vertex is -1 except

$$(a) \begin{array}{c} \overset{-1}{\bullet} \\ \swarrow \quad \searrow \\ \overset{q^{-1}}{\bullet} \quad \overset{q^{-1}}{\bullet} \\ \swarrow \quad \searrow \\ \overset{q}{\bullet} \quad \overset{q^{-1}q}{\bullet} \end{array}, q \in R_3. \text{ GDD 3 of Row 15.}$$

$$(b) \begin{array}{c} \overset{q}{\bullet} \\ \swarrow \quad \searrow \\ \overset{-q}{\bullet} \quad \overset{q^{-1}}{\bullet} \\ \swarrow \quad \searrow \\ \overset{-q}{\bullet} \quad \overset{-q^{-1}1}{\bullet} \end{array}, q \in R_3. \text{ GDD 3 of Row 17.}$$

(XVIII) If $srq = 1$ and $r = q^{-2}$, then $s = q$.

(XIX) The fixed parameter $q \in F^* \setminus \{1, -1\}$ from Row 1 to 10 in Table A_2 except $q \in F^* \setminus \{1\}$ in Row 1; $q \in F^* \setminus \{1, -1\}$ and $q \notin R_4$ in Row 5; $q \in F^* \setminus \{1, -1\}$ and $q \notin R_3$ in Row 7; $q, r, s \in F^* \setminus \{1\}$, $qrs = 1$, $q \neq r$, $q \neq s$, $s \neq r$, in Row 9; $q \in F^* \setminus \{1, -1\}$ and $q \notin R_3$ in Row 10. The fixed parameter $q \in R_3$ from Row 11 to 18 in Table A_2 except $q \in R_3 \cup R_4$ in Row 13; $q \in R_9$ in Row 18

Proof. We only prove (IV) since others are clear. We show it by three cases. Case one. If $\tilde{q}_{21}\tilde{q}_{23} = 1$, then GDD is an arithmetic root system since it is a simple chain. Case two. If $\tilde{q}_{21} = \tilde{q}_{23}$, then GDD is an arithmetic root system by GDD 1 of Row 10 ($q \in F^* \setminus \{1, -1\}$ and $q \notin R_3$) and GDD 1 of Row 11, $q \in R_3$.

Case three. If $\tilde{q}_{21} \neq \tilde{q}_{23}, \tilde{q}_{23}^{-1}$. then GDD is an arithmetic root system by GDD 1 of Row 9. \square

Lemma 3.3. $n = 3$.

(I) GDD is not an arithmetic root system if $\tilde{q}_{23}q_{33} = 1$ and $q_{22} = -1$ with $q_{33} \neq -1$ except

$$\overset{-1}{\bullet} \xrightarrow{q} \overset{-1}{\bullet} \xrightarrow{q^{-1}q} \overset{-1}{\bullet} \quad q^2 \neq 1, \text{ GDD 2 of Row 4 or Type 7.}$$

$$\overset{-q}{\bullet} \xrightarrow{\frac{(a)}{q^2}} \overset{-1}{\bullet} \xrightarrow{q^{-2}q^2} \overset{-1}{\bullet} \quad q^4 \neq 1, \text{ GDD 3 of Row 5 or Type 3.}$$

$$\begin{array}{c} \overset{(b)}{-1} \\ \bullet \text{---} \bullet \text{---} \bullet \\ \underset{q}{-1} \quad \underset{q^{-2}q^2}{-1} \end{array} \quad q^2 \neq 1, \text{ GDD 2 of Row 6 or Type 1.}$$

$$\begin{array}{c} \overset{(c)}{-1} \\ \bullet \text{---} \bullet \\ \underset{q^2}{-1} \quad \underset{q^{-1}q}{q^{-1}} \end{array} \quad q^2 \neq 1, \text{ GDD 3 of Row 6 or Type 6.}$$

$$\begin{array}{c} \overset{(d)}{-1} \\ \bullet \text{---} \bullet \text{---} \bullet \\ \underset{q}{-1} \quad \underset{q^{-3}q^3}{-1} \end{array} \quad q^6 \neq 1, \text{ GDD 2 of Row 7.}$$

$$\begin{array}{c} \overset{(e)}{-1} \\ \bullet \text{---} \bullet \\ \underset{q^3}{-1} \quad \underset{q^{-1}q}{q^{-2}} \end{array} \quad q^3, q^2 \neq 1, \text{ GDD 3 of Row 7.}$$

$$\begin{array}{c} \overset{(f)}{-q} \\ \bullet \text{---} \bullet \text{---} \bullet \\ \underset{q^2}{-q} \quad \underset{q^{-3}q^3}{-1} \end{array} \quad , q^6 \neq 1, \text{ GDD 4 of Row 7.}$$

$$\begin{array}{c} \overset{(g)}{q^{-1}} \\ \bullet \text{---} \bullet \text{---} \bullet \\ \underset{q}{q^{-1}} \quad \underset{q^{-1}q}{-1} \end{array} \quad , q^2 \neq 1, \text{ GDD 1 of Row 8 or Type 7.}$$

$$\begin{array}{c} \overset{(h)}{r} \\ \bullet \text{---} \bullet \text{---} \bullet \\ \underset{r^{-1}}{r^{-1}} \quad \underset{q^{-1}q}{-1} \end{array} \quad , q^2 \neq 1, q \neq r, r^{-1} \text{ GDD 1 of Row 9.}$$

$$\begin{array}{c} \overset{(i')}{q} \\ \bullet \text{---} \bullet \text{---} \bullet \\ \underset{q^{-1}}{q^{-1}} \quad \underset{q^2}{-1} \quad \underset{q^{-2}}{q^{-2}} \end{array} \quad , q^4, q^3 \neq 1. \text{ GDD 2 of Row 10 or Type 1.}$$

$$\begin{array}{c} \overset{(k)}{q} \\ \bullet \text{---} \bullet \text{---} \bullet \\ \underset{q^{-1}}{q^{-1}} \quad \underset{q^{-1}q}{-1} \end{array} \quad q^2, q^3 \neq 1. \text{ GDD 1 of Row 10 or Type 5.}$$

$$\begin{array}{c} \overset{(l)}{q^{-2}} \\ \bullet \text{---} \bullet \text{---} \bullet \\ \underset{q^2}{q^2} \quad \underset{q^{-1}q}{-1} \end{array} \quad q^2, q^3 \neq 1.. \text{ GDD 2 of Row 10 or Type 1.}$$

$$\begin{array}{c} \overset{(m)}{q} \\ \bullet \text{---} \bullet \text{---} \bullet \\ \underset{q^{-1}}{q^{-1}} \quad \underset{q^{-1}q}{-1} \end{array} \quad , q \in R_3. \text{ GDD 1 of Row 11 or Type 5.}$$

$$\begin{array}{c} \overset{(n)}{q^{-1}} \\ \bullet \text{---} \bullet \text{---} \bullet \\ \underset{-q^{-1}}{-q^{-1}} \quad \underset{-q}{-1} \quad \underset{-q^{-1}}{-q^{-1}} \end{array} \quad , q \in R_3. \text{ GDD 3 of Row 14 or Type 4.}$$

(0)

$$\begin{array}{c} q^{-1} q \\ \bullet \quad \bullet \\ \bullet \quad \bullet \\ -1 \quad q^{-1} q \end{array}, q \in R_3. \text{ GDD 3 of Row 15,}$$

$$\begin{array}{c} (p) \\ -1 \quad q \quad -1 - q^{-1} q \\ \bullet \quad \bullet \quad \bullet \end{array}, q \in R_3. \text{ GDD 2 of Row 16.}$$

$$(q) \quad \begin{array}{c} -1 \\ -q \quad \bullet \quad -1 \\ \bullet \quad \bullet \\ -1 \quad q^{-1} q \end{array}, q \in R_3. \text{ GDD 3 of Row 16.}$$

$$(r) \quad \begin{array}{c} q \quad -1 \quad -1 - q^{-1} q \\ \bullet \quad \bullet \quad \bullet \end{array}, q \in R_3. \text{ GDD 4 of Row 16.}$$

$$(s) \quad \begin{array}{c} -q^{-1} q \\ -q^{-1} \quad \bullet \quad -q^{-1} \\ \bullet \quad \bullet \\ -1 \quad q^{-1} q \end{array}, q \in R_3. \text{ GDD 3 of Row 17.}$$

$$(s'') \quad \begin{array}{c} q^{-1} q \\ \bullet \quad \bullet \\ -1 \quad -q^{-1} q \end{array}, q \in R_3. \text{ GDD 3 of Row 17.}$$

$$(t) \quad \begin{array}{c} -1 \\ -1 \quad \bullet \quad -q \\ \bullet \quad \bullet \\ -1 \quad q^{-1} q \end{array}, q \in R_3. \text{ GDD 8 of Row 17.}$$

$$(u) \quad \begin{array}{c} -1 \quad -1 \quad -1 \quad q^{-1} q \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array}, q \in F^* \setminus \{1, -1\}. \text{ GDD 1 of Row 9.}$$

$$(v) \quad \begin{array}{c} q \quad q^{-2} \quad -1 \quad q^2 \quad q^{-2} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array}, q \in F^* \setminus \{1, -1\}. \text{ Type 2}$$

(w) classical GDD satisfied conditions.

(II) GDD is not an arithmetic root system if $\tilde{q}_{23}q_{33} = \tilde{q}_{21}q_{22} = 1$ and $q_{33} \neq -1$ except

$$(a). \quad \begin{array}{c} q \quad q^{-1} \quad q \quad q^{-1} q \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array}, q \in F^* \setminus \{1, -1\}. \text{ GDD 1 of Row 1 or Type 7.}$$

$$(b). \quad \begin{array}{c} q \quad q^{-2} \quad q^2 \quad q^{-2} q^2 \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array}, q \in F^* \setminus \{1, -1\}. \text{ GDD 1 of Row 2 or Type 2.}$$

$$(c). \quad \begin{array}{c} q^2 \quad q^{-2} \quad q \quad q^{-1} q \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array}, q \in F^* \setminus \{1, -1\}. \text{ GDD 1 of Row 3 or Type 1.}$$

(d). $\begin{array}{c} -1 \\ \bullet \end{array} \begin{array}{c} q^{-1} \\ \bullet \end{array} \begin{array}{c} q \\ \bullet \end{array} \begin{array}{c} q^{-1}q \\ \bullet \end{array}$, $q \in F^* \setminus \{1, -1\}$. GDD 1 of Row 4 or Type 7.

(e). $\begin{array}{c} q \\ \bullet \end{array} \begin{array}{c} -q \\ \bullet \end{array} \begin{array}{c} -q^{-1} \\ \bullet \end{array} \begin{array}{c} -q^{-1} \\ \bullet \end{array}$, $q \in R_3$. GDD 1 of Row 12 or Type 4.

(f). $\begin{array}{c} -1 \\ \bullet \end{array} \begin{array}{c} q^{-2} \\ \bullet \end{array} \begin{array}{c} q \\ \bullet \end{array} \begin{array}{c} q^{-1}q \\ \bullet \end{array}$, $q \in R_3 \cup R_6$. GDD 1 of Row 13.

(g). $\begin{array}{c} q^{-1} \\ \bullet \end{array} \begin{array}{c} -1 \\ \bullet \end{array} \begin{array}{c} q^{-1} \\ \bullet \end{array}$, $q \in R_3$. GDD 3 of Row 15.

(h). $\begin{array}{c} q \\ \bullet \end{array} \begin{array}{c} -q^{-1}-q \\ \bullet \end{array} \begin{array}{c} -q^{-1}-q \\ \bullet \end{array}$, $q \in R_3$. GDD 5 of Row 16.

(i). $\begin{array}{c} q^{-3} \\ \bullet \end{array} \begin{array}{c} q^{-1} \\ \bullet \end{array} \begin{array}{c} q \\ \bullet \end{array} \begin{array}{c} q^{-1}q \\ \bullet \end{array}$, $q \in R_9$. GDD 1 of Row 18.

(j). $\begin{array}{c} -q \\ \bullet \end{array} \begin{array}{c} q^2 \\ \bullet \end{array} \begin{array}{c} q^{-2} \\ \bullet \end{array} \begin{array}{c} q^2 \\ \bullet \end{array} \begin{array}{c} q^{-2} \\ \bullet \end{array}$, $q^2 \neq 1$, Type 3.

(k) classical GDD satisfied conditions.

(III) GDD is not an arithmetic root system if this GDD is not classical and every vertex is not -1 , if $q_{22} \neq -1$, $\tilde{q}_{23}q_{22} = 1$ and $q_{33} = -1$ except

(a). $\begin{array}{c} q^2 \\ \bullet \end{array} \begin{array}{c} q^{-2} \\ \bullet \end{array} \begin{array}{c} q \\ \bullet \end{array} \begin{array}{c} q^{-1}-1 \\ \bullet \end{array}$, $q^2 \neq 1$. GDD 1 of Row 6 or Type 1.

(b). $\begin{array}{c} q^{-1} \\ \bullet \end{array} \begin{array}{c} -1 \\ \bullet \end{array} \begin{array}{c} q^2 \\ \bullet \end{array}$, $q^2 \neq 1$. GDD 3 of Row 6 or Type 6.

(c). $\begin{array}{c} q^3 \\ \bullet \end{array} \begin{array}{c} q^{-3} \\ \bullet \end{array} \begin{array}{c} q \\ \bullet \end{array} \begin{array}{c} q^{-1}-1 \\ \bullet \end{array}$, $q \in F^* \setminus \{1, -1\}$, $q \notin R_3$. GDD 1 of Row 7.

(d). $\begin{array}{c} q^{-2} \\ \bullet \end{array} \begin{array}{c} -1 \\ \bullet \end{array} \begin{array}{c} q^3 \\ \bullet \end{array}$, $q \in F^* \setminus \{1, -1\}$, $q \notin R_3$. GDD 3 of Row 7.

(e). $\begin{array}{c} -1 \\ \bullet \end{array} \begin{array}{c} q^{-1} \\ \bullet \end{array} \begin{array}{c} q \\ \bullet \end{array} \begin{array}{c} q^{-1}-1 \\ \bullet \end{array}$, $q \in F^* \setminus \{1, -1\}$. GDD 3 of Row 8 or Type 7.

(g).

$$\begin{array}{c} q \quad -q \quad -q^{-1} \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}, q \in R_3. \text{ GDD 1 of Row 14 or Type 4.}$$

$$\begin{array}{c} (h). \\ -1 \quad q \quad q \quad q^{-1} \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}, q \in R_3. \text{ GDD 1 of Row 15.}$$

$$(i). \quad \begin{array}{c} q^{-1} \quad q \quad q^{-1} \\ \bullet \text{---} \bullet \\ q \quad q^{-1} \quad -1 \end{array}, q \in R_3. \text{ GDD 3 of Row 15.}$$

$$\begin{array}{c} (j). \\ -q \quad -q^{-1} \quad q \quad q^{-1} \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}, q \in R_3. \text{ GDD 1 of Row 16.}$$

$$(k). \quad \begin{array}{c} -1 \\ -1 \quad \bullet \quad -q \\ \bullet \text{---} \bullet \\ q \quad q^{-1} \quad -1 \end{array}, q \in R_3. \text{ GDD 3 of Row 16.}$$

$$\begin{array}{c} (l). \\ -1 \quad -1 \quad q \quad q^{-1} \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}, q \in R_3. \text{ GDD 2 of Row 17.}$$

$$(m). \quad \begin{array}{c} -q^{-1} \quad -q \quad -q^{-1} \\ \bullet \text{---} \bullet \\ q \quad q^{-1} \quad -1 \end{array}, q \in R_3. \text{ GDD 3 of Row 17.}$$

$$(m') \quad \begin{array}{c} -q^{-1} \quad q \quad q^{-1} \\ \bullet \text{---} \bullet \\ -q \quad -q^{-1} \quad -1 \end{array}, q \in R_3. \text{ GDD 3 of Row 17.}$$

$$\begin{array}{c} (n). \\ -1 \quad q \quad -q \quad -q^{-1} \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}, q \in R_3. \text{ GDD 5 of Row 17.}$$

$$\begin{array}{c} (o). \\ -1 \quad -q \quad q \quad q^{-1} \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}, q \in R_3. \text{ GDD 7 of Row 17.}$$

$$(p). \quad \begin{array}{c} -1 \\ -q \quad \bullet \quad -1 \\ \bullet \text{---} \bullet \\ q \quad q^{-1} \quad -1 \end{array}, q \in R_3. \text{ GDD 8 of Row 17.}$$

$$\begin{array}{c} (a'). \\ q \quad q^{-1} \quad q \quad q^{-1} \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}, q^2 \neq 1. \text{ GDD 1 of Row 4 or Type 7.}$$

$$\begin{array}{c} (a''). \\ q \quad q^{-2} \quad q^2 \quad q^{-2} \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}, q^4 \neq 1. \text{ GDD 1 of Row 5 or Type 2.}$$

(q) classical GDD satisfied conditions.

(IV) GDD is not an arithmetic root system if $q_{22} = q_{33} = -1$ and $\tilde{q}_{32} \neq -1$ except

$$q \begin{array}{c} (a') \\ \hline q^{-1} \end{array} \begin{array}{c} -1 \\ \hline q \end{array} \begin{array}{c} -1 \\ \hline -1 \end{array}, q \in F^* \setminus \{1, -1\}. \text{ GDD 2 of Row 4 or Type 7.}$$

$$q \begin{array}{c} (a'') \\ \hline q^{-2} \end{array} \begin{array}{c} -1 \\ \hline q^2 \end{array} \begin{array}{c} -1 \\ \hline -1 \end{array}, q \in F^* \setminus \{1, -1\}. \text{ GDD 2 of Row 5 or Type 2.}$$

$$q^2 \begin{array}{c} (a) \\ \hline q^{-2} \end{array} \begin{array}{c} -1 \\ \hline q \end{array} \begin{array}{c} -1 \\ \hline -1 \end{array}, q \in F^* \setminus \{1, -1\}. \text{ GDD 2 of Row 6 or Type 1.}$$

$$(b). \quad \begin{array}{c} q^{-1} q \\ \diagdown \quad \diagup \\ \bullet \quad \bullet \\ \diagup \quad \diagdown \\ -1 \quad q^2 \end{array} \begin{array}{c} -1 \\ \hline -1 \end{array}, q \in F^* \setminus \{1, -1\}. \text{ GDD 3 of Row 6 or Type 6.}$$

$$q^3 \begin{array}{c} (c) \\ \hline q^{-3} \end{array} \begin{array}{c} -1 \\ \hline q \end{array} \begin{array}{c} -1 \\ \hline -1 \end{array}, q \in F^* \setminus \{1, -1\}. q \notin R_3, \text{ GDD 2 of Row 7.}$$

$$(d). \quad \begin{array}{c} q^{-1} q \\ \diagdown \quad \diagup \\ \bullet \quad \bullet \\ \diagup \quad \diagdown \\ -1 \quad q^3 \end{array} \begin{array}{c} q^{-2} \\ \hline -1 \end{array}, q \in F^* \setminus \{1, -1\}. q \notin R_3, \text{ GDD 3 of Row 7.}$$

$$-1 \begin{array}{c} (e) \\ \hline q^{-1} \end{array} \begin{array}{c} -1 \\ \hline q \end{array} \begin{array}{c} -1 \\ \hline -1 \end{array}, q \in F^* \setminus \{1, -1\}. \text{ GDD 2 of Row 8 or Type 7.}$$

$$(f). \quad \begin{array}{c} -1 \\ \bullet \\ \diagdown \quad \diagup \\ \bullet \quad \bullet \\ \diagup \quad \diagdown \\ -1 \quad q \end{array} \begin{array}{c} r \\ \hline -1 \end{array} \begin{array}{c} -1 \\ \hline -1 \end{array}, q, r, s \in F^* \setminus \{1\}, qrs = 1, q \neq r, q \neq s, s \neq r. \text{ GDD 3 of Row 9.}$$

$$(g). \quad \begin{array}{c} q \\ \diagdown \quad \diagup \\ \bullet \quad \bullet \\ \diagup \quad \diagdown \\ -1 \quad q \end{array} \begin{array}{c} q^{-2} \\ \hline -1 \end{array}, q \in F^* \setminus \{1, -1\}, q^3 \neq 1. \text{ GDD 3 of Row 10 or Type 6.}$$

$$(h). \quad \begin{array}{c} -1 \\ \bullet \\ \diagdown \quad \diagup \\ \bullet \quad \bullet \\ \diagup \quad \diagdown \\ -1 \quad q^{-2} \end{array} \begin{array}{c} q \\ \hline -1 \end{array} \begin{array}{c} -1 \\ \hline -1 \end{array}, q \in F^* \setminus \{1, -1\}, q^3 \neq 1. \text{ GDD 3 of Row 10 or Type 6.}$$

$$(i). \quad \begin{array}{c} -1 \\ \bullet \\ \diagdown \quad \diagup \\ \bullet \quad \bullet \\ \diagup \quad \diagdown \\ -1 \quad q \end{array} \begin{array}{c} q \\ \hline -1 \end{array}, q \in R_3. \text{ GDD 2 of Row 11 or Type 6.}$$

$$q \begin{array}{c} (j) \\ \hline -q \end{array} \begin{array}{c} -1 \\ \hline -q^{-1} \end{array} \begin{array}{c} -1 \\ \hline -1 \end{array}, q \in R_3. \text{ GDD 2 of Row 14 or Type 4.}$$

$$-1 \begin{array}{c} (k) \\ \hline q \end{array} \begin{array}{c} -1 \\ \hline q \end{array} \begin{array}{c} -1 \\ \hline -1 \end{array}, q \in R_3. \text{ GDD 2 of Row 15.}$$

(l).

$$\begin{array}{c} -q \quad -q^{-1} \quad q \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}, q \in R_3. \text{ GDD 2 of Row 16.}$$

$$(m). \begin{array}{c} q^{-1}q \\ \bullet \quad \bullet \\ \diagup \quad \diagdown \\ \bullet \quad \bullet \\ -1 \quad -q \quad -1 \end{array}, q \in R_3. \text{ GDD 3 of Row 16.}$$

$$(n'). \begin{array}{c} -1 \quad -1 \quad q \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}, q \in R_3. \text{ GDD 1 of Row 17.}$$

$$(o). \begin{array}{c} -1 \quad -q \quad -1 \quad q \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}, q \in R_3. \text{ GDD 4 of Row 17.}$$

$$(p). \begin{array}{c} -1 \quad -1 \quad -q \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}, q \in R_3. \text{ GDD 4 of Row 17.}$$

(s)

$$\begin{array}{c} -q^{-1} \quad -1 \quad q^{-2} \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}, q \in R_3. \text{ GDD 4 of Row 17 or Type 3}$$

(t) classical GDD satisfied conditions.

(V) GDD is not an arithmetic root system if $\tilde{q}_{23} = q_{22} = q_{33} = -1$ except

$$(a) \begin{array}{c} -1 \quad -1 \quad -1 \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad q \in R_4, \text{ GDD 2 of Row 6; } q \in R_3, \text{ GDD 1 of Row 17; } q \in R_6, \\ \text{GDD 2 of Row 7; } q \in R_2, \text{ GDD 1 of Row 1.}$$

$$(b) \begin{array}{c} q^{-1} \quad -q^{-1} \quad -1 \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad q \in R_3, \text{ GDD 9 of Row 17;}$$

$$(c) \begin{array}{c} q^{-1}q \\ \bullet \quad \bullet \\ \diagup \quad \diagdown \\ \bullet \quad \bullet \\ -1 \quad -1 \quad -1 \end{array} \quad q \in R_3, \text{ GDD 8 of Row 17.}$$

$$(d) \begin{array}{c} q \\ -q \quad \bullet \quad q^{-1} \\ \diagup \quad \diagdown \\ \bullet \quad \bullet \\ -1 \quad -1 \quad -1 \end{array} \quad q \in R_3, \text{ GDD 8 of Row 17.}$$

(e)

$$\begin{array}{c} -q^{-1} \quad -1 \quad -1 \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad q \in R_6, \text{ GDD 4 of Row 7.}$$

(f) $q \begin{matrix} (f) \\ q^{-1} -1 -1 -1 \end{matrix}$ $q^2 \neq 1$, GDD 1 of Row 9.

(g) $q \begin{matrix} (g) \\ -1 -1 -1 -1 \end{matrix}$ $q \in R_4$, GDD 1 of Row 2 or Type 2.

(h) $\begin{matrix} & -1 & \\ q & \bullet & r \\ & \bullet & \bullet \\ -1 & -1 & -1 \end{matrix}$ $q, q^{-1} \neq r$ with $r, q \neq -1$, GDD 3 of Row 9.

(i) $\begin{matrix} & q & \\ q^{-1} & \bullet & q^{-2} \\ & \bullet & \bullet \\ -1 & -1 & -1 \end{matrix}$ $q \in R_6$. GDD 3 of Row 7.

(j) $q \begin{matrix} (j) \\ q^{-1} -1 -1 -1 \end{matrix}$ $q \in R_4$. GDD 2 of Row 10 or Type 1.

(l) classical GDD satisfied conditions.

3.3 Properties about arithmetic root system with rank 4

Lemma 3.4. $n = 4$.


GDD is not an arithmetic root system if there exist two different vertexes i and j with $q_{ii} \neq -1$ and $\tilde{q}_{ij} \neq 1$ such that $q_{ii}\tilde{q}_{ij} \neq 1$ and i is not a middle vertex (i.e. $i \neq 2, 3$) when GDD is a chain except


(a) $\begin{matrix} -q & -q^{-1} & q & -q^{-1} & q & -q^{-1} & q \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{matrix}$ ($q \in R_3$, see GDD 1 of Row 17 in Table B).

(b) $\begin{matrix} -q & -q^{-1} & q & -q^{-1} & 1 & -1 & q \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{matrix}$ ($q \in R_3$, see GDD 2 of Row 17 in Table B).

(c) $\begin{matrix} & q^{-1} & q & q^{-1} & -1 \\ & \bullet & \bullet & \bullet & \bullet \\ q & \bullet & \bullet & \bullet & \bullet \\ & & q^{-1} & & -1 \end{matrix}$, $q \in R_4$, (see, $q \in R_3$. GDD 5 of Row 18 in Table B.)

(d) $\begin{matrix} & & q & q & q^{-1} \\ & & \bullet & \bullet & \bullet \\ q & \bullet & \bullet & \bullet & \bullet \\ & & q^{-1} & & -1 \end{matrix}$, (see, $q \in R_3$. GDD 10 of Row 20 in Table B.)


(e)  , (see, $q \in R_4$. GDD 4 of Row 22 in Table B.)


(f)  , ($q \in R_3$, see GDD 3 of Row 17 in Table B.)


(g) A classical GDD satisfied conditions.

Lemma 3.5. $n = 4$, and $q^4, q^6 \neq 1$, where q is a fix parameter in Table B

GDD is not an arithmetic root system if there exist two different vertexes i and j with $q_{ii} = -1$, $\tilde{q}_{ij} = -1$ and $q_{jj} = -1$ except

(a)  ($q^2 \neq 1$, see GDD 2 of Row 14 in Table B.)


(b)  , ($q^2 \neq 1$, see GDD 3 of Row 14 in Table B.)

(c)  ($q^2 \neq 1$, see GDD 4 of Row 14 in Table B.)


(d) classical GDD satisfied conditions.

Lemma 3.6. $n = 4$.

GDD is not an arithmetic root system if GDD is




with $\tilde{q}_{12}q_{22} \neq 1$, and $q_{22} \neq -1$, except


(e)  , (see, $q \in R_4$. GDD 4 of Row 22 in Table B.)


Lemma 3.7. $n = 4$.


Chain GDD is not an arithmetic root system if there exists a vertex i with $q_{ii} = -1$, $\tilde{q}_{i,i+1}\tilde{q}_{i,i-1} \neq 1$ and there exists other vertex which does not satisfy simple chain condition


(2) or condition (3). except


(a)  ($q \notin R_3, q \notin R_2$. GDD 6 of Row 9 in Table B).

(b)  ($q \in R_3$. see GDD 2 of Row 17 in Table B).

(c)  ($q \in R_3$, see GDD 5 of Row 17 in Table B).


(d)  ($q \in R_4$, see GDD 6 of Row 22 in Table B).

(e)  ($q \in R_4$. see GDD 7 of Row 22 in Table B).


(f)  ($q^2 \neq 1$. see GDD 4 of Row 14 in Table B).

Lemma 3.8. $n = 4$,


GDD is not an arithmetic root system if GDD is




or




with $q_{ii} \neq -1$ for $i = 1, 3, 4$, except


(a)  ($q \neq 1$, GDD 1 of Row 5 in Table B or Type 5.)


(b)  ($q^2 \neq 1$, see GDD 6 of Row 12 in Table B or Type 5.)


(d) classical GDD satisfied conditions.

Lemma 3.9. $n = 4$, GDD is not an arithmetic root system if $q_{ii} = -1$ for $1 \leq i \leq 4$, except

(a)  $(q \in R_4, \text{ GDD 5 of Row 22 in Table B.})$

(d)  $(q \in R_4, \text{ see GDD 4 of Row 9 in Table B.})$


(f)  $(q \in R_4, \text{ see GDD 2 of Row 9 in Table B.})$

(h)  $(q \in R_3, \text{ see GDD 3 of Row 20 in Table B.})$


(i) classical GDD satisfied conditions.


Lemma 3.10. $n = 4$,

chain GDD is not an arithmetic root system if $q_{44} = q_{22} = -1$, $\tilde{q}_{12}\tilde{q}_{23} \neq 1$ and $q_{11}\tilde{q}_{12} = 1$, or $q_{11} = q_{33} = -1$, $\tilde{q}_{43}\tilde{q}_{23} \neq 1$ and $q_{44}\tilde{q}_{43} = 1$, except

(a)  $(q \in R_4, \text{ see GDD 3 of Row 22 in Table B.})$

(b)  $(q^2 \neq 1, \text{ see GDD 3 of Row 12 in Table B or Type 1.})$

(c)  $(q \in R_4, \text{ GDD 5 of Row 9 in Table B.})$

(d)  $(q \in R_4, \text{ GDD 6 of Row 9 in Table B.})$

(e)  $(\text{ see GDD 2 of Row 12 in Table B or Type 1.})$

(f) $\begin{array}{cccc} -1 & q & q^{-1} & q^2 \\ \bullet & \bullet & \bullet & \bullet \end{array}$ (see GDD 3 of Row 12 in Table B or Type 1).


(g) $\begin{array}{cccc} -1 & -q & q & -q \\ \bullet & \bullet & \bullet & \bullet \end{array}$ ($q \in R_4$, see GDD 6 of Row 22 in Table B).

(h) $\begin{array}{cccc} -1 & -1 & -1 & -q \\ \bullet & \bullet & \bullet & \bullet \end{array}$ ($q \in R_4$, see GDD 7 of Row 22 in Table B).


(i) $\begin{array}{cccc} -1 & -q & -q & -q \\ \bullet & \bullet & \bullet & \bullet \end{array}$ ($q \in R_4$, see GDD 2 of Row 22 in Table B).

(j) classical GDD satisfied conditions.

Lemma 3.11. *GDD is not an arithmetic root system if GDD is*



with $q_{33} \neq -1$ or



with $q_{44} \neq -1$ except

(a) $\begin{array}{cccc} q^2 & q^{-2} & q^2 & -1 \\ \bullet & \bullet & \bullet & \bullet \end{array}$, ($q^2, q^3 \neq 1$, see GDD 3 of Row 9 in Table B.)


(b) $\begin{array}{cccc} -q & -q^{-1} & -q & -1 \\ \bullet & \bullet & \bullet & \bullet \end{array}$, ($q \in R_3$, see GDD 3 of Row 17 in Table B.)

(c) $\begin{array}{cccc} q^{-1} & q^{-1} & q^{-1} & -1 \\ \bullet & \bullet & \bullet & \bullet \end{array}$, ($q \in R_3$, see GDD 3 of Row 18 in Table B.)


(d) $\begin{array}{cccc} -1 & -1 & q^{-1} & -1 \\ \bullet & \bullet & \bullet & \bullet \end{array}$, (see GDD 8 of Row 20 in Table B.)

(e) $\begin{array}{cccc} -q & q^{-1} & -q & q \\ \bullet & \bullet & \bullet & \bullet \end{array}$, (see GDD 8 of Row 22 in Table B.)

Lemma 3.12. *GDD is not an arithmetic root system if GDD is*




Lemma 3.13. *GDD is not an arithmetic root system if GDD is*




except


(b)




(c)




(d)



(e)




(f)




Lemma 3.14. $n = 4$,

Chain GDD is not an arithmetic root system if GDD is $q_{22} = -1$ and $q_{11} \neq -1$ with $\tilde{q}_{12} = \tilde{q}_{23}$ or $q_{33} = -1$ and $q_{44} \neq -1$ with $\tilde{q}_{43} = \tilde{q}_{23}$ except

(a)



(b)



- (c) $\begin{array}{c} q^2 \quad q^{-2}q^2 \quad q^{-2} \quad -1q^3 \quad q^{-3} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array}$ ($q^5 = 1$, see GDD 5 of Row 9 in Table B).
- (d) $\begin{array}{c} q^2 \quad q^{-2}q \quad q^{-1} \quad -1q^3 \quad q^{-3} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array}$ ($q^4 = 1$, $q^2 \neq 1$, see GDD 6 of Row 9 in Table B).
- (e) $\begin{array}{c} -1 \quad q \quad q^{-1}q \quad -1q^{-2} \quad q^2 \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array}$ ($q^3 = 1$, see GDD 3 of Row 12 in Table B or Type 1).
- (f) $\begin{array}{c} q \quad q^{-1}q \quad q^{-1} \quad -1q^2 \quad q^{-2} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array}$ ($q^3 = 1$, see GDD 1 of Row 13 in Table B or Type 1).
- (g) $\begin{array}{c} q \quad q^{-1}q \quad q^{-1} \quad -1q \quad -q^{-1} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array}$ (see, $q^4 = 1$, $q^2 \neq 1$, GDD 1 of Row 14 in Table B).
- (h) $\begin{array}{c} -q^{-1} \quad -q^{-1}q \quad -1q^{-1} \quad q \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array}$ (see, $q \in R_4$, $q^2 \neq 1$, GDD 5 of Row 14 in Table B).
- (i) $\begin{array}{c} -q \quad -q^{-1}q \quad -q^{-1}q \quad -1q^{-1}q \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array}$ (see, $q \in R_3$, GDD 6 of Row 17 in Table B).
- (j) classical GDD satisfied conditions.

Lemma 3.15. $n = 4$,

Chain GDD is not an arithmetic root system if there exist i such that $\tilde{q}_{i+1,i}\tilde{q}_{i,i-1} \neq 1$ and $q_{ii} = -1$ with $q_{i-1,i-1} = -1$, or $q_{i+1,i+1} = -1$, except

- (a) $\begin{array}{c} q \quad q^{-1}q \quad -1q \quad -1q^{-2} \quad q^2 \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array}$ ($q^2 \neq 1$, GDD 3 of Row 8 in Table B or Type 1).
- (b) $\begin{array}{c} -1 \quad -1 \quad -1q \quad -1q^3 \quad q^{-3} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array}$ ($q^4 = 1$, GDD 5 of Row 9 in Table B).
- (c) $\begin{array}{c} -1 \quad q^{-1}q \quad -1q \quad -1q \quad -1 \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array}$ ($q \in R_3$. see GDD 3 of Row 20 in Table B).
- (d) $\begin{array}{c} -1 \quad q \quad q^{-1}q \quad -1q \quad -1 \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array}$ ($q \in R_3$. see GDD 4 of Row 20 in Table B).

$$(f) \begin{array}{c} -1 \quad -q \quad -\frac{1}{q} \quad -\frac{1}{q} \quad -q \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad (q \in R_4. \text{ see GDD 2 of Row 22 in Table B}).$$

$$(g) \begin{array}{c} -1 \quad q \quad -\frac{1}{-1} \quad -\frac{1}{q} \quad -q \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad (q \in R_4, \text{ see GDD 7 of Row 22 in Table B}).$$

$$(h) \begin{array}{c} q^2 \quad q^{-2} q^2 \quad q^{-2} \quad -\frac{1}{q} \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad (q^2, q^3 \neq 1, \text{ see GDD 2 of Row 9 in Table B}).$$

$$(i) \begin{array}{c} -1 \quad q^{-1} \quad -\frac{1}{q} \quad -\frac{1}{q^{-2}} \quad q^2 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad (q^2 \neq 1, \text{ see GDD 2 of Row 12 in Table B or Type 1}).$$

$$(j) \begin{array}{c} q \quad q^{-1} \quad -\frac{1}{-1} \quad -\frac{1}{-q} \quad -q^{-1} \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad (q^2 \neq 1, \text{ see GDD 4 of Row 14 in Table B}).$$

$$(k) \begin{array}{c} q \quad q^{-1} \quad -\frac{1}{q} \quad -\frac{1}{q} \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad (q \in R_3. \text{ see GDD 6 of Row 21 in Table B}).$$

$$(l) \begin{array}{c} -1 \quad q \quad q^{-1} \quad q \quad -\frac{1}{-1} \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad (q \in R_4. \text{ see GDD 3 of Row 12 in Table B or Type 1}).$$

$$(m) \begin{array}{c} q \quad q^{-1} q \quad q^{-1} \quad -\frac{1}{-1} \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad (q \in R_4. \text{ see GDD 1 of Row 13 in Table B or Type 1}).$$

$$(n) \begin{array}{c} q^2 \quad q^{-2} q^2 \quad q^{-2} \quad -\frac{1}{q^3} \quad q^{-3} \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad (q \in R_6. \text{ see GDD 5 of Row 9 in Table B}).$$


$$(o) \begin{array}{c} q^2 \quad q^{-2} q \quad q^{-1} \quad -\frac{1}{q^3} \quad q^{-3} \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad (q \in R_6. \text{ see GDD 6 of Row 9 in Table B}).$$

$$(p) \begin{array}{c} q^{-1} \quad q \quad q^{-1} \quad q \quad -\frac{1}{q} \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad (q \in R_3. \text{ see GDD 2 of Row 18 in Table B}).$$







(q) classical GDD satisfied conditions.

Lemma 3.16. $n = 4$, GDD is not an arithmetic root system if GDD

or



except



(g) classical GDD satisfied conditions.

Lemma 3.17. $n = 4$,

GDD is not an arithmetic root system if there exist three different vertexes i, j and k such that $q_{ii} \neq -1$, $\tilde{q}_{i,j}q_{i,i} \neq 1$ and $\tilde{q}_{i,k}q_{i,i} \neq 1$ with $\tilde{q}_{i,j} \neq 1$ and $\tilde{q}_{i,k} \neq 1$ except

$$(a) \begin{array}{c} -1 \quad -1 \quad -q^{-1} \quad -1 \\ q \quad q^{-1} \quad q^{-1} \quad -1 \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad (q \in R_3. \text{ see GDD 4 of Row 21 in Table B}).$$

$$(b) \begin{array}{c} q^{-1} \quad -1 \quad -q^{-1} \quad -1 \\ q \quad q^{-1} \quad q^{-1} \quad -1 \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad (q \in R_3. \text{ see GDD 2 of Row 20 in Table B}).$$

$$(c) \begin{array}{c} -1 \quad q^{-1} \quad q^{-1} \quad -q^{-1} \quad -1 \\ q \quad q \quad q \quad q^{-1} \quad -1 \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad (q \in R_3. \text{ see GDD 2 of Row 21 in Table B}).$$

Lemma 3.18. $n = 4$,

GDD is not an arithmetic root system if $q_{11} = -1$, $\tilde{q}_{12} = q_{22} \neq -1$ or $q_{44} = -1$, $\tilde{q}_{43} = q_{33}$, except

$$(a) \begin{array}{c} q^{-1} \quad -1 \quad q \quad q \quad -1 \\ q \quad q^{-1} \quad q \quad q \quad -1 \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad (q \in R_3. \text{ see GDD 1 of Row 20 in Table B}).$$

$$(b) \begin{array}{c} -1 \quad q^{-1} \quad q^{-1} \quad q \quad q \quad -1 \\ q \quad q \quad q \quad q \quad q \quad -1 \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad (q \in R_3. \text{ see GDD 1 of Row 21 in Table B}).$$

$$(c) \begin{array}{c} -1 \quad -1 \quad q \quad q \quad -1 \\ q \quad q^{-1} \quad q \quad q \quad -1 \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad (q \in R_3. \text{ see GDD 3 of Row 21 in Table B}).$$

(d) *classical GDD satisfied conditions.*

Lemma 3.19. $n = 4$, *Chain GDD is not an arithmetic root system if $q_{11} = -1$, $\tilde{q}_{12} = -q_{22}$ or $q_{44} = -1$, $\tilde{q}_{43} = -q_{33}$ with $q^4 \neq 1$, except*

$$(a) \begin{array}{c} q^{-1} \quad -1 \quad -q^{-1} \quad -1 \\ q \quad q^{-1} \quad q^{-1} \quad -1 \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad (\text{ see GDD 2 of Row 20 in Table B}).$$


$$(b) \begin{array}{c} -1 \quad q^{-1} \quad q^{-1} \quad -q^{-1} \quad -1 \\ q \quad q \quad q \quad q^{-1} \quad -1 \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad (q \in R_3, \text{ see GDD 2 of Row 21 in Table B}).$$

$$(c) \begin{array}{c} -1 \quad -1 \quad -q^{-1} \quad -1 \\ q \quad q^{-1} \quad q^{-1} \quad -1 \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad (q \in R_3, \text{ see GDD 4 of Row 21 in Table B}).$$


(f) A classical GDD satisfied conditions.

Lemma 3.20. $n = 4$,

GDD is not an arithmetic root system if



or




with $q_{ii} = -1$ for $1 \leq i \leq 4$ and $\tilde{q}_{12} = -\tilde{q}_{23}$ or $\tilde{q}_{24} = -\tilde{q}_{12}$, except

(f) $(q \in R_4, \text{ see GDD 5 of Row 22 in Table B.})$

(g) classical GDD satisfied conditions.

Lemma 3.21. $n = 4$,

GDD is not an arithmetic root system if GDD contains a sub DDG



except


(a) $(q \in R_3, \text{ GDD 3 of Row 20 in Table B.})$


(b) $(\text{ see GDD 6 of Row 21 in Table B.})$


(c) $(q \in R_4, \text{ GDD 1 of Row 2 in Table B.})$


(d) $(q \in R_4, \text{ GDD 2 of Row 9 in Table B.})$

(e) $(q \in R_4 \text{ GDD 5 of Row 9 in Table B.})$

(f)  $(q^2 \neq 1, \text{ GDD 2 of Row 13 in Table B.})$

(g)  $(q \in R_4, \text{ GDD 3 of Row 9 in Table B.})$


(h)  $(q \in R_4, \text{ GDD 4 of Row 9 in Table B.})$


(i)  $(\text{ see , } q \in F^* \setminus \{1, -1\}, \text{ GDD 5 of Row 12 in Table B or Type 6.})$


(j) classical GDD satisfied conditions.


Lemma 3.22. $n = 4,$


Chain GDD is not an arithmetic root system if there exists i such that $q_{ii} = -1$ and $q_{jj} \neq -1$ for $j \neq i$ with $\tilde{q}_{i,i-1} = \tilde{q}_{i,i+1}$, except

(a)  $(\text{ see, } q \in R_5, \text{ GDD 5 of Row 9 in Table B.})$

(c)  $(\text{ see, } q \in R_3, \text{ GDD 1 of Row 13 in Table B or Type 1.})$

(d)  $(\text{ see, } q \in R_4, \text{ GDD 1 of Row 14 in Table B.})$


(e)  $(\text{ see, } q \in R_4, \text{ GDD 5 of Row 14 in Table B.})$

(f)  $(\text{ see, } q \in R_3, \text{ GDD 6 of Row 17 in Table B.})$

(g) A classical GDD satisfied conditions.

Lemma 3.23. $n = 4,$

Chain GDD is not an arithmetic root system if $q_{22} = -1, \tilde{q}_{12}\tilde{q}_{23} \neq 1$ and $\tilde{q}_{12} \neq \tilde{q}_{23}$, or $q_{33} = -1, \tilde{q}_{43}\tilde{q}_{23} \neq 1$ and $\tilde{q}_{34} \neq \tilde{q}_{23}$, except

(a)  $(q \notin R_5. q \in R_2 \cup R_3 \text{ see GDD 5 of Row 9 in Table B.})$

(b)

$$\begin{array}{c} q \quad q^{-1}q \quad q^{-1} \quad -1q^2 \quad q^{-2} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad (q \notin R_3, \text{ see GDD 1 of Row 13 in Table B, or Type 1}).$$

$$(c) \quad \begin{array}{c} q \quad q^{-1}q \quad q^{-1} \quad -1q \quad -q^{-1} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad (q \notin R_4 \cup R_2, \text{ see GDD 1 of Row 14 in Table B}).$$

$$(d) \quad \begin{array}{c} -q^{-1} \quad -q^{-1} \quad -1q^{-1} \quad q \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad (q \notin R_4, q \in R_2, \text{ see GDD 5 of Row 14 in Table B}).$$

$$(e) \quad \begin{array}{c} q \quad q^{-1} \quad -1q \quad -1q^{-2} \quad q^2 \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad (q^2, q^3 \neq 1, \text{ see GDD 3 of Row 8 in Table B or Type 1}).$$

$$(f) \quad \begin{array}{c} q^2 \quad q^{-2}q^2 \quad q^{-2} \quad -1q \quad -1 \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad (q^2, q^3 \neq 1, \text{ see GDD 2 of Row 9 in Table B}).$$

$$(g) \quad \begin{array}{c} q^2 \quad q^{-2}q \quad q^{-1} \quad -1q^3 \quad q^{-3} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad (q^4, q^3 \neq 1, \text{ see GDD 6 of Row 9 in Table B}).$$

$$(h) \quad \begin{array}{c} -1 \quad q^{-1} \quad -1q \quad -1q^{-2} \quad q^2 \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad (q^2, q^3 \neq 1, \text{ see GDD 2 of Row 12 in Table B or Type 4}).$$

$$(i) \quad \begin{array}{c} -1 \quad q \quad q^{-1} \quad -1q^{-2} \quad q^2 \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad (q^2, q^3 \neq 1, \text{ see GDD 3 of Row 12 in Table B or Type 1}).$$

$$(j) \quad \begin{array}{c} -q \quad -q^{-1}q \quad -q^{-1} \quad -1 \quad q \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad (q \in R_3, \text{ see GDD 2 of Row 17 in Table B}).$$

$$(k) \quad \begin{array}{c} -q \quad -q^{-1}q \quad q^{-1} \quad -1q^{-1}q \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad (q \in R_3, \text{ see GDD 5 of Row 17 in Table B}).$$

$$(l) \quad \begin{array}{c} -1 \quad -q \quad q \quad -1 \quad -1q \quad -q \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad (q \in R_4, \text{ see GDD 6 of Row 22 in Table B}).$$


$$(m) \quad \begin{array}{c} -1 \quad q \quad -1 \quad -1 \quad -1q \quad -q \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad (\text{ see GDD 7 of Row 22 in Table B}).$$

$$(n) \quad \begin{array}{c} q \quad q^{-1} \quad -1 \quad -1 \quad -1q \quad -q^{-1} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad (q^2 \neq 1, \text{ see GDD 4 of Row 14 in Table B}).$$

(t) A classical GDD satisfied conditions.

Lemma 3.24. $n = 4$,


GDD



is not an arithmetic root system if $q_{33} \neq -1$ and $q_{44} \neq -1$.


Lemma 3.25. $n = 4$,

GDD




is not an arithmetic root system if there are two among q_{11}, q_{33} and q_{44} which are not equal to -1 , except

(b)



(c)




(d) A classical GDD satisfied conditions.

Lemma 3.26. $n = 4$, GDD is not an arithmetic root system if there exist two different i and j such that $\tilde{q}_{ij} = -1$ and $q^4 \neq 1$, where q is a fix parameter in Table B except


(a)




(b)



(c)



(d)



(e)

- (g) $q \quad q^{-1} \quad -1 \quad -1 \quad -q \quad -q^{-1}$ (see, $q \in R_3$, GDD 4 of Row 14 in Table B).
- (h) $q^2 \quad q^{-2} \quad -1 \quad -1 \quad q^2 \quad -1 \quad -1$ ($q \in R_6$. see GDD 4 of Row 9 in Table B.)
- (i) $q^2 \quad q^{-2} \quad q^2 \quad q^{-2} \quad -1 \quad -1 \quad -1$ ($q \in R_6$. see, GDD 5 of Row 9 in Table B).
- (j) $q^2 \quad q^{-2} \quad q \quad q^{-1} \quad -1 \quad -1 \quad -1$ ($q \in R_6$. see, GDD 6 of Row 9 in Table B).
- (j) A classical GDD satisfied conditions.

Lemma 3.27. $n = 4$. Chain GDD is not an arithmetic root system if $\tilde{q}_{i,i+1}\tilde{q}_{i,i-1} = -1$, and $q_{ii} = -1$, except

- (a) $q^2 \quad q^{-2} \quad q \quad q^{-1} \quad -1 \quad q^3 \quad q^{-3}$ (see, $q \in R_4$, GDD 6 of Row 9 in Table B).
- (b) $q \quad q^{-1} \quad q \quad q^{-1} \quad -1 \quad -q \quad -q^{-1}$ (see, $q^2 \neq 1$, GDD 1 of Row 14 in Table B).
- (c) $-q^{-1} \quad -q \quad -q^{-1} \quad -q \quad -1 \quad q^{-1} \quad q$ (see, $q^2 \neq 1$, GDD 5 of Row 14 in Table B).
- (d) $-1 \quad -q \quad -1 \quad q \quad -1 \quad q \quad -q$ (see, $q \in R_4$, GDD 2 of Row 22 in Table B).
- (e) $-1 \quad q \quad -q \quad q \quad -1 \quad q \quad -q$ (see, $q \in R_4$, GDD 3 of Row 22 in Table B).
- (f) $q^2 \quad q^{-2} \quad q \quad q^{-1} \quad -1 \quad q^3 \quad q^{-3}$ (see, $q \in R_4$, GDD 6 of Row 9 in Table B).

Lemma 3.28. $n = 4$. GDD is not an arithmetic root system if GDD contains sub-GDD

$$q \quad q^{-1} \quad -1 \quad q^{-1} \quad q$$



- except
- (a) $-q \quad -q^{-1} \quad q \quad q^{-1} \quad -1 \quad q^{-1} \quad -q$ (see, $q \in R_3$, GDD 6 of Row 17 in Table B).
- (d) $q^2 \quad q^{-2} \quad q^2 \quad q^{-2} \quad -1 \quad q^3 \quad q^{-3}$ (see, $q \in R_5$, GDD 5 of Row 9 in Table B).

- (e) $\begin{array}{c} q^2 \quad q^{-2}q \quad q^{-1} \quad -\frac{1}{q^3} \quad q^{-3} \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array}$ (see, $q \in R_4$, GDD 6 of Row 9 in Table B).
- (f) $\begin{array}{c} -\frac{1}{q} \quad q^{-1}q \quad -\frac{1}{q^{-2}} \quad q^2 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}$ (see, $q \in R_3$, GDD 3 of Row 12 in Table B or Type 1).
- (g) $\begin{array}{c} q^{-1}q \quad q^{-1}q \quad -\frac{1}{q^{-2}} \quad q^2 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}$ (see, $q \in R_3$, GDD 1 of Row 13 in Table B or Type 1).
- (h) $\begin{array}{c} q \quad q^{-1}q \quad q^{-1} \quad -\frac{1}{q} \quad -q^{-1} \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}$ (see, $q \in R_4$, GDD 1 of Row 14 in Table B).
- (i) $\begin{array}{c} -q^{-1}q \quad -q^{-1}q \quad -\frac{1}{q^{-1}} \quad q \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}$ (see, $q \in R_4$, GDD 5 of Row 14 in Table B).
- (j) $\begin{array}{c} -\frac{1}{q} \quad q \quad -\frac{1}{q} \quad -q \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}$ (see, $q \in R_4$, GDD 3 of Row 22 in Table B).
- (k) $\begin{array}{c} q \quad q^{-1}q \quad \begin{array}{l} q^{-1} \\ -1 \\ q^{-1} \end{array} \\ \bullet \text{---} \bullet \text{---} \begin{array}{l} \bullet \\ \bullet \\ \bullet \end{array} \end{array}$ ($q \in R_3$. see GDD 7 of Row 21 in Table B.)
- (l) classical GDD satisfied conditions.

Lemma 3.29. $n = 4$. GDD is not an arithmetic root system if the fixed parameter $q \notin R_4$ and GDD contains subGDD $\begin{array}{c} -\frac{1}{q} \quad -\frac{1}{q} \quad -\frac{1}{q} \\ \bullet \text{---} \bullet \end{array}$ except

- (c) $\begin{array}{c} q \quad q^{-1}q \quad \begin{array}{l} q \\ -1 \\ -1 \end{array} \\ \bullet \text{---} \bullet \text{---} \begin{array}{l} \bullet \\ \bullet \\ \bullet \end{array} \end{array}$ ($q^2 \neq 1$, see GDD 2 of Row 14 in Table B.)
- (d) $\begin{array}{c} -q^{-1}q \quad -q^{-1}q \quad \begin{array}{l} -q^{-1} \\ -1 \\ -1 \end{array} \\ \bullet \text{---} \bullet \text{---} \begin{array}{l} \bullet \\ \bullet \\ \bullet \end{array} \end{array}$, ($q^2 \neq 1$, see GDD 3 of Row 14 in Table B.)
- (e) $\begin{array}{c} q \quad q^{-1}q \quad -\frac{1}{q} \quad -\frac{1}{q} \quad -q^{-1} \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}$ (see, $q^2 \neq 1$, GDD 4 of Row 14 in Table B).
- (h) $\begin{array}{c} q^2 \quad q^{-2}q^2 \quad q^{-2} \quad -\frac{1}{q} \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}$ ($q \in R_6$. see, GDD 5 of Row 9 in Table B).
- (i) $\begin{array}{c} q^2 \quad q^{-2}q \quad q^{-1} \quad -\frac{1}{q} \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}$ ($q \in R_6$. see, GDD 6 of Row 9 in Table B).
- (j) $\begin{array}{c} q^2 \quad q^{-2}q \quad \begin{array}{l} q^2 \\ -1 \\ q \end{array} \\ \bullet \text{---} \bullet \text{---} \begin{array}{l} \bullet \\ \bullet \\ \bullet \end{array} \end{array}$, ($q \in R_6$. see GDD 4 of Row 9 in Table B.)
- (k) classical GDD satisfied conditions.

Lemma 3.30. $n = 4$, GDD is not an arithmetic root system if GDD is







with $q_{44} = q_{33} \neq -1$.

except




(a) Type 5.

Lemma 3.31. $n = 4$. Chain GDD is not an arithmetic root system if $q_{11} = q_{44} = -1$ and there exist different i, j such that $q_{ii} = -1$ and $q_{jj}^A \neq 1$ with $2 \leq i, j \leq 3$, except



(e) A classical GDD satisfied conditions.

Lemma 3.32. $n = 4$. Chain GDD is not an arithmetic root system if there exists i such that $q_{ii} = -1$ and $\tilde{q}_{i,i+1} = \tilde{q}_{i,i-1} \neq -1$, except



$$(d) \quad \begin{array}{ccccccc} q^{-1} & q^{-1} & -1 & -1 \\ \bullet & \bullet & \bullet & \bullet \end{array} \quad (\text{see, } q \in R_3, \text{ GDD 2 of Row 18 in Table B}).$$

$$(e) \quad \begin{array}{ccccccc} -1 & q^{-1} & -1 & -1 & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet \end{array} \quad (\text{see, } q \in R_3, \text{ GDD 3 of Row 20 in Table B}).$$

$$(f) \quad \begin{array}{ccccccc} -1 & q & q^{-1} & -1 & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet \end{array} \quad (\text{see, } q \in R_3, \text{ GDD 4 of Row 20 in Table B}).$$

$$(g) \quad \begin{array}{ccccccc} q & q^{-1} & -1 & -1 & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet \end{array} \quad (\text{see, } q \in R_3, \text{ GDD 6 of Row 21 in Table B}).$$

$$(h) \quad \begin{array}{ccccccc} -1 & -q & -1 & -1 & -q \\ \bullet & \bullet & \bullet & \bullet & \bullet \end{array} \quad (\text{see, } q \in R_4, \text{ GDD 2 of Row 22 in Table B}).$$

$$(i) \quad \begin{array}{ccccccc} -1 & q & -q & -1 & -q \\ \bullet & \bullet & \bullet & \bullet & \bullet \end{array} \quad (\text{see, } q \in R_4, \text{ GDD 3 of Row 22 in Table B}).$$

$$(k) \quad \begin{array}{ccccccc} q^2 & q^{-2} & q^2 & q^{-2} & -1 & q^3 & q^{-3} \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array} \quad (\text{see, } q \in R_5, \text{ GDD 5 of Row 9 in Table B}).$$

$$(l) \quad \begin{array}{ccccccc} q^2 & q^{-2} & q & q^{-1} & -1 & q^3 & q^{-3} \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array} \quad (\text{see, } q \in R_4, \text{ GDD 6 of Row 9 in Table B}).$$

(m) A classical GDD satisfied conditions.

Lemma 3.33. $n = 4$. chain GDD is not an arithmetic root system if there exists only one place where simple chain condition is not satisfied and $q_{22} \neq -1$, $q_{33} \neq -1$ with $\tilde{q}_{23}q_{22} \neq 1$ or $q_{33}\tilde{q}_{34} \neq 1$, or $q_{22}\tilde{q}_{21} \neq 1$, or $q_{33}\tilde{q}_{32} \neq 1$,

except

$$(a) \quad \begin{array}{ccccccc} q^2 & q^{-2} & q^2 & q^{-2} & q & q^{-1} & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array} \quad (\text{see, } q^2, q^3 \neq 1, \text{ GDD 1 of Row 9 in Table B}).$$

$$(b) \quad \begin{array}{ccccccc} -1 & q^{-1} & q & q^{-1} & q & q & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array} \quad (\text{see, } q \in R_3, \text{ GDD 1 of Row 21 in Table B}).$$

$$(c) \quad \begin{array}{ccccccc} q^{-1} & q & q^{-1} & q & q & q^{-1} & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array} \quad (\text{see, } q \in R_3, \text{ GDD 1 of Row 18 in Table B}).$$

$$(d) \quad \begin{array}{ccccccc} -1 & q & q^{-1} & q & q & q^{-1} & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array} \quad (\text{see, } q \in R_3, \text{ GDD 6 of Row 20 in Table B}).$$

(f) classical GDD satisfied conditions.

Lemma 3.34. $n = 4$. chain GDD is not an arithmetic root system if GDD contains a subGDD

$$\begin{array}{c} -1 \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \\ q \quad q^{-1} \end{array}$$

or

$$\begin{array}{c} q \quad -1 \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \\ q^{-1} \quad q^{-1} \end{array}$$

except

(a) $\begin{array}{c} -1 \quad -1 \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ q^{-1} \quad q^{-1} \quad q^{-1} \end{array}$ (see, $q \in R_3$, GDD 3 of Row 20 in Table B).

(b) $\begin{array}{c} -1 \quad -1 \quad q \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ q^{-1} \quad q^{-1} \quad q^{-1} \quad q^{-1} \end{array}$ (see, $q \in R_3$, GDD 5 of Row 20 in Table B).

(c) $\begin{array}{c} -1 \quad -1 \quad q \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ q \quad q^{-1} \quad q \quad q^{-1} \end{array}$ (see, $q \in R_3$, GDD 3 of Row 21 in Table B).

(d) $\begin{array}{c} -1 \quad -1 \quad -q \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ q \quad q^{-1} \quad q^{-1} \quad q^{-1} \end{array}$ (see, $q \in R_3$, GDD 4 of Row 21 in Table B).

(e) $\begin{array}{c} q \quad -1 \quad -1 \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ q^{-1} \quad q^{-1} \quad q^{-1} \quad q^{-1} \end{array}$ (see, $q \in R_3$, GDD 6 of Row 21 in Table B).

(f) $\begin{array}{c} -1 \quad -1 \quad -1 \quad -q \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ -q \quad -q \quad -q \quad -q \end{array}$ (see, $q \in R_4$, GDD 2 of Row 22 in Table B).

(h) $\begin{array}{c} q^2 \quad q^{-2} \quad q^2 \quad q^{-2} \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ q^{-2} \quad q^{-2} \quad q^{-2} \quad q^{-2} \end{array}$ (see, $q \in R_4$, GDD 1 of Row 9 in Table B).

(i) $\begin{array}{c} q^2 \quad q^{-2} \quad q^2 \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ q^{-2} \quad q^{-2} \quad q^{-2} \quad q^{-2} \end{array}$ (see, $q \in R_4$, GDD 2 of Row 9 in Table B).

(j) $\begin{array}{c} q^2 \quad q^{-2} \quad q^2 \quad -1 \quad q^{-3} \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ q^{-2} \quad q^{-2} \quad q^{-2} \quad q^{-2} \quad q^{-2} \end{array}$ (see, $q \in R_4$. GDD 5 of Row 9 in Table B).

(l) classical GDD satisfied conditions.

Lemma 3.35. $n = 4$. GDD which is is not an arithmetic root system if there exists only one place where simple chain conditions are not satisfied, as well as, $\tilde{q}_{12}\tilde{q}_{23} \neq 1$ and $q_{22} = -1$ with $q_{11}, \tilde{q}_{12}, \tilde{q}_{23} \neq -1$, or $\tilde{q}_{32}\tilde{q}_{34} \neq 1$ and $q_{33} = -1$ with $q_{44}, \tilde{q}_{32}, \tilde{q}_{43} \neq -1$, except

(b)

$$\begin{array}{c} q^2 \quad q^{-2}q^2 \quad q^{-2} \quad -1q^3 \quad q^{-3} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad (\text{ see, } q \notin R_6, \text{ GDD 5 of Row 9 in Table B}).$$

$$(c) \quad \begin{array}{c} -q^{-1} \quad -q^{-1} \quad -q^{-1} \quad -1q^{-1} \quad q \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad (\text{ see, } q^2 \neq 1, \text{ GDD 5 of Row 14 in Table B}).$$

$$(d) \quad \begin{array}{c} -q \quad -q^{-1}q \quad -q^{-1}q \quad -1q^{-1}q \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad (\text{ see, } q \in R_3, \text{ GDD 6 of Row 17 in Table B}).$$

$$(e) \quad \begin{array}{c} -1 \quad -q \quad -1q \quad -1q \quad -q \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad (\text{ see, } q \in R_4, \text{ GDD 2 of Row 22 in Table B}).$$

$$(f) \quad \begin{array}{c} -1 \quad q \quad -q \quad -1q \quad -q \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad (\text{ see, } q \in R_4, \text{ GDD 3 of Row 22 in Table B}).$$

$$(g) \quad \begin{array}{c} q \quad q^{-1}q \quad q^{-1} \quad -1q \quad -q^{-1} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad (\text{ see, } q^2 \neq 1, \text{ GDD 1 of Row 14 in Table B}).$$

(k) classical GDD satisfied conditions.

Lemma 3.36. $n = 4$. GDD is not an arithmetic root system if this GDD is not classical and every vertex is not -1 , except

$$(a) \quad \begin{array}{c} -q \quad -q^{-1} \quad -q \quad -1q \quad -q^{-1}q \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad (q \in R_3, \text{ see GDD 1 of Row 17 in Table B}).$$

Lemma 3.37. $n = 4$. Chain GDD is not an arithmetic root system if $q_{11} = -1$, $q_{22}\tilde{q}_{12} \neq 1$ or $q_{44} = -1$, $q_{33}\tilde{q}_{34} \neq 1$, as well as only one vertex is -1 ,

Lemma 3.38. $n = 4$. chain GDD is not an arithmetic root system if there exists only one place where simple chain condition is not satisfied with $\tilde{q}_{21}q_{22} \neq 1$, $q_{22} \neq -1$ or $q_{33}\tilde{q}_{34} \neq 1$, $q_{33} \neq -1$, or $q_{22}\tilde{q}_{23} \neq 1$, $q_{22} \neq -1$ or $q_{33}\tilde{q}_{32} \neq 1$, $q_{33} \neq -1$. except

$$(a) \quad \begin{array}{c} q^2 \quad q^{-2}q^2 \quad q^{-2} \quad q \quad q^{-1} \quad -1 \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad (\text{ see, } q^2, q^3 \neq 1, \text{ GDD 1 of Row 9 in Table B}).$$

$$(b) \quad \begin{array}{c} -1 \quad q^{-1}q \quad q^{-1} \quad q \quad q \quad -1 \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad (\text{ see, } q \in R_3, \text{ GDD 1 of Row 21 in Table B}).$$

$$(c) \quad \begin{array}{c} q^{-1} \quad q \quad q^{-1} \quad q \quad q^{-1} \quad -1 \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad (\text{ see, } q \in R_3, \text{ GDD 1 of Row 18 in Table B}).$$

$$(d) \quad \begin{array}{c} -1 \quad q \quad q^{-1} \quad q \quad q^{-1} \quad -1 \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad (\text{ see, } q \in R_3, \text{ GDD 6 of Row 20 in Table B}).$$

$$(g) \quad \begin{array}{c} q^{-1} \quad q \quad -1q^{-1} \quad q \quad q \quad -1 \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad (\text{ see, } q \in R_3, \text{ GDD 1 of Row 20 in Table B}).$$

(h) $\begin{array}{ccccccc} -1 & q^{-1} & -1 & q & q^{-1} & -1 & \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$ (see, $q \in R_3$, GDD 5 of Row 20 in Table B).

(i) $\begin{array}{ccccccc} -1 & q & -1 & q^{-1} & q & q & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$ (see, $q \in R_3$, GDD 3 of Row 21 in Table B).

(j) $\begin{array}{ccccccc} q & q^{-1} & -1 & q & q^{-1} & -1 & \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$ (see, $q \in R_3$, GDD 5 of Row 21 in Table B).


(l) $\begin{array}{ccccccc} -q & q & -1 & -q & q & q & -q \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$ (see, $q \in R_4$, GDD 1 of Row 22 in Table B).

(o) classical GDD satisfied conditions.

3.4 Affine with circle


Lemma 3.39. (i)

(a)




are affine if and only if


(a) is an affine GDD over circle



(ii)




are affine, if and only if (b) is an affine GDD over arithmetic root system (c)




(iii)

(g)




is an affine GDD if and only if (g) is an affine GDD over circle

(h)



Lemma 3.40. *GDD*



is affine if and only if 123, 234, 341 and 412 are arithmetic root systems.

4 The proof of Theorem 2.1 from subsection 4.1 to subsection 4.55

4.1 GDD 1 of Row 1

Case (i) Omitting 3 and adding 1.

According to Lemma 3.3 (II) we have to consider following cases.

(a).
 $q \quad q^{-1} \quad q \quad q^{-1} \quad q \quad q^{-1} \quad q$
 $, q \in F^* \setminus \{1\}$. GDD 1 of Row 1.

It is an arithmetic root systems

(b).
 $q \quad q^{-2} \quad q^2 \quad q^{-2} \quad q^2 \quad q^{-2} \quad q^2$
 $, q \in F^* \setminus \{1, -1\}$. GDD 1 of Row 2.

It is an arithmetic root system by Type 2

(c).

$$\begin{array}{c}
q^2 \quad q^{-2} \quad q \quad q^{-1}q \quad q^{-1} \quad q \\
\bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet
\end{array}
, q \in F^* \setminus \{1, -1\}. \text{ GDD 1 of Row 3.}$$

It is an arithmetic root system by Type 1.

$$\begin{array}{c}
\text{(d).} \\
-1 \quad q^{-1} \quad q \quad q^{-1}q \quad q^{-1} \quad q \\
\bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet
\end{array}
, q \in F^* \setminus \{1, -1\}. \text{ GDD 1 of Row 4.}$$

It is an arithmetic root systems

$$\begin{array}{c}
\text{(e).} \\
q \quad -q^{-1} \quad -q^{-1} \quad -q^{-1} \quad -q^{-1} \\
\bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet
\end{array}
, q \in R_3. \text{ GDD 1 of Row 12 or Type 4.}$$

It is an arithmetic root system by Type 4

$$\begin{array}{c}
\text{(f) or (1).} \\
-1 \quad q^{-2} \quad q \quad q^{-1}q \quad q^{-1} \quad q \\
\bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet
\end{array}
, q \in R_3 \cup R_6. \text{ GDD 1 of Row 13.}$$

It is affine by Lemma 3.33.

$$\begin{array}{c}
\text{(h).} \\
q \quad -q^{-1-q} \quad -q^{-1-q} \quad -q^{-1-q} \\
\bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet
\end{array}
, q \in R_3. \text{ GDD 5 of Row 16.}$$

It is arithmetic root system by Lemma 3.4(a).

$$\begin{array}{c}
\text{(i) or (2).} \\
q^{-3} \quad q^{-1} \quad q \quad q^{-1}q \quad q^{-1} \quad q \\
\bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet
\end{array}
, q \in R_9. \text{ GDD 1 of Row 18.}$$

It is affine by Lemma 3.4

Case (ii) Omitting 3 and adding 2.

According to Lemma 3.3 (II) we have to consider following cases.

$$\begin{array}{c}
\text{(a).} \\
\begin{array}{c}
\bullet \quad q \\
\quad \quad \quad | \\
\quad \quad \quad q^{-1} \\
\bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\
q \quad q^{-1} \quad q \quad q^{-1}q
\end{array}
\end{array}
, q \in F^* \setminus \{1\}. \text{ GDD 1 of Row 1.}$$

Omitting 1 in whole is an arithmetic root system since symmetry of this GDD. It is similar in following of Case (i) . It is an arithmetic root system by GDD 1 of Row 5 Table B or Type 5.

$$\begin{array}{c}
\text{(b) or (3).} \\
\begin{array}{c}
\bullet \quad q^2 \\
\quad \quad \quad | \\
\quad \quad \quad q^{-2} \\
\bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\
q \quad q^{-2} \quad q^2 \quad q^{-2}q^2
\end{array}
\end{array}
, q \in F^* \setminus \{1, -1\}. \text{ GDD 1 of Row 2.}$$


It is affine by Lemma 3.4.

(c) or (4).  , $q \in F^* \setminus \{1, -1\}$. GDD 1 of Row 3.

Whole is affine by Lemma 3.4.

(d).  , $q \in F^* \setminus \{1, -1\}$. GDD 1 of Row 4.

It is an arithmetic root system, by GDD 4 of Row 13 or Type 5.

(e) or (5).  , $q \in R_3$. GDD 1 of Row 12.

It is affine by Lemma 3.4.

(f) or (6).  , $q \in R_3 \cup R_6$. GDD 1 of Row 13.

It is affine by Lemma 3.4.

(h) or (7).  , $q \in R_3$. GDD 5 of Row 16.

It is affine by Lemma 3.4.


(i) or (8).  , $q \in R_9$. GDD 1 of Row 18.

affine by Lemma 3.4.


Case (iii) Omitting 1 and adding 3. It is the same as Case (i) .

Case (iv) containing circle with four vertexes.

(a) to (a) or (9).



is arithmetic root system,




is affine.

(a) to (b) empty.

(a) to (c) empty.(a) to (d) empty.(a) to (e) empty.


(a) to (f) empty.(a) to (h) empty.(a) to (i) empty.

(b) to (b)




is not arithmetic root system by Lemma 3.2 (V).

(b) to (c)



and $q^4 = q$ is not arithmetic root system by Lemma 3.2(I).


(b) to (d) empty.(b) to (e)



and $-q^{-2} = q$ is not arithmetic root system by Lemma 3.2(I).


(b) to (f) empty.(b) to (h) empty.(b) to (i) empty.

(c) to (c)




is not arithmetic root system by Lemma 3.2 (I).

(c) to (d)



and $q^2 = -1$ is not arithmetic root system by Lemma 3.2 (I)


(c) to (e).




and $-q^{-1} = q^2$ is not arithmetic root system by Lemma 3.2 (I)

(c) to (f) empty.(c) to (h) empty.(c) to (i) empty.

(d) to (d) or (10)




is arithmetic root system by Lemma 3.2 (IV).



is affine.


(d) to (e) empty.

(d) to (f) or (11).



It is not arithmetic root system when $q \in R_6$ by Lemma 3.2(I). It is an arithmetic


root system when $q \in R_3$ by Type 2.



is affine.

(d) to (h) empty.(d) to (i) empty.


(e) to (e).



is not an arithmetic root system by Lemma 3.2 (V).

(e) to (f) empty.(e) to (h) empty.(e) to (i) empty.


(f) to (f).



is not an arithmetic root system by Lemma 3.2(I).


(f) to (h) empty.(f) to (i) empty.

(h) to (h)



is not arithmetic root system by Lemma 3.2 (V).

(h) to (i) empty. (i) to (i)




is not an arithmetic root system by Lemma 3.2 (V).

4.2 GDD 1 of Row 2


Case (i) Omitting 3 and adding 1.

According to Lemma 3.3 (II) we have to consider following cases.




, $q \in F^* \setminus \{1\}$. GDD 1 of Row 1.

It is an arithmetic root system by Type 2.



, $q \in F^* \setminus \{1, -1\}$. GDD 1 of Row 2.


It is affine by Lemma 3.4



$q^4 \neq 1$, $q \in F^* \setminus \{1, -1\}$. GDD 1 of Row 3.


affine by Lemma 3.4.

(d)




, $q \in F^* \setminus \{1, -1\}$. GDD 1 of Row 4.

It is arithmetic root system by Type 2.




, $q \in R_3$. GDD 1 of Row 12.

affine by Lemma 3.4




, $q \in R_3 \cup R_6$. GDD 1 of Row 13.

It is affine by Lemma 3.4



, $q \in R_3$. GDD 5 of Row 16.

It is affine by Lemma 3.4




, $q^2 \in R_9$. GDD 1 of Row 18.

It is affine by Lemma 3.4

Case (ii) Omitting 3 and adding 2.


According to Lemma 3.3 (II) we have to consider following cases.



(a) or (7).

, $q \in F^* \setminus \{1\}$. GDD 1 of Row 1.


Omitting 1 in whole is an arithmetic root system by Type 2. It is affine by Lemma 3.4.



(b).

, $q \in F^* \setminus \{1, -1\}$. GDD 1 of Row 2.

Omitting 1 is not arithmetic root system by Lemma 3.2 (I).



(c).

, $q \in F^* \setminus \{1, -1\}$. GDD 1 of Row 3.

Omitting 1 is not arithmetic root system by Lemma 3.2 (I).

(d).  , $q \in F^* \setminus \{1, -1\}$. GDD 1 of Row 4.

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(I).

(e).  , $q \in R_3$. GDD 1 of Row 20.

Omitting 1 in whole is not any arithmetic root systems by Lemma 3.2(I).

(f).  , $q \in R_3 \cup R_6$. GDD 1 of Row 13.

Omitting 1 in whole is not any arithmetic root systems by Lemma 3.2(I).

(h).  , $q \in R_3$. GDD 5 of Row 16.


Omitting 1 in whole is not any arithmetic root systems by Lemma 3.2(I).

(i).  , $q \in R_9$. GDD 1 of Row 18.


Omitting 1 is not arithmetic root system by Lemma 3.2(I).

Case (iii) Omitting 1 and adding 3.

According to inversion of Type 2 we have to consider following cases.

(a) or (9)  ($q^2 \neq 1$, see GDD 1 of Row 3 in Table A2).

It is affine by Lemma 3.36

(b)  ($q^2 \neq 1$, see GDD 1 of Row 6 in Table A2).

It is an arithmetic root system by GDD 1 of Row 9.

(c) or (10)

$$\begin{array}{c} q^{-8} \quad q^8 \quad q^{-8} \quad q^8 \quad q^{-4} \quad q^4 \quad q^8 \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad (q \in R_9, \text{ see GDD 2 of Row 18 in Table A2}).$$


It is affine by Lemma 3.4.

Case (iv) Containing circle with four vertexes.

According to Lemma 3.39 (ii) we have to consider following cases.

(a) to (a) empty.


(a) to (b) or (11)



and $q^2 = -1$ is arithmetic root system by Lemma 3.2 (IV).


is affine.

(a) to (c)



is not arithmetic root system by Lemma 3.3(II).

(b) to (a).




is an arithmetic root system by Type 1.

is affine.

(b) to (b) empty. (b) to (c). empty.


(c) to (a)



with $q^4 = q$ is an arithmetic root system by Type 2.

is affine.


(c) to (b)



with $q^4 = -1$ is not arithmetic root system by Lemma 3.2(I).

(c) to (c) empty.

(d) to (a) empty. (d) to (b)



is an arithmetic root system by Lemma 3.2(IV) or Type 1

is affine.

(d) to (c). empty. (e) to (a) empty.(e) to (b) empty.(e) to (c) empty. (f) to (a) empty.

(f) to (b)

$$\begin{array}{ccccccc} & q^2 & & q^{-4} & & -1 & & q^{-1}q \\ & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$

is not an arithmetic root system by Lemma 3.2(I).

(f) to (c) empty. (h) to (a) empty.(h) to (b) empty. (h) to (c) empty.

(i) to (a) empty.(i) to (b) empty. (i) to (c) empty.

4.3 GDD 1 of Row 3

Case (i) Omitting 3 and adding 1.

According to Lemma 3.3 (II) we have to consider following cases.

$$\begin{array}{ccccccc} & \text{(a)} \cdot & & & & & & \\ q & q^{-1}q & & q^{-1}q & & q^{-2} & & q^2 \\ \bullet & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$

, $q \in F^* \setminus \{1\}$. GDD 1 of Row 1.

It is an arithmetic root system by Type 1

$$\begin{array}{ccccccc} & \text{(b) or (1)} \cdot & & & & & & \\ q & q^{-2}q^2 & & q^{-2}q^2 & & q^{-4} & & q^4 \\ \bullet & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$

$q^4 \neq 1$,. GDD 1 of Row 2.

It is affine by Lemma 3.4

$$\begin{array}{ccccccc} & \text{(c) or (2)} \cdot & & & & & & \\ q^2 & q^{-2}q & & q^{-1}q & & q^{-2} & & q^2 \\ \bullet & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$

, $q \in F^* \setminus \{1, -1\}$. GDD 1 of Row 3.

It is affine by Lemma 3.36

$$\begin{array}{ccccccc} & \text{(d)} \cdot & & & & & & \\ -1 & q^{-1}q & & q^{-1}q & & q^{-2} & & q^2 \\ \bullet & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$

, $q \in F^* \setminus \{1, -1\}$. GDD 1 of Row 4.

It is an arithmetic root system by Type 1.

$$\begin{array}{ccccccc} & \text{(e) or (3)} \cdot & & & & & & \\ q & -q & & -q^{-1}q & & -q^{-1}q^2 & & q^{-2} \\ \bullet & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$

, $q \in R_3$. GDD 1 of Row 12.

It is affine by Lemma 3.4

$$\begin{array}{ccccccc} & \text{(f) or (4)} \cdot & & & & & & \\ -1 & q^{-2}q & & q^{-1}q & & q^{-2} & & q^2 \\ \bullet & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$

, $q \in R_3 \cup R_6$. GDD 1 of Row 13.

It is affine when $q \in R_3$. by Lemma 3.18. It is affine when $q \notin R_3$. by Lemma 3.37.

(h) or (5).

$$\begin{array}{c} q \quad -q^{-1}q \quad -q^{-1}q \quad q^{-2} \quad q^2 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad , q \in R_3. \text{ GDD 5 of Row 16.}$$

It is affine by Lemma 3.4

$$\begin{array}{c} q^{-3} \text{ (i) or (6).} \\ q^{-1} \quad q \quad q^{-1}q \quad q^{-2} \quad q^2 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad , q \in R_9. \text{ GDD 1 of Row 18.}$$

It is affine by Lemma 3.4

Case (ii) Omitting 1 and adding 2.

According to Type 1 we have to consider following cases.

(a) or (7)

$$\begin{array}{c} q \quad q^{-1} \quad q \\ \bullet \text{---} \bullet \quad \swarrow \quad \searrow \\ \quad \quad \quad q^{-2} \quad q^2 \end{array} \quad , (q^2 \neq 1, \text{ see GDD 1 of Row 3.})$$

Omitting 3 in whole is an arithmetic root system. Whole is affine by Lemma 3.4.

$$\begin{array}{c} -1 \quad q^{-1} \quad q \\ \bullet \text{---} \bullet \quad \swarrow \quad \searrow \\ \quad \quad \quad q^{-2} \quad q^2 \end{array} \quad , (q^2 \neq 1, \text{ Type 1})$$

Omitting 3 in whole is an arithmetic root system, Whole is affine when $q \notin R_3$ by Lemma 3.4. Whole is arithmetic root system when $q \in R_3$ by Lemma 3.4(d).

Case (iii) Omitting 1 and adding 3.

According to Type 2 we have to consider following cases.

$$\begin{array}{c} (a) \\ q \quad q^{-1}q \quad q^{-2} \quad q^2 \quad q^{-2} \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad (q^2 \neq 1, \text{ see GDD 1 of Row 5 in Table A2 or by Type 2).}$$

$$(b) \quad \begin{array}{c} q \quad q^{-1}q \quad q^{-2} \quad q^2 \quad q^2 \quad q^{-2} \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad (q^2 \neq 1, \text{ by Type 2}).$$


Case (iv) . Containing circle with four vertexes.

According to Lemma 3.39 (ii) we have to consider following cases.

(a) to (a) empty.

(a) to (b) empty. (b) to (a) empty.


(b) to (b) or (9)




and $q = q^4$ is arithmetic root system by Type 1.

is affine.


(c) to (a) or (10)




and $-1 = q^2$ is arithmetic root system by Type 2.




is affine.
(c) to (b) or (11)




is an arithmetic root system by Type 2.




is affine.
(d) to (a) or (12).




is an arithmetic root system by Lemma 3.2 (IV).




is affine.
(d) to (b) or (13)



with $q^2 = -1$ is an arithmetic root system by Lemma 3.2 (IV).



is affine.
(e) to (a) empty.
(e) to (b)



with $-q^{-1} = q^2$ is not an arithmetic root system by Lemma 3.3(II)
(f) to (a)

$$\begin{array}{ccccccc} & q & & q^{-2} & & -1 & & q^{-2}q^2 \\ & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$

is not an arithmetic root system by Lemma 3.2(I).

(f) to (b) empty. (h) to (a) empty. (h) to (b) empty. (i) to (a) empty. (i) to (b) empty.

4.4 GDD 1 of Row 4

Case (i) Omitting 3 and adding 1.

$$\begin{array}{ccccccc} & -1 & & (a') & & -1 & & q^{-1}q & & q^{-1} & & q \\ & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$

$q^2 \neq 1$, GDD 2 of Row 4.

It is an arithmetic root systems

$$\begin{array}{ccccccc} & -q & & (a) & & -1 & & q^{-2}q^2 & & q^{-2} & & q^2 \\ & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$

$q^4 \neq 1$, GDD 3 of Row 5.

It is an arithmetic root system by Type 3.

$$\begin{array}{ccccccc} & -1 & & (b) \text{ or } (1) & & -1 & & q^{-2}q^2 & & q^{-2} & & q^2 \\ & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$

$q^2 \neq 1$, GDD 2 of Row 6.

It is an arithmetic root system when $q \notin R_3$ by GDD 2 of Row 9. It is an affine when $q \in R_3$ by Lemma 3.15.

$$\begin{array}{ccccccc} & -1 & & (2) & & -1 & & q^{-3}q^3 & & q^{-3} & & q^3 \\ & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$

$q^2, q^3 \neq 1$, GDD 2 of Row 7.

It is affine by Lemma 3.15

(f) or (3)

$$\begin{array}{ccccccc} & -q & & q^{-1/2} & & -1 & & q^{-3}q^3 & & q^{-3} & & q^3 \\ & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$

$q^6 \neq 1$, GDD 4 of Row 7.

It is affine by Lemma 3.4.

$$\begin{array}{ccccccc} & q^{-1} & & (g) & & -1 & & q^{-1}q & & q^{-1} & & q \\ & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$

, $q^2 \neq 1$, GDD 1 of Row 8.

It is an arithmetic root system by Type 7.

$$\begin{array}{ccccccc} & r & & (h) \text{ or } (4) & & -1 & & q^{-1}q & & q^{-1} & & q \\ & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$

, GDD 1 of Row 9.

Whole is arithmetic root system when $r = -1$ with $q \in R_3$ by Lemma 3.29(h); when $r = q^{-2}$ by Type 2; when $r = -q^{-1}$ and $q^2 \neq 1$. by Lemma 3.35(c); when $q^3 = r^{-2}$ and $q^6 \neq 1$. by Lemma 3.35(b); when $-q^{-1} = r$ by Lemma 3.35(g). Otherwise whole is affine by Lemma 3.35. That is, whole is affine except $r = -1, q \in R_3; r = q^{-2}; r = -q^{-1}, q^2 \neq 1; q^3 = r^{-2}$ and $q^6 \neq 1; -q^{-1} = r$.

$$\begin{array}{c} (i') \text{ or } (5) \\ q \quad q^{-1} \quad -1 \quad q^2 \quad q^{-2} \quad q^2 \quad q^{-2} \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad , q \notin R_2 , q \notin R_3. \text{ GDD 2 of Row 10. or Type 1}$$

It is an arithmetic root system by Lemma 3.35(b). It is affine when $q \notin R_4$ by Lemma 3.35.

$$\begin{array}{c} (k) \\ q \quad q^{-1} \quad -1 \quad q^{-1}q \quad q^{-1} \quad q \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad q^2, q^3 \neq 1, \text{ GDD 1 of Row 10.}$$

It is an arithmetic root system.

$$\begin{array}{c} (l) \\ q^{-2} \quad q^2 \quad -1 \quad q^{-1}q \quad q^{-1} \quad q \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad q^2, q^3 \neq 1, \text{ GDD 2 of Row 10.}$$

It is an arithmetic root system by GDD 1 of Row 13 or by Type 1.

$$\begin{array}{c} (m) \text{ or } (6) \\ q \quad q^{-1} \quad -1 \quad q^{-1}q \quad q^{-1} \quad q \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad , q \in R_3. \text{ GDD 1 of Row 11.}$$

It is affine by Lemma 3.35.

$$\begin{array}{c} (n) \\ q^{-1} \quad -q^{-1} \quad -1 \quad -q \quad -q^{-1} \quad -q \quad -q^{-1} \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad , q \in R_3. \text{ GDD 3 of Row 14.}$$

It is an arithmetic root system by Type 4.

$$\begin{array}{c} (p) \text{ or } (7) \\ -1 \quad q \quad -1 \quad -q^{-1} \quad -q \quad -q^{-1} \quad -q \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad , q \in R_3. \text{ GDD 2 of Row 16.}$$

It is affine by Lemma 3.27

$$\begin{array}{c} (r) \\ q \quad -1 \quad -1 \quad -q^{-1} \quad -q \quad -q^{-1} \quad -q \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad , q \in R_3. \text{ GDD 4 of Row 16.}$$

It is an arithmetic root system by Lemma 3.4(b).

$$\begin{array}{c} (s') \\ -1 \quad q \quad -1 \quad -1 \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad , q \in R_3. \text{ GDD 1 of Row 17.}$$


is empty.

Case (ii) Omitting 1 and adding 2.


According to Lemma 3.3 (I) we have to consider following cases.

(a).  , $q \in F^* \setminus \{1\}$. GDD 1 of Row 1.


Omitting 3 in whole is not an arithmetic root system by Lemma 3.2(I).

(b).  , $q \in F^* \setminus \{1, -1\}$. GDD 1 of Row 2.

Omitting 3 in whole is not an arithmetic root system by Lemma 3.2(I).

(c).  , $q \in F^* \setminus \{1, -1\}$. GDD 1 of Row 3.

Omitting 3 in whole is not an arithmetic root system by Lemma 3.2(I).

(d).  , $q \in F^* \setminus \{1, -1\}$. GDD 1 of Row 4.


Omitting 3 in whole is not an arithmetic root system by Lemma 3.2(I).

(e).  , $q \in R_3$. GDD 1 of Row 20.

Omitting 3 in whole is not an arithmetic root system by Lemma 3.2(I).

(f).  , $q \in R_3 \cup R_6$. GDD 1 of Row 13.

Omitting 3 in whole is not an arithmetic root system by Lemma 3.2(I).

(h).  , $q \in R_3$. GDD 5 of Row 16.


Omitting 3 in whole is not an arithmetic root system by Lemma 3.2(I).

(i).  , $q \in R_9$. GDD 1 of Row 18.

Omitting 3 in whole is not an arithmetic root system by Lemma 3.2(I).

Case (iii) Omitting 1 and adding 3.

According to Lemma 3.3 (II) we have to consider following cases.

(a).  , $q \in F^* \setminus \{1\}$. GDD 1 of Row 1.


(b).  , $q \in F^* \setminus \{1, -1\}$. GDD 1 of Row 2.

(c).  , $q \in F^* \setminus \{1, -1\}$. GDD 1 of Row 3.

(d).  , $q \in F^* \setminus \{1, -1\}$. GDD 1 of Row 4.

(e).  , $q \in R_3$. GDD 1 of Row 12.

(f).  , $q \in R_3 \cup R_6$. GDD 1 of Row 13.


(h).  , $q \in R_3$. GDD 5 of Row 16.

(i).  , $q \in R_9$. GDD 1 of Row 18.

Case (iv) Containing circle with four vertexes.


According to Lemma 3.39 (ii) we have to consider following cases.

(a') to (a) empty.(a') to (b) empty.(a') to (c)



with $q^2 = -1$, is not an arithmetic root system by Lemma 3.2 (I).


(a') to (d) or (8) .



is an arithmetic root system by Type 7.

is affine.


(a') to (e) empty.(a') to (f)




is not an arithmetic root system by Lemma 3.2 (I).

(a') to (i) empty.

(a) to (a) or (9) .




with $q^2 = -q^{-1}$ is an arithmetic root system by Lemma 3.2(VII)(c) or by Lemma 3.3(II)(f).




is affine.

(a) to (b)



with $q = -q^{-1}$ is not an arithmetic root system by Lemma 3.2 (V).


(a) to (c)




with $q^4 = -q^{-1}$ is not an arithmetic root system by Lemma 3.2(I).

(a) to (d) empty.(a) to (e) empty. (a) to (f) empty.(a) to (i) empty.

(b) to (a) or (10)




with $q^2 = -1$ is an arithmetic root system by Lemma 3.2 (XI)(e).




is affine.

(b) to (b) empty.(b) to (c).



with $q^4 = -1$, is not arithmetic root system by Lemma 3.2(I).


(b) to (d) or (11)




is an arithmetic root system by Type 1.

is affine.

(b) to (e) empty.(b) to (f) or (12) .




is not an arithmetic root system by Lemma 3.2(I) when $q \notin R_3$. Whole is an arithmetic root system by Type 2 when $q \in R_3$.



is affine.

(b) to (i) empty.


(d) to (a) or (13)



with $q^3 = -1$ is an arithmetic root system by Lemma 3.2 (XI)(e).


is affine.

(d) to (b) empty.(d) to (c)




with $q^6 = -1$ is not arithmetic root system by Lemma 3.2(I).

(d) to (d) or (14)




is an arithmetic root system by Lemma 3.2 (II)(b) or by Lemma 3.2 (X)(c).



is affine.

(d) to (e) empty.


(d) to (f)



is not arithmetic root system by Lemma 3.2(I).


(d) to (i) empty (f) to (a) empty.

(f) to (b)



and $q^3 = -q^{-2}$ is not arithmetic root system by Lemma 3.2 (V).


(f) to (c)




with $q^6 = -q^{-1}$ is not an arithmetic root system by Lemma 3.2 (I).

(f) to (d) empty.(f) to (e) empty.(f) to (f) empty.(f) to (i) empty. (g) to (a) empty.

(g) to (b) or (15)




with $q = q^{-2}$ is arithmetic root system by Lemma 3.3 (III)(a) or by Type 1.




is affine.

(g) to (c) or (16)




with $q^{-1} = q^2$ is arithmetic root system by Type 2.




is affine.

(g) to (d) empty.(g) to (e) empty. (g) to (f) empty.(g) to (i). empty. (h) to (a) empty.


(h) to (b) or (17)




with $q = r^2$ is arithmetic root system by Type 1.




is affine.
 (h) to (c) or (18)




with $r = q^2$ is arithmetic root system by Type 2.




is affine.
 (h) to (d) or (19) .




with $-1 = r$ is arithmetic root system by Lemma 3.3 (IV).




is affine.
 (h) to (e) or (20)




with $r = -q^{-1}$ is arithmetic root system by Lemma 3.3 (III)(a) or Type 1.




is affine.
 (h) to (f) or (21).



with $r = -1$ is not an arithmetic root system by Lemma 3.2 (I) when $q \notin R_3$; is an arithmetic root system by Lemma 3.2 (I)(n) when $q \in R_3$.



is affine.
 (h) to (i).




with $r = q^{-3}$ is not an arithmetic root system by Lemma 3.3 (III).
 (i') to (a). empty.

(i') to (b). empty.

(i') to (c) or (22).


with $q^{-1} = q^4$ is an arithmetic root system by Type 2.




is affine.

(i') to (d) or (23) empty. (i') to (e). empty. (i') to (f) empty. (i') to (i). empty.

(k) to (a).



is an arithmetic root system by Type 7.




is affine.

(k) to (b) empty. (k) to (c) empty. (k) to (d) empty. (k) to (e) empty. (k) to (f) empty. (k) to (i) empty. (l) to (a), empty.

(l) to (b) or (24)


with $q = q^{-4}$ is arithmetic root system by Type 1.



is affine.


(l) to (c) or (25)

is an arithmetic root system by GDD 1 of Row 2.



is affine.


(l) to (d).




is an arithmetic root system with $q \in R_4$ by Lemma 3.2 (IV).

(l) to (e) empty.


(l) to (f) or (26)




with $-1 = q^{-2}$ is an arithmetic root system by Type 2.



is affine.
 (l) to (i). empty.
 (m) to (a) or (27).




is arithmetic root system.




is affine.
 (m) to (b) empty.

(m) to (c) empty. (m) to (d) empty. (m) to (e), empty. (m) to (f) empty. (m) to (i).empty. (n) to (a) empty. (n) to (b).empty. (n) to (c). empty. (n) to (d) empty. (n) to (e). empty. (n) to (f) empty. (n) to (i). empty. (p) to (a) empty. (p) to (b) empty. (p) to (c) empty.


(p) to (d) or (28)



is an arithmetic root system with $q \in R_6$ by Lemma 3.2 (X)(d).



is affine.
 (p) to (e) empty.
 (p) to (f)



is not an arithmetic root system by Lemma 3.2 (I).

(p) to (i) empty. (r) to (a) empty. (r) to (b) empty. (r) to (c) empty. (r) to (d) empty. (r) to (e) empty. (r) to (f) empty. (r) to (i) empty.

4.5 GDD 2 of Row 4

Case (i) Omitting 3 and adding 1.

According to Lemma 3.3 (IV) we have to consider following cases.

(a').

$$\bullet \xrightarrow{q} \bullet \xrightarrow{q^{-1}} \bullet \xrightarrow{-1} \bullet \xrightarrow{q} \bullet \xrightarrow{-1} \bullet \xrightarrow{q^{-1}} \bullet \xrightarrow{q} \bullet$$
 , $q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 4.

is an arithmetic root systems.

$$\begin{array}{c} (a''). \\ q \quad q^{-2} \quad -1 \quad q^2 \quad -1 \quad q^{-2} \quad q^2 \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad , q \in F^* \setminus \{1, -1\}. \text{ GDD 2 of Row 5.}$$

is an arithmetic root system by Type 2.

$$\begin{array}{c} (a). \\ q^2 \quad q^{-2} \quad -1 \quad q \quad -1 \quad q^{-1} \quad q \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad , q \in F^* \setminus \{1, -1\}. \text{ GDD 2 of Row 6.}$$

is an arithmetic root system by Type 1.

$$\begin{array}{c} (c) \text{ or } (1). \\ q^3 \quad q^{-3} \quad -1 \quad q \quad -1 \quad q^{-1} \quad q \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad , q \in F^* \setminus \{1, -1\}. \quad q \notin R_3, \text{ GDD 2 of Row 7.}$$

is affine by Lemma 3.35.

$$\begin{array}{c} (e). \\ -1 \quad q^{-1} \quad -1 \quad q \quad -1 \quad q^{-1} \quad q \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad , q \in F^* \setminus \{1, -1\}. \text{ GDD 2 of Row 8.}$$

is an arithmetic root systems.

$$\begin{array}{c} (j). \\ q \quad -q \quad -1 \quad -q^{-1} \quad -1 \quad -q \quad -q^{-1} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad , q \in R_3. \text{ GDD 2 of Row 14.}$$

is an arithmetic root system by Type 4 .

$$\begin{array}{c} (k). \\ -1 \quad q \quad -1 \quad q \quad -1 \quad q^{-1} \quad q \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad , q \in R_3. \text{ GDD 2 of Row 15.}$$

is an arithmetic root system by GDD 6 of Row 21.

$$\begin{array}{c} (l) \text{ or } (2). \\ -q \quad -q^{-1} \quad -1 \quad q \quad -1 \quad q^{-1} \quad q \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad , q \in R_3. \text{ GDD 2 of Row 16.}$$

is affine by Lemma 3.15.

$$\begin{array}{c} (n). \\ -1 \quad q \quad -1 \quad -1 \quad -1 \quad -1 \quad -1 \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad , q \in R_3. \text{ GDD 1 of Row 17.}$$

is empty.

$$\begin{array}{c} (n') \text{ or } (3). \\ -1 \quad -1 \quad -1 \quad q \quad -1 \quad q^{-1} \quad q \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad , q \in R_3. \text{ GDD 1 of Row 17.}$$

is affine by Lemma 3.26.

$$\begin{array}{c} \text{(o) or (4).} \\ \bullet \text{---} \frac{-1}{-q} \bullet \text{---} \frac{-1}{q} \bullet \text{---} \frac{-1}{q^{-1}} \bullet \text{---} q \bullet \end{array} \quad , q \in R_3. \text{ GDD 4 of Row 17.}$$

is affine by Lemma 3.15

$$\begin{array}{c} \text{(p) or (5).} \\ \bullet \text{---} \frac{-1}{q} \bullet \text{---} \frac{-1}{-q} \bullet \text{---} \frac{-1}{-q^{-1}-q} \bullet \end{array} \quad , q \in R_3. \text{ GDD 4 of Row 17.}$$

is affine by Lemma 3.15

$$\begin{array}{c} \text{(r).} \\ \bullet \text{---} \frac{q^{-1}}{-q^{-1}-1} \bullet \text{---} -1 \bullet \text{---} -1 \bullet \text{---} -1 \bullet \end{array} \quad , q \in R_3. \text{ GDD 9 of Row 17.}$$

is empty.

Case (ii) Omitting 3 and adding 2.

According to Lemma 3.3 (IV) we have to consider following cases.

$$\text{(a')} \quad \begin{array}{c} \bullet \frac{q}{q^{-1}} \\ | \\ \bullet \text{---} \frac{q}{q^{-1}} \bullet \text{---} -1 \bullet \text{---} q \bullet \text{---} -1 \bullet \end{array} \quad , q \in F^* \setminus \{1, -1\}. \text{ GDD 2 of Row 4.}$$

is an arithmetic root system by Type 5.

$$\text{(a'') or (6).} \quad \begin{array}{c} \bullet \frac{q^2}{q^{-2}} \\ | \\ \bullet \text{---} \frac{q^2}{q^{-2}} \bullet \text{---} -1 \bullet \text{---} q^2 \bullet \text{---} -1 \bullet \end{array} \quad , q \in F^* \setminus \{1, -1\}. \text{ GDD 2 of Row 5.}$$


Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(I) when $q \notin R_4$.
 Omitting 1 in whole is an arithmetic root system by Type 2 when $q \in R_4$. whole is affine when $q \in R_4$ by Lemma 3.4.

$$\text{(a) or (7).} \quad \begin{array}{c} \bullet \frac{q}{q^{-1}} \\ | \\ \bullet \text{---} \frac{q}{q^{-1}} \bullet \text{---} -1 \bullet \text{---} q \bullet \text{---} -1 \bullet \end{array} \quad , q \in F^* \setminus \{1, -1\}. \text{ GDD 2 of Row 6.}$$

Omitting 1 in whole is an arithmetic root system by Lemma 3.2 (IV). affine by Lemma 3.12

$$\text{(c) or (8).} \quad \begin{array}{c} \bullet \frac{q}{q^{-1}} \\ | \\ \bullet \text{---} \frac{q}{q^{-1}} \bullet \text{---} -1 \bullet \text{---} q \bullet \text{---} -1 \bullet \end{array} \quad , q \in F^* \setminus \{1, -1\}. q \notin R_3, \text{ GDD 2 of Row 7.}$$

Omitting 1 in whole is an arithmetic root system by Lemma 3.2 (IV). affine by Lemma 3.25 and by Lemma 3.12

(e) or (9).  , $q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 8.

Omitting 1 in whole is arithmetic root system when $q = q^{-2}$, $q = q^{-3}$, by Lemma 3.2 (X). Whole is affine when $q \in R_4$ by Lemma 3.16; Whole is an arithmetic root system when $q \in R_3$ by Lemma 3.16(d).

(j).  , $q \in R_3$. GDD 2 of Row 14.

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(I).

(10).  , $q \in R_3$. GDD 2 of Row 15.

Omitting 1 in whole is an arithmetic root system. Whole is affine by Lemma 3.21.

(l) or (11).  , $q \in R_3$. GDD 2 of Row 16.

Omitting 1 in whole is an arithmetic root system by Lemma 3.2 (IV). Whole is affine by Lemma 3.12.

(n).  , $q \in R_3$. GDD 1 of Row 17.
is empty.


(n') or (12).  , $q \in R_3$. GDD 1 of Row 17.

Omitting 1 in whole is an arithmetic root system by Lemma 3.2 (IV). Whole is affine by Lemma 3.29.

(o).  , $q \in R_3$. GDD 4 of Row 17.


Omitting 1 in whole is not arithmetic root system by Lemma 3.2(II) or by Lemma 3.2

(X).



, $q \in R_3$. GDD 4 of Row 17.

Omitting 1 in whole is an arithmetic root system by Lemma 3.2(II) (e). Whole is affine by Lemma 3.16.




, $q \in R_3$. GDD 9 of Row 17.


is empty.

Case (iii) Omitting 1 and adding 3.


According to Lemma 3.3 (III) we have to consider following cases.




, $q \in F^* \setminus \{1, -1\}$. GDD 1 of Row 6.




, $q \in F^* \setminus \{1, -1\}$, $q \notin R_3$. GDD 1 of Row 7.




, $q \in F^* \setminus \{1, -1\}$. GDD 3 of Row 8.




, $q \in R_3$. GDD 1 of Row 14.




, $q \in R_3$. GDD 1 of Row 15.



, $q \in R_3$. GDD 1 of Row 16.



, $q \in R_3$. GDD 2 of Row 17.



, $q \in R_3$. GDD 5 of Row 17.

(o).

$$\begin{array}{ccccccc} -1 & -q & q & q^{-1} & -1 & q & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

, $q \in R_3$. GDD 7 of Row 17.

$$\begin{array}{ccccccc} & (a') & & & & & \\ q & q^{-1} & q & q^{-1} & -1 & q & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

GDD 1 of Row 4.

$$\begin{array}{ccccccc} & (a'') & & & & & \\ q & q^{-2} & q^2 & q^{-2} & -1 & q^2 & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

$q^4 \neq 1$, GDD 1 of Row 5.

$$\begin{array}{ccccccc} & (k') & & & & & \\ -1 & q & -1 & -1 & -1 & -1 & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

GDD 1 of Row 17.

is empty.

$$\begin{array}{ccccccc} & (p') & & & & & \\ q^{-1} & -q^{-1} & -1 & -1 & -1 & -1 & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

$q \notin R_3$ GDD 9 of Row 17.


is empty.

Case (iv) Containing circle with four vertexes.

According to Lemma 3.39 (ii) we have to consider following cases.

(a') to (a) empty.(a') to (c) empty.(a') to (e) empty. (a') to (g) empty.(a') to (h) empty. (a') to (j) empty.(a') to (l) empty.(a') to (n) empty. (a') to (o) empty.(a') to (a') or (14)

is arithmetic root system.



is affine.

(a') to (a'') empty.

(a'') to (a)

$$\begin{array}{ccccccc} -1 & q^{-2} & q & q^{-4} & q^2 \\ \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

with $q = q^4$ is not arithmetic root system by Lemma 3.2 (XIV) or by Lemma 3.2(I).

(a'') to (c)

$$\begin{array}{ccccccc} -1 & q^{-2} & q & q^{-6} & q^2 \\ \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

with $q = q^6$ is not arithmetic root system by Lemma 3.2 (I).


(a'') to (e) empty. (a'') to (g)

$$\begin{array}{ccccccc} -1 & q^{-2} & q & -q^{-2} & q^2 \\ \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

with $q = -q^{-2}$ is not an arithmetic root system by Lemma 3.2 (V).


(a'') to (h) empty. (a'') to (j) empty.(a'') to (l) empty.(a'') to (n) empty. (a'') to (o) empty.(a'') to (a') empty.

(a'') to (a'')




is not an arithmetic root system by Lemma 3.2 (V).

(a) to (a) or (15)




is an arithmetic root system by Type 2.




is affine.

(a) to (c) empty.(a) to (e) or (16)




with $q^2 = -1$ is arithmetic root system by Lemma 3.2 (IV).



is affine.

(a) to (g)




wirh $q^2 = -q^{-1}$ is not arithmetic root system by by Lemma 3.3 (III).

(a) to (h) empty.


(a) to (j) empty.(a) to (l) empty.(a) to (n) empty. (a) to (o) empty.

(a) to (a') empty

(a) to (a'') or (17)




wirh $q^4 = q$ is arithmetic root system by Lemma 3.3 (III)(a) or Type ,



is affine.


(c) to (a) empty.

(c) to (c)




is not an arithmetic root system by Lemma 3.2 (I).

(c) to (e) or (18)




with $-1 = q^3$ is arithmetic root system by Lemma 3.2 (IV).



is affine.

(c) to (g) empty. (c) to (h) empty. (c) to (j) empty. (c) to (l) empty.


(c) to (n)



with $-1 = q^3$ is not an arithmetic root system by Lemma 3.2(I).

(c) to (o) empty. (c) to (a') empty.


(c) to (a'') or (19)



with $q = q^6$ is arithmetic root system by Type 1.


is affine.

(e) to (a)




with $-1 = q^2$ is not arithmetic root system by Lemma 3.2 (I).

(e) to (c)




with $-1 = q^3$ is not an arithmetic root system by Lemma 3.2 (I).

(e) to (e) or (20)




is an arithmetic root system when $q = q^{-2}$ or $q = q^{-3}$ by Lemma 3.2 (X).




is affine.

(e) to (g) empty.

(e) to (h) or (21)




is an arithmetic root system by Type 2.



is affine.


(e) to (j) empty.

(e) to (l)




is not an arithmetic root system by Lemma 3.2 (I).

(e) to (n)



is not an arithmetic root system by Lemma 3.2(I).


(e) to (o)



is not an arithmetic root system by Lemma 3.2 (I).

(e) to (a') empty.(e) to (a'') empty.


(j) to (a)



with $-q^{-1} = q^2$ is not an arithmetic root system by Lemma 3.2 (I).

(j) to (c) empty.(j) to (e) empty.

(j) to (g)




is not an arithmetic root system by Lemma 3.2 (V).


(j) to (h) empty. (j) to (j) empty.(j) to (l) empty.(j) to (n) empty. (j) to (o) empty.

(j) to (a') empty

(j) to (a'') or (22)




with $-q^{-2} = q$ is arithmetic root system by Lemma 3.2 (V)(a).




is affine.

(k) to (a) empty.(k) to (c) empty.

(k) to (e) or (23).




is an arithmetic root system by Type 7.



is affine.

(k) to (g) empty


(k) to (h)



is not an arithmetic root system by Lemma 3.2 (I).

(k) to (j) empty.


(k) to (l)



is not an arithmetic root system by Lemma 3.2 (I).

(k) to (n). empty.

(k) to (o).




is not an arithmetic root system by Lemma 3.2(I).

(k) to (a') empty.(k) to (a'') empty. (l) to (a) empty.(l) to (c). empty. (l) to (e) empty.

(l) to (g) empty.(l) to (h) empty.

(l) to (j).




is not an arithmetic root system by Lemma 3.2(I).

(l) to (l) empty.(l) to (n) empty. (l) to (o) empty.(l) to (a') empty.(l) to (a'') empty.

(n') to (a) empty.(n') to (c) empty.

(n') to (e) or (24)




is an arithmetic root system by Lemma 3.2 (IV).

is affine.

(n') to (g) empty.

(n') to (h) or (25)




is an arithmetic root system by Lemma 3.2 (I)(n).

is affine.

(n') to (j) empty.


(n') to (l)




is not an arithmetic root system by Lemma 3.2 (I).

(n') to (n) empty.

(n') to (o) or (26)




is an arithmetic root system by GDD 9 of Row 17 or by Lemma 3.2(I)(c).



is affine.

n') to (a') empty.(n') to (a'') empty. (o) to (a) empty.(o) to (c) empty.


(o) to (e)



is not an arithmetic root system by Lemma 3.2 (X).

(o) to (g) empty.


(o) to (h)



is not an root system by Lemma 3.2 (I).

(o) to (j) empty.


(o) to (l)



is not an arithmetic root system by Lemma 3.2 (I).

(o) to (n) empty.

(o) to (o)




is not an arithmetic root system by Lemma 3.2 (I).

(o) to (a') empty.(o) to (a'') empty.


(p) to (a). empty.

(p) to (c). empty.


(p) to (e) or (27)




is an arithmetic root system by Lemma 3.2 (X)(d).




is affine.
 (p) to (g) empty.
 (p) to (h)




is not an arithmetic root system by Lemma 3.2 (I).
 (p) to (j) empty.
 (p) to (l)



is not an arithmetic root system by Lemma 3.2 (I).
 (p) to (n)



is not an arithmetic root system by Lemma 3.2 (I). . (p) to (o)



is not an arithmetic root system by Lemma 3.2 (I).
 (p) to (a') empty.(p) to (a'') empty.

4.6 GDD 1 of Row 5

Case (i) Omitting 3 and adding 1.

According to Lemma 3.3 (II) we have to consider following cases.

$$\begin{array}{c} (a') \\ -1 \quad -1 \quad q^{-2} \quad q^{-2} \quad q \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad \text{GDD 2 of Row 4.}$$

is an arithmetic root system by Type 2.

$$(a) \text{ or } (1) \quad \begin{array}{c} -q \quad -1 \quad q^{-2} \quad q^{-2} \quad q \\ q^2 \quad q^2 \quad q^2 \quad q^2 \quad q^2 \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad q^2, q^4 \neq 1, \text{ GDD 3 of Row 5.}$$

is affine by Lemma 3.4.

$$\begin{array}{c} (b) \text{ or } (2) \\ -1 \quad -1 \quad q^{-2} \quad q^{-2} \quad q \\ q \quad q \quad q \quad q \quad q \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad \text{GDD 2 of Row 6.}$$

is affine by Lemma 3.4.

$$\begin{array}{c} (d) \text{ or } (3) \\ -1 \quad -1 \quad q^{-6} \quad q^{-6} \quad q^3 \\ q^2 \quad q^2 \quad q^2 \quad q^2 \quad q^2 \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad q^2 \notin R_3 \cup R_2, \text{ GDD 2 of Row 7.}$$

is affine by Lemma 3.4.

$$(f) \text{ or } (4) \quad \begin{array}{ccccccc} & -q^{-2} & & -1 & & q^{-6}q^6 & & q^{-6} & & q^3 \\ & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$

, $q^2 \notin R_3 \cup R_2$, GDD 4 of Row 7.

is affine by Lemma 3.4.

$$(g) \quad \begin{array}{ccccccc} & q^{-2} & & -1 & & q^{-2}q^2 & & q^{-2} & & q \\ & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$

, GDD 1 of Row 8.

is an arithmetic root system by Type 2.

$$(h) \text{ or } (5) \quad \begin{array}{ccccccc} & r^2 & & r^{-2} & & -1 & & q^{-2}q^2 & & q^{-2} & & q \\ & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$

, GDD 1 of Row 9.

is affine by Lemma 3.4.

$$(i') \text{ or } (6) \quad \begin{array}{ccccccc} & q & & q^{-1} & & -1 & & q^2 & & q^{-2}q^{-2} & & q \\ & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$

$q \notin R_3 \cup R_2$, GDD 2 of Row 10.

is affine by Lemma 3.4.

$$(k) \text{ or } (7) \quad \begin{array}{ccccccc} & q^2 & & q^{-2} & & -1 & & q^{-2}q^2 & & q^{-2} & & q \\ & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$

$q^2 \notin R_3 \cup R_2$, GDD 1 of Row 10.

with $q^2 \neq 1$ is affine by Lemma 3.4.

$$(l) \text{ or } (8) \quad \begin{array}{ccccccc} & q^{-1} & & q & & -1 & & q^{-2}q^2 & & q^{-2} & & q \\ & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$

$q \notin R_3 \cup R_2$, GDD 2 of Row 10.

is affine by Lemma 3.4.

$$(m) \text{ or } (9) \quad \begin{array}{ccccccc} & q^2 & & q^{-2} & & -1 & & q^{-2}q^2 & & q^{-2} & & q \\ & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$

, $q^2 \in R_3$. GDD 1 of Row 11.

is affine by Lemma 3.4.

$$(n) \text{ or } (10) \quad \begin{array}{ccccccc} & q^{-2} & & -q^{-2} & & -1 & & -q^{-2}q^{-2} & & -q^{-2} & & iq \\ & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$

, $q \in R_3$. GDD 3 of Row 14.

is affine by Lemma 3.4 .

$$(p) \text{ or } (11) \quad \begin{array}{ccccccc} & -1 & & q^2 & & -1 & & -q^{-2}q^2 & & -q^{-2}iq \\ & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$

, $q \in R_3$. GDD 2 of Row 16.

is affine by Lemma 3.4.

(r) or (12)

$$\begin{array}{c} q^2 \quad -1 \quad -1 \quad -q^{-2} \quad -q^2 \quad -q^{-2}iq \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad , q \in R_3. \text{ GDD 4 of Row 16.}$$

is affine by Lemma 3.4.

$$\begin{array}{c} (s') \\ -1 \quad q \quad -1 \quad -1 \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}$$

is empty.

Case (ii) Omitting 1 and adding 2.

According to Type 2 we have to consider following cases.

(a) or (13)

$$\begin{array}{c} q^2 \quad q^{-2} \quad q^{-2} \quad -1 \\ \bullet \text{---} \bullet \begin{array}{l} \nearrow \bullet \\ \searrow \bullet \end{array} \\ \quad \quad \quad q^2 \quad q^{-2} \quad q \end{array} \quad , \text{ (see GDD 1 of Row 2 or Type 2.)}$$

Omitting 3 in whole is an arithmetic root system by Type 7. Whole is affine by Lemma 3.4.

(b) or (14)

$$\begin{array}{c} -1 \quad q^{-2} \quad q^{-2} \quad -1 \\ \bullet \text{---} \bullet \begin{array}{l} \nearrow \bullet \\ \searrow \bullet \end{array} \\ \quad \quad \quad q^2 \quad q^{-2} \quad q \end{array} \quad , \text{ (see GDD 1 of Row 5 or Type 2.)}$$

Omitting 3 in whole is an arithmetic root system by Type 7. Whole is affine by Lemma 3.4.

Case (iii) Omitting 1 and adding 3.

According to Type 1 we have to consider following cases.

(a)

$$\begin{array}{c} -1 \quad q^{-2}q^2 \quad q^{-2} \quad q \quad q^{-1} \quad q \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad (q^2, q^4 \neq 1, \text{ GDD 1 of Row 3 in Table A2 or Type 1.)$$

(b)

$$\begin{array}{c} -1 \quad q^{-2}q^2 \quad q^{-2} \quad q \quad q^{-1} \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad (q^2, q^4 \neq 1, \text{ GDD 1 of Row 6 in Table A2 or Type 1).}$$

(c)

$$\begin{array}{c} -1 \quad q^8 \quad q^{-8} \quad q^{-4} \quad q^{-3} \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad (q \in R_9, \text{ GDD 2 of Row 18 in Table A2).}$$

Case (iv) Containing circle with four vertexes.


According to Lemma 3.39 (ii) we have to consider following cases.

(a') to (a) empty.

(a') to (b) or (15)

$$\begin{array}{c} -1 \quad q^2 \quad -1 \quad q^{-1}q \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}$$


with $q = q^4$ or $q = q^6$ is an arithmetic root system by Lemma 3.2 (X).




is affine.

(a') to (c). empty.

(a) to (a) or (16)




with $q = -q^{-1}$ is an arithmetic root system by Type 1.




is affine.

(a) to (b) empty. (a) to (c) empty. (b) to (a) empty.

(b) to (b) or (17)




is an arithmetic root system.




is affine.

(b) to (c) empty.

(d) to (a) or (18).




with $-1 = q^3$ is an arithmetic root system by Lemma 3.2 (XI)(e).



is affine.

(d) to (b) empty. (d) to (c) empty.


(f) to (a)



with $-q^{-2} = q^3$ is not an arithmetic root system by Lemma 3.3(II).


(f) to (b) empty.

(f) to (c)




with $ξ^3 = -q^{-2}$ is not an arithmetic root system by Lemma 3.2 (IV).

(g) to (a) or (19).



with $q = q^{-2}$ is arithmetic root system by Type 7.





is affine.

(g) to (b) empty. (g) to (c) empty.

(h) to (a) or (20)



with $q = r^2$ is an arithmetic root system.



is affine.

(h) to (b) or (21)


with $-1 = r^2$ is an arithmetic root system by Lemma 3.2 (IV).




is affine.

(h) to (c)


with $r^{-8} = q^3$ is not an arithmetic root system by Lemma 3.3 (III).



(i') to (a) or (22)




is an arithmetic root system.



is affine.

(i') to (b) empty. (i') to (c) empty. (k) to (a) empty. (k) to (b) empty. (k) to (c) empty. (l) to (a) empty. (l) to (b) empty. (l) to (c) empty. (m) to (a) empty. (m) to (b) empty. (m) to (c) empty. (n) to (a) empty. (n) to (b) empty. (n) to (c) empty. (p) to (a) empty.

(p) to (b)



is not an arithmetic root system by Lemma 3.2 (X).

(p) to (c) empty. (r) to (a) empty. (r) to (b) empty. (r) to (c) empty.

4.7 GDD 2 of Row 5

Case (i) Omitting 3 and adding 1.

According to Lemma 3.3 (IV) we have to consider following cases.

$$\begin{array}{c}
 (a'). \\
 q^2 \quad -1 \quad q^2 \quad -1 \quad q^{-2} \quad q \\
 \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet
 \end{array}
 \quad , q \in F^* \setminus \{1, -1\}. \text{ GDD 2 of Row 4.}$$

is an arithmetic root system by Type 2.

$$\begin{array}{c}
 (a'') \text{ or } (1). \\
 q \quad -1 \quad q^2 \quad -1 \quad q^{-2} \quad q \\
 \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet
 \end{array}
 \quad , q \in F^* \setminus \{1, -1\}. \text{ GDD 2 of Row 5.}$$

is an affine by Lemma 3.4.

$$\begin{array}{c}
 (a) \text{ or } (1). \\
 q^4 \quad -1 \quad q^2 \quad -1 \quad q^{-2} \quad q \\
 \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet
 \end{array}
 \quad , q \in F^* \setminus \{1, -1\}. \text{ GDD 2 of Row 6.}$$

is affine by Lemma 3.4

$$\begin{array}{c}
 (c) \text{ or } (2). \\
 q^6 \quad -1 \quad q^2 \quad -1 \quad q^{-2} \quad q \\
 \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet
 \end{array}
 \quad , q^2 \in F^* \setminus \{1, -1\}. \quad q^2 \notin R_3, \text{ GDD 2 of Row 7.}$$

is affine by Lemma 3.4.

$$\begin{array}{c}
 (e). \\
 -1 \quad -1 \quad q^2 \quad -1 \quad q^{-2} \quad q \\
 \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet
 \end{array}
 \quad , q^2 \in F^* \setminus \{1, -1\}. \text{ GDD 2 of Row 8.}$$

is an arithmetic root system by Type 2.

$$\begin{array}{c}
 (g) \text{ or } (3). \\
 \begin{array}{c}
 -1 \\
 \bullet \\
 / \quad \backslash \\
 q^2 \quad q^{-4} \quad q \\
 \bullet \quad \bullet \quad \bullet \\
 -1 \quad q^2 \quad -1
 \end{array}
 \end{array}
 \quad , q \in F^* \setminus \{1, -1\}, q^3 \neq 1. \text{ GDD 3 of Row 10.}$$

is affine


$$\begin{array}{c}
 (i) \text{ or } (4). \\
 \begin{array}{c}
 -1 \\
 \bullet \\
 / \quad \backslash \\
 q^2 \quad q^2 \quad q^{-2} \quad q \\
 \bullet \quad \bullet \quad \bullet \quad \bullet \\
 -1 \quad q^2 \quad -1
 \end{array}
 \end{array}
 \quad , q \in R_3. \text{ GDD 2 of Row 11.}$$

is affine.

$$\begin{array}{c}
 (j) \text{ or } (5). \\
 q^2 \quad -q^2 \quad -1 \quad -q^{-2} \quad -1 \quad -q^{-2} \quad i q \\
 \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet
 \end{array}
 \quad , q^2 \in R_3. \text{ GDD 2 of Row 14.}$$


is affine by Lemma 3.4.

(k) or (6).




, $q^2 \in R_3$. GDD 2 of Row 15.

is affine by Lemma 3.4.




, $q^2 \in R_3$. GDD 2 of Row 16.

is affine by Lemma 3.4.




, $q \in R_3$. GDD 1 of Row 17.

is empty.




, $q \in R_3$. GDD 1 of Row 17.

is affine by Lemma 3.4.




, $q \in R_3$. GDD 4 of Row 17.

is affine by Lemma 3.4.



, $q \in R_3$. GDD 4 of Row 17.

is affine by Lemma 3.4.




is empty.

Case (ii) Omitting 1 and adding 2.

According to Type 2 and GDD 4 of Row 7 we have to consider following cases.


(a) or (11)



, (see GDD 2 of Row 5 or Type 2.)


Omitting 3 in whole is an arithmetic root system when $q^6 = 1$ by Lemma 3.2 (XI)(b) or Type 2. Whole is affine by Lemma 3.4.

(b) or (12)



Omitting 3 in whole is an arithmetic root system by Type 7. Whole is affine by Lemma 3.4.



(c) or (13)



Omitting 3 in whole is an arithmetic root system when $q \in R_4 \cup R_5 \cup R_7 \cup R_9$ by Lemma 3.2 (X) (XI). Whole is affine by Lemma 3.4.

Case (iii) Omitting 1 and adding 3.

According to inversion of Type 2 we have to consider following cases.




Case (iv) containing circle with four vertexes.


According to Lemma 3.39 (ii) we have to consider following cases.

(a') to (a) empty.

(a') to (b) or (14)




with $q^2 = -q^{-1}$ is an arithmetic root system when $q \in R_6$ by Type 3.




is affine.

(a'') to (a) or (15)



is an arithmetic root system by Lemma 3.3 (II)(f) or by Lemma 3.3 (III)(h) with $q \in R_3$; by Lemma 3.3 (III)(n) with $q \in R_6$.




is affine.

(a'') to (b) empty.

(a) to (a) or (16)


with $q = q^4$ is an arithmetic root system,



is affine.

(a) to (b) empty (c) to (a). empty. (c) to (b). empty. (e) to (a) empty. (e) to (b). empty. (j) to (a) empty.


(j) to (b).



with $q^2 = -q^{-1}$ is not arithmetic root system by Lemma 3.2(I).

(k) to (a) empty. (k) to (b). empty.

(l) to (a)




with $q = -q^{-2}$ is not an arithmetic root system by Lemma 3.3 (III).

(l) to (b). empty. (n') to (a) empty (n') to (b). empty. (o) to (a) empty. (o) to (b). empty. (p) to (a) empty. (p) to (b) empty.

4.8 GDD 3 of Row 5


Case (i) Omitting 3 and adding 1.

According to Lemma 3.3 (III) we have to consider following cases.




, $q^2 \in F^* \setminus \{1, -1\}$. GDD 1 of Row 6.

is affine by Lemma 3.4.




, $q^2 \in F^* \setminus \{1, -1\}$, $q^2 \notin R_3$. GDD 1 of Row 7.

is affine by Lemma 3.4.



, $q^2 \in F^* \setminus \{1, -1\}$. GDD 3 of Row 8.

is an arithmetic root system by Type 3.



, $q \in R_3$. GDD 1 of Row 14.

is affine by Lemma 3.4.

$$\begin{array}{ccccccc} & & \text{(h) or } (20). & & & & \\ -1 & q^2 & q^2 & q^{-2}-1 & q^2 & & -q^{-1} \\ \bullet & \bullet & \bullet & \bullet & \bullet & & \bullet \end{array}$$

, $q \in R_3$. GDD 1 of Row 15.

is affine by Lemma 3.4.

$$\begin{array}{ccccccc} & & \text{(j) or } (21). & & & & \\ -q^2 & -q^{-2} & q^2 & q^{-2}-1 & q^2 & & -q^{-1} \\ \bullet & \bullet & \bullet & \bullet & \bullet & & \bullet \end{array}$$

, $q \in R_3$. GDD 1 of Row 16.

is affine by Lemma 3.4.

$$\begin{array}{ccccccc} & & \text{(l) or } (22). & & & & \\ -1 & -1 & q^2 & q^{-2}-1 & q^2 & & -q^{-1} \\ \bullet & \bullet & \bullet & \bullet & \bullet & & \bullet \end{array}$$

, $q^2 \in R_3$. GDD 2 of Row 17.

is affine by Lemma 3.4.

$$\begin{array}{ccccccc} & & \text{(n) or } (23). & & & & \\ -1 & q^2 & -q^2 & -q^{-2}-1 & -q^2 & & -(iq)^{-1} \\ \bullet & \bullet & \bullet & \bullet & \bullet & & \bullet \end{array}$$

, $q^2 \in R_3$. GDD 5 of Row 17.

is affine by Lemma 3.4.

$$\begin{array}{ccccccc} & & \text{(o) or } (24). & & & & \\ -1 & -q^2 & q^2 & q^{-2}-1 & q^2 & & -q^{-1} \\ \bullet & \bullet & \bullet & \bullet & \bullet & & \bullet \end{array}$$

, $q^2 \in R_3$. GDD 7 of Row 17.

is affine by Lemma 3.4.

$$\begin{array}{ccccccc} & & \text{(a')} & & & & \\ q^2 & q^{-2} & q^2 & q^{-2}-1 & q^2 & & -q^{-1} \\ \bullet & \bullet & \bullet & \bullet & \bullet & & \bullet \end{array}$$

GDD 1 of Row 4.

is an arithmetic root system by Type 3.

$$\begin{array}{ccccccc} & & \text{(a'')} \text{ or } (25) & & & & \\ q & q^{-2} & q^2 & q^{-2}-1 & q^2 & & -q^{-1} \\ \bullet & \bullet & \bullet & \bullet & \bullet & & \bullet \end{array}$$

GDD 1 of Row 5.

is affine by Lemma 3.4.

$$\begin{array}{ccccccc} & & \text{(k')} & & & & \\ -1 & q & -1 & -1 & -1 & & \\ \bullet & \bullet & \bullet & \bullet & \bullet & & \bullet \end{array}$$

GDD 1 of Row 17.

is empty.

(p')

$$\begin{array}{c} q^{-1} \quad -q^{-1} \quad -1 \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}$$

GDD 9 of Row 17.

is empty.

Case (ii) Omitting 1 and adding 2.

According to Type 3 and GDD 4 of Row 7 we have to consider following cases.

(a) or (26)

$$\begin{array}{c} q^2 \quad q^{-2} \quad q^2 \\ \bullet \text{---} \bullet \begin{array}{l} \nearrow \bullet \\ \searrow \bullet \end{array} \\ \quad \quad \quad -1 \quad \quad \quad q^2 \quad -q^{-1} \end{array}, (q^4 \neq 1, \text{ see GDD 3 of Row 5.})$$

Omitting 3 in whole is an arithmetic root system by Lemma 3.2 (IV). Whole is affine by Lemma 3.4.

(b) or (27)

$$\begin{array}{c} -1 \quad q^{-2} \quad q^2 \\ \bullet \text{---} \bullet \begin{array}{l} \nearrow \bullet \\ \searrow \bullet \end{array} \\ \quad \quad \quad -1 \quad \quad \quad q^2 \quad -q^{-1} \end{array}, (q^2 \neq 1, \text{ see Type 3.})$$

Omitting 3 in whole is an arithmetic root system when $q \in R_4 \cup R_6 \cup R_8$ by Lemma 3.2 (X). Whole is affine by Lemma 3.4.

(c) or (28)

$$\begin{array}{c} q^3 \quad q^{-3} \quad q^2 \\ \bullet \text{---} \bullet \begin{array}{l} \nearrow \bullet \\ \searrow \bullet \end{array} \\ \quad \quad \quad -1 \quad \quad \quad q^2 \quad -q^{-1} \end{array}, (\text{ see GDD 4 of Row 7.})$$

Omitting 3 in whole is an arithmetic root system by Lemma 3.2(I V). Whole is affine by Lemma 3.4.

Case (iii) Omitting 1 and adding 3.

According to inversion of Type 3 we have to consider following cases.

(a)

$$\begin{array}{c} q^2 \quad q^{-2} \quad -1 \quad q^2 \quad -q^{-1} \quad q \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad (q \in R_3 \cup R_6, \text{ see GDD 2 of Row 13 in Table A2}).$$

(b)

$$\begin{array}{c} q^2 \quad q^{-2} \quad -1 \quad q^2 \quad -q^{-1} \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad (q \in R_3, \text{ see GDD 4 of Row 15 in Table A2}).$$

(c)

$$\begin{array}{c} q^2 \quad q^{-2} \quad -1 \quad q^2 \quad -q^{-1} \quad -q \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad (q \in R_3, \text{ see GDD 5 of Row 17 in Table A2}).$$

Case (iv) containing circle with four vertexes.

According to Lemma 3.39 (ii) we have to consider following cases.


(a) to (a)

$$\begin{array}{c} q^2 \quad q^{-4} \quad q^4 \quad q^{-1} \quad -q^{-1} \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}$$


with $q = q^4$ is not an arithmetic root system by Lemma 3.2 (I).

(a) to (b). empty. (a) to (c) empty. (c) to (a) empty. (c) to (b) empty. (c) to (c) empty. (e) to (a) empty.

(e) to (b) or (29)




is an arithmetic root system by Type 3.




is affine.

(e) to (c) or (30)




is an arithmetic root system by Lemma 3.2 (IV).



is affine.


(g) to (a) empty. (g) to (b) empty. (g) to (c) empty. (h) to (a) empty.

(h) to (b)



is not an arithmetic root system by Lemma 3.2(I).


(h) to (c)



is not an arithmetic root system by Lemma 3.2(I).

(j) to (a) empty.


(j) to (b)



with $-q^{-1} = q^2$ is not an arithmetic root system by Lemma 3.2(I).


(j) to (c) empty.

(l) to (a)



with $-q^{-2} = q$ is not an arithmetic root system by Lemma 3.3 (I)II.


(l) to (b)



is not an arithmetic root system by Lemma 3.2 (I).

(l) to (c) or (31)


is an arithmetic root system by Lemma 3.2(I)(j).



affine.


(n) to (a) empty. (n) to (b) empty. (n) to (c) empty.

(o) to (a) empty. (o) to (b)




is not an arithmetic root system by Lemma 3.2(I).

(o) to (c) or (32)




is an arithmetic root system by Type 4 with $q \in R_3$.



is affine.

(a') to (a) empty. (a') to (b) empty. (a') to (c) empty.

(a'') to (a)




is not an arithmetic root system by Lemma 3.2 (I).

(a'') to (b) empty. (a'') to (c) empty.

4.9 GDD 1 of Row 6


Case (i) Omitting 3 and adding 1.

According to Lemma 3.3 (I) we have to consider following cases.




GDD 2 of Row 4.

is an arithmetic root system by Type 1.



$q^4 \neq 1$, GDD 3 of Row 5.

is an affine by Lemma 3.4.



$q^4 \neq 1$, GDD 2 of Row 6.

is an affine by Lemma 3.7.

$$\begin{array}{c} \text{(d) or (3)} \\ -1 \quad q \quad -1 \quad q^{-3}q^3 \quad q^{-6} \quad q^6 \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad , q \notin R_3 \cup R_2 \quad q^6 \neq 1, \text{ GDD 2 of Row 7.}$$

is affine by Lemma 3.7.

$$\text{(f) or (4)} \quad \begin{array}{c} -q^{-\frac{1}{2}} \quad -1 \quad q^{-3}q^3 \quad q^{-6} \quad q^6 \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad , q \notin R_3, q^3 \notin R_2 \text{ GDD 4 of Row 7.}$$

is affine by Lemma 3.7.

$$\begin{array}{c} \text{(g)} \\ q^{-1} \quad q \quad -1 \quad q^{-1}q \quad q^{-2} \quad q^2 \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad , \text{ GDD 1 of Row 8.}$$

is an arithmetic root system by Type 1.

$$\begin{array}{c} \text{(h) or (5)} \\ r \quad r^{-1} \quad -1 \quad q^{-1}q \quad q^{-2} \quad q^2 \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad , q, q^{-1} \neq r, \text{ GDD 1 of Row 9.}$$

is an arithmetic root system when $r = q^{-3}$ and $q^2, q^3 \neq 1$, by GDD 6 of Row 9 or by Lemma 3.27(a). Whole is affine when $r \neq q^{-3}$ by Lemma 3.7.

$$\begin{array}{c} \text{(i') or (6)} \\ q \quad q^{-1} \quad -1 \quad q^2 \quad q^{-2}q^4 \quad q^{-4} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad q^4 \neq 1, , q \notin R_3. \text{ GDD 2 of Row 10.}$$

is affine when $q^5 \neq 1$ by Lemma 3.7. Whole is an arithmetic root system when $q^5 = 1$ by Lemma 3.7(a).

$$\begin{array}{c} \text{(k) or (7)} \\ q \quad q^{-1} \quad -1 \quad q^{-1}q \quad q^{-2} \quad q^2 \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad , q \notin R_3. \text{ GDD 1 of Row 10.}$$

is affine when $q^4 \neq 1$ by Lemma 3.7. Whole is an arithmetic root system when $q^4 = 1$ by Lemma 3.7(a).

$$\begin{array}{c} \text{(l) or (8)} \\ q^{-2} \quad q^2 \quad -1 \quad q^{-1}q \quad q^{-2} \quad q^2 \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad , q \notin R_3. \text{ GDD 2 of Row 10.}$$

is affine by Lemma 3.7.

$$\begin{array}{c} \text{(m) or (9)} \\ q \quad q^{-1} \quad -1 \quad q^{-1}q \quad q^{-2} \quad q^2 \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad , q \in R_3. \text{ GDD 1 of Row 11.}$$

is affine by Lemma 3.7.

(n) or (10)

$$\begin{array}{c}
 q^{-1} \quad -q^{-1} \quad -q \quad -q^{-1} \quad q^{-2} \\
 \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet
 \end{array}
 , q \in R_3. \text{ GDD 3 of Row 14.}$$

is affine by Lemma 3.4.

$$\begin{array}{c}
 \text{(p) or (11)} \\
 -1 \quad -1 \quad -q^{-1} \quad q^{-2} \quad q^2 \\
 \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet
 \end{array}
 , q \in R_3. \text{ GDD 2 of Row 16.}$$

is affine by Lemma 3.7.


$$\begin{array}{c}
 \text{(r) or (12)} \\
 q \quad -1 \quad -1 \quad -q^{-1} \quad q^{-2} \quad q^2 \\
 \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet
 \end{array}
 , q \in R_3. \text{ GDD 4 of Row 16.}$$

is affine by Lemma 3.4.

$$\begin{array}{c}
 \text{(s')} \\
 -1 \quad -1 \quad -1 \\
 \bullet \text{---} \bullet \text{---} \bullet
 \end{array}$$

is empty.

(t) or (13)




GDD 8 of Row 17.

is affine.

Case (ii) Omitting 1 and adding 2.

According to inversion of Type 2 we have to consider following cases.

(a) or (14)




Omitting 3 in whole is an arithmetic root system. Whole is an arithmetic root system when $q \in R_3$ by Lemma 3.4(d). Whole is affine when $q \notin R_3$ by Lemma 3.4.

(b) or (15)



Omitting 3 in whole is an arithmetic root system. Whole is affine by Lemma 3.4 with $q \notin R_3$. Whole is an arithmetic root system by Lemma 3.4 (c) with $q \in R_3$.

(c)



Omitting 3 in whole is not an arithmetic root system by Lemma 3.2(I).

Case (iii) Omitting 1 and adding 3.

According to Type 2 we have to consider following cases.


(b) $\begin{array}{c} -1 \quad q^{-1}q \quad q^{-2}q^2 \quad q^{-2} \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad (q^2 \neq 1, \text{ see Type 2}).$

(c) $\begin{array}{c} -1 \quad q^{-1}q \quad q^{-2}q^2 \quad q^{-2} \quad q^2 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad (q^2 \neq 1, \text{ see Type 2}).$


Case (iv) Containing circle with four vertexes.

According to Lemma 3.39 (ii) we have to consider following cases.

(a') to (b) or (15).




is an arithmetic root system by Lemma 3.2 (X)(a) or by Type 1.




is affine.

(a') to (c) or (16).




with $q^2 = -1$ is an arithmetic root system by Type 1 or by Lemma 3.2 (X)(b).



is affine.


(a) to (b) empty.

(a) to (c)




with $q^4 = -q^{-1}$ is not an arithmetic root system by Lemma 3.3(II).

(b) to (b) or (17)




is an arithmetic root system by Lemma 3.2 (X)(b) with $q \in R_3$.




is affine.

(b) to (c)




with $q^4 = -1$ is not an arithmetic root system by Lemma 3.2 (XI)(e).

(d) to (b) or (18).



is an arithmetic root system when $q \in R_5 \cup R_4$ by Lemma 3.2 (X).




is affine.

(d) to (c)

with $q^6 = -1$ is not an arithmetic root system by Lemma 3.2 (XI)(e) and Lemma 3.2 (X).

(f) to (b) empty.


(f) to (c)




with $q^6 = -q^{-1}$ is not an arithmetic root system by Lemma 3.3 (II).

(g) to (b) empty.

(g) to (c) or (19)




with $q^2 = q^{-1}$ is an arithmetic root system by Type 7.




is affine.

(h) to (b) or (20)




with $r = -1$ is an arithmetic root system by Lemma 3.2 (IV).




is affine.

(h) to (c) or (21)




with $r = q^2$ is an arithmetic root system.




is affine.

(i') to (b) empty. (i') to (c) empty. (k) to (b) empty. (k) to (c) empty.

(l) to (b) or (22)




with $-1 = q^{-2}$ is an arithmetic root system.




is affine.
 (l) to (c) or (23)

with $q^2 = q^{-2}$ is an arithmetic root system by Type 7.



is affine.
 (m) to (b) empty. (m) to (c) empty. (n) to (b) empty. (n) to (c) empty.
 (p) to (b) or (24)



is an arithmetic root system by Lemma 3.2 (X)(b).

is affine.
 (p) to (c) empty.
 (r) to (b) empty.(r) to (c) empty.

4.10 GDD 2 of Row 6

Case (i) Omitting 3 and adding 1.

According to Lemma 3.3 (IV) we have to consider following cases.

$$\begin{array}{c}
 (a') \\
 q \quad q^{-1} \quad -1 \quad q \quad -1 \quad q^{-2} \quad q^2 \\
 \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet
 \end{array}
 \quad , q \in F^* \setminus \{1, -1\}. \text{ GDD 2 of Row 4.}$$

is an arithmetic root system by Type 1.

$$\begin{array}{c}
 (a'') \\
 q \quad q^{-2} \quad -1 \quad q^2 \quad -1 \quad q^{-4} \quad q^4 \\
 \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet
 \end{array}
 \quad , q \in F^* \setminus \{1, -1\}. \text{ GDD 2 of Row 5.}$$

is affine by Lemma 3.4.

$$\begin{array}{c}
 (a) \text{ or } (1) \\
 q^2 \quad q^{-2} \quad -1 \quad q \quad -1 \quad q^{-2} \quad q^2 \\
 \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet
 \end{array}
 \quad , q \in F^* \setminus \{1, -1\}. \text{ GDD 2 of Row 6.}$$

is affine by Lemma 3.7.
 (c) or (2).

$$\begin{array}{ccccccc} q^3 & q^{-3} & -1 & q & -1 & q^{-2} & q^2 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

, $q \in F^* \setminus \{1, -1\}$. $q \notin R_3$, GDD 2 of Row 7.

is affine by Lemma 3.7.

$$\begin{array}{ccccccc} \text{(e).} & & & & & & \\ -1 & q^{-1} & -1 & q & -1 & q^{-2} & q^2 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

, $q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 8.

is an arithmetic root system arithmetic root system by Type 1.

$$\begin{array}{ccccccc} \text{(j) or (4).} & & & & & & \\ q & -q & -1 & -q^{-1} & 1 & q^2 & q^{-2} \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

, $q \in R_3$. GDD 2 of Row 14.

is affine by Lemma 3.4.

$$\begin{array}{ccccccc} \text{(k) or (5).} & & & & & & \\ -1 & q & -1 & q & -1 & q^{-2} & q^2 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

, $q \in R_3$. GDD 2 of Row 15.

is affine by Lemma 3.7.

$$\begin{array}{ccccccc} \text{(l) or (6).} & & & & & & \\ -q & -q^{-1} & 1 & q & -1 & q^{-2} & q^2 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

, $q \in R_3$. GDD 2 of Row 16.

is affine by Lemma 3.7.

$$\begin{array}{ccccccc} \text{(n).} & & & & & & \\ -1 & q & -1 & -1 & -1 & q^{-2} & q^2 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

, $q \in R_3$. GDD 1 of Row 17.

is empty.

$$\begin{array}{ccccccc} \text{(n') or (7).} & & & & & & \\ -1 & -1 & -1 & q & -1 & q^{-2} & q^2 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

, $q \in R_3$. GDD 1 of Row 17.

is affine by Lemma 3.7.

$$\begin{array}{ccccccc} \text{(o) or (8).} & & & & & & \\ -1 & -q & -1 & q & -1 & q^{-2} & q^2 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

, $q \in R_3$. GDD 4 of Row 17.

is affine by Lemma 3.7.

$$\begin{array}{ccccccc} \text{(p) or (9).} & & & & & & \\ -1 & q & -1 & -q & -1 & q^{-2} & q^2 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

, $q \in R_3$. GDD 4 of Row 17.

is affine by Lemma 3.7.

(r).

$$\begin{array}{c} q^{-1} -q^{-1} -1 -1 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}$$

is empty.

Case (ii) Omitting 3 and adding 2.

According to Lemma 3.3 (IV) we have to consider following cases.

$$(a') \text{ or } (10). \quad \begin{array}{c} q^2 \\ \bullet \\ | \\ \bullet \\ q \text{---} q^{-1} -1 q \text{---} -1 \end{array} \quad , q \in F^* \setminus \{1, -1\}. \text{ GDD 2 of Row 4.}$$

Omitting 1 in whole is an arithmetic root system by Lemma 3.2 (IV). Whole is an arithmetic root system when $q \in R_3$ by GDD 9 of Row 20. Whole is affine when $q \notin R_3$ by Lemma 3.12.

$$(a'') \text{ or } (11). \quad \begin{array}{c} q^4 \\ \bullet \\ | \\ \bullet \\ q \text{---} q^{-2} -1 q^2 \text{---} -1 \end{array} \quad , q \in F^* \setminus \{1, -1\}. \text{ GDD 2 of Row 5.}$$

Omitting 1 in whole is an arithmetic root system when $q \in R_7$ by Lemma 3.2(I)(h). Omitting 1 in whole is an arithmetic root system when $q \in R_6$ by Type 2. Whole is affine when $q \in R_6 \cup R_7$ by Lemma 3.4.

$$(a) \text{ or } (12). \quad \begin{array}{c} q^2 \\ \bullet \\ | \\ \bullet \\ q^2 \text{---} q^{-2} -1 q \text{---} -1 \end{array} \quad , q \in F^* \setminus \{1, -1\}. \text{ GDD 2 of Row 6.}$$

Omitting 1 in whole is an arithmetic root system by Lemma 3.2 (IV). Whole is affine checked step by step and by Lemma 3.12.

$$(c) \text{ or } (13). \quad \begin{array}{c} q^2 \\ \bullet \\ | \\ \bullet \\ q^3 \text{---} q^{-3} -1 q \text{---} -1 \end{array} \quad , q \in F^* \setminus \{1, -1\}. \quad q \notin R_3, \text{ GDD 2 of Row 7.}$$

Omitting 1 in whole is an arithmetic root system by Lemma 3.2 (IV). Whole is affine checked step by step or by Lemma 3.12.

$$(e) \text{ or } (14). \quad \begin{array}{c} q^2 \\ \bullet \\ | \\ \bullet \\ -1 \text{---} q^{-1} -1 q \text{---} -1 \end{array} \quad , q \in F^* \setminus \{1, -1\}. \text{ GDD 2 of Row 8.}$$

Omitting 1 in Whole is arithmetic root system when $q \in R_4 \cup R_5 \cup R_3$ by Lemma 3.2 (X). Whole is an arithmetic root system when $q \in R_3$ by Lemma 3.16(d). Whole is affine when $q \notin R_3$ by Lemma 3.12.

(j).  , $q \in R_3$. GDD 2 of Row 14.

Omitting 1 in whole is not arithmetic root system by Lemma 3.2(I).

(k) or (15).  , $q \in R_3$. GDD 2 of Row 15.

Omitting 1 in whole is an arithmetic root system by Type 1. Whole is affine by Lemma 3.16.

(l) or (16).  , $q \in R_3$. GDD 2 of Row 16.

Omitting 1 in whole is an arithmetic root system by Lemma 3.3 (IV). Whole is affine by Lemma 3.12.

(n).  , $q \in R_3$. GDD 1 of Row 17.

is empty.

(n') or (17).  , $q \in R_3$. GDD 1 of Row 17.

Omitting 1 in whole is an arithmetic root system by Lemma 3.3 (IV). Whole is affine by Lemma 3.12.

(o).  , $q \in R_3$. GDD 4 of Row 17.

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2 (X).

(p) or (18).  , $q \in R_3$. GDD 4 of Row 17.

Omitting 1 in whole is an arithmetic root system by Lemma 3.2 (X)(b). Whole is affine by Lemma 3.16.

$$(r). \quad \begin{array}{ccccccc} & q^{-1} & -q^{-1} & 1 & -1 & -1 & \\ & \bullet & \text{---} & \bullet & \text{---} & \bullet & \end{array} \quad , q \in R_3. \text{ GDD 9 of Row 17.}$$

is empty.

Case (iii) Omitting 1 and adding 3.

According to Lemma 3.3 (III) we have to consider following cases.

$$(a). \quad \begin{array}{ccccccc} q^4 & & & & & & \\ q^{-4} & q^2 & q^{-2} & -1 & q & -1 & \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array} \quad , q \in F^* \setminus \{1, -1\}. \text{ GDD 1 of Row 6.}$$

$$(c). \quad \begin{array}{ccccccc} q^6 & & & & & & \\ q^{-6} & q^2 & q^{-2} & -1 & q & -1 & \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array} \quad q^2 \neq 1, q^2 \notin R_3 \cup R_2. \text{ GDD 1 of Row 7.}$$

$$(e). \quad \begin{array}{ccccccc} -1 & & & & & & \\ q^{-2} & q^2 & q^{-2} & -1 & q & -1 & \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array} \quad , q \in F^* \setminus \{1, -1\}. \text{ GDD 3 of Row 8.}$$

$$(g). \quad \begin{array}{ccccccc} q^2 & & & & & & \\ -q^2 & -q^{-2} & -q^2 & -1 & (iq)^{-1} & -1 & \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array} \quad , q \in R_3. \text{ GDD 1 of Row 14.}$$

$$(h). \quad \begin{array}{ccccccc} -1 & & & & & & \\ q^2 & q^2 & q^{-2} & -1 & q & -1 & \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array} \quad , q \in R_3. \text{ GDD 1 of Row 15.}$$

$$(j). \quad \begin{array}{ccccccc} -q^2 & & & & & & \\ q^{-2} & q^2 & q^{-2} & -1 & q & -1 & \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array} \quad , q \in R_3. \text{ GDD 1 of Row 16.}$$

$$(l). \quad \begin{array}{ccccccc} -1 & & & & & & \\ -1 & q^2 & q^{-2} & -1 & q & -1 & \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array} \quad , q \in R_3. \text{ GDD 2 of Row 17.}$$

$$(n). \quad \begin{array}{ccccccc} -1 & & & & & & \\ q^2 & -q^2 & -q^{-2} & -1 & iq & -1 & \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array} \quad , q \in R_3. \text{ GDD 5 of Row 17.}$$

$$(o). \quad \begin{array}{ccccccc} -1 & & & & & & \\ -q^2 & q^2 & q^{-2} & -1 & q & -1 & \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array} \quad , q \in R_3. \text{ GDD 7 of Row 17.}$$

(a').

$$\begin{array}{c} q^2 \quad q^{-2} \quad q^2 \quad q^{-2}-1 \quad q \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array}$$

$q^2 \neq 1$, GDD 1 of Row 4.

$$\begin{array}{c} (a'') \\ q \quad q^{-2} \quad q^2 \quad q^{-2}-1 \quad q \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array}$$

GDD 1 of Row 5.

$$\begin{array}{c} (k') \\ -1 \quad q \quad -1 \quad -1 \quad -1 \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array}$$

GDD 1 of Row 17.

empty

$$\begin{array}{c} (p') \\ q^{-1} \quad -q^{-1}-1 \quad -1 \quad -1 \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array}$$

GDD 9 of Row 17.

is empty.

Case (iv) containing circle with four vertexes.

According to Lemma 3.39 (ii) we have to consider following cases.

(a') to (a) or (19)

$$\begin{array}{c} -1 \quad q^{-1} \quad q \quad q^{-4}q^2 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}$$

with $q = q^4$ is an arithmetic root system by Lemma 3.2(I) or Type 2.

$$\begin{array}{c} q \\ q^{-1} \quad q^{-4} \\ -1 \quad q \quad -1 \quad q^{-2}q^2 \end{array}$$

is affine.

(a') to (c)

$$\begin{array}{c} -1 \quad q^{-1} \quad q \quad q^{-6}q^2 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}$$

with $q = q^6$ is not an arithmetic root system by Lemma 3.2(I).

(a') to (e) empty.(a') to (g) empty.(a') to (h) empty. (a') to (j) empty.(a') to (l)

empty.(a') to (n) empty.(a') to (o) empty. (a') to (a') empty

(a') to (a'') or (20)

$$\begin{array}{c} -1 \quad q^{-1} \quad q \quad q^{-2}q^2 \\ \bullet \text{---} \bullet \text{---} \bullet \\ q \\ q^{-1} \quad q^{-2} \\ -1 \quad q \quad -1 \quad q^{-2}q^2 \end{array}$$

is an arithmetic root system by Type 1.

is affine.

(a'') to (a)

$$\begin{array}{c} -1 \quad q^{-2} \quad q \quad q^{-8}q^4 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}$$

with $q = q^8$ is not an arithmetic root system by Lemma 3.2(I).


(a'') to (c)

$$\begin{array}{c} -1 \quad q^{-2} \quad q \quad q^{-12}q^4 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}$$

with $q = q^{12}$ is not an arithmetic root system by Lemma 3.2(I).

(a'') to (e) empty. (a'') to (g) empty.(a'') to (h) empty. (a'') to (j) empty.(a'') to (l) empty.(a'') to (n) empty. (a'') to (o) empty.

(a'') to (a') or (21)




with $q^4 = q$ is an arithmetic root system by Lemma 3.3(II)(f).

is affine.

(a'') to (a'') empty. (a) to (a) empty.


(a) to (c) or (22).



with $q^6 = q^2$ is an arithmetic root system.

is affine.

(a) to (e) or (23).




with $q^2 = -1$ is an arithmetic root system.

is affine.

(a) to (g) empty.(a) to (h) empty. (a) to (j) empty.(a) to (l) empty.(a) to (n) empty.
(a) to (o) empty.

(a) to (a') or (24)




is arithmetic root system,


is affine.

(a) to (a'') empty. (c) to (a) empty.(c) to (c) empty.

(c) to (e) or (25)



with $q^3 = -1$ is an arithmetic root system by Lemma 3.2 (IV).




is affine.

(c) to (g) empty.(c) to (h) empty. (c) to (j) empty.(c) to (l) empty.(c) to (n) empty.

(c) to (o) empty.

(c) to (a') or (26).




is an arithmetic root system by Type 7.

is affine.


(c) to (a'') empty.

(e) to (a)




with $-1 = q^4$ is not an arithmetic root system by Lemma 3.2 (I).

(e) to (c)




with $-1 = q^6$ is not an arithmetic root system by Lemma 3.2(I).

(e) to (e) or (27)




is an arithmetic root system when $q \in R_4 \cup R_5 \cup R_3$ by Lemma 3.2 (X).



is affine.


(e) to (g) empty.(e) to (h)



is not an arithmetic root system by Lemma 3.2 (I).


(e) to (j) empty.

(e) to (l)




is not an arithmetic root system by Lemma 3.2 (I).

(e) to (n)




with $q \in R_3$ is not an arithmetic root system by Lemma 3.2 (I).

(e) to (o)




is not an arithmetic root system by Lemma 3.2 (I).

(e) to (a') or (28)




with $-1 = q^2$ is an arithmetic root system by Lemma 3.2 (XI)(e).



is affine.


(e) to (a'') empty. (j) to (a) empty. (j) to (c) empty. (j) to (e) empty. (j) to (g) empty.
(j) to (h) empty. (j) to (j) empty. (j) to (l) empty. (j) to (n) empty. (j) to (o) empty.
(j) to (a')




with $-q^{-1} = q^2$ is not an arithmetic root system by Lemma 3.3(II).

(j) to (a'') empty. (k) to (a) empty. (k) to (c) empty.

(k) to (e) or (29)




is an arithmetic root system by Lemma 3.2 (X) or by Type 1.




is affine.

(k) to (g) empty.

(k) to (h) or (30)




is an arithmetic root system by Type 2.




(k) to (j) empty.

(k) to (l)




is not an arithmetic root system by Lemma 3.2(I).

(k) to (n)




is not an arithmetic root system by Lemma 3.2 (I).

(k) to (o)




is not an arithmetic root system by Lemma 3.2 (I).

(k) to (a') empty. (k) to (a'') empty. (l) to (a) empty. (l) to (c) empty. (l) to (e) empty. (l) to (g) empty. (l) to (h) empty. (l) to (j) empty. (l) to (l) empty. (l) to (n) empty. (l) to (o) empty. (l) to (a') empty. (l) to (a'') empty. (n') to (a) empty. (n') to (c) empty. (n') to (e) or (30)1




is an arithmetic root system by Lemma 3.2 (IV).




is affine.

(n') to (g) empty.

(n') to (h) or (32)




is an arithmetic root system by Lemma 3.2(I)(n).



is affine.


(n') to (j) empty.

(n') to (l)




is not an arithmetic root system by Lemma 3.2(I).

(n') to (n)




is not an arithmetic root system by Lemma 3.2(I).

(n') to (o) or (33)




is an arithmetic root system by Lemma 3.2(I)(n).




is affine. . (n') to (a') empty. (n') to (a'') empty. (o) to (a) empty. (o) to (c) empty.


(o) to (e) or (34)




is an arithmetic root system by Lemma 3.2 (X)(b).




is affine.
 (o) to (g) empty.
 (o) to (h)




is not an arithmetic root system by Lemma 3.2 (I).
 (o) to (j) empty.
 (o) to (l)




is not an arithmetic root system by Lemma 3.2 (I).
 (o) to (n)




is not an arithmetic root system by Lemma 3.2 (I).
 (o) to (o) or (35)




is an arithmetic root system by Type 4.




is affine.
 (o) to (a') empty. (o) to (a'') empty. (p) to (a) empty. (p) to (c) empty.
 (p) to (e) or (36)



is an arithmetic root system by Lemma 3.2 (X)(b).




is affine.
 (p) to (g) empty.
 (p) to (h)



is not an arithmetic root system by Lemma 3.2 (I).
 (p) to (j) empty.

p) to (l)



is not an arithmetic root system by Lemma 3.2 (I).

(p) to (n) empty. (p) to (o) empty. (p) to (a') empty. (p) to (a'') empty.

4.11 GDD 3 of Row 6


Case (i) Omitting 3 and adding 1.

According to Lemma 3.3 (III) we have to consider following cases.



Omitting 2 in whole is an arithmetic root system by symmetry of this GDD. It is similar in following of Case (i) . Whole is affine by Lemma 3.4.

(b) or (2),




Omitting 2 in whole is an arithmetic root system by Type 7. Omitting 1 in whole is an arithmetic root system when $q \in R_3 \cup R_6$ by Lemma 3.3 (IV)(k). Whole is affine when $q \in R_3$.




is affine by Lemma 3.4.

(d).




Omitting 2 in whole is not an arithmetic root system by Lemma 3.3 (III).




, $q \in F^* \setminus \{1, -1\}$. GDD 3 of Row 8.

is an arithmetic root system by GDD 4 of Row 12, or Type 6.




, $q \in R_3$. GDD 1 of Row 14.

is affine by Lemma 3.4.




, $q \in R_3$. GDD 1 of Row 15.

is affine by Lemma 3.4.



, $q \in R_3$. GDD 3 of Row 15.


Omitting 1 in whole is an arithmetic root system by Lemma 3.2 (X)(b). Omitting 2 in whole is an arithmetic root system by Type 7. Whole is affine.



, $q \in R_3$. GDD 1 of Row 16.


is affine by Lemma 3.4.

(k).



, $q \in R_3$. GDD 3 of Row 16.


Omitting 1 in whole is not an arithmetic root system by Lemma 3.3 (XI).



, $q \in R_3$. GDD 2 of Row 17.

is affine by Lemma 3.4.

(m) or (9),



Omitting 1 in whole is an arithmetic root system by Lemma 3.2 (XI)(c). Omitting 2 in whole is an arithmetic root system by Lemma 3.3 (III)(j). Whole is affine.




is affine by Lemma 3.4.



is affine by Lemma 3.4.


(p) or (12),




Omitting 2 in whole is an arithmetic root system by Lemma 3.3 (III)(o). Omitting 1 in whole is an arithmetic root system by Lemma 3.3 (V)(a). Whole is affine.



is an arithmetic root system by GDD 4 of Row 8 or by Type 6.




is affine by Lemma 3.4.



GDD 1 of Row 17.

is empty.




GDD 9 of Row 17.

is empty.


Case (ii) Omitting 3 and adding 2.

According to Lemma 3.3 (I) we have to consider following cases.



GDD 2 of Row 4.


Omitting 1 in whole is not arithmetic root system when $q^3 \neq 1$ by Lemma 3.3 (IV). Whole is an arithmetic root system when $q^3 = 1$ by GDD 8 of Row 20 or by Lemma 3.11(d).



(a) or (14)

$q^2 \neq -1$, GDD 3 of Row 5.

Omitting 1 is an arithmetic root system when $q \in R_6$ by Type 3. Whole is affine by Lemma 3.4.




(b) or (15)

GDD 2 of Row 6.

Omitting 1 in whole is an arithmetic root system when $q \in R_3$ by Lemma 3.2 (XI) (a), Whole is affine by Lemma 3.11.

(c) or (16)




GDD 3 of Row 6.

Omitting 1 in whole is an arithmetic root system when $q \in R_3 \cup R_5$ by Lemma 3.2 (XI)(a). Omitting 2 in whole is an arithmetic root system by Type 7. Whole is affine.


(d) or (17)  $q^3 \notin R_2$, GDD 2 of Row 7.

Omitting 1 in whole is an arithmetic root system when $q \in R_7$ by Lemma 3.2 (XI). Whole is affine by Lemma 3.11.

(e) or (18)

 $q \notin R_3$, GDD 3 of Row 7.

Omitting 1 in whole is an arithmetic root system when $q \in R_6 \cup R_4 \cup R_5$ by Lemma 3.2 (XI). Omitting 2 in whole is an arithmetic root system when $q \in R_4 \cup R_3 \cup R_6$ by Type 1 and by Type 7 and by Lemma 3.3 (III)(n). Whole is affine when $q \in R_4 \cup R_3 \cup R_6$.

(f)  , $q \notin R_3$, GDD 4 of Row 7.

Omitting 1 in whole is an arithmetic root system when $q \in R_8$ by Type 3. Omitting 1 in whole is not an arithmetic root system when $q \notin R_8$ by Lemma 3.2(I)(I). Whole is affine when $q \in R_8$ by Lemma 3.4.


(g)  , GDD 1 of Row 8.

Omitting 1 in whole is an arithmetic root system when $q \in R_3 \cup R_5 \cup R_7 \cup R_6 \cup R_4$ by Lemma 3.2 (X) and by Lemma 3.2 (XI)(e). Whole is affine when $q \in R_4 \cup R_5 \cup R_7 \cup R_6$ by Lemma 3.11; Whole is an arithmetic root system when $q \in R_3$ by Lemma 3.11(c).

(h)  , GDD 1 of Row 9.


Omitting 1 in whole is an arithmetic root system when $q^2 = r$, $q \notin R_3$ or $q^4 = r$, $q \notin R_3 \cup R_5$ or $q^6 = r$, $q \notin R_5 \cup R_7 \cup R_6$ or $-q^2 = r$, $q \in R_6$, or $r = -1$, $q \in R_4 \cup R_6 \cup R_8 \cup R_{12} \cup R_3$, or $q \in R_4$ by Lemma 3.2 (IV), by Lemma 3.2 (X) and by Lemma 3.2 (XI)(e). Whole is affine when $r \neq q^2$ by Lemma 3.11. Whole is an arithmetic root system when $r = q^2$ by Lemma 3.11(a)(c).

(i')




, $q \notin R_3$. GDD 2 of Row 10.

Omitting 1 in whole is an arithmetic root system when $q \in R_5 \cup R_9 \cup R_8 \cup R_{13}$ by Lemma 3.3(I) and by Lemma 3.2 (IV). Whole is an arithmetic root system when $q \in R_5$ by Lemma 3.11(a). Whole is affine when $q \notin R_5$ by Lemma 3.11.




, $q \notin R_3$. GDD 1 of Row 10.

Omitting 1 in whole is an arithmetic root system when $q \in R_5 \cup R_4$ by Lemma 3.3(I)(d) and by Lemma 3.2 (IV). Whole is affine by Lemma 3.11.




, $q \notin R_3$. GDD 2 of Row 10.

Omitting 1 in whole is an arithmetic root system when $q \in R_4 \cup R_6 \cup R_8$ by Lemma 3.2 (X). Whole is an arithmetic root system when $q \in R_4$ by Lemma 3.11 (a). Whole is affine when $q \notin R_4$. by Lemma 3.11.



, $q \in R_3$. GDD 1 of Row 11.


Omitting 1 in whole is an arithmetic root system when $q \in R_3$. Whole is affine by Lemma 3.11.



, $q \in R_3$. GDD 3 of Row 14.


Omitting 1 in whole is not arithmetic root system by Lemma 3.2(I).

(o)



$q \in R_3$, GDD 3 of Row 15.


Omitting 2 in whole is an arithmetic root system by Type 7. Omitting 1 in whole is an arithmetic root system by Lemma 3.2 (X). Whole is affine.



, $q \in R_3$. GDD 2 of Row 16.

Omitting 1 in whole is an arithmetic root system when $q \in R_3$ by Lemma 3.2 (XI).
Whole is affine by Lemma 3.11.

(q)



Omitting 1 in whole is not an arithmetic root system by Lemma 3.2 (XI).




Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(I).




is empty.

(s)



Omitting 1 in whole is not an arithmetic root system by Lemma 3.2 (X).

(t)



Omitting 1 in whole is an arithmetic root system by Lemma 3.3 (V)(a). Omitting 2 in whole is an arithmetic root system by Lemma 3.3 (III)(o). Whole is affine.

Case (iii) Omitting 1 and adding 3 is the same as Case (ii) .


Case (iv) Containing circle with four vertexes.

This case can be ignored since it is over a cycle.

Case (v) complete diagram containing circle with four vertexes.


According to Lemma 3.40 we have to consider following cases.

(b) in Case (i)




If 134 is arithmetic root system, then $\tilde{q}_{34} = q^2$ checked step by step. 234 is an arithmetic root system when $q^2 \in R_3$ by GDD 2 of Row 11. Whole is affine when $\tilde{q}_{34} = q^2$ and $q^2 \in R_3$.

(d) in Case (i)




If 134 is an arithmetic root system, then $\tilde{q}_{34} = q^3$ and $q \notin R_3$. by GDD 3 of Row 7 or Lemma 3.2(I). 234 is an arithmetic root system when $q^8 = 1$ by GDD 3 of Row 10. Whole is affine when $\tilde{q}_{34} = q^3$ and $q^8 = 1$.

(i) in Case (i)




If 134 is an arithmetic root system, then $\tilde{q}_{34} = q^{-1}$ by GDD 3 of Row 15 and Lemma 3.2(I). 234 is an arithmetic root system by GDD 3 of Row 6. Whole is affine when $\tilde{q}_{34} = q^{-1}$.

(k) in Case (i)



If 134 is an arithmetic root system, then $\tilde{q}_{34} = -q$ by GDD 3 of Row 16 and by Lemma 3.2(I). 234 is not an arithmetic root system checked step by step.


(m) in Case (i)



with $q \in R_3$.


If 134 is an arithmetic root system, then $\tilde{q}_{34} = -q^{-1}$ by Lemma 3.2(I)(I)(1), and 134 is an arithmetic root system by Lemma 3.2(I)(1). 234 is arithmetic root system by GDD 3 of Row 6 or by Type 6. Whole is affine when $\tilde{q}_{34} = -q^{-1}$ and $q \in R_3$.

(p) in Case (i)



If 134 is an arithmetic root system, then $\tilde{q}_{34} = -1$ by Lemma 3.2(I). 234 is not an arithmetic root system checked step by step.


(c) in Case (ii)



If 234 is an arithmetic root system, then $\tilde{q}_{34} = q^{-4}$ by Type 6. If 134 is an arithmetic

root system when $q \in R_6$ by Type 6. Whole is affine then $q \in R_6$.


(e) in Case (ii)



with $q \notin R_3$.

If 134 is an arithmetic root system, then $\tilde{q}_{34} = q^3$ by Lemma 3.2(I)(m) and by Lemma 3.2(I) when $q^2 = q^{-6}$. 234 is an arithmetic root system by Type 6. Whole is affine.


(o) in Case (ii)



with $q \in R_3$.


If 134 is an arithmetic root system, then $\tilde{q}_{34} = q^{-1}$ by GDD 3 of Row 15 and by Lemma 3.2(I). Thus 234 is an arithmetic root system by Type 6. Whole is affine when $\tilde{q}_{34} = q^{-1}$ and $q^2 \neq 1$.

(q) in Case (ii)



If 134 is an arithmetic root system, then $\tilde{q}_{34} = -q$ by Lemma 3.2(I)(k). If 234 is an arithmetic root system, then $-q = q^2$, which is a contradiction.


(s) in Case (ii)



with $q \in R_3$.

If 234 is arithmetic root system, then $\tilde{q}_{34} = -q^{-1}$ by Lemma 3.2(I). 234 and 134 are arithmetic root system by Lemma 3.2(I)(l) and by Type 6. Whole is affine when $\tilde{q}_{34} = -q^{-1}$.

(t) in Case (ii)



with $q \in R_3$.

If 134 is arithmetic root system, then $\tilde{q}_{34} = -1$ by Lemma 3.2(I)(g). 234 is not arithmetic root system checked step by step.

4.12 GDD 1 of Row 7

Case (i) Omitting 3 and adding 1.

According to Lemma 3.3 (I) we have to consider following cases.

$$\begin{array}{c} \overset{(a') \text{ or } (1)}{-1} \\ \bullet \text{---} \frac{q}{q} \text{---} \frac{-1}{q^{-1}q} \text{---} \frac{q^{-3}}{q^{-3}} \text{---} \frac{q^3}{q^3} \bullet \end{array} \quad q^2 \neq 1, \text{ GDD 2 of Row 4.}$$

affine by Lemma 3.34.

$$\begin{array}{c} \overset{(a) \text{ or } (2)}{-q} \\ \bullet \text{---} \frac{q^2}{q^2} \text{---} \frac{-1}{q^{-2}q^2} \text{---} \frac{q^{-6}}{q^{-6}} \text{---} \frac{q^6}{q^6} \bullet \end{array} \quad q^4 \neq 1, \text{ GDD 3 of Row 5.}$$

is affine by Lemma 3.4.

$$\begin{array}{c} \overset{(b) \text{ or } (3)}{-1} \\ \bullet \text{---} \frac{q}{q} \text{---} \frac{-1}{q^{-2}q^2} \text{---} \frac{q^{-6}}{q^{-6}} \text{---} \frac{q^6}{q^6} \bullet \end{array} \quad q^2 \notin R_3, q \notin R_2, \text{ GDD 2 of Row 6.}$$

is affine by Lemma 3.7 and by Lemma 3.15.

$$\begin{array}{c} \overset{(d) \text{ or } (4)}{-1} \\ \bullet \text{---} \frac{q}{q} \text{---} \frac{-1}{q^{-3}q^3} \text{---} \frac{q^{-9}}{q^{-9}} \text{---} \frac{q^9}{q^9} \bullet \end{array} \quad q \notin R_2 \cup R_3, q^3 \notin R_2 \cup R_3, \text{ GDD 2 of Row 7.}$$

is affine by Lemma 3.15.

(f) or (5)

$$\begin{array}{ccccccc} -q^{-1} & -1 & q^{-3}q^3 & q^{-9} & q^9 \\ \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

, $q \notin R_2 \cup R_3$, $q^2 \notin R_2 \cup R_3$, GDD 4 of Row 7.

is affine by Lemma 3.4.

$$\begin{array}{ccccccc} q^{-1} \begin{array}{c} (g) \\ \text{or} \\ (6) \end{array} & -1 & q^{-1}q & q^{-3} & q^3 \\ \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

, $q \notin R_2 \cup R_3$, GDD 1 of Row 8.

is an arithmetic root system when $q \in R_4$ by GDD 1 of Row 22. Whole is affine $q \notin R_4$ by Lemma 3.38.

$$\begin{array}{ccccccc} r \begin{array}{c} (h) \\ \text{or} \\ (7) \end{array} & -1 & q^{-1}q & q^{-3} & q^3 \\ \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

, GDD 1 of Row 9.

is affine by Lemma 3.7.

$$\begin{array}{ccccccc} q \begin{array}{c} (i') \\ \text{or} \\ (8) \end{array} & -1 & q^2 & q^{-2}q^{-3} & q^3 \\ \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

, $q \notin R_3 \cup R_2$. GDD 2 of Row 10.

is affine by Lemma 3.7.

$$\begin{array}{ccccccc} q \begin{array}{c} (k) \\ \text{or} \\ (9) \end{array} & -1 & q^{-1}q & q^{-3} & q^3 \\ \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

, $q \notin R_3 \cup R_2$. GDD 1 of Row 10.

is affine by Lemma 3.7.

$$\begin{array}{ccccccc} q^{-2} \begin{array}{c} (l) \\ \text{or} \\ (10) \end{array} & -1 & q^{-1}q & q^{-3} & q^3 \\ \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

, $q \notin R_3 \cup R_2$. GDD 2 of Row 10.

is affine by Lemma 3.7.

$$\begin{array}{ccccccc} q \begin{array}{c} (m) \end{array} & -1 & q^{-1}q & q^{-3} & q^3 \\ \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

, $q \in R_3$. GDD 1 of Row 11.

is empty.

$$\begin{array}{ccccccc} q^{-1} \begin{array}{c} (n) \\ \text{or} \\ (11) \end{array} & -1 & -q & -q^{-1}q^{-3} & q^3 \\ \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

, $q \in R_3$. GDD 3 of Row 14.

is affine by Lemma 3.4.

$$\begin{array}{ccccccc} -1 \begin{array}{c} (p) \\ \text{or} \\ (12) \end{array} & -1 & -q^{-1}q & -q^{-3}q^3 \\ \bullet & \bullet & \bullet & \bullet \end{array}$$

, $q \in R_3$. GDD 2 of Row 16.

is affine by Lemma 3.7.

(r) or (13)

$$\begin{array}{c}
 q \quad -1 \quad -1 \quad -q^{-1} \quad -q^{-3} \quad -q^3 \\
 \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet
 \end{array}
 , q \in R_3. \text{ GDD 4 of Row 16.}$$

is affine by Lemma 3.4.

$$\begin{array}{c}
 (s') \\
 -1 \quad q \quad -1 \quad -1 \quad -1 \quad -1 \\
 \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet
 \end{array}
 , q \in R_3. \text{ GDD 1 of Row 17.}$$

is empty.

Case (ii) Omitting 1 and adding 2. It is checked step by step.

$$\begin{array}{c}
 \text{(a) or (14)} \\
 -1 \quad q^{-1} \quad q \quad q^{-1} \quad -1 \\
 \bullet \text{---} \bullet \begin{array}{l} \nearrow \bullet \\ \searrow \bullet \end{array} \\
 \qquad \qquad \qquad q^{-3} \quad q^3
 \end{array}
 , \text{ (see GDD 1 of Row 7 in Table A2.)}$$

Omitting 3 in whole is an arithmetic root system by Type 7. affine by Lemma 3.4.

Case (iii) Omitting 1 and adding 3.

It is empty checked step by step.

Case (iv) containing circle with four vertexes.

It is empty checked step by step.

4.13 GDD 2 of Row 7

Case (i) Omitting 3 and adding 1.

According to Lemma 3.3 (IV) we have to consider following cases.

$$\begin{array}{c}
 (a') \text{ or } (1). \\
 q \quad q^{-1} \quad -1 \quad q \quad -1 \quad q^{-3} \quad q^3 \\
 \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet
 \end{array}
 , q \in F^* \setminus \{1, -1\}. \text{ GDD 2 of Row 4.}$$

It is affine by Lemma 3.15.

$$\begin{array}{c}
 (a'') \text{ or } (2). \\
 q \quad q^{-2} \quad -1 \quad q^2 \quad -1 \quad q^{-6} \quad q^6 \\
 \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet
 \end{array}
 , q^2 \notin R_3 \cup R_2, q^4 \neq 1, \text{ GDD 2 of Row 5.}$$

It is affine when $q \notin R_4$ by Lemma 3.4.

$$\begin{array}{c}
 \text{(a) or } (3). \\
 q^2 \quad q^{-2} \quad -1 \quad q \quad -1 \quad q^{-3} \quad q^3 \\
 \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet
 \end{array}
 , q \in F^* \setminus \{1, -1\}. \text{ GDD 2 of Row 6.}$$

It is affine by Lemma 3.15.

$$\begin{array}{c}
 \text{(c) or } (4). \\
 q^3 \quad q^{-3} \quad -1 \quad q \quad -1 \quad q^{-3} \quad q^3 \\
 \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet
 \end{array}
 , q \in F^* \setminus \{1, -1\}. q \notin R_3, \text{ GDD 2 of Row 7.}$$

It is affine by Lemma 3.15.

$$\begin{array}{ccccccc} & \text{(e) or (5).} & & & & & \\ -1 & q^{-1} & -1 & q & -1 & q^{-3} & q^3 \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$

$q \notin R_3, q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 8.

It is affine by Lemma 3.15.

$$\begin{array}{ccccccc} & \text{(j) or (6).} & & & & & \\ q & -q & -1 & -q^{-1} & -1 & -q^3 & -q^{-3} \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$

$, q \in R_3$. GDD 2 of Row 14.

It is affine by Lemma 3.4.

$$\begin{array}{ccccccc} & \text{(k).} & & & & & \\ -1 & q & -1 & q & -1 & q^{-3} & q^3 \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$

$, q \in R_3$. GDD 2 of Row 15.

It is empty.

$$\begin{array}{ccccccc} & \text{(l).} & & & & & \\ -q & -q^{-1} & -1 & q & -1 & q^{-3} & q^3 \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$

$, q \in R_3$. GDD 2 of Row 16.

is empty.

$$\begin{array}{ccccccc} & \text{(n).} & & & & & \\ -1 & q & -1 & -1 & -1 & -1 & -1 \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$

$, q \in R_3$. GDD 1 of Row 17.

is empty.

$$\begin{array}{ccccccc} & \text{(n').} & & & & & \\ -1 & -1 & -1 & q & -1 & q^{-3} & q^3 \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$

$, q \in R_3$. GDD 1 of Row 17.

is empty.

$$\begin{array}{ccccccc} & \text{(o).} & & & & & \\ -1 & -q & -1 & q & -1 & q^{-3} & q^3 \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$

$, q \in R_3$. GDD 4 of Row 17.

is empty.

$$\begin{array}{ccccccc} & \text{(p).} & & & & & \\ -1 & q & -1 & -q & -1 & -q^{-3} & -q^3 \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$

$, q \in R_3$. GDD 4 of Row 17.

is empty.

(r).

$$\begin{array}{c}
 q^{-1} \quad -q^{-1} \quad -1 \quad -1 \quad -1 \quad -1 \\
 \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet
 \end{array}
 \quad , q \in R_3. \text{ GDD 9 of Row 17.}$$

is empty.

Case (ii) Omitting 3 and adding 2.

According to Lemma 3.3 (IV) we have to consider following cases.

$$\begin{array}{c}
 q^3 \\
 \bullet \\
 | \\
 q^{-3} \\
 \bullet \\
 | \\
 \bullet \text{---} \bullet \text{---} \bullet \\
 q \quad q^{-1} \quad -1 \quad q \quad -1
 \end{array}
 \quad , q \in F^* \setminus \{1, -1\}. \text{ GDD 2 of Row 4.}$$

(a') or (7).

Omitting 1 in whole is an arithmetic root system by Lemma 3.2 (IV) (IV). Whole is affine when $q \notin R_4$ by Lemma 3.12. Whole is affine when $q \in R_4$ by Lemma 3.25.

$$\begin{array}{c}
 q^6 \\
 \bullet \\
 | \\
 q^{-6} \\
 \bullet \\
 | \\
 \bullet \text{---} \bullet \text{---} \bullet \\
 q \quad q^{-2} \quad -1 \quad q^2 \quad -1
 \end{array}
 \quad q^2 \in R_2 \cup R_3, q^2, q^4 \neq 1, \text{ GDD 2 of Row 5.}$$

(a'') or (8).

Omitting 1 in whole is an arithmetic root system when $q \in R_8$ by Type 2. Omitting 1 in whole is an arithmetic root system when $q \in R_{18}$ by Lemma 3.2(I)(h). Whole is affine when $q \in R_8 \cup R_{18}$ by Lemma 3.4.

$$\begin{array}{c}
 q^3 \\
 \bullet \\
 | \\
 q^{-3} \\
 \bullet \\
 | \\
 \bullet \text{---} \bullet \text{---} \bullet \\
 q^2 \quad q^{-2} \quad -1 \quad q \quad -1
 \end{array}
 \quad , q \in F^* \setminus \{1, -1\}. \text{ GDD 2 of Row 6.}$$

(a) or (9).

Omitting 1 in whole is an arithmetic root system by Lemma 3.2 (IV). Whole is affine by Lemma 3.12.

$$\begin{array}{c}
 q^3 \\
 \bullet \\
 | \\
 q^{-3} \\
 \bullet \\
 | \\
 \bullet \text{---} \bullet \text{---} \bullet \\
 q^3 \quad q^{-3} \quad -1 \quad q \quad -1
 \end{array}
 \quad , q \in F^* \setminus \{1, -1\}. \quad q \notin R_3, \text{ GDD 2 of Row 7.}$$


(c) or (10).

Omitting 1 in whole is an arithmetic root system by Lemma 3.2 (IV). Whole is affine by Lemma 3.25.

$$\begin{array}{c}
 q^3 \\
 \bullet \\
 | \\
 q^{-3} \\
 \bullet \\
 | \\
 \bullet \text{---} \bullet \text{---} \bullet \\
 -1 \quad q^{-1} \quad -1 \quad q \quad -1
 \end{array}
 \quad , q \in F^* \setminus \{1, -1\}. \text{ GDD 2 of Row 8.}$$


(e) or (11).

Omitting 1 in whole is an arithmetic root system when $q \in R_4 \cup R_5 \cup R_6$ by Lemma 3.2 (X). Whole is affine by Lemma 3.16.




, $q \in R_3$. GDD 2 of Row 14.

Omitting 1 in whole is an arithmetic root system by Lemma 3.2(I)(c). Whole is affine by Lemma 3.4.




, $q \in R_3$. GDD 2 of Row 15.




, $q \in R_3$. GDD 2 of Row 16.




, $q \in R_3$. GDD 3 of Row 16.




, $q \in R_3$. GDD 1 of Row 17.



, $q \in R_3$. GDD 1 of Row 17.




, $q \in R_3$. GDD 4 of Row 17.



, $q \in R_3$. GDD 4 of Row 17.

Omitting 1 in whole is an arithmetic root system by Lemma 3.3 (V)(a). Whole is affine by Lemma 3.12.



, $q \in R_3$. GDD 9 of Row 17.

is empty.

Case (iii) Omitting 1 and adding 3.

According to Lemma 3.3 (III) we have to consider following cases.

$$\begin{array}{ccccccc} q^6 & \text{(a).} & & & & & \\ q^{-6} & q^3 & & q^{-3} & -1 & q & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

GDD 1 of Row 6.

$$\begin{array}{ccccccc} q^9 & \text{(c).} & & & & & \\ q^{-9} & q^3 & & q^{-3} & -1 & q & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

, $q \in F^* \setminus \{1, -1\}$, $q^3 \notin R_3$. GDD 1 of Row 7.

$$\begin{array}{ccccccc} -1 & \text{(e).} & & & & & \\ q^{-3} & q^3 & & q^{-3} & -1 & q & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

, $q^3 \in F^* \setminus \{1, -1\}$. GDD 3 of Row 8.

$$\begin{array}{ccccccc} q^3 & \text{(g).} & & & & & \\ -q^3 & -q^{-3} & & -q^3 & -1 & -q^{-1} & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

, $q^3 \in R_3$. GDD 1 of Row 14.

$$\begin{array}{ccccccc} -1 & \text{(h).} & & & & & \\ q^3 & q^3 & & q^{-3} & -1 & q & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

, $q \in R_3$. GDD 1 of Row 15.

is empty.

$$\begin{array}{ccccccc} -q^3 & \text{(j).} & & & & & \\ -q^{-3} & q^3 & & q^{-3} & -1 & q & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

, $q \in R_3$. GDD 1 of Row 16.

is empty.

$$\begin{array}{ccccccc} -1 & \text{(l).} & & & & & \\ -1 & q^3 & & q^{-3} & -1 & q & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

, $q \in R_3$. GDD 2 of Row 17.

is empty.

$$\begin{array}{ccccccc} -1 & \text{(n).} & & & & & \\ q^3 & -q^3 & & -q^{-3} & -1 & -q & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

, $q^3 \in R_3$. GDD 5 of Row 17.

$$\begin{array}{ccccccc} -1 & \text{(o).} & & & & & \\ -q^3 & q^3 & & q^{-3} & -1 & q & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

, $q \in R_3$. GDD 7 of Row 17.

is empty.

(a').

$$\begin{array}{ccccccc} q^3 & q^{-3} & q^3 & q^{-3-1} & q & & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

GDD 1 of Row 4.

$$\begin{array}{ccccccc} q^3 & (a'') & q^6 & q^{-6-1} & q^2 & & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

$q^3 \notin R_2 \cup R_4$, GDD 1 of Row 5.

$$\begin{array}{ccccccc} -1 & (k') & -1 & -1 & -1 & -1 & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

GDD 1 of Row 17.

is empty.

$$\begin{array}{ccccccc} q^{-1} & (p') & -q^{-1-1} & -1 & -1 & -1 & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

GDD 9 of Row 17.

is empty.

Case (iv) containing circle with four vertexes.

According to Lemma 3.39 (ii) we have to consider following cases.

(a') to (a)

$$\begin{array}{ccccccc} -1 & q^{-1} & q & q^{-6}q^3 & & & \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

with $q = q^6$ is not arithmetic root system by Lemma 3.2 (I).

(a') to (c)

$$\begin{array}{ccccccc} -1 & q^{-1} & q & q^{-9}q^3 & & & \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

with $q = q^9$ is not arithmetic root system by Lemma 3.2 (I).

(a') to (e) empty. (a') to (g) empty. (a') to (n) empty. (a') to (a') empty. (a') to (a'')

empty.

(a'') to (a)

$$\begin{array}{ccccccc} -1 & q^{-2} & q & q^{-12}q^6 & & & \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

with $q = q^{12}$ is not an arithmetic root system by Lemma 3.2 (I).

(a'') to (c)

$$\begin{array}{ccccccc} -1 & q^{-2} & q & q^{-18}q^6 & & & \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

with $q = q^{18}$ is not an arithmetic root system by Lemma 3.2 (I).

(a'') to (e) empty. (a'') to (g) empty. (a'') to (n) empty.

(a'') to (a')

$$\begin{array}{ccccccc} -1 & q^{-2} & q & q^{-6}q^6 & & & \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$


with $q = q^6$ is not arithmetic root system by Lemma 3.3 (II) or by Lemma 3.2 (V).

(a'') to (a'') empty.

(a) to (a) or (14)

$$\begin{array}{ccccccc} -1 & q^{-2} & q^2 & q^{-6}q^3 & & & \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$


with $q^2 = q^6$ is an arithmetic root system by Type 2.




is affine.
(a) to (c)

with $q^9 = q^2$ is not an arithmetic root system by Lemma 3.2 (I).

(a) to (e) or (15)




with $-1 = q^2$ is an arithmetic root system by Lemma 3.2 (IV).




is affine.

(a) to (g) empty. (a) to (n) empty. (a) to (a') empty. (a) to (a'') empty. (c) to (a).
empty.

(c) to (c) or (16)




with $q^3 = q^9$ is an arithmetic root system by Lemma 3.2 (IV).




is affine.

(c) to (e) or (17)




with $-1 = q^3$ is an arithmetic root system.




is affine.

(c) to (g) empty. (c) to (n) empty.

(c) to (a') or (18)




is an arithmetic root system by Lemma 3.2 (XI)(e).




is affine.

(c) to (a'') or (19).




is an arithmetic root system by Type 1.




is affine.
(e) to (a)


with $-1 = q^6$ is not an arithmetic root system by Lemma 3.2 (I).
(e) to (c)



with $-1 = q^9$ is not arithmetic root system by Lemma 3.2 (I).
(e) to (e) or (20)




is an arithmetic root system when $1 = q^4$ or $-1 = q^{-3}$ or $q^3 = q^{-2}$ by Lemma 3.2 (X).




is affine. (e) to (g) empty.
(e) to (n)


is not an arithmetic root system by Lemma 3.2 (I).
(e) to (a') empty. (e) to (a'') empty.
(j) to (a)




with $-q^{-1} = q^6$ is not an arithmetic root system by Lemma 3.2 (I).
(j) to (c)



with $-q^{-1} = q^9$ is not an arithmetic root system by Lemma 3.2 (I).
(j) to (e) empty. (j) to (g) empty. (j) to (n) empty.
(j) to (a')



and $-q^{-1} = q^3$ is not arithmetic root system by Lemma 3.3 (II).
(j) to (a'')




with $q^3 = -q^{-2}$ is not an arithmetic root system by Lemma 3.2 (V).

4.14 GDD 3 of Row 7


Case (i) Omitting 3 and adding 1.

According to Lemma 3.3 (III) we have to consider following cases.

(a).  , $q \in F^* \setminus \{1, -1\}$. GDD 1 of Row 6.


Omitting 2 in whole is not an arithmetic root system by Lemma 3.2 (V).

(b) or (1).




$q^{-2}, q \in F^* \setminus \{1, -1\}$. GDD 3 of Row 6.

Omitting 2 in whole is an arithmetic root system by Lemma 3.3 (III)(n) with $q \in R_6$.
 Omitting 2 in whole is not an arithmetic root system by Lemma 3.3 (III) with $q \notin R_6$.
 Omitting 1 in whole is an arithmetic root system by Lemma 3.2 (XI)(e) with $q \in R_6$.
 Whole is affine with $q \in R_6$.

(c).  , $q \in F^* \setminus \{1, -1\}$, $q \notin R_3$. GDD 1 of Row 7.


Omitting 2 in whole is not an arithmetic root system by Lemma 3.2 (V).

(d) or (2).




$q^{-2}, q \in F^* \setminus \{1, -1\}$, $q \notin R_3$. GDD 3 of Row 7.

Omitting 2 in whole is an arithmetic root system by Lemma 3.2 (V)(b) when $q \in R_6$.
 Omitting 1 in whole is an arithmetic root system by Type 7. Whole is affine by Lemma 3.4.


(e) or (3).  , $q \in F^* \setminus \{1, -1\}$. GDD 3 of Row 8.

Omitting 2 in whole is arithmetic root system by Lemma 3.3 (III)(n). Whole is affine by Lemma 3.4.



, $q \in R_3$. GDD 1 of Row 14.


Omitting 2 in whole is not an arithmetic root system by Lemma 3.2(I).



, $q \in R_3$. GDD 1 of Row 15.


is empty.

(i).




, $q \in R_3$. GDD 3 of Row 15.

is empty.



, $q \in R_3$. GDD 1 of Row 16.


is empty.



, $q \in R_3$. GDD 2 of Row 17.

is empty.

(m).



, $q \in R_3$. GDD 3 of Row 17.

is empty.


(n) or (4).  , $q \in R_3$. GDD 5 of Row 17.

Omitting 2 in whole is an arithmetic root system by Lemma 3.2 (V)(b). Whole is affine by Lemma 3.4.

(o).  , $q \in R_3$. GDD 7 of Row 17.

is empty.

(p).

 , $q \in R_3$. GDD 8 of Row 17.

is empty.

(a') or (5).  GDD 1 of Row 4.

Omitting 2 in whole is arithmetic root system when $q \in R_6$ by Lemma 3.3(II)(f). Whole is affine when $q \in R_6$ by Lemma 3.4.

(a'')  GDD 1 of Row 5.

Omitting 2 in whole is not an arithmetic root system by Lemma 3.2(I).

(k')  GDD 1 of Row 17.

is empty.

(p')  GDD 9 of Row 17.

is empty.

Case (ii) Omitting 3 and adding 2.

According to Lemma 3.3 (I) we have to consider following cases.

(a') or (6)  GDD 2 of Row 4.

Omitting 1 in whole is arithmetic root system when $q^3 = -1$ by Lemma 3.2 (XI)(e). Omitting 1 in whole is an arithmetic root system when $q \in R_4$ by Type 7. Whole is affine by Lemma 3.4.


(a) or (7)  GDD 3 of Row 5.

Omitting 1 in whole is arithmetic root system when $q \in R_8$ by Type 3. Omitting 1 in whole is not arithmetic root system when $q \notin R_8$ by Lemma 3.2(I). Whole is affine by Lemma 3.4 when $q \in R_8$.


(b) or (8)  GDD 2 of Row 6.

Omitting 1 in whole is an arithmetic root system when $q \in R_7$ by Lemma 3.2 (XI). Whole is affine when $q \in R_7$ by Lemma 3.4.

(c) or (9)


 GDD 3 of Row 7.

Omitting 1 in whole is an arithmetic root system when $q \in R_4 \cup R_6 \cup R_5$ by Lemma 3.2 (XI) and by Type 7. Omitting 2 in whole is an arithmetic root system when $q \in R_3 \cup R_6$ by Lemma 3.3 (III) (h) (n). Whole is affine when $q \in R_6$.

(d) or (10)  GDD 2 of Row 7.

Omitting 1 in whole is an arithmetic root system $q \in R_{10}$ by Lemma 3.2 (XI). Whole is affine by Lemma 3.4.

(e)




GDD 3 of Row 7.


Omitting 1 in whole is an arithmetic root system when $q \in R_9$ by Lemma 3.2 (XI)(b). Omitting 2 in whole is not an arithmetic root system by Lemma 3.2 (V).



Omitting 1 in whole is an arithmetic root system when $q \in R_{11}$ by Type 3. Omitting 1 in whole is not an arithmetic root system when $q \notin R_{11}$ by Lemma 3.2(I). Whole is affine by Lemma 3.4.



Omitting 1 in whole is an arithmetic root system $q \in R_4 \cup R_7 \cup R_{10} \cup R_6$. by Lemma 3.3 (IV) and by Lemma 3.2 (IV). Whole is affine by Lemma 3.4.



Omitting 1 in whole is an arithmetic root system when $r = q^3$, $q \notin R_6$ or $r = q^6$, $q \notin R_{12}$ or $r = -q^{-3}$, $q \in R_9$ or $q^3 = -1$, or $r = -1$ by Lemma 3.2 (IV) (X) (XI). Whole is affine by Lemma 3.4.




Omitting 1 in whole is an arithmetic root system when $q \in R_7 \cup R_{13} \cup R_{19} \cup R_{12}$ by Lemma 3.2 (IV) (X). It is affine by Lemma 3.4.



Omitting 1 in whole is arithmetic root system when $q \in R_5 \cup R_8 \cup R_6$ by Lemma 3.3 (IV) and by Lemma 3.2 (IV). Whole is affine by Lemma 3.4.




Omitting 1 in whole is an arithmetic root system when $q \in R_4 \cup R_5 \cup R_8 \cup R_{11} \cup R_6$ by Lemma 3.3 (IV) and by Lemma 3.2 (IV). Whole is affine by Lemma 3.4.



, $q \in R_3$. GDD 1 of Row 11.


is empty.



, $q \in R_3$. GDD 3 of Row 14.


Omitting 1 in whole is an arithmetic root system by Lemma 3.2(I)(C). Whole is affine by Lemma 3.4.

(o)



GDD 3 of Row 15.


is empty.




, $q \in R_3$. GDD 2 of Row 16.

Omitting 1 in whole is an arithmetic root system by Lemma 3.3V)(a). Whole is affine by Lemma 3.4.

(q)




is empty.



, $q \in R_3$. GDD 4 of Row 16.


is empty.



, $q \in R_3$. GDD 1 of Row 17.


is empty.

(s)



is empty.


(t)




is empty.

Case (iii) Omitting 1 and adding 3.


According to Lemma 3.3 (IV) we have to consider following cases.



Omitting 2 in whole is arithmetic root system when $q \in R_5$ by Lemma 3.2 (I) or Type 2. Omitting 2 in whole is arithmetic root system when $q \in R_4$ by Lemma 3.2(I)(o). Whole is affine by Lemma 3.4.




Omitting 2 in whole is not an arithmetic root system by Lemma 3.2(I).




Omitting 2 in whole is an arithmetic root system when $q^8 = 1$ by Lemma 3.2(I). Omitting 2 in whole is an arithmetic root system when $q \in R_{18}$ by Lemma 3.2(I)(h). Whole is affine by Lemma 3.4.

(b) or (21).




Omitting 2 in whole is an arithmetic root system when $q \in R_5$ by Lemma 3.2(I) or

Type 2. Omitting 2 in whole is an arithmetic root system when $q \in R_{12}$ by Lemma 3.2(I)(h). Omitting 3 in whole is an arithmetic root system by Lemma 3.2 (IV). Whole is affine.

(c) or (22).  , $q^3 \in F^* \setminus \{1, -1\}$. $q \notin R_3$, GDD 2 of Row 7.

Omitting 2 in whole is an arithmetic root system when $q^{11} = 1$ or $q^{12} = -1$ by Type 2 and by Lemma 3.2(I)(h). Whole is affine by Lemma 3.4.

(d) or (23).


 , $q^3 \in F^* \setminus \{1, -1\}$. $q \notin R_3$, GDD 3 of Row 7.

Omitting 2 in whole is an arithmetic root system by Lemma 3.2(I)(h). Omitting 1 in whole is the same as Omitting 1 in whole. Whole is affine.

(e) or (24).  , $q^3 \in F^* \setminus \{1, -1\}$. GDD 2 of Row 8.

Omitting 2 in whole is an arithmetic root system when $q \in R_5$ by Type 2. Whole is affine when $q \in R_5$ by Lemma 3.4.

(f) or (25).




, $q^3, s^3, r^3 \in F^* \setminus \{1\}$, $q^3 s^3 r^3 = 1$, $q^3 \neq s^3$, $q^3 \neq r^3$, $r^3 \neq s^3$. GDD 3 of Row 9.

Omitting 3 in whole is arithmetic root system when $q = r^3$, or $r^6 = q$, or $q = r^9$, or $q = -r^3$, or $q^3 \in R_3$, or $r^3 = -1$.. Omitting 2 in whole is arithmetic root system when

$s^3 = q^2$, $q \notin R_5$ by Type 2. Whole is affine.


(g).



, $q^3 \in F^* \setminus \{1, -1\}$. $q^3 \notin R_2 \cup R_3$, GDD 3 of Row 10.

Omitting 2 in whole is not an arithmetic root system by Lemma 3.2(I).


(h).



, $q^2, q^3 \neq 1$. GDD 3 of Row 10.

Omitting 2 in whole is not an arithmetic root system by Lemma 3.2(I).


(i).



, $q \in R_3$. GDD 2 of Row 11.

is empty.


(j). $q^{-2} q$ q^{-1}



, $q \in R_3$. GDD 2 of Row 14.


Omitting 2 in whole is not an arithmetic root system by Lemma 3.2 (XIV) or by Lemma 3.2(I)

(k). $q^{-2} q$ q^{-1}



, $q \in R_3$. GDD 2 of Row 15.


is empty.



, $q \in R_3$. GDD 2 of Row 16.


is empty.

(m).




, $q \in R_3$. GDD 3 of Row 16.

Omitting 2 in whole is an arithmetic root system by Lemma 3.2 (X)(b). Omitting 3 in whole is not an arithmetic root system by Lemma 3.2(I).




, $q \in R_3$. GDD 1 of Row 17.

is empty.




, $q \in R_3$. GDD 1 of Row 17.

is empty.



, $q \in R_3$. GDD 4 of Row 17.


is empty.



, $q \in R_3$. GDD 4 of Row 17.

Omitting 2 in whole is not an arithmetic root system by Lemma 3.2 (XI).

(q).




, $q \in R_3$. GDD 8 of Row 17.

is empty.

Case (v) Complete diagram containing circle with four vertexes.

According to Lemma 3.40 we have to consider following cases.


(b) in Case (i)



with $q \notin R_3$.


134 is not an arithmetic root system by Lemma 3.2(I),

(d) in Case (i)




134 is not an arithmetic root system by Lemma 3.2(I).

(c) in Case (ii)




with $q \notin R_3$. 134 is not an arithmetic root system by Lemma 3.2(I).

(e) in Case (ii)




134 is not arithmetic root system by Lemma 3.2(I).

(b) in Case (iii)




134 is not arithmetic root system by Lemma 3.2(I).

(d) in Case (iii) or (26)




If 134 is arithmetic root system, then $q^{-2} = q^3$ and $\tilde{q}_{14} = q^{-2}$ by Lemma 3.2(I).
 214 is arithmetic root system by Lemma 3.2(I)(m), Whole is affine when $\tilde{q}_{14} = q^{-2}$ and $q \in R_5$.

(f) in Case (iii)




124 is not arithmetic root system by Lemma 3.2(I).

(g) in Case (iii)




134 is not arithmetic root system by Lemma 3.2(I).

(h) in Case (iii)



with $q^6 \neq 1$. 134 is not arithmetic root system by Lemma 3.2(I).


(m) in Case (iii)




If 124 is arithmetic root system, then $q^{-2} = -q$ by Lemma 3.2(I), which is a contradiction.

4.15 GDD 4 of Row 7


Case (i) Omitting 3 and adding 1. According to Lemma 3.3 (III) we have to consider following cases.




$q \notin R_3 \cup R_2, q^3 \in F^* \setminus \{1, -1\}$. GDD 1 of Row 6.



$q^3 \notin R_3 \cup R_2, q \notin R_3 \cup R_2$, GDD 1 of Row 7.



$q \notin R_3 \cup R_2, q^3 \in F^* \setminus \{1, -1\}$. GDD 3 of Row 8.



$q \notin R_3 \cup R_2, -q^3 \in R_3$. GDD 1 of Row 14.

$$\begin{array}{ccccccc} & \text{(h) or }_3(5). & & & & & \\ -1 & q^3 & q^3 & q^{-3}-1 & q^2 & & -q^{-1} \\ \bullet & \bullet & \bullet & \bullet & \bullet & & \bullet \end{array}$$

is affine by Lemma 3.4.

$q \notin R_3 \cup R_2, q^3 \in R_3$. GDD 1 of Row 15.

$$\begin{array}{ccccccc} & \text{(j) or }_3(6). & & & & & \\ -q^3 & -q^{-3} & q^3 & q^{-3}-1 & q^2 & & -q^{-1} \\ \bullet & \bullet & \bullet & \bullet & \bullet & & \bullet \end{array}$$

is affine by Lemma 3.4.

$q \notin R_3 \cup R_2, q^3 \in R_3$. GDD 1 of Row 16.

$$\begin{array}{ccccccc} & \text{(l) or }_3(7). & & & & & \\ -1 & -1 & q^3 & q^{-3}-1 & q^2 & & -q^{-1} \\ \bullet & \bullet & \bullet & \bullet & \bullet & & \bullet \end{array}$$

is affine by Lemma 3.4.

$q \notin R_3 \cup R_2, q^3 \in R_3$. GDD 2 of Row 17.

$$\begin{array}{ccccccc} & \text{(n) or }_3(8). & & & & & \\ -1 & q^3 & -q^{-3} & -1 & q^2 & & q^{-1} \\ \bullet & \bullet & \bullet & \bullet & \bullet & & \bullet \end{array}$$

is affine by Lemma 3.4.

$q \notin R_3 \cup R_2, q^3 \in R_3$. GDD 5 of Row 17.

$$\begin{array}{ccccccc} & \text{(o) or }_3(9). & & & & & \\ -1 & -q^3 & q^3 & q^{-3}-1 & q^2 & & -q^{-1} \\ \bullet & \bullet & \bullet & \bullet & \bullet & & \bullet \end{array}$$

is affine by Lemma 3.4.

$q \notin R_3 \cup R_2, q^3 \in R_3$. GDD 7 of Row 17.

$$\begin{array}{ccccccc} & \text{(a') or }_3(10). & & & & & \\ q^3 & q^{-3} & q^3 & q^{-3}-1 & q^2 & & -q^{-1} \\ \bullet & \bullet & \bullet & \bullet & \bullet & & \bullet \end{array}$$

is affine by Lemma 3.4.

$q \notin R_3 \cup R_2, q^3 \notin R_2$, GDD 1 of Row 4.

$$\begin{array}{ccccccc} & \text{(a'') or }_3(11). & & & & & \\ q^3 & q^{-6} & q^6 & q^{-6}-1 & q^4 & & -q^{-2} \\ \bullet & \bullet & \bullet & \bullet & \bullet & & \bullet \end{array}$$

is affine by Lemma 3.4.

$q^2 \notin R_3 \cup R_2, q^3 \notin R_2 \cup R_4$, GDD 1 of Row 5.

$$\begin{array}{ccccccc} & \text{(k') or }_3(12). & & & & & \\ -1 & q & -1 & -1 & -1 & & \\ \bullet & \bullet & \bullet & \bullet & \bullet & & \bullet \end{array}$$

is empty.

$$\begin{array}{ccccccc} & \text{(p')} & & & & & \\ q^{-1} & -q^{-1} & -1 & -1 & -1 & & \\ \bullet & \bullet & \bullet & \bullet & \bullet & & \bullet \end{array}$$


is empty.

Case (ii) Omitting 1 and adding 2. According to Type 3 we have to consider following cases.


(a) or (13)

$$\begin{array}{ccccccc} & & & & q^3 & & \\ & & & & / & & \\ q^3 & q^{-3} & -1 & q^{-3} & q^3 & & \\ & \bullet & \bullet & \bullet & \bullet & & \\ & & & & \backslash & & \\ & & & & q^2 & & \\ & & & & \bullet & & \\ & & & & \backslash & & \\ & & & & -q^{-1} & & \\ & & & & \bullet & & \end{array}, (q \notin R_3 \cup R_2, \text{ see GDD 4 of Row 7 in Table A2.})$$

Omitting 2 in whole is an arithmetic root system by Lemma 3.2 (IV). affine by Lemma 3.4.


(b) or (14)  ($q \notin R_3 \cup R_2$, see Type 3.)


Omitting 2 in whole is an arithmetic root system when $q \in R_5 \cup R_7 \cup R_9 \cup R_6 \cup R_4$ by Lemma 3.2 (IV) (X) (XI)(e). Whole is affine by Lemma 3.4.


(c) or (15)  ($q \notin R_3 \cup R_2$, see Type 3.)

Omitting 2 in whole is an arithmetic root system by Lemma 3.2 (IV). Whole is affine by Lemma 3.4.

Case (iii) Omitting 1 and adding 3. According to inversion of Type 3 we have to consider following cases.

(a)  ($q \notin R_3 \cup R_2$, $q \in R_6$, see GDD 2 of Row 13 in Table A2).

(b)  ($q \in R_3$, see GDD 4 of Row 15 in Table A2).
is empty.

(c)  ($q \in R_6$, see GDD 5 of Row 17 in Table A2).
is empty.

Case (iv) Containing circle with four vertexes.

According to Lemma 3.39 (ii) we have to consider following cases.

(a) to (a) empty. (c) to (a) empty. (e) to (a) empty. (g) to (a) empty. (h) to (a) empty. (j) to (a) empty. (l) to (a) empty. (n) to (a) empty. (o) to (a) empty. (a') to (a) empty. (a'') to (a) empty.

4.16 GDD 1 of Row 8

Case (i) Omitting 3 and adding 1.

According to Lemma 3.3 (III) we have to consider following cases.

(a).  , $q \in F^* \setminus \{1, -1\}$. GDD 1 of Row 6.

is an arithmetic root system by Type 1.

(c) or (1).

$$\begin{array}{c} q^3 \quad q^{-3} \quad q \quad q^{-1}-1 \quad q \quad q^{-1} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad , q \in F^* \setminus \{1, -1\} , q \notin R_3. \text{ GDD 1 of Row 7.}$$

is an arithmetic root system when $q \in R_4$ by GDD 1 of Row 22; Otherwise it is affine by Lemma 3.38.

$$\begin{array}{c} \text{(e).} \\ -1 \quad q^{-1} \quad q \quad q^{-1}-1 \quad q \quad q^{-1} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad , q \in F^* \setminus \{1, -1\}. \text{ GDD 3 of Row 8.}$$

is an arithmetic root system by Type 7.

$$\begin{array}{c} \text{(g).} \\ q \quad -q \quad -q^{-1} \quad -q \quad -1 \quad -q^{-1}-q \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad , q \in R_3. \text{ GDD 1 of Row 14.}$$

is an arithmetic root system by Type 4.

$$\begin{array}{c} \text{(h).} \\ -1 \quad q \quad q \quad q^{-1}-1 \quad q \quad q^{-1} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad , q \in R_3. \text{ GDD 1 of Row 15.}$$

is an arithmetic root system by GDD 1 of Row 20.

$$\begin{array}{c} \text{(j) or (2).} \\ -q \quad -q^{-1}q \quad q^{-1}-1 \quad q \quad q^{-1} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad , q \in R_3. \text{ GDD 1 of Row 16.}$$

is affine by Lemma 3.38.

$$\begin{array}{c} \text{(l) or (3).} \\ -1 \quad -1 \quad q \quad q^{-1}-1 \quad q \quad q^{-1} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad , q \in R_3. \text{ GDD 2 of Row 17.}$$

is affine by Lemma 3.26.

$$\begin{array}{c} \text{(n) or (4).} \\ -1 \quad q \quad -q \quad -q^{-1}-1 \quad -q \quad -q^{-1} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad , q \in R_3. \text{ GDD 5 of Row 17.}$$

is affine by Lemma 3.7.

$$\begin{array}{c} \text{(o) or (5).} \\ -1 \quad -q \quad q \quad q^{-1}-1 \quad q \quad q^{-1} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad , q \in R_3. \text{ GDD 7 of Row 17.}$$

is affine by Lemma 3.38.

(a').

$$\begin{array}{c}
 q \quad q^{-1} \quad q \quad q^{-1}-1 \quad q \quad q^{-1} \\
 \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet
 \end{array}
 \quad \text{GDD 1 of Row 4.}$$

is an arithmetic root system by Type 7.

$$\begin{array}{c}
 (a'') \\
 q \quad q^{-2} \quad q^2 \quad q^{-2}-1 \quad q^2 \quad q^{-2} \\
 \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet
 \end{array}
 \quad \text{GDD 1 of Row 5.}$$

is an arithmetic root system by Type 2.

$$\begin{array}{c}
 (k') \\
 -1 \quad q \quad -1 \quad -1 \quad -1 \\
 \bullet \text{---} \bullet \text{---} \bullet
 \end{array}
 \quad \text{GDD 1 of Row 17.}$$

is empty.

$$\begin{array}{c}
 (p') \\
 q^{-1} \quad -q^{-1}-1 \quad -1 \quad -1 \\
 \bullet \text{---} \bullet \text{---} \bullet
 \end{array}
 \quad \text{GDD 9 of Row 17.}$$

is empty.

Case (ii) Omitting 3 and adding 2.

According to Lemma 3.3 (I) we have to consider following cases.

$$\begin{array}{c}
 (a') \text{ or } (6) \quad \begin{array}{c}
 q^{-1} \\
 \bullet \\
 q \\
 \bullet \\
 -1 \quad q \quad -1 \quad q^{-1} \quad q \\
 \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet
 \end{array}
 \quad \text{GDD 2 of Row 4.}
 \end{array}$$

Omitting 1 in whole is arithmetic root system when $q \in R_3 \cup R_4$ by Lemma 3.2 (X).

Whole is affine by Lemma 3.25.

$$\begin{array}{c}
 (a) \quad \begin{array}{c}
 q^{-2} \\
 \bullet \\
 q^2 \\
 \bullet \\
 -q^{-2} \quad -1 \quad q^{-2} \quad q^2 \\
 \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet
 \end{array}
 \quad q \notin R_2 \cup R_4, \text{ GDD 3 of Row 5.}
 \end{array}$$

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(I).

$$\begin{array}{c}
 (b) \text{ or } (7) \quad \begin{array}{c}
 q^{-2} \\
 \bullet \\
 q^2 \\
 \bullet \\
 -1 \quad q \quad -1 \quad q^{-2} \quad q^2 \\
 \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet
 \end{array}
 \quad \text{GDD 2 of Row 6.}
 \end{array}$$

Omitting 1 in whole is an arithmetic root system when $q \in R_3 \cup R_4 \cup R_5$ by Lemma 3.2 (X). Whole is affine when $q \notin R_3$ by Lemma 3.12 or by Lemma 3.25. Whole is an arithmetic root system when $q \in R_3$ by Lemma 3.25(b).

(d) or (8)  GDD 2 of Row 7.


Omitting 1 in whole is an arithmetic root system when $q \in R_4 \cup R_6 \cup R_5$ by Lemma 3.3 (X). Whole is affine by Lemma 3.25.

(f) or (9)  , $q \notin R_2 \cup R_3$, GDD 4 of Row 7.


Omitting 1 in whole is an arithmetic root system when $q \in R_5 \cup R_6$ by Type 3 and by Lemma 3.2(I)(h). Omitting 1 in whole is not an arithmetic root system when $q \notin R_5 \cup R_6$ by Lemma 3.2(I). Whole is affine by Lemma 3.4.

(g)  , GDD 1 of Row 8.

Omitting 1 in whole is an arithmetic root system by Lemma 3.2 (IV). Whole is an arithmetic root system by GDD 6 Row 12 or Type 5.

(h) or (10)  , GDD 1 of Row 9.


Omitting 1 in whole is an arithmetic root system by Lemma 3.2 (IV). Whole is affine by Lemma 3.8.

(i') or (11)  , $q \notin R_3$. GDD 2 of Row 10.


Omitting 1 in whole is an arithmetic root system by Lemma 3.2 (IV). Whole is affine by Lemma 3.8.

(k)  , $q \notin R_3$. GDD 1 of Row 10.

Whole is an arithmetic root system by Lemma 3.2 (IV). Whole is an arithmetic root system by GDD 6 of Row 12 or Type 5.

(l) or (12)  , $q \notin R_3$. GDD 2 of Row 10.

Omitting 1 in whole is an arithmetic root system by Lemma 3.2 (IV). Whole is affine when $q^3 \neq 1$ by Lemma 3.8; Whole is an arithmetic root system when $q^3 = 1$ by Type 5.

(m)  , $q \in R_3$. GDD 1 of Row 11.


Whole is an arithmetic root system by Type 5.

(n)  , $q \in R_3$. GDD 3 of Row 14.

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(I).

(p)  , $q \in R_3$. GDD 2 of Row 16.

Omitting 1 in whole is not an arithmetic root system by Lemma 3.3 (X).

(r)  , $q \in R_3$. GDD 4 of Row 16.

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(I),


(s')  , $q \in R_3$. GDD 1 of Row 17.
is empty.

Case (iii) Omitting 1 and adding 3.


It is the same as Case (i) .

Case (iv) containing circle with four vertexes.


According to Lemma 3.39 (ii) we have to consider following cases.

(a) to (a) 

with $q^{-2} = q^2$ is not an arithmetic root system by Lemma 3.2 (I).

(a) to (c) 


with $q^{-3} = q^2$ is not an arithmetic root system by Lemma 3.2 (I).

(a) to (e) 

with $-1 = q^2$ is not an arithmetic root system by Lemma 3.2 (I).

(a) to (g) empty. (a) to (h) empty. (a) to (j) empty (a) to (l) empty. (a) to (n) empty.
 (a) to (o) empty.


(a) to (a') or (13)



with $q^{-1} = q^2$ is an arithmetic root system by Type 2.


is affine.

(a) to (a'')




with $q^{-1} = q^4$ is not an arithmetic root system by Lemma 3.2 (I).

(c) to (c)




with $q^{-3} = q^3$ is not an arithmetic root system by Lemma 3.2 (I).

(c) to (e)



with $-1 = q^3$ is not an arithmetic root system by Lemma 3.2 (I).


(c) to (g) empty. (c) to (h) empty. (c) to (j) empty. (c) to (l) empty.
 (c) to (n)



with $-1 = q^3$ is not an arithmetic root system by Lemma 3.2 (I).


(c) to (o) empty.

(c) to (a').




with $q^{-1} = q^3$ is not an arithmetic root system by Lemma 3.2 (I).

(c) to (a'')




with $q^6 = q^{-1}$ is not an arithmetic root system by Lemma 3.2 (I).

(e) to (e) or (14)




is an arithmetic root system.



is affine.

(e) to (g) empty.


(e) to (h)



is not an arithmetic root system by Lemma 3.2 (I).


(e) to (j) empty

(e) to (l)




is not an arithmetic root system by Lemma 3.2 (I).

(e) to (n)



is not an arithmetic root system by Lemma 3.2 (I).


(e) to (o)



is not an arithmetic root system by Lemma 3.2 (I).

(e) to (a') empty. (e) to (a'') empty. (g) to (g) empty. (g) to (h) empty. (g) to (j) empty. (g) to (l) empty. (g) to (n) empty. (g) to (o) empty. (g) to (a') empty. (g) to (a'') empty.


(h) to (h)



is not an arithmetic root system by Lemma 3.2 (I).


(h) to (j) empty.

(h) to (l)




is not an arithmetic root system by Lemma 3.2(I).

(h) to (n)



is not an arithmetic root system by Lemma 3.2 (I).


(h) to (o)



is not an arithmetic root system by Lemma 3.2 (I).

(h) to (a') empty. (h) to (a'') empty. (j) to (j) empty. (j) to (l) empty. (j) to (n) empty. (j) to (o) empty. (j) to (a') empty. (j) to (a'') empty.

(l) to (l)



is not an arithmetic root system by Lemma 3.2 (I).



(l) to (n) empty.

(l) to (o)

is not an arithmetic root system by Lemma 3.2 (I).


(l) to (a') empty. (l) to (a'') empty.

(n) to (n)



is not an arithmetic root system by Lemma 3.2 (I).


(n) to (o)



is not an arithmetic root system by Lemma 3.2 (I).

(n) to (a') empty. (n) to (a'') empty.


(o) to (o)




is not an arithmetic root system by Lemma 3.2 (I).

(o) to (a') empty. (o) to (a'') empty. (a') to (a') empty.

(a') to (a'') or (15)



and $q^2 = q^{-1}$ is an arithmetic root system by Lemma 3.3 (II)(c) or by Type 1.




is affine.

(a'') to (a'') empty.

4.17 GDD 2 of Row 8


Case (i) Omitting 3 and adding 1.

According to Lemma 3.3 (IV) we have to consider following cases.



, $q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 4.

is an arithmetic root system by Type 7.



, $q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 5.

is an arithmetic root system by Type 2.

(a).

$$\begin{array}{c} q^2 \quad -1 \quad -1 \\ q^{-2} \quad q \quad q^{-1} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad , q \in F^* \setminus \{1, -1\}. \text{ GDD 2 of Row 6.}$$

is an arithmetic root system by Type 1.

$$\begin{array}{c} (c) \text{ or } (1). \\ q^3 \quad -1 \quad -1 \\ q^{-3} \quad q \quad q^{-1} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad , q \in F^* \setminus \{1, -1\}. \quad q \notin R_3, \text{ GDD 2 of Row 7.}$$

is an arithmetic root system when $q \in R_4$ by GDD 2 of Row 22; Whole is affine when $q \notin R_4$ by Lemma 3.15.

$$\begin{array}{c} (e). \\ -1 \quad -1 \quad -1 \\ q^{-1} \quad q \quad q^{-1} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad , q \in F^* \setminus \{1, -1\}. \text{ GDD 2 of Row 8.}$$

is an arithmetic root system by Type 7.

$$\begin{array}{c} (j). \\ q \quad -1 \quad -1 \\ -q \quad -q^{-1} \quad -q \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad , q \in R_3. \text{ GDD 2 of Row 14.}$$

is an arithmetic root system by Type 4.

$$\begin{array}{c} (k). \\ -1 \quad -1 \quad -1 \\ q \quad q \quad q^{-1} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad , q \in R_3. \text{ GDD 2 of Row 15.}$$

is an arithmetic root system by GDD 3 of Row 20 or by Lemma 3.9(h).

$$\begin{array}{c} (l) \text{ or } (2). \\ -q \quad -1 \quad -1 \\ -q^{-1} \quad q \quad q^{-1} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad , q \in R_3. \text{ GDD 2 of Row 16.}$$

is affine by Lemma 3.27.

$$\begin{array}{c} (n). \\ -1 \quad -1 \quad -1 \quad -1 \\ q \quad -1 \quad -1 \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad , q \in R_3. \text{ GDD 1 of Row 17.}$$

is empty.

$$\begin{array}{c} (n') \text{ or } (3). \\ -1 \quad -1 \quad -1 \\ -1 \quad q \quad q^{-1} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad , q \in R_3. \text{ GDD 1 of Row 17.}$$

is affine by Lemma 3.9.

$$\begin{array}{c} (o) \text{ or } (4). \\ -1 \quad -1 \quad -1 \\ -q \quad q \quad q^{-1} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad , q \in R_3. \text{ GDD 4 of Row 17.}$$

is affine by Lemma 3.9.

(p) or (5).

$$\begin{array}{c} -1 \quad q \quad -1 \quad -q \quad -1 \quad -q^{-1} \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad , q \in R_3. \text{ GDD 4 of Row 17.}$$

is affine by Lemma 3.9.

$$\begin{array}{c} q^{-1} \quad (r) \\ -q^{-1} \quad -1 \quad -1 \quad -1 \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad , q \in R_3. \text{ GDD 9 of Row 17.}$$

is empty.

Case (ii) Omitting 3 and adding 2.

According to Lemma 3.3 (IV) we have to consider following cases

$$(a') \text{ or } (6). \quad \begin{array}{c} -1 \\ \bullet \\ q^{-1} \\ \bullet \text{---} \bullet \text{---} \bullet \\ q \quad q^{-1} \quad -1 \quad q \quad -1 \end{array} \quad , q \in F^* \setminus \{1, -1\}. \text{ GDD 2 of Row 4.}$$

Omitting 1 in whole is an arithmetic root system when $q \in R_3 \cup R_4$ by Lemma 3.2 (X). Whole is affine when $q \in R_4$ by Lemma 3.16. Whole is an arithmetic root system when $q \in R_3$ by Lemma 3.16(d).

$$(a''). \quad \begin{array}{c} -1 \\ \bullet \\ q^{-2} \\ \bullet \text{---} \bullet \text{---} \bullet \\ q \quad q^{-2} \quad -1 \quad q^2 \quad -1 \end{array} \quad q \notin R_4, q \in F^* \setminus \{1, -1\}. \text{ GDD 2 of Row 5.}$$

Omitting 1 in whole is not an arithmetic root system when $q \notin R_4$ by Lemma 3.2(II).

$$(a) \text{ or } (7). \quad \begin{array}{c} -1 \\ \bullet \\ q^{-1} \\ \bullet \text{---} \bullet \text{---} \bullet \\ q^2 \quad q^{-2} \quad -1 \quad q \quad -1 \end{array} \quad , q \in F^* \setminus \{1, -1\}. \text{ GDD 2 of Row 6.}$$

Omitting 1 in whole is an arithmetic root system when $q \in R_4 \cup R_5 \cup R_3$ by Lemma 3.2 (X). Whole is an arithmetic root system when $q \in R_3$ by Lemma 3.16(d). Whole is affine when $q \in R_4 \cup R_5$ by Lemma 3.16.

$$(c) \text{ or } (8). \quad \begin{array}{c} -1 \\ \bullet \\ q^{-1} \\ \bullet \text{---} \bullet \text{---} \bullet \\ q^3 \quad q^{-3} \quad -1 \quad q \quad -1 \end{array} \quad , q \in F^* \setminus \{1, -1\}. q \notin R_3, \text{ GDD 2 of Row 7.}$$

Omitting 1 in whole is an arithmetic root system when $q \in R_4 \cup R_5 \cup R_6$ by Lemma 3.2 (X). Whole is affine when $q \in R_4 \cup R_5 \cup R_6$ by Lemma 3.16.

$$(e) \text{ or } (9). \quad \begin{array}{c} -1 \\ \bullet \\ q^{-1} \\ \bullet \text{---} \bullet \text{---} \bullet \\ -1 \quad q^{-1} \quad -1 \quad q \quad -1 \end{array} \quad , q \in F^* \setminus \{1, -1\}. \text{ GDD 2 of Row 8.}$$

Omitting 1 in whole is an arithmetic root system when $q \in R_3$ by GDD 2 of Row 15 or by Lemma 3.2 (XI)(b). Whole is affine by Lemma 3.16.

(j).  , $q \in R_3$. GDD 2 of Row 14.

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(I).

(k) or (10).  , $q \in R_3$. GDD 2 of Row 15.

Omitting 1 in whole is an arithmetic root system by Type 7. Whole is affine by Lemma 3.9.

(l).  , $q \in R_3$. GDD 2 of Row 16.

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(II) or by Lemma 3.2 (X).

(n).  , $q \in R_3$. GDD 1 of Row 17.
is empty.

(n') or (11).  , $q \in R_3$. GDD 1 of Row 17.


Omitting 1 in whole is an arithmetic root system by GDD 1 of Row 17 or by Lemma 3.3 (V) or by Lemma 3.2 (XI). Whole is affine by Lemma 3.9.

(o).  , $q \in R_3$. GDD 4 of Row 17.

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2 (XI).

(p).  , $q \in R_3$. GDD 4 of Row 17.

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2 (XI).



, $q \in R_3$. GDD 9 of Row 17.

is empty.

Case (iii) Omitting in whole 1 and adding 3.


It is the same as Case (i) when we change q in Case (i) into q^{-1} .

Case (iv) Containing circle with four vertexes.


According to Lemma 3.39 (ii) we have to consider following cases.

(a') to (a') empty.

(a') to (a'') or (12)




with $q^{-1} = q^2$ is an arithmetic root system by Lemma 3.2 (XI)(h) or Lemma 3.3 (III)(h).




is affine.

(a') to (a) or (13)




with $q = q^{-2}$ is an arithmetic root system.




is affine.

(a') to (c) or (14)




with $q = q^{-3}$ is an arithmetic root system,



is affine.


(a') to (e) empty. (a') to (j) empty. (a') to (k) empty. (a') to (l) empty. (a') to (n') empty. (a') to (o) empty. (a') to (p) empty. (a'') to (a'') empty.

(a'') to (a)



with $q = q^{-4}$ is not an arithmetic root system by Lemma 3.3 (III).


(a'') to (c)



with $q^{-6} = q$ is not an arithmetic root system by Lemma 3.3 (III).

(a'') to (e) empty. (a'') to (j) empty. (a'') to (k) empty. (a'') to (l) empty. (a'') to (n') empty. (a'') to (o) empty. (a'') to (p) empty.


(a) to (a) or (15)



with $q^2 = q^{-2}$ is an arithmetic root system.

is affine.


(a) to (c) or (16)




with $q^2 = q^{-3}$ is arithmetic root system.

is affine.

(a) to (e) or (17)




with $q^2 = -1$ is arithmetic root system by Lemma 3.2 (XI)(e).



is affine.

(a) to (j) empty. (a) to (k) empty. (a) to (l) empty. (a) to (n') empty. (a) to (o) empty. (a) to (p) empty.


(c) to (c) or (18)




with $q^3 = q^{-3}$ is an arithmetic root system.

is affine.

(c) to (e) or (19)




with $q^3 = -1$ is arithmetic root system by Lemma 3.2 (XI)(e).




is affine.

(c) to (j) empty. (c) to (k) empty. (c) to (l) empty. (c) to (n') empty. (c) to (o) empty.

(c) to (p) or (20)




with $q^3 = -1$ is arithmetic root system by Lemma 3.2 (XI)(e).




is affine.

(e) to (e) or (21)




is arithmetic root system by Lemma 3.2 (XI).




is affine. (e) to (j) empty.

(e) to (k) or (22)




is an arithmetic root system by Lemma 3.3 (IV)(k) or by Lemma 3.2 (XI)(b).




is affine.

(e) to (l) empty.

(e) to (n') or (23)




is an arithmetic root system by Lemma 3.2 (XI)(e).




is affine.

(e) to (o) or (24)




is an arithmetic root system by Lemma 3.2 (XI)(c) or Lemma 3.3 (IV)(o).



is affine.
 (e) to (p) or (25)


is an arithmetic root system by Lemma 3.2 (XI)(c) or Lemma 3.3 (IV)(o).



is affine with $q \in R_6$.

(j) to (j) empty. (j) to (k) empty. (j) to (l) empty. (j) to (n') empty. (j) to (o) empty. (j) to (p) empty.

(k) to (k) or (26)




is arithmetic root system.

is affine.

(k) to (l) empty.


(k) to (n') or (27)



is an arithmetic root system by Lemma 3.2 (XI)(e).

is affine.


(k) to (o)



is not an arithmetic root system by Lemma 3.2 (XI) or Lemma 3.3 (IV).

(k) to (p) empty. (l) to (l) empty. (l) to (n') empty. (l) to (o) empty. (l) to (p) empty.


(n') to (n') or (28)




is arithmetic root system.

is affine.

(n') to (o) or (29)




is arithmetic root system by Lemma 3.2 (XI)(e)




is affine.

(n') to (p) empty.

(o) to (o) or (30)




is arithmetic root system.




is affine.

(o) to (p) empty.

(p) to (p) or (31)



is arithmetic root system,




is affine with $q \in R_6$.

4.18 GDD 3 of Row 8


Case (i) Omitting 3 and adding 1.

According to Lemma 3.3 (I) we have to consider following cases.




GDD 2 of Row 4.

is an arithmetic root system by Type 7.




$q \notin R_4$, GDD 3 of Row 5.

is an arithmetic root system by Type 3.



$q^2 \neq 1$, GDD 2 of Row 6.

is affine by Lemma 3.15 when $q \notin R_3$. It is an arithmetic root system by Lemma 3.15(d) when $q \in R_3$.



GDD 2 of Row 7.

is affine by Lemma 3.15.

$$\begin{array}{c} \text{(f) or (3)} \\ -q \frac{1}{q^2} \quad -1 \quad q^{-3}q^3 \quad q^{-3} \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array}$$

, $q \notin R_2 \cup R_3$, GDD 4 of Row 7.

is affine by Lemma 3.4.

$$\begin{array}{c} \text{(g)} \\ q^{-1} \frac{1}{q} \quad -1 \quad q^{-1}q \quad q^{-1} \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}$$

, GDD 1 of Row 8.

is an arithmetic root system by Type 7.

$$\begin{array}{c} \text{(h) or (4)} \\ r \quad r^{-1} \quad -1 \quad q^{-1}q \quad q^{-1} \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}$$

, GDD 1 of Row 9.

is arithmetic root system when $r = q^{-2}$ with $q \notin R_3$. by Type 1. Otherwise it is affine by Lemma 3.35 when $r \neq -1$; It is affine by Lemma 3.23 when $r = -1$ and $q \notin R_4$.

$$\begin{array}{c} \text{(i')} \text{ or (5)} \\ q^{-1} \frac{1}{q} \quad -1 \quad q^{-2}q^2 \quad q^{-2} \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}$$

, $q \notin R_3 \cup R_2$. GDD 2 of Row 10.

is affine by Lemma 3.35.

$$\begin{array}{c} \text{(k) or (6)} \\ q \quad q^{-1} \quad -1 \quad q^{-1}q \quad q^{-1} \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}$$

, $q \notin R_3$. GDD 1 of Row 10.

is an arithmetic root system whrn $q \in R_4$ by GDD 3 of Row 22 or by Lemma 3.35(f). Otherwise it is affine by Lemma 3.35.

$$\begin{array}{c} \text{(l) or (7)} \\ q^{-2} \frac{1}{q^2} \quad -1 \quad q^{-1}q \quad q^{-1} \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}$$

, $q \notin R_3$. GDD 2 of Row 10.

is an arithmetic root system by Type 1.

$$\begin{array}{c} \text{(m)} \\ q \quad q^{-1} \quad -1 \quad q^{-1}q \quad q^{-1} \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}$$

, $q \in R_3$. GDD 1 of Row 11.

is an arithmetic root system by GDD 3 of Row 12 or Type 1.

$$\begin{array}{c} \text{(n)} \\ q^{-1} \quad -q^{-1-1} \quad -q \quad -q^{-1}q \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}$$

, $q \in R_3$. GDD 3 of Row 14.

is an arithmetic root system by Type 4.

$$\begin{array}{c} \text{(p)} \\ -1 \quad q \quad -1 \quad -q^{-1}-q \quad -q^{-1}-1 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}$$

, $q \in R_3$. GDD 2 of Row 16.

is affine by Lemma 3.15.

$$\begin{array}{c} \text{(r) or (8)} \\ q \quad -1 \quad -1 \quad -q^{-1}-q \quad -q^{-1}-1 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}$$

, $q \in R_3$. GDD 4 of Row 16.

is affine by Lemma 3.4.


$$\begin{array}{c} \text{(s')} \\ -1 \quad q \quad -1 \quad -1 \quad -1 \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}$$

, $q \in R_3$. GDD 1 of Row 17.


is empty.

Case (ii) Omitting 3 and adding 2.

According to Lemma 3.3 (III) we have to consider following cases.

(a) or (9).  , $q \in F^* \setminus \{1, -1\}$. GDD 1 of Row 6.

Omitting 1 in whole is arithmetic root system by symmetry. It is similar in following. Whole is affine when $q \notin R_3$ by Lemma 3.4. Whole is arithmetic root system when $q \in R_3$ by Lemma 3.4(c).

(c) or (10).  , $q \in F^* \setminus \{1, -1\}$, $q \notin R_3$. GDD 1 of Row 7.

is affine by Lemma 3.4.

(e) or (11).  , $q \in F^* \setminus \{1, -1\}$. GDD 3 of Row 8.

is affine checked step by step.

(g) or (12).  , $q \in R_3$. GDD 1 of Row 14.

is affine by Lemma 3.4.

(h) or (13).  , $q \in R_3$. GDD 1 of Row 15.


is affine by Lemma 3.4.

(j) or (14).  , $q \in R_3$. GDD 1 of Row 16.

is affine by Lemma 3.4.


(l) or (15).  , $q \in R_3$. GDD 2 of Row 17.

is affine by Lemma 3.4.




, $q \in R_3$. GDD 5 of Row 17.

is affine by Lemma 3.4.




, $q \in R_3$. GDD 7 of Row 17.

is affine by Lemma 3.4.




GDD 1 of Row 4.

is an arithmetic root system when $q \in R_3$ by GDD 6 of Row 18. It is an affine when $q \notin R_3$ checked step by step.




GDD 1 of Row 5.

is affine by Lemma 3.4.



GDD 1 of Row 17.




GDD 9 of Row 17.

Case (iii) Omitting 1 and adding 3 is the same as Case (i)

Case (iv) containing circle with four vertexes.


(a') to (a') or (20)



is an arithmetic root system when $q \in R_3$ by Lemma 3.2 (XI)(b).

is affine.

- (a') to (a) empty.
- (a') to (b) or (21)




is an arithmetic root system when $q \in R_3 \cup R_4$ by Lemma 3.2 (XI)(a) (e).

is affine.

- (a') to (d) or (22)


is an arithmetic root system when $q^3 = q^{-1}$ or $q^3 = -1$ by Lemma 3.2 (XI).



is affine.

- (a') to (f) empty.(a') to (g) empty.
- (a') to (h) or (23)


with $r = -1$ is an arithmetic root system when $q \in R_3 \cup R_4 \cup R_6$ by Lemma 3.2 (XI)(e).



is affine.

- (a') to (i') empty. (a') to (k) empty.
- (a') to (l) or (24)


with $-1 = q^{-2}$ is an arithmetic root system by Lemma 3.2 (XI)(e).




is affine.

- (a') to (m) empty.(a') to (n) empty.
- (a') to (p) or (25)

with $q \in R_6$ is an arithmetic root system by Lemma 3.2 (XI)(d).




is affine.



(a') to (r) empty.

(a) to (a) or (26)

is an arithmetic root system when $q^3 = 1$ by Lemma 3.2 (V)(b).




is affine.

(a) to (b) empty. (a) to (d) empty. (a) to (f) empty. (a) to (g) empty.

(a) to (h) or (27)

with $r^2 = -q^{-1}$ is an arithmetic root system when $q \in R_6$ by Lemma 3.3 (III)(h).





is affine.

(a) to (i') empty. (a) to (k) empty. (a) to (l) empty. (a) to (m) empty. (a) to (n)

empty. (a) to (p) empty. (a) to (r) empty.

(b) to (b) or (28)



is an arithmetic root system by Lemma 3.2 (XI)(b).



is affine.

(b) to (d) or (29)

is an arithmetic root system when $q^3 = q^{-2}$.





is affine.

(b) to (f) empty.

(b) to (g) or (30)


with $-1 = q^{-2}$ is arithmetic root system by Lemma 3.2 (XI)(e).



is affine.

(b) to (h) or (31)

with $-1 = r^2$ is an arithmetic root system when $q \in R_3 \cup R_4 \cup R_6$ by Lemma 3.2 (XI)(e).




is affine.

(b) to (i') empty.

(b) to (k) or (32)

with $-1 = q^2$ is an arithmetic root system by Lemma 3.2 (XI)(e).



is affine.


(b) to (l)

with $-1 = q^{-4}$ is not an arithmetic root system by Lemma 3.2 (XI)(e).

(b) to (m) empty. (b) to (n) empty.

(b) to (p) or (33)

is an arithmetic root system.




is affine.

(b) to (r) empty.

(d) to (d)


is not an arithmetic root system by Lemma 3.2 (XI).



(d) to (f) empty.

(d) to (g) or (34)


with $-1 = q^{-3}$ is an arithmetic root system by Lemma 3.2 (XI)(e).



is affine.


(d) to (h) or (35)


with $-1 = r^3$ is an arithmetic root system with $q \in R_4 \cup R_6$ by Lemma 3.2 (XI)(e).




is affine.

(d) to (i')






with $-1 = q^{-3}$ is an arithmetic root system by Lemma 3.2 (XI)(e).



is affine.

(d) to (k) or (36)


with $-1 = q^3$ is an arithmetic root system by Lemma 3.2 (XI)(e).



is affine.

(d) to (l) or (37)

with $-1 = q^{-6}$ is an arithmetic root system by Lemma 3.2 (XI)(e).




is affine.

(d) to (m) empty. (d) to (n) empty. (d) to (p) empty. (d) to (r) empty.


(f) to (f)

is not an arithmetic root system by Lemma 3.2 (V).

(f) to (g) or (38)




with $q^{-3} = -q^{-1}$ is arithmetic root system by Type 1.



is affine.

(f) to (h) or (39)

with $r^3 = -q^{-1}$ is arithmetic root system when $q \in R_{18} \cup R_6$ by Lemma 3.3 (III).




is affine.

(f) to (i') empty.

(f) to (k)

with $-q^{-1} = q^3$ is not arithmetic root system by Lemma 3.3 (III).

(f) to (l)





with $-q^{-1} = q^{-6}$ is not arithmetic root system by Lemma 3.3 (III)

(f) to (m) empty.

(f) to (n) empty. (f) to (p) empty. (f) to (r) empty.

(g) to (g) or (40)



is arithmetic root system.



is affine.

(g) to (h) empty. (g) to (i') empty. (g) to (k) empty. (g) to (l) empty. (g) to (m) empty. (g) to (n) empty. (g) to (p) empty. (g) to (r) empty.

(h) to (h) or (41)





with $r' = r$ is arithmetic root system.

is affine.

(h) to (i') or (42)


with $q^{-1} = r^2$ is arithmetic root system.




is affine.

(h) to (k) empty.

(h) to (l) or (43)




with $r = q^{-2}$ is arithmetic root system.



is affine.

(h) to (m) empty.

(h) to (n) or (44)




with $-q = r$, $q \in R_6$ is arithmetic root system by Lemma 3.3 (III)(o).

is affine.

(h) to (p) or (45)


with $r = -1$ is arithmetic root system by Lemma 3.2 (XI)(e).



is affine.

(h) to (r) or (46)

with $-q = r$ is arithmetic root system when $q \in R_6$ by Lemma 3.3 (III)(l); when $q \in R_4$ by Type 1.



is affine.

(i') to (i') or (47)





is arithmetic root system,

is affine.

(i') to (k) empty.(i') to (l) empty.(i') to (m) empty.

(i') to (n) or (48)

with $-q^2 = q^{-1}$ is arithmetic root system by Lemma 3.3 (III)(n).




is affine.

(i') to (p) empty.



(i') to (r)

with $-q^2 = q^{-1}$, $q \in R_6$ is not an arithmetic root system by Lemma 3.3 (III).



(k) to (k) or (49)


is arithmetic root system.




is affine.

(k) to (l) empty.(k) to (m) empty.(k) to (n) empty. (k) to (p) empty.(k) to (r) empty.

(l) to (l) or (50)




is arithmetic root system.




is affine.

(l) to (m) empty.(l) to (n) empty. (l) to (p) empty.

(l) to (r) or (51)




with $-q = q^{-2}$ is arithmetic root system by Lemma 3.3 (III)(1).



is affine.

(m) to (m) or (52)

is arithmetic root system.



is affine.

(m) to (n) empty. (m) to (p) empty.(m) to (r) empty.

(n) to (n)



is not arithmetic root system by Lemma 3.2 (V).

(n) to (p)


empty.(n) to (r)

is not arithmetic root system by Lemma 3.2 (V).

(p) to (p) or (53)




is arithmetic root system by Lemma 3.2 (XI)(b).




is affine.

(p) to (r) empty.



(r) to (r)



is not arithmetic root system by Lemma 3.2 (V).


4.19 GDD 4 of Row 8

It is the same as GDD 3 of Row 8.

4.20 GDD 1 of Row 9


Case (i) Omitting 3 and adding 1.

According to Lemma 3.3 (III) we have to consider following cases.




$q^2 \neq 1$, GDD 1 of Row 6.

is an arithmetic root system when $r = q^{-3}$ and $q^4 \neq 1$ by GDD 6 of Row 9 table B or by Lemma 3.7(a). It is affine when $r \neq q^{-3}$ by Lemma 3.7.




, $q \in F^* \setminus \{1, -1\}$, $q \notin R_3$. GDD 1 of Row 7.

is affine by Lemma 3.7.




$q \in F^* \setminus \{1, -1\}$. GDD 3 of Row 8.

is an arithmetic root system when $r = q^{-2}$ and $q \notin R_3$ by Type 1. It is affine when $r \neq q^{-2}$ by Lemma 3.35.




, $q \in R_3$. GDD 1 of Row 14.

is affine by Lemma 3.4.




, $q \in R_3$. GDD 1 of Row 15.

is affine by Lemma 3.7.



, $q \in R_3$. GDD 1 of Row 16.

is an arithmetic root system when $r = -q$ by GDD 5 of Row 17 or by Lemma 3.7(c). Otherwise it is affine by Lemma 3.7.



, $q \in R_3$. GDD 2 of Row 17.

is affine by Lemma 3.7.

$$\begin{array}{c} \text{(n) or (8).} \\ -1 \quad -q \quad -q^{-1} \quad -1 \quad -r^{-1} \quad -r \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \\ \hline \end{array}$$

, $q \in R_3$. GDD 5 of Row 17.

is affine by Lemma 3.7.

$$\begin{array}{c} \text{(o) or (9).} \\ -1 \quad -q \quad q \quad q^{-1} \quad -1 \quad r^{-1} \quad r \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \\ \hline \end{array}$$

, $q \in R_3$. GDD 7 of Row 17.

is affine by Lemma 3.7.

$$\begin{array}{c} \text{(a') or (10).} \\ q^{-1} \quad q \quad q^{-1} \quad q \quad -1 \quad r \quad r^{-1} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \\ \hline \end{array}$$

GDD 1 of Row 4.

Whole is arithmetic root system when $q = -r^{-1}$ and $q^2 \neq 1$ by Lemma 3.35(g). Whole is an arithmetic root system when $q^3 = r^2$ and $q^6 \neq 1$ by Lemma 3.35(b). Whole is arithmetic root system when $q^2 = r^{-1}$ by Type 1. Otherwise Whole is affine by Lemma 3.35.

$$\begin{array}{c} \text{(a'') or (11)} \\ q \quad q^{-2} \quad q^2 \quad q^{-2} \quad -1 \quad r^{-2} \quad r^2 \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \\ \hline \end{array}$$

$q^4 \neq 1$, GDD 1 of Row 5.

is affine by Lemma 3.4.

$$\begin{array}{c} \text{(k') or (12)} \\ -1 \quad q \quad -1 \quad -1 \quad -1 \quad r^{-1} \quad r \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \\ \hline \end{array}$$

$q \in R_3$, GDD 1 of Row 17.

is affine when $r \neq -1$ by Lemma 3.26.

$$\begin{array}{c} \text{(p') or (13)} \\ q^{-1} \quad -q^{-1} \quad -1 \quad -1 \quad -1 \quad r^{-1} \quad r \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \\ \hline \end{array}$$

$q \in R_3$, GDD 9 of Row 17.

is affine when $r \neq -1$ by Lemma 3.26 or by Lemma 3.4.

(ii) Omitting 3 and adding 2.

According to Lemma 3.3 (I) we have to consider following cases.

$$\begin{array}{c} \text{(a') or (14)} \\ \begin{array}{c} r \\ \bullet \\ r^{-1} \\ | \\ \bullet \\ -1 \quad q \quad -1 \quad q^{-1} \quad q \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \\ \hline \end{array} \end{array}$$

GDD 2 of Row 4.

Omitting 1 in whole is arithmetic root system when $r = -q$, and $q \in R_3$ or $r = q^3$, and $q^4 \neq 1$ or $r = q^2$ and $r \notin R_3$ by Lemma 3.2 (X). Whole is affine by Lemma 3.12.


$$\begin{array}{c} \text{(a) or (15)} \\ \begin{array}{c} r^2 \\ \bullet \\ r^{-2} \\ | \\ \bullet \\ -q^{-1}q^2 \quad -1 \quad q^{-2}q^2 \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \\ \hline \end{array} \end{array}$$

GDD 3 of Row 5.


Omitting 1 in whole is arithmetic root system when $r^{-2} = q^{-3}$ or $r^{-2} = q^2$ by Lemma 3.2(I)(h) and Type 3 with $q \notin R_3$. Whole is affine by Lemma 3.4.

(b) or (16)  GDD 2 of Row 6.

Omitting 1 in whole is arithmetic root system when $r^2 = q$, with $q^2, q^3 \neq 1$; or $r^2 = q^3$, $q^3, q^5 \neq 1$; or $r^2 = -q, q \in R_3$; or $r^2 = -1, q \in R_3 \cup R_6$. Whole is affine by Lemma 3.12.

(d) or (17)  $q^2, q^3 \neq 1, q^3 \neq -1$, GDD 2 of Row 7.

Omitting 1 in whole is arithmetic root system when $r^3 = q$ or $r^3 = q^2, q^5 \neq 1$ or $q^5 = -1, q \in R_4$ by Lemma 3.2 (X) (XI). Whole is affine when $r^3 = q$ or $r^3 = q^2, q^5 \neq 1$ or $q^5 = -1, q \in R_4$ by Lemma 3.12.

(f) or (18)  $q^3 \neq 1$, GDD 4 of Row 7.

Omitting 1 in whole is arithmetic root system when $r^3 = q^2, q \notin R_5$ by Lemma 3.2(I)(h) or Type 3. Whole is affine by Lemma 3.4

(g) or (19)  , GDD 1 of Row 8.

Omitting 1 in whole is arithmetic root system by Lemma 3.2 (IV). Whole is affine by Lemma 3.8 and by Lemma 3.5.

(h) or (20)  , GDD 1 of Row 9.

Omitting 1 in whole is arithmetic root system by Lemma 3.2 (IV). Whole is affine by Lemma 3.8 and by Lemma 3.5.

(i') or (21)  $q \notin R_3$. GDD 2 of Row 10.

Omitting 1 in whole is arithmetic root system by Lemma 3.2 (IV). Whole is affine by Lemma 3.8 and by Lemma 3.5.

(k) or (22)  $q \notin R_3$. GDD 1 of Row 10.


Omitting 1 in whole is arithmetic root system by Lemma 3.2 (IV). Whole is affine by Lemma 3.8 and by Lemma 3.5.

(l) or (23)  , $q \notin R_3$. GDD 2 of Row 10.

Omitting 1 in whole is arithmetic root system by Lemma 3.2 (IV). Whole is affine by Lemma 3.8 and by Lemma 3.5.

(m) or (24)  , $q \in R_3$. GDD 1 of Row 11.

Omitting 1 in whole is arithmetic root system by Lemma 3.2 (IV). Whole is affine by Lemma 3.8 and by Lemma 3.5.

(n) or (25)  , $q \in R_3$. GDD 3 of Row 14.

Omitting 1 in whole is arithmetic root system when $-r = -1$ by Lemma 3.2(I)(c). Whole is affine by Lemma 3.4.

(p) or (26)  , $q \in R_3$. GDD 2 of Row 16.

Omitting 1 in whole is arithmetic root system when $r = -q$; or $r = -q^2$; or $r = 1$, $q \in R_3$ by Lemma 3.2 (X) (XI). affine by Lemma 3.25 and by Lemma 3.5.

(r)  , $q \in R_3$. GDD 4 of Row 16.

Omitting 1 in whole is not arithmetic root system by Lemma 3.2(I).

(s') or (27)  , $q \in R_3$. GDD 1 of Row 17.

Omitting 1 in whole is arithmetic root system when $r = -q$, $q \in R_3$ or $r = -1$ by Lemma 3.2 (X) (XI). Whole is affine by Lemma 3.12.

Case (iii) Omitting 1 and adding 3.


It is the same as Case (i) .

Case (iv) containing circle with four vertexes.

According to Lemma 3.39 (ii) we have to consider following cases.

(a) to (a) or (28)

with $r^2 = q^2$ is arithmetic root system by Type 2.



is affine.

(a) to (c)

with $q^2 = r^3$ is not arithmetic root system by Lemma 3.2 (I).

(a) to (e)

with $q^2 = -1$ is not arithmetic root system by Lemma 3.2 (I).

(a) to (g)

with $-r^{-1} = q^2$ is not an arithmetic root system by


Lemma 3.2 (I).

(a) to (h) empty.

(a) to (j)

with $-r = q^2$ is not arithmetic root system by Lemma 3.2 (I).

(a) to (l)



with $-1 = q^2$ is not an arithmetic root system by Lemma 3.2 (I).


(a) to (n)

with $q^2 = -1$ is not an arithmetic root system by Lemma 3.2 (I).

(a) to (o) empty.

(a) to (a') or (29)

with $r = q^2$ is an arithmetic root system by Type 2.




is affine.

(a) to (a'')

with $r = q^4$ is not an arithmetic root system by Lemma 3.2 (I).

(a) to (k')

and $q^2 = -1$ with $s \in R_3$ is not an root system by Lemma 3.2(I).



(a) to (p') $\begin{array}{c} q \quad q^{-2} \quad q^2 \quad -s^{-1}-1 \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array}$
 with $q^2 = s^{-1}$, $s \in R_3$ is not an arithmetic root system by Lemma 3.2(I).

(c) to (c) $\begin{array}{c} q \quad q^{-3} \quad q^3 \quad r^{-3}r \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array}$
 with $r^3 = q^3$ is not an arithmetic root system by Lemma 3.2 (I).

(c) to (e) $\begin{array}{c} q \quad q^{-3} \quad q^3 \quad r^{-1}r \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array}$
 with $-1 = q^3$ is not an arithmetic root system by Lemma 3.2 (I).

(c) to (g) $\begin{array}{c} q \quad q^{-3} \quad q^3 \quad r^{-1}r \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array}$
 with $q^3 = -r^{-1}$ with $r \in R_6$ is not an arithmetic root system by Lemma 3.2 (I).

(c) to (h) $\begin{array}{c} q \quad q^{-3} \quad q^3 \quad r \quad r \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array}$
 with $-1 = q^3$, $r \in R_3$ is not an arithmetic root system by Lemma 3.2 (I).

(c) to (j) $\begin{array}{c} q \quad q^{-3} \quad q^3 \quad -r^{-1}r \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array}$
 with $-r = q^3$, $r \in R_3$ is not an arithmetic root system by Lemma 3.2 (I).

(c) to (l) $\begin{array}{c} q \quad q^{-3} \quad q^3 \quad -1 \quad r \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array}$
 with $-1 = q^3$, $r \in R_3$ is not an arithmetic root system by Lemma 3.2 (I).

(c) to (n) $\begin{array}{c} q \quad q^{-3} \quad q^3 \quad -r \quad r \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array}$

with $-1 = q^3$ is not an arithmetic root system by Lemma 3.2 (I).

(c) to (o) $\begin{array}{c} q \quad q^{-3} \quad q^3 \quad -r \quad r \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array}$
 with $-1 = q^3$, $r \in R_3$ is not an arithmetic root system by Lemma 3.2 (I).

(c) to (a') $\begin{array}{c} q \quad q^{-3} \quad q^3 \quad r^{-1}r \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array}$

with $r = q^3$, $r \in R_3$ is not an arithmetic root system by Lemma 3.2 (I).
 (c) to (a'') $\begin{array}{c} q^2 \quad q^{-6} \quad q^6 \quad r^{-2}r^2 \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array}$

with $r = q^6$ is not an arithmetic root system by Lemma 3.2 (I).
 (c) to (k') $\begin{array}{c} q \quad q^{-3} \quad q^3 \quad s \quad -1 \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array}$


with $q^3 = -1$, $s \in R_3$ is not an arithmetic root system by Lemma 3.2(I).

(c) to (p') $\begin{array}{c} q \quad q^{-3} \quad q^3 \quad -s^{-1}-1 \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array}$

with $q^3 = s^{-1}$, $s \in R_3$ is not an arithmetic root system by Lemma 3.2(I) .

(e) to (e) or (30)

is an arithmetic root system by Lemma 3.2 (IV).



is affine.

(e) to (g) empty.


(e) to (h) or (32)

with $r \in R_3$ is not an arithmetic root system by Lemma 3.2(I).

(e) to (j) empty.

(e) to (l) or (31)


is an arithmetic root system when $q = -r$ and $r \in R_3$ by Lemma 3.2(I)(j).



is affine.

(e) to (n)


with $r \in R_6$ is arithmetic root system when $-r^{-1} = q^{-1}$ by Type 3.



is affine.

(e) to (o) or (33)

is arithmetic root system when $r = -q$ by Type 4.




is affine.

(e) to (a') empty. (e) to (a'') empty.

(e) to (k') or (34)

with $s \in R_3$ is an root system when $q = s^2$ or $q = s$ or $q = s^3$ or $q = -s$ by Lemma 3.2 (X).



is affine.

(e) to (p') empty. (g) to (g) empty. (g) to (h) empty. (g) to (j) empty. (g) to (l) empty. (g) to (n) empty. (g) to (o) empty.

(g) to (a')

with $-q^{-1} = r$ is not an arithmetic root system by Lemma 3.3(II).

(g) to (a'')

with $r = -q^{-2}$ is not arithmetic root system by Lemma 3.2 (V).

(g) to (k') empty.

(g) to (p')

with $s^{-1} = -q^{-1}$ with $s \in R_3$ is not arithmetic root system by Lemma 3.2 (V).

(h) to (h)

is not an arithmetic root system by Lemma 3.2(I).

(h) to (j) empty.

(h) to (l)

is not an arithmetic root system with $r, q \in R_3$.

(h) to (n)

is not an arithmetic root system with $r \in R_6$ by Lemma 3.2 (I).

(h) to (o)

is not an arithmetic root system by Lemma 3.2 (I).

(h) to (a') empty. (h) to (a'') empty.

(h) to (k') or (35)

is an arithmetic root system when $s = q^{-1}$ with $q, s \in R_3$. by Type 1. Whole is not an arithmetic root system when $s \neq q^{-1}$ by Lemma 3.2(I).

is affine.

(h) to (p') empty. (j) to (j) empty. (j) to (l) empty. (j) to (n) empty. (j) to (o) empty.

(j) to (a') or (36)

with $-q = r$ is an arithmetic root system by Lemma 3.2 (I)(b).

is affine.

(j) to (a'')














with $r = -q^2$ is not an arithmetic root system by Lemma 3.2(I),

(j) to (k') empty. (j) to (p') empty.

(l) to (l)


is not an arithmetic root system with $q \in R_3$ by Lemma 3.2 (I).

(l) to (n)



is not an arithmetic root system by Lemma 3.2 (I).


(l) to (o)



is not an arithmetic root system by Lemma 3.2 (I).

(l) to (a') empty. (l) to (a'') empty.


(l) to (k')



is not an arithmetic root system by Lemma 3.2(I).


(l) to (p') empty.

(n) to (n)



is not an arithmetic root system with $q \in R_3$ by Lemma 3.2(I).


(n) to (o)



is not an arithmetic root system with $r \in R_3, q \in R_6$ by Lemma 3.2 (I).

(n) to (a') empty. (n) to (a'') empty.


(n) to (k')



is not an arithmetic root system by Lemma 3.2(I) with $s \in R_3$. by Lemma 3.2.


(n) to (p') empty.

(o) to (o)



is not an arithmetic root system by Lemma 3.2 (I).(o) to (a') empty. (o) to (a'') empty.


(o) to (k')




is not an arithmetic root system by Lemma 3.2(I).

(o) to (p') empty. (a') to (a') empty.

(a') to (a'') or (37)




with $r = q^2$ is an arithmetic root system by Type 1.




is affine.

(a') to (k') empty.

(a') to (p') or (38)




with $s^{-1} = q$ is an arithmetic root system by Lemma 3.3 (II)(f).



is affine.


(a'') to (a'') empty. (a'') to (k') empty.

(a'') to (p')




with $s^{-1} = q, s \in R_3$ is not an arithmetic root system by Lemma 3.2 (V).

(k') to (k') or (39)



is an arithmetic root system when $s = t$ or $s^{-1} = t \in R_3$. by Lemma 3.2 (XI).



is affine.

(k') to (p')empty.

(p') to (p')

with $t^{-1} = s^{-1}$ is not an arithmetic root system by Lemma 3.2 (V).


4.21 GDD 2 of Row 9

GDD 2 of Row 9 as GDD 1 of Row 9.

4.22 GDD 3 of Row 9


Case (i) Omitting 3 and adding 1.

According to Lemma 3.3 (IV) we have to consider following cases.




, $q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 4.

Omitting 2 in whole is an arithmetic root system when $q = s^2$, $s^3 \neq 1$, or $q = s^3$, with $s^2, s^3, s^4 \neq 1$, or $q = -s$, $s \in R_3$ or $s = -1$, $q^2, r^2 \neq 1$ by Lemma 3.2 (X). Whole is affine by Lemma 3.12.



, $q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 5.


Omitting 2 in whole is an arithmetic root system when $s = -q$ by Type 2. Whole is affine by Lemma 3.4.




, $q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 6.

Omitting 2 in whole is an arithmetic root system when $q^2 = s$, or $s^2 = q^2$, or $s^3 = q^2$, or $q^2 = -s$, $s \in R_3$ or $s = -1$ or $q^2 = -1$, $s \in R_2 \cup R_3 \cup R_4 \cup R_6$ by Lemma 3.2 (X), (IV) and (XI)(e). Whole is affine by Lemma 3.16.

(b) or (4).




Omitting 2 in whole is an arithmetic root system when $q = s^2$, $1 \neq s^3$, or $q = s^3$ with $s^2, s^3, s^4 \neq 1$, or $s = -1$ or $s = -q$, $s \in R_3$ by Lemma 3.2 (X). Omitting 1 in whole is an arithmetic root system when $q = r^2$, $1 \neq r^3$, or $q = r^3$ with $r^2, r^3, r^4 \neq 1$, or $r = -1$ or $r = -q$, $r \in R_3$ by Lemma 3.2 (X). Whole is affine.



Omitting 2 in whole is an arithmetic root system when $q^3 = s$ or $s^2 = q^3$, or $q^3 = s^3$ or $-s = q^3$, $s \in R_3$ or $q^3 = -1$, $s \in R_2 \cup R_3 \cup R_4 \cup R_6$ by Lemma 3.2 (X)(XI) and by Lemma 3.3 (V). Whole is affine by Lemma 3.16.


(d) or (6).



Omitting 2 in whole is an arithmetic root system when $s^3 = q$ or $s^6 = q$ or $s^9 = q$ or $-s^3 = q$, $s \in R_9$ or $s^3 = -1$ by Lemma 3.2 (X)(IV). Omitting 1 in whole is an arithmetic root system when $r^3 = q^2$ by Type 2. Whole is affine.



Omitting 2 in whole is an arithmetic root system when $s = -1$, $q \in R_3 \cup R_4 \cup R_6$ or $s = -q$ with $s \in R_3$ or $q \in R_3$ by Lemma 3.2 (XI). Whole is affine by Lemma 3.16.



$q, r, s \in F^* \setminus \{1\}$, $qrs = 1$, $q \neq r$, $q \neq s$, $s \neq r$. GDD 3 of Row 9.


$q', r', s' \in F^* \setminus \{1\}$, $q'r's' = 1$, $q' \neq r'$, $q' \neq s'$, $s' \neq r'$. $q = q'$

Omitting 2 in whole is an arithmetic root system when $s' = s$, $s \in R_3$ or $s' = s^{-1}$, or

$s' = -s$ with $s \in R_3 \cup R_6$, or $s' = -1, s \in R_2 \cup R_3 \cup R_4 \cup R_6$ or $s = -1, s' \in R_2 \cup R_3 \cup R_4 \cup R_6$ by Lemma 3.2 (XI).


Omitting 1 in whole is an arithmetic root system when $r' = r, r \in R_3$ or $r' = r^{-1}$ or $r' = -r$ with $r \in R_3 \cup R_6$, or $r' = -1, r \in R_2 \cup R_3 \cup R_4 \cup R_6$ or $r = -1, r' \in R_2 \cup R_3 \cup R_4 \cup R_6$ by Lemma 3.2 (XI). Whole is affine.

(g) or (9).



Omitting 2 in whole is an arithmetic root system when $q = -s$ with $q \in R_3 \cup R_6$, or $s = -1, q \in R_3 \cup R_4 \cup R_6$. Omitting 1 in whole is an arithmetic root system when $r = q^{-2}$ with $r \in R_3$, or $r^{-1} = q^{-2}$ or $-r = q^{-2}$ with $r \in R_3 \cup R_6$, or $-1 = q^{-2}, r \in R_2 \cup R_3 \cup R_4 \cup R_6$. or $-1 = r, q^{-2} \in R_2 \cup R_3 \cup R_4 \cup R_6$ by Lemma 3.2 (XI). Whole is affine.


(h) or (10).



Omitting 2 in whole is an arithmetic root system when $q = s^2$ or $q = -s^{-2}, q \in R_6$ or $-1 = s^{-2}, q \in R_4 \cup R_6$ by Lemma 3.2 (XI).

Omitting 1 in whole is an arithmetic root system when $q = r^2$ or $q = -r^{-2}, q \in R_6$ or $-1 = r^{-2}, q \in R_4 \cup R_6$ by Lemma 3.2 (XI). affine.

(i) or (11).



Omitting 2 in whole is an arithmetic root system when $q = -s$ or $s = -1$.

Omitting 1 in whole is an arithmetic root system when $-r = q$ or $r = -1$. Whole is affine.

(j) or (12).  , $q \in R_3$. GDD 2 of Row 14.


Omitting 2 in whole is an arithmetic root system when $q = -s^{-1}$, $q = s^{-2}$, $-s^{-3} = q$ by Lemma 3.2 (X). Whole is affine by Lemma 3.4.

(k) or (13).  , $q \in R_3$. GDD 2 of Row 15.

Omitting 2 in whole is an arithmetic root system when $q = s$ (omitting) or $s = q^{-1}$ (omitting), or $s = -q$ ($q \in R_3$) or $s = -1$ ($q \in R_3$) by Lemma 3.2 (XI). Whole is affine by Lemma 3.16.

(l) or (14).  , $q \in R_3$. GDD 2 of Row 16.

Omitting 2 in whole is an arithmetic root system when $s = -q$ or $-q = s^2$ or $-q = s^3$, $s^3 \neq 1$, or $s = -1$ by Lemma 3.2 (XI)(IV). Whole is affine by Lemma 3.16

(m).  , $q \in R_3$. GDD 3 of Row 16.

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(I).

(n) or (15).  , $t \in R_3$. GDD 1 of Row 17.

Omitting 2 in whole is an arithmetic root system when $t = s$, or $t = s^{-1}$, or $t = -s$, or $s = -1$. Whole is affine by Lemma 3.16.

(n') or (16).  , $q \in R_3$. GDD 1 of Row 17.


Omitting 2 in whole is an arithmetic root system when $s \in R_2 \cup R_3 \cup R_4 \cup R_6$. Whole is affine by Lemma 3.16

(o) or (17).  , $q \in R_3$. GDD 4 of Row 17.

Omitting 2 in whole is an arithmetic root system when $q = -s^{-1}$ or $s = -1$ by Lemma 3.2 (XI). Whole is affine by Lemma 3.16.



Omitting 2 in whole is an arithmetic root system when $q = -s$ or $q = -s^{-1}$ or $s = -1$ by Lemma 3.2 (XI). Whole is affine by Lemma 3.16.



Omitting 2 in whole is an arithmetic root system when $q = s^2$ or $q = s^3$ by Lemma 3.2 (X). Omitting 1 in whole is an arithmetic root system when $r = -q^{-1}$ by Type 4. Whole is affine.



Omitting 2 in whole is an arithmetic root system when $s = -q$ by Type 4. Whole is affine by Lemma 3.4.

Case (ii) Omitting 3 and adding 2 as Case (i)

Case (iii) Omitting 1 and adding 3 as Case (i)


Case (iv) containing circle with four vertexes.

This case can be ignored since it is over a cycle.

Case (v) complete diagram containing circle with four vertexes.

According to Lemma 3.40 we have to consider following cases.

(b) in Case (i)



If 134 is an arithmetic root system then considering Lemma 3.2(I) we have $a_1, \tilde{q}_{34} =$

q^{-1} , $s = q^2$ or a_2 . $\tilde{q}_{34} = -1$. $s = -q$, $q \in R_3$, by Lemma 3.2(I)(h). or a_3 . $\tilde{q}_{34} = q^{-2}$, $s = q^3$ by Lemma 3.2(I)(m).


a_4 . $\tilde{q}_{34} = -q$. $s = -1$, $q \in R_3$.

For a_3 , if 234 is an arithmetic root system, then $r = q^3$ by Lemma 3.2(I)(m). empty.

For a_1 , if 234 is an arithmetic root system, then $r = q^2$. empty. For a_2 . if 234 is an arithmetic root system, then $r = -q$. empty.

For a_4 , $r = -1$, empty.


(d) in Case (i)



with $q^2, q^3 \neq 1$.

If 234 is an arithmetic root system, then $\tilde{q}_{34} = q^{-1}$, $q^3 = r^3$ by Lemma 3.2(I). empty.

(f) in Case (i) or (21)



a_2 . 134 is GDD 3 of Row 9. then $\tilde{q}_{34} = s^{-1}(s')^{-1}$ and $s \neq (s')^{-1}$.

a_{22} . If 234 is by Type 6. then $r' = r$, $s' = s$. $\tilde{q}_{34} = r^{-2} = s^{-2}$, $r = -s$. empty.

a_{23} . 234 is GDD 3 of Row 9. then $s^{-1}(s')^{-1}r(r') = 1$, i.e. $(r')^2 = s^2$, $(s')^2 = r^2$. $r' = \pm s$, $s' = \pm r$.

a_{231} . $r' = s, s' = r$. $\tilde{q}_{34} = q$, affine. a_{232} . $r' = s, s' = -r$. empty since $r's'q = s(-r)q = -1$,

a_{233} . $r' = -s, s' = r$. empty.

a_{234} . $r' = -s, s' = -r$. Whole is affine when $\tilde{q}_{34} = -q$.

a_3 . 134 is Type 6.

a_{31} . $s' = s$, then $\tilde{q}_{34} = s^{-2}$. Then $r = r'$. If 234 is an arithmetic root system, then $s^{-2} = r^{-2}$, $s = -r$. affine.

a_{32} . $\tilde{q}_{34} = s$. $s' = s^{-2}$,

a_{322} . 234 is GDD 3 of Row 9, $r' = q$, empty.

a_{323} . if 234 is Type 6, then $r' = r$, $s = r^{-2}$. empty by Lemma 3.2(XVIII).

a_{33} . $\tilde{q}_{34} = s'$. $s = (s')^{-2}$,

a_{332} . 234 is GDD 3 of Row 9, then $s' = (r')^{-1}r^{-1}$, i.e. $r = (s')^{-1}(r')^{-1} = q$, which is a contradiction.


a_{333} . if 234 is Type 6. a_{3331} . $r = r'$, $s' = r^{-2}$, $rs = r's'$. $rs = r^{-2}r$, empty by Lemma 3.2(XVIII).

a_{3332} . If $r = s'$, $r' = r^{-2}$. empty by Lemma 3.2(XVIII).

a_{3333} . $r' = s'$, $r = (r')^{-2}$. $s = (r')^{-2} = r$. empty.

Whole is affine when a_{231} . $\tilde{q}_{34} = r^{-1}s^{-1}$. $s = r'$. $s' = r$. a_{234} . $\tilde{q}_{34} = -r^{-1}s^{-1}$. $r' = -s$, $s' = -r$. a_{31} . $s' = s$, $r' = r$. $\tilde{q}_{34} = s^{-2}$. $s = -r$.


(g) in Case (i) or (22)



with $q \notin R_3$. If 134 is an arithmetic^s root system, then $\tilde{q}_{34} = r$ by Lemma 3.2(XVIII).

Whole is affine when $\tilde{q}_{34} = r$. $q = -r$. $q^3, q^2 \neq 1$,

(h) in Case (i) or (23)



with $q^3, q^2 \neq 1$.

a_1 . If 134 is GDD 3 of Row 9, then $\tilde{q}_{34} = q^{-1}s^2$.

a_{11} . If 324 is GDD 3 of Row 9, then $q^{-1}s^2 = q^{-1}r^2$. i.e. $s^2 = r^2$. which is a contradiction.

a_{12} . If 324 is by Type 6, then

a_{121} . $q = r^{-2}$ and $\tilde{q}_{34} = q^{-2} = q^{-1}s^2$, i.e. $s^{-2} = r^{-2}$, which is a contradiction.

a_{122} . $\tilde{q}_{34} = q$, i.e. $q = q^{-1}s^2$, and $r^{-2} = q^{-2}$, which is a contradiction.

a_{122} . $\tilde{q}_{34} = q$, i.e. $q = q^{-1}s^2$, and $r^{-2} = q^{-2}$, which is a contradiction.

a_{123} . $\tilde{q}_{34} = r^{-2} = q^{-1}s^2$, and $r^4 = q$. $q = r^2s^2 = q^{-2}$, i.e. $1 = q^3$, which is a contradiction.

a_2 . If 134 is Type 6.

a_{21} . $q^{-2} = \tilde{q}_{34}$ and $s^{-2} = q$. 234 is an arithmetic root system by GDD 3 of Row 9. a_{22} . If $q = \tilde{q}_{34}$ and $s^{-2} = q^{-2}$, which is a contradiction.

a_{23} . If $\tilde{q}_{34} = s^{-2}$, and $q = s^4$.

a_{231} . If 234 is Type 6. a_{2311} . If $q = s^{-2}$ and $r^{-2} = q^{-2}$ empty.


a_{2312} . If $q = r^{-2}$ and $s^{-2} = q^{-2}$, empty.

a_{2313} . If $s^{-2} = r^{-2}$. empty.

a_{232} . If 234 is GDD 3 of Row 9, then $\tilde{q}_{34} = q^{-1}r^2 = s^{-2}$, i.e. $q = r^2s^2$. On other hand, $q^{-2} = r^2s^2$. Thus $q = q^{-2}$, which is a contradiction.

Whole is affine when $q^{-2} = \tilde{q}_{34}$ and $s^{-2} = q$.

(i) in Case (i)



with $q \in R_3$.

a_1 . 134 is Type 6.


a_{11} . $\tilde{q}_{34} = q$ and $s = q^2$. If 234 is Type 6, then $r = q^{-2}$, which is a contradiction.

a_{12} . $\tilde{q}_{34} = s$, and $s^{-2} = q$. empty.

Thus 234 has to be GDD 3 of Row 9.


a_2 . If 134 is GDD 3 of Row 9, then $\tilde{q}_{34} = r$, 234 is Type 6. $r^{-2} = q$, empty.

(m) in Case (i)



with $q \in R_3$. 234 is not an arithmetic root system by Lemma 3.2(I).

(q) in Case (i)



with $q \in R_3$. If 234 is an arithmetic root system, then 234 has to be Lemma 3.2(I)(g), i.e. $\tilde{q}_{34} = q^{-1}$. $r = -1$. 134 is Type 6. $s = q^2 = q^{-1}$, empty.

4.23 GDD 4 of Row 9

GDD 4 of Row 9 as GDD 1 of Row 9

4.24 GDD 1 of Row 10

Case (i) Omitting 3 and adding 1.

According to Lemma 3.3 (III) we have to consider following cases.

(a) or (1).

$$\begin{array}{ccccccc} q^2 & & & & & & \\ q^{-2} & q & & q^{-1}-1 & q^{-1} & q & \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$
, $q \in F^* \setminus \{1, -1\}$, GDD 1 of Row 6.
is an arithmetic root system when $q \in R_4$ by Lemma 3.7(a). Whole is affine when $q \notin R_4 \cup R_2 \cup R_3$ by Lemma 3.7.

(c) or (2).

$$\begin{array}{ccccccc} q^3 & & & & & & \\ q^{-3} & q & & q^{-1}-1 & q^{-1} & q & \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$
, $q \in F^* \setminus \{1, -1\}$, $q \notin R_3$. GDD 1 of Row 7.
is affine $q \notin R_2 \cup R_3$ by Lemma 3.7.

(e) or (3).

$$\begin{array}{ccccccc} -1 & & & & & & \\ q^{-1} & q & & q^{-1}-1 & q^{-1} & q & \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$
, $q \in F^* \setminus \{1, -1\}$. GDD 3 of Row 8.
is an arithmetic root system when $q \in R_4$ by GDD 3 of Row 22; when $q \in R_3$ by GDD 3 of Row 12. Otherwise whole is affine by Lemma 3.28.

(g) or (4).

$$\begin{array}{ccccccc} q & & & & & & \\ -q & -q & -q & -1 & -q & -q^{-1} & \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$
, $q \in R_3$. GDD 1 of Row 14.
is affine by Lemma 3.4.

(h).

$$\begin{array}{ccccccc} -1 & & & & & & \\ q & q & & q^{-1}-1 & q^{-1} & q & \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$
, $q \in R_3$. GDD 1 of Row 15.
is empty.

(j).

$$\begin{array}{ccccccc} -q & & & & & & \\ -q^{-1} & q & & q^{-1}-1 & q^{-1} & q & \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$
, $q \in R_3$. GDD 1 of Row 16.
is empty.

(l).


$$\begin{array}{ccccccc} -1 & & & & & & \\ -1 & q & & q^{-1}-1 & q^{-1} & q & \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$
, $q \in R_3$. GDD 2 of Row 17.

empty

(n) or (5).


$$\begin{array}{ccccccc} -1 & & & & & & \\ q & -q & -q^{-1}-1 & -q^{-1}-q & & & \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$
, $q \in R_3$. GDD 5 of Row 17.

Whole is affine by Lemma 3.7




, $q \in R_3$. GDD 7 of Row 17.

empty




GDD 1 of Row 4.

arithmetic root system when $q \in R_3$ by GDD 1 of Row 13; when $q \in R_6$ by GDD 6 of Row 17; when $q \in R_{10}$ by GDD 5 of Row 9 or by Lemma 3.28(d); when $q \in R_4$ by Lemma 3.28(h). Otherwise whole is affine when $q \notin R_{10} \cup R_4 \cup R_3 \cup R_6$ by Lemma 3.28.




$q \notin R_4 \cup R_2$, GDD 1 of Row 5.

Whole is affine when $q \notin R_4$ by Lemma 3.4.



GDD 1 of Row 17.

empty




GDD 9 of Row 17.

empty

Case (ii) Omitting 3 and adding 2.


According to Lemma 3.3 (I) we have to consider following cases.

Omitting 1 in whole is an arithmetic root system in this case by symmetry.




GDD 2 of Row 4.

Omitting 1 in whole is an arithmetic root system by symmetry of this GDD. It is similar in following of Case (ii) . whole is an arithmetic root system by GDD 3 of Row 13 or by Type 5.




GDD 3 of Row 5.

Whole is affine by Lemma 3.4.




GDD 2 of Row 6.

Whole is affine by Lemma 3.28.



GDD 2 of Row 7.

Whole is affine by Lemma 3.28.




Whole is affine $q \notin R_2 \cup R_3$ by Lemma 3.4.



is an arithmetic root system by Type 5.




Whole is affine by Lemma 3.28.



Whole is affine by Lemma 3.28.




Whole is affine by Lemma 3.28.




, $q \notin R_3$. GDD 2 of Row 10.

Whole is affine by Lemma 3.28.




, $q \in R_3$. GDD 1 of Row 11.




, $q \in R_3$. GDD 3 of Row 14.

Whole is affine by Lemma 3.4.




, $q \in R_3$. GDD 2 of Row 16.

Whole is affine by Lemma 3.28.



, $q \in R_3$. GDD 4 of Row 16.

Whole is affine by Lemma 3.4.




, $q \in R_3$. GDD 1 of Row 17.

Case (iii) Omitting 1 and adding 3 as Case (i)

Case (iv) Containing circle with four vertexes.

According to Lemma 3.39 (ii) we have to consider following cases.


(a) to (a)



is not an arithmetic root system by Lemma 3.2 (I).


(a) to (c) empty.

(a) to (e)



with $q^2 = -1$ is not an arithmetic root system by Lemma 3.2 (I).


(a) to (g)



with $-q^{-1} = q^2$ is not an arithmetic root system by Lemma 3.2 (I).


(a) to (n) empty.

(a) to (a') empty.(a) to (a'')




with $q = q^4$ is not an arithmetic root system by Lemma 3.2(I)

(c) to (c)



is not an arithmetic root system by Lemma 3.2 (I).


(c) to (e)



with $-1 = q^3$ is not an arithmetic root system by Lemma 3.2 (I).

(c) to (g) empty.


(c) to (n)



with $q^3 = -1$ is not an arithmetic root system by Lemma 3.2 (I).


(c) to (a') empty.

(c) to (a'')




with $q = q^6$ is not an arithmetic root system by Lemma 3.2 (I).

(e) to (e) or (19)




is an arithmetic root system by Lemma 3.2 (IV).



is affine. (e) to (g) empty.


(e) to (n)



with $q \in R_6$ is not an arithmetic root system by Lemma 3.2 (I).

(e) to (a') empty.(e) to (a'')

(g) to (g)




with $q \in R_6$ is not an arithmetic root system by Lemma 3.2 (V).

(g) to (n) empty.


(g) to (a') empty.

(g) to (a'')



with $q = -q^{-2}$ is not an arithmetic root system by Lemma 3.2 (V).


(n) to (n)



with $q \in R_6$ is not an arithmetic root system by Lemma 3.2 (I).

(n) to (a') empty.

(n) to (a'') empty (a') to (a') or (20)




is an arithmetic root system.

is affine.

(a') to (a'') empty.

(a'') to (a'')




is not an arithmetic root system by Lemma 3.2 (V).

4.25 GDD 2 of Row 10


Case (i) Omitting 3 and adding 1.

According to Lemma 3.3 (III) we have to consider following cases.




, $q \in F^* \setminus \{1, -1\}$. GDD 1 of Row 6.

Whole is affine by Lemma 3.7




, $q \in F^* \setminus \{1, -1\}$, $q \notin R_3$. GDD 1 of Row 7.

is affine by Lemma 3.7




, $q \in F^* \setminus \{1, -1\}$. GDD 3 of Row 8.

is arithmetic root system by Type 1



, $q \in R_3$. GDD 1 of Row 14.

Whole is affine by Lemma 3.4



, $q \in R_3$. GDD 1 of Row 15.

is empty.

$$\begin{array}{c} \text{(j).} \\ -q \quad -q^{-1}q \quad q^{-1}-1 \quad q^2 \quad q^{-2} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array}$$

, $q \in R_3$. GDD 1 of Row 16.

is empty.

$$\begin{array}{c} \text{(l).} \\ -1 \quad -1 \quad q \quad q^{-1}-1 \quad q^2 \quad q^{-2} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array}$$

, $q \in R_3$. GDD 2 of Row 17.

is empty.

$$\begin{array}{c} \text{(n) or (4).} \\ -1 \quad q \quad -q \quad -q^{-1}-1 \quad q^2 \quad q^{-2} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array}$$

, $q \in R_3$. GDD 5 of Row 17.

Whole is affine by Lemma 3.7.

$$\begin{array}{c} \text{(o).} \\ -1 \quad -q \quad q \quad q^{-1}-1 \quad q^2 \quad q^{-2} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array}$$

, $q \in R_3$. GDD 7 of Row 17.

empty.

$$\begin{array}{c} \text{(a').} \\ q \quad q^{-1} \quad q \quad q^{-1}-1 \quad q^2 \quad q^{-2} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array}$$

GDD 1 of Row 4.

is an arithmetic root system by Type 1.

$$\begin{array}{c} \text{(a'') or (5)} \\ q \quad q^{-2} \quad q^2 \quad q^{-2}-1 \quad q^4 \quad q^{-4} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array}$$

$q^4 \neq 1$, GDD 1 of Row 5.

Whole is affine by Lemma 3.4.

$$\begin{array}{c} \text{(k')} \\ -1 \quad q \quad -1 \quad -1 \quad -1 \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array}$$

empty.

$$\begin{array}{c} \text{(p')} \\ q^{-1} \quad -q^{-1}-1 \quad -1 \quad -1 \\ \bullet \quad \bullet \quad \bullet \end{array}$$

empty.

Case (ii) Omitting 3 and adding 2.

According to Lemma 3.3 (I) we have to consider following cases.

$$\begin{array}{c} \text{(a') or (6)} \\ \begin{array}{c} q^{-2} \\ \bullet \\ q^2 \\ \bullet \\ -1 \quad q \quad -1 \quad q^{-1}q \end{array} \end{array}$$

GDD 2 of Row 4.


Omitting 1 in whole is an arithmetic root system when $q \in R_4 \cup R_5$ by Lemma 3.2 (X). Whole is affine by Lemma 3.12.

$$\begin{array}{c} \text{(a) or (7)} \\ \begin{array}{c} q^{-4} \\ \bullet \\ q^4 \\ \bullet \\ -q^{-2}q^2 \quad -1 \quad q^{-2}q^2 \end{array} \end{array}$$

$q^2 \notin R_2 \cup R_3$, GDD 3 of Row 5.

Omitting 1 in whole is an arithmetic root system when $q \in R_6$ by Type 3; when $q \in R_7$ by Lemma 3.2(I)(h). Whole is affine when $q \in R_7$ by Lemma 3.4.


(b) or (8)



GDD 2 of Row 6.

Omitting 1 in whole is an arithmetic root system when $q \in R_5 \cup R_6 \cup R_7$ by Lemma 3.2 (X). Whole is affine by Lemma 3.12.


(d) or (9)



$q^3 \notin R_2 \cup R_3$, GDD 2 of Row 7.

Omitting 1 in whole is an arithmetic root system when $q \in R_7 \cup R_8$ by Lemma 3.2 (X). Whole is affine by Lemma 3.12.


(f) or (10)



$q^3 \notin R_2 \cup R_3$, GDD 4 of Row 7.

Omitting 1 in whole is an arithmetic root system when $q \in R_8$ by Type 3. Omitting 1 in whole is an arithmetic root system when $q^{-6} = q^3$ by Lemma 3.2(I)(h). Whole is affine by Lemma 3.4.


(g) or (11)



, GDD 1 of Row 8.

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2 (IV). Whole is affine by Lemma 3.8.


(h) or (12)



, GDD 1 of Row 9.

Omitting 1 in whole is an arithmetic root system by Lemma 3.2 (IV). Whole is affine by Lemma 3.8.

(i') or (13)




$q^2 \notin R_3 \cup R_2$. GDD 2 of Row 10.

Omitting 1 in whole is an arithmetic root system by Lemma 3.2 (IV). Whole is affine by Lemma 3.8 when $q^4 \neq -1$. Whole is affine by Lemma 3.25 when $q^4 = -1$.

(k) or (14)  , $q \notin R_3$. GDD 1 of Row 10.

Omitting 1 in whole is an arithmetic root system by Lemma 3.2 (IV). Whole is affine by Lemma 3.8.

(l) or (15)  , $q \notin R_3$. GDD 2 of Row 10.


Omitting 1 in whole is an arithmetic root system by Lemma 3.2 (IV). Whole is affine by Lemma 3.8.

(m)  , $q \in R_3$. GDD 1 of Row 11.

is empty

(n)  , $q \in R_3$. GDD 3 of Row 14.

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(I).

(p) or (16)  , $q \in R_3$. GDD 2 of Row 16.

Omitting 1 in whole is an arithmetic root system by Lemma 3.2 (X). Whole is affine by Lemma 3.12.

(r)  , $q \in R_3$. GDD 4 of Row 16.

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(I).

(s')  , $q \in R_3$. GDD 1 of Row 17.

is empty.

Case (iii) Omitting 1 and adding 3.

According to Lemma 3.3 (III) we have to consider following cases.

$$q^4 \begin{matrix} \text{(a).} \\ q^{-4} \end{matrix} \begin{matrix} q^2 & q^{-2}-1 & q & q^{-1} \end{matrix}$$

, $q^4 \neq 1$. GDD 1 of Row 6.

$$q^6 \begin{matrix} \text{(c).} \\ q^{-6} \end{matrix} \begin{matrix} q^2 & q^{-2}-1 & q & q^{-1} \end{matrix}$$

, $q \in F^* \setminus \{1, -1\}$, $q \notin R_3$. GDD 1 of Row 7.

$$-1 \begin{matrix} \text{(e).} \\ q^{-2} \end{matrix} \begin{matrix} q^2 & q^{-2}-1 & q & q^{-1} \end{matrix}$$

, $q \in F^* \setminus \{1, -1\}$. GDD 3 of Row 8.

$$q^2 \begin{matrix} \text{(g).} \\ -q^2 \end{matrix} \begin{matrix} -q^{-2} & -q^{-2}-1 & -iq & (iq)^{-1} \end{matrix}$$

, $q^2 \in R_3$. GDD 1 of Row 14.

$$-1 \begin{matrix} \text{(h).} \\ q^2 \end{matrix} \begin{matrix} q^2 & q^{-2}-1 & q & q^{-1} \end{matrix}$$

, $q \in R_3$. GDD 1 of Row 15.

empty

$$-q^2 \begin{matrix} \text{(j).} \\ -q^{-2} \end{matrix} \begin{matrix} q^2 & q^{-2}-1 & q & q^{-1} \end{matrix}$$

, $q \in R_3$. GDD 1 of Row 16.

empty

$$-1 \begin{matrix} \text{(l).} \\ -1 \end{matrix} \begin{matrix} q^2 & q^{-2}-1 & q & q^{-1} \end{matrix}$$

, $q \in R_3$. GDD 2 of Row 17.

empty

$$-1 \begin{matrix} \text{(n).} \\ q^2 \end{matrix} \begin{matrix} -q^2 & -q^{-2}-1 & iq & (iq)^{-1} \end{matrix}$$

, $q \in R_3$. GDD 5 of Row 17.

$$-1 \begin{matrix} \text{(o).} \\ -q^2 \end{matrix} \begin{matrix} q^2 & q^{-2}-1 & q & q^{-1} \end{matrix}$$

, $q \in R_3$. GDD 7 of Row 17.

is empty.

$$q^2 \begin{matrix} \text{(a')} \\ q^{-2} \end{matrix} \begin{matrix} q^2 & q^{-2}-1 & q & q^{-1} \end{matrix}$$

GDD 1 of Row 4.

$$q \begin{matrix} \text{(a'')} \\ q^{-2} \end{matrix} \begin{matrix} q^2 & q^{-2}-1 & q & q^{-1} \end{matrix}$$

GDD 1 of Row 5.

(k')

$$\begin{array}{c} -1 \quad q^2 \quad -1 \quad -1 \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}$$

is empty.

$$\begin{array}{c} q^{-2} \quad (p') \\ -q^{-2} \quad -1 \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}$$

is empty.

Case (iv) containing circle with four vertexes.

According to Lemma 3.39 (ii) we have to consider following cases.

(a) to (a)

$$\begin{array}{c} q^{-1} \quad q^2 \quad q^{-2} \quad q^{-4} \quad q^2 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}$$

with $q^4 = q^{-2}$ is not an arithmetic root system by Lemma 3.2(I)

(a) to (c)

$$\begin{array}{c} q^{-1} \quad q^2 \quad q^{-2} \quad q^{-6} \quad q^2 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}$$

with $q^6 = q^{-2}$ is not an arithmetic root system by Lemma 3.2(I)

(a) to (e) or (17)

$$\begin{array}{c} q^{-1} \quad q^2 \quad q^{-2} \quad q^{-2} \quad q^2 \\ \bullet \text{---} \bullet \text{---} \bullet \\ \quad \quad \quad \bullet \\ \quad \quad \quad q^{-2} \end{array}$$

with $-1 = q^{-2}$ is an arithmetic root system by Type 2.

$$\begin{array}{c} \quad \quad \quad \bullet \\ \quad \quad \quad q^{-2} \\ \quad \quad \quad \bullet \\ q^2 \quad \quad \quad q^{-2} \\ \bullet \text{---} \bullet \text{---} \bullet \\ q^{-1} \quad q \quad -1 \quad q^{-2} \quad q^2 \end{array}$$

is affine.

(a) to (g) empty.

(a) to (n) empty.

(a) to (a') or (18)

$$\begin{array}{c} q^{-1} \quad q^2 \quad q^{-2} \quad q^{-2} \quad q^2 \\ \bullet \text{---} \bullet \text{---} \bullet \\ \quad \quad \quad \bullet \\ \quad \quad \quad q^{-2} \end{array}$$

with $q^{-2} = q^2$ is an arithmetic root system by Type 2.

$$\begin{array}{c} \quad \quad \quad \bullet \\ \quad \quad \quad q^{-2} \\ \quad \quad \quad \bullet \\ q^2 \quad \quad \quad q^{-2} \\ \bullet \text{---} \bullet \text{---} \bullet \\ q^{-1} \quad q \quad -1 \quad q^{-2} \quad q^2 \end{array}$$

is affine.

(a) to (a'')

$$\begin{array}{c} q^{-1} \quad q^2 \quad q^{-2} \quad q^{-2} \quad q^2 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}$$

with $q = q^{-2}$ is not an arithmetic root system by Lemma 3.2 (I)

(c) to (a)

$$\begin{array}{c} q^{-1} \quad q^3 \quad q^{-3} \quad q^{-4} \quad q^2 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}$$

with $q^4 = q^{-3}$ is not an arithmetic root system by Lemma 3.2(I).

(c) to (c)

$$\begin{array}{c} q^{-1} \quad q^3 \quad q^{-3} \quad q^{-6} \quad q^2 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}$$

with $q^{-3} = q^6$ is not an arithmetic root system by Lemma 3.2(I).

(c) to (e)

$$\begin{array}{c} q^{-1} q^3 \quad q^{-3} q^{-2} q^2 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}$$

with $-1 = q^{-3}$ is not an arithmetic root system by Lemma 3.2 (I).

(c) to (g) empty

(c) to (n) empty

(c) to (a')

$$\begin{array}{c} q^{-1} q^3 \quad q^{-3} q^{-2} q^2 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}$$

with $q^2 = q^{-3}$ is not an arithmetic root system by Lemma 3.2 (I).

(c) to (a'') empty.

(e) to (a)

$$\begin{array}{c} q^{-1} q \quad -1 q^{-4} q^2 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}$$

with $-1 = q^4$ is not an arithmetic root system by Lemma 3.2 (I).

(e) to (c)

$$\begin{array}{c} q^{-1} q \quad -1 q^{-6} q^2 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}$$

with $-1 = q^6$ is not an arithmetic root system by Lemma 3.2 (I).

(e) to (e) or (19)

$$\begin{array}{c} q^{-1} q \quad -1 q^{-2} q^2 \\ \bullet \text{---} \bullet \text{---} \bullet \\ \quad \quad \quad \bullet \\ \quad \quad \quad -1 \end{array}$$

is an arithmetic root system by Lemma 3.2 (IV).

$$\begin{array}{c} q \quad \quad \quad q^{-2} \\ \bullet \quad \quad \bullet \\ \bullet \text{---} \bullet \text{---} \bullet \\ q^{-1} q \quad -1 q^{-2} q^2 \end{array}$$

is affine.

(e) to (g) empty. (e) to (n)

$$\begin{array}{c} (iq)^{-1} \quad -1 \quad q^2 \quad -q^2 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}$$

is not an arithmetic root system by Lemma 3.2 (I).

(e) to (a') or (20)

$$\begin{array}{c} q^{-1} q \quad -1 q^{-2} q^2 \\ \bullet \text{---} \bullet \text{---} \bullet \\ \quad \quad \quad \bullet \\ \quad \quad \quad -1 \end{array}$$

with $-1 = q^2$ is an arithmetic root system by Lemma 3.2 (IV).

$$\begin{array}{c} q \quad \quad \quad q^{-2} \\ \bullet \quad \quad \bullet \\ \bullet \text{---} \bullet \text{---} \bullet \\ q^{-1} q \quad -1 q^{-2} q^2 \end{array}$$

is affine.

(e) to (a'') empty.

(g) to (a)

$$\begin{array}{c} q^{-1} q \quad -q q^{-4} q^2 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}$$

with $q^4 = -q$ is not an arithmetic root system by Lemma 3.2(I).

(g) to (c) empty.

(g) to (e) empty. (g) to (g) empty. (g) to (n) empty. (g) to (a') empty.

(g) to (a'') empty.


(n) to (a) empty.(n) to (c) empty.

(n) to (e) or (21)

is an arithmetic root system by Type 3.

is affine.

(n) to (g) empty.(n) to (n)



is not an arithmetic root system by Lemma 3.2(I)


(n) to (a') empty.(n) to (a'') empty.

(a') to (a) or (22)

with $q^4 = q^{-1}$ is an arithmetic root system by Type 2.

is affine.

(a') to (c)



with $q^6 = q^{-1}$ is not an arithmetic root system by Lemma 3.2(I).


(a') to (e) empty.(a') to (g) empty.(a') to (n) empty.

(a') to (a') or (23)

with $q^{-1} = q^2$ is an arithmetic root system by Type 7.

is affine.


(a') to (a'') empty. We ignore the case (a'') since it is considered in previous.



4.26 GDD 3 of Row 10

Case (i) Omitting 3 and adding 1.

According to Lemma 3.3 (IV) we have to consider following cases.



, $q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 4.

Omitting 2 in whole is an arithmetic root system when $q \in R_5 \cup R_7 \cup R_6 \cup R_4$ by Lemma 3.2 (X). Whole is affine by Lemma 3.16.


(a'') or (2).  , $q^2 \in F^* \setminus \{1, -1\}$. GDD 2 of Row 5.

Omitting 2 in whole is an arithmetic root system when $q \in R_6$ by Type 2; Omitting 2 in whole is not an arithmetic root system when $q \notin R_6$ by Lemma 3.2(I). Whole is affine by Lemma 3.4.


(a) or (3).  , $q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 6.

Omitting 2 in whole is an arithmetic root system when $q \in R_4 \cup R_6 \cup R_8$ by Lemma 3.2 (XI)(e) and by Lemma 3.2 (X). Whole is affine by Lemma 3.16.


(b) or (4).

 , $q \in F^* \setminus \{1, -1\}$. GDD 3 of Row 6.

Omitting 2 in whole is an arithmetic root system when $q \in R_5 \cup R_6 \cup R_7 \cup R_4$ by Lemma 3.2 (X). Omitting 1 in whole is an arithmetic root system by Type 7. Whole is affine.

(c) or (5).  , $q \in F^* \setminus \{1, -1\}$. $q \notin R_3$, GDD 2 of Row 7.

Omitting 2 in whole is an arithmetic root system when $q \in R_5 \cup R_7 \cup R_9 \cup R_6 \cup R_4$ by Lemma 3.2 (X) and Lemma 3.3 (V). Whole is affine by Lemma 3.16.

(d). 


(, $q \in F^* \setminus \{1, -1\}$. $q \notin R_3$, $q^2 \notin R_3 \cup R_2$, GDD 3 of Row 7.)

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(I).

(e) or (6).  , $q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 8.


Omitting 2 in whole is an arithmetic root system when $q \in R_4$ by Lemma 3.3 (V)(a). Whole is affine by Lemma 3.16.

(f) or (7).




, $q, r, s \in F^* \setminus \{1\}$, $qrs = 1$, $q \neq r$, $q \neq s$, $s \neq r$. GDD 3 of Row 9.

Omitting 2 in whole is an arithmetic root system when $s = q^2$ or $s = q^{-2}$, $q \in R_3 \cup R_6$ or $s = -q^{-2}$, $s \in R_3 \cup R_6$ or $s = -1$, $q^{-2} \in R_2 \cup R_3 \cup R_4 \cup R_6$ or $q^{-2} = -1$, $s \in R_2 \cup R_3 \cup R_4 \cup R_6$ by Lemma 3.2 (XI). Omitting 1 in whole is an arithmetic root system when $r = -q$, $q \in R_6$, or $r = -1$, $q \in R_4 \cup R_6$ by Lemma 3.2 (XI). Whole is affine.

(g) or (8). 


, $q \in F^* \setminus \{1, -1\}$, $q^3 \neq 1$. GDD 3 of Row 10.

Omitting 2 in whole is an arithmetic root system when $q \in R_6 \cup R_4$ by Lemma 3.2 (XI). Omitting 1 in whole is an arithmetic root system when $q \in R_4 \cup R_6$ by Lemma 3.2 (XI). Whole is affine.

(h) or (9).  , $q^2, q^3 \neq 1$. GDD 3 of Row 10.

Omitting 2 in whole is an arithmetic root system when $q \in R_5 \cup R_6$ by Lemma 3.2 (XI). Omitting 1 in whole is an arithmetic root system when $q \in R_4 \cup R_6$ by Lemma 3.2 (XI). Whole is affine.

(i).



is empty.



Omitting 2 in whole is not an arithmetic root system by Lemma 3.2(I).




is empty.



is empty.

(m).




Omitting 2 in whole is not an arithmetic root system by Lemma 3.2 (XI).



is empty.




is empty.




, $q \in R_3$. GDD 4 of Row 17.

is empty.




, $q \in R_3$. GDD 4 of Row 17.

Omitting 2 in whole is an arithmetic root system by Lemma 3.2 (XI), Whole is affine by Lemma 3.16.



, $q \in R_3$. GDD 8 of Row 17.

is empty.




, $q \in R_3$. GDD 9 of Row 17.

is empty.


Case (ii) Omitting 3 and adding 2.

According to Lemma 3.3 (IV) we have to consider following cases.




, $q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 4.

Omitting 1 in whole is an arithmetic root system since symmetry of this GDD. It is similar in following of Case (ii) . Whole is an arithmetic root system by Type 6.



, $q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 5.


Whole is affine by Lemma 3.4



, $q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 6.


is affine by Lemma 3.16.

(b) or (13).




, $q \in F^* \setminus \{1, -1\}$. GDD 3 of Row 6.

Omitting 1 in whole is an arithmetic root system by Type 7. Omitting 2 in whole is an arithmetic root system when $q \in R_5 \cup R_7 \cup R_6 \cup R_4$ by Lemma 3.2 (X) (X) (e). Whole is affine.

(c) or (14).  , $q \in F^* \setminus \{1, -1\}$. $q \notin R_3$, GDD 2 of Row 7.

Whole is affine by Lemma 3.16.

(d) or (15).




$q^6, q^9 \neq 1$, GDD 3 of Row 7.

Omitting 1 in whole is an arithmetic root system when $q \in R_5 \cup R_8$ by Lemma 3.2 (X). Omitting 2 in whole is an arithmetic root system when $q \in R_8 \cup R_4$ by Type 2. Whole is affine.

(e).  , $q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 8.

whole is an arithmetic root system by Type 6.

(f) or (16).




, $q, r, s \in F^* \setminus \{1\}$, $qrs = 1$, $q \neq r$, $q \neq s$, $s \neq r$. GDD 3 of Row 9.

Omitting 1 in whole is an arithmetic root system when $s = -q$, $q \in R_6$ or $-1 = s$, $q \in R_4 \cup R_6$ by Lemma 3.2 (XI). Omitting 2 in whole is an arithmetic root system when $r = q^2$ or $r = -q^{-2}$ $q \in R_6$ or $-1 = q^{-2}$, $r \in R_2 \cup R_3 \cup R_4 \cup R_6$ or $-1 = r$,


$q^{-2} \in R_2 \cup R_3 \cup R_4 \cup R_6$. Whole is affine.

(g).




Omitting 1 in whole is not an arithmetic root system by Lemma 3.2 (XI).

(h) or (17).



Omitting 1 in whole is an arithmetic root system by Lemma 3.2 (XI) when $q \in R_6 \cup R_4$.
 Omitting 2 in whole is an arithmetic root system when $q \in R_6 \cup R_5$. Whole is affine when $q \in R_6$.

(i).



is empty.




Whole is affine by Lemma 3.4.




is empty.

(l).




, $q \in R_3$. GDD 2 of Row 16.

is empty.
(m).




, $q \in R_3$. GDD 3 of Row 16.

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2 (X).




, $q \in R_3$. GDD 1 of Row 17.

is empty.




, $q \in R_3$. GDD 1 of Row 17.

is empty.




, $q \in R_3$. GDD 4 of Row 17.

is empty




, $q \in R_3$. GDD 4 of Row 17.

Whole is affine by Lemma 3.16.



, $q \in R_3$. GDD 8 of Row 17.

is empty.



, $q \in R_3$. GDD 9 of Row 17.

is empty.

Case (iii) Omitting 1 and adding 3 is the same as Case (i) .


Case (iv) containing circle with four vertexes.

This case can be ignored since it is over a cycle.

Case (v) complete diagram containing circle with four vertexes.

According to Lemma 3.40 we have to consider following cases.


(b) in Case (i)



with $q^2, q^3 \neq 1$,


If 324 is Type 6, then $q = q^2$, which is a contradiction. If 324 is GDD 3 of Row 7, and $q = q^3$, which is a contradiction.

(d) in Case (i) or (20)



If 134 is an arithmetic root system, then a_1 . $\tilde{q}_{34} = q^{-1}$ and $q^{-6} = q^2$ by Type 6 or a_2 . $\tilde{q}_{34} = q^{-2}$ and $q^{-6} = q^3$ by Lemma 3.2(I)(m). For a_1 , 234 is an arithmetic root system by Lemma 3.2(I)(m), For a_2 , 234 is not an arithmetic root system by Lemma 3.2(I). Whole is affine when $\tilde{q}_{34} = q^{-1}$ and $q^8 = 1$.

(f) in Case (i) or (21)



a_1 , assume 314 is Type 6.

a_{11} . $\tilde{q}_{34} = q^{-2}$ and $s = q^4$.

a_{111} . Assume 324 is Type 6, then $r = q^{-2}$ and $q = r^{-2}$. Consequently $q = q^4$, empty.

a_{112} . 234 is GDD 3 of Row 9. then $r = q$, empty.

a_{12} . $\tilde{q}_{34} = s$. $q^{-2} = s^{-2}$. $q = -s$. 234 is an arith by GDD 3 of Row 9.


a_2 . 134 is GDD 3 of Row 9. $\tilde{q}_{34} = q^2 s^{-1}$.

a_{21} . 234 is GDD 3 of Row 9, then $q^2 s^{-1} = r^{-1} q^{-1}$, i.e, $q^3 = s r^{-1}$.

a_{22} . 234 is Type 6, then $q^2s^{-1} = r$ and $q = r^{-2}$.

Whole is affine when $\tilde{q}_{34} = s$. $q = -s$. or $\tilde{q}_{34} = q^2s^{-1}$. $q^3 = sr^{-1}$. or $\tilde{q}_{34} = q^2s^{-1}$.
 $q^2s^{-1} = r$. $q = r^{-2}$ with $, q, r, s \in F^* \setminus \{1\}, qrs = 1, q \neq r, q \neq s, s \neq r..$

(g) in Case (i) or (22)




a_1 . 134 is Type 6.

a_{11} . $\tilde{q}_{34} = q$. 234 is Type 6.

a_{12} . $\tilde{q}_{34} = q^{-2}$ and $q = q^4$, empty.

a_2 . 134 is GDD 3 of Row 9, $\tilde{q}_{34} = q^{-1}q^2$, empty. Whole is affine when $\tilde{q}_{34} = q$ with $q^2, q^3 \neq 1$.

(h) in Case(i) or (23)



with $q^2, q^3 \neq 1$.

a_1 . If 314 is Type 6. a_{11} . $\tilde{q}_{34} = q$ and $q^4 = q^{-2}$. 324 is Type 6.


a_2 . If 134 is GDD 3 of Row 9, then $\tilde{q}_{34} = q^{-5}$.

a_{21} . If 324 is Type 6, then $q = q^{-5}$, empty; or $q^{-5} = q^{-2}$, empty.

a_{22} . If 324 is GDD 3 of Row 9, then $\tilde{q}_{34} = q$, i.e. $q = q^4$, which is a contradiction.

Whole is affine when $\tilde{q}_{34} = q$ and $q \in R_6$.

(m) in Case(i)




with $q^2, q^3 \neq 1$.

If 324 is an arithmetic root system, then $\tilde{q}_{34} = q^{-1}$ by Lemma 3.2(I)(k). 314 is not an


arithmetic root system checked step by step.

(b) in Case (ii) or (24)




If 234 is an arithmetic root system, then it is Type 6 and $\tilde{q}_{34} = q^{-1}$, as well as $q = q^2$, empty; or $\tilde{q}_{34} = q^{-2}$ and $q = q^3$, empty.

(d) in Case (ii)



If 134 is an arithmetic root system, then a_1 . $\tilde{q}_{34} = q^{-1}$ and $q^{-6} = q^3$ by GDD 3 of Row 7, or a_2 . $\tilde{q}_{34} = q^{-2}$, $q^{-2} = q^{-1}$ and $q^{-6} = q^3$, empty. For a_1 , if 234 is an arithmetic root system, then $q^3 = q^2$ by Type 6, empty,

(f) in Case (ii) or (25)



a_1 . If 234 is GDD 3 of Row 9, then $\tilde{q}_{34} = r$.

a_{11} . 134 is Type 6. then $q^{-2} = r^{-2}$, i.e. $q = -r$.

a_2 . If 234 is Type 6.

a_{21} . $\tilde{q}_{34} = s$ and $q = s^{-2}$.

a_{211} . If 134 is Type 6, then $s = q^{-2}$ and $r = s^{-2}$, empty.


a_{212} . If 134 is GDD 3 of Row 9, then $rsq^{-2} = 1$, i.e. $sr = q^2$. $sr = s^{-4}$. $r = s^{-5}$.

a_{22} . $\tilde{q}_{34} = q$, $q^{-2} = s$. a_{221} . If 134 is Type 6, empty.

a_{222} . If 134 is GDD 3 of Row 9, then $rq^{-1} = 1$, empty.


Whole is affine when $\tilde{q}_{34} = r$ and $q = -r$, or $\tilde{q}_{34} = s$ and $q = s^{-2}$. $rsq^{-2} = 1$.

(g) in Case (ii)



If 234 is Type 6, then $\tilde{q}_{34} = q^{-2}$. 134 is not an arithmetic root system checked step by step.

(h) in Case (ii) or (26)




a_1 . If 234 is Type 6, then $\tilde{q}_{34} = q$.

a_{11} . $\tilde{q}_{34} = q$. Then 134 is Type 6 and $q^4 = q^{-2}$, i.e. $q^6 = 1$. Whole is affine when $\tilde{q}_{34} = q$, $q \in R_6$.

If 234 is by GDD 3 of Row 9, then $\tilde{q}_{34} = q$, empty.

(m) in Case (ii).



134 is not an arithmetic root system by Lemma 3.2(I).

4.27 GDD 1 of Row 11

Case (i) Omitting 3 and adding 1.

According to Lemma 3.3 (III) we have to consider following cases.

(a) or (1).

$$\begin{array}{c} q^2 \quad q^{-2} \quad q \quad q^{-1} - 1 \quad q^{-1} \quad q \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad , q \in F^* \setminus \{1, -1\}. \text{ GDD 1 of Row 6.}$$

Whole is an arithmetic root system when $q \in R_4$ by Lemma 3.7 (a) Whole is affine when $q \notin R_4$ by Lemma 3.7.

$$\begin{array}{c} q^3 \quad (c). \\ q^{-3} \quad q \quad q^{-1} - 1 \quad q^{-1} \quad q \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad , q \in F^* \setminus \{1, -1\} , q \notin R_3. \text{ GDD 1 of Row 7.}$$

is empty.

$$\begin{array}{c} (e). \\ -1 \quad q^{-1} \quad q \quad q^{-1} - 1 \quad q^{-1} \quad q \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad , q \in F^* \setminus \{1, -1\}. \text{ GDD 3 of Row 8.}$$

is an arithmetic root system by Lemma 3.28 (f).

$$\begin{array}{c} (g). \\ q \quad -q \quad -q^{-1} \quad -1 \quad -q \quad -q^{-1} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad , q \in R_3. \text{ GDD 1 of Row 14.}$$

is empty .

$$\begin{array}{c} (h) \text{ or } (2). \\ -1 \quad q \quad q \quad q^{-1} - 1 \quad q^{-1} \quad q \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad , q \in R_3. \text{ GDD 1 of Row 15.}$$

Whole is affine by Lemma 3.28.

$$\begin{array}{c} (j) \text{ or } (3). \\ -q \quad -q^{-1} \quad q \quad q^{-1} - 1 \quad q^{-1} \quad q \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad , q \in R_3. \text{ GDD 1 of Row 16.}$$

Whole is affine by Lemma 3.28.

$$\begin{array}{c} (l) \text{ or } (4). \\ -1 \quad -1 \quad q \quad q^{-1} - 1 \quad q^{-1} \quad q \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad , q \in R_3. \text{ GDD 2 of Row 17.}$$

Whole is affine by Lemma 3.28.

$$\begin{array}{c} (n). \\ -1 \quad q \quad -q \quad -q^{-1} - 1 \quad -q^{-1} - q \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad , q \in R_3. \text{ GDD 5 of Row 17.}$$

is empty .

$$\begin{array}{c} (o) \text{ or } (5). \\ -1 \quad -q \quad q \quad q^{-1} - 1 \quad q^{-1} \quad q \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad , q \in R_3. \text{ GDD 7 of Row 17.}$$

Whole is affine by Lemma 3.7.

(a').

$$\begin{array}{ccccccc} q & q^{-1} & q & q^{-1} & -1 & q^{-1} & q \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$
 $q^2 \neq 1$, GDD 1 of Row 4.
 is an arithmetic root system by Lemma 3.28(g).

$$\begin{array}{ccccccc} (a'') & q^2 & (6) \\ q & q^{-2} & q^2 & q^{-2} & -1 & q^{-2} & q^2 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$
 $q^4 \neq 1$, GDD 1 of Row 5.

Whole is affine by Lemma 3.28.

$$\begin{array}{ccccccc} (k') \\ -1 & q & -1 & -1 & -1 & -1 & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$
 GDD 1 of Row 17.

is empty.

$$\begin{array}{ccccccc} (p') \\ q^{-1} & -q^{-1} & -1 & -1 & -1 & -1 & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$
 GDD 9 of Row 17.

is empty.

Case (ii) Omitting 3 and adding 2.

According to Lemma 3.3 (I) we have to consider following cases.

$$\begin{array}{c} (a') \\ q \\ q^{-1} \\ \bullet \\ -1 \quad q \quad -1 \quad q^{-1} \quad q \end{array}$$
 GDD 2 of Row 4.

Omitting 1 in whole is an arithmetic root system since symmetry of this GDD. It is similar in following of Case (ii) . Whole is arithmetic root system by GDD 3 of Row 13 or Type 5.

$$(a) \text{ or } (6) \quad \begin{array}{c} q^2 \\ q^{-2} \\ \bullet \\ -q \quad q^2 \quad -1 \quad q^{-2} \quad q^2 \end{array}$$
 GDD 3 of Row 5.


Whole is affine by Lemma 3.28 or by Lemma 3.4.

$$(b) \quad \begin{array}{c} q^2 \\ q^{-2} \\ \bullet \\ -1 \quad q \quad -1 \quad q^{-2} \quad q^2 \end{array}$$
 GDD 2 of Row 6.

is an arithmetic root system by Lemma 3.28.


$$(d) \quad \begin{array}{c} q^3 \\ q^{-3} \\ \bullet \\ -1 \quad q \quad -1 \quad q^{-3} \quad q^3 \end{array}$$
 GDD 2 of Row 7.

is empty.




$q \notin R_3$, GDD 4 of Row 7.

Whole is affine by Lemma 3.28 with $q^3 \in R_3$.




, GDD 1 of Row 8.

is an arithmetic root system by GDD 6 of Row 12 or Type 5.




, GDD 1 of Row 9.

Whole is affine by Lemma 3.30 or Lemma 3.28,




, $q \notin R_3$. GDD 2 of Row 10.

Whole is affine Lemma 3.28 with $q^2 \in R_3$.




, $q \notin R_3$. GDD 1 of Row 10.

is empty.




, $q \notin R_3$. GDD 2 of Row 10.

is empty.




, $q \in R_3$. GDD 1 of Row 11.

Whole is affine by Lemma 3.30 or Lemma 3.28,




, $q \in R_3$. GDD 3 of Row 14.

Whole is affine by Lemma 3.30 or by Lemma 3.4.




, $q \in R_3$. GDD 2 of Row 16.

Whole is affine by Lemma 3.30 or Lemma 3.28.



, $q \in R_3$. GDD 4 of Row 16.

Whole is affine by Lemma 3.30 or Lemma 3.28.




, $q \in R_3$. GDD 1 of Row 17.

Case (iii) Omitting 1 and adding 3 is the same as Case (i) .

Case (iv) containing circle with four vertexes.

According to Lemma 3.39 (ii) we have to consider following cases.


(a) to (a).



is not an arithmetic root system by Lemma 3.2 (I).


(a) to (c). empty. (a) to (e). empty. (a) to (h). empty. (a) to (j). empty. (a) to (l). empty. (a) to (n). empty. (a) to (o). empty. (a) to (a'). empty.

(a) to (a'').



with $q = q^4$ is not an arithmetic root system by Lemma 3.2(I).


(e) to (e) or (15).



is an arithmetic root system by Lemma 3.4 (IV).

is affine.

(e) to (h) or (16).




is an arithmetic root system by Type 2.

is affine.


(e) to (j). is empty.

(e) to (l)



is not an arithmetic root system by Lemma 3.2 (I).

(e) to (o) or (17).




is an arithmetic root system by Type 4.

is affine.


(e) to (a'). empty. (e) to (a''). empty.

(h) to (h).




is not an arithmetic root system by Lemma 3.2 (I).

(h) to (l).



is not an arithmetic root system by Lemma 3.2(I).


(h) to (o).



is not an arithmetic root system by Lemma 3.2(I).

(h) to (a') empty. (h) to (a'') empty.


(j) to (j)



is not an arithmetic root system by Lemma 3.2 (I).


(j) to (l) empty. (j) to (o) empty. (j) to (a') empty. (j) to (a'') empty.

(l) to (l)



is not an arithmetic root system by Lemma 3.2 (I).


(l) to (o)



is not an arithmetic root system by Lemma 3.2 (I).

(l) to (a') empty. (l) to (a'') empty.


(o) to (o)



is not an arithmetic root system by Lemma 3.2(I).

(o) to (a') empty. (o) to (a'') empty.

(a') to (a') or (18).




is an arithmetic root system,

is affine.

(a') to (a'') empty.

(a'') to (a'')




is not an arithmetic root system by Lemma 3.2 (V).

4.28 GDD 2 of Row 11


Case (i) Omitting 3 and adding 1.

According to Lemma 3.3 (IV) we have to consider following cases.



, $q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 4.

Omitting 2 in whole is an arithmetic root system since symmetry of this GDD. It is similar in following of Case (i) . Whole is an arithmetic root system by GDD 2 of Row 13 or Type 6.



(a'') or (1).


, $q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 5.

Whole is affine by Lemma 3.4.

(a) or (2).  , $q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 6.

Whole is affine by Lemma 3.16.


(b) or (3).



, $q \in F^* \setminus \{1, -1\}$. GDD 3 of Row 6.


Omitting 1 or 2 is an arithmetic root system by Lemma 3.2 (III) or Lemma 3.2 (X)(b).

Whole is affine.

(c).  , $q \in F^* \setminus \{1, -1\}$. $q \notin R_3$, GDD 2 of Row 7.

is empty.

(d).




, $q \in F^* \setminus \{1, -1\}$. $q \notin R_3$, GDD 3 of Row 7. It is empty.

(e).  , $q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 8.

is an arithmetic root system by GDD 5 of Row 12 or Type 6.


(f) or (4).



, $q, r, s \in F^* \setminus \{1\}$, $qrs = 1$, $q \neq r$, $q \neq s$, $s \neq r$. GDD 3 of Row 9.

Omitting 1 is an arithmetic root system when $-q = r$ or $r = -1$ by Lemma 3.2 (XI).
 Omitting 2 is an arithmetic root system when $-q = s$ or $s = -1$ by Lemma 3.2 (XI).
 Whole is affine.


(g).



, $q^2, q^3 \neq 1$. GDD 3 of Row 10.

is empty.


(h).



, $q^2, q^3 \neq 1$. GDD 3 of Row 10.


is empty.

(i) or (5).



, $q \in R_3$. GDD 2 of Row 11.

Omitting 1 and Omitting 2 are arithmetic root systems by Lemma 3.2 (XI)(b). Whole is affine.



, $q \in R_3$. GDD 2 of Row 14.

is empty.

(k) or (6). , $q \in R_3$. GDD 2 of Row 15.

Whole is affine by Lemma 3.20.

(l) or (7). , $q \in R_3$. GDD 2 of Row 16.

Whole is affine by Lemma 3.16.

(m). , $q \in R_3$. GDD 3 of Row 16.

is empty.

(n). , $q \in R_3$. GDD 1 of Row 17.

is empty.

(n') or (8). , $q \in R_3$. GDD 1 of Row 17.

Whole is affine by Lemma 3.16.

(o) or (9). , $q \in R_3$. GDD 4 of Row 17.

Whole is affine by Lemma 3.16.

(p). , $q \in R_3$. GDD 4 of Row 17.

is empty.

(q). , $q \in R_3$. GDD 8 of Row 17.

is empty.

(r). , $q \in R_3$. GDD 9 of Row 17.

is empty.

Case (ii) Omitting 3 and adding 2 is the same as Case (i) .

Case (iii) Omitting 1 and adding 3 is the same as Case (i) .


Case (iv) containing circle with four vertexes.

This case can be ignored since it is over a cycle.

Case (v) complete diagrams containing circle with four vertexes.


According to Lemma 3.40 we have to consider following cases.

(b) in Case (i) or (10)




If 234 is an arithmetic root system, then $\tilde{q}_{34} = q^{-1}$ by Lemma 3.2(I). 134 is an arithmetic root system by Type 6. Whole is affine when $\tilde{q}_{34} = q^{-1}$.

(f) in Case (i)



If 134 is an arithmetic root system, then a_1 . $\tilde{q}_{34} = r$. or a_2 . $\tilde{q}_{34} = s$, $q = s^{-2}$, which is a contradiction by Lemma 3.2(XVIII) or a_3 . $\tilde{q}_{34} = q$. $s = q^{-2}$, which is a contradiction by Lemma 3.2(XVIII). a_1 . If 234 is an arithmetic root system, then $q = r^{-2}$, which is a contradiction by Lemma 3.2(XVIII).

(i) in Case (i) or (11)



with $q \in R_3$.

If 134 is an arithmetic root system, then $\tilde{q}_{34} = q^{-2}$ by Type 6. 234 is an arithmetic root system by Type 6. Whole is affine when $\tilde{q}_{34} = q^{-2}$.

4.29 GDD 1 of Row 12

Case (i) Omitting 3 and adding 1.

According to Lemma 3.3 (II) we have to consider following cases.

$$\begin{array}{c} \text{(a).} \\ q \quad q^{-1} \quad q \quad q^{-1} \quad q \quad q^{-1} \quad -q^{-1} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array}$$

$$q = -\xi, \xi \in R_3, \text{ GDD 1 of Row 1.}$$

is an arithmetic root system by Type 4.

$$\begin{array}{c} \text{(b) or (1).} \\ q \quad q^{-2} \quad q^2 \quad q^{-2} \quad q^2 \quad q^{-2} \quad -q^{-2} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array}$$

$$q = -\xi, \xi \in R_3, q \in F^* \setminus \{1, -1\}. \text{ GDD 1 of Row 2.}$$

Whole is affine by Lemma 3.4.

$$\begin{array}{c} \text{(c) or (2).} \\ q^2 \quad q^{-2} \quad q \quad q^{-1} \quad q \quad q^{-1} \quad -q^{-1} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array}$$

$$q = -\xi, \xi \in R_3, q \in F^* \setminus \{1, -1\}. \text{ GDD 1 of Row 3.}$$

Whole is affine by Lemma 3.4.

$$\begin{array}{c} \text{(d).} \\ -1 \quad q^{-1} \quad q \quad q^{-1} \quad q \quad q^{-1} \quad -q^{-1} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array}$$

$$q = -\xi, \xi \in R_3, q \in F^* \setminus \{1, -1\}. \text{ GDD 1 of Row 4.}$$

is an arithmetic root system Type 4.

$$\begin{array}{c} \text{(e) or (3).} \\ q \quad -q \quad -q^{-1} \quad -q \quad -q^{-1} \quad q \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array}$$

$$, q \in R_3. \text{ GDD 1 of Row 12.}$$

Whole is affine by Lemma 3.4.

$$\begin{array}{c} \text{(f) or (4).} \\ -1 \quad q^{-2} \quad q \quad q^{-1} \quad q \quad q^{-1} \quad -q^{-1} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array}$$

$$q = -\xi, \xi \in R_3, q \in R_3 \cup R_6. \text{ GDD 1 of Row 13.}$$

Whole is affine by Lemma 3.4.

$$\begin{array}{c} \text{(h) or (5).} \\ q \quad -q^{-1} \quad q \quad -q^{-1} \quad q \quad -q^{-1} \quad q^{-1} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array}$$

$$, q \in R_3. \text{ GDD 5 of Row 16.}$$

Whole is affine by Lemma 3.4.


$$\begin{array}{c} \text{(i).} \\ q^{-3} \quad q^{-1} \quad q \quad q^{-1} \quad q \quad q^{-1} \quad -q^{-1} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array}$$

$$q = -\xi, \xi \in R_3, q \in R_9. \text{ GDD 1 of Row 18.}$$


is empty.

Case (ii) Omitting 3 and adding 2.


According to Lemma 3.3 (II) we have to consider following cases.

(a) or (6).  $q = -\xi, \xi \in R_3, q \in F^* \setminus \{1\}$. GDD 1 of Row 1.

Omitting 1 in whole is an arithmetic root system by Type 3. Whole is affine by Lemma 3.4.

(b).  $, q \in F^* \setminus \{1, -1\}$. GDD 1 of Row 2.

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(I).

(c).  $, q \in F^* \setminus \{1, -1\}$. GDD 1 of Row 3.


Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(I).

(d) or (7).  $q \in R_6$. GDD 1 of Row 4.


Omitting 1 in whole is an arithmetic root system by Type 4. Whole is affine by Lemma 3.4.

(e).  $, q \in R_3$. GDD 1 of Row 12.


Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(I).

(f).  $, q \in R_6$. GDD 1 of Row 13.

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(I).

(h).  $, q \in R_3$. GDD 5 of Row 16.

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(I).




, $q \in R_6, q \in R_9$. GDD 1 of Row 18.

is empty.

Case (iii) Omitting 1 and adding 3.

According to inversion of Type 4 we have to consider following cases.




Case (iv) containing circle with four vertexes.

According to Lemma 3.39 (ii) we have to consider following cases.

(a) to (a). empty. (b) to (a) empty. (c) to (a) empty.


(d) to (a)



is not an arithmetic root system by Lemma 3.2 (I).

(e) to (a) empty.

(f) to (a)



is not an arithmetic root system by Lemma 3.2 (I).

(h) to (a) empty.


4.30 GDD 1 of Row 13

Case (i) Omitting 3 and adding 1.


According to Lemma 3.3 (II) we have to consider following cases.



Whole is affine by Lemma 3.37.



Whole is affine by Lemma 3.4.



Whole is affine when $q \in R_3$ by Lemma 3.18. Whole is affine when $q \in R_6$ by Lemma 3.37.

$$\begin{array}{c} \text{(d) or (4),} \\ -1 \quad q^{-1} \quad q \quad q^{-1}q \quad q^{-2} \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad , q \in F^* \setminus \{1, -1\}. \text{ GDD 1 of Row 4.}$$

is an arithmetic root system by Lemma 3.18(b) with $q \in R_3$. Whole is affine by Lemma 3.38 with $q \in R_6$.

$$\begin{array}{c} \text{(e) or (5),} \\ q \quad -q \quad -q \quad -q \quad -q \quad q^2 \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad , q \in R_3. \text{ GDD 1 of Row 12.}$$

Whole is affine by Lemma 3.38.

$$\begin{array}{c} \text{(f) or (6),} \\ -1 \quad q^{-2} \quad q \quad q^{-1}q \quad q^{-2} \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad , q \in R_3 \cup R_6. \text{ GDD 1 of Row 13.}$$

Whole is affine when $q \in R_3$ by Lemma 3.18. Whole is affine when $q \in R_6$ by Lemma 3.38.

$$\begin{array}{c} \text{(h) or (7),} \\ q \quad -q^{-1}q \quad -q^{-1}q \quad q^{-2} \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad , q \in R_3. \text{ GDD 5 of Row 16.}$$

Whole is affine by Lemma 3.37 or by Lemma 3.4.

$$\begin{array}{c} \text{(i).} \\ q^{-3} \quad q^{-1} \quad q \quad q^{-1}q \quad q^{-2} \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad , q \in R_9. \text{ GDD 1 of Row 18.}$$

is empty.

Case (ii) Omitting 3 and adding 2.


According to Lemma 3.3 (II) we have to consider following cases.

$$\text{(a) or (8), } \begin{array}{c} \bullet \\ | \\ q^{-2} \\ | \\ \bullet \\ \text{---} \bullet \text{---} \bullet \\ q \quad q^{-1} \quad q \quad q^{-1}q \end{array} \quad , q \in F^* \setminus \{1\}. \text{ GDD 1 of Row 1.}$$


Omitting 1 in whole is an arithmetic root system by Lemma 3.2 (XI)(a). Whole is affine by Lemma 3.4.

$$\text{(b). } \begin{array}{c} \bullet \\ | \\ q^{-4} \\ | \\ \bullet \\ \text{---} \bullet \text{---} \bullet \\ q \quad q^{-2} \quad q^2 \quad q^{-2} \quad q^2 \end{array} \quad , q^2 \in F^* \setminus \{1, -1\}. \text{ GDD 1 of Row 2.}$$

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(I).

(c).  , $q \in F^* \setminus \{1, -1\}$. GDD 1 of Row 3.


Omitting 1 in whole is not an arithmetic root system by Lemma 3.2 (V).

(d) or (9),  , $q \in F^* \setminus \{1, -1\}$. GDD 1 of Row 4.


Omitting 1 in whole is an arithmetic root system when $q \in R_3$ by Lemma 3.3 (III)(h).

Omitting 1 in whole is not an arithmetic root system when $q \in R_6$ by Lemma 3.3 (III).

Whole is affine by Lemma 3.4.

(e).  , $q \in R_3$. GDD 1 of Row 12.

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(I).

(f) or (10),  , $q \in R_3 \cup R_6$. GDD 1 of Row 13.

Omitting 1 in whole is an arithmetic root system by Lemma 3.2 (V)(b) with $q \in R_6$.

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2 (V) with $q \in R_3$.

Whole is affine by Lemma 3.4.


(h).  , $q \in R_3$. GDD 5 of Row 16.


Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(I).

(i).  , $q \in R_9$. GDD 1 of Row 18.
is empty.

Case (iii) Omitting 1 and adding 3.

According to Type 2 we have to consider following cases.


(a)  (see, $q \notin R_4$, GDD 2 of Row 5 in Table A2 or Type 2).

(b)  (see, Type 2).

Case (iv) containing circle with four vertexes.

(a) to (a) empty.

(a) to (b) or (11)




with $q = q^{-2}$ is an arithmetic root system by Type 7.


is affine.

(b) to (a) empty. (b) to (b) empty. (c) to (a) empty. (c) to (b) empty.

(d) to (a) or (12)




is an arithmetic root system by Lemma 3.2(II)(a) with $q \in R_3$; is not an arithmetic root system by Lemma 3.2 (X) with $q \in R_6$.




is affine.

(d) to (b) empty. (e) to (a) empty. (e) to (b) empty.

(f) to (a) or (13)




is an arithmetic root system by Type 2.




is affine.

(f) to (b) empty. (h) to (a) empty.

(h) to (b) or (14)



with $-q = q^{-2}$ is an arithmetic root system by Lemma 3.3 (III)(j).



is affine.

4.31 GDD 2 of Row 13

Case (i) Omitting 3 and adding 1.

According to Lemma 3.3 (II) we have to consider following cases.

(b) or (1) $\overset{q}{\bullet} \overset{q^{-1}q}{\bullet} \overset{q^{-1}}{\bullet} \overset{-q^{-1}q^2}{\bullet} \overset{-1}{\bullet}$ (see, when $q \in R_3$, Type 3; when $q \in R_6$, Type 4).

Whole is affine by Lemma 3.17.

(c) or (2) $\overset{-1}{\bullet} \overset{q^{-1}q}{\bullet} \overset{q^{-1}}{\bullet} \overset{-q^{-1}q^2}{\bullet} \overset{-1}{\bullet}$ (see, when $q \in R_3$, Type 3; when $q \in R_6$, Type 4).

Whole is affine when $q \in R_6$. by Lemma 3.17. Whole is an arithmetic root system when $q \in R_3$. by Lemma 3.17(c).

Case (ii) Omitting 3 and adding 2.

Checking step by step we have to consider following cases

(a) or (3) $\overset{-1}{\bullet} \overset{q^2}{\bullet} \overset{-1}{\bullet} \overset{-q^{-1}q^2}{\bullet} \overset{q^{-1}}{\bullet} \overset{q}{\bullet}$, (see GDD 2 of Row 13.)

Omitting 1 in whole is an arithmetic root system when $q \in R_3$ by Lemma 3.2 (V)(b).

Whole is affine by Lemma 3.4.

Case (iii) Omitting 1 and adding 3.

According to Type 3 and GDD 4 of Row 7 we have to consider following cases.

(a) $\overset{q^2}{\bullet} \overset{q^{-2}-1}{\bullet} \overset{q^2}{\bullet} \overset{-q^{-1}q^2}{\bullet} \overset{q}{\bullet}$ (see, $q^4, q^2 \neq 1$ GDD 3 of Row 5 in Table A2, or Type 2).

(b) $\overset{q^3}{\bullet} \overset{q^{-3}-1}{\bullet} \overset{q^2}{\bullet} \overset{-q^{-1}q^2}{\bullet} \overset{q}{\bullet}$ (see, $q^2, q^3 \neq 1$, GDD 4 of Row 7 in Table A2).


(c) $\overset{-1}{\bullet} \overset{q^{-2}-1}{\bullet} \overset{q^2}{\bullet} \overset{-q^{-1}q^2}{\bullet} \overset{q}{\bullet}$ (see, Type 3).

Case (iv) containing circle with four vertexes.


According to Lemma 3.39 (ii) we have to consider following cases.

(b) to (a) empty. (b) to (b) empty. (b) to (c) empty. (c) to (a) empty.

(c) to (b) or (4)




with $q^3 = -1$ is an arithmetic root system by Lemma 3.3 (I)(h) or by Lemma 3.2 (IV).




is affine.

(c) to (c) or (5)



is an arithmetic root system when $q \in R_3 \cup R_6$ by Lemma 3.2 (X).



is affine.

4.32 GDD 1 of Row 14

Case (i) Omitting 3 and adding 1.

According to Lemma 3.3 (I) we have to consider following cases.

$$\begin{array}{c} (a') \\ -1 \\ q \end{array} \begin{array}{c} -1 \\ q^{-1}q \end{array} \begin{array}{c} q^{-1} \\ q^{-1} \end{array} \begin{array}{c} -q^{-1} \\ -q^{-1} \end{array}$$

$q = -\xi, \xi \in R_3$, GDD 2 of Row 4.

is an arithmetic root system by GDD 2 of Row 4 or Type 4.

$$\begin{array}{c} (a) \text{ or } (1) \\ -q \\ q^2 \end{array} \begin{array}{c} -1 \\ q^{-2}q^2 \end{array} \begin{array}{c} q^{-2} \\ q^{-2} \end{array} \begin{array}{c} -q^{-2} \\ -q^{-2} \end{array}$$

$q^2 = -\xi, \xi \in R_3$. GDD 3 of Row 5.

is affine by by Lemma 3.4.

$$\begin{array}{c} (b) \text{ or } (2) \\ -1 \\ q \end{array} \begin{array}{c} -1 \\ q^{-2}q^2 \end{array} \begin{array}{c} q^{-2} \\ q^{-2} \end{array} \begin{array}{c} -q^{-2} \\ -q^{-2} \end{array}$$

GDD 2 of Row 6.

Whole is affine by Lemma 3.4, where $q^2 = -\xi, \xi \in R_3$.

$$\begin{array}{c} (d) \text{ or } (3) \\ -1 \\ q \end{array} \begin{array}{c} -1 \\ q^{-3}q^3 \end{array} \begin{array}{c} q^{-3} \\ q^{-3} \end{array} \begin{array}{c} -q^{-3} \\ -q^{-3} \end{array}$$

$q \notin R_3, q^3 \in R_6$, GDD 2 of Row 7.

Whole is affine by Lemma 3.4.

(f) or (4)

$$\begin{array}{c} -q^{-1}q^2 \quad -1 \quad q^{-3}q^3 \quad q^{-3} \quad -q^{-3} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad q \notin R_3 \cup R_2, q^3 \in R_6, \text{ GDD 4 of Row 7.}$$

Whole is affine when $q \in R_{18}$ by Lemma 3.4.

$$\begin{array}{c} q^{-1} \binom{\text{(g)}}{q} \quad -1 \quad q^{-1}q \quad q^{-1} \quad -q^{-1} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad , q = -\xi, \xi \in R_3.. \text{ GDD 1 of Row 8.}$$

is an arithmetic root system by Type 4,

$$\begin{array}{c} r \quad \binom{\text{(h)} \text{ or } \text{(5)}}{r^{-1} \quad -1} \quad q^{-1}q \quad q^{-1} \quad -q^{-1} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad , \text{ GDD 1 of Row 9.}$$

Whole is affine by Lemma 3.4.

$$\begin{array}{c} q \quad \binom{\text{(i')} \text{ or } \text{(6)}}{q^{-1} \quad -1} \quad q^2 \quad q^{-2} \quad q^2 \quad -q^2 \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad q \notin R_3 \cup R_2, \text{ GDD 2 of Row 10.}$$

Whole is affine by Lemma 3.4.

$$\begin{array}{c} q \quad \binom{\text{(k)} \text{ or } \text{(7)}}{q^{-1} \quad -1} \quad q^{-1}q \quad q^{-1} \quad -q^{-1} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad q \notin R_3 \cup R_2, \text{ GDD 1 of Row 10.}$$

Whole is affine $q \notin R_3 \cup R_2$ by Lemma 3.4.

$$\begin{array}{c} q^{-2} \binom{\text{(l)} \text{ or } \text{(8)}}{q^2 \quad -1} \quad q^{-1}q \quad q^{-1} \quad -q^{-1} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad q \notin R_3 \cup R_2, \text{ GDD 2 of Row 10.}$$

Whole is affine $q \notin R_3 \cup R_2$ by Lemma 3.4.

(m) empty.

$$\begin{array}{c} q^{-1} \binom{\text{(n)} \text{ or } \text{(9)}}{-q^{-1} \quad -1} \quad -q \quad -q^{-1} \quad q \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad , q \in R_3. \text{ GDD 3 of Row 14.}$$

Whole is affine by Lemma 3.4.


$$\begin{array}{c} -1 \quad \binom{\text{(p)} \text{ or } \text{(10)}}{q \quad -1} \quad -q^{-1}q \quad -q^{-1}q^{-1} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad , q \in R_3. \text{ GDD 2 of Row 16.}$$

Whole is affine by Lemma 3.4.

$$\begin{array}{c} q \quad \binom{\text{(r)} \text{ or } \text{(11)}}{-1 \quad -1} \quad -q^{-1}q \quad -q^{-1}q^{-1} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad , q \in R_3. \text{ GDD 4 of Row 16.}$$

Whole is affine by Lemma 3.4.

(s')




is empty.

Case (ii) Omitting 1 and adding 2.


According to Type 4 and Lemma 3.2(I) we have to consider following cases.

(a) or (12)



Omitting 3 in whole is an arithmetic root system by Type 7. Whole is affine by Lemma 3.4.


(b) or (13)



Omitting 3 in whole is an arithmetic root system by Type 7. Whole is affine by Lemma 3.4.

Case (iii) Omitting 1 and adding 3.


According to inversion of Type 4 we have to consider following cases.



Case (iv) containing circle with four vertexes.

According to Lemma 3.39 (ii) we have to consider following cases.


(a') to (a).



is not an arithmetic root system by Lemma 3.2(I)(I) with $q \in R_6$.


(a) to (a) empty.

(b) to (a)



is not an arithmetic root system by Lemma 3.2 (I) with $q \in R_6$.


(d) to (a).




with $q \in R_{18}$ is not an arithmetic root system by Lemma 3.2 (I).

(f) to (a) empty. (g) to (a) empty.

(h) to (a) or (14)




with $r = -1$ is an arithmetic root system by Lemma 3.2(I) with $q \in R_6$.




is affine.

(i') to (a) empty. (k) to (a) empty. (l) to (a) empty. (m) to (a) empty. (n) to (a) empty.

(p) to (a) or (15),



is an arithmetic root system by Lemma 3.2(I) with $q \in R_6$.




is affine.

(r) to (a) empty.

4.33 GDD 2 of Row 14


Case (i) Omitting 3 and adding 1.

According to Lemma 3.3 (IV) we have to consider following cases.




$q = -\xi, \xi \in R_3, q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 4.

is an arithmetic root system by Type 4.




$q^2 = -\xi, \xi \in R_3, q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 5.

Whole is affine by Lemma 3.4.




$q = -\xi, \xi \in R_3, q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 6.

Whole is affine by Lemma 3.4.



$q = -\xi, \xi \in R_3, q \in F^* \setminus \{1, -1\}$. $q \notin R_3$, GDD 2 of Row 7.


Whole is affine by Lemma 3.4.



$q = -\xi, \xi \in R_3, q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 8.


is an arithmetic root system by Type 4.

(j) or (4),




$q = -\xi, \xi \in R_3$. GDD 2 of Row 14.

Whole is affine by Lemma 3.4.




$q = -\xi, \xi \in R_3, q \in R_3$. GDD 2 of Row 15.

is empty.




$q = -\xi, \xi \in R_3, q \in R_3$. GDD 2 of Row 16.

is empty.




is empty.



$q = -\xi, \xi \in R_3$. by Lemma 3.2 (XI)(e).


Whole is affine by Lemma 3.4.

(o). empty.



$q \in R_3$. GDD 4 of Row 17.


Whole is affine by Lemma 3.4.



is empty.

Case (ii) Omitting 3 and adding 2.

According to Lemma 3.3 (IV) we have to consider following cases.



$q = -\xi, \xi \in R_3, q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 4.


Omitting 1 in whole is an arithmetic root system by Lemma 3.2(I). Whole is affine by Lemma 3.4.

(a'').  , $q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 5.


Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(I).

(a).  , $q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 6.

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(I).

(c).  , $q \in F^* \setminus \{1, -1\}$. $q \notin R_3$, GDD 2 of Row 7.


Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(I).

(e).  , $q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 8.

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(I).


(j).  , $q \in R_3$. GDD 2 of Row 14.

Omitting 1 is not an arithmetic root system by Lemma 3.2 (I).


(k).  , $q \in R_3$. GDD 2 of Row 15.

(l).  , $q \in R_3$. GDD 2 of Row 16.


(n).  , $q \in R_3$. GDD 1 of Row 17.



, $q \in R_3$. GDD 1 of Row 17.




, $q \in R_3$. GDD 4 of Row 17.



, $q \in R_3$. GDD 4 of Row 17.



Omitting 1 is not an arithmetic root system by Lemma 3.2 (I).



, $q \in R_3$. GDD 9 of Row 17.

Case (iii) Omitting 1 and adding 3.

According to inversion of Type 4 we have to consider following cases.




Case (iv) containing circle with four vertexes.

According to Lemma 3.39 (ii) we have to consider following cases.

(a') to (c). empty. (a') to (d). empty. (a'') to (c). empty. (a'') to (d). empty. (a) to (c). empty. (a) to (d). empty.

(c) to (c) or (8)




with $q \in R_6$ is an arithmetic root system by Lemma 3.2(I)(n).

is affine with $q \in R_6$.


(c) to (d) or (9)

with $q \in R_6$ is an arithmetic root system by Lemma 3.2 (IV).




is affine with $q \in R_6$.

(e) to (c).



is not an arithmetic root system by Lemma 3.2 (I)


(e) to (d).



is not an arithmetic root system by Lemma 3.2 (X)


(j) to (c). empty. (j) to (d). empty.

(k) to (c).




is not an arithmetic root system by Lemma 3.2(I).

(k) to (d) or (10),




is an arithmetic root system by Lemma 3.2 (X).




is affine with $q \in R_6$.

(l) to (c). empty. (l) to (d). empty.

(n') to (c) or (11),




is an arithmetic root system by Lemma 3.2 (I)(n).




is affine.

(n') to (d) or (12),




is an arithmetic root system by Lemma 3.2 (X)(b).



is affine.


(p) to (c) or (13),



is an arithmetic root system by Type 2.

is affine with $q \in R_6$.

(p) to (d) or (14),




is an arithmetic root system by Lemma 3.2 (X)(b).

is affine.

4.34 GDD 3 of Row 14


Case (i) Omitting 3 and adding 1.

According to Lemma 3.3 (III) we have to consider following cases.




$q = -\xi, \xi \in R_3, q \in R_6$. GDD 1 of Row 6.

Whole is affine by Lemma 3.4.




$q = -\xi, \xi \in R_3, q \in F^* \setminus \{1, -1\}, q \notin R_3$. GDD 1 of Row 7.

Whole is affine by Lemma 3.4.



$q = -\xi, \xi \in R_3, q \in F^* \setminus \{1, -1\}$. GDD 3 of Row 8.

is an arithmetic root system by Type 4.



$, q \in R_3$. GDD 1 of Row 14.

Whole is affine by Lemma 3.4.

$$\begin{array}{ccccccc} & & \text{(h).} & & & & \\ -1 & & q & & q^{-1}-1 & & q & & -q \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$

is empty.

$$q = -\xi, \xi \in R_3, q \in R_3. \text{ GDD 1 of Row 15.}$$

$$\begin{array}{ccccccc} & & \text{(j).} & & & & \\ -q & & -q^{-1}q & & q^{-1}-1 & & q & & -q \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$

is empty.

$$q = -\xi, \xi \in R_3, q \in R_3. \text{ GDD 1 of Row 16.}$$

$$\begin{array}{ccccccc} & & \text{(l).} & & & & \\ -1 & & -1 & & q & & q^{-1}-1 & & q & & -q \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$

is empty.

$$q = -\xi, \xi \in R_3, q \in R_3. \text{ GDD 2 of Row 17.}$$

$$\begin{array}{ccccccc} & & \text{(n) or (4),} & & & & \\ -1 & & q & & -q & & -q^{-1}-1 & & -q & & q \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$

$$, q \in R_3. \text{ GDD 5 of Row 17.}$$

Whole is affine by Lemma 3.4.

$$\begin{array}{ccccccc} & & \text{(o).} & & & & \\ -1 & & -q & & q & & q^{-1}-1 & & q & & -q \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$

is empty.

$$q = -\xi, \xi \in R_3, q \in R_3. \text{ GDD 7 of Row 17.}$$

$$\begin{array}{ccccccc} & & \text{(a').} & & & & \\ q & & q^{-1} & & q & & q^{-1}-1 & & q & & -q \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$

is an arithmetic root system by Type 4.

$$q = -\xi, \xi \in R_3 \text{ GDD 1 of Row 4.}$$

$$\begin{array}{ccccccc} & & \text{(a'') or (5)} & & & & \\ q & & q^{-2} & & q^2 & & q^{-2}-1 & & q & & -q \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$

$$q = -\xi, \xi \in R_3 \text{ GDD 1 of Row 5.}$$

Whole is affine by Lemma 3.4.

$$\begin{array}{ccccccc} & & \text{(k')} & & & & \\ -1 & & q & & -1 & & -1 & & -1 \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$

is empty.

$$\begin{array}{ccccccc} & & \text{(p')} & & & & \\ q^{-1} & & -q^{-1}-1 & & -1 & & -1 \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$

is empty.

Case (ii) Omitting 1 and adding 2.

According to Type 4 and GDD 9 of Row 17 or by Lemma 3.2(I) we have to consider following cases.

(a) or (6)

$$\begin{array}{c}
 -q^{-1} \quad -\Gamma q \quad -q^{-1} \\
 \bullet \text{---} \bullet \begin{array}{l} \nearrow \bullet \\ \searrow \bullet \end{array} \\
 \quad \quad \quad -q^{-1} \quad q^{-1}
 \end{array}
 , \text{ (see GDD 3 of Row 14 in Table A2 or Type 4.)}$$

Omitting 3 in whole is an arithmetic root system by GDD 3 of Row 14 or typ7. whole is affine by Lemma 3.4.

(b) or (7)

$$\begin{array}{c}
 -1_{-1} \quad -\Gamma q \quad -q^{-1} \\
 \bullet \text{---} \bullet \begin{array}{l} \nearrow \bullet \\ \searrow \bullet \end{array} \\
 \quad \quad \quad -q^{-1} \quad q^{-1}
 \end{array}
 , q \in R_3. \text{ (see GDD 9 of Row 17 in Table A2.)}$$

Omitting 3 in whole is an arithmetic root system GDD 1 of Row 9 in Table A2 or by Lemma 3.2 (IV). Whole is affine by Lemma 3.4.

(c)

$$\begin{array}{c}
 -1_{-q} \quad -\Gamma q \quad -q^{-1} \\
 \bullet \text{---} \bullet \begin{array}{l} \nearrow \bullet \\ \searrow \bullet \end{array} \\
 \quad \quad \quad -q^{-1} \quad q^{-1}
 \end{array}
 , (q \in R_3. \text{ see GDD 2 of Row 14 in Table A2 or Type 4.)$$

Omitting 3 in whole is not an arithmetic root system by Lemma 3.2(II).

Case (iii) Omitting 1 and adding 3.

According to inversion of Type 3 we have to consider following cases.

(a)

$$\begin{array}{c}
 q \quad q^{-1}-1 \quad -q \quad q \quad -1 \\
 \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet
 \end{array}
 , \text{ (see, } q \in R_3, \text{ GDD 4 of Row 15.)}$$

empty.

(c)

$$\begin{array}{c}
 q \quad q^{-1}-1 \quad -q \quad q^{-1}-1 \\
 \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet
 \end{array}
 \text{ (see, } q \in R_3, \text{ GDD 5 of Row 17 in Table A2).}$$

is empty.

(d)

$$\begin{array}{c}
 q \quad q^{-1}-1 \quad -q \quad q \quad -1 \\
 \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet
 \end{array}
 \text{ (see, } q \in R_6, \text{ GDD 6 of Row 17 in Table A2).}$$

(g)

$$\begin{array}{c}
 q \quad q^{-1}-1 \quad -q \quad q^{-1}-1 \\
 \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet
 \end{array}
 , \text{ (see, } q \in R_6, \text{ GDD 7 of Row 17.)}$$

Case (iv) Containing circle with four vertexes.

According to Lemma 3.39 (ii) we have to consider following cases.


(a) to (d) empty. (a) to (g) empty.

(c) to (d).

$$\begin{array}{c}
 q \quad q^{-3} \quad q^3 \quad -q \quad -q \\
 \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet
 \end{array}$$


with $q^3 = -1$ is not an arithmetic root system by Lemma 3.2 (I).

(c) to (g).




is not an arithmetic root system by Lemma 3.2 (I) with $q \in R_6$.

(e) to (d)




is not an arithmetic root system by Lemma 3.2 (I).

(e) to (g) or (8),




is an arithmetic root system by Lemma 3.2 (IV) with $q \in R_6$.



is affine.


(g) to (d) empty. (g) to (g) empty.

(n) to (d)




is not an arithmetic root system by Lemma 3.2 (I) with $q \in R_6$.

(n) to (g) or (9)



is an arithmetic root system by Type 3 with $q \in R_6$.



is affine.

(a') to (d) empty. (a') to (g) empty. (a'') to (d) empty. (a'') to (g) empty.

4.35 GDD 1 of Row 15

Case (i) Omitting 3 and adding 1.

According to Lemma 3.3 (I) we have to consider following cases.



is an arithmetic root system when $q \in R_3$ by GDD 3 of Row 21.

(a) or (1)

$$\begin{array}{c} -q^{-1/2} \quad -1 \quad q^{-2}q^2 \quad q^2 \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad q^4 \neq 1, \text{ GDD 3 of Row 5.}$$

Whole is affine by Lemma 3.4.

$$\begin{array}{c} \text{(b) or (2)} \\ -1 \quad q \quad -1 \quad q^{-2}q^2 \quad q^2 \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad q^2 \neq 1, \text{ GDD 2 of Row 6.}$$

Whole is affine by Lemma 3.7.

$$\begin{array}{c} \text{(d) or (3)} \\ -1 \quad q \quad -1 \quad q^{-3}q^3 \quad q^3 \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad q^3 \in R_3, \text{ GDD 2 of Row 7.}$$

Whole is affine when $q^3 \in R_3$ by Lemma 3.15.

$$\begin{array}{c} \text{(f) or (4)} \\ -q \quad q^2 \quad -1 \quad q^{-3}q^3 \quad q^3 \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad , \text{ GDD 4 of Row 7.}$$

Whole is affine when $q^3 \in R_3$ by by Lemma 3.4.

$$\begin{array}{c} \text{(g)} \\ q^{-1} \quad q \quad -1 \quad q^{-1}q \quad q \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad , \text{ GDD 1 of Row 8.}$$

is an arithmetic root system by GDD 1 of Row 20 or by Lemma 3.18.

$$\begin{array}{c} \text{(h) or (5)} \\ r \quad r^{-1} \quad -1 \quad q^{-1}q \quad q \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad , \text{ GDD 1 of Row 9.}$$

Whole is affine by Lemma 3.7.

$$\begin{array}{c} \text{(i') or (6)} \\ q \quad q^{-1} \quad -1 \quad q^2 \quad q^{-2} \quad q^{-2} \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad , q \notin R_3. \text{ GDD 2 of Row 10.}$$

Whole is affine by Lemma 3.7 with $q \in R_6$.

$$\begin{array}{c} \text{(k)} \\ q \quad q^{-1} \quad -1 \quad q^{-1}q \quad q \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad , q \notin R_3. \text{ GDD 1 of Row 10.}$$

is empty.

$$\begin{array}{c} \text{(l)} \\ q^{-2} \quad q^2 \quad -1 \quad q^{-1}q \quad q \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad , q \notin R_3. \text{ GDD 2 of Row 10.}$$

is empty.

$$\begin{array}{c} \text{(m) or (7)} \\ q \quad q^{-1} \quad -1 \quad q^{-1}q \quad q \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad , q \in R_3. \text{ GDD 1 of Row 11.}$$

Whole is affine by Lemma 3.7.

$$\begin{array}{c}
 q^{-1} \text{ (n)} \\
 -q^{-1} -1 \quad -q \quad -q^{-1} -q^{-1} -1 \\
 \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet
 \end{array}
 \quad , q \in R_3. \text{ GDD 3 of Row 14.}$$

is empty.

$$\begin{array}{c}
 \text{(p)} \\
 -1 \quad q \quad -1 \quad -q^{-1} -q \quad -q \quad -1 \\
 \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet
 \end{array}
 \quad , q \in R_3. \text{ GDD 2 of Row 16.}$$

is empty.

$$\begin{array}{c}
 \text{(r)} \\
 q \quad -1 \quad -1 \quad -q^{-1} -q \quad -q \quad -1 \\
 \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet
 \end{array}
 \quad , q \in R_3. \text{ GDD 4 of Row 16.}$$

is empty.

$$\begin{array}{c}
 \text{(s')} \\
 -1 \quad q \quad -1 \quad -1 \quad -1 \quad -1 \\
 \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet
 \end{array}
 \quad , q \in R_3. \text{ GDD 1 of Row 17.}$$

is empty.

Case (ii) Omitting 3 and adding 2.

According to Lemma 3.3 (III) we have to consider following cases.

$$\begin{array}{c}
 \text{(a).} \\
 \begin{array}{c}
 -1 \\
 \bullet \\
 q \\
 \bullet \text{---} \bullet \text{---} \bullet \\
 q^2 \quad q^{-2} \quad q \quad q^{-1} -1
 \end{array}
 \end{array}
 \quad , q \in R_3. \text{ GDD 1 of Row 6.}$$

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2 (XI).

$$\begin{array}{c}
 \text{(c).} \\
 \begin{array}{c}
 -1 \\
 \bullet \\
 q \\
 \bullet \text{---} \bullet \text{---} \bullet \\
 q^3 \quad q^{-3} \quad q \quad q^{-1} -1
 \end{array}
 \end{array}
 \quad , q \in F^* \setminus \{1, -1\} , q \notin R_3. \text{ GDD 1 of Row 7.}$$

is empty.

$$\begin{array}{c}
 \text{(e) or (8),} \\
 \begin{array}{c}
 -1 \\
 \bullet \\
 q \\
 \bullet \text{---} \bullet \text{---} \bullet \\
 -1 \quad q^{-1} \quad q \quad q^{-1} -1
 \end{array}
 \end{array}
 \quad , q \in F^* \setminus \{1, -1\}. \text{ GDD 3 of Row 8.}$$


Omitting 1 in whole is an arithmetic root system by GDD 1 of Row 15, or by Lemma 3.2 (XI)(h). Whole is affine by Lemma 3.4.

$$\begin{array}{c}
 \text{(g).} \\
 \begin{array}{c}
 -1 \\
 \bullet \\
 -q^{-1} \\
 \bullet \text{---} \bullet \text{---} \bullet \\
 q \quad -q \quad -q^{-1} q \quad -1
 \end{array}
 \end{array}
 \quad , q \in R_3. \text{ GDD 1 of Row 14.}$$

is empty.

$$\begin{array}{c}
 \text{(h).} \\
 \begin{array}{c}
 -1 \\
 \bullet \\
 q \\
 \bullet \text{---} \bullet \text{---} \bullet \\
 -1 \quad q \quad q \quad q^{-1} -1
 \end{array}
 \end{array}
 \quad , q \in R_3. \text{ GDD 1 of Row 15.}$$

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2 (XI).

(j).  , $q \in R_3$. GDD 1 of Row 16.

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2 (XI).


(l).  , $q \in R_3$. GDD 2 of Row 17.

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2 (XI).

(n).  , $q \in R_3$. GDD 5 of Row 17.
is empty.

(o) or (9),  , $q \in R_3$. GDD 7 of Row 17.

Omitting 1 is an arithmetic root system by Lemma 3.2 (XI)(f). Whole is affine by Lemma 3.4.


(a') or (10),  GDD 1 of Row 4.

Omitting 1 is an arithmetic root system by Lemma 3.3 (III)(h). Whole is affine by Lemma 3.4.

(a'')  GDD 1 of Row 5.

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2 (XI).



(k')  GDD 1 of Row 17.
is empty.



GDD 9 of Row 17.

Case (iii) Omitting 1 and adding 3.


According to Type 2 we have to consider following cases.



Case (iv) Containing circle with four vertexes.

According to Type 2 we have to consider following cases.

(a') to (a) or (11),




is an arithmetic root system by Lemma 3.2 (XI) or Type 7.

is affine.


(a') to (b) empty.

(a) to (a) empty. (a) to (b) empty.

(b) to (a) or (12)




is an arithmetic root system by Lemma 3.2 (XI)(b). .



is affine.

(b) to (b) empty.


(d) to (a).




is not an arithmetic root system by Lemma 3.2 (XI).

(d) to (b) empty. (f) to (a) empty. (f) to (b) empty. (g) to (a) empty. (g) to (b) empty.

(h) to (a) or (13).




with $r = -1$. Whole is an arithmetic root system by Lemma 3.2 (XI)(e).




is affine.

(h) to (b) or (14)




with $r = q^{-2}$ is an arithmetic root system by Type 7.




is affine.

(i') to (a) empty.

(i') to (b) or (15)




with $q = q^4$ is an arithmetic root system by Type 7.




is affine.

(m) to (a) empty.

(m) to (b) or (16)



with $q = q^{-2}$ is arithmetic root system by Type 7.




is affine.

4.36 GDD 2 of Row 15

Case (i) Omitting 3 and adding 1.

According to Lemma 3.3 (IV) we have to consider following cases.



, $q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 4.

is an arithmetic root system by GDD 6 of Row 21, or by Lemma 3.21(b).

(a'') or (1).

$$\begin{array}{ccccccc} q & q^{-2} & -1 & q^2 & -1 & q^2 & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

, $q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 5.

Whole is affine by Lemma 3.4.

$$\begin{array}{ccccccc} q^2 & \text{(a) or (2).} & & & & & \\ q^{-2} & -1 & q & -1 & q & & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

, $q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 6.

is affine by Lemma 3.21.

$$\begin{array}{ccccccc} q^3 & \text{(c).} & & & & & \\ q^{-3} & -1 & q & -1 & q & & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

, $q \in F^* \setminus \{1, -1\}$. $q \notin R_3$, GDD 2 of Row 7.

is empty.

$$\begin{array}{ccccccc} -1 & \text{(e).} & & & & & \\ q^{-1} & -1 & q & -1 & q & & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

, $q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 8.

is an arithmetic root system by GDD 3 of Row 20 or by Lemma 3.21(a).

$$\begin{array}{ccccccc} q & \text{(j).} & & & & & \\ -q & -1 & -q^{-1} & -1 & -q^{-1} & & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

, $q \in R_3$. GDD 2 of Row 14.

is empty.

$$\begin{array}{ccccccc} -1 & \text{(k) or (3).} & & & & & \\ q & -1 & q & -1 & q & & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

, $q \in R_3$. GDD 2 of Row 15.

is affine by Lemma 3.21.

$$\begin{array}{ccccccc} -q & \text{(l) or (4).} & & & & & \\ -q^{-1} & -1 & q & -1 & q & & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

, $q \in R_3$. GDD 2 of Row 16.

is affine by Lemma 3.21.

$$\begin{array}{ccccccc} -1 & \text{(n).} & & & & & \\ q & -1 & -1 & -1 & -1 & & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

, $q \in R_3$. GDD 1 of Row 17.

is empty.

$$\begin{array}{ccccccc} -1 & \text{(n') or (5).} & & & & & \\ -1 & -1 & q & -1 & q & & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$


, $q \in R_3$. GDD 1 of Row 17.

is affine by Lemma 3.21.

$$\begin{array}{ccccccc} -1 & \text{(o) or (6).} & & & & & \\ -q & -1 & q & -1 & q & & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$


, $q \in R_3$. GDD 4 of Row 17.

is affine by Lemma 3.21.



, $q \in R_3$. GDD 4 of Row 17.

is empty.




, $q \in R_3$. GDD 9 of Row 17.

is empty.

Case (ii) Omitting 3 and adding 2.


According to Lemma 3.3 (IV) we have to consider following cases.



(a') or (7).

, $q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 4.


Omitting 1 in whole is an arithmetic root system because of symmetry of this GDD. It is similar in following of Case (ii) . Whole is affine by Lemma 3.21.



(a'') or (8).

, $q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 5.


is affine by Lemma 3.21.



(a) or (9).

, $q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 6.


is affine by Lemma 3.21.



(c).

, $q \in F^* \setminus \{1, -1\}$. $q \notin R_3$, GDD 2 of Row 7.


is empty.



(e) or (10).

, $q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 8.


is affine by Lemma 3.21.



(j).


, $q \in R_3$. GDD 2 of Row 14.

is empty.




, $q \in R_3$. GDD 2 of Row 15.

is affine by Lemma 3.21.




, $q \in R_3$. GDD 2 of Row 16.

is affine by Lemma 3.21.




, $q \in R_3$. GDD 1 of Row 17.




, $q \in R_3$. GDD 1 of Row 17.

is affine by Lemma 3.21.




, $q \in R_3$. GDD 4 of Row 17.

is affine by Lemma 3.21.



, $q \in R_3$. GDD 4 of Row 17.




, $q \in R_3$. GDD 9 of Row 17.

Case (iii) Omitting 1 and adding 3 is the same as (i).


Case (iv) containing circle with four vertexes.

According to Lemma 3.39 (ii) we have to consider following cases.

(a') to (a') or (15).



is an arithmetic root system.




is affine.

(a') to (a'').

empty.


(a') to (a). empty. (a') to (e) empty. (a') to (k) empty. (a') to (l) empty. (a') to (n') empty. (a') to (o) empty.

(a'') to (a'').




is not an arithmetic root system by Lemma 3.2 (V).

(a'') to (a) or (16).




with $q = q^4$ is an arithmetic root system by Lemma 3.3 (III)(h).




is affine.

(a'') to (e) empty. (a'') to (k) empty. (a'') to (l). empty. (a'') to (n') empty. (a'') to (o) empty.

(a) to (a) or (17).




is an arithmetic root system,




is affine.

(a) to (e) empty. (a) to (k) empty. (a) to (l). empty. (a) to (n') empty. (a) to (o) empty. (e) to (e) or (18).




is an arithmetic root system by Lemma 3.2 (XI)(b).



is affine.

(e) to (k) or (19).

is an arithmetic root system by Lemma 3.2 (XI) or Type 7.




is affine.

(e) to (l). empty.

(e) to (n') or (20).

is arithmetic root system by Lemma 3.2 (XI)(e).




is affine.


(e) to (o).

is not an arithmetic root system by Lemma 3.2 (XI).

(k) to (k) or (21).



is an arithmetic root system by Lemma 3.2 (XI)(b).




is affine.

(k) to (l). empty.


(k) to (n') or (22).

is arithmetic root system by Lemma 3.2 (XI).




is affine.


(k) to (o) or (23).



is an arithmetic root system by Lemma 3.2 (XI)(d).




is affine.
 (l) to (l) or (24).




is an arithmetic root system Type 7.

is affine.
 (l) to (n'). empty. (l) to (o) empty.
 (n') to (n') or (25)




is an arithmetic root system,

is affine.
 (n') to (o) or (26).



is an arithmetic root system by Lemma 3.2 (XI)(e).

is affine.
 (o) to (o).




is not an arithmetic root system by Lemma 3.2 (XI).

4.37 GDD 3 of Row 15

Case (i) Omitting 3 and adding 1.


According to Lemma 3.3 (III) we have to consider following cases.




, $q \in R_3$. GDD 1 of Row 6.

Omitting 2 in whole is an arithmetic root system by Type 1. Whole is affine by Lemma 3.4.

(b) or (2).




Omitting 1 in whole is not an arithmetic root system by Lemma 3.2 (X)(b). Omitting 2 in whole is an arithmetic root system by Type 7. Whole is affine.



is empty.

(d).



$, q \in F^* \setminus \{1, -1\}$, $q \notin R_3$. GDD 3 of Row 7. It is empty.



is an arithmetic root system by GDD 7 of Row 20.




is empty.



Omitting 2 in whole is an arithmetic root system Lemma 3.3(II)(f). Whole is affine by Lemma 3.4.

(i) or (4).




Omitting 2 in whole is an arithmetic root system by Lemma 3.2 (IV). Omitting 1 in whole is an arithmetic root system by Type 7. Whole is affine,



Omitting 2 in whole is not an arithmetic root system by Lemma 3.3(II).

(k).




Omitting 2 in whole is not an arithmetic root system by Lemma 3.3(II).



Omitting 2 in whole is not an arithmetic root system by Lemma 3.3(II).


(m).



Omitting 2 in whole is not an arithmetic root system by Lemma 3.3(II).




is empty.



Omitting 2 in whole is not an arithmetic root system by Lemma 3.2 (IX).


(p).



Omitting 2 in whole is not an arithmetic root system by Lemma 3.2 (IX).



is an arithmetic root system by GDD 7 of Row 21.




Omitting 2 in whole is an arithmetic root system by Type 2. is affine by Lemma 3.4.



is empty.

(p')




GDD 9 of Row 17.

is empty.


Case (ii) Omitting 3 and adding 2.

According to Lemma 3.3 (I) we have to consider following cases.




GDD 2 of Row 4.

Omitting 1 in whole is an arithmetic root system because of symmetry of this GDD. It is similar in following of Case (ii) . is affine by Lemma 3.24.



GDD 3 of Row 5.


is affine by Lemma 3.4.



GDD 2 of Row 6.


is affine by Lemma 3.24.

(c)



GDD 3 of Row 7.


is empty.



GDD 2 of Row 7.

is empty.

(e)



GDD 3 of Row 7.

is empty.

(f)
$$\begin{array}{c} q^{-3} q^3 \\ \bullet \quad \bullet \\ \text{---} \bullet \quad \bullet \text{---} \\ -q^{-1}q^2 \quad -1 \quad q^{-3}q^3 \end{array}$$

, GDD 4 of Row 7.

is empty.

(g) or (9)
$$\begin{array}{c} q^{-1} q \\ \bullet \quad \bullet \\ \text{---} \bullet \quad \bullet \text{---} \\ q^{-1}q \quad -1 \quad q^{-1}q \end{array}$$

, GDD 1 of Row 8.

is affine by Lemma 3.24.

(h) or (10)
$$\begin{array}{c} q^{-1} q \\ \bullet \quad \bullet \\ \text{---} \bullet \quad \bullet \text{---} \\ r \quad r^{-1} \quad -1 \quad q^{-1}q \end{array}$$

, GDD 1 of Row 9.

is affine by Lemma 3.24.

(i')
$$\begin{array}{c} q^{-2} q^2 \\ \bullet \quad \bullet \\ \text{---} \bullet \quad \bullet \text{---} \\ q \quad q^{-1} \quad -1 \quad q^2 \quad q^{-2} \end{array}$$

, $q \notin R_3$. GDD 2 of Row 10.

is empty.

(k)
$$\begin{array}{c} q^{-1} q \\ \bullet \quad \bullet \\ \text{---} \bullet \quad \bullet \text{---} \\ q \quad q^{-1} \quad -1 \quad q^{-1}q \end{array}$$

, $q \notin R_3$. GDD 1 of Row 10.

is empty.

(l)
$$\begin{array}{c} q^{-1} q \\ \bullet \quad \bullet \\ \text{---} \bullet \quad \bullet \text{---} \\ q^{-2}q^2 \quad -1 \quad q^{-1}q \end{array}$$

, $q \notin R_3$. GDD 2 of Row 10.

is empty.

(m) or (11)
$$\begin{array}{c} q^{-1} q \\ \bullet \quad \bullet \\ \text{---} \bullet \quad \bullet \text{---} \\ q \quad q^{-1} \quad -1 \quad q^{-1}q \end{array}$$

, $q \in R_3$. GDD 1 of Row 11.


is affine by Lemma 3.24.

(n)
$$\begin{array}{c} -q^{-1} \\ \bullet \quad \bullet \\ \text{---} \bullet \quad \bullet \text{---} \\ q^{-1} \quad -q^{-1} \quad -1 \quad -q \quad -q^{-1} \end{array}$$

, $q \in R_3$. GDD 3 of Row 14.


is empty.

(o) or (12)



GDD 3 of Row 15.


Omitting 1 in whole is an arithmetic root system by Lemma 3.2 (IV). Omitting 2 in whole is an arithmetic root system by Type 7. Whole is affine.



, $q \in R_3$. GDD 2 of Row 16.


is empty.

(q)




GDD 3 of Row 16.

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2 (X).



, $q \in R_3$. GDD 4 of Row 16.


is affine by Lemma 3.4 or by Lemma 3.24.



, $q \in R_3$. GDD 1 of Row 17.

is empty.

(s) or (14)




GDD 3 of Row 17.

Omitting 1 in whole is an arithmetic root system by Lemma 3.2 (IV). Omitting 2 in

whole is not an arithmetic root system by Lemma 3.3(II). Whole is affine.

(t)



Omitting 1 in whole is an arithmetic root system by Lemma 3.2 (IV) or by Lemma 3.2 (V). Omitting 2 in whole is not an arithmetic root system by Lemma 3.3(II).


Case (iii) Omitting 1 and adding 3 is the same as Case (i) .

Case (iv) Containing circle with four vertexes. empty since it is a circle.

Case (v) complete diagram containing circle with four vertexes.

According to Lemma 3.40 we have to consider following cases.


(b) in Case (i) or (14)



with $q \in R_3$.


If 134 is an arithmetic root system, then $\tilde{q}_{43} = q^{-1}$ by Lemma 3.2(I). 234 is an arithmetic root system by GDD 3 of Row 6 . Whole is affine when $\tilde{q}_{43} = q^{-1}$.

(i) in Case (i)




134 is not an arithmetic root system by Lemma 3.2 (XVI).


(k) in Case (i)




134 is not an arithmetic root system by Lemma 3.2(I).
 (m) in Case (i)




134 is not an arithmetic root system by Lemma 3.2(XVI).
 (p) in Case(i)



134 is not an arithmetic root system by Lemma 3.2(I).
 (o) in Case (ii)




134 is not an arithmetic root system by Lemma 3.2(XVI).
 (q) in Case (ii)




134 is not an arithmetic root system by Lemma 3.2(I).

(s) in Case (ii)



134 is not an arithmetic root system by Lemma 3.2(XVI).

(t) in Case (ii)



134 is not an arithmetic root system by Lemma 3.2(I).

4.38 GDD 4 of Row 15

Case (i) Omitting 3 and adding 1.

According to Type 3 we have to consider following cases.

(a) $q^2 \quad q^{-2} - 1 \quad q^{-1} \quad -q \quad q^{-1} \quad -1$ (see, $q \in R_3$, GDD 3 of Row 5 in Table A2 or Type 3).

Whole is an arithmetic root system by GDD 2 of Row 20 in Table B or by Lemma 3.17(b)


(b) $-1 \quad q^{-2} - 1 \quad q^{-1} \quad -q \quad q^{-1} \quad -1$ (Type 3).

Whole is an arithmetic root system by GDD 4 of Row 21 in Table B or by Lemma 3.17(a).

Case (ii) Omitting 3 and adding 2.

Checking step by step we have to consider following cases.


(a) or (1)



Omitting 1 in whole is an arithmetic root system because of symmetry of this GDD.


It is similar in following of Case (ii) . Whole is affine by Lemma 3.4.

(b) or (2)



is affine by Lemma 3.4.

(c) or (3)



is affine by Lemma 3.4


Case (iii) Omitting 1 and adding 3 is the same as Case (i) .

There are not any affine.

Case (iv) Containing circle with four vertexes.

According to Lemma 3.39 (ii) we have to consider following cases.

(a) to (a) or (4).




is an arithmetic root system,

is affine.

(a) to (b). empty

(b) to (b) or (5).



is an arithmetic root system by Lemma 3.2 (XI)(b).

is affine.

4.39 GDD 1 of Row 16

Case (i) Omitting 3 and adding 1.

According to Lemma 3.3 (I) we have to consider following cases.

$$\begin{array}{c} (a') \text{ or } (1) \\ -1 \quad -1 \\ q \quad q^{-1}q \quad -q^{-1-q} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad \text{GDD 2 of Row 4.}$$

is affine by Lemma 3.34.

$$\begin{array}{c} (a) \text{ or } (2) \\ -q \quad -1 \\ q^2 \quad q^{-2}q^2 \quad -q^{-2-q^2} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad \text{GDD 3 of Row 5.}$$

is affine by Lemma 3.4.

$$\begin{array}{c} (b) \text{ or } (3) \\ -1 \quad -1 \\ q \quad q^{-2}q^2 \quad -q^{-2-q^2} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad \text{GDD 2 of Row 6.}$$

is affine by Lemma 3.7.

$$\begin{array}{c} (d) \text{ or } (4) \\ -1 \quad -1 \\ q \quad q^{-3}q^3 \quad -q^{-3-q^3} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad \text{GDD 2 of Row 7.}$$

is affine by Lemma 3.7 with $q^3 \in R_3$.

(f) or (5)

$$\begin{array}{c} -q^{-1} \quad -1 \\ q^2 \quad q^{-3}q^3 \quad -q^{-3-q^3} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad , \text{GDD 4 of Row 7.}$$

is affine by Lemma 3.4 with $q^3 \in R_3$.

$$\begin{array}{c} (g) \text{ or } (6) \\ q^{-1} \quad -1 \\ q \quad q^{-1}q \quad -q^{-1-q} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad , \text{GDD 1 of Row 8.}$$

is affine by Lemma 3.38.

$$\begin{array}{c} (h) \text{ or } (7) \\ r \quad -1 \\ r^{-1} \quad q^{-1}q \quad -q^{-1-q} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad , \text{GDD 1 of Row 9.}$$

Whole is an arithmetic root system when $r = -q$ by Lemma 3.7(c).

Whole is affine when $r \neq -q$ by Lemma 3.7.

$$\begin{array}{c} (i') \text{ or } (8) \\ q \quad -1 \\ q^{-1} \quad q^2 \quad q^{-2} \quad -q^{-2-q^2} \\ \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad , q \notin R_3. \text{GDD 2 of Row 10.}$$

is affine by Lemma 3.7 with $q^2 \in R_3$.

(k)

$$\begin{array}{ccccccc} q & q^{-1} & -1 & q^{-1}q & -q^{-1-q} & & \\ \bullet & \bullet & \bullet & \bullet & \bullet & & \end{array}$$
 , $q \notin R_3$. GDD 1 of Row 10.
 is empty.

$$\begin{array}{ccccccc} q^{-2} & \overset{(l)}{q^2} & -1 & q^{-1}q & -q^{-1-q} & & \\ \bullet & \bullet & \bullet & \bullet & \bullet & & \end{array}$$
 , $q \notin R_3$. GDD 2 of Row 10.
 is empty.

$$\begin{array}{ccccccc} q & \overset{(m)}{q^{-1}} & \overset{(9)}{-1} & q^{-1}q & -q^{-1-q} & & \\ \bullet & \bullet & \bullet & \bullet & \bullet & & \end{array}$$
 , $q \in R_3$. GDD 1 of Row 11.

is affine by Lemma 3.7.

$$\begin{array}{ccccccc} q^{-1} & \overset{(n)}{-q^{-1}} & -1 & -q & -q^{-1} & q^{-1} & \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \end{array}$$
 , $q \in R_3$. GDD 3 of Row 14.
 is empty.

$$\begin{array}{ccccccc} -1 & \overset{(p)}{q} & -1 & -q^{-1-q} & -q^{-1-q} & & \\ \bullet & \bullet & \bullet & \bullet & \bullet & & \end{array}$$
 , $q \in R_3$. GDD 2 of Row 16.
 is empty.

$$\begin{array}{ccccccc} q & \overset{(r)}{-1} & -1 & -q^{-1-q} & q^{-1} & q & \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \end{array}$$
 , $q \in R_3$. GDD 4 of Row 16.
 is empty.

$$\begin{array}{ccccccc} -1 & \overset{(s')}{q} & -1 & -1 & -1 & & \\ \bullet & \bullet & \bullet & \bullet & \bullet & & \end{array}$$
 is empty.

Case (ii) Omitting 3 and adding 2.

According to Lemma 3.3 (III) we have to consider following cases.

$$(a). \begin{array}{ccccccc} & & & -q & & & \\ & & & -q^{-1} & & & \\ & & \bullet & & & & \\ \bullet & \bullet & \bullet & \bullet & \bullet & & \\ q^2 & q^{-2} & q & q^{-1} & -1 & & \end{array}$$
 , $q \in R_3$. GDD 1 of Row 6.

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2 (V).


$$(c). \begin{array}{ccccccc} & & & -q & & & \\ & & & -q^{-1} & & & \\ & & \bullet & & & & \\ \bullet & \bullet & \bullet & \bullet & \bullet & & \\ q^3 & q^{-3} & q & q^{-1} & -1 & & \end{array}$$
 , $q \in F^* \setminus \{1, -1\}$, $q \notin R_3$. GDD 1 of Row 7.

is empty.

$$(e) \text{ or } (10). \begin{array}{ccccccc} & & & -q & & & \\ & & & -q^{-1} & & & \\ & & \bullet & & & & \\ \bullet & \bullet & \bullet & \bullet & \bullet & & \\ -1 & q^{-1} & q & q^{-1} & -1 & & \end{array}$$
 , $q \in F^* \setminus \{1, -1\}$. GDD 3 of Row 8.

Omitting 1 in whole is an arithmetic root system by Lemma 3.3 (III)(j). Whole is affine by Lemma 3.4.

(g).  , $q \in R_3$. GDD 1 of Row 14.
is empty.

(h).  , $q \in R_3$. GDD 1 of Row 15.

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2 (V).

(j).  , $q \in R_3$. GDD 1 of Row 16.

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2 (V).

(l).  , $q \in R_3$. GDD 2 of Row 17.

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2 (V).

(n).  , $q \in R_3$. GDD 5 of Row 17.
is empty.

(o).  , $q \in R_3$. GDD 7 of Row 17.

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2 (V).

(a').  GDD 1 of Row 4.

Omitting 1 in whole is not an arithmetic root system by Lemma 3.3(II).

(a'')  GDD 1 of Row 5.

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(I).

(k') $\overset{-1}{\bullet} \overset{q}{\bullet} \overset{-1}{\bullet} \overset{-1}{\bullet} \overset{-1}{\bullet}$ GDD 1 of Row 17.
 is empty.

(p') $\overset{q^{-1}}{\bullet} \overset{-q^{-1}}{\bullet} \overset{-1}{\bullet} \overset{-1}{\bullet}$ GDD 9 of Row 17.
 is empty.

Case (iii) Omitting 1 and adding 3.

checking step by step we have to consider following cases


(a) $\overset{-1}{\bullet} \overset{q^{-1}q}{\bullet} \overset{-q^{-1}q}{\bullet} \overset{-q^{-1}q}{\bullet}$ (see GDD 5 of Row 16 in Table A2 ur by Lemma 3.2 (I)(b)).

Case (iv) Containing circle with four vertexes.


According to Lemma 3.39 (ii) we have to consider following cases.

(a') to (a). empty.

(a) to (a) or (11)




with $-q^{-1} = -q^2$ is an arithmetic root system by Lemma 3.3 (II)(f).




is affine.

(b) to (a) empty. (d) to (a) empty. (f) to (a) empty. (g) to (a). empty.

(h) to (a) or (12)



with $-q = r$ is an arithmetic root system by Type 7.




is affine.

(m) to (a), empty.

4.40 GDD 2 of Row 16


Case (i) Omitting 3 and adding 1.

According to Lemma 3.3 (IV) we have to consider following cases.

$$q \begin{array}{c} (a') \text{ or } (1). \\ q^{-1} \quad -1 \quad q \quad -1 \quad -q^{-1}-q \end{array}$$



, $q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 4.

is affine by Lemma 3.15.

$$q \begin{array}{c} (a'') \text{ or } (2). \\ q^{-2} \quad -1 \quad q^2 \quad -1 \quad -q^{-2}-q^2 \end{array}$$



$q^2 \in R_3, q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 5.

is affine by Lemma 3.4.

$$q^2 \begin{array}{c} (a) \text{ or } (3). \\ q^{-2} \quad -1 \quad q \quad -1 \quad -q^{-1}-q \end{array}$$



, $q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 6.

is affine by Lemma 3.7.

$$q^3 \begin{array}{c} (c). \\ q^{-3} \quad -1 \quad q \quad -1 \quad -q^{-1}-q \end{array}$$



, $q \in F^* \setminus \{1, -1\}$. $q \notin R_3$, GDD 2 of Row 7.

is empty.

$$-1 \begin{array}{c} (e) \text{ or } (4). \\ q^{-1} \quad -1 \quad q \quad -1 \quad -q^{-1}-q \end{array}$$



, $q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 8.

is affine by Lemma 3.15.

$$q \begin{array}{c} (j). \\ -q \quad -1 \quad -q^{-1} \quad 1 \quad q \quad q^{-1} \end{array}$$



, $q \in R_3$. GDD 2 of Row 14.

is empty.

$$-1 \begin{array}{c} (k) \text{ or } (5). \\ q \quad -1 \quad q \quad -1 \quad -q^{-1}-q \end{array}$$



, $q \in R_3$. GDD 2 of Row 15.

is affine by Lemma 3.15.

$$-q \begin{array}{c} (l) \text{ or } (6). \\ -q^{-1} \quad 1 \quad q \quad -1 \quad -q^{-1}-q \end{array}$$


, $q \in R_3$. GDD 2 of Row 16.

is affine by Lemma 3.15.

$$-1 \begin{array}{c} (n). \\ q \quad -1 \quad -1 \quad -1 \end{array}$$


is empty.

(n') or (7).

$$\begin{array}{c} -1 \quad -1 \quad -1 \quad q \quad -1 \quad -q^{-1}-q \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad , q \in R_3. \text{ GDD 1 of Row 17.}$$

is affine by Lemma 3.15.

$$\begin{array}{c} \text{(o) or (8).} \\ -1 \quad -q \quad -1 \quad q \quad -1 \quad -q^{-1}-q \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad , q \in R_3. \text{ GDD 4 of Row 17.}$$

is affine by Lemma 3.15.

$$\begin{array}{c} \text{(p).} \\ -1 \quad q \quad -1 \quad -q \quad -1 \quad q^{-1} \quad q \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad , q \in R_3. \text{ GDD 4 of Row 17.}$$

is empty.

$$\begin{array}{c} \text{(r).} \\ q^{-1} \quad -q^{-1}-1 \quad -1 \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}$$

is empty.

Case (ii) Omitting 3 and adding 2.

According to Lemma 3.3 (IV) we have to consider following cases.

$$\text{(a')} \text{ or (9).} \quad \begin{array}{c} \bullet \\ | \\ -q \\ -q^{-1} \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ q \quad q^{-1} \quad -1 \quad q \quad -1 \end{array} \quad , q \in F^* \setminus \{1, -1\}. \text{ GDD 2 of Row 4.}$$

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2 (IV). Whole is affine by Lemma 3.12.

$$\text{(a'')} \quad \begin{array}{c} \bullet \\ | \\ -q^2 \\ -q^{-2} \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ q \quad q^{-2} \quad -1 \quad q^2 \quad -1 \end{array} \quad , q \in F^* \setminus \{1, -1\}. \text{ GDD 2 of Row 5.}$$

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(I).

$$\text{(a)} \text{ or (10).} \quad \begin{array}{c} \bullet \\ | \\ -q \\ -q^{-1} \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ q^2 \quad q^{-2} \quad -1 \quad q \quad -1 \end{array} \quad , q \in F^* \setminus \{1, -1\}. \text{ GDD 2 of Row 6.}$$

Omitting 1 in whole is an arithmetic root system by Lemma 3.2 (IV). Whole is affine by Lemma 3.12.

$$\text{(c).} \quad \begin{array}{c} \bullet \\ | \\ -q \\ -q^{-1} \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \\ q^3 \quad q^{-3} \quad -1 \quad q \quad -1 \end{array} \quad , q \in F^* \setminus \{1, -1\}. \quad q \notin R_3, \text{ GDD 2 of Row 7.}$$

is empty.

(e).  , $q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 8.

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(II).

(j).  , $q \in R_3$. GDD 2 of Row 14.
is empty.

(k) or (11).  , $q \in R_3$. GDD 2 of Row 15.

Omitting 1 in whole is an arithmetic root system by Lemma 3.2(II)(e). Whole is affine by Lemma 3.12.

(l) or (12).  , $q \in R_3$. GDD 2 of Row 16.

Omitting 1 in whole is an arithmetic root system by Lemma 3.2 (IV). Whole is affine by Lemma 3.12


(n).  , $q \in R_3$. GDD 1 of Row 17.
is empty.

(n') or (13).  , $q \in R_3$. GDD 1 of Row 17.


Omitting 1 in whole is an arithmetic root system by GDD 1 of Row 9 or by Lemma 3.2 (IV). Whole is affine by Lemma 3.12.

(o) or (14).  , $q \in R_3$. GDD 4 of Row 17.

Omitting 1 in whole is an arithmetic root system by Type 7. Whole is affine by Lemma 3.12.




$q \in R_3$, GDD 4 of Row 17.




, $q \in R_3$. GDD 9 of Row 17.

Case (iii) Omitting 1 and adding 3.


According to Lemma 3.3 (III) we have to consider following cases.




$q = -\xi, \xi \in R_3, q \in R_6$. GDD 1 of Row 6.




$q = -\xi, \xi \in R_3, q \in F^* \setminus \{1, -1\}, q \notin R_3$. GDD 1 of Row 7.



$q = -\xi, \xi \in R_3, q \in F^* \setminus \{1, -1\}$. GDD 3 of Row 8.




, $q \in R_3$. GDD 1 of Row 14.




, $q \in R_3$. GDD 1 of Row 15.

is empty.




, $q \in R_3$. GDD 1 of Row 16.

is empty.




, $q \in R_3$. GDD 2 of Row 17.

is empty.



, $q \in R_3$. GDD 5 of Row 17.



, $q \in R_3$. GDD 7 of Row 17.

is empty.

$$\begin{array}{ccccccc} & (a') & & & & & \\ q & q^{-1} & q & q^{-1}-1 & -q & -1 & \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

GDD 1 of Row 4.

$$\begin{array}{ccccccc} & (a'') & & & & & \\ q & q^{-2} & q^2 & q^{-2}-1 & -q^2 & -1 & \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

GDD 1 of Row 5.

$$\begin{array}{ccccccc} & (k') & & & & & \\ -1 & q & -1 & -1 & -1 & & \\ \bullet & \bullet & \bullet & \bullet & \bullet & & \bullet \end{array}$$

GDD 1 of Row 17.

is empty.

$$\begin{array}{ccccccc} & (p') & & & & & \\ q^{-1} & -q^{-1} & 1 & -1 & -1 & & \\ \bullet & \bullet & \bullet & \bullet & \bullet & & \bullet \end{array}$$

GDD 9 of Row 17.

is empty.

Case (iv) Containing circle with four vertexes.

According to Lemma 3.39 (ii) we have to consider following cases.

(a') to (a). empty. (a') to (c). empty. (a') to (e) empty. (a') to (g). empty. (a') to (n) empty. (a') to (a'). empty. (a') to (a'') empty.

(a'') to (a)

$$\begin{array}{ccccccc} & & & & & & \\ -1 & q^{-2} & q & q^{-4}-q^2 & & & \\ \bullet & \bullet & \bullet & \bullet & & & \bullet \end{array}$$

with $q = q^4$ is not an arithmetic root system by Lemma 3.2(I).

(a'') to (c) empty. (a'') to (e) empty. (a'') to (g) empty. (a'') to (n) empty. (a'') to (a'). empty. (a'') to (a''), empty.

(a) to (a) or (15).

$$\begin{array}{ccccccc} & & & & & & \\ -1 & q^{-2} & q^2 & q^{-2}-q & & & \\ \bullet & \bullet & \bullet & \bullet & & & \bullet \\ & & q^2 & & & & \\ & q^{-2} & & q^{-2} & & & \\ \bullet & \bullet & \bullet & \bullet & & & \bullet \\ -1 & q & -1 & -q^{-1}q & & & \end{array}$$

is an arithmetic root system by Type 2.

is affine.

(a) to (c) empty. (a) to (e) empty. (a) to (g) empty. (a) to (n) empty. (a) to (a'). empty. (a) to (a''). empty.

(e) to (a) empty. (e) to (c)

$$\begin{array}{ccccccc} & & & & & & \\ -1 & q^{-1}-1 & -q^{-3}q & & & & \\ \bullet & \bullet & \bullet & \bullet & & & \bullet \end{array}$$

is not an arithmetic root system by Lemma 3.2 (I).


(e) to (e)

$$\begin{array}{ccccccc} & & & & & & \\ -1 & q^{-1}-1 & -q^{-1}q & & & & \\ \bullet & \bullet & \bullet & \bullet & & & \bullet \end{array}$$

is not an arithmetic root system by Lemma 3.2 (X).

(e) to (g) empty.

(e) to (n) or (16)




is an arithmetic root system by Type 3.

is affine.

(e) to (a') empty. (e) to (a'') empty.


(k) to (a') empty. (k) to (a'') empty.

(k) to (a) empty. (k) to (c)




with $-q^3 = -1$ is not an arithmetic root system by Lemma 3.2 (I).

(k) to (e) or (17).




is an arithmetic root system by Lemma 3.2 (II)(e) or by Lemma 3.2 (X)(d).



is affine.

(k) to (g) empty.


(k) to (n)




is not an arithmetic root system by Lemma 3.2(I).

(k) to (a') empty. (k) to (a''). empty. (l) to (a) empty. (l) to (c) empty. (l) to (e) empty. (l) to (g) empty. (l) to (n) empty.

(l) to (a') or (18).




is an arithmetic root system,



is affine.


(l) to (a''). empty. (n') to (a) empty.

(n') to (c).




with $-q^{-3} = -1$ is not an arithmetic root system by Lemma 3.2 (I).

(n') to (e) or (19).




is an arithmetic root system by Lemma 3.2 (IV).



is affine.

(n') to (g) empty.

(n') to (n).




is not an arithmetic root system by Lemma 3.2 (I).

(n') to (a') empty. (n') to (a'') empty.

(o) to (a).


empty.

(o) to (c)




with $-1 = -q^3$ is not an arithmetic root system by Lemma 3.2(I).

(o) to (e) or (20)




is an arithmetic root system by Type 7.



is affine.

(o) to (g) empty.

(o) to (n).




is not an arithmetic root system by Lemma 3.2(I).

(o) to (a') empty. (o) to (a'') empty.

4.41 GDD 3 of Row 16

Case (i) Omitting 3 and adding 1.


According to Lemma 3.3 (III) we have to consider following cases.



, $q \in R_3$. GDD 1 of Row 6.


Omitting 2 in whole is not an arithmetic root system by Lemma 3.2 (V).

(b).




, $q \in R_3$. GDD 3 of Row 6.

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(II) or by Lemma 3.2 (XI).



, $q \in F^* \setminus \{1, -1\}$, $q \notin R_3$. GDD 1 of Row 7.


is empty.



, $q \in F^* \setminus \{1, -1\}$. GDD 3 of Row 8.


Omitting 2 in whole is an arithmetic root system by GDD 2 of Row 17 or by Lemma 3.3 (III)(1). Whole is affine by Lemma 3.4.

(g).



, $q \in R_3$. GDD 1 of Row 14.


is empty.



, $q \in R_3$. GDD 1 of Row 15.


Omitting 2 in whole is not an arithmetic root system by Lemma 3.2 (V).

(i).



, $q \in R_3$. GDD 3 of Row 15.


Omitting 2 in whole is not an arithmetic root system by Lemma 3.3(II).



, $q \in R_3$. GDD 1 of Row 16.


Omitting 2 in whole is not an arithmetic root system by Lemma 3.2 (V).

(k).



, $q \in R_3$. GDD 3 of Row 16.


Omitting 2 in whole is not an arithmetic root system by Lemma 3.2 (V).



, $q \in R_3$. GDD 2 of Row 17.

Omitting 2 in whole is not an arithmetic root system by Lemma 3.2 (V).

(m).



Omitting 2 in whole is not an arithmetic root system by Lemma 3.2 (V).




is empty.



Omitting 2 in whole is not an arithmetic root system by Lemma 3.2 (V),

(p).



Omitting 2 in whole is not an arithmetic root system by Lemma 3.2 (V).



Omitting 2 in whole is not an arithmetic root system by Lemma 3.3(II),




Omitting 2 in whole is not an arithmetic root system by Lemma 3.2 (XIV).



(p')

is empty.




GDD 9 of Row 17.

is empty.


Case (ii) Omitting 3 and adding 2.

According to Lemma 3.3 (I) we have to consider following cases.




GDD 2 of Row 4.

Omitting 1 in whole is an arithmetic root system by Lemma 3.2(II)(g),
Whole is affine by Lemma 3.4.



GDD 3 of Row 5.


Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(I).



GDD 2 of Row 6.


Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(II).

(c)




GDD 3 of Row 6.

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(II).




GDD 2 of Row 7.

is empty.



, GDD 4 of Row 7.

is empty.



, GDD 1 of Row 8.

Omitting 1 in whole is an arithmetic root system by Lemma 3.2 (X)(b). Whole is affine by Lemma 3.4.

(h) or (4)  , GDD 1 of Row 9.

Omitting 1 in whole is an arithmetic root system when $(-q)^3 = r$ or $(-q)^2 = r$ or $-q = r$ by Lemma 3.2(II) and Type 7. Whole is affine when $(-q)^3 = r$ or $q^2 = r$ by Lemma 3.4. whole is an arithmetic root system when $-q = r$ by Lemma 3.4(f).

(i')  , $q \notin R_3$. GDD 2 of Row 10.

is empty.

(k)  , $q \notin R_3$. GDD 1 of Row 10.

is empty.

(l)  , $q \notin R_3$. GDD 2 of Row 10.

is empty.


(m)  , $q \in R_3$. GDD 1 of Row 11.

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(II).

(n)  , $q \in R_3$. GDD 3 of Row 14.

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(I).

(o)


 GDD 3 of Row 15.

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(II).

(p) or (5)  , $q \in R_3$. GDD 2 of Row 16.

Omitting 1 in whole is an arithmetic root system by Lemma 3.2(II)(d). Whole is affine by Lemma 3.4.


(q)




Omitting 2 in whole is not an arithmetic root system by Lemma 3.2 (V).

(r)  , $q \in R_3$. GDD 4 of Row 16.

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(I).


(s')  , $q \in R_3$. GDD 1 of Row 17.
is empty.

(s)



Omitting 2 in whole is not an arithmetic root system by Lemma 3.2 (V).

(t)




Omitting 2 in whole is not an arithmetic root system by Lemma 3.2 (V).

Case (iii) Omitting 2 and adding 3.


Checking step by step we have to consider following cases.

(a) or (6)



Omitting 1 in whole is an arithmetic root system by Type 7. Whole is affine by Lemma 3.4.

(b)



Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(II).


Case (iv) Containing circle with four vertexes.

This case can be ignored since it is over a cycle.

Case (v) complete diagram containing circle with four vertexes.


According to Lemma 3.40 we have to consider following cases.

(b) in Case (i)




with $q \in R_3$. If 134 is an arithmetic root system then $\tilde{q}_{34} = -q$ by Lemma 3.2(I)(k). 234 is not an arithmetic root system since it is not GDD 3 of Row 9 and by Type 6.


(i) in Case (i)




134 is not an arithmetic root system by Lemma 3.2(I).
 (k) in Case (i)




134 is not an arithmetic root system by Lemma 3.2(I).
 (m) in Case (i)



134 is not an arithmetic root system by Lemma 3.2(I).
 (p) in Case (i)




134 is not an arithmetic root system by Lemma 3.2(I).
 (c) in Case (ii)



with $q \in R_3$.


If 134 is an arithmetic root system, then $\tilde{q}_{34} = -q$ by Lemma 3.2(I). 234 is not an arithmetic root system checked step by step.

(o) in Case (ii)




134 is not an arithmetic root system by Lemma 3.2(I).

(q) in Case (ii)




134 is not an arithmetic root system by Lemma 3.2(I).

(s) in Case (ii)




134 is not an arithmetic root system by Lemma 3.2(I).

(t) in Case (ii)



134 is not an arithmetic root system by Lemma 3.2(I).

(b) in Case (iii)



with $q \in R_3$.

If 234 is an arithn then $\tilde{q}_{24} = (-q)^{-2}$ by Type 6. If 124 is an arithmetic root system, then $\tilde{q}_{24} = q^2$. Consequently, $q^4 = 1$, which implies a contradiction.

4.42 GDD 4 of Row 16

Case (i) Omitting 3 and adding 1.


checking step by step we have to consider following cases



is affine by Lemma 3.7.

Case (ii) Omitting 3 and adding 2.

According to Lemma 3.2 (I) we have to consider following cases.



Omitting 1 in whole is an arithmetic root system by Lemma 3.2 (IV). Whole is affine by Lemma 3.4.

Case (iii) Omitting 1 and adding 3.

According to Lemma 3.3 (III) we have to consider following cases.

$$\begin{array}{ccccccc} q^2 & \text{(a).} & & & & & \\ q^{-2} & q & q^{-1} & -1 & -1 & -q & \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

$$q = -\xi, \xi \in R_3, \text{ GDD 1 of Row 6.}$$

$$\begin{array}{ccccccc} q^3 & \text{(c).} & & & & & \\ q^{-3} & q & q^{-1} & -1 & -1 & -q & \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

$$q = -\xi, \xi \in R_3, , q \notin R_3. \text{ GDD 1 of Row 7.}$$

$$\begin{array}{ccccccc} -1 & \text{(e).} & & & & & \\ q^{-1} & q & q^{-1} & -1 & -1 & -q & \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

$$q = -\xi, \xi \in R_3, \text{ GDD 3 of Row 8.}$$

$$\begin{array}{ccccccc} q & \text{(g).} & & & & & \\ -q & -q^{-1} & -1 & -1 & q^{-1} & & \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

$$, q \in R_3. \text{ GDD 1 of Row 14.}$$

$$\begin{array}{ccccccc} -1 & \text{(h).} & & & & & \\ q & q & q^{-1} & -1 & -1 & -q & \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

$$q = -\xi, \xi \in R_3, \text{ GDD 1 of Row 15.}$$

is empty.

$$\begin{array}{ccccccc} -q & \text{(j).} & & & & & \\ -q^{-1} & q & q^{-1} & -1 & -1 & -q & \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

$$q = -\xi, \xi \in R_3, \text{ GDD 1 of Row 16.}$$

is empty.

$$\begin{array}{ccccccc} -1 & \text{(l).} & & & & & \\ -1 & q & q^{-1} & -1 & -1 & -q & \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

$$q = -\xi, \xi \in R_3, \text{ GDD 2 of Row 17.}$$

is empty.

$$\begin{array}{ccccccc} -1 & \text{(n).} & & & & & \\ q & -q & -q^{-1} & -1 & -1 & q & \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

$$q \in R_3, \text{ GDD 5 of Row 17.}$$

$$\begin{array}{ccccccc} -1 & \text{(o).} & & & & & \\ -q & q & q^{-1} & -1 & -1 & -q & \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$


$$q = -\xi, \xi \in R_3, \text{ GDD 7 of Row 17.}$$

is empty.


$$\begin{array}{ccccccc} q & \text{(a').} & & & & & \\ q^{-1} & q & q^{-1} & -1 & -1 & -q & \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

$$q = -\xi, \xi \in R_3, \text{ GDD 1 of Row 4.}$$


(a'')



$q = -\xi, \xi \in R_3$, GDD 1 of Row 5.



is empty.




is empty.

Case (iv) containing circle with four vertexes.

According to Lemma 3.39 (ii) we have to consider following cases.


(a) to (a) empty.

(a) to (c)




with $-1 = -q^3$ is not an arithmetic root system by Lemma 3.2(I).

(a) to (e) or (3)




is an arithmetic root system by Lemma 3.2 (IV).




is affine.

(a) to (g) empty.

(a) to (n) or (4).



is an arithmetic root system by Type 3.



is affine,

(a) to (a') empty. (a) to (a'') empty.

4.43 GDD 5 of Row 16

Case (i) Omitting 3 and adding 1.

checking step by step we have to consider following cases

(a) or (1)

$$\begin{array}{c} -1 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad \begin{array}{l} q^{-1}q \quad -q^{-1}q \quad q^{-1}q \\ \text{(see, } q \in R_3, \text{ GDD 1 of Row 16 in Table A2).} \end{array}$$

is affine by Lemma 3.33

Case (ii) Omitting 3 and adding 2.

Checking step by step we have to consider following cases.

(a)

$$\begin{array}{c} \bullet \text{---} \bullet \begin{array}{l} \nearrow \bullet \\ \searrow \bullet \end{array} \\ \begin{array}{l} -q^{-1} \quad -q^{-1}q \\ -q \quad -q^{-1} \end{array} \end{array} \quad \begin{array}{l} \bullet \\ \bullet \end{array} \quad \begin{array}{l} -q^{-1}q \\ -q \end{array} \quad \text{(see GDD 5 of Row 16.)}$$

Omitting 1 in whole is an arithmetic root system by GDD 5 of Row 16. Whole is affine by Lemma 3.4.

Case (iii) Omitting 1 and adding 3.

According to Lemma 3.3 (II) we have to consider following cases.

(a).

$$\begin{array}{c} q \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad \begin{array}{l} q^{-1}q \quad q^{-1}q \quad q^{-1} \quad -q \\ q = -\xi, \xi \in R_3, q \in F^* \setminus \{1\}. \text{ GDD 1 of Row 1.} \end{array}$$

(b).

$$\begin{array}{c} q \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad \begin{array}{l} q^{-2}q^2 \quad q^{-2}q^2 \quad q^{-2} \quad -q^2 \\ q^2 = -\xi, \xi \in R_3, q^2 \in R_6. \text{ GDD 1 of Row 2.} \end{array}$$

(c).

$$\begin{array}{c} q^2 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad \begin{array}{l} q^{-2}q \quad q^{-1}q \quad q^{-1} \quad -q \\ q = -\xi, \xi \in R_3, q \in F^* \setminus \{1, -1\}. \text{ GDD 1 of Row 3.} \end{array}$$

(d).

$$\begin{array}{c} -1 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad \begin{array}{l} q^{-1}q \quad q^{-1}q \quad q^{-1} \quad -q \\ q = -\xi, \xi \in R_3, q \in F^* \setminus \{1, -1\}. \text{ GDD 1 of Row 4.} \end{array}$$


(e).

$$\begin{array}{c} q \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad \begin{array}{l} -q^{-1} \quad -q^{-1} \quad -q^{-1} \quad q^{-1} \\ , q \in R_3. \text{ GDD 1 of Row 12.} \end{array}$$

(f).

$$\begin{array}{c} -1 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad \begin{array}{l} q^{-2}q \quad q^{-1}q \quad q^{-1} \quad -q \\ q = -\xi, \xi \in R_3, q \in R_3 \cup R_6. \text{ GDD 1 of Row 13.} \end{array}$$


(g).



$q = -\xi, \xi \in R_3, q \in R_6$. GDD 3 of Row 15.


empty

(h).



$q \in R_3$. GDD 5 of Row 16.

(i).



$q = -\xi, \xi \in R_9$. GDD 1 of Row 18.


empty

Case (iv) containing circle with four vertexes.


According to Lemma 3.39 (ii) we have to consider following cases.

(a) to (a). empty.

(a) to (b) empty.(a) to (c) empty.(a) to (d) or (2).




is an arithmetic root system by Lemma 3.2 (IV).




is affine.

(a) to (e) empty.

(a) to (f) or (3)



is an arithmetic root system by Type 3.



is affine.


(a) to (h) empty.

4.44 GDD 1 of Row 17

Case (i) Omitting 3 and adding 1.


According to Lemma 3.3 (V) we have to consider following cases.

(a) or (1)




($r \in R_4$, GDD 2 of Row 6; $r \in R_3$, GDD 1 of Row 17;
 $r \in R_6$, GDD 2 of Row 7; $r \in R_2$, GDD 1 of Row 1;).


is affine by Lemma 3.5.




is affine by Lemma 3.4



is affine by Lemma 3.5




is affine by Lemma 3.5




is affine by Lemma 3.5

Case (ii) Omitting 3 and adding 2.


According to Lemma 3.3 (V) we have to consider following cases.




Omitting 1 in whole is an arithmetic root system when $r = q$ and $q \in R_3$, or $r = q^{-1}$, or $r = -q$ and $r \in R_6$ by Lemma 3.2 (XI) or $r = -1$. Whole is affine by Lemma 3.9.




Omitting 1 in whole is an arithmetic root system when $q = -r$ by Type 4. Whole is affine when $q = -r$ by Lemma 3.9.

(e) or (8)  (see, $r \in R_6$, GDD 4 of Row 7 in Table A2).

Omitting 1 in whole is an arithmetic root system when $q = r^{-2}$ by Type 2. Omitting 1 in whole is not an arithmetic root system when $q \neq r^{-2}$ by Lemma 3.2(I). Whole is affine when $q = r^{-2}$, $r \in R_6$ and $q \in R_3$, by Lemma 3.26.

(f) or (9)  (see, $r^2 \neq 1$, GDD 1 of Row 9 in Table A2).


Omitting 1 in whole is an arithmetic root system when $r = q$ or $r = q^2$ or $r = -q$ by Lemma 3.2 (X). Whole is affine by Lemma 3.26.

(g)  (see, $q \in R_3$, $r \in R_4$, GDD 1 of Row 2 in Table A2).

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(I).


Case (iii) Omitting 1 and adding 3.

According to Lemma 3.3 (IV) we have to consider following cases.

(a').  , $q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 4.

(a'').  , $q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 5.

(a).  , $q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 6.

(c).  , $q \in F^* \setminus \{1, -1\}$. $q \notin R_3$, GDD 2 of Row 7.

empty

(e).  , $q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 8.

(j).

$$\begin{array}{ccccccc}
 q & -q & -1 & -q^{-1} & -1 & -1 & -1 \\
 \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet
 \end{array}$$

, $q \in R_3$. GDD 2 of Row 14.

empty

$$\begin{array}{ccccccc}
 (k). \\
 -1 & -1 & -1 & -1 & -1 & -1 & -1 \\
 q & q & q & q & q & q & q \\
 \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet
 \end{array}$$

, $q \in R_3$. GDD 2 of Row 15.

$$\begin{array}{ccccccc}
 (l). \\
 -q & -q^{-1} & -1 & -1 & -1 & -1 & -1 \\
 q & q & q & q & q & q & q \\
 \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet
 \end{array}$$

, $q \in R_3$. GDD 2 of Row 16.

$$\begin{array}{ccccccc}
 (n). \\
 -1 & -1 & -1 & -1 & -1 & -1 & -1 \\
 q & q & q & q & q & q & q \\
 \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet
 \end{array}$$

, $q \in R_3$. GDD 1 of Row 17.

empty

$$\begin{array}{ccccccc}
 (n'). \\
 -1 & -1 & -1 & -1 & -1 & -1 & -1 \\
 -1 & -1 & -1 & -1 & -1 & -1 & -1 \\
 q & q & q & q & q & q & q \\
 \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet
 \end{array}$$

, $q \in R_3$. GDD 1 of Row 17.

$$\begin{array}{ccccccc}
 (o). \\
 -1 & -1 & -1 & -1 & -1 & -1 & -1 \\
 -q & -q & -q & -q & -q & -q & -q \\
 q & q & q & q & q & q & q \\
 \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet
 \end{array}$$

, $q \in R_3$. GDD 4 of Row 17.

$$\begin{array}{ccccccc}
 (p). \\
 -1 & -1 & -1 & -1 & -1 & -1 & -1 \\
 q & q & q & q & q & q & q \\
 -q & -q & -q & -q & -q & -q & -q \\
 \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet
 \end{array}$$

, $q \in R_3$. GDD 4 of Row 17.

empty

$$\begin{array}{ccccccc}
 (r). \\
 q^{-1} & -q^{-1} & -1 & -1 & -1 & -1 & -1 \\
 -q^{-1} & -q^{-1} & -q^{-1} & -q^{-1} & -q^{-1} & -q^{-1} & -q^{-1} \\
 \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet
 \end{array}$$

, $q \in R_3$. GDD 9 of Row 17.

empty


Case (iv) containing circle with four vertexes.

According to Lemma 3.39 (ii) we have to consider following cases.

(a) to (a') empty. (a) to (a'') empty. (a) to (a) empty. (a) to (e) or (10).


$$\begin{array}{ccccccc}
 -1 & -1 & -1 & -1 & -1 & -1 & -1 \\
 r & r & r & r & r & r & r \\
 \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet
 \end{array}$$

is an arithmetic root system when $r = q^{-1}$, or $r = q$, or $r = -q^{-1}$, or $r = -1$ by Lemma 3.2 (XI) .




is affine.
 (a) to (k) or (11).


is an arithmetic root system when $r = q$, or $r = q^{-1}$, or $r = -q$, or $r = -1$ by Lemma 3.2 (XI).




is affine.
 (a) to (l) empty. (a) to (n') or (12)




is an arithmetic root system when $r \in R_2 \cup R_3 \cup R_4 \cup R_6$ by Lemma 3.2 (XI)(e).




is affine.
 (a) to (o) or (13).




is an arithmetic root system when $r = -q^{-1}$, or $r = q$ by Lemma 3.2 (XI).



is affine.
 (b) to (a') or (14).




and $q = r^{-1}$ is an arithmetic root system by Lemma 3.2 (IX)(f) or by Lemma 3.3 (III)(o).



is affine.


(b) to (a'') or (15).




with $q = r^{-2}$ is an arithmetic root system by Lemma 3.2 (V)(c).

is affine.

(b) to (a) or (16)




with $q^2 = r^{-1}$ is an arithmetic root system by Lemma 3.2 (IX)(f) or by Lemma 3.3 (III)(o).



is affine.

(b) to (e) empty. (b) to (k) empty.


(b) to (l).




with $-q = r^{-1}$ is empty because of $q \in R_3$ and $r \in R_3$.

(b) to (n') empty. (b) to (o) empty.

(e) to (a') or (17)




with $q = -r^{-1}$ is an arithmetic root system by Lemma 3.3 (III)(h).




is affine.

(e) to (a'') empty.

(e) to (a)



with $q^2 = -r^{-1}$ is an arithmetic root system by Lemma 3.3 (III). (h) or (1) or (18)




is affine.


(e) to (e) empty.(e) to (k) empty.(e) to (l) empty.

(e) to (n') empty.(e) to (o) empty.

(f) to (a') or (19)




with $q = r$ is an arithmetic root system.




is affine.

(f) to (a'') or (20)




with $q = r^2$ is an arithmetic root system by Lemma 3.3 (III)(h).




is affine.

(f) to (a) or (21)



with $r = q^2$ is an arithmetic root system by Type 7.




is affine.


(f) to (e) empty.

(f) to (k) empty.

(f) to (l) or (22)



with $r = -q$ is an arithmetic root system by Type 7.



is affine.

(f) to (n') empty.

(f) to (o) empty.

(g) to (a') empty.(g) to (a'') empty.

(g) to (a) empty.

(g) to (e) empty.(g) to (k) empty.(g) to (l) empty.(g) to (n') empty. (g) to (o) empty.

4.45 GDD 2 of Row 17

Case (i) Omitting 3 and adding 1.

Checking step by step we have to consider following cases.

(a) or (1)

$$\begin{array}{ccccccc} -q & -q^{-1} & -1 & -1 & q & q^{-1} & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array} \quad (\text{ see } q \in R_3, \text{ GDD 4 of Row 16 in Table A2}).$$

is affine by Lemma 3.7

(b)

$$\begin{array}{ccccccc} -1 & -1 & -1 & -1 & q & q^{-1} & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array} \quad (\text{ see, } q \in R_4, \text{ GDD 1 of Row 2 in Table A2}).$$

is empty.

Case (ii) Omitting 3 and adding 2.

checking step by step we have to consider following cases

(a) or (2)

$$\begin{array}{ccccccc} & & & & q^{-1} & -1 & \\ & & & & \bullet & \bullet & \\ -1 & q^{-1} & & & / & & \\ \bullet & \bullet & & & \bullet & & \\ q & & & & \backslash & & \\ & & & & -1 & -1 & \\ & & & & \bullet & \bullet & \end{array} \quad , (q \in R_3, \text{ see GDD 2 of Row 17}).$$

Omitting 1 in whole is an arithmetic root system by Type 7. Whole is affine by Lemma 3.4.

(b)

$$\begin{array}{ccccccc} -1 & q^{-1} & q^{-1} & -1 & & & \\ \bullet & \bullet & \bullet & \bullet & & & \\ -1 & q^{-1}q & & & / & & \\ \bullet & \bullet & & & \bullet & & \\ & & & & \backslash & & \\ & & & & -1 & -1 & \\ & & & & \bullet & \bullet & \end{array} \quad (q \in R_4, \text{ see GDD 1 of Row 6 in Table A2 or Type 1.})$$

is empty.

(c)

$$\begin{array}{ccccccc} & & & & q^{-1} & -1 & \\ & & & & \bullet & \bullet & \\ q & q^{-1}q & & & / & & \\ \bullet & \bullet & & & \bullet & & \\ & & & & \backslash & & \\ & & & & -1 & -1 & \\ & & & & \bullet & \bullet & \end{array} \quad (q \in R_4, \text{ see Type 1.})$$

is empty.

Case (iii) Omitting 1 and adding 3.

According to Lemma 3.3 (I) we have to consider following cases.

(a')

$$\begin{array}{ccccccc} -1 & q & -1 & q^{-1}q & -1 & -1 & \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array} \quad \text{GDD 2 of Row 4.}$$

(a)

$$\begin{array}{ccccccc} -q & q^2 & -1 & q^{-2}q^2 & -1 & -1 & \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array} \quad \text{GDD 3 of Row 5.}$$

(b)

$$\begin{array}{ccccccc} -1 & q & -1 & q^{-2}q^2 & -1 & -1 & \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array} \quad \text{GDD 2 of Row 6.}$$

(d)

$$\begin{array}{ccccccc} -1 & q & -1 & q^{-3}q^3 & -1 & -1 & \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array} \quad \text{GDD 2 of Row 7.}$$

is empty.

$$\begin{array}{ccccccc} -q & \overset{(f)}{q^2} & -1 & q^{-3}q^3 & -1 & -1 & \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$
 , GDD 4 of Row 7.
 is empty.

$$\begin{array}{ccccccc} q^{-1} & \overset{(g)}{q} & -1 & q^{-1}q & -1 & -1 & \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$
 , GDD 1 of Row 8.

$$\begin{array}{ccccccc} r & \overset{(h)}{r^{-1}} & -1 & q^{-1}q & -1 & -1 & \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$
 , GDD 1 of Row 9.

$$\begin{array}{ccccccc} q & \overset{(i')}{q^{-1}} & -1 & q^2 & q^{-2} & -1 & -1 \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$
 , $q \notin R_3$. GDD 2 of Row 10.
 is empty.

$$\begin{array}{ccccccc} q & \overset{(k)}{q^{-1}} & -1 & q^{-1}q & -1 & -1 & \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$
 , $q \notin R_3$. GDD 1 of Row 10.
 is empty.

$$\begin{array}{ccccccc} q^{-2} & \overset{(l)}{q^2} & -1 & q^{-1}q & -1 & -1 & \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$
 , $q \notin R_3$. GDD 2 of Row 10.
 is empty.

$$\begin{array}{ccccccc} q & \overset{(m)}{q^{-1}} & -1 & q^{-1}q & -1 & -1 & \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$
 , $q \in R_3$. GDD 1 of Row 11.

$$\begin{array}{ccccccc} q^{-1} & \overset{(n)}{-q^{-1}q} & -1 & -q & -q^{-1} & -1 & -1 \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$
 , $q \in R_3$. GDD 3 of Row 14.
 is empty.

$$\begin{array}{ccccccc} -1 & \overset{(p)}{q} & -1 & -q^{-1}q & -1 & -1 & \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$
 , $q \in R_3$. GDD 2 of Row 16.
 is empty.

$$\begin{array}{ccccccc} q & \overset{(r)}{-1} & -1 & -q^{-1}q & -1 & -1 & \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$
 , $q \in R_3$. GDD 4 of Row 16.
 is empty.


$$\begin{array}{ccccccc} -1 & \overset{(s')}{q} & -1 & -1 & -1 & -1 & -1 \\ \bullet & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \end{array}$$
 , $q \in R_3$. GDD 1 of Row 17.
 is empty.

Case (iv) containing circle with four vertexes.


According to Lemma 3.39 (ii) we have to consider following cases.

(a) to (a') empty.

(a) to (a) or (3).




with $-q^{-1} = -q^2$ is an arithmetic root system by Lemma 3.3 (III)(n).




is affine.

(a) to (b) empty. (a) to (g) empty.

(a) to (h) or (4).



with $r = -q$ is an arithmetic root system by Type 7.



is affine.


(a) to (m) .empty.

4.46 GDD 3 of Row 17

Case (i) Omitting 3 and adding 1.


Checking step by step we have to consider following cases.

(a) or (1)



Omitting 2 in whole is an arithmetic root system by Type 7. Whole is affine by Lemma 3.4.

(b) or (2)




, (see $q \in R_3$, GDD 3 of Row 17 in Table A2.)

Omitting 1 in whole is an arithmetic root system. Omitting 2 in whole is an arithmetic root system. Whole is affine.

Case (ii) Omitting 3 and adding 2.

Checking step by step we have to consider following cases.

(b) or (3)




, (see $q \in R_3$, GDD 3 of Row 17 in Table A2.)

Omitting 2 in whole is an arithmetic root system. Omitting 1 in whole is an arithmetic root system. Whole is affine.

Case (iii) Omitting 2 and adding 3.

According to Lemma 3.3 (I) we have to consider following cases.


(a') or (4)



$q = -\xi, \xi \in R_3$, GDD 2 of Row 4.

Omitting 2 in whole is an arithmetic root system by Lemma 3.2(II)(e) or by Lemma 3.2 (X)(d). Whole is affine by Lemma 3.4.


(a)



$q^2 = -\xi^2, \xi^2 \in R_3$, GDD 3 of Row 5.

Omitting 2 in whole is not an arithmetic root system by Lemma 3.2(I).


(b)



$q^2 = -\xi^2, \xi^2 \in R_3$, GDD 2 of Row 6.


Omitting 2 in whole is not an arithmetic root system by Lemma 3.2(II).

(c)



$q = -\xi, \xi \in R_3$, GDD 3 of Row 7.


Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(I).



$q^3 \in R_6$, GDD 2 of Row 7.


Omitting 2 in whole is not an arithmetic root system by Lemma 3.2 (X).

(e)




$q = -\xi, \xi \in R_3$, GDD 3 of Row 7.

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(I).




$q = -\xi, \xi \in R_3$, GDD 4 of Row 7.

is empty.




$q = -\xi, \xi \in R_3$, GDD 1 of Row 8.

Omitting 2 in whole is an arithmetic root system by Lemma 3.2 (IV). Whole is affine by Lemma 3.4.




$q = -\xi, \xi \in R_3$, GDD 1 of Row 9.

Omitting 2 in whole is an arithmetic root system by Lemma 3.2 (IV). is affine by Lemma 3.4.



$q^2 = -\xi^2, \xi \in R_3$, GDD 2 of Row 10.

Omitting 2 in whole is an arithmetic root system by Lemma 3.2 (IV). Whole is affine by Lemma 3.4.




$q = -\xi, \xi \in R_3$, GDD 1 of Row 10.

Omitting 2 in whole is an arithmetic root system by Lemma 3.2 (IV). Whole is affine

by Lemma 3.4.

(l) or (9)  $q = -\xi, \xi \in R_3, \text{ GDD 2 of Row 10.}$

Omitting 2 in whole is an arithmetic root system by Lemma 3.2 (IV). Whole is affine by Lemma 3.4.


(m)  $q = -\xi, \xi \in R_3, \text{ GDD 1 of Row 11.}$

is empty.

(n)  $, q \in R_3. \text{ GDD 3 of Row 14.}$

Omitting 2 in whole is not an arithmetic root system by Lemma 3.2(I).

(o)


 $q = -\xi, \xi \in R_3, \text{ GDD 3 of Row 15.}$

is empty.


(p) or (10)  $, q \in R_3. \text{ GDD 2 of Row 16.}$

Omitting 2 in whole is an arithmetic root system Type 7. Whole is affine by Lemma 3.4.


(q)

 $q = -\xi, \xi \in R_3, \text{ GDD 3 of Row 16.}$

is empty.

(r)  $, q \in R_3. \text{ GDD 4 of Row 16.}$


Omitting 2 in whole is not an arithmetic root system by Lemma 3.2(I).



, $q \in R_3$. GDD 1 of Row 17.

is empty.


(s)



$q = -\xi, \xi \in R_3$, GDD 3 of Row 17.

is empty.

(t)



$q = -\xi, \xi \in R_3$, GDD 8 of Row 17.

is empty.


Case (iv) containing circle with four vertexes.

This case can be ignored since it is over a cycle.

Case (v) Complete diagram containing circle with four vertexes.


According to Lemma 3.40 we have to consider following cases.

(b) in Case (i)




234 is not an arithmetic root system by Lemma 3.2(I).

(b) in Case (ii)



If 134 is an arithmetic root system, then $\tilde{q}_{34} = q^{-1}$ by Lemma 3.3(I). 234 is not an arithmetic root system by Lemma 3.3(I).

(e) in Case (iii)



with $q \in R_6$.

124 is not an arithmetic root system by Lemma 3.2(I).

4.47 GDD 4 of Row 17

Case (i) Omitting 3 and adding 1.

According to Lemma 3.3 (IV) we have to consider following cases.

$$\begin{array}{c}
 (a') \text{ or } (1) \\
 q \quad q^{-1} \quad -1 \quad q \quad -1 \quad -q \quad -1 \\
 \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet
 \end{array}
 , q \in F^* \setminus \{1, -1\}. \text{ GDD 2 of Row 4.}$$

is affine by Lemma 3.15.

$$\begin{array}{c}
 (a'') \text{ or } (2) \\
 q \quad q^{-2} \quad -1 \quad q^2 \quad -1 \quad -q^2 \quad -1 \\
 \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet
 \end{array}
 , q \in F^* \setminus \{1, -1\}. \text{ GDD 2 of Row 5.}$$

is affine by Lemma 3.4.


$$\begin{array}{c}
 (a) \text{ or } (3) \\
 q^2 \quad q^{-2} \quad -1 \quad q \quad -1 \quad -q \quad -1 \\
 \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet
 \end{array}
 , q \in F^* \setminus \{1, -1\}. \text{ GDD 2 of Row 6.}$$

is affine by Lemma 3.15.

$$\begin{array}{c}
 (c) \\
 q^3 \quad q^{-3} \quad -1 \quad q \quad -1 \quad -q \quad -1 \\
 \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet
 \end{array}
 , q \in F^* \setminus \{1, -1\}. \quad q \notin R_3, \text{ GDD 2 of Row 7.}$$


is empty.

(e) or (4).




, $q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 8.

is affine by Lemma 3.15.




, $q \in R_3$. GDD 2 of Row 14.

is empty.




, $q \in R_3$. GDD 2 of Row 15.

is affine by Lemma 3.15.




, $q \in R_3$. GDD 2 of Row 16.

is affine by Lemma 3.15.




is empty.




, $q \in R_3$. GDD 1 of Row 17.

is affine by Lemma 3.15.



, $q \in R_3$. GDD 4 of Row 17.


is affine by Lemma 3.15.



is empty.

Case (ii) Omitting 3 and adding 2.

According to Lemma 3.3 (IV) we have to consider following cases.



, $q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 4.


Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(II).

(a'').  , $q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 5.

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(I).

(a).  , $q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 6.

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(II).

(c).  , $q \in F^* \setminus \{1, -1\}$. $q \notin R_3$, GDD 2 of Row 7.
is empty.


(e).  , $q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 8.

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(II).

(j).  , $q \in R_3$. GDD 2 of Row 14.
is empty.

(k) or (9).  , $q \in R_3$. GDD 2 of Row 15.


Omitting 1 in whole is an arithmetic root system by Lemma 3.2(II) or by Lemma 3.2 (XI)(d). Whole is affine by Lemma 3.20.

(l) or (10).  , $q \in R_3$. GDD 2 of Row 16.

Omitting 1 in whole is an arithmetic root system by Type 7. Whole is affine by Lemma 3.16.


(n).  , $q \in R_3$. GDD 1 of Row 17.

is empty.




, $q \in R_3$. GDD 1 of Row 17.

Omitting 1 in whole is an arithmetic root system by Lemma 3.3 (V)(a). Whole is affine by Lemma 3.20.




, $q \in R_3$. GDD 4 of Row 17.

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(II) or by Lemma 3.2 (XI).



, $q \in R_3$. GDD 4 of Row 17.

is empty.




, $q \in R_3$. GDD 9 of Row 17.


is empty.

Case (iii) Omitting 1 and adding 3.


According to Lemma 3.3 (IV) we have to consider following cases.




$q = -\xi, \xi \in R_3, q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 4.




$q^2 = -\xi, \xi \in R_3, q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 5.



$q = -\xi, \xi \in R_3, q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 6.



$q = -\xi, \xi \in R_3, q \in F^* \setminus \{1, -1\}$. $q \notin R_3$, GDD 2 of Row 7.



$q = -\xi, \xi \in R_3, q \in F^* \setminus \{1, -1\}$. GDD 2 of Row 8.

$$\begin{array}{c}
 \text{(j).} \\
 q \quad -q \quad -1 \quad -q^{-1} \quad 1 \quad q^{-1} \quad -1 \\
 \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet
 \end{array}$$

, $q \in R_3$. GDD 2 of Row 14.

$$\begin{array}{c}
 \text{(k).} \\
 -1 \quad q \quad -1 \quad q \quad -1 \quad -q \quad -1 \\
 \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet
 \end{array}$$

$q = -\xi, \xi \in R_3, q \in R_3$. GDD 2 of Row 15.

empty

$$\begin{array}{c}
 \text{(l).} \\
 -q \quad -q^{-1} \quad 1 \quad q \quad -1 \quad -q \quad -1 \\
 \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet
 \end{array}$$

$q = -\xi, \xi \in R_3, q \in R_3$. GDD 2 of Row 16.

is empty.

$$\begin{array}{c}
 \text{(n).} \\
 -1 \quad q \quad -1 \quad -1 \quad -1 \\
 \bullet \text{---} \bullet \text{---} \bullet
 \end{array}$$

is empty.

$$\begin{array}{c}
 \text{(n').} \\
 -1 \quad -1 \quad -1 \quad q \quad -1 \quad -q \quad -1 \\
 \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet
 \end{array}$$

$q = -\xi, \xi \in R_3, q \in R_3$. GDD 1 of Row 17.

is empty.

$$\begin{array}{c}
 \text{(o).} \\
 -1 \quad -q \quad -1 \quad q \quad -1 \quad -q \quad -1 \\
 \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet
 \end{array}$$

$q = -\xi, \xi \in R_3, q \in R_3$. GDD 4 of Row 17.

is empty .

$$\begin{array}{c}
 \text{(r).} \\
 q^{-1} \quad -q^{-1} \quad 1 \quad -1 \quad -1 \\
 \bullet \text{---} \bullet \text{---} \bullet
 \end{array}$$

is empty.

Case (iv) containing circle with four vertexes.


According to Lemma 3.39 (ii) we have to consider following cases.

(a') to (a'). empty. (a') to (a'') empty. (a') to (a) empty. (a') to (c) empty. (a') to (e) empty. (a') to (j) empty. (a'') to (a'). empty. (a'') to (a''). empty.

(a'') to (a) or (12).

$$\begin{array}{c}
 -1 \quad q^{-2} \quad q \quad q^{-4} \quad -1 \\
 \bullet \text{---} \bullet \text{---} \bullet
 \end{array}$$

with $q^4 = q$ is an arithmetic root system by Lemma 3.3 (III)(h).




is affine.

(a'') to (c) empty. (a'') to (e) empty.

(a'') to (j) or (13).

$$\begin{array}{c}
 -1 \quad q^{-2} \quad q \quad -q^{-2} \quad -1 \\
 \bullet \text{---} \bullet \text{---} \bullet
 \end{array}$$


with $q^{-2} = q$ is an arithmetic root system by Lemma 3.2 (V)(c).




is affine.

(a) to (a') empty. (a) to (a'') empty.

(a) to (a) or (14).




is an arithmetic root system.




is affine.

(a) to (c) empty. (a) to (e) empty.

(a) to (j) or (15).




with $q^{-1} = q^2$ is an arithmetic root system by Lemma 3.3 (III)(o) or by Lemma 3.2 (IX)(f) or by Lemma 3.3 (III)(o).




is affine.

(e) to (a') empty. (e) to (a'') empty. (e) to (a) empty.

(e) to (c) or (16).




with $-1 = -q^3$ is an arithmetic root system by GDD 2 of Row 6 or Lemma 3.3 (XI)(e).




is affine.

(e) to (e) or (17).




is an arithmetic root system by Lemma 3.2 (XI)(d).




is affine.

(k) to (a') empty. (k) to (a'') empty. (k) to (a) empty.

(k) to (c) or (18)



with $-q^3 = -1$ is an arithmetic root system by Type 7.





is affine.

(k) to (j) empty.

(l) to (a') or (19).

is an arithmetic root system.





is affine.

(l) to (a''). empty. (l) to (a). empty. (l) to (c) empty. (l) to (e) empty. (l) to (j). empty. (n') to (a') empty. (n') to (a'') empty. (n') to (a) empty.


(n') to (c) or (20).

with $-q^3 = -1$ is an arithmetic root system.




is affine.

(n') to (e) or (21)




is an arithmetic root system by Lemma 3.2 (XI)(e).



is affine.


(n') to (j) empty. (o) to (a') empty. (o) to (a'') empty. (o) to (a) empty.

(o) to (c) or (22)




with $-q^3 = -1$ is an arithmetic root system by Lemma 3.2 (XI)(e).

is affine.
 (o) to (e) or (23)



is an arithmetic root system by Type 7.




is affine.
 (o) to (j) empty.


4.48 GDD 5 of Row 17

Case (i) Omitting 3 and adding 1.

According to Type 3 we have to consider following cases.



is affine by Lemma 3.7 or by Type 3.




is affine by Lemma 3.38.

Case (ii) Omitting 1 and adding 2.


According to Lemma 3.3 (III) we have to consider following cases.



Omitting 3 is an arithmetic root system by Lemma 3.2 (V)(a). Whole is affine by Lemma 3.4.




Omitting 3 in whole is not an arithmetic root system by Lemma 3.2 (V).



, $q \in F^* \setminus \{1, -1\}$. $q \in R_6$, GDD 3 of Row 8.


Omitting 3 is an arithmetic root system by GDD 5 of Row 17 or by Lemma 3.3 (III)(n).

Whole is affine by Lemma 3.4.




, $q \in R_3$. GDD 1 of Row 14.

Omitting 3 is not an arithmetic root system by Lemma 3.2(I).




is empty.

, $q \in R_3$. GDD 1 of Row 15.




is empty.

$q \in R_3$. GDD 1 of Row 16.



is empty.


$q \in R_3$. GDD 2 of Row 17.



, $q \in R_3$. GDD 5 of Row 17.


Omitting 3 is an arithmetic root system by GDD 4 of 15 Row or by Lemma 3.2 (V)(b).

Whole is affine by Lemma 3.4.




is empty.

, $q \in R_3$. GDD 7 of Row 17.




GDD 1 of Row 4.

Omitting 3 is not an arithmetic root system by Lemma 3.2 (IX).




$q^2 \in R_6$, GDD 1 of Row 5.

Omitting 3 is not an arithmetic root system by Lemma 3.2 (IX).



GDD 1 of Row 17.

is empty.




GDD 9 of Row 17.


is empty.

Case (iii) Omitting 1 and adding 3


According to Lemma 3.3 (I) we have to consider following cases.




$q = -\xi, \xi \in R_3$, GDD 2 of Row 4.




$q^2 = -\xi, \xi \in R_3$, GDD 3 of Row 5.




$q^2 = -\xi, \xi \in R_3$, GDD 2 of Row 6.




GDD 3 of Row 7.




$q^3 = -\xi, \xi \in R_3, q \notin R_3$. GDD 2 of Row 7.



$-q^3 \in R_3, q \notin R_3$. GDD 4 of Row 7.



$-q \in R_3$, GDD 1 of Row 8.



$-q \in R_3$, GDD 1 of Row 9.

$$\begin{array}{ccccccc}
 & (i') & & & & & \\
 q & q^{-1} & -1 & q^2 & q^{-2} & -q^{-2} & -1 \\
 \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet
 \end{array}$$

$-q^2 \in R_3, q \in R_6, q \notin R_3$. GDD 2 of Row 10.

$$\begin{array}{ccccccc}
 & (k) & & & & & \\
 q & q^{-1} & -1 & q^{-1}q & -q & -1 & \\
 \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet
 \end{array}$$

$-q \in R_3, q \notin R_3$. GDD 1 of Row 10.

$$\begin{array}{ccccccc}
 & (l) & & & & & \\
 q^{-2} & q^2 & -1 & q^{-1}q & -q & -1 & \\
 \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet
 \end{array}$$

$, q \in R_6, q \notin R_3$. GDD 2 of Row 10.

$$\begin{array}{ccccccc}
 & (m) & & & & & \\
 q & q^{-1} & -1 & q^{-1}q & -q & -1 & \\
 \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet
 \end{array}$$

$, q \in R_3$. GDD 1 of Row 11.

is empty.

$$\begin{array}{ccccccc}
 & (n) & & & & & \\
 q^{-1} & -q^{-1} & -1 & -q & -q^{-1} & -1 & \\
 \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet
 \end{array}$$

$, q \in R_3$. GDD 3 of Row 14.

$$\begin{array}{ccccccc}
 & (p) & & & & & \\
 -1 & q & -1 & -q^{-1}q & q & -1 & \\
 \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet
 \end{array}$$

$, q \in R_3$. GDD 2 of Row 16.

$$\begin{array}{ccccccc}
 & (r) & & & & & \\
 q & -1 & -1 & -q^{-1}q & q & -1 & \\
 \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet
 \end{array}$$

$, q \in R_3$. GDD 4 of Row 16.

$$\begin{array}{ccccccc}
 & (s') & & & & & \\
 -1 & q & -1 & -1 & -1 & & \\
 \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet
 \end{array}$$

is empty.

Case (iv) containing circle with four vertexes.

According to Lemma 3.39 (ii) we have to consider following cases.

(a) to (a') empty. (a) to (a). empty.

(a) to (b) empty. (a) to (d) empty.


(a) to (f) empty.

(a) to (g) empty.

(a) to (h) or (6).

$$\begin{array}{ccccccc}
 & & & & & & \\
 -1 & q^2 & q^{-2} & -r^{-1} & -1 & & \\
 \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet
 \end{array}$$


with $-r = q^{-2}$ is an arithmetic root system.




is affine.

(a) to (i'). empty. (a) to (k). empty

(a) to (l) or (7).




is an arithmetic root system.




is affine.

(a) to (n) or (8)




is an arithmetic root system by Lemma 3.3 (III)(o).




is affine.

(a) to (p) empty.

(a) to (r) or (9)




with $q^{-2} = q$, is an arithmetic root system by Lemma 3.3 (III)(l).




is affine.

(b) to (a')




is not an arithmetic root system by Lemma 3.2 (XI) with $q \in R_3$.

(b) to (a) empty. (b) to (b)




with $\xi^2 = -q$ is not an arithmetic root system by Lemma 3.3 (XI) with $q \in R_3$.

(b) to (d)




with $\xi^3 = -q$ is not an arithmetic root system by Lemma 3.3 (XI)

(b) to (f) empty. (b) to (g) empty. (b) to (h) or (10)




with $-r = -1$, is an arithmetic root system by Lemma 3.3 (XI)(e).




is affine.

(b) to (i') empty. (b) to (k) empty. (b) to (l) empty. (b) to (n) empty.

(b) to (p) or (11)



is an arithmetic root system by Type 7.



is affine.

(b) to (r) empty.

4.49 GDD 6 of Row 17


Case (i) Omitting 3 and adding 1.

checking step by step we have to consider following cases empty by Lemma 3.2

Case (ii) Omitting 3 and adding 2.

checking step by step we have to consider following cases


(a)



, ($q \in R_3$, see GDD 6 of Row 17.)

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2 (V).


(b) or (1)



Omitting 1 in whole is an arithmetic root system by Lemma 3.3 (III)(o) Whole is affine by Lemma 3.4.

Case (iii) Omitting 1 and adding 3.

According to Type 4 and GDD 8,9 of Row 17 we have to consider following cases.




Case (iv) containing circle with four vertexes.

It is empty.

4.50 GDD 7 of Row 17

Case (i) Omitting 3 and adding 1.


According to Lemma 3.3 (I) we have to consider following cases.



is affine by Lemma 3.19.



is affine by Lemma 3.19 or by Lemma 3.4.



is affine by Lemma 3.19.

(d) or (4)

$$\begin{array}{ccccccc} -1 & q & -1 & q^{-3}q^3 & -q^3 & -1 & \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

$q^3 \in R_3$. GDD 2 of Row 7.

is affine by Lemma 3.19.

$$\begin{array}{ccccccc} \text{(f) or (5)} & & & & & & \\ -q & q^2 & -1 & q^{-3}q^3 & -q^3 & -1 & \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

$q^3 \in R_3$. GDD 4 of Row 7.

is affine by Lemma 3.19.

$$\begin{array}{ccccccc} \text{(g) or (6)} & & & & & & \\ q^{-1} & q & -1 & q^{-1}q & -q & -1 & \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

, GDD 1 of Row 8.

is affine by Lemma 3.19.

$$\begin{array}{ccccccc} \text{(h) or (7)} & & & & & & \\ r & r^{-1} & -1 & q^{-1}q & -q & -1 & \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

, GDD 1 of Row 9.

is affine by Lemma 3.19.

$$\begin{array}{ccccccc} \text{(i')} & & & & & & \\ q & q^{-1} & -1 & q^2 & q^{-2} & -q^{-2} & -1 \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

$q^2 = \xi$, $\xi \in R_3$, $q \in R_6$. GDD 2 of Row 10.

is empty.

$$\begin{array}{ccccccc} \text{(k)} & & & & & & \\ q & q^{-1} & -1 & q^{-1}q & -q & -1 & \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

$q \notin R_3$. GDD 1 of Row 10.

is empty.

$$\begin{array}{ccccccc} \text{(l)} & & & & & & \\ q^{-2} & q^2 & -1 & q^{-1}q & -q & -1 & \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

$q \notin R_3$. GDD 2 of Row 10.

is empty.

$$\begin{array}{ccccccc} \text{(m) or (8)} & & & & & & \\ q & q^{-1} & -1 & q^{-1}q & -q & -1 & \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

$q \in R_3$. GDD 1 of Row 11.

is affine by Lemma 3.19.

$$\begin{array}{ccccccc} \text{(n)} & & & & & & \\ q^{-1} & -q^{-1} & -1 & -q & -q^{-1} & -1 & \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

$q \in R_3$. GDD 3 of Row 14.

is empty.

$$\begin{array}{ccccccc} \text{(p)} & & & & & & \\ -1 & q & -1 & -q^{-1}q & q & -1 & \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$


$q \in R_3$. GDD 2 of Row 16.

is empty.

$$\begin{array}{ccccccc} \text{(r)} & & & & & & \\ q & -1 & -1 & -q^{-1}q & q & -1 & \\ \bullet & \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \end{array}$$

$q \in R_3$. GDD 4 of Row 16.

is empty.





is empty.

Case (ii) Omitting 3 and adding 2.

According to Lemma 3.3 (III) we have to consider following cases.



Omitting 1 in whole is not an arithmetic root system, by Lemma 3.2 (V).



Omitting 1 in whole is an arithmetic root system by GDD 7 of Row 17 or by Lemma 3.2 (IX)(f). Whole is affine by Lemma 3.4.




Omitting 3 is an arithmetic root system by GDD 6 of Row 17 or by Lemma 3.2 (IX)(e). Whole is affine by Lemma 3.4.



Omitting 1 in whole is not an arithmetic root system by Lemma 3.2 (V) or by Lemma 3.2 (IX).

(l).  , $q \in R_3$. GDD 2 of Row 17.

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2 (V).

(n).  , $q \in R_3$. GDD 5 of Row 17.
is empty.

(o).  , $q \in R_3$. GDD 7 of Row 17.

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2 (V).

(a').  GDD 1 of Row 4.

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2 (IX),

(a'').  GDD 1 of Row 5.


Omitting 1 in whole is not an arithmetic root system by Lemma 3.2 (IX).


(k').  GDD 1 of Row 17.
is empty.

(p').  GDD 9 of Row 17.
is empty.

Case (iii) Omitting 1 and adding 3.

According to Type 4 we have to consider following cases.

(a)  (see, $q \in R_3$, GDD 2 of Row 14 in Table A2).

(b)  (see, $q \in R_3$, GDD 3 of Row 14 in Table B).

(c) $\begin{array}{c} -1 \quad -1 \quad -1 \quad -q^{-1} \quad q^{-1} \quad q \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array}$ (see, $q \in R_3$, GDD 9 of Row 17 in Table B).

Case (iv) Containing circle with four vertexes.

According to Lemma 3.39 (ii) we have to consider following cases.

(a') to (a) $\begin{array}{c} -1 \quad q \quad -1 \quad -q^{-1} \quad 1 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}$


is not an arithmetic root system by Lemma 3.2 (XI) or by Lemma 3.2 (II).

(a') to (b) empty.

(a') to (c) or (11).

$$\begin{array}{c} -1 \quad q \quad -1 \quad -1 \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}$$

is an arithmetic root system by Lemma 3.2(II)(f) or by Lemma 3.2 (XI)(e).




is affine.

(a) to (a) empty.(a) to (b) or (12)

$$\begin{array}{c} -1 \quad q^2 \quad -q^{-1} \quad -q^{-2} \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}$$

with $-q^2 = -q^{-1}$ is an arithmetic root system by Lemma 3.2 (IX)(d) or by Lemma 3.3 (III)(d).




is affine.

(a) to (c) empty.

(b) to (a) or (13)

$$\begin{array}{c} -1 \quad q \quad -1 \quad -q^{-2} \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}$$

is an arithmetic root system by Lemma 3.2(II)(g) or by Lemma 3.2 (XI)(d).




is affine.

(b) to (b). empty.

(b) to (c) or (14).

$$\begin{array}{c} -1 \quad q \quad -1 \quad -1 \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}$$

is an arithmetic root system by Lemma 3.2 (II)(f) and by Lemma 3.2 (XI)(e).




is affine.

(d) to (a)

is not an arithmetic root system by Lemma 3.2 (XI).

(d) to (b). empty.

(d) to (c) or (15)




is an arithmetic root system by Lemma 3.2 (XI)(e).

is affine.

(f) to (a) empty. (f) to (b). empty. (f) to (c) empty. (g) to (a) empty. (g) to (b). empty.

(g) to (c) empty.


(h) to (a) or (16)



with $r = -1$ is an arithmetic root system by Lemma 3.2 (XI)(e).

is affine.


(h) to (b) or (17)



with $r = -q$ is an arithmetic root system.

is affine.

(h) to (c) or (18)



with $r = -1$. Whole is an arithmetic root system.

is affine.


(m) to (a) empty.(m) to (b). empty. (m) to (c) empty.

4.51 GDD 8 of Row 17

Case (i) Omitting 3 and adding 1.


According to Lemma 3.3 (V) we have to consider following cases.

(a) or (1)




Omitting 2 in whole is arithmetic root system when $r = -q^{-1}$ by Type 4. Omitting 2 in whole is not an arithmetic root system when $r \neq -q^{-1}$ by Lemma 3.2(I). Whole is affine when $r = -q^{-1}$ by Lemma 3.4.

(b)




Omitting 2 in whole is not an arithmetic root system by Lemma 3.2(I).

(c) or (2)




Omitting 1 in whole is an arithmetic root system when $q = -r$ by Type 4. Omitting 2 in whole is an arithmetic root system when $q = -r$ by Type 3. Whole is affine when $q = -r$.


(c')




Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(I).

(e)  $(r \in R_6, \text{ see GDD 4 of Row 7 in Table A2.})$

Omitting 2 in whole is not an arithmetic root system by Lemma 3.2(I).

(f) or (3)  $(r^2 \neq 1, \text{ see GDD 1 of Row 9 in Table A2.})$


Omitting 2 in whole is not an arithmetic root system when $r \neq -q$ by Lemma 3.2 (IV). Omitting 2 in whole is an arithmetic root system when $r = -q$ by Type 4. Whole is affine by Lemma 3.4.

(g)  $(r \in R_4, \text{ see GDD 1 of Row 2 in Table A2.})$


Omitting 2 in whole is not an arithmetic root system by Lemma 3.2(I).

Case (ii) Omitting 3 and adding 2.


According to Lemma 3.3 (V) we have to consider following cases.

(a) or (4)  $(r \in R_3, \text{ see GDD 1 of Row 17 in Table A2.})$


Omitting 1 in whole is an arithmetic root system when $q = r$ or $q = r^2$ by Lemma 3.2 (X). Whole is affine by Lemma 3.4.

(b)  $(r \in R_3, \text{ see GDD 9 of Row 17 in Table A2.})$


Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(I).




Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(I).




Omitting 2 in whole is not an arithmetic root system by Lemma 3.2(I).



Omitting 1 in whole is not an arithmetic root system when $q \neq r^2$ by Lemma 3.2(I).
 Omitting 1 in whole is an arithmetic root system when $q = r^2$ by Type 3. Whole is affine.
 by Lemma 3.4.




Omitting 1 in whole is an arithmetic root system by Lemma 3.2 (IV). Whole is affine
 by Lemma 3.4.

(g)  $(r \in R_4, \text{ see GDD 1 of Row 2 in Table B.})$


Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(I)

Case (iii) Omitting 2 and adding 3.


Checking step by step we have to consider following cases.

(a)  $(q \in R_6, \text{ see GDD 2 of Row 13.})$


Omitting 1 in whole is not an arithmetic root system by Lemma 3.3 (III).

(b)  $(q \in R_3, \text{ see GDD 2 of Row 13.})$


Omitting 1 in whole is not an arithmetic root system by Lemma 3.3 (III).

(c) or (7)  $(q \in R_3, \text{ see GDD 4 of Row 15.})$


Omitting 1 in whole is an arithmetic root system by GDD 7 of Row 17 or by Lemma 3.3 (III)(o). Whole is affine by Lemma 3.4.

(d)  $(q \in R_3, \text{ see GDD 5 of Row 17.})$

is empty.

(e) or (8)  $(q \in R_3, \text{ see GDD 6 of Row 17.})$

Omitting 1 in whole is an arithmetic root system by Lemma 3.3 (III)(h). Whole is affine by Lemma 3.4.

(f) or (9)  $(q \in R_3, \text{ see GDD 7 of Row 17.})$

Omitting 1 in whole is an arithmetic root system. Whole is affine by Lemma 3.4.


Case (iv) Containing circle with four vertexes.

This case can be ignored since it is over a cycle.

Case (v) Complete diagram containing circle with four vertexes.


According to Lemma 3.40 we have to consider following cases.

(c') in Case (i)



234 is not an arithmetic root system by Lemma 3.2(I).

(c) in Case (i)



234 is not an arithmetic root system by Lemma 3.2(I).

(c) in Case (ii) and (c') in Case (ii) are the same as (c) in Case (i) and (c') in Case (i).

4.52 GDD 9 of Row 17

(i) Omitting 3 and adding 1.

According to Lemma 3.3 (V) we have to consider following cases.

(a) or (1)

$$\overset{-1}{r} \quad -1_{-1} \quad -1_{-q^{-1}q^{-1}}$$

($q \in R_3, r \in R_2 \cup R_3 \cup R_4 \cup R_6$, see GDD 1 of Row 17 in Table A2). It is affine by Lemma 3.4.

(b) or (2)

$$\overset{r^{-1}}{-r^{-1}} \quad -1_{-1} \quad -1_{-q^{-1}q^{-1}}$$

($r \in R_3$. see GDD 9 of Row 17 in Table A2).

is affine by Lemma 3.4.

(e) or (3) $\begin{array}{c} -r^{-1} \quad -1 \quad -1 \\ r^2 \quad -1 \quad -q^{-1}q^{-1} \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array}$ (see, $r \in R_6$, GDD 4 of Row 7 in Table A2).

is affine by Lemma 3.4.

(f) or (4) $\begin{array}{c} r \quad -1 \quad -1 \\ r^{-1} \quad -1 \quad -q^{-1}q^{-1} \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array}$ (see, $r^2 \neq 1$, GDD 1 of Row 9 in Table A2).

is affine by Lemma 3.4.

(g) or (5) $\begin{array}{c} r \\ -1 \quad -1 \quad -1 \\ -q^{-1}q^{-1} \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array}$ (see, $r \in R_4$, GDD 1 of Row 2 in Table A2).

is affine by Lemma 3.4.

Case (ii) Omitting 1 and adding 2.

According to Type 4 we have to consider following cases.

(a) or (6) $\begin{array}{c} -1 \quad -1 \\ -q \quad -1 \\ \bullet \text{---} \bullet \begin{array}{l} \nearrow -1 \\ \searrow -q^{-1} \end{array} \bullet \end{array}$, Type 4

Omitting 3 in whole is an arithmetic root system by Lemma 3.3 (V)(a). Whole is affine by Lemma 3.4.

(b) or (7) $\begin{array}{c} -q^{-1} \quad -1 \\ -q \quad -1 \\ \bullet \text{---} \bullet \begin{array}{l} \nearrow -1 \\ \searrow -q^{-1} \end{array} \bullet \end{array}$, Type 4

Omitting 3 in whole is an arithmetic root system by Lemma 3.3 (IV). Whole is affine by Lemma 3.4.

(c) or (8) $\begin{array}{c} -1 \quad -1 \\ -1 \quad -1 \\ \bullet \text{---} \bullet \begin{array}{l} \nearrow -1 \\ \searrow -q^{-1} \end{array} \bullet \end{array}$, (see, $q \in R_3$, GDD 9 of Row 17 in Table A2).

Omitting 3 in whole is an arithmetic root system. Whole is affine by Lemma 3.16 or by Lemma 3.4.

Case (iii) Omitting 1 and adding 3.

According to inversion of Type 4 we have to consider following cases.

(a) $\begin{array}{c} -1 \quad -1 \\ q^{-1}q^{-1} \quad -q^{-1}q^{-1} \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array}$ (see GDD 6 of Row 17 in Table A2).

(b)

$$\begin{array}{c} -1 \quad q^{-1}q \quad -q \quad -1 \quad -1 \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad (\text{ see GDD 7 of Row 17 in Table A2}).$$

(c)
$$\begin{array}{c} q \quad q^{-1}-q \quad q^{-1} \quad -1 \quad -1 \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad (q \in R_3. \text{ see GDD 2 of Row 13 in Table A2}).$$

is empty.

(e)
$$\begin{array}{c} -1 \quad q^{-1}-q \quad q^{-1} \quad -1 \quad -1 \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad (q \in R_3. \text{ see GDD 4 of Row 15 in Table A2}).$$

is empty.

(f)
$$\begin{array}{c} -1 \quad -q^{-1}q \quad q \quad -1 \quad -1 \quad -1 \\ \bullet \text{---} \bullet \text{---} \bullet \text{---} \bullet \end{array} \quad (q \in R_3. \text{ see GDD 5 of Row 17 in Table A2}).$$


is empty.

Case (iv) containing circle with four vertexes.

According to Lemma 3.39 (ii) we have to consider following cases.

(a) to (a) or (9)
$$\begin{array}{c} -1 \quad r \quad -1 \quad q^{-1}q^{-1} \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}$$


is not an arithmetic root system by Lemma 3.2 (I) when $q \neq r$. is an arithmetic root system by Type 2 when $q = r$.



is affine.

(a) to (b) or (10)
$$\begin{array}{c} -1 \quad r \quad -1 \quad q \quad q^{-1} \\ \bullet \text{---} \bullet \text{---} \bullet \end{array}$$

is an arithmetic root system when $r = q^{-1}$ or $q^{-1} = r^2$ by Lemma 3.2 (XI).



is affine.

(b) to (a) empty. (b) to (b) empty. (e) to (a) empty. (e) to (b) empty. (f) to (a) empty. (f) to (b) empty. (g) to (a) empty. (g) to (b) empty.

4.53 GDD 1 of Row 18

Case (i) Omitting 3 and adding 1.

According to Lemma 3.3 (II) we have to consider following cases.

(a) or (1).
 $q \quad q^{-1} q \quad q^{-1} q \quad q^{-1} q^{-3}$

, $q \in F^* \setminus \{1\}$. GDD 1 of Row 1.

is affine by Lemma 3.4.

(b) or₂(2).
 $q \quad q^{-2} q^2 \quad q^{-2} q^2 \quad q^{-2} q^{-6}$

, $q \in F^* \setminus \{1, -1\}$. GDD 1 of Row 2.

is affine by Lemma 3.4.

(c) or (3).
 $q^2 \quad q^{-2} q \quad q^{-1} q \quad q^{-1} q^{-3}$

, $q \in F^* \setminus \{1, -1\}$. GDD 1 of Row 3.

is affine by Lemma 3.4.

(d) or (4).
 $-1 \quad q^{-1} q \quad q^{-1} q \quad q^{-1} q^{-3}$

, $q \in F^* \setminus \{1, -1\}$. GDD 1 of Row 4.

is affine by Lemma 3.4.

(f).
 $-1 \quad q^{-2} q \quad q^{-1} q \quad q^{-1} q^{-3}$

, $q \in R_3 \cup R_6$. GDD 1 of Row 13.

is empty.

(h).
 $q \quad -q^{-1}q \quad -q^{-1}q \quad -q^{-1}q^{-3}$

, $q \in R_3$. GDD 5 of Row 16.

is empty.

(i) or (5).
 $q^{-3} \quad q^{-1} q \quad q^{-1} q \quad q^{-1} q^{-3}$

, $q \in R_9$. GDD 1 of Row 18.

is affine by Lemma 3.4.

Case (ii) Omitting 3 and adding 2.

According to Lemma 3.3 (II) we have to consider following cases.

(a) or (6).


, $q \in R_9$. GDD 1 of Row 1.

Omitting 1 in whole is an arithmetic root system by GDD 1 of Row 18. Whole is affine by Lemma 3.4.

(b).


, $q \in F^* \setminus \{1, -1\}$. GDD 1 of Row 2.

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(I).




, $q \in F^* \setminus \{1, -1\}$. GDD 1 of Row 3.

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(I).




, $q \in F^* \setminus \{1, -1\}$. GDD 1 of Row 4.


Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(I).




, $q \in R_3$. GDD 1 of Row 12.



, $q \in R_3 \cup R_6$. GDD 1 of Row 13.



, $q \in R_3$. GDD 5 of Row 16.



, $q \in R_9$. GDD 1 of Row 18.

Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(I).

Case (iii) Omitting 1 and adding 3.

Checking step by step we have to consider following cases.

There are not affine.

Case (iv) containing circle with four vertexes.

empty.

4.54 GDD 2 of Row 18

Case (i) Omitting 3 and adding 1.


Checking step by step we have to consider following cases.

There are not affine.

Case (ii) Omitting 3 and adding 2.

Checking step by step we have to consider following cases.

(a)



Omitting 1 in whole is not an arithmetic root system by Lemma 3.2(I).

Case (iii) Omitting 1 and adding 3.

Checking step by step we have to consider following cases.


There are not affine.

Case (iv) containing circle with four vertexes.

It is empty.


4.55 More about GDD 3 of Row 15

Lemma 4.1. *or (15)*



is affine.

Proof. Omitting 2 and adding 1 in GDD 3 of Row 15.



Omitting 1 in whole is an arithmetic root system. Omitting 3 in whole is an arithmetic root system. Thus whole is affine.

5 Appendix

5.1 Affine over connected Generalized Dynkin Diagram with Rank 1 or 2

Proposition 5.1. $n = 2$. *A connected GDD is affine if and only if it is not an arithmetic root system*

Proposition 5.2. $n = 3$. *A connected GDD is affine if and only if it is not an arithmetic root system and omitting every vertex still is an arithmetic root system.*

References

- [AS02] N. Andruskiewitsch, H.-J. Schneider, Pointed Hopf algebras, in *New Directions in Hopf Algebras*, MSRI Series, pp.1-68, Cambridge Univ. Press, 2002.
- [AS10] N. Andruskiewitsch, H.-J. Schneider, On the classification of finite-dimensional pointed Hopf algebras, *Ann. Math.* **171** (2010), 375-417.
- [AHS08] N. Andruskiewitsch, I. Heckenberger and H.J. Schneider, The Nichols algebra of a semisimple Yetter-Drinfeld module, *Amer. J. Math.* **132** (2010), 1493-1547.
- [An11] I. Angiono, Nichols algebras of unidentified diagonal Type, arXiv:1108.5157.
- [Ha] F. Harary, *Graph Theory*, AddisonCWesley, USA, 1969.
- [He06a] I. Heckenberger, Classification of arithmetic root systems, *Adv. Math.* **220** (2009), 59-124.
- [He06b] I. Heckenberger, The Weyl-Brandt groupoid of a Nichols algebra of diagonal Type, *Invent. Math.* **164** (2006), 175–188.
- [He05] I. Heckenberger, Nichols algebras of diagonal Type and arithmetic root systems, *Habilitation*, 2005.
- [He07] I. Heckenberger, Examples of finite-dimensional rank 2 Nichols algebras of diagonal Type, *Compos. Math.*, **143** (2007) 1, 165-190.
- [HS08] I. Heckenberger, H.-J. Schneider, Root systems and Weyl groupoids for Nichols algebras, arXiv:0807.0691.
- [Kh99] V. K. Kharchenko, A Quantum analog of the poincaré-Birkhoff-Witt theorem, *Algebra and Logic*, **38** (1999), 259-276

- [Ka85] V. G. Kac. Infinite dimensional Lie Algebras. London:Cambridge Univ, 1985.
- [Lo83] M. Lothaire, Combinatorics on words. London:Cambridge University Press, 1983.
- [Wa02] Z. Wan. The introduction of Kac-Moody algebras. Beijing: Science Press, 2002(in Chinese).
- [WZZ15a] W. Wu, S. Zhang and Y.-Z. Zhang, Relationship between Nichols braided Lie algebras and Nichols algebras, J. Lie Theory **25** (2015), 45-63.
- [WZZ15b] W. Wu, S. Zhang and Y.-Z. Zhang, On Nichols (braided) Lie algebras, Int. J. Math. **26** (2015), 1550082.
- [ZWTZ] S. Zhang, W. Wu, Z. Tan and Y.-Z. Zhang, Nichols algebras over classical Weyl groups, Fomin-Kirillov algebras and Lyndon basis, arXiv:1307.8227