

Local Existence of Strong Solutions to the 3D Zakharov-Kuznestov Equation in a Bounded Domain

Chuntian Wang

*Department of Mathematics and The Institute for Scientific Computing and Applied Mathematics
Indiana University, Bloomington, IN 47405
email: wang211@uemail.iu.edu*

April 21, 2019

Contents

1	Introduction	2
2	The ZK equation in a rectangle	3
3	Parabolic regularization	4
3.1	Global bounds independent of ϵ for u^ϵ	5
3.2	Local bounds independent of ϵ for u^ϵ	5
3.3	A singular perturbation argument	13
4	Passage to the limit	16
5	Appendix: a trace result	17

Abstract

We consider here the local existence of strong solutions for the Zakharov-Kuznestov (ZK) equation posed in a limited domain $\mathcal{M} = (0, 1)_x \times (-\pi/2, \pi/2)^d$, $d = 1, 2$. We prove that in space dimensions 2 and 3, there exists a strong solution on a short time interval, whose length only depends on the given data. We use the parabolic regularization of the ZK equation as in [STW12] to derive the global and local bounds independent of ϵ for various norms of the solution. In particular, we derive the local bound of the nonlinear term by a singular perturbation argument. Then we can pass to the limit and hence deduce the local existence of strong solutions.

Keywords: Zakharov-Kuznetsov equation, Korteweg-de Vries equation, Local existence of strong solutions in 3D

1 Introduction

The Zakharov-Kuznetsov equation

$$\frac{\partial u}{\partial t} + \Delta \frac{\partial u}{\partial x} + c \frac{\partial u}{\partial x} + u \frac{\partial u}{\partial x} = f, \quad (1.1)$$

where $u = u(x, x^\perp, t)$, $x^\perp = y$ or $x^\perp = (y, z)$, describes the propagation of nonlinear ionic-sonic waves in a plasma submitted to a magnetic field directed along the x -axis. Here $c > 0$ is the sound velocity. It has been derived formally in a long wave, weakly nonlinear regime from the Euler-Poisson system in [ZK74] and [LS82]. A rigorous derivation is provided in [LLS13]. For more general physical references, see [BPS81] and [BPS83]. When u depends only on x and t , (1.1) reduces to the classical Korteweg-de Vries (KdV) equation.

Concerning the initial and boundary value problems of the Korteweg-de Vries equation posed on a bounded interval $(0, L)$, we refer the interested readers to e.g. [BSZ03], [CG01a] and [CG01b].

The initial and boundary value problem associated with (1.1) has been studied in the half space $\{(x, y) : x > 0\}$ ([Fam06]), on a strip like $\{(x, y) : x \in \mathbb{R}, 0 < y < L\}$ ([BF13]) or $\{(x, x^\perp) : 0 < x < 1, x^\perp \in \mathbb{R}^d, d = 1, 2\}$ ([Fam08] and [ST10]), and in a rectangle $\{(x, x^\perp) : 0 < x < 1, x^\perp \in (-\pi/2, \pi/2)^d, d = 1, 2\}$ ([STW12]). Specifically in [STW12], the authors have established, for arbitrary large initial data, the existence of global weak solutions in space dimensions 2 and 3 ($d = 1$ and 2 respectively) and a result of uniqueness of such solutions in the two-dimensional case.

As for the existence of strong solutions, the global existence in space dimension 2 has been proven in a half strip $\{(x, y) : x > 0, y \in (0, L)\}$ in [LT13]. The existence and exponential decay of regular solutions to the linearized ZK equation in a rectangle $\{(x, y) : x \in (0, L), y \in (0, B)\}$ has been studied in [DL13].

However, to the best of our knowledge, there has been no result so far for the local existence of strong solutions in 3D in a limited domain. In the present article we prove the short time existence of strong solutions in a 3D rectangular domain as in [STW12].

The article is organized as follows.

Firstly we introduce the basic settings of the equation and the related functional spaces in Section 2.

Secondly we introduce the parabolic regularization as in [ST10] and [STW12]. We derive the global bounds on u^ϵ independent of ϵ in Section 3.1. Then we derive the local bounds on $[0, T_*)$, where $T_* > 0$ depends only on the data (Sections 3.2 and 3.3). In particular, we use a singular perturbation argument in the x direction to deduce the local bounds on the nonlinear term in Section 3.3.

Finally we pass to the limit on the regularized equation and obtain the local existence of a strong solution (Section 4).

In the Appendix, we recall from [ST10] and [STW12] a trace result concerning a singular perturbation problem in the x direction.

2 The ZK equation in a rectangle

We aim to study the ZK equation:

$$\frac{\partial u}{\partial t} + \Delta \frac{\partial u}{\partial x} + c \frac{\partial u}{\partial x} + u \frac{\partial u}{\partial x} = f, \quad (2.1)$$

in a rectangle or parallelepiped domain in \mathbb{R}^n with $n = 2$ or 3 , denoted as $\mathcal{M} = (0, 1)_x \times (-\pi/2, \pi/2)^d$, with $d = 1$ or 2 , $\Delta u = u_{xx} + \Delta^\perp u$, $\Delta^\perp u = u_{yy}$ or $u_{yy} + u_{zz}$. In the sequel we will use the notations $I_x = (0, 1)_x$, $I_y = (-\pi/2, \pi/2)_y$, $I_z = (-\pi/2, \pi/2)_z$, and $I_{x^\perp} = I_y$ or $I_y \times I_z$. We assume the boundary conditions on $x = 0, 1$ to be

$$u(0, x^\perp, t) = u(1, x^\perp, t) = u_x(1, x^\perp, t) = 0, \quad (2.2)$$

and the initial condition reads:

$$u(x, x^\perp, 0) = u_0(x, x^\perp). \quad (2.3)$$

We also need suitable boundary conditions in the y and z directions. As in [STW12], we will choose either the Dirichlet boundary conditions

$$u = 0 \text{ at } y = \pm \frac{\pi}{2} \text{ (} z = \pm \frac{\pi}{2} \text{)}, \quad (2.4)$$

or the periodic boundary conditions

$$u|_{y=\frac{\pi}{2}}^{y=-\frac{\pi}{2}} = u_y|_{y=\frac{\pi}{2}}^{y=-\frac{\pi}{2}} = 0 \text{ (} u|_{z=\frac{\pi}{2}}^{z=-\frac{\pi}{2}} = u_z|_{z=\frac{\pi}{2}}^{z=-\frac{\pi}{2}} = 0 \text{)}. \quad (2.5)$$

We will study the initial and boundary value problem (2.1)-(2.3) supplemented with the boundary condition (2.4), that is, in the Dirichlet case, and we will make some remarks on the extension to the periodic boundary condition case.

We will denote by $|\cdot|$ and (\cdot, \cdot) the norm and the inner product of $L^2(\mathcal{M})$.

We use the following functional space in the sequel:

$$Au := \Delta u_x + cu_x, \quad \forall u \in D(A),$$

where $D(A) = \{u \in L^2(\mathcal{M}) : Au \in L^2(\mathcal{M}), u = 0 \text{ on } \partial\mathcal{M}, u_x(1, x^\perp, t) = 0\}$. Note that the trace theorem proven in [STW12] shows that if $u \in L^2(\mathcal{M})$ and $Au \in L^2(\mathcal{M})$ then the traces of u on $\partial\mathcal{M}$, and of u_x at $x = 1$ make sense.

We also consider the space

$$\Xi = \{u \in H^2(\mathcal{M}) \cap H_0^1(\mathcal{M}), u_x|_{x=1} = 0\}, \quad (2.6)$$

and endow this space with the scalar product and norm $[\cdot, \cdot]_2$ and $[\cdot]_2$,

$$\begin{aligned} [u, v]_2 &= (u_{xx}, v_{xx}) + (u_{yy}, v_{yy}) + (u_{zz}, v_{zz}), \\ [u]_2^2 &= |u_{xx}|^2 + |u_{yy}|^2 + |u_{zz}|^2, \end{aligned} \quad (2.7)$$

which make it a Hilbert space, thanks to the Dirichlet boundary condition and the Poincaré inequality and elliptic boundary regularity (see [ST10] and [STW12]). Furthermore by a result proven in [STW12], we know that $D(A) \subset H^2(\mathcal{M}) \cap H_0^1(\mathcal{M})$, and hence

$$D(A) \subset \Xi. \quad (2.8)$$

3 Parabolic regularization

For the sake of simplicity we will only treat the more complicated case when $d = 2$; the case when $d = 1$ is easier. To begin with, we recall the parabolic regularization introduced in [ST10] and [STW12], that is, for $\epsilon > 0$ “small”, we consider the parabolic equation,

$$\begin{cases} \frac{\partial u^\epsilon}{\partial t} + \Delta \frac{\partial u^\epsilon}{\partial x} + c \frac{\partial u^\epsilon}{\partial x} + u^\epsilon \frac{\partial u^\epsilon}{\partial x} + \epsilon Lu^\epsilon = f, \\ u^\epsilon(0) = u_0, \end{cases} \quad (3.1)$$

where

$$Lu^\epsilon := \frac{\partial^4 u^\epsilon}{\partial x^4} + \frac{\partial^4 u^\epsilon}{\partial y^4} + \frac{\partial^4 u^\epsilon}{\partial z^4},$$

supplemented with the boundary conditions (2.2), (2.4) and the additional boundary conditions

$$u_{yy}^\epsilon|_{y=\pm\frac{\pi}{2}} = u_{zz}^\epsilon|_{z=\pm\frac{\pi}{2}} = 0, \quad (3.2)$$

$$u_{xx}^\epsilon|_{x=0} = 0. \quad (3.3)$$

Note that since $u_{yy}^\epsilon = u_{zz}^\epsilon = 0$ at $x = 0$, (3.3) is equivalent to

$$\Delta u^\epsilon|_{x=0} = 0.$$

It is a classical result (see e.g. [Lio69], [LSU68] or also [STW12]) that there exists a unique solution to the parabolic problem which is sufficiently regular for all the subsequent calculations to be valid; in particular,

$$u^\epsilon \in L^2(0, T; H^4(\mathcal{M})) \cap C^1([0, T]; \Xi). \quad (3.4)$$

3.1 Global bounds independent of ϵ for u^ϵ

Firstly, we prove the following global bounds:

Lemma 3.1. *We assume that*

$$u_0 \in L^2(\mathcal{M}), \quad (3.5)$$

$$f \in L^2(0, T; L^2(\mathcal{M})), \quad (3.6)$$

then, for every $T > 0$ the following estimates independent of ϵ hold:

$$\begin{cases} u^\epsilon \text{ is bounded in } L^\infty(0, T; L^2(\mathcal{M})), \\ u^\epsilon \text{ is bounded in } L^2(0, T; H_0^1(\mathcal{M})), \\ u_x^\epsilon(0, \cdot, \cdot) \text{ is bounded in } L^2(0, T; L^2(I_{x^+})). \end{cases} \quad (3.7)$$

Proof. As in [STW12], we multiply (3.1) with u^ϵ and xu^ϵ , integrate over \mathcal{M} and integrate by parts. \square

3.2 Local bounds independent of ϵ for u^ϵ

We first introduce a useful result:

Lemma 3.2. *Under the same assumptions as in Lemma 3.1, if we further suppose that*

$$f \in L^\infty(0, T; L^2(\mathcal{M})), \quad (3.8)$$

then we have

$$|u_x^\epsilon(t)|^2 \leq |u_t^\epsilon(t)|^2 + \kappa, \quad 0 \leq t \leq T, \quad (3.9)$$

where κ is a constant depending only on $|u_0|$, $|f|_{L^\infty(0, T; L^2(\mathcal{M}))}$ and T .

Proof. We rewrite (3.1)₁ as

$$\Delta \frac{\partial u^\epsilon}{\partial x} + c \frac{\partial u^\epsilon}{\partial x} + u^\epsilon \frac{\partial u^\epsilon}{\partial x} + \epsilon Lu^\epsilon = -\frac{\partial u^\epsilon}{\partial t} + f. \quad (3.10)$$

We multiply (3.10) by $(1+x)u^\epsilon$, integrate over \mathcal{M} , integrate by parts, and follow the same calculations as in [STW12]; we find when $\epsilon \leq \frac{1}{4}$,

$$\begin{aligned} |u_x^\epsilon|^2 &\leq - \int_{\mathcal{M}} u_t^\epsilon (1+x)u^\epsilon d\mathcal{M} + |f|^2 + |(1+x)u^\epsilon|^2 + c|u^\epsilon|^2 + c'|u^\epsilon|^6 \\ &\leq |u_t^\epsilon|^2 + |f|^2 + c'|u^\epsilon|^2 + c'|u^\epsilon|^6, \end{aligned} \quad (3.11)$$

Here and below c' indicates an absolute constant which may be different at each occurrence. Hence if we call ν a bound of $|u^\epsilon|_{L^\infty(0, T; L^2(\mathcal{M}))}$ as in (3.7)₁, we can set

$$\kappa = |f|_{L^\infty(0, T; L^2(\mathcal{M}))}^2 + c'\nu^2 + c'\nu^6 + |u_{0x}|^2; \quad (3.12)$$

and by (3.8) we obtain (3.9). Thus we have completed the proof of Lemma 3.2. \square

Now we are ready to prove the following result giving the local bounds on u^ϵ independent of ϵ :

Proposition 3.1. *Under the same assumptions as in Lemma 3.2, if we further suppose that*

$$f_t \in L^\infty(0, T; L^2(\mathcal{M})), \quad (3.13)$$

$$f \in L^2(0, T; L^2(I_x; H^2(I_{x^\perp}))), \quad (3.14)$$

$$Lu_0 \in L^2(\mathcal{M}), \quad (3.15)$$

$$\Delta u_{0x} + u_0 u_{0x} + cu_{0x} - f(0) \in L^2(\mathcal{M}), \quad (3.16)$$

$$\nabla^\perp u_0, u_{0yy}, u_{0zz} \in L^2(\mathcal{M}), \quad (3.17)$$

then there exists $T_* = \min(T, T_1)$,

$$T_1 = \frac{c_3}{\mu^4}, \quad (3.18)$$

$$\mu = \mu(\kappa, |f_t|_{L^\infty(0, T; L^2(\mathcal{M}))}, |Lu_0|, |\Delta u_{0x} + u_0 u_{0x} + cu_{0x} - f(0)|),$$

such that for every t , $0 \leq t \leq T_*$,

$$|u_t^\epsilon(t)| \lesssim \mu, \quad (3.19)$$

$$\int_0^{T_*} |\nabla u_t^\epsilon(s)|^2 ds \lesssim \mu, \quad (3.20)$$

$$|\nabla u^\epsilon(t)| \leq C(\mu), \quad (3.21)$$

$$\int_0^{T_*} |\nabla u_y^\epsilon(s)|^2 ds \leq C(\mu), \quad \int_0^{T_*} |\nabla u_z^\epsilon(s)|^2 ds \leq C(\mu), \quad (3.22)$$

$$|u_{yy}^\epsilon(t)| \leq C(\mu), \quad |u_{zz}^\epsilon(t)| \leq C(\mu), \quad (3.23)$$

$$\int_0^{T_*} |\nabla u_{yy}^\epsilon(s)|^2 ds \leq C(\mu), \quad \int_0^{T_*} |\nabla u_{zz}^\epsilon(s)|^2 ds \leq C(\mu), \quad (3.24)$$

$$\epsilon \int_0^{T_*} [u_{yy}^\epsilon]_2^2 ds \leq C(\mu), \quad \epsilon \int_0^{T_*} [u_{zz}^\epsilon]_2^2 ds \leq C(\mu), \quad (3.25)$$

where \lesssim means \leq up to a multiplicative constant independent of ϵ , the constant c_3 depends only on the data, and the constant $C(\mu)$ depends only on μ and the data and may be different at each occurrence.

Proof. We differentiate (3.1) in t , write $u_t^\epsilon = v^\epsilon$ and we find:

$$\begin{cases} \frac{\partial v^\epsilon}{\partial t} + \Delta \frac{\partial v^\epsilon}{\partial x} + c \frac{\partial v^\epsilon}{\partial x} + u^\epsilon \frac{\partial v^\epsilon}{\partial x} + v^\epsilon \frac{\partial u^\epsilon}{\partial x} + \epsilon Lv^\epsilon = f_t, \\ v^\epsilon(0) = u_{t0}^\epsilon = -\epsilon Lu_0 - \Delta u_{0x} - u_0 u_{0x} - cu_{0x} + f(0). \end{cases} \quad (3.26)$$

Thus when $\epsilon \leq 1$,

$$|u_{t0}^\epsilon| \leq |Lu_0| + |\Delta u_{0x} + u_0 u_{0x} + cu_{0x} - f(0)|. \quad (3.27)$$

From (3.15) and (3.16), we obtain

$$u_{t0}^\epsilon \text{ is bounded independently of } \epsilon \text{ in } L^2(\mathcal{M}). \quad (3.28)$$

Multiplying (3.26) by $(1+x)v^\epsilon$, integrating over \mathcal{M} and integrating by parts, dropping ϵ for the moment we find

- $\int_{\mathcal{M}} \frac{\partial v}{\partial t} (1+x)v d\mathcal{M} = \frac{1}{2} \frac{d}{dt} |\sqrt{1+x}v|^2,$
- $\int_{\mathcal{M}} \Delta \frac{\partial v}{\partial x} v d\mathcal{M} = \frac{1}{2} \int_{I_{x^\perp}} (v_x|_{x=0})^2 dx^\perp,$
- $\int_{\mathcal{M}} \Delta \frac{\partial v}{\partial x} xv d\mathcal{M} = \frac{3}{2} |v_x|^2 + \frac{1}{2} |\nabla^\perp v|^2,$
- $\int_{\mathcal{M}} cv_x(1+x)v d\mathcal{M} = -\frac{c}{2} \int_{\mathcal{M}} v^2 d\mathcal{M},$
- $\int_{\mathcal{M}} uv_x(1+x)v d\mathcal{M} = -\frac{1}{2} \int_{\mathcal{M}} (1+x)u_x v^2 d\mathcal{M} - \frac{1}{2} \int_{\mathcal{M}} uv^2 d\mathcal{M},$
- $\int_{\mathcal{M}} vu_x(1+x)v d\mathcal{M} = \int_{\mathcal{M}} (1+x)u_x v^2 d\mathcal{M},$
- $\int_{\mathcal{M}} f_t(1+x)v d\mathcal{M} \leq \frac{1}{2} |f_t|^2 + \frac{1}{2} |(1+x)v|^2 \leq \frac{1}{2} |f_t|^2 + |v|^2,$
- $\epsilon \int_{\mathcal{M}} \frac{\partial^4 v}{\partial x^4} (1+x)v dx dx^\perp = \epsilon |\sqrt{1+x}v_{xx}|^2 - \epsilon \int_{I_{x^\perp}} (v_x|_{x=0})^2 dx^\perp,$
- $\epsilon \int_{\mathcal{M}} \left(\frac{\partial^4 v}{\partial y^4} + \frac{\partial^4 v}{\partial z^4} \right) (1+x)v d\mathcal{M} = \epsilon \left(|\sqrt{1+x}v_{yy}|^2 + |\sqrt{1+x}v_{zz}|^2 \right).$

Hence we arrive, when $\epsilon \leq \frac{1}{4}$, at

$$\begin{aligned}
& \frac{d}{dt} |\sqrt{1+x}u_t^\epsilon|^2 + |\nabla u_t^\epsilon|^2 + \frac{1}{4} |u_{tx}^\epsilon|_{x=0}|_{L^2(I_{x^\perp})}^2 \\
& \quad + 2\epsilon \left(|\sqrt{1+x}u_{txx}^\epsilon|^2 + |\sqrt{1+x}u_{tyy}^\epsilon|^2 + |\sqrt{1+x}u_{tzz}^\epsilon|^2 \right) \\
& \leq \left| \int_{\mathcal{M}} ((1+x)u_x^\epsilon - u^\epsilon) (u_t^\epsilon)^2 d\mathcal{M} \right| + (c+2) |u_t^\epsilon|^2 + |f_t|^2.
\end{aligned} \tag{3.29}$$

For the first term on the right-hand-side of (3.29), we have

$$\begin{aligned}
\left| \int_{\mathcal{M}} ((1+x)u_x^\epsilon - u^\epsilon) (u_t^\epsilon)^2 d\mathcal{M} \right| & \lesssim (\text{with } \sigma^\epsilon(t) := |u_x^\epsilon| + |u^\epsilon|) \\
& \lesssim \sigma^\epsilon(t) |u_t^\epsilon|_{L^4(\mathcal{M})}^2 \\
& \lesssim (\text{by } H^{3/4} \subset L^4 \text{ in } 3D) \\
& \lesssim \sigma^\epsilon(t) |u_t^\epsilon|^{1/2} |\nabla u_t^\epsilon|^{3/2} \\
& \leq c' (\sigma^\epsilon(t))^4 |u_t^\epsilon|^2 + \frac{1}{8} |\nabla u_t^\epsilon|^2 \\
& \leq (\text{by (3.9)}) \\
& \leq c' (|u_t^\epsilon|^6 + \kappa^2 |u_t^\epsilon|^2 + |u^\epsilon|^4 |u_t^\epsilon|^2) + \frac{1}{8} |\nabla u_t^\epsilon|^2.
\end{aligned} \tag{3.30}$$

Applying (3.30) to (3.29), we obtain

$$\begin{aligned}
\frac{d}{dt}|\sqrt{1+x}u_t^\epsilon|^2 + \frac{7}{8}|\nabla u_t^\epsilon|^2 + \frac{1}{4}|u_{tx}^\epsilon|_{x=0}|_{L^2(I_{x^\perp})}^2 \\
+ 2\epsilon(|\sqrt{1+x}u_{yxx}^\epsilon|^2 + |\sqrt{1+x}u_{yyy}^\epsilon|^2 + |\sqrt{1+x}u_{yzz}^\epsilon|^2) \\
\leq c_1(|u_t^\epsilon|^2 + 1)^3 + |f_t|^2 \\
\leq c_1(|\sqrt{1+x}u_t^\epsilon|^2 + 1)^3 + |f_t|^2.
\end{aligned} \tag{3.31}$$

where c_1 depends only on κ . Setting $|\sqrt{1+x}u_t^\epsilon|^2 + 1 := \mathcal{Y}^\epsilon$, (3.31) implies that

$$\frac{d}{dt}\mathcal{Y}^\epsilon \leq c_2(\mathcal{Y}^\epsilon)^3, \tag{3.32}$$

with $c_2 := c_1 + |f_t|_{L^\infty(0,T;L^2(\mathcal{M}))}^2$. Thus

$$\mathcal{Y}^\epsilon(t) \leq 2\mu_0^2, \quad 0 \leq t \leq \frac{3}{8c_2\mu_0^4}, \tag{3.33}$$

where μ_0 is a bound of $\sqrt{\mathcal{Y}^\epsilon(0)}$ independent of ϵ as provided by (3.28). Now (3.33) implies that

$$|u_t^\epsilon(t)| \lesssim \mu_0, \quad 0 \leq t \leq \frac{3}{8c_2\mu_0^4}. \tag{3.34}$$

Then by (3.34) and (3.9) we deduce that

$$|u_x^\epsilon(t)| \lesssim \mu, \quad 0 \leq t \leq T_*. \tag{3.35}$$

with $\mu := \mu_0 + \sqrt{\kappa}$, and $T_* = \min(T, T_1)$,

$$T_1 = \frac{c_3}{\mu^4} \leq \frac{3}{8c_2\mu_0^4}. \tag{3.36}$$

By (3.34), (3.36) and (3.31) we obtain (3.20).

We multiply (3.1) by $(1+x)u_{yy}^\epsilon$, integrate over \mathcal{M} and integrate by parts, dropping ϵ for the moment, we find

$$\begin{aligned}
& \bullet \int_{\mathcal{M}} u_t (1+x)u_{yy} d\mathcal{M} = -\frac{1}{2} \frac{d}{dt} |\sqrt{1+x}u_y|^2, \\
& \bullet \int_{\mathcal{M}} u_{xxx} (1+x)u_{yy} d\mathcal{M} = -\frac{3}{2} |u_{xy}^2| - \frac{1}{2} |u_{xy}^2|_{x=0}|_{L^2(I_y)}, \\
& \bullet \int_{\mathcal{M}} u_{xyy} (1+x)u_{yy} d\mathcal{M} = -\frac{1}{2} |u_{yy}^2|, \\
& \bullet \int_{\mathcal{M}} u_{xzz} (1+x)u_{yy} d\mathcal{M} = -\frac{1}{2} |u_{zy}^2|,
\end{aligned}$$

- $\int_{\mathcal{M}} cu_x(1+x)u_{yy} d\mathcal{M} = \frac{c}{2} \int_{\mathcal{M}} u_y^2 d\mathcal{M},$
- $\int_{\mathcal{M}} uu_x(1+x)u_{yy} d\mathcal{M} = \frac{1}{2} \int_{\mathcal{M}} u_y^2 (u - (1+x)u_x) d\mathcal{M},$
- $\begin{aligned} \epsilon \int_{\mathcal{M}} u_{xxxx}(1+x)u_{yy} d\mathcal{M} &= -\epsilon \int_{\mathcal{M}} u_{xxxxy}(1+x)u_y d\mathcal{M} \\ &= \epsilon \int_{\mathcal{M}} u_{xxxxy}u_y d\mathcal{M} + \epsilon \int_{d\mathcal{M}} u_{xxxxy}(1+x)u_{xy} d\mathcal{M} \\ &= -2\epsilon \int_{\mathcal{M}} u_{xxy}u_{xy}d\mathcal{M} - \epsilon \int_{\mathcal{M}} (1+x)u_{xxy}^2 d\mathcal{M} \\ &= \epsilon \int_{\mathcal{M}} u_{xy}^2|_{x=0} d\mathcal{M} - \epsilon \int_{\mathcal{M}} (1+x)u_{xxy}^2 d\mathcal{M}, \end{aligned}$
- $\epsilon \int_{\mathcal{M}} u_{zzzz}(1+x)u_{yy} d\mathcal{M} = -\epsilon \int_{\mathcal{M}} (1+x)u_{zzz}^2 d\mathcal{M},$
- $\epsilon \int_{\mathcal{M}} u_{yyyy}(1+x)u_{yy} d\mathcal{M} = -\epsilon \int_{\mathcal{M}} (1+x)u_{yyy}^\epsilon d\mathcal{M},$
- $\int_{\mathcal{M}} f(1+x)u_{yy} d\mathcal{M} = - \int_{\mathcal{M}} f_y(1+x)u_y d\mathcal{M} \leq \frac{1}{2}|(1+x)u_y|^2 + \frac{1}{2}|f_y|^2.$

Hence when $\epsilon \leq \frac{1}{4}$, we have

$$\begin{aligned}
&\frac{d}{dt}|\sqrt{1+x}u_y^\epsilon|^2 + |\nabla u_y^\epsilon|^2 + \frac{1}{4}|u_{xy}^\epsilon|_{x=0}|_{L^2(I_y)}^2 \\
&\quad + 2\epsilon(|\sqrt{1+x}u_{yxx}^\epsilon|^2 + |\sqrt{1+x}u_{yyy}^\epsilon|^2 + |\sqrt{1+x}u_{yzz}^\epsilon|^2) \\
&\leq \left| \int_{\mathcal{M}} (u_y^\epsilon)^2 ((1+x)u_x^\epsilon - u^\epsilon) d\mathcal{M} \right| + (c+2)|u_y^\epsilon|^2 + |f_y|^2.
\end{aligned} \tag{3.37}$$

For the first term on the right-hand-side of (3.37), we find

$$\begin{aligned}
\left| \int_{\mathcal{M}} (u_y^\epsilon)^2 ((1+x)u_x^\epsilon - u^\epsilon) d\mathcal{M} \right| &\lesssim (\text{with } \sigma^\epsilon(t) := |u_x^\epsilon| + |u^\epsilon|) \\
&\lesssim \sigma^\epsilon(t)|u_y^\epsilon|_{L^4(\mathcal{M})}^2 \\
&\leq c'(\sigma^\epsilon(t))^4|u_y^\epsilon|^2 + \frac{1}{8}|\nabla u_y^\epsilon|^2 \\
&\leq (\text{by (3.35)}) \\
&\leq c'(\mu^4 + |u^\epsilon|^4)|u_y^\epsilon|^2 + \frac{1}{8}|\nabla u_y^\epsilon|^2, \quad 0 \leq t \leq T_*.
\end{aligned} \tag{3.38}$$

Applying (3.38) to (3.37), we find

$$\begin{aligned}
&\frac{d}{dt}|\sqrt{1+x}u_y^\epsilon|^2 + \frac{7}{8}|\nabla u_y^\epsilon|^2 + \frac{1}{4}|u_{xy}^\epsilon|_{x=0}|_{L^2(I_y)}^2 \\
&\quad + 2\epsilon(|\sqrt{1+x}u_{yxx}^\epsilon|^2 + |\sqrt{1+x}u_{yyy}^\epsilon|^2 + |\sqrt{1+x}u_{yzz}^\epsilon|^2) \\
&\leq c'\mu^4|u_y^\epsilon|^2 + |f_y|^2 \\
&\leq c'\mu^4|\sqrt{1+x}u_y^\epsilon|^2 + |f_y|^2, \quad 0 \leq t \leq T_*.
\end{aligned} \tag{3.39}$$

We can then close the Gronwall inequality on the time interval $(0, T_*)$, and obtain

$$\begin{aligned} |\sqrt{1+x} u_y^\epsilon(t)|^2 &\leq C(\mu) \left(|\sqrt{1+x} u_{0y}|^2 + \int_0^{T_*} |f_y(s)|^2 ds \right) \\ &\leq C(\mu, |u_{0y}|, |f_y|_{L^2(0, T; L^2(\mathcal{M}))}), \quad 0 \leq t \leq T_*, \end{aligned}$$

which implies

$$|u_y^\epsilon(t)| \leq C(\mu), \quad 0 \leq t \leq T_*. \quad (3.40)$$

By (3.40) and (3.39) we obtain

$$\int_0^{T_*} |\nabla u_y^\epsilon(s)|^2 ds \leq C(\mu). \quad (3.41)$$

Similarly, we can obtain the same kind of estimates for u_z^ϵ , ∇u_z^ϵ , that is

$$|u_z^\epsilon(t)| \leq C(\mu), \quad 0 \leq t \leq T_*, \quad (3.42)$$

$$\int_0^{T_*} |\nabla u_z^\epsilon(s)|^2 ds \leq C(\mu). \quad (3.43)$$

From (3.35), (3.40) and (3.42) we obtain (3.21).

We then multiply (3.1) by $(1+x)u_{yyyy}^\epsilon$, integrate over \mathcal{M} and integrate by parts, to find

$$\begin{aligned} &\bullet \int_{\mathcal{M}} u_t (1+x) u_{yyyy} d\mathcal{M} = \frac{1}{2} \frac{d}{dt} |\sqrt{1+x} u_{yy}|^2, \\ &\bullet \int_{\mathcal{M}} u_{xxx} (1+x) u_{yyyy} d\mathcal{M} = \frac{3}{2} |u_{xyy}|^2 + \frac{1}{2} |u_{xyy}|_{x=0}^2_{L^2(I_y)}, \\ &\bullet \int_{\mathcal{M}} u_{xyy} (1+x) u_{yyyy} d\mathcal{M} = \frac{1}{2} |u_{yyy}^2|^2, \\ &\bullet \int_{\mathcal{M}} u_{xzz} (1+x) u_{yyyy} d\mathcal{M} = \frac{1}{2} |u_{zyy}^2|^2, \\ &\bullet \int_{\mathcal{M}} cu_x (1+x) u_{yyyy} d\mathcal{M} = -\frac{c}{2} |u_{yy}|^2, \\ &\bullet \int_{\mathcal{M}} uu_x (1+x) u_{yyyy} d\mathcal{M} \\ &\quad = - \int_{\mathcal{M}} u_y u_x (1+x) u_{yyy} d\mathcal{M} - \int_{\mathcal{M}} uu_{xy} (1+x) u_{yyy} d\mathcal{M} \\ &\quad = \int_{\mathcal{M}} u_{yy}^2 u_x (1+x) d\mathcal{M} + 2 \int_{\mathcal{M}} u_y u_{xy} (1+x) u_{yy} d\mathcal{M} + \int_{\mathcal{M}} uu_{xyy} (1+x) u_{yy} d\mathcal{M} \end{aligned}$$

$$\begin{aligned}
&= \int_{\mathcal{M}} u_{yy}^2 u_x (1+x) d\mathcal{M} + \int_{\mathcal{M}} \frac{\partial (u_y^2)}{\partial x} (1+x) u_{yy} d\mathcal{M} + \frac{1}{2} \int_{\mathcal{M}} u \frac{\partial (u_{yy}^2)}{\partial x} (1+x) d\mathcal{M} \\
&= \frac{1}{2} \int_{\mathcal{M}} u_{yy}^2 u_x (1+x) d\mathcal{M} - \int_{\mathcal{M}} u_y^2 u_{yy} d\mathcal{M} - \int_{\mathcal{M}} u_y^2 (1+x) u_{xyy} d\mathcal{M} - \frac{1}{2} \int_{\mathcal{M}} u u_{yy}^2 d\mathcal{M} \\
&= \frac{1}{2} \int_{\mathcal{M}} (u_x (1+x) - u) u_{yy}^2 d\mathcal{M} - \int_{\mathcal{M}} u_y^2 u_{yy} d\mathcal{M} - \int_{\mathcal{M}} u_y^2 (1+x) u_{xyy} d\mathcal{M} \\
\bullet \int_{\mathcal{M}} u_{xxxx} (1+x) u_{yyyy} d\mathcal{M} &= -\epsilon \int_{\mathcal{M}} u_{xxxxy} (1+x) u_{yyy} d\mathcal{M} \\
&= -\epsilon \int_{\mathcal{M}} u_{xxxxy} u_{yyy} d\mathcal{M} + \epsilon \int_{\mathcal{M}} u_{xxxxy} (1+x) u_{xyyy} d\mathcal{M} \\
&= -\epsilon \int_{I_{x^\perp}} u_{xyy}^2|_{x=0} dI_{x^\perp} + \epsilon \int_{\mathcal{M}} (1+x) u_{xxyy}^2 d\mathcal{M}, \\
\bullet \int_{\mathcal{M}} u_{yyyy} (1+x) u_{yyyy} d\mathcal{M} &= \epsilon |\sqrt{1+x} u_{yyyy}|^2, \\
\bullet \int_{\mathcal{M}} u_{zzzz} (1+x) u_{yyyy} d\mathcal{M} &= \epsilon |\sqrt{1+x} u_{zzyy}|^2, \\
\bullet \int_{\mathcal{M}} f(1+x) u_{yyyy} d\mathcal{M} &= - \int_{\mathcal{M}} f_{yy} (1+x) u_{yy} d\mathcal{M} \leq \frac{1}{2} |(1+x) u_{yy}|^2 + \frac{1}{2} |f_{yy}|^2.
\end{aligned}$$

Hence when $\epsilon \leq \frac{1}{4}$,

$$\begin{aligned}
&\frac{d}{dt} |\sqrt{1+x} u_{yy}^\epsilon|^2 + |\nabla u_{yy}^\epsilon|^2 + \frac{1}{4} |u_{xxy}^\epsilon|_{x=0}^2|_{L^2(I_y)} \\
&\quad + 2\epsilon (|\sqrt{1+x} u_{yyxx}^\epsilon|^2 + |\sqrt{1+x} u_{yyyy}^\epsilon|^2 + |\sqrt{1+x} u_{yyzz}^\epsilon|^2) \\
&\quad \leq \left| \int_{\mathcal{M}} ((1+x) u_x^\epsilon - u^\epsilon) (u_{yy}^\epsilon)^2 d\mathcal{M} \right| + 2 \left| \int_{\mathcal{M}} u_y^2 u_{yy} d\mathcal{M} \right| \\
&\quad \quad + 2 \left| \int_{\mathcal{M}} u_y^2 (1+x) u_{xyy} d\mathcal{M} \right| + (c+1) |u_{yy}^\epsilon|^2 + |f_{yy}|^2 \\
&\quad := I_1^\epsilon + I_2^\epsilon + I_3^\epsilon + (c+1) |u_{yy}^\epsilon|^2 + |f_{yy}|^2.
\end{aligned} \tag{3.44}$$

For I_1^ϵ , by the similar calculations in (3.38) we deduce

$$I_1^\epsilon \leq c' (\mu^4 + |u^\epsilon|^4) |u_{yy}^\epsilon|^2 + \frac{1}{8} |\nabla u_{yy}^\epsilon|^2, \quad 0 \leq t \leq T_*. \tag{3.45}$$

For I_2^ϵ we have

$$\begin{aligned}
I_2^\epsilon &\leq 2 |u_y^\epsilon|_{L^4(\mathcal{M})}^2 |u_{yy}^\epsilon| \\
&\lesssim |u_y^\epsilon|^{1/2} |\nabla u_y^\epsilon|^{3/2} |u_{yy}^\epsilon| \\
&\leq (\text{by (3.40)}) \\
&\leq C(\mu) |\nabla u_y^\epsilon|^{3/2} |u_{yy}^\epsilon| \\
&\leq C(\mu) |\nabla u_y^\epsilon|^{3/2} |u_{yy}^\epsilon|^2 + C(\mu) |\nabla u_y^\epsilon|^{3/2}, \quad 0 \leq t \leq T_*.
\end{aligned} \tag{3.46}$$

For I_3^ϵ we have

$$\begin{aligned}
I_3^\epsilon &\leq 2|u_y^\epsilon|_{L^4(\mathcal{M})}^2 |u_{xyy}^\epsilon| \\
&\lesssim |u_y^\epsilon|^{1/2} |\nabla u_y^\epsilon|^{3/2} |u_{xyy}^\epsilon| \\
&\leq \frac{1}{8} |u_{xyy}^\epsilon|^2 + c' |u_y^\epsilon| |\nabla u_y^\epsilon|^3 \\
&\leq (\text{by Jensen's inequality, } |\nabla u_y^\epsilon|^3 \lesssim |u_{xy}^\epsilon|^3 + |u_{yy}^\epsilon|^3 + |u_{yz}^\epsilon|^3) \\
&\leq \frac{1}{8} |u_{xyy}^\epsilon|^2 + c' |u_y^\epsilon| |u_{xy}^\epsilon|^3 + c' |u_y^\epsilon| |u_{yy}^\epsilon|^3 + c' |u_y^\epsilon| |u_{yz}^\epsilon|^3 \\
&:= \frac{1}{8} |u_{xyy}^\epsilon|^2 + J_4^\epsilon + J_5^\epsilon + J_6^\epsilon.
\end{aligned} \tag{3.47}$$

We now estimate J_4^ϵ . We observe that since $u_x^\epsilon = 0$ at $y = \pm \frac{\pi}{2}$,

$$|u_{xy}^\epsilon|^2 = \int_{\mathcal{M}} (u_{xy}^\epsilon)^2 d\mathcal{M} = - \int_{\mathcal{M}} u_x^\epsilon u_{xyy}^\epsilon d\mathcal{M} \leq c' |u_x^\epsilon| |u_{xyy}^\epsilon|. \tag{3.48}$$

Thus we have

$$\begin{aligned}
J_4^\epsilon &\leq c' |u_y^\epsilon| |u_x^\epsilon|^{3/2} |u_{xyy}^\epsilon|^{3/2} \\
&\leq c' |\nabla u^\epsilon|^{5/2} |u_{xyy}^\epsilon|^{3/2} \\
&\leq (\text{by (3.21) which is already proven}) \\
&\leq C(\mu)^{5/2} |u_{xyy}^\epsilon|^{3/2} \\
&\leq C(\mu)^{10} + \frac{1}{8} |u_{xyy}^\epsilon|^2, \quad 0 \leq t \leq T_*.
\end{aligned} \tag{3.49}$$

Similarly for J_6^ϵ , since $u_z^\epsilon = 0$ at $y = \pm \frac{\pi}{2}$, we can apply the intermediate derivative theorem to u_z^ϵ , and deduce that $|u_{zyy}^\epsilon|^2 \leq c' |u_z^\epsilon| |u_{zyy}^\epsilon|$. Hence by estimates similar as in (3.49) we have

$$J_6^\epsilon \leq C(\mu)^{10} + \frac{1}{8} |u_{zyy}^\epsilon|^2, \quad 0 \leq t \leq T_*. \tag{3.50}$$

To estimate J_5^ϵ , by (3.40) we have

$$J_5^\epsilon \leq C(\mu) |u_{yy}^\epsilon|^3, \quad 0 \leq t \leq T_*. \tag{3.51}$$

Collecting the estimates in (3.49), (3.51) and (3.50), along with (3.47) we obtain

$$I_3^\epsilon \leq \frac{3}{8} |\nabla u_{yy}^\epsilon|^2 + C(\mu) + C(\mu) |u_{yy}^\epsilon|^3, \quad 0 \leq t \leq T_*. \tag{3.52}$$

Collecting the estimates in (3.45), (3.46) and (3.52), along with (3.44) we obtain

$$\begin{aligned}
&\frac{d}{dt} |\sqrt{1+x} u_{yy}^\epsilon|^2 + \frac{1}{2} |\nabla u_{yy}^\epsilon|^2 + 2\epsilon (|\sqrt{1+x} u_{yyxx}^\epsilon|^2 + |\sqrt{1+x} u_{yyy}^\epsilon|^2 + |\sqrt{1+x} u_{yyzz}^\epsilon|^2) \\
&\leq c' \left(\mu^4 + |u^\epsilon|^4 + C(\mu) |\nabla u_y^\epsilon|^{3/2} + C(\mu) |u_{yy}^\epsilon| + c + 1 \right) |u_{yy}^\epsilon|^2 \\
&\quad + C(\mu) |\nabla u_y^\epsilon|^{3/2} + C(\mu) + |f_{yy}^\epsilon|^2, \quad 0 \leq t \leq T_*.
\end{aligned} \tag{3.53}$$

In particular, setting $\eta^\epsilon(t) = c'(\mu^4 + |u^\epsilon|^4 + C(\mu)|\nabla u_y^\epsilon|^{3/2} + C(\mu)|u_{yy}^\epsilon| + c + 1)$, from (3.53) we infer that

$$\frac{d}{dt}|\sqrt{1+x}u_{yy}^\epsilon|^2 \leq \eta^\epsilon(t)|\sqrt{1+x}u_{yy}^\epsilon|^2 + C(\mu)|\nabla u_y^\epsilon|^{3/2} + C(\mu) + |f_{yy}|^2, \quad 0 \leq t \leq T_*. \quad (3.54)$$

Since $|\nabla u_y^\epsilon|^{3/2} \leq |\nabla u_y^\epsilon|^2 + c'$, along with (3.41) we deduce

$$\int_0^{T_*} \eta^\epsilon(s) ds \leq C(\mu).$$

We can then close the Gronwall inequality on the time interval $(0, T_*)$ in (3.53), and obtain

$$|\sqrt{1+x}u_{yy}^\epsilon(t)| \leq C(\mu, |u_{0yy}|, |f_{yy}|_{L^2(0,T;L^2(\mathcal{M}))}), \quad 0 \leq t \leq T_*,$$

which implies

$$|u_{yy}^\epsilon(t)| \leq C(\mu), \quad 0 \leq t \leq T_*. \quad (3.55)$$

By (3.55) and (3.44) we obtain

$$\int_0^{T_*} |\nabla u_{yy}^\epsilon(s)|^2 ds \leq C(\mu), \quad (3.56)$$

$$\epsilon \int_0^{T_*} [u_{yy}^\epsilon]_2^2 ds \leq C(\mu), \quad 0 \leq t \leq T_*. \quad (3.57)$$

Similarly we can obtain the same kind of estimates for u_{zz}^ϵ , ∇u_{zz}^ϵ and $\epsilon[u_{zz}^\epsilon]_2^2$.

Combining all the previous local bounds, we obtain (3.19)-(3.25). Hence we have completed the proof of Proposition 3.1. \square

3.3 A singular perturbation argument

We are now ready to show the local estimates for u_{xx}^ϵ and $u^\epsilon u_x^\epsilon$ by singular perturbation.

Proposition 3.2. *Under the same assumptions as in Proposition 3.1, we have*

$$u_{xx}^\epsilon \text{ is bounded independently of } \epsilon \text{ in } L^2(0, T_*, L^2(\mathcal{M})), \quad (3.58)$$

$$u^\epsilon u_x^\epsilon \text{ is bounded independently of } \epsilon \text{ in } L^2(0, T_*; L^2(\mathcal{M})). \quad (3.59)$$

Remark 3.1. *Note that by (3.58) and (3.23) we deduce that*

$$u^\epsilon \text{ is bounded independently of } \epsilon \text{ in } L^2(0, T_*, \Xi). \quad (3.60)$$

Remark 3.2. *We know that*

$$\begin{aligned} \int_{\mathcal{M}} (u^\epsilon u_x^\epsilon)^{3/2} d\mathcal{M} &\leq \left(\int_{\mathcal{M}} (u^\epsilon)^6 d\mathcal{M} \right)^{1/4} \left(\int_{\mathcal{M}} (u_x^\epsilon)^2 d\mathcal{M} \right)^{3/4} \\ &= |u^\epsilon|_{L^6(\mathcal{M})}^{3/2} |u_x^\epsilon|^{3/2} \\ &\leq (\text{by } H^1(\mathcal{M}) \subset L^6(\mathcal{M}) \text{ in } 3D) \\ &\lesssim |\nabla u^\epsilon|^3. \end{aligned}$$

Hence

$$\operatorname{ess\,sup}_{t \in (0, T_*)} |u^\epsilon u_x^\epsilon(t)|_{L^{3/2}(\mathcal{M})} \lesssim \operatorname{ess\,sup}_{t \in (0, T_*)} |\nabla u^\epsilon(t)|^2 \lesssim (\text{by (3.21)}) \lesssim C(\mu)^2,$$

which implies that

$$\begin{aligned} u^\epsilon u_x^\epsilon &\text{ is bounded independently of } \epsilon \text{ in } L^\infty(0, T_*; L^{3/2}(\mathcal{M})), \\ &\text{ and hence in } L^{3/2}(I_x; L^{3/2}((0, T_*) \times I_{x^\perp})). \end{aligned}$$

Thus we can apply Lemma 5.1 in the Appendix with $p = 3/2$ and $Y = L^{3/2}((0, T_*) \times I_{x^\perp})$, and obtain

$$u_{xx}^\epsilon \text{ is bounded independently of } \epsilon \text{ in } L^\infty(I_x; L^{3/2}((0, T_*) \times I_{x^\perp})). \quad (3.61)$$

However, to obtain more useful estimates as in (3.58) and (3.59), we need to use the following proof which provides a stronger result.

Proof of Proposition 3.2. We rewrite the regularized equation (3.1) as follows:

$$\begin{cases} u_{xxx}^\epsilon + u^\epsilon u_x^\epsilon + \epsilon u_{xxxx}^\epsilon = g^\epsilon, \\ u^\epsilon(0) = u^\epsilon(1) = u_x^\epsilon(1) = u_{xx}^\epsilon(0) = 0, \end{cases} \quad (3.62)$$

where $g^\epsilon := -u_t^\epsilon - \Delta^\perp u_x^\epsilon - cu_x^\epsilon - \epsilon u_{yyyy}^\epsilon - \epsilon u_{zzzz}^\epsilon + f$. Hence by (3.19), (3.24) and (3.25), we know that each term in g^ϵ is bounded independently of ϵ in $L^2(0, T_*, L^2(\mathcal{M}))$, and thus

$$g^\epsilon \text{ is bounded independently of } \epsilon \text{ in } L^2(0, T_*, L^2(\mathcal{M})). \quad (3.63)$$

Multiplying (3.62) by x and integrating in x from 0 to 1, we find

$$\begin{aligned} \bullet & \int_0^1 x u_{xxx} dx = - \int_0^1 u_{xx} dx + u_{xx} x \Big|_{x=0}^{x=1} = u_x \Big|_{x=0} + u_{xx} \Big|_{x=1}, \\ \bullet & \int_0^1 x u u_x dx = \int_0^1 \frac{\partial}{\partial x} \left(\frac{u^2}{2} \right) dx = - \frac{1}{2} \int_0^1 u^2 dx, \\ \bullet & \epsilon \int_0^1 x u_{xxxx} dx = - \epsilon \int_0^1 u_{xxx} dx + \epsilon u_{xxx} x \Big|_{x=0}^{x=1} = - \epsilon u_{xx} \Big|_{x=1} + \epsilon u_{xxx} \Big|_{x=1}. \end{aligned}$$

Hence

$$u_x^\epsilon \Big|_{x=0} + u_{xx}^\epsilon \Big|_{x=1} - \frac{1}{2} \int_0^1 (u^\epsilon)^2 dx - \epsilon u_{xx}^\epsilon \Big|_{x=1} + \epsilon u_{xxx}^\epsilon \Big|_{x=1} = \int_0^1 g^\epsilon x dx. \quad (3.64)$$

Integrating (3.62) in x from \tilde{x} to 1, we obtain

$$\begin{aligned} \bullet & \int_{\tilde{x}}^1 u_{xxx} dx = u_{xx} \Big|_{x=1} - u_{xx}, \\ \bullet & \int_{\tilde{x}}^1 u u_x dx = - \frac{1}{2} u^2, \\ \bullet & \epsilon \int_{\tilde{x}}^1 u_{xxxx} dx = \epsilon u_{xxx} \Big|_{x=1} - \epsilon u_{xxx}. \end{aligned}$$

Hence

$$u_{xx}^\epsilon|_{x=1} - u_{xx}^\epsilon - \frac{1}{2}(u^\epsilon)^2 + \epsilon u_{xxx}^\epsilon|_{x=1} - \epsilon u_{xxx}^\epsilon = \int_{\tilde{x}}^1 g^\epsilon dx. \quad (3.65)$$

Then (3.64) and (3.65) imply

$$u_x^\epsilon|_{x=0} - \frac{1}{2} \int_0^1 (u^\epsilon)^2 dx - \epsilon u_{xx}^\epsilon|_{x=1} + u_{xx}^\epsilon + \frac{1}{2}(u^\epsilon)^2 + \epsilon u_{xxx}^\epsilon = \int_0^1 g^\epsilon x dx - \int_{\tilde{x}}^1 g^\epsilon dx,$$

which we rewrite as

$$u_{xx}^\epsilon + \epsilon u_{xxx}^\epsilon = \epsilon u_{xx}^\epsilon|_{x=1} + h^\epsilon, \quad (3.66)$$

where

$$h^\epsilon = -u_x^\epsilon|_{x=0} + \frac{1}{2} \int_0^1 (u^\epsilon)^2 dx - \frac{1}{2}(u^\epsilon)^2 + \int_0^1 g^\epsilon x dx - \int_{\tilde{x}}^1 g^\epsilon dx. \quad (3.67)$$

Now we estimate the term $(u^\epsilon)^2$ in (3.67). Since

$$|(u^\epsilon)^2|^2 \leq |u^\epsilon|_{L^4(\mathcal{M})}^4 \lesssim (\text{by } H^{3/4}(\mathcal{M}) \subset L^4(\mathcal{M}) \text{ in 3D}) \lesssim |\nabla u^\epsilon|^3 |u^\epsilon|,$$

we have

$$\int_0^{T_*} |(u^\epsilon)^2|^2 ds \lesssim \int_0^{T_*} |\nabla u^\epsilon|^3 |u^\epsilon| ds \leq (\text{by (3.21) and the Poincaré inequality}) \lesssim C(\mu)^4 T_*. \quad (3.68)$$

Thus

$$(u^\epsilon)^2 \text{ is bounded independently of } \epsilon \text{ in } L^2(0, T_*; L^2(\mathcal{M})). \quad (3.69)$$

Applying (3.7)₄, (3.63) and (3.69) to (3.67) we find

$$h^\epsilon \text{ is bounded independently of } \epsilon \text{ in } L^2(0, T_*, L^2(\mathcal{M})). \quad (3.70)$$

Multiplying (3.66) by u_{xx}^ϵ , integrating in x from 0 to 1, we obtain

- $\epsilon \int_0^1 u_{xxx}^\epsilon u_{xx}^\epsilon dx = \frac{\epsilon}{2} u_{xx}^2|_{x=1},$
- $\epsilon \int_0^1 u_{xx}^\epsilon|_{x=1} u_{xx}^\epsilon dx = \epsilon u_{xx}^\epsilon|_{x=1} \int_0^1 u_{xx}^\epsilon dx = -\epsilon u_{xx}^\epsilon|_{x=1} u_x^\epsilon|_{x=0};$

hence we arrive at

$$\begin{aligned} \int_0^1 (u_{xx}^\epsilon)^2 dx + \frac{\epsilon}{2} (u_{xx}^\epsilon)^2|_{x=1} &= -\epsilon u_{xx}^\epsilon|_{x=1} u_x^\epsilon|_{x=0} + \int_0^1 u_{xx}^\epsilon h^\epsilon dx, \\ &\leq \frac{\epsilon}{4} (u_{xx}^\epsilon|_{x=1})^2 + c' \epsilon (u_x^\epsilon|_{x=0})^2 + \frac{1}{2} |u_{xx}^\epsilon|_{L^2(I_x)}^2 + \frac{1}{2} |h^\epsilon|_{L^2(I_x)}^2. \end{aligned}$$

Thus

$$\frac{1}{2} \int_0^1 (u_{xx}^\epsilon)^2 dx + \frac{\epsilon}{4} (u_{xx}^\epsilon|_{x=1})^2 \leq c' \epsilon (u_x^\epsilon|_{x=0})^2 + \frac{1}{2} |h^\epsilon|_{L^2(I_x)}^2. \quad (3.71)$$

We integrate both sides of (3.71) in I_{x^\perp} and then in time from 0 to T_* ; by (3.70) and (3.7)₄ we obtain (3.58). As in Remark 3.1, we thus have (3.60).

Now since

$$\begin{aligned} |u^\epsilon u_x^\epsilon|^2 &\leq |u^\epsilon|_{L^4(\mathcal{M})}^2 |u_x^\epsilon|_{L^4(\mathcal{M})}^2 \\ &\lesssim (\text{by } H^{3/4}(\mathcal{M}) \subset L^4(\mathcal{M}) \text{ in 3D}) \\ &\lesssim |u^\epsilon|^{1/2} |\nabla u^\epsilon|^{3/2} |u_x^\epsilon|^{1/2} [u^\epsilon]_2^{3/2} \\ &\lesssim |u^\epsilon|^2 |\nabla u^\epsilon|^4 + [u^\epsilon]_2^2, \end{aligned}$$

hence we obtain

$$\int_0^{T_*} |u^\epsilon u_x^\epsilon|^2 ds \lesssim \int_0^{T_*} |u^\epsilon|^2 |\nabla u^\epsilon|^4 ds + \int_0^{T_*} [u^\epsilon]_2^2 ds.$$

This together with (3.21) and (3.60) implies (3.59). \square

4 Passage to the limit

Using a compactness argument, we can pass to the limit in (3.1). Hence we obtain (2.1), with a function $u \in \mathcal{C}^1([0, T_*]; L^2(\mathcal{M})) \cap L^2(0, T; H_0^1(\mathcal{M}))$. Then we rewrite (2.1) as

$$u_{xxx} = -u_t - \Delta^\perp u_x - cu_x - uu_x - f. \quad (4.1)$$

From (3.19), (3.24) and (3.59), we infer that each term in the right-hand-side of (4.1) belongs to $L^2(0, T_*; L^2(\mathcal{M}))$, and hence

$$u_{xxx} \in L^2(0, T_*; L^2(\mathcal{M})). \quad (4.2)$$

Now we are ready to state the main result: the local existence of strong solutions.

Theorem 4.1. *The assumptions are the same as in Proposition 3.1, that is (3.5), (3.6), (3.8) and (3.13)-(3.17). We suppose also that the following compatibility conditions hold:*

$$u_0 = 0 \text{ on } \partial\mathcal{M}, \quad u_{0x}|_{x=1} = 0, \quad u_{0yy}|_{y=\pm\frac{\pi}{2}} = u_{0zz}|_{z=\pm\frac{\pi}{2}} = 0, \quad (4.3)$$

$$u_{t0} = 0 \text{ on } \partial\mathcal{M}, \quad \frac{\partial u_{t0}}{\partial x}|_{x=1} = 0, \quad \frac{\partial^2 u_{t0}}{\partial y^2}|_{y=\pm\frac{\pi}{2}} = \frac{\partial^2 u_{t0}}{\partial z^2}|_{z=\pm\frac{\pi}{2}} = 0, \quad (4.4)$$

where $u_{t0} = -\Delta u_{0x} - u_0 u_{0x} - cu_{0x} + f(0)$. Then there exists a local strong solution to (2.1)-(2.4) on some time interval $[0, T_*)$, $T_* > 0$ depending only on the data as in Proposition 3.1, such that

$$\nabla u, u_{yy}, u_{zz}, u_t \in L^\infty(0, T_*; L^2(\mathcal{M})), \quad (4.5)$$

$$u \in L^2(0, T_*; D(A)), \quad (4.6)$$

$$u \in L^2(0, T_*; \Xi), \quad (4.7)$$

$$u \in L^2(0, T_*; H^3(I_x; L^2(I_{x^\perp})) \cap H^3(I_{x^\perp}; L^2(I_x))), \quad (4.8)$$

$$u_t \in L^2(0, T_*; H^1(\mathcal{M})). \quad (4.9)$$

Moreover, we have for every $t \in (0, T_*)$,

$$u_{yy}(t)|_{y=\pm\frac{\pi}{2}} = u_{zz}(t)|_{z=\pm\frac{\pi}{2}} = 0. \quad (4.10)$$

Remark 4.1. We have proven that all the spatial derivatives of the third order of u are in $L^2(0, T_*; L^2(\mathcal{M}))$, except for u_{xxy} and u_{xxz} .

Proof. We rewrite (2.1) as

$$Au = -u_t - uu_x - f; \quad (4.11)$$

from (3.19) and (3.59) we know that each term on the right-hand side of (4.11) belongs to $L^2(0, T_*; L^2(\mathcal{M}))$. Hence Au belongs to the same space. We also know that $u_x(1, x^\perp, t) = 0$, $t \in [0, T]$ using the same argument as in [STW12]. Hence we obtain (4.6). Now by (2.8) and (4.6), we deduce (4.7).

By (3.22), we know that u_{yyy} , u_{zzz} both belong to $L^2(0, T_*; L^2(\mathcal{M}))$. Hence we can apply the trace theorem and pass to the limit on the boundary conditions in (3.2) to obtain (4.10).

The other results can be deduced directly from (3.19)-(3.24) and (4.2). \square

Remark 4.2. As for the periodic case, that is, (2.1) and the boundary and initial conditions (2.2), (2.3) and (2.5), the results are exactly the same as in the Dirichlet case discussed above. The reasoning is totally the same and therefore we skip it.

5 Appendix: a trace result

We recall a trace result from [STW12], which is used in the article.

Lemma 5.1. Let Y be a (not necessarily reflexive) Banach space and let $p \geq 1$. Assume that two sequences of functions $u^\epsilon, g^\epsilon \in L_x^p(I_x; Y)$ satisfy

$$\begin{cases} u_{xxx}^\epsilon + \epsilon u_{xxxx}^\epsilon = g^\epsilon, \\ u^\epsilon(0) = u^\epsilon(1) = u_x^\epsilon(1) = u_{xx}^\epsilon(0) = 0, \end{cases} \quad (5.1)$$

with g^ϵ bounded in $L_x^p(I_x; Y)$ as $\epsilon \rightarrow 0$. Then u_{xx}^ϵ (and hence u_x^ϵ , and u^ϵ) is bounded in $L_x^\infty(I_x; Y)$ as $\epsilon \rightarrow 0$. Furthermore if Y is reflexive, then for any subsequence $u^\epsilon \rightarrow u$ converging strongly or weakly in $L_x^q(I_x; Y)$, $1 < q < \infty$, $u_x^\epsilon(1)$ converges to $u_x(1)$ in Y (weakly at least), and hence $u_x(1) = 0$.

Acknowledgments

This work was partially supported by the National Science Foundation under the grants, DMS-0906440 and DMS 1206438, and by the Research Fund of Indiana University.

The author would like to thank Professor Roger Temam, Professor Jean-Claude Saut, and Professor Nathan Glatt-Holtz for their encouragements and suggestions.

References

- [BF13] E.S. Baykova and A. Faminskii, *On initial-boundary value problem in a strip for generalized two-dimensional Zakharov-Kuznetsov equation*, Advances Diff. Equations (2013), no. 18(7–8), 663–686.
- [BPS81] J. L. Bona, W. G. Pritchard, and L. R. Scott, *An evaluation of a model equation for water waves*, Philos. Trans. Roy. Soc. London Ser. A **302** (1981), no. 1471, 457–510. MR 633485 (83a:35088)
- [BPS83] ———, *A comparison of solutions of two model equations for long waves*, Fluid dynamics in astrophysics and geophysics (Chicago, Ill., 1981), Lectures in Appl. Math., vol. 20, Amer. Math. Soc., Providence, R.I., 1983, pp. 235–267. MR 716887 (84j:76011)
- [BSZ03] Jerry L. Bona, Shu Ming Sun, and Bing-Yu Zhang, *A nonhomogeneous boundary-value problem for the Korteweg-de Vries equation posed on a finite domain*, Comm. Partial Differential Equations **28** (2003), no. 7-8, 1391–1436. MR 1998942 (2004h:35195)
- [CG01a] Thierry Colin and Jean-Michel Ghidaglia, *An initial-boundary value problem for the Korteweg-de Vries equation posed on a finite interval*, Adv. Differential Equations **6** (2001), no. 12, 1463–1492. MR 1858429 (2002i:35160)
- [CG01b] Thierry Colin and Marguerite Gisclon, *An initial-boundary-value problem that approximate the quarter-plane problem for the Korteweg-de Vries equation*, Nonlinear Anal. **46** (2001), no. 6, Ser. A: Theory Methods, 869–892. MR 1859802 (2002i:35161)
- [DL13] Gleb G. Doronin and Nikolai A. Larkin, *Exponential decay for the linear Zakharov-Kuznetsov equation without critical domains restrictions*, arXiv:1305.6236 (2013).
- [Fam06] A. V. Faminskii, *On the nonlocal well-posedness of a mixed problem for the Zakharov-Kuznetsov equation*, Sovrem. Mat. Prilozh. (2006), no. 38, 135–148. MR 2469478 (2010a:35215)
- [Fam08] Andrei V. Faminskii, *Well-posed initial-boundary value problems for the Zakharov-Kuznetsov equation*, Electron. J. Differential Equations (2008), No. 127, 23. MR 2443150 (2009g:35261)
- [Lio69] J.-L. Lions, *Quelques méthodes de résolution des problèmes aux limites non linéaires*, Dunod, 1969. MR 0259693 (41 #4326)
- [LLS13] F. Linares, D. Lannes, and Jean-Claude Saut, *The Cauchy problem for the Euler-Poisson system and derivation of the Zakharov-Kuznetsov equation*, Progress in Nonlinear Differential Equations and their Applications, M. Cicognani, FL. Colombini and D. Del Santo Eds. **Vol 84** (2013), 183–215.
- [LS82] E.W. Laedke and K.H. Spatschek, *Growth rates of bending solitons*, J. Plasma Phys. **26** (1982), no. 3, 469–484.
- [LSU68] O. A. Ladyzenskaja, V. A. Solonnikov, and N. N. Ural'ceva, *Linear and quasilinear equations of parabolic type*, Translated from the Russian by S. Smith. Translations of Mathematical Monographs, Vol. 23, American Mathematical Society, Providence, R.I., 1968. MR 0241822 (39 #3159b)
- [LT13] Nikolai A. Larkin and Eduardo Tronco, *Regular solutions of the 2D Zakharov-Kuznetsov equation on a half-strip*, J. Differential Equations **254** (2013), no. 1, 81–101. MR 2983044
- [ST10] Jean-Claude Saut and Roger Temam, *An initial boundary-value problem for the Zakharov-Kuznetsov equation*, Adv. Differential Equations **15** (2010), no. 11-12, 1001–1031.
- [STW12] Jean-Claude Saut, Roger Temam, and Chuntian Wang, *An initial and boundary-value problem for the Zakharov-Kuznetsov equation in a bounded domain*, J. Math. Phys. **53** (2012), no. 11, 115612, 29. MR 3026557

[ZK74] V.E. Zakharov and E.A. Kuznetsov, *On three-dimensional solitons*, Sov. Phys. JETP **30** (1974), 285–286.