THE DISTRIBUTION OF 2-SELMER RANKS OF QUADRATIC TWISTS OF ELLIPTIC CURVES WITH PARTIAL TWO-TORSION

ZEV KLAGSBRUN AND ROBERT J. LEMKE OLIVER

Abstract. This paper presents a new result concerning the distribution of 2-Selmer ranks in the quadratic twist family of an elliptic curve over an arbitrary number field K with a single point of order two that does not have a cyclic 4-isogeny defined over its twodivision field. We prove that at least half of all the quadratic twists of such an elliptic curve have arbitrarily large 2-Selmer rank, showing that the distribution of 2-Selmer ranks in the quadratic twist family of such an elliptic curve differs from the distribution of 2-Selmer ranks in the quadratic twist family of an elliptic curve having either no rational two-torsion or full rational two-torsion.

1. INTRODUCTION

1.1. Distributions of Selmer Ranks. Let E be an elliptic curve defined over a number field K and let $\text{Sel}_2(E/K)$ be its [2](#page-2-0)-Selmer group (see Section 2 for its definition). We define the 2-Selmer rank of E/K , denoted $d_2(E/K)$, by

$$
d_2(E/K) = \dim_{\mathbb{F}_2} \mathrm{Sel}_2(E/K) - \dim_{\mathbb{F}_2} E(K)[2].
$$

In 1994, Heath-Brown proved that the 2-Selmer ranks of all of the quadratic twists of the congruent number curve E/\mathbb{Q} given by $y^2 = x^3 - x$ had a particularly nice distribution. In particular, he showed that there are explicits constants $\alpha_0, \alpha_1, \alpha_2, \ldots$ summing to one such that

$$
\lim_{X \to \infty} \frac{|\{d \text{ squarefree } |d| < X : d_2(E^d/\mathbb{Q}) = r\}|}{|\{d \text{ squarefree } |d| < X\}|} = \alpha_r
$$

for every $r \in \mathbb{Z}^{\geq 0}$, where E^d is the quadratic twist of E by d [\[HB94\]](#page-9-0). This result was extended by Swinnerton-Dyer and Kane to all elliptic curves E over Q with $E(\mathbb{Q})[2] \simeq \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ that do not have a cyclic 4-isogeny defined over K $|$ Kan10, $|$ SD08.

A similar result was obtained by Klagsbrun, Mazur, and Rubin for elliptic curves E over a general number field K with Gal $(K(E[2])/K) \simeq S_3$, where squarefree d are replaced by quadratic characters of K and a suitable ordering of all such characters is taken [\[KMR13\]](#page-9-2).

In this work we show that this type of result does not hold when $E(K)[2] \simeq \mathbb{Z}/2\mathbb{Z}$. In particular, we prove the following:

Theorem 1.1. For $d \in \mathcal{O}_K$, let χ_d be the quadratic character of K that cuts out the extension $K(\sqrt{d})$ and define

$$
C(K,X):=\{\chi_d:|\mathbf{N}_{K/\mathbb{Q}}d|
$$

Let E be an elliptic curve defined over K with $E(K)[2] \simeq \mathbb{Z}/2\mathbb{Z}$ that does not have a cyclic isogeny defined over $K(E[2])$. Then for any fixed r,

$$
\liminf_{X \to \infty} \frac{|\{\chi \in C(K, X) : d_2(E^{\chi}/K) \ge r\}|}{|C(K, X)|} \ge \frac{1}{2}
$$

where E^{χ} is the quadratic twist of E by any $d \in \mathcal{O}_K$ with $\chi_d = \chi$.

In particular, this shows that there is not a distribution function on 2-Selmer ranks within the quadratic twist family of E.

Theorem [1.1](#page-0-0) is an easy consequence of the following result.

Theorem 1.2. Let E be an elliptic curve defined over K with $E(K)[2] \simeq \mathbb{Z}/2\mathbb{Z}$ that does not have a cyclic isogeny defined over $K(E[2])$. Then the normalized distribution

$$
\frac{P_r(\mathcal{T}(E/E'), X)}{\sqrt{\frac{1}{2}\log\log X}}
$$

converges weakly to the Gaussian distribution

$$
G(z) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{z} e^{\frac{-w^2}{2}} dw,
$$

where

$$
P_r(\mathcal{T}(E/E'), X) = \frac{|\{\chi \in C(K, X) : \text{ord}_2 \mathcal{T}(E^{\chi}/E'^{\chi}) \le r\}|}{|C(K, X)|}
$$

for $X \in \mathbb{R}^+$, $r \in \mathbb{Z}^{\geq 0}$, and $\mathcal{T}(E^{\chi}/E'^{\chi})$ as defined in Section [2.](#page-2-0)

In turn, Theorem [1.2](#page-1-0) follows from a variant of the Erdős-Kac theorem for quadratic characters of number fields. Let $C(K)$ denote the set of all quadratic characters of K. For any $\chi \in C(K)$, we can associate a unique squarefree ideal D_{χ} to χ by taking D_{χ} to be the squarefree part of the ideal $\langle d \rangle$, where d is any element of \mathcal{O}_K such that $\chi_d = \chi$. We say that a function f on the ideals of \mathcal{O}_K is **additive** if $f(\mathfrak{a}\mathfrak{a}') = f(\mathfrak{a}) + f(\mathfrak{a}')$ whenever the ideals \mathfrak{a} and \mathfrak{a}' are relatively prime; we define $f(\chi)$ for $\chi \in C(K)$ to be $f(D_{\chi})$.

To an additive function f, we attach quantities $\mu_f(X)$ and $\sigma_f(X)$, defined to be

$$
\mu_f(X) := \sum_{\mathbf{N}\mathfrak{p} < X} \frac{f(\mathfrak{p})}{\mathbf{N}\mathfrak{p}}, \quad \sigma_f(X) := \left(\sum_{\mathbf{N}\mathfrak{p} < X} \frac{f(\mathfrak{p})^2}{\mathbf{N}\mathfrak{p}}\right)^{1/2},
$$

and we prove the following.

Theorem 1.3. Suppose that f is an additive function such that $0 \leq f(\mathfrak{p}) \leq 1$ for every prime p. If $\sigma_f(X) \to \infty$ as $X \to \infty$, then

$$
\lim_{X \to \infty} \frac{|\{\chi \in C(K, X) : f(\chi) - \mu_f(X) \le z \cdot \sigma_f(X)\}|}{|C(K, X)|} = G(z),
$$

In the special case where $K = \mathbb{Q}$, Theorem [1.1](#page-0-0) follows from results recently obtained by Xiong about the distribution of $\dim_{\mathbb{F}_2} \text{Sel}_{\phi}(E/\mathbb{Q})$ in quadratic twist families [\[Xio13\]](#page-10-1). We are able to obtain results over general number fields by applying different methods to the coarser question about the distribution of $\text{ord}_2T(E/E')$ in twist families (see Theorem [1.2\)](#page-1-0).

Remark 1.4. The methods in the paper can easily be adapted to studying cubic twists of the j-invariant 0 curve $y^2 = x^3 + k$ for any number field K with $K(\sqrt{k}, \sqrt{-3})/K$ biquadratic. In that case, Theorem [1.1](#page-0-0) holds with 2-Selmer rank replaced by 3-Selmer rank and quadratic characters replaced by cubic characters.

1.2. Layout. We begin in Section [2](#page-2-0) by recalling the definitions of the 2-Selmer group and the Selmer groups associated with a 2-isogeny ϕ and presenting some of the connections between them. In Section [3,](#page-3-0) we examine the behavior of the local conditions for the ϕ -Selmer group under quadratic twist and show how that quantity $\mathcal{T}(E^{\chi}/E^{\prime\chi})$ can be related to the value $f(\chi)$ of an additive function f on $C(K)$. Theorem [1.3](#page-1-1) is proved in Section [4](#page-4-0) and we conclude with the proofs of Theorems [1.1](#page-0-0) and [1.2](#page-1-0) in Section [5.](#page-8-0)

Acknowledgement. The first author would like to express his thanks to Karl Rubin for his helpful comments and suggestions, to Ken Kramer for a series of valuable discussions, and to Michael Rael and Josiah Sugarman for helpful conversations regarding the Erdős-Kac theorem. The first author was supported by NSF grants DMS-0457481, DMS-0757807, and DMS-0838210.

2. Selmer Groups

We begin by recalling the definition of the 2-Selmer group. If E is an elliptic curve defined over a field K, then $E(K)/2E(K)$ maps into $H^1(K, E[2])$ via the Kummer map. The following diagram commutes for every place v of K, where δ is the Kummer map.

$$
E(K)/2E(K) \xrightarrow{\delta} H^1(K, E[2])
$$

\n
$$
\downarrow Res_v
$$

\n
$$
E(K_v)/2E(K_v) \xrightarrow{\delta} H^1(K_v, E[2])
$$

We define a distinguished local subgroup $H^1_f(K_v, E[2]) \subset H^1(K_v, E[2])$ as the image $\delta(E(K_v)/2E(K_v)) \subset H^1(K_v, E[2])$ for each place v of K and we define the 2-Selmer group of E/K , denoted $\text{Sel}_2(E/K)$, by

$$
\mathrm{Sel}_2(E/K) = \ker\left(H^1(K, E[2]) \xrightarrow{\sum res_v} \bigoplus_{v \text{ of } K} H^1(K_v, E[2]) / H^1_f(K_v, E[2])\right).
$$

The 2-Selmer group is a finite dimensional \mathbb{F}_2 -vector space that sits inside the exact sequence of \mathbb{F}_2 -vector spaces

$$
0 \to E(K)/2E(K) \to {\rm Sel}_2(E/K) \to {\rm III}(E/K)[2] \to 0
$$

where $III(E/K)$ is the Tate-Shafaravich group of E.

If $E(K)$ has a single point of order two, then there is a two-isogeny $\phi : E \to E'$ between E and E' with kernel $C = E(K)[2]$. This isogeny gives rise to two Selmer groups.

We have a short exact sequence of G_K modules

(1)
$$
0 \to C \to E(\overline{K}) \xrightarrow{\phi} E'(\overline{K}) \to 0
$$

which gives rise to a long exact sequence of cohomology groups

$$
0 \to C \to E(K) \xrightarrow{\phi} E'(K) \xrightarrow{\delta} H^1(K, C) \to H^1(K, E) \to H^1(K, E') \dots
$$

The map δ is given by $\delta(Q)(\sigma) = \sigma(R) - R$ where R is any point on $E(\overline{K})$ with $\phi(R) = Q$.

This sequence remains exact when we replace K by its completion K_v at any place v, which gives rise to the following commutative diagram.

$$
E'(K)/\phi(E(K)) \xrightarrow{\delta} H^1(K, C)
$$

\n
$$
\downarrow \qquad \qquad \downarrow Res_v
$$

\n
$$
E'(K_v)/\phi(E(K_v)) \xrightarrow{\delta} H^1(K_v, C)
$$

In a manner similar to how we defined the 2-Selmer group, we define distinguished local subgroups $H^1_{\phi}(K_v, C) \subset H^1(K_v, C)$ as the image of $E'(K_v)/\phi(E(K_v))$ under δ for each place v of K. We define the ϕ -**Selmer group of E**, denoted $\text{Sel}_{\phi}(E/K)$ as

$$
\mathrm{Sel}_{\phi}(E/K) = \ker \left(H^1(K, C) \xrightarrow{\sum res_v} \bigoplus_{v \text{ of } K} H^1(K_v, C) / H^1_{\phi}(K_v, C) \right).
$$

The isogeny ϕ on E gives gives rise to a dual isogeny $\hat{\phi}$ on E' with kernel $C' = \phi(E[2])$. Exchanging the roles of (E, C, ϕ) and $(E', C', \hat{\phi})$ in the above defines the $\hat{\phi}$ -Selmer group, $\mathrm{Sel}_{\hat{\phi}}(E'/K)$, as a subgroup of $H^1(K, C')$. The groups $\mathrm{Sel}_{\phi}(E/K)$ and $\mathrm{Sel}_{\hat{\phi}}(E'/K)$ are finite dimensional \mathbb{F}_2 -vector spaces and we can compare the sizes of the ϕ -Selmer group, the ϕ -Selmer group, and the 2-Selmer group using the following two theorems.

Theorem 2.1. The φ-Selmer group, the $\hat{\phi}$ -Selmer group, and the 2-Selmer group sit inside the exact sequence

(2)
$$
0 \to E'(K)[2]/\phi(E(K)[2]) \to \mathrm{Sel}_{\phi}(E/K) \to \mathrm{Sel}_{2}(E/K) \xrightarrow{\phi} \mathrm{Sel}_{\hat{\phi}}(E'/K).
$$

Proof. This is a well known diagram chase. See Lemma 2 in [\[FG08\]](#page-9-3) for example. \Box

The **Tamagawa ratio** $\mathcal{T}(E/E')$ defined as $\mathcal{T}(E/E')$ = $\left|\text{Sel}_{\phi}(E/K)\right|$ $\left|\operatorname{Sel}_{\hat{\phi}}(E'/K)\right|$ gives a second relationship between the \mathbb{F}_2 -dimensions of $\mathrm{Sel}_{\phi}(E/K)$ and $\mathrm{Sel}_{\hat{\phi}}(E'/K)$.

Theorem 2.2 (Cassels). The Tamagawa ratio $\mathcal{T}(E/E')$ is given by

$$
\mathcal{T}(E/E') = \prod_{v \text{ of } K} \frac{|H^1_\phi(K_v, C)|}{2}
$$

.

Proof. This is a combination of Theorem 1.1 and equations (1.22) and (3.4) in $|Cas65|$. \Box

Stepping back, we observe that if $\mathcal{T}(E/E') \geq 2^{r+2}$, then $d_{\phi}(E/K) \geq r+2$, and therefore by Theorem [2.1,](#page-3-1) $d_2(E/K) \geq r$. (If E does not have a cyclic 4-isogeny defined over K then we can in fact show that $\mathcal{T}(E/E') \geq 2^r$ implies that $d_2(E/K) \geq r$, but this is entirely unnecessary for our purposes.)

3. Local Conditions at Twisted Places

For the remainder of this paper, we will let E be an elliptic curve with $E(K)[2] \simeq \mathbb{Z}/2\mathbb{Z}$ and let $\phi : E \to E'$ be the isogeny with kernel $C = E(K)[2]$.

If $\mathfrak{p} \nmid 2$ is a prime where E has good reduction, then $H^1_\phi(K_{\mathfrak{p}}, C)$ is a 1-dimensional \mathbb{F}_2 subspace of $H^1(K_{\mathfrak{p}}, C)$ equal to the unramified local subgroup $H^1_u(K_{\mathfrak{p}}, C)$. If such a \mathfrak{p} is ramified in the extension F/K cut out by a character χ , then the twisted curve E^{χ} will have bad reduction at **p**. The following lemma addresses the size of $H^1_\phi(K_\mathfrak{p}, C^{\chi})$.

Lemma 3.1. Suppose $\mathfrak{p} \nmid 2$ is a prime where E has good reduction and \mathfrak{p} is ramified in the extension F/K cut out by χ .

- (i) If $E(K_{\mathfrak{p}})[2] \simeq \mathbb{Z}/2\mathbb{Z} \simeq E'(K_{\mathfrak{p}})[2]$, then $\dim_{\mathbb{F}_2} H^1_{\phi}(K_{\mathfrak{p}}, C^{\chi}) = 1$.
- (ii) If $E(K_{\mathfrak{p}})[2] \simeq \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z} \simeq E'(K_{\mathfrak{p}})[2]$, then $\dim_{\mathbb{F}_2} H^1_{\phi}(K_{\mathfrak{p}}, C^{\chi}) = 1$.
- (iii) If $E(K_{\mathfrak{p}})[2] \simeq \mathbb{Z}/2\mathbb{Z}$ and $E'(K_{\mathfrak{p}})[2] \simeq \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$, then $\dim_{\mathbb{F}_2} H^1_{\phi}(K_{\mathfrak{p}}, C^{\chi}) = 2$.
- (iv) If $E(K_{\mathfrak{p}})[2] \simeq \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ and $E'(K_{\mathfrak{p}})[2] \simeq \mathbb{Z}/2\mathbb{Z}$, then $\dim_{\mathbb{F}_2} H^1_{\phi}(K_{\mathfrak{p}}, C^{\chi}) = 0$.

Proof. From Lemma 3.7 in [\[Kla11\]](#page-9-5), we have

$$
E'^{\chi}(K_{\mathfrak{p}})[2^{\infty}]/\phi(E^{\chi}(K_{\mathfrak{p}})[2^{\infty}]) = E'^{\chi}(K_{\mathfrak{p}})[2]/\phi(E^{\chi}(K_{\mathfrak{p}})[2]).
$$

All four results then follow immediately.

Remark 3.2. Suppose that Δ and Δ' are discriminants of any integral models of E and E' respectively. The condition that $E(K_{\mathfrak{p}})[2] \simeq \mathbb{Z}/2\mathbb{Z}$ (resp. $E'(K_{\mathfrak{p}})[2] \simeq \mathbb{Z}/2\mathbb{Z}$) is equivalent to the condition that Δ (resp. Δ') is not a square in $K_{\mathfrak{p}}$. Which case of Lemma [3.1](#page-4-1) we are is therefore determined by the Legendre symbols $\left(\frac{\Delta}{n}\right)$ $\frac{\Delta}{\mathfrak{p}}$ and $\left(\frac{\Delta'}{\mathfrak{p}}\right)$ $\frac{\Delta'}{p}\bigg).$

We use Theorem [2.2](#page-3-2) and Lemma [3.1](#page-4-1) to relate $\text{ord}_2 \mathcal{T}(E^{\chi}/E^{\gamma})$ to the value $g(\chi)$ of an additive function g on $C(K)$ defined as follows: For $\chi \in C(K)$ cutting out F/K , let

(3)
$$
g(\chi) = \sum_{\substack{\mathfrak{p} \text{ ramified in } F/K}} \frac{\left(\frac{\Delta'}{\mathfrak{p}}\right) - \left(\frac{\Delta}{\mathfrak{p}}\right)}{2}
$$

That is, $g(\chi)$ roughly counts the difference between the number of primes ramified in F/K where condition *(iii)* of Proposition [3.1](#page-4-1) is satisfied and the number of primes ramified in F/K where condition (iv) is satisfied. We then have the following:

Proposition 3.3. The order of 2 in the Tamagawa ratio $\mathcal{T}(E^{\chi}/E^{\prime\chi})$ is given by

$$
\text{ord}_2 \mathcal{T}(E^{\chi}/E'^{\chi}) = g(\chi) + \sum_{v \mid 2\Delta \infty} \left(\text{dim}_{\mathbb{F}_2} H^1_{\phi}(K_v, C^{\chi}) - 1\right).
$$

Proof. By Thereom [2.2,](#page-3-2) $\text{ord}_2 \mathcal{T}(E^{\chi}/E^{\prime \chi})$ is given by

$$
\text{ord}_2 \mathcal{T}(E^{\chi}/E'^{\chi}) = \sum_{v|2\Delta\Delta_{F/K}\infty} \left(\text{dim}_{\mathbb{F}_2} H^1_{\phi}(K_v, C^d) - 1\right),
$$

where $\Delta_{F/K}$ is the relative discriminant of the extension F/K cut out by χ . By Remark [3.2,](#page-4-2) Lemma [3.1](#page-4-1) gives us that

$$
\dim_{\mathbb{F}_2} H_{\phi}^1(K_{\mathfrak{p}}, C^{\chi}) - 1 = \frac{\left(\frac{\Delta'}{\mathfrak{p}}\right) - \left(\frac{\Delta}{\mathfrak{p}}\right)}{2}
$$

for places $\mathfrak{p} \mid \Delta_{F/K}$ with $\mathfrak{p} \nmid 2\Delta\infty$ and the result follows.

4. The Erdős-Kac Theorem For Quadratic Characters

Because the sum

$$
\sum_{v|2\Delta\infty} \left(\dim_{\mathbb{F}_2} H^1_{\phi}(K_v, C^{\chi}) - 1\right)
$$

can be bounded uniformly for a fixed elliptic curve, Proposition [3.3](#page-4-3) suggests that we should study the distribution of the additive function $g(\chi)$ in order to understand the distribution of $\mathcal{T}(E^{\chi}/E'^{\chi})$ as χ varies.

When $f : \mathbb{N} \to \mathbb{C}$ is an additive function on the integers, then under mild hypotheses, the classical Erdős-Kac Theorem tells us that the distribution of f on natural numbers less than X approaches a normal distribution with mean $\mu(X)$ and variance $\sigma^2(X)$ as $X \to \infty$, where $\mu(X)$ and $\sigma(X)$ are the rational analogues of $\mu_f(X)$ and $\sigma_f(X)$ defined in the introduction. Theorem [1.3](#page-1-1) is the statement that the same type of result holds for additive functions on quadratic characters of a number field.

We begin by observing that if $\chi_{d_1} = \chi_{d_2}$, then, if $(d_1) = \mathfrak{a} \mathfrak{b}_1^2$ and $(d_2) = \mathfrak{a} \mathfrak{b}_2^2$, \mathfrak{b}_1 and \mathfrak{b}_2 lie in the same ideal class of \mathcal{O}_K . Conversely, if \mathfrak{b}_1 and \mathfrak{b}_2 lie in the same class, then $\chi_{d_1} = \chi_{\varepsilon d_2}$ for some $\varepsilon \in \mathcal{O}_K^{\times}$. Thus, we see that elements of $C(K, X)$ correspond to triples $(\mathfrak{b}, \mathfrak{a}, \varepsilon)$ with \mathfrak{b} a representative of its ideal class of minimal norm, \mathfrak{a} a squarefree ideal of norm $\mathbf{N}\mathfrak{a} < X/\mathbf{N}\mathfrak{b}^2$ such that $\mathfrak{a}\mathfrak{b}^2$ is principal, and ε an element of $\mathcal{O}_K^{\times}/(\mathcal{O}_K^{\times})^2$. From this discussion, it is apparent, for a prime ideal $\mathfrak p$ and a fixed choice $\mathfrak b_0$ of $\mathfrak b$, that

$$
\lim_{X \to \infty} \frac{|\{\chi \in C(K, X) \leftrightarrow (\mathfrak{b}_0, \mathfrak{a}, \varepsilon) : \mathfrak{p} | \mathfrak{a} \}|}{|C(K, X)|} = \Pr(\mathfrak{p} | \mathfrak{a} : \mathfrak{a} \text{ is squarefree}, \mathfrak{a} \mathfrak{b}_0^2 \text{ is principal})
$$
\n
$$
= \frac{1}{\mathbf{N}\mathfrak{p} + 1},
$$

whence, ranging over all χ , the probablity that $\mathfrak{p} \mid \mathfrak{a}$ is $\frac{1}{N\mathfrak{p}+1}$.

Thus, treating the events $\mathfrak{p}_1 \mid \mathfrak{a}$ and $\mathfrak{p}_2 \mid \mathfrak{a}$ as independent, we might predict that

$$
\frac{1}{|C(K,X)|} \sum_{\chi \in C(K,X)} f(\chi) \approx \tilde{\mu}_f(X),
$$

where

$$
\tilde{\mu}_f(X) := \sum_{\mathbf{N}\mathfrak{p} < X} \frac{f(\mathfrak{p})}{\mathbf{N}\mathfrak{p} + 1} = \mu_f(X) + O(1).
$$

In fact, this prediction is true, which we establish using the method of moments, following the blueprint of Granville and Soundararajan [\[GS07\]](#page-9-6). This requires the following technical result about the function $g_{\mathfrak{p}}(\chi)$, defined to be

$$
g_{\mathfrak{p}}(\chi) := \begin{cases} f(\mathfrak{p}) \left(1 - \frac{1}{N\mathfrak{p}+1}\right) & \text{if } \mathfrak{p} \mid \mathfrak{a}, \text{ and} \\ f(\mathfrak{p}) \left(-\frac{1}{N\mathfrak{p}+1}\right) & \text{if } \mathfrak{p} \nmid \mathfrak{a}. \end{cases}
$$

Theorem 4.1. With notation as above, uniformly for $k \leq \sigma_f(z)^{2/3}$, we have that

$$
\frac{1}{|C(K,X)|}\sum_{\chi \in C(K,X)} \left(\sum_{\mathbf{N}\mathfrak{p} < z} g_{\mathfrak{p}}(\chi)\right)^k = c_k \sigma_f(z)^k \left(1 + O\left(\frac{k^3}{\sigma_f(z)^2}\right)\right) + O\left(X^{\lambda-1}3^k \pi_K(z)^k\right)
$$

if k is even, and

$$
\frac{1}{|C(K,X)|}\sum_{\chi \in C(K,X)} \left(\sum_{\mathbf{N}\mathfrak{p} < z} g_{\mathfrak{p}}(\chi)\right)^k \ll c_k \sigma_f(z)^{k-1} k^{3/2} + X^{\lambda-1} 3^k \pi_K(z)^k
$$

if k is odd. Here, $c_k = \Gamma(k+1)/2^{k/2}\Gamma(\frac{k}{2}+1)$ and $\lambda < 1$ depends only on the degree of K.

The proof of Theorem [4.1](#page-5-0) relies upon a result about the distribution of squarefree ideals. To this end, given ideals c and q and squarefree $\mathfrak{d} \mid \mathfrak{q}$, define $N^{\text{sf}}(X; \mathfrak{c}, \mathfrak{q}, \mathfrak{d})$ to be the number of squarefree ideals $\mathfrak a$ of norm up to X in the same class as c and such that $(\mathfrak a, \mathfrak q) = \mathfrak d$. We then have:

Lemma 4.2. With notation as above, we have that

$$
N^{\mathrm{sf}}(X;\mathfrak{c},\mathfrak{q},\mathfrak{d})=\frac{1}{|\mathrm{Cl}(K)|}\frac{\mathrm{res}_{s=1}\zeta_K(s)}{\zeta_K(2)}\phi(\mathfrak{q},\mathfrak{d})X+O(X^{\lambda}3^{\omega(\mathfrak{q})}),
$$

where $\lambda = \frac{\deg K - 1}{\deg K + 1}$ if $\deg K \geq 3$ and $\lambda = 1/2$ otherwise, $\omega(\mathfrak{q})$ denotes the number of distinct primes dividing q, and

$$
\phi(\mathfrak{q},\mathfrak{d})=\prod_{\mathfrak{p}\mid \mathfrak{d}}\frac{1}{\mathbf{N}\mathfrak{p}+1}\prod_{\mathfrak{p}\mid \mathfrak{q},\mathfrak{p}\nmid \mathfrak{d}}\frac{\mathbf{N}\mathfrak{p}}{\mathbf{N}\mathfrak{p}+1}.
$$

Proof. This follows from elementary considerations and the classical estimate

$$
\sum_{\substack{\mathbf{N}\mathfrak{a}\leq X \\ \mathfrak{a}\mathfrak{c}^{-1} \text{ prim.}}} 1 = \frac{1}{|\mathrm{Cl}(K)|} \mathrm{res}_{s=1} \zeta_K(s) \cdot X + O(X^{\frac{\deg K - 1}{\deg K + 1}}).
$$

Proof of Theorem [4.1.](#page-5-0) For any ideal **q**, define $g_{\mathfrak{q}}(\chi) := \prod_{\mathfrak{p}^{\alpha}||\mathfrak{q}} g_{\mathfrak{p}}(\chi)^{\alpha}$. We then have that

$$
\sum_{\chi \in C(K,X)} \left(\sum_{\mathbf{N}\mathfrak{p} < z} g_{\mathfrak{p}}(\chi) \right)^k = \left| \mathcal{O}_K^{\times}/(\mathcal{O}_K^{\times})^2 \right| \cdot \sum_{\mathfrak{b}} \sum_{\substack{\mathbf{N}\mathfrak{a} < \frac{X}{N\mathfrak{b}^2} \\ \mathfrak{a}\mathfrak{b}^2 \text{ prin.}}} \left(\sum_{\substack{\mathbf{N}\mathfrak{p} < z \\ \mathfrak{a}\mathfrak{b}^2 \text{ prin.}}} g_{\mathfrak{p}}(\mathfrak{a}) \right)^k
$$

$$
= \left| \mathcal{O}_K^{\times}/(\mathcal{O}_K^{\times})^2 \right| \cdot \sum_{\mathfrak{b}} \sum_{\substack{\mathbf{N}\mathfrak{p}_1, \dots, \mathbf{N}\mathfrak{p}_k < z \\ \mathfrak{a}\mathfrak{b}^2 \text{ principal}}} \sum_{\substack{\mathbf{N}\mathfrak{a} < \frac{X}{N\mathfrak{b}^2} \\ \mathfrak{a}\mathfrak{b}^2 \text{ principal}}} g_{\mathfrak{p}_1 \dots \mathfrak{p}_k}(\mathfrak{a}),
$$

where, as expected, the summation over $\mathfrak b$ is taken to be over representatives of minimal norm for each ideal class and the prime on the summation over a indicates it is to be taken over squarefree ideals. Given an ideal \mathfrak{c} , we now consider for any \mathfrak{q} the more general summation

$$
\sum_{\mathfrak{a}\in\mathcal{Y}\atop \mathfrak{a}\in\mathcal{I}\atop \text{principal}} g_{\mathfrak{q}}(\mathfrak{a}) = \sum_{\mathfrak{d}\mid\sqrt{\mathfrak{q}}} g_{\mathfrak{q}}(\mathfrak{d}) N^{\mathrm{sf}}(Y; \mathfrak{c}, \mathfrak{q}, \mathfrak{d})
$$
\n
$$
= \frac{Y}{|\mathrm{Cl}(K)|} \frac{\mathrm{res}_{s=1}\zeta_K(s)}{\zeta_K(2)} \sum_{\mathfrak{d}\mid\sqrt{\mathfrak{q}}} g_{\mathfrak{q}}(\mathfrak{d}) \phi(\mathfrak{q}, \mathfrak{d}) + O\left(Y^{\lambda}3^{\omega(\mathfrak{q})}\sum_{\mathfrak{d}\mid\sqrt{\mathfrak{q}}} |g_{\mathfrak{q}}(\mathfrak{d})|\right),
$$
\n
$$
=: \frac{Y}{|\mathrm{Cl}(K)|} \frac{\mathrm{res}_{s=1}\zeta_K(s)}{\zeta_K(2)} G(\mathfrak{q}) + O\left(Y^{\lambda}3^{\omega(\mathfrak{q})}\right),
$$

say, where $\sqrt{\mathfrak{q}} = \prod_{\mathfrak{p}|\mathfrak{q}} \mathfrak{p}$. We note that $G(\mathfrak{q})$ is multiplicative, and is given by

$$
G(\mathfrak{q}) = \prod_{\mathfrak{p}^{\alpha}||\mathfrak{q}} \frac{f(\mathfrak{p})^{\alpha}}{N\mathfrak{p}+1} \left(\left(1 - \frac{1}{N\mathfrak{p}+1}\right)^{\alpha} + N\mathfrak{p} \cdot \left(-\frac{1}{N\mathfrak{p}+1}\right)^{\alpha} \right).
$$

 \Box

Thus, $G(q) = 0$ unless each α is at least two, i.e. q is square-full.

Returning to the original problem, we find that

$$
\sum_{\chi \in C(K,X)} \left(\sum_{\mathbf{N}\mathfrak{p} < z} g_{\mathfrak{p}}(\chi) \right)^k = c(K) \cdot X \sum_{\mathbf{N}\mathfrak{p}_1,\dots,\mathbf{N}\mathfrak{p}_k < z} G(\mathfrak{p}_1 \dots \mathfrak{p}_k) + O\left(X^{\lambda} 3^k \pi_K(z)^k\right),
$$

where $\pi_K(z) := \#\{\mathfrak{p} : \mathbb{N}\mathfrak{p} < z\}$ and

$$
c(K) := \left| \mathcal{O}_K^{\times}/(\mathcal{O}_K^{\times})^2 \right| \frac{1}{|\mathrm{Cl}(K)|} \frac{\mathrm{res}_{s=1} \zeta_K(s)}{\zeta_K(2)} \sum_{\mathfrak{b}} \frac{1}{\mathbf{N} \mathfrak{b}^2}.
$$

Noting that the above discussion also proves that $|C(K, X)| = c(K) \cdot X + O(X^{\lambda})$, the goal is to estimate the summation over $\mathfrak{p}_1, \ldots, \mathfrak{p}_k$. Since $G(\mathfrak{q}) = 0$ unless \mathfrak{q} is square-full, we have that

$$
\sum_{\mathbf{N}\mathfrak{p}_1,\dots,\mathbf{N}\mathfrak{p}_k < z} G(\mathfrak{p}_1 \dots \mathfrak{p}_k) = \sum_{s \leq k/2} \sum_{\substack{\alpha_1 + \dots + \alpha_s = k, \\ \text{each } \alpha_i \geq 2}} \frac{k!}{\alpha_1! \dots \alpha_s!} \sum_{\substack{\mathfrak{p}_1 < \dots < \mathfrak{p}_s \\ \mathbf{N}\mathfrak{p}_s < z}} G(\mathfrak{p}_1^{\alpha_1} \dots \mathfrak{p}_s^{\alpha_s}),
$$

where $\mathfrak{p} < \mathfrak{p}'$ is determined by a norm-compatible linear ordering on the prime ideals of \mathcal{O}_K . We note that, since $G(\mathfrak{p}^{\alpha}) \leq \frac{f(\mathfrak{p})^2}{N\mathfrak{p}}$ $\frac{(\mathfrak{p})^2}{N\mathfrak{p}}$, the inner summation contributes no more than $O(\sigma_f(z)^{2s})$, which will be an error term unless $s=\frac{k}{2}$ $\frac{k}{2}$; the dependence of this error on k can be sussed out exactly as in Granville and Soundararajan's work. In fact, the handling of the main term, arising when k is even and $s = \frac{k}{2}$ $\frac{k}{2}$, is also essentially the same. In particular, the inner summation is equal to

$$
\frac{1}{(\frac{k}{2})!} \sum_{\substack{\mathbf{N}\mathfrak{p}_1,\dots,\mathbf{N}\mathfrak{p}_{\frac{k}{2}} < z}} G(\mathfrak{p}_1^2 \dots \mathfrak{p}_{\frac{k}{2}}^2) = \frac{1}{(\frac{k}{2})!} \left(\sum_{\substack{\mathbf{N}\mathfrak{p} < z}} G(\mathfrak{p}^2) + O(\log \log k) \right)^{k/2}
$$

$$
= \frac{1}{(\frac{k}{2})!} \left(\sigma_f(z)^2 + O(\log \log k) \right)^{k/2}.
$$

This yields Theorem [4.1.](#page-5-0)

We are now ready to prove Theorem [1.3.](#page-1-1)

Proof of Theorem [1.3.](#page-1-1) Recall that we wish to show that the quantity

$$
\frac{f(\chi) - \mu_f(X)}{\sigma_f(X)}, \ \chi \in C(K, X),
$$

is normally distributed as $X \to \infty$. As remarked above, we will do so using the method of moments. In particular, we have that

$$
\frac{1}{|C(K,X)|}\sum_{\chi \in C(K,X)}(f(\chi) - \mu_f(X))^k = \frac{1}{|C(K,X)|}\sum_{\chi \in C(K,X)}\left(\sum_{\mathfrak{p}|\mathfrak{a}}f(\mathfrak{p}) - \sum_{\mathbf{N}\mathfrak{p} < X}\frac{f(\mathfrak{p})}{\mathbf{N}\mathfrak{p} + 1} + O(1)\right)^k
$$
\n
$$
= \frac{1}{|C(K,X)|}\sum_{\chi \in C(K,X)}\left(\sum_{\mathbf{N}\mathfrak{p} < z}g_{\mathfrak{p}}(\mathfrak{a}) + O\left(\frac{\log X}{\log z}\right)\right)^k.
$$

Considering the error term in Theorem [4.1,](#page-5-0) we take $z = X^{\frac{1-\lambda}{k}}$. With this choice, the inner summation becomes

$$
\left(\sum_{\mathbf{N}\mathfrak{p}
$$

hence Theorem [4.1,](#page-5-0) the Cauchy-Schwarz inequality, and the fact that $\sigma_f(z) = \sigma_f(X) +$ $O(\log k)$ yield that

$$
\frac{1}{|C(K,X)|} \sum_{\chi \in C(K,X)} (f(\chi) - \mu_f(X))^k = c_k \sigma_f(X)^k \left(1 + O\left(\frac{k^{3/2}}{\sigma_f(X)}\right)\right)
$$

if k is even, and

$$
\frac{1}{|C(K,X)|} \sum_{\chi \in C(K,X)} (f(\chi) - \mu_f(X))^k \ll c_k \sigma_f(X)^{k-1} k^{3/2}
$$

if k is odd. This proves the theorem.

5. Proof of Main Theorems

In order to apply Theorem [1.3](#page-1-1) to the additive function $g(\chi)$ defined in [\(3\)](#page-4-4), we need to evaluate the quantities $\mu_g(X)$ and $\sigma_g(X)$. We begin with the following

Proposition 5.1. Let $c \in K^{\times}$ be non-square. Then

$$
\sum_{\mathbf{N}\mathfrak{p}\leq X} \frac{1+\left(\frac{c}{\mathfrak{p}}\right)}{\mathbf{N}\mathfrak{p}} = \log\log X + O(1).
$$

Proof. This is a consequence of the prime ideal theorem and the fact that the Hecke Lfunction attached to the non-trivial character of $Gal(K(\sqrt{c})/K)$ is analytic and non-vanishing on the line $\Re(s) = 1$.

We now decompose $\mu_g(X)$ as

$$
\mu_g(X) = \frac{1}{2} \sum_{\substack{\mathbf{N}\mathfrak{p} < X \\ p \nmid 2\Delta\infty}} \frac{1 + \left(\frac{\Delta'}{\mathfrak{p}}\right)}{\mathbf{N}\mathfrak{p}} - \frac{1}{2} \sum_{\substack{\mathbf{N}\mathfrak{p} < X \\ \mathfrak{p} \nmid 2\Delta\infty}} \frac{1 + \left(\frac{\Delta}{\mathfrak{p}}\right)}{\mathbf{N}\mathfrak{p}},
$$

and it immediately follows from Proposition [5.1](#page-8-1) that $\mu_g(X) = O(1)$. We also rewrite $\sigma_g(X)$ as

$$
\sigma_g(X) = \left(\sum_{\substack{\mathbf{N}\mathfrak{p}\leq X, \mathfrak{p}\nmid 2\Delta\infty \\ \left(\frac{\Delta}{\mathfrak{p}}\right)\neq \left(\frac{\Delta'}{\mathfrak{p}}\right)}}\frac{1}{\mathbf{N}\mathfrak{p}}\right)^{1/2} = \left(\frac{1}{2}\sum_{\substack{\mathbf{N}\mathfrak{p}\leq X \\ \mathfrak{p}\nmid 2\Delta\infty}}\frac{1-\left(\frac{\Delta\Delta'}{\mathfrak{p}}\right)}{\mathbf{N}\mathfrak{p}}\right)^{1/2}
$$

.

In order to apply Proposition [5.1](#page-8-1) to $\sigma_g(X)$, we therefore need $\Delta\Delta'$ to be non-square in K. This will be the case when E does not have a cyclic 4-isogeny defined over $K(E[2])$.

Proposition 5.2. If E is an elliptic curve with $E(K)[2] \simeq \mathbb{Z}/2\mathbb{Z}$ that does not have a cyclic 4 -isogeny defined over $K(E[2])$, then $\Delta\Delta' \notin (K^{\times})^2$

Proof. Let $Q' \in E'[2] - C'$, $C = \langle P \rangle$, and take $Q \in E[4]$ with $\phi(Q) = Q'$. Since $Q' \in E$ $E'(K)[2] - C'$, and both $\phi \circ \hat{\phi} = [2]_{E'}$ and $\hat{\phi} \circ \phi = [2]_E$, it follows that $2Q = \hat{\phi}(Q') = P$. Let $M = K(E[2])$. Since E has no cyclic 4-isogeny defined over M, there exists $\sigma \in G_M$ such that $\sigma(Q) \notin \langle Q \rangle = \{0, Q, P, Q + P\}$. In particular, since $\phi^{-1}(Q') \subset \langle Q \rangle$, we get that $\phi(\sigma(Q)) \neq Q'$. We then get that

$$
\sigma(Q') = \sigma(\phi(Q)) = \phi(\sigma(Q)) \neq Q',
$$

showing that Q' is not defined over M, and therefore that $K(E'[2]) \not\subset M$. It then follows that $K(E[2])$ and $K(E'[2])$ are disjoint quadratic extensions of K and that $E'(K)[2] \simeq \mathbb{Z}/2\mathbb{Z}$. As $K(E[2])$ and $K(E'[2])$ are given by $K(\sqrt{\Delta})$ and $K(\sqrt{\Delta'})$ respectively, it follows that $\Delta\Delta' \notin (K^\times)^2$.

By Proposition [5.1,](#page-8-1) we therefore get that $\sigma_g(X) = \sqrt{\frac{1}{2}}$ $\frac{1}{2}$ log log $X + O(1)$ whenever E does not have a cyclic 4-isogeny defined over $K(E[2])$.

Proof of Theorem [1.2.](#page-1-0) Applying Theorem [1.3](#page-1-1) to $q(\chi)$, we get that

(4)
$$
\lim_{X \to \infty} \frac{\left| \left\{ \chi \in C(K, X) : g(\chi) - O(1) \le z \left(\sqrt{\frac{1}{2} \log \log X} + O(1) \right) \right\} \right|}{|C(K, X)|} = G(z)
$$

for every $z \in \mathbb{R}$. By Proposition [3.3,](#page-4-3) there is some contant C, independent of χ , such that $|g(\chi) - \text{ord}_2 \mathcal{T}(E^{\chi}/E^{\prime \chi})| < C$, so in fact [\(4\)](#page-9-7) holds with $g(\chi)$ replaced by $\text{ord}_2 \mathcal{T}(E^{\chi}/E^{\prime \chi})$ and the result follows. \Box

Proof of Theorem [1.1.](#page-0-0) By Theorem [1.2,](#page-1-0)

$$
\lim_{X \to \infty} \frac{|\{\chi \in C(K, X) : \text{ord}_2 \mathcal{T}(E^{\chi}/E'^{\chi}) \ge r\}|}{|C(K, X)|} = \frac{1}{2}
$$

for any fixed $r \ge 0$. As $d_2(E^{\chi}/K) \ge \text{ord}_2 \mathcal{T}(E^{\chi}/E'^{\chi}) - 2$, this shows that for any $\epsilon > 0$,

$$
\frac{|\{\chi \in C(K, X) : d_2(E^{\chi}/K) \ge r\}|}{|C(K, X)|} \ge \frac{1}{2} - \epsilon
$$

for sufficiently large X.

REFERENCES

- [Cas65] J.W.S. Cassels. Arithmetic on curves of genus 1. VIII: On the conjectures of Birch and Swinnerton-Dyer. Journal für die reine und angewandte Mathematik (Crelles Journal), 1965(217):180–199, 1965.
- [FG08] E.V. Flynn and C. Grattoni. Descent via isogeny on elliptic curves with large rational torsion subgroups. Journal of Symbolic Computation, 43(4):293–303, 2008.
- [GS07] A. Granville and K. Soundararajan. Sieving and the Erdős–Kac theorem. Equidistribution in number theory, an introduction, pages 15–27, 2007.
- [HB94] D.R. Heath-Brown. The size of Selmer groups for the congruent number problem, II. Inventiones mathematicae, 118(1):331–370, 1994.
- [Kan10] D.M. Kane. On the ranks of the 2-Selmer groups of twists of a given elliptic curve. Preprint available at http://arxiv. org/abs/1009.1365, 2010.
- [Kla11] Z. Klagsbrun. Selmer ranks of quadratic twists of elliptic curves with partial rational two-torsion. Preprint available at http://arxiv.org/abs/1201.5408, 2011.
- [KMR13] Z. Klagsbrun, B Mazur, and K. Rubin. A Markov model for Selmer ranks in families of twists. Preprint available at http://arxiv.org/pdf/1303.6507v1.pdf, 2013.

- [SD08] P. Swinnerton-Dyer. The effect of twisting on the 2-Selmer group. In Mathematical Proceedings of the Cambridge Philosophical Society, volume 145, pages 513–526. Cambridge Univ Press, 2008.
- [Xio13] Maosheng Xiong. On Selmer groups of quadratic twists of elliptic curve a two-torsion over q. Mathematika, 2013.

E-mail address: zev.klagsbrun@gmail.com

E-mail address: rjlo@stanford.edu