CONSTRUCTIONS OF ROUND FOLD MAPS ON CIRCLE BUNDLES

NAOKI KITAZAWA

ABSTRACT. (Stable) fold maps are fundamental tools in a generalization of the theory of Morse functions on smooth manifolds and its application to studies of algebraic and differential topological properties of smooth manifolds. Round fold maps were introduced as stable fold maps with singular value sets, or the set consisting of all the singular values, of concentric spheres by the author in 2013; for example, some special generic maps on spheres are regarded as round fold maps whose singular value sets are connected.

To construct explicit fold maps has been a fundamental and difficult problem. In 2014, the author succeeded in constructing explicit round fold maps into a Euclidean space of dimension larger than 1 on bundles over the standard sphere whose dimension is equal to that of the Euclidean space and connected sums of bundles over the standard sphere with fibers diffeomorphic to standard spheres by constructing maps locally and gluing them together properly. Later, the author constructed such maps on bundles over spheres or more general manifolds including families of bundles whose fibers are circles over given manifolds by applying operations compatiable with the structures of bundles (*P-operations*).

In this paper, we obtain new round fold maps on closed smooth manifolds having the structures of bundles whose fibers are circles over given closed manifolds by applying P-operations under weaker conditions. We also study algebraic and differential topological properties of the resulting maps and the source manifolds.

1. Introduction and terminologies

Fold maps are important in generalizing the theory of Morse functions and its application to studies of geometry of manifolds. Studies of such maps were started by Whitney ([24]) and Thom ([23]) in the 1950's. A fold map is a smooth map whose singular points are of the form

$$(x_1, \dots, x_m) \mapsto (x_1, \dots, x_{n-1}, \sum_{k=n}^{m-i} x_k^2 - \sum_{k=m-i+1}^m x_k^2)$$

for two positive integers $m \ge n$ and an integer $0 \le i \le m-n+1$ (for each singular point, we can determine the integer i uniquely and the integer is said to be the *index* of the point). A Morse function is regarded as a fold map (n=1). For a fold map from a closed smooth manifold of dimension m into a smooth manifold of dimension $n \ge 1$ (without boundary), the followings hold.

 $^{2010\ \}textit{Mathematics Subject Classification}.\ \text{Primary } 57\text{R}45.\ \text{Secondary } 57\text{N}15.$

Key words and phrases. Singularities of differentiable maps; singular sets, fold maps. Differential topology.

- (1) The *singular set*, which is defined as the set of all the singular points, and the set of all the singular points whose indices are i are closed smooth submanifolds of dimension n-1 of the source manifold.
- (2) The restriction map to the singular set is a smooth immersion of codimension 1.

We also note that if the restriction map to the singular set is an immersion with normal crossings, then it is stable. Stable maps are important in the theory of global singularity; see [8] for example).

Since around the 1990s, some classes of fold maps easy to handle have been actively studied. For example, in [2], [7], [19], [20] and [22], special generic maps were studied. A special generic map is defined as a fold map such that the indices of singular points are always 0. From Morse functions and their singular points (singular sets), we can know some topological properties such as Euler numbers and homology groups and from special generic maps and their singular points (singular sets), we can often know more precise information of their source manifolds such as the homotopy types and the homeomorphism and diffeomorphism types. For example, homotopy spheres are characterized as manifolds admitting Morse functions with two singular points except the case where the sphere is 4-dimensional and special generic maps often restrict the diffemorphism types of homotopy spheres of source manifolds (Standard spheres always admit special generic maps into Euclidean spaces whose dimensions are not higher than those of the source manifolds). As another example, the diffeomorphism types of manifolds admitting special generic maps were determined under appropriate conditions. For such facts, see [19] and [20] for example. In [22], Sakuma studied simple fold maps. A simple fold map is defined as a fold map such that the inverse image of each singular value does not have any connected component with more than one singular points (see also [18]) and special generic maps are simple, for example. In [15], Kobayashi and Saeki investigated topology of stable maps including fold maps which are stable into the plane. In [21], Saeki and Suzuoka found good properties of manifolds admitting stable maps such that the inverse images of regular values are always disjoint unions of spheres. Later, in [11], round fold maps, which will be mainly studied in this paper, were introduced. A round fold map is defined as a fold map satisfying the followings.

- (1) The singular set is a disjoint union of standard spheres.
- (2) The restriction map to the singular set is an embedding.
- (3) The *singular value set*, which is defined as the set of all the singular values of the map, is a disjoint union of spheres embedded concentrically.

The m-dimensional standard sphere S^m ($m \ge 2$) admits a round fold map whose singular set is connected into \mathbb{R}^n for $2 \le n \le m$. Such a map is also a special generic map and some special generic maps on homotopy spheres are round fold maps whose singular sets are connected as discussed in [19].

In [11], homology groups and homotopy groups of manifolds admitting round fold maps are studied. Some examples of round fold maps and the diffeomorphism types of their source manifolds are given by the author in [10], [11], [13] and [14]. For example, in the first two papers, we have obtained round fold maps on m-dimensional closed smooth manifolds having the structures of bundles over the n-dimensional $(n \geq 2)$ standard sphere S^n whose fibers are closed smooth manifolds and whose structure groups consist of diffeomorphisms. It has been a very difficult problem to construct explicit fold maps, although the existence problems for fold maps have

been solved under a lot of situations (see [5] and [6] for example). The author has given a lot of answers for the construction problems, which will make it easier to study topological properties of a lot of smooth manifolds by using fold maps being not so complex.

Here, we introduce terminologies on (fiber) bundles, which are important in this paper. In this paper, a bundle means a bundle whose fibers are diffeomorphic closed smooth manifolds and whose structure group consists of diffeomorphisms on the fibers. We call a bundle whose fiber is a closed smooth manifold X an X-bundle. We return to precise presentations about explicit round fold maps. In [11] and [13], the author constructed round fold maps into \mathbb{R}^n $(n \geq 2)$ also on m-dimensional (m > n) closed smooth manifolds represented as connected sums of bundles over the n-dimensional standard sphere S^n with fibers diffemorphic to the (m-n)dimensional standard sphere S^{m-n} under the assumption that m > 2n holds. In [14], as new answers, we construct new round fold maps on closed smooth manifolds having the structures of smooth bundles over (exotic) spheres or more general manifolds. More precisely, we considered round fold maps having good algebraic and differential topological properties and bundles over the source manifolds of the maps with good topological properties and constructed new round fold maps on the bundles by using operations compatiable with the structures of bundles (Poperations). For example, we constructed round fold maps on manifolds having the structures of S^1 -bundles (circle bundles) over some smooth manifolds and obtain ones into \mathbb{R}^n $(n \geq 2)$ on S^1 -bundles over manifolds represented as connected sums of smooth bundles over S^n with fibers diffemorphic to S^{m-n} under the assumption that $m \ge 2n$ and m - n > 2 hold.

In this paper, we extend results of [14] and give some applications. More precisely, we construct new examples of round fold maps on smooth manifolds having the structures of S^1 -bundles over some smooth manifolds under weaker conditions by using methods similar to ones used in [14]. In this paper, we also study algebraic and differential topological properties of obtained maps and their source manifolds precisely.

This paper is organized as the following.

In section 2, we recall round fold maps and some terminologies on round fold maps such as axes and proper cores. We also introduce classes of round fold maps such as C^{∞} L-trivial round fold maps and C^{∞} S-trivial ones. A C^{∞} L-trivial round fold map is defined as a round fold map satisfying a kind of triviality on the topological type around each connected component of the singular value set and a C^{∞} S-trivial round fold map is defined as one satisfying a kind of triviality on the topological type around the singular value set. Such classes were first introduced in [11] and [12] by the author and the original definitions are a bit different from those of the present paper. We introduce examples of round fold maps in Example 1.

In section 3, we recall P-operations defined in [14], which are operations used to construct new round fold maps on manifolds having the structures of bundles over smooth manifolds admitting C^{∞} L-trivial round fold maps. We apply these operations to construct new round fold maps on S^1 -bundles over manifolds admitting C^{∞} L-trivial round fold maps and prove Theorems 1, 2, 3 and 4, which are regarded as extensions of results in [14], which are introduced as Propositions 3 and 4 in the present paper.

In section 4, as applications, we construct new round fold maps by applying Propositions and Theorems on P-operations obtained in section 3 to round maps introduced in Example 1. Furthermore, we study the local and global structures of the obtained maps. For example, we study problems whether these maps being L-trivial are S-trivial or not. We also study the diffeomorphism types of the soruce manifolds of obtained maps.

In this paper, for a smooth map c, the fiber of a point in the target manifold means the inverse image of the point by the map c. A regular fiber means a fiber of a regular value of a smooth map. We define the singular set of a smooth map c as the set consisting of all the singular points of c and denote the set by S(c). For the map c, the set c(S(c)) is said to be the singular value set of c and the set $\mathbb{R}^n - c(S(c))$ is said to be the regular value set of c.

Throughout this paper, all the manifolds and maps between them are assumed to be smooth and of class C^{∞} unless otherwise stated. Last, in this paper, We assume that M is a closed manifold of dimension m, that N is a manifold of dimension nwithout boundary and that $m \ge n \ge 1$ holds.

2. Round fold maps

In this section, we review round fold maps. See also [11] and [12] for example.

Definition 1 (round fold maps ([12])). $f: M \to \mathbb{R}^n$ $(m \ge n \ge 2)$ is said to be a round fold map if f is C^{∞} equivalent to a fold map $f_0: M \to \mathbb{R}^n$ such that the followings hold.

- (1) The singular set $S(f_0)$ is a disjoint union of (n-1)-dimensional standard spheres and consists of $l \in \mathbb{N}$ connected components.
- (2) The restriction map $f_0|_{S(f_0)}$ is an embedding. (3) Let $D^n_r:=\{(x_1,\cdots,x_n)\in\mathbb{R}^n\mid\sum_{k=1}^nx_k^2\leq r\}$. Then $f_0(S(f_0))=$ $\sqcup_{k=1}^{l} \partial D^{n}_{k}$ holds.

We call f_0 a normal form of f. We call a ray L from $0 \in \mathbb{R}^n$ an axis of f_0 and $D^{n}_{\frac{1}{2}}$ the proper core of f_{0} . Suppose that for a round fold map f, its normal form f_0 and diffeomorphisms $\Phi: M \to M$ and $\phi: \mathbb{R}^n \to \mathbb{R}^n$, the relation $\phi \circ f = f_0 \circ \Phi$ holds. Then for an axis L of f_0 , we also call $\phi^{-1}(L)$ an axis of f and for the proper core $D^n_{\frac{1}{2}}$ of f_0 , we also call $\phi^{-1}(D^n_{\frac{1}{2}})$ a proper core of f.

For a round fold map $f: M \to \mathbb{R}^n$ and a connected component of the singular value set C, there exists a closed tubular neighborhood N(C), which is regarded as a trivial [-1,1]-bundle over C (C corresponds to the image $C \times \{0\} \subset C \times [-1,1]$ of the section of the trivial bundle N(C) regarded as $C \times [-1,1]$, such that the composition of the map $f|_{f^{-1}(N(C))}: f^{-1}(N(C)) \to N(C)$ and the projection to each connected component of the boundary $\partial N(C)$ gives $f^{-1}(N(C))$ the structure of a bundle. In the case where the inverse image of a connected component C' of the boundary $\partial N(C)$ is non-empty, the inverse image is a disjoint union of connected components of the boundary $\partial f^{-1}(N(C))$ and if we restrict the map giving $f^{-1}(N(C))$ the structure of a bundle over the connected component C' to the set $f^{-1}(C')$, then it coincides with the map f on $f^{-1}(C')$.

For example, if a connected component C is the image of a connected component of the singular set consisting of singular points of index 0, then the composition of the map $f|_{f^{-1}(N(C))}: f^{-1}(N(C)) \to N(C)$ and the projection to each connected component of the boundary $\partial N(C)$ gives $f^{-1}(N(C))$ the structure of a bundle whose fiber is a standard closed disc and whose structure group is a subgroup of linear transformations on the disc. See also [19] for example.

Definition 2. Let $f: M \to \mathbb{R}^n$ be a round fold map. Assume that for any connected component C of f(S(f)) and a small closed tubular neighborhood N(C) of C such that $\partial N(C)$ is the disjoint union of two connected components C_a and C_b , $f^{-1}(N(C))$ has the structures of isomorphic trivial bundles whose fibers are diffeomorphic to a closed manifold F_C over C_a and C_b and $f|_{f^{-1}(C_a)}: f^{-1}(C_a) \to C_a$ and $f|_{f^{-1}(C_b)}: f^{-1}(C_b) \to C_b$ give the structures of subbundles of the bundles $f^{-1}(N(C))$ ($f^{-1}(C_a)$ or $f^{-1}(C_b)$ may be empty). Then, f is said to be C^{∞} L-trivial. We call a fiber F_C of the bundles $f^{-1}(N(C))$ a normal fiber of C corresponding to the bundles $f^{-1}(N(C))$.

Let f be a normal form of a round fold map and $P_1:=D^n_{\frac{1}{2}}$. We set $E:=f^{-1}(P_1)$ and $E':=M-f^{-1}(\operatorname{Int} P_1)$. We set $F:=f^{-1}(p)$ for $p\in\partial P_1$. We put $P_2:=\mathbb{R}^n-\operatorname{Int} P_1$. Let $f_1:=f|_E:E\to P_1$ if F is non-empty and let $f_2:=f|_{E'}:E'\to P_2$.

 f_1 gives the structure of a trivial bundle over P_1 and $f_1|_{\partial E}: \partial E \to \partial P_1$ gives the structure of a trivial bundle over ∂P_1 if F is non-empty. $f_2|_{\partial E'}: \partial E' \to \partial P_2$ gives the structure of a trivial bundle over ∂P_2 if F is non-empty.

We can give E' the structures of bundles over ∂P_2 as follows.

Since for $\pi_P(x) := \frac{1}{2} \frac{x}{|x|}$ $(x \in P_2)$, $\pi_P \circ f|_{E'}$ is a proper submersion, this map gives E' the structure of a C^{∞} $f^{-1}(L)$ -bundle over ∂P_2 (apply Ehresmann's fibration theorem [4]).

Considering these observations, we introduce the following classes. For the classes, see also [11], in which such classes were introduced first. These classes are a bit different from ones here.

Definition 3. Let $f: M \to \mathbb{R}^n$ be a round fold map.

For a proper core P and an axis L of f, We give the manifold $f^{-1}(\mathbb{R}^n - \text{Int}P)$ the structure of a bundle over ∂P whose fiber is diffeomorphic to $f^{-1}(L)$ such that $f|_{f^{-1}(\partial P)}: f^{-1}(\partial P) \to \partial P$ gives the structure of a subbundle of the previous bundle $f^{-1}(\mathbb{R}^n - \text{Int}P)$ ($f^{-1}(\partial P)$ may be empty). We call such a bundle a surrounding bundle of f.

If a surrounding bundle is a trivial bundle, then f is said to be C^{∞} S-trivial.

We easily obtain a round fold map which is C^{∞} L-trivial and C^{∞} S-trivial as shown in the following (see [11] and [12] for example).

Let \bar{M} be a compact manifold with non-empty boundary $\partial \bar{M}$. Let $a \in \mathbb{R}$. Then, there exists a Morse function $\tilde{f}: \bar{M} \to [a, +\infty)$ satisfying the followings.

- (1) a is the minimum of \tilde{f} and $\tilde{f}^{-1}(a) = \partial \bar{M}$ holds.
- (2) All the singular points of $\tilde{f}: \bar{M} \to [a, +\infty)$ are in $\bar{M} \partial \bar{M}$ and at distinct singular points, the values are always distinct.

Let $\Phi: \partial(\bar{M} \times \partial(\mathbb{R}^n - \operatorname{Int}D^n)) \to \partial(\partial\bar{M} \times D^n)$ and $\phi: \partial(\mathbb{R}^n - \operatorname{Int}D^n) \to \partial D^n$ be diffeomorphisms. Let $p_1: \partial\bar{M} \times \partial(\mathbb{R}^n - \operatorname{Int}D^n) \to \partial(\mathbb{R}^n - \operatorname{Int}D^n)$ and $p_2: \partial\bar{M} \times \partial D^n \to \partial D^n$ be the canonical projections. Suppose that the following diagram commutes.

$$\partial \bar{M} \times \partial (\mathbb{R}^n - \mathrm{Int} D^n) \xrightarrow{\Phi} \partial \bar{M} \times \partial D^n$$

$$\downarrow^{p_1} \qquad \qquad \downarrow^{p_2}$$

$$\partial (\mathbb{R}^n - \mathrm{Int} D^n) \xrightarrow{\phi} \partial D^n$$

By using the diffeomorphism Φ , we construct $M:=(\partial \bar{M}\times D^n)\bigcup_{\Phi}(\bar{M}\times\partial(\mathbb{R}^n-\mathrm{Int}D^n))$. Let $p:\partial \bar{M}\times D^n\to D^n$ be the canonical projection. Then gluing the two maps p and $\tilde{f}\times\mathrm{id}_{S^{n-1}}$ together by using the two diffeomorphisms Φ and ϕ , we obtain a round fold map $f:M\to\mathbb{R}^n$.

If \bar{M} is a compact manifold without boundary, then there exists a Morse function $\tilde{f}: \bar{M} \to [a, +\infty)$ such that $\tilde{f}(\bar{M}) \subset (a, +\infty)$ and that at distinct singular points, the values are always distinct. We are enough to consider $\tilde{f} \times \mathrm{id}_{S^{n-1}}$ and embed $[a, +\infty) \times S^{n-1}$ into \mathbb{R}^n to construct a round fold map whose source manifold is $\bar{M} \times S^{n-1}$.

We call this construction of a round fold map a trivial spinning construction.

We introduce some known examples of round fold maps with their source manifolds.

- **Example 1.** (1) Special generic maps with singular sets diffeomorphic to standard spheres into Euclidean spaces whose dimensions are larger than 1 such that the restriction maps to the singular sets are embeddings are round fold maps. Such maps are C^{∞} L-trivial and C^{∞} S-trivial. The source manifolds of such maps are always homotopy spheres. Let M be a closed m-dimensional manifold admitting such a map $f: M \to \mathbb{R}^n \ (m \ge n \ge 2)$. For a small closed tubular neighborhood N(f(S(f))) of the set f(S(f)), $f^{-1}(N(f(S(f))))$ has the structures of trivial bundles as in Definiiton 2 such that a normal fiber $F_{f(S(f))}$ of f(S(f)) corresponding to the bundles $f^{-1}(N(f(S(f))))$ is diffeomorphic to the standard closed disc D^{m-n+1} . The m-dimensional standard sphere S^m admits such a map into \mathbb{R}^n for $m \ge n \ge 2$. In section 5 of [19], on every homotopy sphere whose dimension is $m \ (2 \le m < 4, m > 4)$ and the 4-dimensional standard sphere S^4 , such maps into \mathbb{R}^2 are constructed.
 - (2) [[11], [13], [14]] Let $F \neq \emptyset$ be a closed and connected manifold. Let M have the structure of an F-bundle over S^n . Then, M admits a round fold map $f: M \to \mathbb{R}^n$ such that the followings hold.
 - (a) f is C^{∞} S-trivial.
 - (b) For an axis L of f, $f^{-1}(L)$ is diffeomorphic to $F \times [0,1]$.
 - (c) Two connected components of the fiber of a point in a proper core of f is regarded as fibers of the F-bundle over S^n .
 - (3) [[10], [11], [13]] Let $m, n \in \mathbb{N}$, $n \geq 2$ and $m \geq 2n$. Let M be an m-dimensional manifold represented as the connected sum of $l \in \mathbb{N}$ closed oriented manifolds having the sturctures of S^{m-n} -bundles over S^n admits a C^∞ L-trivial round fold map f into \mathbb{R}^n such that the followings hold.
 - (a) All the regular fibers of f are disjoint unions of finite copies of S^{m-n} .
 - (b) The number of connected components of the singular set S(f) and the number of connected components of the fiber of a point in a proper core of f are l.

- (c) For any connected component C of f(S(f)) and a small closed tubular neighborhood N(C) of C, $f^{-1}(N(C))$ has the structures of trivial bundles as in Definition 2 such that a normal fiber F_C of C corresponding to the bundles $f^{-1}(N(C))$ is diffeomorphic to a disjoint union of a finite number of the following manifolds.
 - (i) The (m-n+1)-dimensional standard closed disc D^{m-n+1} .
 - (ii) The standard (m-n+1)-dimensional sphere S^{m-n+1} with the interior of a union of disjoint three (m-n+1)-dimensional standard closed discs removed.
- (d) All the connected components of the fiber of a point in a proper core of f are regarded as fibers of the S^{m-n} -bundles over S^n and a fiber of any S^{m-n} -bundle over S^n appeared in the connected sum is regarded as a connected component of the fiber of a point in a proper core of f.
- (4) Special generic maps into \mathbb{R}^n whose singular sets are disjoint unions of two standard spheres such that the restriction maps to the singular sets are embeddings are round fold maps. Let M be a closed m-dimensional manifold admitting such a map $f: M \to \mathbb{R}^n$ ($m \ge n \ge 2$). Since f is special generic, for each connected component C of f(S(f)), there exists a small closed tubular neighborhood N(C) such that $f^{-1}(N(C))$ has the structures of isomorphic trivial bundles whose fibers are diffeomorphic to the standard closed disc D^{m-n+1} as in Definition 2.

We easily know that the m-dimensional manifolds admitting such maps into \mathbb{R}^n ($m \geq n \geq 2$) being C^{∞} L-trivial are always homeomorphic to $S^{n-1} \times S^{m-n+1}$ and that $S^{n-1} \times S^{m-n+1}$ admits such a map being C^{∞} S-trivial and C^{∞} L-trivial by virtue of a trivial spinning construction. Moreover, for example, in the case where m-n=0,1,2,3 holds, the source manifold of such a map is diffeomorphic to the product $S^{n-1} \times S^{m-n+1}$. It easily follows from the fact that the diffeomorphism groups of the spheres S^1 , S^2 and S^3 are homotopy equivalent to the groups of all the linear transformations on the spheres ([9], [17]).

For explicit theories of such special generic maps, see also [19].

- 3. P-operations and constructions of round fold maps of new types on manifolds having the structures of S^1 -bundles
- 3.1. **P-operations.** The following proposition has been shown in [14].

Proposition 1 ([14]). Let M be a closed manifold of dimension m and $f: M \to \mathbb{R}^n$ be a C^{∞} L-trivial round fold map $(m \ge n \ge 2)$. Let $F \neq \emptyset$ be a closed manifold and M' be a closed manifold having the structure of an F-manifold over M such that for any connected component C of f(S(f)) and a small closed tubular neighborhood N(C) of C, the restriction to $f^{-1}(N(C))$ is a trivial bundle. Then on M' there exists a C^{∞} L-trivial round fold map $f': M' \to \mathbb{R}^n$.

In the proof of this proposition, we use the notation $D^n_r:=\{(x_1,\cdots,x_n)\in\mathbb{R}^n\mid \sum_{k=1}^n x_k^2\leq r\}$ appeared in Definition 1.

Proof of Proposition 3. We assume that f(M) is diffeomorphic to D^n . We can prove the theorem similarly if f(M) is not diffeomorphic to D^n .

We may assume that $f: M \to \mathbb{R}^n$ is a normal form. Let S(f) consist of l connected components. Set $P_0 := D^n_{\frac{1}{2}}$ and $P_k := D^n_{k+\frac{1}{2}} - \operatorname{Int} D^n_{k-\frac{1}{2}}$ for an integer $1 \le k \le l$.

Then $f^{-1}(P_k)$ has the structure of a trivial C^{∞} bundle over $\partial D^n_{k-\frac{1}{2}}$ or $\partial D^n_{k+\frac{1}{2}}$ with fibers diffeomorphic to a compact C^{∞} manifold, which we denote by E_k , such that $f|_{f^{-1}(\partial D^n_{k-\frac{1}{2}})}$ and $f|_{f^{-1}(\partial D^n_{k+\frac{1}{2}})}$ give the structures of subbundles (we denote the fibers of two subbundles by $E_k^1 \subset E_k$ and $E_k^2 \subset E_k$, respectively) for $1 \leq k \leq l$. For any integer $1 \le k \le l$ and a diffeomorphism ϕ_k from $f^{-1}(\partial D^n_{k-\frac{1}{2}}) \subset f^{-1}(P_k)$ onto $f^{-1}(\partial D^n_{k-\frac{1}{2}}) \subset f^{-1}(P_{k-1})$ regarded as a bundle isomorphism between the two trivial bundles over standard spheres inducing the identification between the base spaces, M is regarded as $(\cdots((f^{-1}(D^n_{\frac{1}{2}}))\bigcup_{\phi_1}f^{-1}(P_1))\cdots)\bigcup_{\phi_l}f^{-1}(P_l)$ and for any integer $1 \le k \le l$ and a diffeomorphism Φ_k from $f^{-1}(\partial D^n_{k-\frac{1}{\alpha}}) \times F \subset f^{-1}(P_k) \times I$ F onto $f^{-1}(\partial D^n_{k-\frac{1}{2}}) \times F \subset f^{-1}(P_{k-1}) \times F$ regarded as a bundle isomorphism between the two trivial F-bundles inducing ϕ_k , M' is regarded as $(\cdots ((f^{-1}(D^n_{\frac{1}{n}})\times$ $F)\bigcup_{\Phi_1}(f^{-1}(P_1)\times F))\cdots\bigcup_{\Phi_l}(f^{-1}(P_l)\times F)$. We construct a map on $f^{-1}(P_k)\times F$. This manifold has the structure of a trivial bundle over $\partial D^n_{k-\frac{1}{2}}$ or $\partial D^n_{k+\frac{1}{2}}$ with fibers diffeomorphic to $E_k \times F$. On $E_k \times F$ there exists a Morse function f_k such that the followings hold.

- (1) $\tilde{f}_k(E_k \times F) \subset [k \frac{1}{2}, k + \frac{1}{2}]$ and $\tilde{f}_k(\operatorname{Int}(E_k \times F)) \subset (k \frac{1}{2}, k + \frac{1}{2})$ hold. (2) $\tilde{f}_k(E_k^1 \times F) = \{k \frac{1}{2}\}$ holds if $E_k^1 \times F$ is non-empty. (3) $\tilde{f}_k(E_k^2 \times F) = \{k + \frac{1}{2}\}$ holds if $E_k^2 \times F$ is non-empty.

generic map from S^n into \mathbb{R}^n presented in Example 1 (1).

(4) Singular points of \tilde{f}_k are in the interior of $E_k \times F$ and at two distinct singular points, the values are always distinct.

We obtain a map $\mathrm{id}_{S^{n-1}} \times \tilde{f}_k : S^{n-1} \times E_k \times F \to S^{n-1} \times [k-\frac{1}{2},k+\frac{1}{2}]$. We can identify $S^{n-1} \times [k-\frac{1}{2},k+\frac{1}{2}]$ with $P_k = D^n_{k+\frac{1}{2}} - \mathrm{Int} D^n_{k-\frac{1}{2}}$ by identifying $(p,t) \in S^{n-1} \times [k-\frac{1}{2},k+\frac{1}{2}]$ with $tp \in P_k$ where we regard S^{n-1} as the unit sphere of dimension n-1. By gluing the composition of the projection from $f^{-1}(D^n_{\frac{1}{2}})\times F$ onto $f^{-1}(D^n_{\frac{1}{2}})$ and $f|_{f^{-1}(D^n_{\frac{1}{2}})}: f^{-1}(D^n_{\frac{1}{2}}) \to D^n_{\frac{1}{2}}$ and the family $\{\mathrm{id}_{S^{n-1}} \times \tilde{f}_k\}$ together by using the family $\{\Phi_k\}$ and the family of identifications in the target manifold \mathbb{R}^n , we obtain a new round fold map $f': M' \to \mathbb{R}^n$.

In the proof, from $f: M \to \mathbb{R}^n$, we obtain $f': M' \to \mathbb{R}^n$. We call the operation of constructing f' from f a P-operation by F to f. For example, for any closed manifold $F \neq \emptyset$ and on any manifold admitting the structure of a F-bundle over the n-dimensional $(n \geq 2)$ standard sphere, we obtain a round fold map as introduced in Example 1 (2) by a P-operation by F to a special

3.2. P-operations by S^1 to round fold maps and their applications. We apply P-operations by S^1 to obtain round fold maps which do not appear in [14]. Before this, we recall fundamental terms and facts on S^1 -bundles. In this paper, for a topological space X, we denote the k-th (co)homology group of X whose coefficient ring is R by $H_k(X;R)$ (resp. $H^k(X;R)$).

It is well-known that S^1 -bundles are regarded as bundles whose structure groups consist of linear transformations on the circle and regarded as the 2nd orthogonal group O(2). An S^1 -bundle is said to be *orientable* if the structure group consist of orientation preserving linear transformations and regarded as the 2nd rotation group $SO(2) \subset O(2)$. We can consider Stiefel-Whitney classes of S^1 -bundles, which are cohomology classes of the base spaces whose coefficient rings are $\mathbb{Z}/2\mathbb{Z}$, including their 1st Stiefel-Whitney classes (2nd Stiefel-Whitney classes), which are 1st cohomology classes (resp. 2nd cohomology classes) of the base spaces with coefficients $\mathbb{Z}/2\mathbb{Z}$. If an S^1 -bundle is orientable, then we can orient the bundle and consider its Euler class, which is a 2nd cohomology class of the base space whose coefficient ring is \mathbb{Z} .

For an oriented S^1 -bundle over a 2-dimensional oriented closed and connected manifold, we can consider the pairing of the Euler class and the fundamental class of the base space and call the obtained integer the *Euler number* of the bundle. We introduce known facts on classifications of S^1 -bundles without proofs.

Proposition 2. Let X be a topological space.

- (1) The 1st Stiefel-Whitney class $\alpha \in H^1(X; \mathbb{Z}/2\mathbb{Z})$ of an S^1 -bundle over X vanishes if and only if the bundle is orientable.
- (2) For any $\alpha \in H^2(X; \mathbb{Z})$, there exists an oriented S^1 -bundle over X whose Euler class is α .
- (3) Two oriented S^1 -bundles on X are isomorphic if the Euler classes are same.
- (4) Two oriented S^1 -bundles over a 2-dimensional oriented closed and connected manifold are isomorphic if and only if the Euler numbers of the bundles coincide. Furthermore, manifolds admitting the structures of S^1 bundles over a 2-dimensional oriented closed and connected manifolds are homeomorphic (diffeomorphic) if and only if the absolute values of the Euler numbers of the bundles coincide. The 1st homology group and the 2nd cohomology group of the manifold admitting an oriented S^1 -bundle such that the absolute value of the Euler number is $k \in \mathbb{N} \setminus \{0\}$ with coefficient ring \mathbb{Z} are isomorphic to the direct sum of the 1st (co)homology group of the base space with coefficient ring \mathbb{Z} and $\mathbb{Z}/k\mathbb{Z}$. The latter part of this direct sum is generated by the 1st homology class represented by a fiber of the bundle. Furthermore, for any section of the restriction of the bundle to any 1-skelton representing some element of 1st homology group of the base space with coefficient ring \mathbb{Z} , the image of the section represents an element of the 1st homology group before and if we consider the direct sum here, the 1st homology class is of the form (x.0).

For the theory of bundles whose structure groups consist of linear transformations including S^1 -bundles and vector bundles and their characteristic classes including Stiefel-Whitney classes and Euler classes, see also [16] for example.

Let M be a closed manifold of dimension m and let $f: M \to \mathbb{R}^n$ ($m \ge n \ge 2$) be a C^{∞} L-trivial round fold map. For any connected component C of f(S(f)), we denote by N(C) a small closed tubular neighborhood of C such that the inverse image has the structures of trivial bundles as in Definition 2 and by F_C a normal fiber of C corresponding to the bundles $f^{-1}(N(C))$.

We now construct a graph corresponding to $f_F := (f, \{F_C\}_{C \in \pi_0(S(f))})$, where $C \in \pi_0(f(S(f)))$ represents the element of the set $\pi_0(f(S(f)))$ of all the connected components of f(S(f)).

Let V_{f_F} be the set of all the closures of connected components of the regular value set $\mathbb{R}^n - f(S(f))$. Let R be a commutative ring. We define the set $E_{f_F}^{(H_k,R)} \subset V_{f_F} \times V_{f_F}$ so that the followings hold.

- (1) The set $E_{f_F}^{(H_k,R)}$ consists of elements (P_1,P_2) of the set $V_{f_F} \times V_{f_F}$ such that the intersection $P_1 \cap P_2$ is not empty.
- (2) A pair (P_1, P_2) of closures of connected components of the space $\mathbb{R}^n f(S(f))$ such that $P_1 \cap P_2$ is not empty and that P_1 is in the bounded connected component of $\mathbb{R}^n \operatorname{Int} P_2$, $(P_1, P_2) \in E_{f_F}$ holds if and only if for the connected component $C = P_1 \cap P_2$ of f(S(f)), the homomorphism from $H_k(\partial_i F_C; R)$ into $H_k(F_C; R)$ induced by the natural inclusion is surjective.
- (3) A pair (P_1, P_2) of closures of connected components of the space $\mathbb{R}^n f(S(f))$ such that $P_1 \cap P_2$ is not empty and that P_2 is in the bounded connected component of $\mathbb{R}^n \operatorname{Int} P_1$, $(P_1, P_2) \in E_{f_F}$ holds if and only if for the connected component $C := P_1 \cap P_2$ of f(S(f)), the homomorphism from $H_k(\partial_o F_C; R)$ into $H_k(F_C; R)$ induced by the natural inclusion is surjective.

Thus, we obtained a graph $(V_{f_F}, E_{f_F}^{(H_k, R)})$ and we call it the (H_k, R) -graph of $f_F = (f, \{F_C\}_{C \in \pi_0(f(S(f)))})$.

Definition 4. For the (H_k, R) -graph of $f_F = (f, \{F_C\}_{C \in \pi_0(f(S(f)))})$, a non-empty subset $S \subset V_{f_F}$ is said to be a *starting set* of the graph if for any $x \in V_{f_F}$, there exists a path from a vertex in S to x.

Let R_i be a commutative ring and let k_i be an integer for i=1,2. For the (H_{k_1},R_1) -graph and the (H_{k_2},R_2) -graph of $f_F=(f,\{F_C\}_{C\in\pi_0(f(S(f)))})$, the (H_{k_1},R_1) -graph and the (H_{k_2},R_2) -graph are start-equivalent on the set S if for any vertex $s\in V_{f_F}$, there exists a vertex $s'\in S$ such that there exists a path of both of these graphs from s' to s.

We show the following theorem.

Theorem 1. Let M be a closed manifold of dimension m and $f: M \to \mathbb{R}^n$ $(m \ge n \ge 2)$ be a C^∞ L-trivial round fold map. For any connected component C of f(S(f)), we denote by N(C) a small closed tubular neighborhood of C such that the inverse image has the structures of trivial bundles as in Definition 2 and by F_C a normal fiber of C corresponding to the bundles $f^{-1}(N(C))$ over C. Let F_C satisfy $H^2(F_C; \mathbb{Z}) \cong \{0\}$. Let S be a starting set of the $(H_1, \mathbb{Z}/2\mathbb{Z})$ -graph of $f_F = (f, \{F_C\}_{C \in \pi_0(f(S(f)))})$. Let F_s be the regular fiber of a point in a connected component of $\mathbb{R}^n - f(S(f))$ whose closure is represented as a vertex $s \in S$ of the graph above and let $i_s: F_s \to M$ be the natural inclusion. Then, we have the followings.

- (1) Let $n \geq 4$ hold. Then, by a P-operation to f, for any closed manifold M' having the structure of an S^1 -bundle over M whose 1st Stiefel-Whitney class vanishes on $i_{s*}(H_1(F_s; \mathbb{Z}/2\mathbb{Z})) \subset H_1(M; \mathbb{Z}/2\mathbb{Z})$ for any $s \in S$, we can obtain a round fold map $f': M' \to \mathbb{R}^n$.
- (2) Let n=3 hold. Furthermore, we assume that for any commutative ring R, the $(H_1, \mathbb{Z}/2\mathbb{Z})$ -graph of $f_F = (f, \{F_C\}_{C \in \pi_0(f(S(f)))})$ and the (H_0, R) -graph of $f_F = (f, \{F_C\}_{C \in \pi_0(f(S(f)))})$ are start-equivalent on the set S. Then, by a P-operation to f, for any closed manifold M' having the structure of an S^1 -bundle over M such that the followings hold, we can obtain a round fold map $f': M' \to \mathbb{R}^n$.
 - (a) The 1st Stiefel-Whitney class vanishes on $i_{s_*}(H_1(F_s; \mathbb{Z}/2\mathbb{Z})) \subset H_1(M; \mathbb{Z}/2\mathbb{Z})$ for any $s \in S$.
 - (b) For $s \in S$, let C_s' be a 2-dimensional sphere in a closure of a connected component of $\mathbb{R}^3 f(S(f))$ represented by $s \in S$ such that for

a connected component C_s of f(S(f)) and the small closed tubular neighborhood $N(C_s)$ defined before, C_s' is a connected component of $\partial N(C_s)$. The restriction of the bundle to the image of a section of the trivial bundle given by $f|_{f^{-1}(C_s')}: f^{-1}(C_s') \to C_s'$ is trivial.

(3) Let n=2 hold. Furthermore, we assume that for any commutative ring R, the $(H_1, \mathbb{Z}/2\mathbb{Z})$ -graph of $f_F = (f, \{F_C\}_{C \in \pi_0(f(S(f)))})$ and the (H_0, R) -graph of $f_F = (f, \{F_C\}_{C \in \pi_0(f(S(f)))})$ are start-equivalent on the set S and that the $(H_1, \mathbb{Z}/2\mathbb{Z})$ -graph of $f_F = (f, \{F_C\}_{C \in \pi_0(f(S(f)))})$ and the (H_1, \mathbb{Z}) -graph of $f_F = (f, \{F_C\}_{C \in \pi_0(f(S(f)))})$ are start-equivalent on the set S. For $s \in S$, let C_s' be a circle in a closure of a connected component of $\mathbb{R}^2 - f(S(f))$ represented by $s \in S$ such that for a connected component C_s of f(S(f)) and the small closed tubular neighborhood $N(C_s)$ defined before, C_s' is a connected component of $\partial N(C_s)$. The restriction of the bundle to the total space of the trivial bundle given by $f|_{f^{-1}(C_s')}: f^{-1}(C_s') \to C_s'$ is orientable and trivial.

Then, by a P-operation to f, for any closed manifold M' having the structure of an S^1 -bundle over M, we can obtain a round fold map $f': M' \to \mathbb{R}^n$.

Proof. In all the cases, it suffices to prove that for any connected component $C \subset f(S(f))$, given S^1 -bundles over M are trivial on $f^{-1}(N(C))$. Let $C_0 \subset f(S(f))$ be a connected component which is in the closures of two connected component of $\mathbb{R}^n - f(S(f))$ represented by vertices $s_1, s_2 \in V_{f_F}$ of the graph and let $(s_1, s_2) \in E_{f_F}^{(H_1, \mathbb{Z}/2\mathbb{Z})}$. Let the boundary $\partial N(C_0)$ of $N(C_0)$ defined in the assumption of this theorem consist of two connected components C_1 and C_2 as in Definition 2 and let C_j be in the closure of a connected component of $\mathbb{R}^n - f(S(f))$ represented by s_j . In addition, we denote the fiber of the bundle $f^{-1}(C_j)$ over C_j as in Definition 2 by $\partial_i F_{C_0}$.

We prove the statement in the first case. The homomorphism from $H_1(f^{-1}(C_1); \mathbb{Z}/2\mathbb{Z}) \cong H_1(\partial_1 F_{C_0}; \mathbb{Z}/2\mathbb{Z})$ into $H_1(f^{-1}(N(C_0)); \mathbb{Z}/2\mathbb{Z}) \cong H_1(F_{C_0}; \mathbb{Z}/2\mathbb{Z})$ induced by the natural inclusion is surjective, since the homomorphism from $H_1(\partial_1 F_{C_0}; \mathbb{Z}/2\mathbb{Z})$ into $H_1(F_{C_0}; \mathbb{Z}/2\mathbb{Z})$ induced by the natural inclusion is assumed to be surjective. The kernel of the homomorphism from $H^1(f^{-1}(N(C_0)); \mathbb{Z}/2\mathbb{Z})$ into $H^1(f^{-1}(C_1); \mathbb{Z}/2\mathbb{Z})$ induced by the natural inclusion is $\{0\}$. $H^2(f^{-1}(N(C_0)); \mathbb{Z}) \cong H^2(C_0 \times F_{C_0}; \mathbb{Z}) \cong \{0\}$ holds since $H^2(F_{C_0}; \mathbb{Z}) \cong \{0\}$ is assumed. Thus, if the restriction of an S^1 -bundle over M to $f^{-1}(C_1)$ is orientable, then the restriction of the original bundle to $f^{-1}(N(C_0))$ is orientable and it is a trivial bundle.

Consider an (n-1)-dimensional sphere C_s' in a closure of a connected component of $\mathbb{R}^n - f(S(f))$ represented by $s \in S$ such that for a connected component C_s of f(S(f)) and a small closed tubular neighborhood $N(C_s)$ as in Definition 2, C_s' is a connected component of $\partial N(C_s)$. $f^{-1}(C_s')$ is diffeomorphic to $C_s' \times F_s$ and $C_s \times F_s$. $H^1(f^{-1}(C_s'); \mathbb{Z}/2\mathbb{Z}) \cong H^1(C_s \times F_s; \mathbb{Z}/2\mathbb{Z}) \cong H^1(F_s; \mathbb{Z}/2\mathbb{Z})$ holds and if the 1st Stiefel-Whitney class of an S^1 -bundle over M vanishes on F_s , then the restriction of the S^1 -bundle over M to $f^{-1}(C_s')$ is trivial. Thus, by the discussion in the previous paragraph together with an induction based on the fact that S is a starting set of the graph, for any connected component C of f(S(f)), the restriction of any S^1 -bundle over M whose 1st Stiefel-Whitney class vanishes on $i_*(H_1(F_s; \mathbb{Z}/2\mathbb{Z})) \subset H_1(M; \mathbb{Z}/2\mathbb{Z})$ ($s \in S$) to $f^{-1}(N(C))$ is orientable and it is a trivial bundle. This completes the proof of the first case.

We prove the statement in the second case. As in the first case, the homomorphism from $H_1(f^{-1}(C_1); \mathbb{Z}/2\mathbb{Z}) \cong H_1(\partial_1 F_{C_0}; \mathbb{Z}/2\mathbb{Z})$ into $H_1(f^{-1}(N(C_0)); \mathbb{Z}/2\mathbb{Z}) \cong$ $H_1(F_{C_0}; \mathbb{Z}/2\mathbb{Z})$ induced by the natural inclusion is surjective, since the homomorphism from $H_1(\partial_1 F_{C_0}; \mathbb{Z}/2\mathbb{Z})$ into $H_1(F_{C_0}; \mathbb{Z}/2\mathbb{Z})$ induced by the natural inclusion is assumed to be surjective. The kernel of the homomorphism from $H^1(f^{-1}(N(C_0)); \mathbb{Z}/2\mathbb{Z})$ into $H^1(f^{-1}(C_1); \mathbb{Z}/2\mathbb{Z})$ induced by the natural inclusion is $\{0\}$. We have $H^2(f^{-1}(N(C_0)); \mathbb{Z}) \cong$ $H^2(C_0 \times F_{C_0}; \mathbb{Z}) \cong H^2(C_0; \mathbb{Z}) \otimes H^0(F_{C_0}; \mathbb{Z}) \cong \mathbb{Z} \otimes H^0(F_{C_0}; \mathbb{Z}) \text{ since } H^2(F_{C_0}; \mathbb{Z}) \cong \{0\}.$ We also have $H^2(f^{-1}(C_1); \mathbb{Z}) \cong H^2(C_0 \times \partial_1 F_{C_0}; \mathbb{Z}) \cong (H^2(C_0; \mathbb{Z}) \otimes H^0(\partial_1 F_{C_0}; \mathbb{Z})) \oplus (H^2(C_0; \mathbb{Z}) \otimes H^0(\partial_1 F_{C_0}; \mathbb{Z}))$ $(H^0(C_0; \mathbb{Z}) \otimes H^2(\partial_1 F_{C_0}; \mathbb{Z})) \cong (\mathbb{Z} \otimes H^0(\partial_1 F_{C_0}; \mathbb{Z})) \oplus (H^0(C_0; \mathbb{Z}) \otimes H^2(\partial_1 F_{C_0}; \mathbb{Z}))$ since $H^2(F_{C_0}; \mathbb{Z}) \cong \{0\}$. Moreover, we have $H_2(f^{-1}(C_1); \mathbb{Z}) \cong H_2(\partial_1 F_{C_0}; \mathbb{Z}) \oplus$ $(H_0(\partial_1 F_{C_0}; \mathbb{Z}) \otimes \mathbb{Z})$ and $H_2(f^{-1}(N(C_0)); \mathbb{Z}) \cong H_2(C_0 \times F_{C_0}; \mathbb{Z}) \cong \mathbb{Z} \otimes H_0(F_{C_0}; \mathbb{Z}).$ The homomorphism from $H_0(\partial_1 F_{C_0}; \mathbb{Z})$ into $H_0(F_{C_0}; \mathbb{Z})$ induced by the natural inclusion is assumed to be surjective. Thus, by the discussion here, the kernel of the homomorphism from $H^2(f^{-1}(N(C_0));\mathbb{Z})$ into $H^2(f^{-1}(C_1);\mathbb{Z})$ induced by the natural inclusion is $\{0\}$. Thus, if the restriction of a bundle over M to $f^{-1}(C_1)$ is trivial, then the restriction of the original bundle to $f^{-1}(N(C_0))$ is orientable by a discussion similar to that of the proof of the first case and it is a trivial bundle. Consider a 2-dimensional sphere C_s in a closure of a connected component of $\mathbb{R}^n - f(S(f))$ represented by $s \in S$ such that for a connected component C_s of f(S(f)) and a small closed tubular neighborhood $N(C_s)$ defined in the assumption of this theorem, C_s is a connected component of $\partial N(C_s)$. $f^{-1}(C_s)$ is diffeomorphic to $C_s' \times F_s$ and $C_s \times F_s$. $H^1(f^{-1}(C_s'); \mathbb{Z}/2\mathbb{Z}) \cong H^1(C_s \times F_s; \mathbb{Z}/2\mathbb{Z}) \cong H^1(F_s; \mathbb{Z}/2\mathbb{Z})$ holds. $H^2(f^{-1}(N(C));\mathbb{Z}) \cong H^2(C \times F_C;\mathbb{Z}) \cong H^2(C;\mathbb{Z}) \otimes H^0(F_C;\mathbb{Z}) \cong \mathbb{Z} \otimes$ $H^0(F_C;\mathbb{Z})$ and $H^2(f^{-1}(C_s');\mathbb{Z}) \cong H^2(C_s \times F_s;\mathbb{Z}) \cong (H^2(C_s;\mathbb{Z}) \otimes H^0(F_s;\mathbb{Z})) \oplus$ $(\mathbb{Z} \otimes H^2(F_s;\mathbb{Z}))$ also hold. Thus, by the structures of these cohomology groups, if the 1st Stiefel-Whitney class of an S^1 -bundle over M vanishes on F_s and the restriction of the bundle to the image of a section of the trivial bundle given by $f|_{f^{-1}(C_{s'})}: f^{-1}(C_{s'}) \to C_{s'}$ is trivial, then the restriction of the S^1 -bundle over M to $f^{-1}(C_s')$ is trivial. Thus, by the discussion in the previous paragraph together with an induction based on the assumptions that S is a starting set of the $(H_1,\mathbb{Z}/2\mathbb{Z})$ -graph of $f_F=(f,\{F_C\}_{C\in\pi_0(f(S(f)))})$ and that the graph and the (H_0,R) -graph of $f_F=(f,\{F_C\}_{C\in\pi_0(f(S(f)))})$ are start-equivalent on the set S, for any connected component C of f(S(f)), if we restrict any S^1 -bundle over M whose 1st Stiefel-Whitney class vanishes on $i_*(H_1(F_s; \mathbb{Z}/2\mathbb{Z})) \subset H_1(M; \mathbb{Z}/2\mathbb{Z})$ $(s \in S)$ and whose restriction of the bundle to the image of a section of the trivial bundle given by $f|_{f^{-1}(C_{s'})}: f^{-1}(C_{s'}) \to C_{s'}$ is trivial to $f^{-1}(N(C))$, then the resulting bundle is orientable and trivial. This completes the proof of the second case. We prove the statement in the last case. The homomorphism from $H_1(f^{-1}(C_1); \mathbb{Z}/2\mathbb{Z})$ $H_1(\partial_1 F_{C_0}; \mathbb{Z}/2\mathbb{Z}) \oplus (H_1(C_0; \mathbb{Z}/2\mathbb{Z}) \otimes H_0(\partial_1 F_{C_0}; \mathbb{Z}/2\mathbb{Z}))$ into $H_1(f^{-1}(N(C_0)); \mathbb{Z}/2\mathbb{Z}) \cong H_0(\partial_1 F_{C_0}; \mathbb{Z}/2\mathbb{Z})$ $H_1(F_{C_0}; \mathbb{Z}/2\mathbb{Z}) \oplus (H_1(C_0; \mathbb{Z}/2\mathbb{Z}) \otimes H_0(F_{C_0}; \mathbb{Z}/2\mathbb{Z}))$ induced by the natural inclusion is surjective, since the homomorphism from $H_k(\partial_1 F_{C_0}; \mathbb{Z}/2\mathbb{Z})$ into $H_k(F_{C_0}; \mathbb{Z}/2\mathbb{Z})$ induced by the natural inclusion is assumed to be surjective for k=0,1. It follows that the kernel of the homomorphism from $H^1(f^{-1}(N(C_0)); \mathbb{Z}/2\mathbb{Z})$ into $H^1(f^{-1}(C_1); \mathbb{Z}/2\mathbb{Z})$ induced by the natural inclusion is $\{0\}$. The homomorphism from $H_k(\partial_1 F_{C_0}; \mathbb{Z})$ into $H_k(F_{C_0}; \mathbb{Z})$ and the homomorphism from $H_k(\partial_1 F_{C_0}; \mathbb{Z})$ into $H_k(F_{C_0}; \mathbb{Z})$ induced by the natural inclusions are assumed to be surjective for k=0,1. Thus, $H_2(f^{-1}(N(C_0));\mathbb{Z})\cong (H_1(C_0;\mathbb{Z})\otimes H_1(F_{C_0};\mathbb{Z}))\oplus$ $H_2(F_{C_0}; \mathbb{Z})$ is regarded also as the direct sum of the following two groups.

- (1) The image of the restriction of the homomorphism from $H_2(f^{-1}(C_1); \mathbb{Z}) \cong (H_1(C_0; \mathbb{Z}) \otimes H_1(\partial_1 F_{C_0}; \mathbb{Z})) \oplus H_2(\partial_1 F_{C_0}; \mathbb{Z})$ into $H_2(f^{-1}(N(C_0)); \mathbb{Z})$ induced by the natural inclusion to the part $H_1(C_0; \mathbb{Z}) \otimes H_1(\partial_1 F_{C_0}; \mathbb{Z})$.
- (2) $H_2(F_{C_0}; \mathbb{Z})$.

Hence, the kernel of the homomorphism from $H^2(f^{-1}(N(C_0)); \mathbb{Z}) \cong (H^1(C_0; \mathbb{Z}) \otimes H^1(F_{C_0}; \mathbb{Z})) \oplus H^2(F_{C_0}; \mathbb{Z})$ into $H^2(f^{-1}(C_1); \mathbb{Z}) \cong (H^1(C_0; \mathbb{Z}) \otimes H^1(\partial_1 F_{C_0}; \mathbb{Z})) \oplus H^2(\partial_1 F_{C_0}; \mathbb{Z})$ induced by the natural inclusion is $\{0\}$.

By the arguments before, if the restriction of an S^1 -bundle over M to $f^{-1}(C_1)$ is orientable and the Euler class of the (oriented) bundle obtained by the restriction vanishes, then the restriction of the S^1 -bundle over M to $f^{-1}(N(C_0))$ is orientable and the Euler class of the (oriented) bundle obtained by the restriction vanishes. Consider a circle C_s' in a closure of a connected component of $\mathbb{R}^2 - f(S(f))$ represented by $s \in S$ such that for a connected component C_s of f(S(f)) and a small closed tubular neighborhood $N(C_s)$ as in Definition 2, C_s' is a connected component of $\partial N(C)$. $f^{-1}(C_s)$ is diffeomorphic to $C_s \times F_s$ and $C_s \times F_s$. We have $H^1(f^{-1}(C_s'); \mathbb{Z}/2\mathbb{Z}) \cong H^1(C_s \times F_s; \mathbb{Z}/2\mathbb{Z}) \cong \mathbb{Z}/2\mathbb{Z} \otimes H^1(F_s; \mathbb{Z}/2\mathbb{Z}), H^2(f^{-1}(N(C)); \mathbb{Z}) \cong H^1(f^{-1}(C_s'); \mathbb{Z}/2\mathbb{Z}) \cong H^1(C_s \times F_s; \mathbb{Z}/2\mathbb{Z}) \cong H^1(C$ $H^2(C \times F_C; \mathbb{Z}) \cong H^1(C; \mathbb{Z}) \otimes H^1(F_C; \mathbb{Z}) \cong \mathbb{Z} \otimes H^1(F_C; \mathbb{Z}) \text{ and } H^2(f^{-1}(C_s'); \mathbb{Z}) \cong \mathbb{Z}$ $H^2(C_s \times F_s; \mathbb{Z}) \cong (H^1(C_s; \mathbb{Z}) \otimes H^1(F_s; \mathbb{Z})) \oplus (H^0(C_s; \mathbb{Z}) \otimes H^2(F_s; \mathbb{Z})) \cong (\mathbb{Z} \otimes H^1(F_s; \mathbb{Z})) \oplus (H^1(C_s; \mathbb{Z}) \otimes H^1(F_s; \mathbb{Z})) \cong (\mathbb{Z} \otimes H^1(F_s; \mathbb{Z})) \oplus (H^1(C_s; \mathbb{Z}) \otimes H^1(F_s$ $H^1(F_s;\mathbb{Z})) \oplus (\mathbb{Z} \otimes H^2(F_s;\mathbb{Z}))$. S is a starting set of the $(H_1,\mathbb{Z}/2\mathbb{Z})$ -graph of $f_F =$ $(f, \{F_C\}_{C \in \pi_0(f(S(f)))})$. This graph and the (H_0, R) -graph of $f_F = (f, \{F_C\}_{C \in \pi_0(f(S(f)))})$ are start-equivalent on the set S. This graph and the (H_1,\mathbb{Z}) -graph of f_F $(f, \{F_C\}_{C \in \pi_0(f(S(f)))})$ are also start-equivalent on the set S. Thus, by the discussion in the previous paragraph, for any connected component C of f(S(f)), if we restrict any S^1 -bundle over M whose restriction of the bundle to the total space of the trivial bundle given by $f|_{f^{-1}(C_s')}: f^{-1}(C_s') \to C_s'$ is trivial to $f^{-1}(N(C))$, then the resulting bundle is orientable and trivial. This completes the proof of the last case.

This completes the proof of all the statements.

We show Theorem 1 in the case where the manifold M is connected and n=2,3

Theorem 2. Let M be a closed and connected manifold of dimension m and $f: M \to \mathbb{R}^n$ ($m \ge n \ge 2$) be a C^∞ L-trivial round fold map. For any connected component C of f(S(f)), we denote by N(C) a small closed tubular neighborhood of C such that the inverse image has the structures of trivial bundles as in Definition 2 and by F_C a normal fiber of C corresponding to the bundles $f^{-1}(N(C))$ over C. Let F_C satisfy $H^2(F_C; \mathbb{Z}) \cong \{0\}$. Let S be a starting set of the $(H_1, \mathbb{Z}/2\mathbb{Z})$ -graph of $f_F = (f, \{F_C\}_{C \in \pi_0(f(S(f)))})$. Let F_s be the regular fiber of a point in a connected component of $\mathbb{R}^n - f(S(f))$ whose closure is represented as a vertex $s \in S$ of the graph above and let $i_s: F_s \to M$ be the natural inclusion. Then, we have the followings.

- (1) Let n=3 hold. Then, by a P-operation to f, for any closed manifold M' having the structure of an S^1 -bundle over M such that the followings hold, we can obtain a round fold map $f': M' \to \mathbb{R}^3$.
 - (a) The 1st Stiefel-Whitney class vanishes on $i_{s_*}(H_1(F_s; \mathbb{Z}/2\mathbb{Z})) \subset H_1(M; \mathbb{Z}/2\mathbb{Z})$ for any $s \in S$.
 - (b) For a 2-dimensional sphere C_0' in a connected component of $(\mathbb{R}^3 f(S(f))) \cap f(M)$ such that for a connected component C_0 of f(S(f)) and the small closed tubular neighborhood $N(C_0)$ defined before, C_0' is

a connected component of $\partial N(C_0)$, the restriction of the bundle to the image of a section of the trivial bundle given by $f|_{f^{-1}(C_0')}: f^{-1}(C_0') \to C_0'$ is trivial.

(2) Let n=2 hold. Furthermore, we assume that the $(H_1, \mathbb{Z}/2\mathbb{Z})$ -graph of $f_F = (f, \{F_C\}_{C \in \pi_0(f(S(f)))})$ and the (H_1, \mathbb{Z}) -graph of $f_F = (f, \{F_C\}_{C \in \pi_0(f(S(f)))})$ are start-equivalent on the set S.

For $s \in S$, let C_s' be a circle in a closure of a connected component of $\mathbb{R}^2 - f(S(f))$ represented by $s \in S$ such that for a connected component C_s of f(S(f)) and the small closed tubular neighborhood $N(C_s)$ defined before, C_s' is a connected component of $\partial N(C_s)$. Let M' be a manifold having the structure of an oriented S^1 -bundle over M such that the restriction of the bundle to the total space of the trivial bundle given by $f|_{f^{-1}(C_s')}: f^{-1}(C_s') \to C_s'$ is orientable and trivial.

Then, by a P-operation to f, for any closed manifold M' having the structure of an S^1 -bundle over M, we can obtain a round fold map $f': M' \to \mathbb{R}^2$.

Proof. We prove the first part. For any connected component C of f(S(f)) and the small tubular neighborhood N(C) of C, we have $H^2(f^{-1}(N(C)); \mathbb{Z}) \cong H^2(F_C; \mathbb{Z}) \oplus (H^0(F_C; \mathbb{Z}) \otimes \mathbb{Z}) \cong H^0(F_C; \mathbb{Z}) \otimes \mathbb{Z}$. By an argument similar to that of Theorem 1, the restriction of any S^1 -bundle over M such that the 1st Stiefel-Whitney class vanishes on $i_{s*}(H_1(F_s; \mathbb{Z}/2\mathbb{Z})) \subset H_1(M; \mathbb{Z}/2\mathbb{Z})$ for any $s \in S$ to $f^{-1}(N(C))$ is orientable. Thus, by a discussion similar to that of the proof of Theorem 1, we have only to show that for any connected component C of f(S(f)) and each connected component C' of the boundary $\partial N(C)$ of N(C), the restriction of the bundle over M satisfying the conditions mentioned in the assumption of this theorem to $f^{-1}(C')$ is trivial. Since the bundle $f^{-1}(N(C))$ over C as in Definition 2 is trivial and represented by $C \times F_C$, the restriction of the bundle over M to $f^{-1}(C_0')$ is trivial by the second condition and M is connected, it immediately follows.

We prove the second part. Let C be a connected component of f(S(f)). For the small tubular neighborhood N(C) of C, let C' be a connected component of the boundary of the tubular neighborhood N(C) and let F_C' be a fiber of the bundle given by $f|_{f^{-1}(C')}:f^{-1}(C')\to C'$. We have $H^1(f^{-1}(C');\mathbb{Z}/2\mathbb{Z})\cong H^1(C'\times \mathbb{Z})$ $F_C'; \mathbb{Z}/2\mathbb{Z}) \cong \mathbb{Z}/2\mathbb{Z} \otimes H^1(F_C'; \mathbb{Z}/2\mathbb{Z}) \text{ and } H^1(f^{-1}(N(C)); \mathbb{Z}/2\mathbb{Z}) \cong (H^1(C; \mathbb{Z}/2\mathbb{Z}) \otimes H^1(C; \mathbb{Z}/2\mathbb{Z}))$ $H^0(F_C; \mathbb{Z}/2\mathbb{Z})) \oplus ((H^0(C; \mathbb{Z}/2\mathbb{Z}) \otimes H^1(F_C; \mathbb{Z}/2\mathbb{Z}))) \cong (\mathbb{Z}/2\mathbb{Z} \otimes H^0(F_C; \mathbb{Z}/2\mathbb{Z})) \oplus$ $(\mathbb{Z}/2\mathbb{Z} \otimes H^1(F_C; \mathbb{Z}/2\mathbb{Z}))$. We also have $H^2(f^{-1}(C'); \mathbb{Z}) \cong H^2(C' \times F_C'; \mathbb{Z}) \cong (H^1(C'; \mathbb{Z}) \otimes H^1(F_C'; \mathbb{Z})) \oplus (H^0(C'; \mathbb{Z}) \otimes H^2(F_C'; \mathbb{Z})) \cong (\mathbb{Z} \otimes H^1(F_C'; \mathbb{Z})) \oplus (\mathbb{Z} \otimes H^$ $H^2(F_C';\mathbb{Z})$ and $H^2(f^{-1}(N(C));\mathbb{Z}) \cong H^2(C \times F_C;\mathbb{Z}) \cong H^1(C;\mathbb{Z}) \otimes H^1(F_C;\mathbb{Z}) \cong$ $\mathbb{Z} \otimes H^1(F_C; \mathbb{Z})$. For any connected component C of f(S(f)), the restriction of the bundle over M satisfying the conditions mentioned in the assumption of this theorem to a fiber F_C of the bundle $f^{-1}(N(C))$ over C is orientable by the structures of the groups $H^1(f^{-1}(N(C)); \mathbb{Z}/2\mathbb{Z})$ and $H^1(f^{-1}(C'); \mathbb{Z}/2\mathbb{Z})$ together with the assumption that the $(H_1, \mathbb{Z}/2\mathbb{Z})$ -graph of $f_F = (f, \{F_C\}_{C \in \pi_0(f(S(f)))})$ has a starting set S. Furthermore, the restriction of the bundle over M satisfying the conditions mentioned in the assumption of this theorem to $f^{-1}(N(C))$ is orientable by the assumptions that the restriction of the bundle to the total space of the trivial bundle given by $f|_{f^{-1}(C_s')}: f^{-1}(C_s') \to C_s'$ is orientable and that the manifold M is connected. Thus, by a discussion similar to that of the proof of Theorem 1, for any connected component C of f(S(f)), the restriction of the bundle to $f^{-1}(N(C))$ is trivial. This completes the proof of the desired fact.

We immediately have the following corollary to Theorem 2.

Corollary 1. Let M be a closed and connected manifold of dimension $m \geq 3$ and $f: M \to \mathbb{R}^3$ be a C^∞ L-trivial round fold map such that f(M) is diffeomorphic to D^3 . For any connected component C of f(S(f)), we denote by N(C) a small closed tubular neighborhood of C such that the inverse image has the structures of trivial bundles as in Definition 2 and by F_C the normal fiber of C corresponding to the bundles $f^{-1}(N(C))$ over C. Let F_C satisfy $H^2(F_C; \mathbb{Z}) \cong \{0\}$. Let S be a starting set of the $(H_1, \mathbb{Z}/2\mathbb{Z})$ -graph of $f_F = (f, \{F_C\}_{C \in \pi_0(f(S(f)))})$. Let F_s be the regular fiber of a point in a connected component of $\mathbb{R}^n - f(S(f))$ whose closure is represented as a vertex $s \in S$ of the graph above and let $i_s: F_s \to M$ be the natural inclusion.

Then, by a P-operation to f, for any closed manifold M' having the structure of an S^1 -bundle over M such that the 1st Stiefel-Whitney class vanishes on $i_{s_*}(H_1(F_s; \mathbb{Z}/2\mathbb{Z})) \subset H_1(M; \mathbb{Z}/2\mathbb{Z})$ for any $s \in S$, we can obtain a round fold map $f': M' \to \mathbb{R}^3$.

We review the following propositions, which have been shown in [14] and which are regarded as specific cases of Theorems 1 and 2.

Proposition 3. Let M be a closed manifold of dimension m, $f: M \to \mathbb{R}^n$ be a C^{∞} L-trivial round fold map and $m \ge n \ge 3$. Assume that either of the followings holds.

- (1) $n \ge 4$ holds.
- (2) n = 3 holds, f(M) is diffeomorphic to D^3 and M is connected.

Assume that for any connected component C of f(S(f)), a normal fiber F_C of C corresponding to a bundle satisfies $H^2(F_C; \mathbb{Z}) \cong \{0\}$. Then for any closed manifold having the structure of an orientable S^1 -bundle over M, by a P-operation by S^1 on f, we can obtain a round fold map $f': M' \to \mathbb{R}^n$.

Furthermore, for any connected component C of f(S(f)), let the normal fiber F_C of C satisfy $H^1(F_C; \mathbb{Z}_2) \cong \{0\}$. Then for any closed manifold having the structure of an S^1 -bundle over M, by a P-operation by S^1 on f, we can obtain a C^{∞} L-trivial round fold map $f': M' \to \mathbb{R}^n$.

Proposition 4. Let M be a closed manifold of dimension $m \geq 2$ and $f: M \to \mathbb{R}^2$ be a C^{∞} L-trivial round fold map.

- (1) For any connected component C of f(S(f)), we denote by N(C) a small closed tubular neighborhood of C such that f⁻¹(N(C)) has the structures of trivial bundles as in Definition 3 and by F_C a normal fiber of C corresponding to the bundles f⁻¹(N(C)) over C. Let F_C satisfy H¹(F_C; Z) ≅ H²(F_C; Z) ≅ {0}. Then, by a P-operation by S¹ on f, for any closed manifold having the structure of an S¹-bundle over M such that for any connected component C of f(S(f)), the restriction of the bundle to f⁻¹(∂N(C)) is an orientable bundle, we can obtain a round fold map f': M' → R².
- (2) Furthermore, if f(M) is diffeomorphic to D^2 and M is connected, then by a P-operation by S^1 on f, for any closed manifold having the structure of an S^1 -bundle over M, we can obtain a round fold map $f': M' \to \mathbb{R}^2$.

Example 2. For any pair of positive integers m and n satisfying $m > n \ge 2$, we can consider round fold maps from m-dimensional closed manifolds into \mathbb{R}^n as in Example 1 (3). A special generic map from the m-dimensional homotopy sphere

into \mathbb{R}^n presented in Example 1 (1) are simplest examples of such maps. If for the two integers, m-n>2 holds, then the maps satisfy the assumption of Theorem 1. In this case, the maps satisfy the assumption of Proposition 3 or 4. In this case, the $(H_1; \mathbb{Z}/2\mathbb{Z})$ -graph of $f_F = (f, \{F_C\}_{C \in \pi_0(f(S(f)))})$ is a graph satisfying the followings.

- (1) The set $E_{f_F}^{(H_1,\mathbb{Z}/2\mathbb{Z})} \subset V_{f_F} \times V_{f_F}$ consist of all the pairs of (s_1, s_2) such that the closures of connected components of $\mathbb{R}^n f(S(f))$ represented by s_1 and s_2 intersect.
- (2) The set consisting of just one vertex $s \in V_{f_F}$ representing the closure of the connected component of $\mathbb{R}^n f(S(f))$ including a proper core of f is a starting set of the graph.

For any non-negative integer $k \neq 0, m-n$ and any commutative ring R, the $(H_k; R)$ -graph of $f_F = (f, \{F_C\}_{C \in \pi_0(f(S(f)))})$ and the graph above coincide. The $(H_0; R)$ -graph of $f_F = (f, \{F_C\}_{C \in \pi_0(f(S(f)))})$ satisfies the followings.

- (1) The set $E_{f_F}^{(H_0,R)} \subset V_{f_F} \times V_{f_F}$ consist of all the pairs of (s_1, s_2) such that the closures of connected components of $\mathbb{R}^n f(S(f))$ represented by s_1 and s_2 intersect but a pair (s_1, s_2) where s_1 or s_2 represents the closure of the unbounded connected component of $\mathbb{R}^n f(S(f))$.
- (2) The set consisting of just one vertex $s \in V_{f_F}$ representing the closure of the connected component of $\mathbb{R}^n f(S(f))$ including a proper core of f is a starting set of the graph.

If m-n=1 holds, then the maps do not satisfy the assumption of Propositions 3 or 4, either. However, the maps satisfy the assumptions of Theorems 1 and 2. In this case, for any commutative ring R, the (H_1, R) -graph of $f_F = (f, \{F_C\}_{C \in \pi_0(f(S(f)))})$ is a graph satisfying the followings.

- (1) The set $E_{f_F}^{(H_1,R)} \subset V_{f_F} \times V_{f_F}$ consist of the followings.
 - (a) All the pairs $(s_1, s_2) \in V_{f_F} \times V_{f_F}$ such that the closures of connected components of $\mathbb{R}^n f(S(f))$ represented by s_1 and s_2 intersect and that the closure of the connected component of $\mathbb{R}^n f(S(f))$ represented by s_1 is in the bounded connected component of the complement set of the connected component of $\mathbb{R}^n f(S(f))$ whose closure is represented by s_2 in \mathbb{R}^n .
 - (b) The pair $(s_1, s_2) \in V_{f_F} \times V_{f_F}$ such that s_1 represents the closure of the unbounded connected component of $\mathbb{R}^n f(S(f))$ and that the previous closed set and the closure of the connected component of $\mathbb{R}^n f(S(f))$ represented by s_2 intersect.
- (2) The set consisting of just one vertex $s \in V_{f_F}$ representing the closure of the connected component of $\mathbb{R}^n f(S(f))$ including a proper core of f is a starting set of the graph.

For any non-negative integer $k \neq 1$ and any commutative ring R, the $(H_k; R)$ -graph of $f_F = (f, \{F_C\}_{C \in \pi_0(f(S(f)))})$ is a graph mentined in the case where m - n > 2 holds.

By analogy of the proof of Theorem 1, we have the following theorem.

Theorem 3. Let M be a closed manifold of dimension m and $f: M \to \mathbb{R}^n$ $(m \ge n \ge 2)$ be a C^{∞} L-trivial round fold map. For any connected component C of f(S(f)), we denote by N(C) a small closed tubular neighborhood of C such that the inverse image has the structures of trivial bundles as in Definition 2 and by F_C a normal fiber of C corresponding to the bundles $f^{-1}(N(C))$ over C. Let $H_1(F_C; \mathbb{Z})$ be torsion-free. Let S be a starting set of the (H_2, \mathbb{Z}) -graph of $f_F = (f, \{F_C\}_{C \in \pi_0(f(S(f)))})$. Let F_s be the regular fiber of a point in a connected component of $\mathbb{R}^n - f(S(f))$ whose closure is represented as a vertex $s \in S$ of the graph above and let $i_s: F_s \to M$ be the natural inclusion. Then, we have the followings.

- (1) Let $n \geq 4$ hold. Then, by a P-operation on f, for any closed manifold M' having the structure of an oriented S^1 -bundle over M whose Euler class vanishes on $i_{s*}(H_2(F_s;\mathbb{Z})) \subset H_2(M;\mathbb{Z})$ for any $s \in S$, we can obtain a round fold map $f': M' \to \mathbb{R}^n$.
- (2) Let n=3 hold. Furthermore, for any commutative ring R, the (H_2,\mathbb{Z}) graph of $f_F = (f, \{F_C\}_{C \in \pi_0(f(S(f)))})$ and the (H_0, R) -graph of $f_F = (f, \{F_C\}_{C \in \pi_0(f(S(f)))})$ are start-equivalent on the set S. Then, by a P-operation on f, for any
 closed manifold M' having the structure of an oriented S^1 -bundle over Msuch that the followings hold, we can obtain a round fold map $f': M' \to \mathbb{R}^n$.
 - (a) The Euler class vanishes on $i_{s*}(H_2(F_s; \mathbb{Z})) \subset H_2(M; \mathbb{Z})$ for any $s \in S$.
 - (b) For $s \in S$, let C_s' be a 2-dimensional sphere in a closure of a connected component of $\mathbb{R}^n f(S(f))$ represented by $s \in S$ such that for a connected component C_s of f(S(f)) and the small closed tubular neighborhood $N(C_s)$ defined before, C_s' is a connected component of $\partial N(C)$. The restriction of the bundle to the image of a section of the trivial bundle given by $f|_{f^{-1}(C_s')}: f^{-1}(C_s') \to C_s'$ is trivial.
- (3) Let n = 2 hold. Furthermore, we also assume that for any commutative ring R, the (H₂; Z)-graph of f_F = (f, {F_C}_{C∈π₀(f(S(f)))}) and the (H₁; Z)-graph of f_F = (f, {F_C}_{C∈π₀(f(S(f)))}) are start-equivalent on the set S. For s ∈ S, let C_s' be a circle in a closure of a connected component of R² f(S(f)) represented by s ∈ S such that for a connected component C_s of f(S(f)) and the small closed tubular neighborhood N(C_s) defined before, C_s' is a connected component of ∂N(C_s). Let M' be a manifold having the structure of an oriented S¹-bundle over M such that the restriction of the bundle to the total space of the trivial bundle given by f|_{f-1(C_s')}: f⁻¹(C_s') → C_s' is orientable and trivial.

Then, by a P-operation on f, we can obtain a round fold map $f': M' \to \mathbb{R}^n$.

Proof. Let C_0 be a connected component of f(S(f)) which is in the closures of two connected component of $\mathbb{R}^n - f(S(f))$ represented by two vertices $s_1, s_2 \in V_{f_F}$ and let $(s_1, s_2) \in E_{f_F}$. We use C_j and $\partial_j F_{C_0}$ as in the proof of Theorem 1. The homomorphism from $H_2(f^{-1}(C_1); \mathbb{Z}) \cong H_2(C_1 \times \partial_1 F_{C_0}; \mathbb{Z}) \cong (H_0(\partial_1 F_{C_0}; \mathbb{Z}) \otimes H_2(C_0; \mathbb{Z})) \oplus (H_1(C_0; \mathbb{Z}) \otimes H_1(\partial_1 F_{C_0}; \mathbb{Z})) \oplus H_2(\partial_1 F_{C_0}; \mathbb{Z})$ into $H_2(f^{-1}(N(C)); \mathbb{Z}) \cong (H_0(F_C; \mathbb{Z}) \otimes H_2(C; \mathbb{Z})) \oplus (H_1(C; \mathbb{Z}) \otimes H_1(F_C; \mathbb{Z})) \oplus H_2(F_C; \mathbb{Z})$ induced by the natural inclusion is assumed to be surjective in the first case, since the homomorphism from $H_2(\partial_1 F_{C_0}; \mathbb{Z})$ into $H_2(F_C; \mathbb{Z})$ induced by the natural inclusion is assumed to be surjective. In the second case, the homomorphism from $H_k(\partial_1 F_{C_0}; \mathbb{Z})$ into $H_k(F_C; \mathbb{Z})$ induced by the natural inclusion is assumed to be surjective for k = 0, 2

and as a result, the homomorphism above is surjective. In the last case, the homomorphism from $H_k(\partial_1 F_{C_0}; \mathbb{Z})$ into $H_k(F_C; \mathbb{Z})$ induced by the natural inclusion is assumed to be surjective for k=1,2 and as a result, the homomorphism above is surjective. Thus, the kernel of the homomorphism from $H^2(f^{-1}(N(C)); \mathbb{Z})$ into $H^2(f^{-1}(C_1); \mathbb{Z})$ induced by the natural inclusion is $\{0\}$ in all the three cases by virtue of the universal coefficient theorem together with the assumption that the group $H_1(F_C; \mathbb{Z})$ is torsion-free for any connected component C of f(S(f)). So, if the Euler class of the restriction of an oriented S^1 -bundle over M to $f^{-1}(C_1)$ vanishes, then the Euler classes of the restrictions of the bundle over M to $f^{-1}(N(C_0))$ vanish and the obtained bundles are trivial bundles.

By arguments similar to ones in the proofs of Theorem 1, for each case, for any connected component C of f(S(f)), the restriction of any oriented S^1 -bundle over M satisfying the given condition to $f^{-1}(N(C))$ is a trivial bundle. This completes the proof of the statement for all the three cases.

By analogy of the proof of Theorem 2, we have the following theorem.

Theorem 4. Let M be a closed and connected manifold of dimension $m \geq 3$ and $f: M \to \mathbb{R}^3$ be a C^{∞} L-trivial round fold map. For any connected component C of f(S(f)), we denote by N(C) a small closed tubular neighborhood of C such that the inverse image has the structures of trivial bundles as in Definition 2 and by F_C a normal fiber of C corresponding to the bundles $f^{-1}(N(C))$ over C. Let $H_1(F_C; \mathbb{Z})$ be torsion-free. Let S be a starting set of the (H_2, \mathbb{Z}) -graph of $f_F = (f, \{F_C\}_{C \in \pi_0(f(S(f)))})$. Let F_S be the regular fiber of a point in a connected component of $\mathbb{R}^n - f(S(f))$ whose closure is represented as a vertex $s \in S$ of the graph above and let $i_s: F_s \to M$ be the natural inclusion. Then, by a P-operation to f, for any closed manifold M' having the structure of an oriented S^1 -bundle over M such that the followings hold, we can obtain a round fold map $f': M' \to \mathbb{R}^3$.

- (1) The Euler class vanishes on $i_{s*}(H_2(F_s;\mathbb{Z})) \subset H_2(M;\mathbb{Z})$ for any $s \in S$.
- (2) For a 2-dimensional sphere C₀' in a connected component of (R³-f(S(f))) ∩ f(M) such that for a connected component C₀ of f(S(f)) and the small closed tubular neighborhood defined before, C₀' is a connected component of ∂N(C₀), the restriction of the bundle to the image of a section of the trivial bundle given by f|_{f⁻¹(C₀')}: f⁻¹(C₀') → C₀' is trivial.

Proof. Let C be a connected component of f(S(f)). For the small tubular neighborhood N(C) of C, let C' be a connected component of the boundary of the tubular neighborhood N(C) and let $F_{C'}$ be a fiber of the bundle given by $f|_{f^{-1}(C')}: f^{-1}(C') \to C'$. We have $H^2(f^{-1}(C'); \mathbb{Z}) \cong H^2(C' \times F_{C'}; \mathbb{Z}) \cong (H^2(C'; \mathbb{Z}) \otimes H^0(F_{C'}; \mathbb{Z})) \oplus (H^0(C'; \mathbb{Z}) \otimes H^2(F_{C'}; \mathbb{Z})) \cong (\mathbb{Z} \otimes H^0(F_{C'}; \mathbb{Z})) \oplus (\mathbb{Z} \otimes H^2(F_{C'}; \mathbb{Z}))$ and $H^2(f^{-1}(N(C)); \mathbb{Z}) \cong H^2(C \times F_{C}; \mathbb{Z}) \cong (H^2(C; \mathbb{Z}) \otimes H^0(F_{C}; \mathbb{Z})) \oplus (H^0(C; \mathbb{Z}) \otimes H^2(F_{C}; \mathbb{Z})) \cong \mathbb{Z} \otimes H^0(F_{C}; \mathbb{Z}) \oplus (\mathbb{Z} \otimes H^2(F_{C}; \mathbb{Z}))$. For any connected component C of f(S(f)), the restriction of an oriented S^1 -bundle over M satisfying the conditions mentioned in the assumption of this theorem to a fiber F_C of the bundle $f^{-1}(N(C))$ over C is trivial by the structures of the groups $H^2(f^{-1}(N(C)); \mathbb{Z})$ and $H^2(f^{-1}(C'); \mathbb{Z})$ together with the assumption that the (H_2, \mathbb{Z}) -graph of $f_F = (f, \{F_C\}_{C \in \pi_0(f(S(f)))})$ has a starting set S. Furthermore, the restriction of the bundle over M satisfying the conditions mentioned in the assumption of this theorem to $f^{-1}(N(C))$ is trivial by the assumptions that the restriction of the bundle to the total space of the trivial bundle given by $f|_{f^{-1}(C_0')}: f^{-1}(C_0') \to C_0'$ is trivial

and that the manifold M is connected. Thus, for any connected component C of f(S(f)), the restriction of the bundle to $f^{-1}(N(C))$ is trivial. This completes the proof of the desired fact.

We immediately have the following corollary to Theorem 4.

Corollary 2. Let M be a closed and connected manifold of dimension $m \geq 3$ and $f: M \to \mathbb{R}^3$ be a C^{∞} L-trivial round fold map such that f(M) is diffeomorphic to D^3 . For any connected component C of f(S(f)), we denote by N(C) a small closed tubular neighborhood of C such that the inverse image has the structures of trivial bundles as in Definition 2 and by F_C the normal fiber of C corresponding to the bundles $f^{-1}(N(C))$ over C. Let $H_1(F_C; \mathbb{Z})$ be torsion-free. Let S be a starting set of the (H_2, \mathbb{Z}) -graph of $f_F = (f, \{F_C\}_{C \in \pi_0(f(S(f)))})$. Let F_s be the regular fiber of a point in a connected component of $\mathbb{R}^n - f(S(f))$ whose closure is represented as a vertex $s \in S$ of the graph above and let $i_s : F_s \to M$ be the natural inclusion. Then, by a P-operation on f, for any closed manifold M' having the structure of an oriented S^1 -bundle over M such that the Euler class vanishes on $i_{s*}(H_2(F_s; \mathbb{Z})) \subset H_2(M; \mathbb{Z})$ for any $s \in S$, we can obtain a round fold map $f': M' \to \mathbb{R}^3$.

Example 3. A round fold map f presented in the beginning of Exmaple 2 satisfies the assumption of Theorems 3 and 4. In the case where for the two integers, m-n=2 holds, the map satisfies the assumptions of Theorems 3 and 4 but does not satisfy the assumption of Theorem 1 or that of Theorem 2, either. In this case, the map does not satisfy the assumption of Proposition 3 or Proposition 4, either.

4. Algebraic and differential topological properties of round fold maps given by P-operations by S^1 and their source manifolds

We explicitly apply P-operations by S^1 to some round fold maps satisfying the assumption of Propositions, Theorems and Corollaries in section 3. In this section, we use the following terminologies and notations.

For a freely generated commutative group A of rank $r \geq 1$, $x \in A$ is said to be *primitive* if for the subgroup generated by x (we denote the subgroup by < x >), the quotient group A/< x > is a freely generated commutative group of rank r-1. The diffeomorphism type of a manifold having the structure of a non-trivial S^k -bundle (k = 2, 3) over S^2 is unique. We denote the manifold by $S^k \tilde{\times} S^2$.

4.1. Applications of Propositions and Theorems in section 3 to round fold maps in Example 1 (1) and (3). We may apply Propositions, Theorems and Corollaries in section 3 to obtain a new round fold map by a P-operation by S^1 to a round fold map $f: M \to \mathbb{R}^n$ in Example 1 (1) (see also [12]). In fact, by any P-operation by S^1 to the map f, we obtain the product of S^1 and the original source manifold as the resulting source manifold if the dimension of the original source manifold is larger than 2. Moreover, by a P-operation by S^1 to such a map from the 2-dimensional standard sphere S^2 into the plane, we obtain a manifold having the structure of an S^1 -bundle over S^2 as the resulting source manifold and any manifold having the structure of an S^1 -bundle over S^2 is realized as the resulting source manifold by a P-operation by S^1 to every such map from S^2 into the plane. If for the round fold map $f: M \to \mathbb{R}^n$, the dimension of the source manifold M and n coincide, then the obtained maps are maps as in Example 1 (2)

with $F := S^1$.

We consider P-operations by S^1 to round fold maps $f: M \to \mathbb{R}^n \ (n \ge 2)$ in Example 1 (3).

First, in the case where $n \geq 3$ and $m \geq 2n$ hold in this example, the manifold having the structure of an S^1 -bundle over the original source manifold M is always the product of S^1 and M. By any P-operation by S^1 , we always obtain a source manifold diffeomorphic to the product of the two manifolds.

In the case where n=2 and m>2n=4 hold in this example, on any manifold having the structure of an S^1 -bundle over the original source manifold M, we can construct a round fold map by a P-operation by S^1 to the original map f by Propositions 3 and 4 (we can apply Theorems 1, 2, 3 and 4 too). The author studied the diffeomorphism types of the resulting source manifolds of the resulting round fold maps in [14].

In the case where (m, n) = (4, 2) holds in this example, we cannot apply Proposition 4. Moreover, we cannot apply Theorem 1 or 2, either. On the other hand, we can apply Theorems 3 and 4. As a result on the diffeomorphism types of the source manifolds of round fold maps obtained by P-operations by S^1 to the original round fold map f, we have the following theorem.

Theorem 5. Let M be a connected sum of $l \in \mathbb{N}$ closed manifolds having the structure of an S^2 -bundles over S^2 . By a P-operation by S^1 to the map in Example 3, for any cohomology class $\alpha \in H^2(M; \mathbb{Z})$ vanishing on all the cycles corresponding to the fibers of the bundles, we can construct a round fold map from any closed manifold $M(\alpha)$ having the structure of an oriented S^1 -bundle over M whose Euler class is α into \mathbb{R}^2 . Furthermore, we have the followings.

- (1) $\alpha \in H^2(M; \mathbb{Z})$ is chosen to be primitive and if it is primitive, then $M(\alpha)$ is simply-connected.
- (2) If $M(\alpha)$ is simply-connected, then $\alpha \in H^2(M; \mathbb{Z})$ must be primitive and $M(\alpha)$ is diffeomorphic to a connected sum of 2l-2 copies of $S^2 \times S^3$ (we replace this by S^5 in the case where l=1 holds) and a closed manifold having the structure of an S^3 -bundle over S^2 .
- (3) If M is represented as a connected sum of finite copies of $S^2 \times S^2$ and $M(\alpha)$ is simply-connected, then $M(\alpha)$ is represented as a connected sum of 2l-1 copies of $S^2 \times S^3$.
- (4) If M is represented as a connected sum of more than one closed manifolds having the structures of S²-bundles over S² (l > 1 must hold) and not represented as a connected sum of finite copies of S² × S², then for any closed manifold represented as a connected sum of 2l − 2 copies of S² × S³ and a closed manifold having the structure of an S³-bundle over S², we can choose α ∈ H²(M; Z) so that M(α) is diffeomorphic to the manifold.
- (5) If M is a closed manifold having the structure of an S^2 -bundle over S^2 and $M(\alpha)$ is simply-connected, then $M(\alpha)$ is diffeomorphic to $S^2 \times S^3$.

For the proof of Theorem 5, we need the following proposition, which is a main theorem of [3]. See also [1], in which Barden classified closed and simply-connected 5-dimensional manifolds completely.

Proposition 5 ([3]). Let X be a closed and simply-connected manifold which may not be smooth of dimension 4. Let l > 1 be an integer and let $H^2(X; \mathbb{Z}) \cong \mathbb{Z}^l$ hold.

- (1) For any primitive element $\alpha \in H^2(X; \mathbb{Z})$, let $X(\alpha)$ be a manifold having the structure of an oriented S^1 -bundle whose Euler class is α . Then for any α , $X(\alpha)$ is simply-connected and have a differentiable structure and one of the followings holds.
 - (a) $X(\alpha)$ is diffeomorphic to a connected sum of l-1 copies of $S^2 \times S^3$
 - (b) $X(\alpha)$ is diffeomorphic to a connected sum of l-2 copies of $S^2 \times S^3$ (we replace this by S^5 in the case where l=2 holds) and $S^3 \tilde{\times} S^2$.
- (2) Let X be smooth manifold and let $\alpha \in H^2(X; \mathbb{Z})$ be primitive. A manifold $X(\alpha)$ having the structure of an oriented S^1 -bundle whose Euler class is α is not diffeomorphic to a connected sum of l-1 copies of $S^2 \times S^3$ if and only if the followings hold.
 - (a) The 2nd Stiefel-Whitney class β of the tangent bundle TX over X does not vanish.
 - (b) α is not equal to β mod 2.

Example 4. Let X be a connected sum of a finite number of closed smooth manifolds having the structures of S^2 -bundles over S^2 .

Assume that all the bundles appeared in the connected sum are trivial. Then the 2nd Stiefel-Whitney class β of the tangent bundle TX over X vanishes and $X(\alpha)$ in Proposition 5 is diffeomorphic to a connected sum of finite copies of $S^2 \times S^3$ by the proposition.

Assume that there exist a bundle which is not trivial and a trivial bundle in the connected sum. Then the 2nd Stiefel-Whitney class β of the tangent bundle TX over X vanishes on cycles corresponding to fibers of all the S^2 -bundles in this family. On the other hand, the class does not vanish on the cycle corresponding to the base space of an S^2 -bundle in the family which is not trivial. We can take $\alpha \in H^2(X; \mathbb{Z})$ so that α vanishes on cycles corresponding to fibers of all the S^2 -bundles appeared in the connected sum and that the class does not vanish only on the cycle corresponding to the base space of just one trivial S^2 -bundle in the family. In this case, α is not equal to β mod 2. Then $X(\alpha)$ in Proposition 5 is not diffeomorphic to a connected sum of finite copies of $S^2 \times S^3$ and is diffeomorphic to a connected sum of finite copies of $S^2 \times S^3$ by the proposition. We can also take $\alpha \in H^2(X; \mathbb{Z})$ so that α is equal to β mod 2. Then, $X(\alpha)$ in Proposition 5 is diffeomorphic to a connected sum of finite copies of $S^2 \times S^3$.

Proof of Theorem 5. A round fold map into \mathbb{R}^2 as in Example 1 (3) exists on M. We can apply Theorem 2 to this map. This completes the proof of the former part. We prove five statements in the latter part.

We can choose $\alpha \in H^2(M; \mathbb{Z})$ so that it vanishes on all the cycles corresponding to the fibers of the C^{∞} S^2 -bundles over S^2 and on all but one cycles corresponding to the base spaces of the C^{∞} S^2 -bundles over S^2 and that the value is 1 on the cycle corresponding to the base space of the remained C^{∞} S^2 -bundle over S^2 . Then α is primitive and we obtain a round fold map from $M(\alpha)$ into \mathbb{R}^2 by a P-operation by S^1 . By Proposition 5, we have the first statement.

For the proof of the second statement, we review and apply the method used in the proof of Lemma 3 of [3], which is a key lemma to Theorem 2 of [3] or Proposition 5 of the present paper.

From the homotopy sequence of the bundle $M(\alpha)$ over M, $\pi_1(M(\alpha))$ is cyclic. It follows that $\pi_1(M(\alpha)) \cong H_1(M(\alpha); \mathbb{Z})$. Substituting $H^k(M; \mathbb{Z}) \cong \{0\}$ for any odd integer k and any integer k not smaller than 5 in the Gysin sequence of the bundle

 $M(\alpha)$ over M yields the following two exact sequences where $p:M(\alpha)\to M$ is the projection of the bundle and where $\bigcup \alpha$ denotes the operation of taking cup products with α .

$$\{0\} \qquad \xrightarrow{p^*} H^3(M(\alpha); \mathbb{Z}) \longrightarrow H^2(M; \mathbb{Z}) \xrightarrow{\bigcup \alpha} H^4(M; \mathbb{Z}) \cong \mathbb{Z} \xrightarrow{p^*} H^4(M(\alpha); \mathbb{Z}) \longrightarrow \{0\}$$

$$\{0\} \xrightarrow{p^*} H^5(M(\alpha); \mathbb{Z}) \longrightarrow H^4(M; \mathbb{Z}) \cong \mathbb{Z} \xrightarrow{\bigcup \alpha} \{0\}$$

By the second sequence, we have $H^5(M(\alpha); \mathbb{Z}) \cong \mathbb{Z}$. The homomorphism in the first sequence $\bigcup \alpha: H^2(M; \mathbb{Z}) \to H^4(M; \mathbb{Z})$ is surjective if and only if α is primitive. Furthermore, $\bigcup \alpha: H^2(M; \mathbb{Z}) \to H^4(M; \mathbb{Z})$ is surjective if and only if $H^4(M(\alpha); \mathbb{Z}) \cong H_1(M(\alpha); \mathbb{Z}) \cong \pi_1(M(\alpha)) \cong \{0\}$ holds (we can apply Poincare duality theorem since $H^5(M(\alpha); \mathbb{Z}) \cong \mathbb{Z}$ holds). Furthermore, by the first sequence, $H^3(M(\alpha); \mathbb{Z}) \cong \mathbb{Z}^{2l-1}$ holds if α is primitive. By virtue of Poincare duality theorem, $H^3(M(\alpha); \mathbb{Z}) \cong H_2(M(\alpha); \mathbb{Z})$ always holds. Thus, by the theory of [1] or Proposition 5, the second statement of the five statements holds.

By the second statement of the five statements and Example 4 (or Proposition 5), the third and fourth statements of the five statements hold.

The 2nd Stiefel-Whitney class of the tangent bundle of $S^2 \tilde{\times} S^2$ vanishes on the cycle corresponding to the fiber and does not vanish on one corresponding to the base space. So, if in the situation of the last statement of the five statements, M is diffeomorphic to $S^2 \tilde{\times} S^2$ and $M(\alpha)$ is simply-connected, then α must be primitive and as a result it equals the 2nd Stiefel-Whitney class of $S^2 \tilde{\times} S^2$ mod 2. Furthermore, by Proposition 5, $M(\alpha)$ is diffeomorphic to $S^2 \times S^3$ under this condition and the same fact follows by the same proposition if $M = S^2 \times S^2$. Thus this completes the proof of the last one of the five statements.

Note that also in [10], [11] and [13], round fold maps from the simply-connected 5-dimensional manifolds represented as connected sums of manifolds having the structures of S^3 -bundles over S^2 into the plane are obtained. Regular fibers of these maps are disjoint unions of finite copies of S^3 . Theorem 5 shown above gives new explicit examples of round fold maps from such simply-connected 5-dimensional manifolds into the plane.

- 4.2. Applications of Propositions, Theorems and Corollaries mentioned in section 3 to round fold maps in Example 1 (4). We consider a round fold map f from a closed m-dimensional manifold M into \mathbb{R}^n ($m \ge n \ge 2$) in Example 1 (4).
- 4.2.1. The case where $n \geq 4$ holds. Let $n \geq 4$. Then, for any manifold having the structure of an S^1 -bundle over the manifold M, we can obtain a new round fold map by a P-operation by S^1 on the map f. If $m-n \geq 2$, then the resulting manifold is always diffeomorphic to the product of S^1 and the original source manifold M. If m-n=1, then the resulting manifold is homeomorphic to the product of S^{n-1} and a 3-dimensional manifold having the structure of an S^1 -bundle over S^2 and conversely, such a manifold is realized as the resulting source manifold of a round fold map obtained by P-operation by S^1 on the map f. If m=n, then the resulting manifold is homeomorphic to the product of S^{n-1} and the 2-dimensional torus or

the Klein Bottle and conversely, these two manifolds are realized as the resulting source manifolds of round fold maps obtained by P-operations by S^1 to the original round fold map f.

4.2.2. The case where n=3 holds. Let n=3. In this case, for any manifold having the structure of an S^1 -bundle over the source manifold M, the restriction of the bundle to the inverse image of any 2-dimensional sphere smoothly embedded in the interior of the image of the map is orientable. If the bundle obtained by the restriction is oriented and the Euler class vanishes, then, from Theorem 1 or Theorem 2, on the manifold having the structure of an S^1 -bundle over M, we obtain a round fold map by a P-operation by S^1 to the original map f.

We consider the resulting source manifold of the round fold map given by a P-operation in the case where n=3 holds. If $m-n=m-3\geq 2$ or $m\geq 5$ holds, then the resulting manifold must be diffeomorphic to the product of S^1 and the original source manifold. If m-n=m-3=1 or m=4 holds, then the resulting manifold must be diffeomorphic to the product of S^2 and a 3-dimensional manifold having the structure of an S^1 -bundle over S^2 and conversely, we obtain any such manifold as the source manifold of the resulting round fold map given by a P-operation by S^1 to the original round fold map f. If m=n=3 holds, then the resulting manifold must be diffeomorphic to the product of S^2 and the 2-dimensional torus or the Klein Bottle and conversely, these manifolds are obtained by P-operations by S^1 to the original map f as the source manifolds of resulting round fold maps.

4.2.3. The case where n=2 holds. Let n=2. In this case, for a manifold having the structure of an S^1 -bundle over the source manifold of such a round fold map, if the restriction of the bundle to the inverse image of any circle smoothly embedded in the regular value set of the map is orientable, then, by Theorem 3 or 4, on the original manifold having the structure of an S^1 -bundle, we obtain a round fold map by a P-operation by S^1 to the original round fold map.

We consider the resulting source manifolds given by these P-operations in the case where n=2 holds. If $m-n=m-2\geq 2$ or $m\geq 4$ holds, then the resulting manifold must be diffeomorphic to the product of S^1 and the original source manifold. If m-n=m-2=1 or m=3 holds, then the resulting manifold must be diffeomorphic to the product of S^1 and a 3-dimensional manifold having the structure of an S^1 -bundle over S^2 and conversely, any such manifold is obtained by a P-operation by S^1 to the original round fold map.

Last, we consider the case where (m,n)=(2,2) holds. On any manifold having the structure of an oriented S^1 -bundle over the oriented torus $S^1\times S^1$, we obtain a round fold map into \mathbb{R}^2 by a P-operation by S^1 to the original map. The resulting source manifolds are diffeomorphic (homeomorphic) if and only if the absolute values of the Euler numbers of the bundles coincide. Of course, for any integer k, we can obtain an oriented S^1 -bundle whose Euler number is k in this case. Note that by P-operations by S^1 to the original map, we obtain a family $\{M_k\}_{k\in\mathbb{Z}}$ of 3-dimensional oriented closed and connected manifolds and a family of round fold maps $\{f_k: M_k \to \mathbb{R}^2\}_{k\in\mathbb{Z}}$ such that the followings hold.

- (1) M_k has the structure of an oriented S^1 -bundle over the oriented 2-dimensional torus $S^1 \times S^1$ with Euler number $k \in \mathbb{Z}$.
- (2) f_k is C^{∞} L-trivial.
- (3) The inverse image of the axis of f_k is the 2-dimensional torus $S^1 \times S^1$.

Furthermore, f_k is not C^{∞} S-trivial for $k \neq 0$, which follows from Proposition 2 (4).

We can also obtain a round fold map from the product of S^{n-1} and the Klein Bottle by a P-operation by S^1 .

4.3. Further applications.

- 4.3.1. Case 1. Let $m, n \in \mathbb{N}$ and $m > n \ge 2$. Let M be an m-dimensional closed manifold admitting a round fold map $f: M \to \mathbb{R}^n$ such that the followings hold.
 - (1) f is C^{∞} L-trivial.
 - (2) The image f(M) is diffeomorphic to $S^{n-1} \times [-1, 1]$.
 - (3) The inverse image of an axis of the map f is homeomorphic to $S^1 \times S^{m-n}$
 - (4) S(f) consists of four connected components.
 - (5) The regular fiber of a point in a connected component of the set \mathbb{R}^n f(S(f)) consists of the disjoint union of two spheres.

We easily obtain such a map by a trivial spinning construction mentioned before Example 1. The source manifold is diffeomorphic to the product of S^{n-1} and a manifold homeomorphic to $S^1 \times S^{m-n}$. Moreover, for each connected component C of f(S(f)), there exists a small closed tubular neighborhood N(C) such that $f^{-1}(N(C))$ has the structures of isomorphic trivial bundles whose fibers are diffeomorphic to the standard closed disc D^{m-n+1} or homeomorphic to the standard (m-n+1)-dimensional sphere S^{m-n+1} with the interior of a union of disjoint three (m-n+1)-dimensional standard closed discs removed as in Definition 2.

We apply Propositions and Theorems introduced in section 3.

Let n > 4. In the case where m - n > 3 holds, any manifold having the structure of an orientable S^1 -bundle over M is diffeomorphic to $M \times S^1$ and there exists a manifold having the structure of an S^1 -bundle over M diffeomorphic to the product of S^{n-1} and a manifold homeomorphic to the product of the Klein Bottle and S^{m-n} too. In this case, on any manifold having the structure of an S^1 -bundle over M, we obtain a round fold map by a P-operation by S^1 to the map f. In the case where m-n=2 holds, on any manifold having the structure of an orientable S^1 -bundle over M such that the restriction of the bundle to a regular fiber of the map f with two connected components is trivial, we can construct a new round fold map by a P-operation by S^1 to the map f and moreover, the bundle is a trivial S^1 bundle. In this case, there exists a manifold having the structure of an S^1 -bundle over M diffeomorphic to the product of S^{n-1} and a manifold homeomorphic to the product of the Klein Bottle and S^2 too and on the manifold, we can construct a new round fold map similarly. In the case where m-n=1 holds, any manifold having the structure of an orientable S^1 -bundle over M is diffeomorphic to the product of M and a manifold having the structure of an orientable S^1 -bundle over the 2-dimensional torus and on any manifold having the structure of an orientable S^1 -bundle over M, we can construct a new round fold map by a P-operation by S^1 to the map f. There exists a manifold having the structure of an S^1 -bundle over M diffeomorphic to the product of S^{n-1} and a manifold diffeomorphic to the product of the Klein Bottle and S^1 and on the manifold, we can construct a new round fold map similarly.

Let n=3. In the case where $m-n=m-3\geq 3$ $(m\geq 6)$ holds, on any manifold having the structure of an orientable S^1 -bundle over M such that the restriction

of the bundle to any 2-dimensional sphere in the set $\mathbb{R}^3 - f(S(f))$ is trivial, we can construct a new round fold map by a P-operation by S^1 to the map f and moreover, the bundle is a trivial S^1 -bundle. There exists a manifold having the structure of an S^1 -bundle over M diffeomorphic to the product of $S^{n-1} = S^2$ and a manifold homeomorphic to the product of the Klein Bottle and S^{m-n} and on the manifold, we can construct a new round fold map similarly. In the case where m-n=m-3=2 (m=5) holds, on any manifold having the structure of an orientable S^1 -bundle over M such that the restriction of the bundle to any 2-dimensional sphere in the set $\mathbb{R}^n - f(S(f))$ and a regular fiber of the map f with two connected components are trivial, we can construct a new round fold map by a P-operation by S^1 to the map f and moreover, the bundle is a trivial S^1 -bundle. There exists a manifold having the structure of an S^1 -bundle over M diffeomorphic to the product of $S^{n-1} = S^2$ and a manifold homeomorphic to the product of the Klein Bottle and $S^{m-n} = S^3$ and on the manifold, we can construct a new round fold map similarly. In the case where m-n=m-3=1 (m=4) holds, any manifold having the structure of an orientable S^1 -bundle over M such that the restriction of the bundle to any 2-dimensional sphere in the set $\mathbb{R}^3 - f(S(f))$ is trivial is diffeomorphic to the product of M and a manifold having the structure of an orientable S^1 -bundle over the 2-dimensional torus and on any such manifold, we can construct a new round fold map by a P-operation by S^1 to the map f. There exists a manifold having the structure of an S^1 -bundle over M diffeomorphic to the product of $S^{n-1} = S^2$ and a manifold diffeomorphic to the product of the Klein Bottle and S^1 and on the manifold, we can construct a new round fold map similarly.

Let n=2. For simplicity, we only consider the case where $m-n=m-2\geq 3$ $(m\geq 5)$ holds.

Any manifold having the structure of an orientable S^1 -bundle over M is homeomorphic to the product of a 3-dimensional manifold having the structure of an orientable S^1 -bundle over the 2-dimensional torus and S^{m-n} and for each product of a 3-dimensional manifold having the structure of an orientable S^1 -bundle over the 2-dimensional torus and $S^{m-n} = S^{m-2}$, there exists a manifold having the structure of an orientable S^1 -bundle over M homeomorphic to the product. Furthermore, by Proposition 2 (4), the 1st (co)homology group of a 3-dimensional manifold having the structure of an oriented S^1 -bundle over the 2-dimensional torus with Euler number k is isomorphic to $\mathbb{Z} \oplus \mathbb{Z} \oplus \mathbb{Z}/|k|\mathbb{Z}$. From the discussion here, we have the following theorem.

Theorem 6. By P-operations by S^1 to the map $f: M \to \mathbb{R}^2$ here, we have a family of (m+1)-dimensional oriented manifolds $\{M_k\}_{k\in\mathbb{Z}}$ and round fold maps $\{f_k: M_k \to \mathbb{R}^2\}_{k\in\mathbb{Z}}$. Furthermore, the followings hold.

- (1) By any P-operation by S^1 to the map f such that the resulting S^1 -bundle over M is orientable, the resulting source manifold of the resulting map is diffeomorphic to a manifold belonging to the family $\{M_k\}$.
- (2) M_k is homeomorphic to the product of a 3-dimensional manifold having the structure of an oriented S^1 -bundle over the 2-dimensional oriented torus $S^1 \times S^1$ with Euler number k and $S^{m-n} = S^{m-2}$
- (3) The inverse image of an axis of f_k is homeomorphic to $S^1 \times S^{m-2} \times S^1$
- (4) f_k is C^{∞} L-trivial.
- (5) f_k is not C^{∞} S-trivial if k is not 0.

In this case, there exists a manifold having the structure of an S^1 -bundle over M diffeomorphic to the product of $S^{n-1} = S^1$ and a manifold homeomorphic to the product of the Klein Bottle and $S^{m-n} = S^{m-2}$ too and on this manifold, we can construct a new round fold map similarly.

4.3.2. Case 2. Let $m, n \in \mathbb{N}$ and $m > n \ge 2$. Let M be an m-dimensional closed manifold admitting a round fold map $f: M \to \mathbb{R}^n$ such that the followings hold.

- (1) f is C^{∞} L-trivial.
- (2) The image f(M) is diffeomorphic to D^n .
- (3) The inverse image of an axis of the map f is homeomorphic to $S^1 \times S^{m-n}$ with the interior of an (m-n+1)-dimensional standard closed disc removed.
- (4) S(f) consists of three connected components.
- (5) The regular fiber of a point in a connected component of the set \mathbb{R}^n f(S(f)) consists of the disjoint union of two spheres.

We easily obtain such a map by a trivial spinning construction mentioned before Example 1. For a proper core P of f, the manifold $f^{-1}(\mathbb{R}^n-\mathrm{Int}P)$ is diffeomorphic to the product of S^{n-1} and a manifold homeomorphic to $S^1\times S^{m-n}$ with the interior of an (m-n+1)-dimensional standard closed disc removed. Moreover, for each connected component C of f(S(f)), there exists a small closed tubular neighborhood N(C) such that $f^{-1}(N(C))$ has the structures of isomorphic trivial bundles whose fibers are diffeomorphic to the standard closed disc D^{m-n+1} or homeomorphic to the standard (m-n+1)-dimensional sphere S^{m-n+1} with the interior of a union of disjoint three (m-n+1)-dimensional standard closed discs removed as in Definition 2.

We consider the case where $m - n \ge 3$ holds for simplicity.

Let $n \geq 3$ hold. In this case, the 2nd homology groups $H_2(M; \mathbb{Z})$ and $H_2(f^{-1}(\mathbb{R}^n - \text{Int}P); \mathbb{Z})$ and the 2nd cohomology groups $H^2(M; \mathbb{Z})$ and $H^2(f^{-1}(\mathbb{R}^n - \text{Int}P); \mathbb{Z})$ are zero and any orientable S^1 -bundle over M is trivial. We can construct a round fold map from $M \times S^1$ into \mathbb{R}^n by a P-operation by S^1 to f. In this case, according to a discussion similar to that of Case 1, there exists a manifold having the structure of an S^1 -bundle over M which is not orientable admitting a round fold map into \mathbb{R}^n obtained by a P-operation by S^1 to f.

Let n=2 hold. In this case, the 2nd homology groups $H_2(M;\mathbb{Z})$ and $H_2(f^{-1}(\mathbb{R}^n-\operatorname{Int}P);\mathbb{Z})$ are $\operatorname{Int}P(f^{-1}(\mathbb{R}^n-\operatorname{Int}P);\mathbb{Z})$ are all \mathbb{Z} . Moreover, the homomorphisms between the 2nd homology and cohomology groups induced by the inclusions of $f^{-1}(\mathbb{R}^n-\operatorname{Int}P)$ into M are isomorphisms. Let $v\in H^2(M;\mathbb{Z})\cong \mathbb{Z}$ be a generator of the group and $v'\in H_2(f^{-1}(\mathbb{R}^n-\operatorname{Int}P);\mathbb{Z})$ be the generator of the group defined as the inverse image of v of the isomorphism induced by the natural inclusion. Let M_k be an oriented manifold having the structure of an oriented S^1 -bundle over M whose Euler class is kv. The restriction of the bundle M_k to $f^{-1}(\mathbb{R}^n-\operatorname{Int}P)$ is an oriented S^1 -bundle whose Euler class is kv'. Since we have $H_1(f^{-1}(\mathbb{R}^n-\operatorname{Int}P);\mathbb{Z})\cong \mathbb{Z}\oplus \mathbb{Z}$, by Proposition 2 and the homeomorphism type of the manifold $f^{-1}(\mathbb{R}^n-\operatorname{Int}P)$, the 1st homology group of the total space of this obtained bundle whose coefficient ring is \mathbb{Z} is isomorphic to $\mathbb{Z}\oplus \mathbb{Z}\oplus \mathbb{Z}\oplus$

Theorem 7. By P-operations by S^1 to the map $f: M \to \mathbb{R}^2$ here, we have a family of (m+1)-dimensional oriented manifolds $\{M_k\}_{k\in\mathbb{Z}}$ and round fold maps $\{f_k: M_k \to \mathbb{R}^2\}_{k\in\mathbb{Z}}$. Furthermore, we have the followings.

- (1) By any P-operation by S^1 to the map f such that the resulting S^1 -bundle over M is orientable, the resulting source manifold of the resulting map is diffeomorphic to a manifold belonging to the family $\{M_k\}$.
- (2) M_k has the structure of an oriented S^1 -bundle over M such that the 1st homology group of the total space of the bundle obtained by the restriction of the bundle M_k to $f^{-1}(\mathbb{R}^n \operatorname{Int} P)$ with coefficient ring \mathbb{Z} is isomorphic to $\mathbb{Z} \oplus \mathbb{Z} \oplus (\mathbb{Z}/|k|\mathbb{Z})$ and that the 1st homology group $H_1(M_k; \mathbb{Z})$ is isomorphic to $\mathbb{Z} \oplus (\mathbb{Z}/|k|\mathbb{Z})$.
- (3) The inverse image of an axis of f_k is diffeomorphic to the product of the inverse image of an axis of f and S^1 .
- (4) f_k is C^{∞} L-trivial.
- (5) f_k is not C^{∞} S-trivial if k is not 0.

4.3.3. Case 3. Let $m, n \in \mathbb{N}$ and $m > n \geq 2$. We also assume that $m \geq 2n$ and $m - n \geq 3$ hold. Let M_i be a closed and connected oriented manifold admitting a round fold map $f_i: M_i \to \mathbb{R}^n$ satisfying the conditions similar to those mentioned in the beginning of Case 2 (i = 1, 2). We also assume that the fiber of a proper core of f_1 is diffeomorphic to the (m - n)-dimensional standard sphere S^{m-n} and that f_2 is obtained by a trivial spinning construction. Let us construct a round fold map from the connected sum M of the two manifolds M_1 and M_2 into \mathbb{R}^n by applying a method used in [10], [11] and [13].

Let P_i be a proper core of f_i (i=1,2) and let Q be a small closed tubular neighborhood of the connected component of $f_2(S(f_2))$ which is the boundary of $f_2(M_2)$. It easily follows that $f_1|_{f_1^{-1}(P_1)}: f_1^{-1}(P_1) \to P_1$ gives the structure of a trivial S^{m-n} -bundle over P_1 and that $f_2^{-1}(Q)$ is a closed tubular neighborhood of $f_2^{-1}(C) \subset M_2$.

Since the map f_2 is obtained by a trivial spinning construction, the inclusion of $f_2^{-1}(C)$ into M_2 is homotopic to a section of the trivial bundle over the proper core P_2 of f_2 above given by $f_2|_{f_2^{-1}(\partial P_2)}: f_2^{-1}(\partial P_2) \to \partial P_2$ and the section is null-homotopic in the manifolds $f_2^{-1}(P_2)$ and M_2 since the fiber of this trivial bundle is a sphere of dimension m-n>n-1 and (n-1)-connected. From the inequality $m \geq 2n = 2(n-1)+2$, the inclusion of $f^{-1}(C)$ into M_2 is unknot in the C^{∞} category and $f_2^{-1}(Q)$ has the structure of a trivial D^{m-n+1} -bundle over an (n-1)-dimensional sphere $\partial Q \cap f_2(M)$. More precisely, $f_2|_{\partial f_2^{-1}(Q)}: \partial f_2^{-1}(Q) \to \partial Q \cap f_2(M)$ gives the structure of a subbundle of the bundle.

Let $V_1:=f_1^{-1}(P_1)$ and $V_2:=f_2^{-1}(Q)$. From the discussion above, we may regard that the following holds for a diffeomorphism $\Phi:\partial V_2\to\partial V_1$ regarded as a bundle isomorphism between the two trivial S^{m-n+1} -bundles over S^{n-1} inducing a diffeomorphism between the base spaces and an orientation reversing diffeomorphism $\Psi:\partial D^m\to\partial D^m$ extending to a diffeomorphism on D^m or from $M_2-(M_2-D^m)$ onto $M_1-(M_1-D^m)$, where for two manifolds X_1 and X_2 , $X_1\cong X_2$ means that X_1 and X_2 are diffeomorphic.

$$(M_{1} - \operatorname{Int}V_{1}) \bigcup_{\Phi} (M_{2} - \operatorname{Int}V_{2})$$

$$\cong (M_{1} - \operatorname{Int}V_{1}) \bigcup_{\Phi} ((D^{m} - \operatorname{Int}V_{2}) \bigcup (M_{2} - \operatorname{Int}D^{m}))$$

$$\cong (M_{1} - \operatorname{Int}V_{1}) \bigcup_{\Phi} ((S^{m} - (\operatorname{Int}V_{2} \sqcup \operatorname{Int}D^{m})) \bigcup_{\Psi} (M_{2} - \operatorname{Int}D^{m}))$$

$$\cong (M_{1} - \operatorname{Int}D^{m}) \bigcup_{\Psi} (M_{2} - \operatorname{Int}D^{m})$$

This means that the resulting manifold M, which is the connected sum of M_1 and M_2 , admits a round fold map into \mathbb{R}^n . More precisely, the resulting map is obtained by gluing the two maps $f_1|_{M_1-\operatorname{Int}V_1}$ and $f_2|_{M_2-\operatorname{Int}V_2}$. The resulting map $f:M\to\mathbb{R}^n$ is C^∞ L-trivial and the inverse image of the axis of f is represented as a manifold diffeomorphic to the connected sum of two manifolds homeomorphic to $S^1\times S^{n-1}$ with the interior of an (m-n+1)-dimensional standard closed disc removed.

The cohomology group $H^2(M;\mathbb{Z})$ is zero $(n \neq 2)$ or $\mathbb{Z} \oplus \mathbb{Z}$ (n = 2). The manifolds $f_1^{-1}(\mathbb{R}^n - \operatorname{Int} P_1)$ and $f_2^{-1}(\mathbb{R}^n - (P_2 \bigcup Q))$ are diffeomorphic to products of S^{n-1} and manifolds homeomorphic to $S^1 \times S^{m-n}$ with the interior of an (m-n+1)dimensional standard closed disc removed. For a proper core P of f, $f_1^{-1}(\mathbb{R}^n \operatorname{Int} P_1 \bigcup f_2^{-1}(\mathbb{R}^n - (P_2 \bigcup Q))$ is regarded as the inverse image $f^{-1}(\mathbb{R}^n - \operatorname{Int} P)$. In the case where $n \geq 3$ is assumed, any orientable S^1 -bundle over M is trivial since the cohomology group $H^2(M;\mathbb{Z})$ is zero and we can construct a round fold map from $M \times S^1$ into \mathbb{R}^n by a P-operation by S^1 . In the case where n=2 is assumed, we have $H^2(f^{-1}(\mathbb{R}^2 - \operatorname{Int} P); \mathbb{Z}) \cong \mathbb{Z} \oplus \mathbb{Z}$ and the homomorphism between the 2nd cohomology groups induced by the natural inclusion of $f^{-1}(\mathbb{R}^2 - \text{Int}P)$ into M is an isomorphism. In this case, we can set ν_1 as a generator of the group $H^2(f_1^{-1}(\mathbb{R}^n - \operatorname{Int}P_1); \mathbb{Z}) \cong \mathbb{Z}$ and ν_2 as a generator of the group $H^2(f_2^{-1}(\mathbb{R}^n - (P_2 \cup Q)); \mathbb{Z}) \cong \mathbb{Z}$ and by regarding $H^2(f_1^{-1}(\mathbb{R}^n - \operatorname{Int}P_1); \mathbb{Z}) \oplus H^2(f_2^{-1}(\mathbb{R}^n - \mathbb{Z}))$ $(P_2[JQ));\mathbb{Z})$ as $H^2(f^{-1}(\mathbb{R}^2-\mathrm{Int}P);\mathbb{Z})$ and using the isomorphism between the cohomology groups before, we can define the class $\nu_i' \in H^2(M;\mathbb{Z})$ corresponding to ν_i (i=1,2). There exists an oriented manifold $M_{(k_1,k_2)}$ having the structure of an oriented S^1 -bundle over M whose Euler class is $k_1\nu_1' + k_2\nu_2'$ admitting a round fold map into \mathbb{R}^2 obtained by a P-operation by S^1 to f. Thus, we have the following theorem.

Theorem 8. By P-operations by S^1 to the map $f: M \to \mathbb{R}^2$ here, we have a family of (m+1)-dimensional oriented manifolds $\{M_{(k_1,k_2)}\}_{(k_1,k_2)\in\mathbb{Z}\oplus\mathbb{Z}}$ and round fold maps $\{f_{(k_1,k_2)}: M_{(k_1,k_2)}\to\mathbb{R}^2\}_{(k_1,k_2)\in\mathbb{Z}\oplus\mathbb{Z}}$. Furthermore, we have the followings.

- (1) By any P-operation by S^1 to the map f such that the resulting S^1 -bundle over M is orientable, the resulting source manifold of the resulting map is diffeomorphic to a manifold belonging to the family $\{M_{(k_1,k_2)}\}$.
- (2) $M_{(k_1,k_2)}$ has the structure of an oriented S^1 -bundle over M and the followings hold.
 - (a) The 1st homology group of the total space of the bundle obtained by the restriction of the bundle $M_{(k_1,k_2)}$ to $f_1^{-1}(\mathbb{R}^2 \text{Int}P_1) \subset M$ with coefficient ring \mathbb{Z} is isomorphic to $\mathbb{Z} \oplus \mathbb{Z} \oplus (\mathbb{Z}/|k_1|\mathbb{Z})$.

- (b) The 1st homology group of the total space of the bundle obtained by the restriction of the bundle $M_{(k_1,k_2)}$ to $f_2^{-1}(\mathbb{R}^n-(P_2\bigcup Q))\subset M$ is isomorphic to $\mathbb{Z} \oplus \mathbb{Z} \oplus (\mathbb{Z}/|k_2|\mathbb{Z})$.
- (c) Let $k_1 \neq 0$ and $k_2 \neq 0$ hold and let k_0 be the LCM of $|k_1|$ and $|k_2|$. Then the 1st homology group of $M_{(k_1,k_2)}$ with coefficient ring $\mathbb Z$ is isomorphic to $\mathbb{Z} \oplus \mathbb{Z} \oplus (\mathbb{Z}/k_0\mathbb{Z})$.
- (d) Let $k_1 = 0$ or $k_2 = 0$ hold. Then the 1st homology group of $M_{(k_1,k_2)}$ with coefficient ring \mathbb{Z} is isomorphic to $\mathbb{Z} \oplus \mathbb{Z}$.
- (3) The inverse image of an axis of $f_{(k_1,k_2)}$ is diffeomorphic to the product of the inverse image of an axis of f and S^1 .
- (4) $f_{(k_1,k_2)}$ is C^{∞} L-trivial.
- (5) If $k_1 \neq 0$ and $k_2 \neq 0$ hold, then $f_{(k_1,k_2)}$ is not C^{∞} S-trivial. (6) $f_{(k_1,k_2)}$ and $f_{(k_3,k_4)}$ are not C^{∞} equivalent if $|k_1| \neq |k_3|$ or $|k_2| \neq |k_4|$ holds.

References

- [1] D. Barden, Simply Connected Five-Manifolds, Ann. of Math., 3 (82) (Nov., 1965), 365–385.
- [2] O. Burlet G. de Rham, Sur certaines applications generiques d'une variete close a 3 dimensions dans le plan, Enseign. Math. 20 (1974). 275-292.
- [3] H. Duan C. Liang, Circle bundles over 4-manifolds, Archiv der Mathematik 3 (85) (September 2005), 278-282.
- [4] C. Ehresmann, Les connexions infinitesimales dans un espace fibre differentiable, Colloque de Topologie, Bruxelles (1950), 29–55.
- [5] Y. Eliashberg, On singularities of folding type, Math. USSR Izv. 4 (1970), 1119–1134.
- [6] Y. Eliashberg, Surgery of singularities of smooth mappings, Math. USSR Izv. 6 (1972), 1302-
- [7] Y. K. S. Furuya, P. Porto, On special generic maps from a closed manifold into the plane, Topology Appl. 35 (1990), 41–52.
- [8] M. Golubitsky, V. Guillemin, Stable Mappings and Their Singularities, Graduate Texts in Mathematics (14), Springer-Verlag (1974).
- [9] A. E. Hatcher, A proof of the Smale conjecture, Ann. of Math. 117 (1983), 553-607.
- [10] N. Kitazawa, On round fold maps (in Japanese), to appear in RIMS Kokyuroku Bessatsu
- [11] N. Kitazawa, On manifolds admitting fold maps with singular value sets of concentric spheres, Doctoral Dissertation, Tokyo Institute of Technology (2014).
- [12] N. Kitazawa, Fold maps with singular value sets of concentric spheres, to appear in Hokkaido Mathematical Journal.
- [13] N. Kitazawa, Differential topology of manifolds admitting round fold maps, arXiv:1304.0618 (April 2013).
- [14] N. Kitazawa, Constructions of round fold maps on smooth bundles, to appear in Tokyo J. of Math. Volume 37, Number 2 or later.
- [15] M. Kobayashi, O. Saeki, Simplifying stable mappings into the plane from a global viewpoint, Trans. Amer. Math. Soc. 348 (1996), 2607-2636.
- [16] J. Milnor, J. Stasheff, Characteristic classes, Annals of Mathematics Studies, No. 76. Princeton, N. J; Princeton University Press (1974).
- [17] S. Smale, Diffeomorphisms of the 2-sphere, Proc. Amer. Math. Soc. 10 (1959) 621?-626.
- [18] O. Saeki, Notes on the topology of folds, J. Math. Soc. Japan Volume 44, Number 3 (1992),
- [19] O. Saeki, Topology of special generic maps of manifolds into Euclidean spaces, Topology Appl. 49 (1993), 265–293.
- [20] O. Saeki, K. Sakuma, On special generic maps into R³, Pacific J. Math. 184 (1998), 175–193.
- [21] O. Saeki, K. Suzuoka, Generic smooth maps with sphere fibers J. Math. Soc. Japan Volume 57. Number 3 (2005), 881–902.
- [22] K. Sakuma, On the topology of simple fold maps, Tokyo J. of Math. Volume 17, Number 1 (1994), 21-32.

- [23] R. Thom, Les singularites des applications differentiables, Ann. Inst. Fourier (Grenoble) 6 (1955-56), 43–87.
- [24] H. Whitney, On singularities of mappings of Euclidean spaces: I, mappings of the plane into the plane, Ann. of Math. 62 (1955), 374–410.

Department of Mathematics, Tokyo Institute of Technology, 2-12-1 Ookayama, Meguroku, Tokyo 152-8551, JAPAN, Tel+81-(0)3-5734-2205, Fax +81-(0)3-5734-2738,

 $E\text{-}mail\ address: \verb+kitazawa.n.aa@m.titech.ac.jp+$