LINEAR DIFFERENTIAL EQUATIONS ON THE RIEMANN SPHERE AND REPRESENTATIONS OF QUIVERS

KAZUKI HIROE

ABSTRACT. Our interest in this paper is a generalization of the additive Deligne-Simpson problem which is originally defined for Fuchsian differential equations on the Riemann sphere. We shall extend this problem to differential equations having an arbitrary number of unramified irregular singular points and determine the existence of solutions of the generalized additive Deligne-Simpson problems. Moreover we apply this result to the geometry of the moduli spaces of stable meromorphic connections of trivial bundles on the Riemann sphere. Namely, open embedding of the moduli spaces into quiver varieties is given and the non-emptiness condition of the moduli spaces is determined. Furthermore the connectedness of the moduli spaces is shown.

INTRODUCTION

The additive Deligne-Simpson problem asks the existence of an irreducible Fuchsian differential equation on the Riemann sphere with prescribed local data. In this paper we shall consider analogous problems for differential equations on the Riemann sphere with unramified irregular singular points.

First of all let us recall the classical case, i.e., the additive Deligne-Simpson problem for systems of linear Fuchsian differential equations (see [26]). A system of first order linear differential equations is called *Fuchsian* if it is of the form

$$\frac{d}{dz}Y = \sum_{i=1}^{p} \frac{A_i}{z - a_i}Y \quad (A_i \in M(n, \mathbb{C}), i = 1, \dots, p).$$

Here we call each A_i the residue matrix at the singular point a_i . Also $A_0 := -\sum_{i=1}^p A_i$ is called the residue matrix at ∞ . We say that $\frac{d}{dz}Y = \sum_{i=1}^p \frac{A_i}{z-a_i}Y$ is *irreducible* if A_0, \ldots, A_p have no nontrivial simultaneous invariant vector subspace of \mathbb{C}^n , i.e., if there exists $W \subsetneq \mathbb{C}^n$ such that $A_i W \subset W$ for all $i = 0, \ldots, p$, then $W = \{0\}$.

Definition 0.1 (additive Deligne-Simpson problem (classical case)). The additive Deligne-Simpson problem consists of points a_1, \ldots, a_p in \mathbb{C} and conjugacy classes C_0, C_1, \ldots, C_p in $M(n, \mathbb{C})$. A solution of the problem is an *irreducible* Fuchsian differential equation

$$\frac{dY}{dz} = \sum_{i=1}^{p} \frac{A_i}{z - a_i} Y \quad (A_i \in M(n, \mathbb{C}), i = 1, \dots, p)$$

whose residue matrices $A_i \in C_i$ for $i = 0, \ldots, p$.

This problem is developed by V. Kostov as an analogy of the problem studied by P. Deligne and C. Simpson (see [31]), so called multiplicative

Deligne-Simpson problem. After Kostov who gave an necessary and sufficient condition of the existence of solutions under generic conditions in [26], a complete necessary and sufficient condition was given by W. Crawley-Boevey [9].

As a generalization of this problem, it seems to be natural to consider similar problems for non-Fuchsian equations (see for example [4],[5], [27]). Our generalization in this paper is an extension of the work of P. Boalch in [4] and Kostov's way of generalization given in [27] is somewhat different from ours. Before formulating the generalized problem precisely, let us recall some facts of local formal theory around irregular singular points of differential equations. Let M(n, R) be the ring of $n \times n$ matrices with coefficients in a commutative ring R and $\operatorname{GL}(n, R)$ the group which consists of all invertible elements in M(n, R) of the multiplication. Set $\mathbb{C}((z)) :=$ $\{\sum_{i=r}^{\infty} c_i z^i \mid c_i \in \mathbb{C}, r \in \mathbb{Z}\}$ and $\mathbb{C}[\![z]\!] := \{\sum_{i=0}^{\infty} c_i z^i \mid c_i \in \mathbb{C}\}$. Let us attach to $C = \sum_{i=r}^{\infty} c_i z^i \in M(n, \mathbb{C}((z)))$ the integer $\operatorname{ord}(C) := \min\{i \mid c_i \neq 0\}$ called the order. For $A \in M(n, \mathbb{C}((z)))$ and $X \in \operatorname{GL}(n, \mathbb{C}((z)))$ the gauge transformation of A by X is

$$X[A] := XAX^{-1} + \left(\frac{d}{dz}X\right)X^{-1}.$$

Definition 0.2 (Hukuhara-Turrittin-Levelt normal form). If an element $B \in M(n, \mathbb{C}((z)))$ is of the form

$$B = \operatorname{diag}(q_1(z^{-1})I_{n_1} + R_1 z^{-1}, \dots, q_m(z^{-1})I_{n_m} + R_m z^{-1})$$

with $q_i(s) \in s^2 \mathbb{C}[s]$ satisfying $q_i \neq q_j$ if $i \neq j$ and $R_i \in M(n_i, \mathbb{C})$, then B is called the Hukuhara-Turrittin-Levelt normal form or the HTL normal form shortly. Here I_m is the identity matrix of $M(m, \mathbb{C})$.

Now let us introduce truncated orbits which play the same role as the conjugacy classes C_i of residue matrices in the Deligne-Simpson problem of Fuchsian systems. Let us define $G_k := \operatorname{GL}(n, \mathbb{C}[\![z]\!]/z^k \mathbb{C}[\![z]\!]), k \geq 1$, which can be identified with

$$\left\{ A_0 + A_1 x + \dots + A_{k-1} z^{k-1} \middle| \begin{array}{l} A_0 \in \operatorname{GL}(n, \mathbb{C}), \ A_i \in M(n, \mathbb{C}), \\ i = 1, \dots, k-1 \end{array} \right\}.$$

Also define

$$\mathfrak{g}_{k} := M(n, \mathbb{C}[\![z]\!]/z^{k}\mathbb{C}[\![z]\!])$$

= $\left\{ A_{0} + A_{1}z + \dots + A_{k-1}z^{k-1} \mid A_{i} \in M(n, \mathbb{C}), i = 0, \dots, k-1 \right\}.$

The dual vector space $\mathfrak{g}_k^* := \operatorname{Hom}_{\mathbb{C}}(\mathfrak{g}_k, \mathbb{C})$ is identified with

$$M(n, z^{-k}\mathbb{C}[\![z]\!]/\mathbb{C}[\![z]\!]) = \left\{\frac{A_k}{z^k} + \dots + \frac{A_1}{z} \mid A_i \in M(n, \mathbb{C})\right\}$$

by the nondegenerate bilinear form $\mathfrak{g}_k \times \mathfrak{g}_k^* \ni (A, B) \mapsto \operatorname{Res}(\operatorname{tr}(AB)) \in \mathbb{C}$. Here we set $\operatorname{Res}(\sum_{i=r}^{\infty} A_i z^i) := A_{-1}$.

Then an HTL normal form B with $\operatorname{ord}(B) \geq -k$ can be seen as an element in \mathfrak{g}_k^* . Thus we can consider the G_k -orbit $\mathcal{O}_B := \{gBg^{-1} \in \mathfrak{g}_k^* \mid g \in G_k\}$ of B in \mathfrak{g}_k^* called the *truncated orbit* of B. Now let us formulate a generalization of Deligne-Simpson problem. **Definition 0.3** (generalized additive Deligne-Simpson problem). The generalized additive Deligne-Simpson problem consists of a collection of points a_1, \ldots, a_p in \mathbb{C} , of nonzero positive integers k_0, \ldots, k_p and of HTL normal forms $B_i \in \mathfrak{g}_{k_i}^* \subset M(n, \mathbb{C}((z)))$ for $i = 0, \ldots, p$. A solution of the generalized additive Deligne-Simpson problem is an *irreducible* differential equation

$$\frac{d}{dz}Y = \left(\sum_{i=1}^{p}\sum_{j=1}^{k_i}\frac{A_{i,j}}{(z-a_i)^j} + \sum_{2\le j\le k_0}A_{0,j}z^{j-2}\right)Y$$

satisfying that $A^{(i)}(z) \in \mathcal{O}_{B_i}$ for i = 0, ..., p. Here $A^{(i)}(z) := \sum_{j=1}^{k_i} A_{i,j} z^{-j}$ for i = 0, ..., p and $A_{0,1} := -\sum_{i=1}^{p} A_{i,1}$. We say

$$\frac{d}{dz}Y = \left(\sum_{i=1}^{p}\sum_{j=1}^{k_i}\frac{A_{i,j}}{(z-a_i)^j} + \sum_{2 \le j \le k_0}A_{0,j}x^{j-2}\right)Y$$

is *irreducible* if the collection $(A_{i,j})_{\substack{0 \le i \le p \\ 1 \le j \le k_i}}$ of coefficient matrices is irreducible.

This can be seen as a natural generalization of additive Deligne-Simpson problems which contains the original problems for Fuchsian equations as the special case $k_0 = \cdots = k_p = 1$. In the case $k_1 = \cdots = k_p = 1$ and $k_0 \leq 3$, P. Boalch obtains a necessary and sufficient condition for the existence of a solution of the generalized additive Deligne-Simpson problem in [4] and for an arbitrary k_0 , see [16].

In order to determine the existence condition of a solution of the additive Deligne-Simpson problem for Fuchsian systems, Crawley-Boevey [9] shows that Fuchsian systems can be realized as representations of quivers, and applies the existence theorem of irreducible representations of deformed preprojective algebras associated with the quivers to the existence of solutions of additive Deligne-Simpson problems. One can find a review of his work in Section 2 and 3.

As a generalization of his work, we can associate our generalized additive Deligne-Simposn problem with a quiver defined as follows. The detail of the construction shall be explained in Sections 4.1, 4.2 and 4.3. Let us suppose that HTL normal forms B_0, \ldots, B_p are written by

$$B_{i} = \operatorname{diag}\left(q_{[i,1]}(z^{-1})I_{n_{[i,1]}} + R_{[i,1]}z^{-1}, \dots, q_{[i,m_{i}]}(z^{-1})I_{n_{[i,m_{i}]}} + R_{[i,m_{i}]}z^{-1}\right)$$

and choose complex numbers $\xi_1^{[i,j]}, \ldots, \xi_{e_{[i,j]}}^{[i,j]}$ so that

$$\prod_{k=1}^{e_{[i,j]}} (R_{[i,j]} - \xi_k^{[i,j]}) = 0$$

for $i = 0, \ldots, p$ and $j = 1, \ldots, m_i$. Set $I_{irr} := \{i \in \{0, \ldots, p\} \mid m_i > 1\} \cup \{0\}$ and $I_{reg} := \{0, \ldots, p\} \setminus I_{irr}$.

Then let ${\sf Q}$ be the quiver with the set of vertices

$$Q_0 := \left\{ [i, j] \left| \begin{array}{c} i \in I_{irr}, \\ j = 1, \dots, m_i \end{array} \right\} \cup \left\{ [i, j, k] \left| \begin{array}{c} i = 0, \dots, p, \\ j = 1, \dots, m_i, \\ k = 1, \dots, e_{[i, j]} - 1 \end{array} \right\} \right\}$$

and the set of arrows

$$\begin{aligned} \mathsf{Q}_{1} &= \left\{ \rho_{[i,j']}^{[0,j]} \colon [0,j] \to [i,j'] \middle| \begin{array}{l} j = 1, \dots, m_{0}, \\ i \in I_{\mathrm{irr}} \setminus \{0\}, \\ j' = 1, \dots, m_{i} \end{array} \right\} \\ &\cup \left\{ \rho_{[i,j],[i,j']}^{[k]} \colon [i,j] \to [i,j'] \middle| \begin{array}{l} i \in I_{\mathrm{irr}}, \ 1 \leq j < j' \leq m_{i}, \\ 1 \leq k \leq d_{i}(j,j') \end{array} \right\} \\ &\cup \left\{ \rho_{[i,j,1]} \colon [i,j,1] \to [i,j] \mid i \in I_{\mathrm{irr}}, \ j = 1, \dots, m_{i} \right\} \\ &\cup \left\{ \rho_{[0,j]}^{[i,1,1]} \colon [i,1,1] \to [0,j] \mid i \in I_{\mathrm{reg}}, \ j = 1, \dots, m_{0} \right\} \\ &\cup \left\{ \rho_{[i,j,k]} \colon [i,j,k] \to [i,j,k-1] \middle| \begin{array}{l} i = 0, \dots, p, \\ j = 1, \dots, m_{i}, \\ k = 2, \dots, e_{[i,j]} - 1 \end{array} \right\}. \end{aligned}$$

Here $d_i(j, j') := \deg_{\mathbb{C}[z]}(q_{[i,j]}(z) - q_{[i,j']}(z)) - 2$. To each vector $\beta \in \mathbb{Z}^{Q_0}$, we associate integers

$$q(\beta) := \sum_{a \in \mathsf{Q}_0} \beta_a^2 - \sum_{\rho \in \mathsf{Q}_1} \beta_{s(\rho)} \beta_{t(\rho)}, \qquad p(\beta) := 1 - q(\beta)$$

Here $s(\rho)$ and $t(\rho)$ are the source and target of the arrow ρ respectively. Let $\alpha = (\alpha_a)_{a \in \mathbb{Q}_0} \in \mathbb{Z}^{\mathbb{Q}_0}$ be the vector defined by setting $\alpha_{[i,j]} := n_{[i,j]}$ and $\alpha_{[i,j,k]} := \operatorname{rank} \prod_{l=1}^{k} (R_j^{(i)} - \xi_l^{[i,j]})$. Also define $\lambda = (\lambda_a)_{a \in \mathbb{Q}_0} \in \mathbb{C}^{\mathbb{Q}_0}$ by $\lambda_{[i,j]} := -\xi_1^{[i,j]}$ for $i \in I_{\operatorname{irr}} \setminus \{0\}, j = 1, \ldots, m_i, \lambda_{[0,j]} := -\xi^{[0,j]} - \sum_{i \in I_{\operatorname{reg}}} \xi_1^{[i,1]}$ for $j = 1, \ldots, m_0$, and $\lambda_{[i,j,k]} := \xi_k^{[i,j]} - \xi_{k+1}^{[i,j]}$ for $i = 0, \ldots, p, j = 1, \ldots, m_i$ and $k = 1, \ldots, e_{[i,j]} - 1$. The following sublattice of $\mathbb{Z}^{\mathbb{Q}_0}$ plays an essential role in this paper,

$$\mathcal{L} := \left\{ \beta \in \mathbb{Z}^{\mathbf{Q}_0} \left| \sum_{j=1}^{m_0} \beta_{[0,j]} = \sum_{j=1}^{m_i} \beta_{[i,j]} \text{ for all } i \in I_{\mathrm{irr}} \setminus \{0\} \right\}.$$

Set $\mathcal{L}^+ := \mathcal{L} \cap (\mathbb{Z}_{>0})^{\mathsf{Q}_0}$.

Then the following is the main theorem of this paper.

Theorem 0.4 (see Theorem 7.12). Let us consider the generalized additive Deligne-Simpson problem consisting of positive integers k_0, \ldots, k_p and HTL normal forms $B_i \in \mathfrak{g}_{k_i}^*$ for $i = 0, \ldots, p$. Let us define $\mathbf{Q} = (\mathbf{Q}_0, \mathbf{Q}_1), \alpha \in (\mathbb{Z}_{\geq 0})^{\mathbf{Q}_0}$ and $\lambda \in \mathbb{C}^{\mathbf{Q}_0}$ as above. Then the generalized additive Deligne-Simpson problem has a solution if and only if the following are satisfied,

- (1) α is a positive root of Q and $\alpha \cdot \lambda = \sum_{a \in Q_0} \alpha_a \lambda_a = 0$,
- (2) for any decomposition $\alpha = \beta_1 + \cdots + \beta_r$ where $\beta_i \in \mathcal{L}^+$ are positive roots of Q satisfying $\beta_i \cdot \lambda = 0$, we have

$$p(\alpha) > p(\beta_1) + \dots + p(\beta_r).$$

Here we note that the condition 2 is weaker than the corresponding condition of Crawley-Boevey's theorem (see Theorem 1.2 in [8] and see also Theorem 2.6) since we deal only with positive roots in \mathcal{L} . If, however, $I_{\rm irr} = \{0\}$ which contains known cases by Crawley-Boevey [9], Boalch [4], and [16], then $\mathcal{L} = \mathbb{Z}^{Q_0}$ and the conditions in the above theorem coincide with Crawley-Boevey's one. Thus the theorem covers the above preceding known results for the additive Delinge-Simpson problems.

Let us discuss moduli spaces of meromorphic connections. In the theory of isomonodromic deformation, the Riemann-Hilbert problem of moduli spaces of meromorphic connections not only with regular singularities but also irregular singularities plays a central role, see [22] for instance. Thus many researchers are interested in the geometry of the moduli spaces of the meromorphic connections with irregular singularities, see [3] [7],[17] for instance. As an application of our main theorem, we shall discuss some geometric properties of the moduli spaces. Precisely to say, following the Boalch's paper [3], we define the moduli space $\mathfrak{M}(\mathbf{B})$ of meromorphic connections on trivial bundles associated with the collection of HTL normal forms $\mathbf{B} = (B_i)_{0 \le i \le p}$, see Section 1.2. Then we shall show that $\mathfrak{M}(\mathbf{B})$ can be embedded onto an open subset of a quiver variety $\mathfrak{M}_{\lambda}^{\mathrm{reg}}(\mathbf{Q}, \alpha)$.

Theorem 0.5 (see Theorem 5.15). Let us take $\mathbf{B} = (B_i)_{0 \le i \le p}$, the collection of HTL normal forms, the quiver \mathbf{Q} , $\alpha \in (\mathbb{Z}_{\ge 0})^{\mathbf{Q}_0}$ and $\lambda \in \mathbb{C}^{\mathbf{Q}_0}$ as in Theorem 0.4. Then there exists $\lambda' \in \mathbb{C}^{\mathbf{Q}_0}$ and an injection

$$\Phi \colon \mathfrak{M}(\mathbf{B}) \hookrightarrow \mathfrak{M}_{\lambda'}^{reg}(\mathsf{Q},\alpha)$$

such that

$$\Phi(\mathfrak{M}(\mathbf{B})) = \left\{ x \in \mathfrak{M}_{\lambda'}^{reg}(\mathbf{Q}, \alpha) \middle| \det \left(x_{\rho_{[i,j']}^{[0,j]}} \right)_{\substack{1 \le j \le m_0 \\ 1 \le j' \le m_i}} \neq 0, \ i \in I_{irr} \setminus \{0\} \right\}.$$

In particular if $I_{irr} = \{0\}$, then $\lambda' = \lambda$ and Φ is bijective.

As a corollary of this embedding theorem, we can show the connectedness of $\mathfrak{M}(\mathbf{B})$.

Theorem 0.6 (see Theorem 5.14). If $\mathfrak{M}(\mathbf{B}) \neq \emptyset$, then $\mathfrak{M}(\mathbf{B})$ has a structure of connected complex manifold.

Let us give a remark of the theorems which is already obtained by many researchers under several restrictions. If $\#I_{irr} = 1$, it is known that $\mathfrak{M}(\mathbf{B})$ is isomorphic to the quiver variety $\mathfrak{M}_{\lambda}^{\mathrm{reg}}(\mathbf{Q}, \alpha)$ by Crawley-Boevey for the Fuchsian case in [9] and by Boalch in [4] and the work of D. Yamakawa with the author in [16] for the case $\#I_{irr} = 1$. However for the case $\#I_{irr} > 1$, Boalch gave an example of a moduli space which is not isomorphic to any quiver varieties in [4]. To avoid the difficulty, Yamakawa [36] defines a generalization of quiver varieties which can be realized as $\mathfrak{M}(\mathbf{B})$ for general I_{irr} under the restriction $m_i \leq 2$ for all $i \in I_{irr}$. In the above theorem we impose no restriction to **B** and show that although the moduli space $\mathfrak{M}(\mathbf{B})$ may not be isomorphic to a quiver variety itself as Boalch suggested, there always exists the embedding onto the open subset of the quiver variety.

Furthermore, Theorem 0.4 determines the non-emptiness of the moduli space $\mathfrak{M}(\mathbf{B})$ as a generalization of the results for the case $\#I_{irr} \leq 1$ by Crawley-Boevey [9], Boalch [4] and Yamakawa and the author [16].

Theorem 0.7. (see Corollary 7.13) The moduli space $\mathfrak{M}(\mathbf{B})$ is non-empty if and only if the following are satisfied,

- (1) α is a positive root of \mathbf{Q} and $\alpha \cdot \lambda = \sum_{a \in \mathbf{Q}_0} \alpha_a \lambda_a = 0$, (2) for any decomposition $\alpha = \beta_1 + \dots + \beta_r$ where $\beta_i \in \mathcal{L}^+$ are positive roots of Q satisfying $\beta_i \cdot \lambda = 0$, we have

$$p(\alpha) > p(\beta_1) + \dots + p(\beta_r).$$

Finally let us mention the multiplicative Deligne-Simpson problem. Simpson and also Kostov gave necessary and sufficient conditions for the existence of a solution of the problem under some generic conditions, see [31] and [26]. In [11], Crawley-Boevey and P. Shaw gave a correspondence between the space of solutions of the problem and the so-called multiplicative quiver variety (see also [35] for multiplicative quiver varieties) and obtained a sufficient condition for the existence of a solution. Furthermore in [6], Boalch considered a generalization of the multiplicative Deligne-Simpson problem for differential equations with irregular singular points and gave correspondence between the space of solutions of the problem and the multiplicative quiver variety which is a further generalization of the multiplicative quiver variety considered by Crawley-Boevey and Shaw in [11].

Acknowledgment.

The author expresses his gratitude to Toshio Oshima and Daisuke Yamakawa. This project would not have completed without the collaborations with them. He also thank Philip Boalch for reading the earlier version of this paper and giving him helpful comments. This work is influenced by many other mathematicians, in particular the participants of "Workshop on Accessory Parameters" mainly organized by Yoshishige Haraoka. Most work of this project was done during his stay in Research Institute for Mathematical Sciences in Kyoto University. He thanks for the hospitality and the support.

1. Additive Delinge-Simpson problem

In this section, we shall define a generalization of the additive Delinge-Simpson problem for differential equations with at most unramified irregular singularities on the Riemann sphere. Also we recall moduli spaces of meromorphic connections on trivial vector bundles over the Riemann sphere studied by Boalch in [3] and moreover see that the additive Deligne-Simpson problem is related to the non-emptiness problem of the moduli spaces.

1.1. A generalization of the additive Deligne-Simpson problem. As we saw in Introduction, the original additive Delinge-Simpson problem for Fuchsian differential equations consists of a collection of conjugacy classes of $M(n,\mathbb{C})$. The counterparts for irregular singular cases of the conjugacy classes are Hukuhara-Turrittin-Levelt normal forms of $M(n, \mathbb{C}((z)))$. We shall recall the definition of Hukuhara-Turrittin-Levelt normal forms and give a definition of the additive Deline-Simpson problem for differential equations with at most unramified irregular singularities.

Let us consider a differential equation

$$\frac{d}{dz}Y = AY, \quad (A \in M(n, \mathbb{C}((z)))).$$

For $X \in \operatorname{GL}(n, \mathbb{C}((z)))$, we define a new differential equation $\frac{d}{dz}\tilde{Y} = B\tilde{Y}$ by

$$B := XAX^{-1} + \left(\frac{d}{dz}X\right)X^{-1}.$$

We write B =: X[A] and call this operation the gauge transform of A by X. Let $\mathbb{C}((t))$ be a finite field extension of $\mathbb{C}((z))$, namely there exists $r \in \mathbb{Z}_{\geq 1}$ such that $t^r = z$. Then the differential equation $\frac{d}{dz}Y = AY$ over $\mathbb{C}((z))$ defines the differential equation $\frac{d}{dt}Z = \overline{A}Z$ over $\mathbb{C}((t))$ where $\overline{A} := rt^{r-1}AZ$.

Definition 1.1 (HTL normal form). By Hukuhara-Turrittin-Levelt normal form or HTL normal form for short, we mean an element in $M(n, \mathbb{C}((t)))$ of the form

diag
$$(q_1(t^{-1})I_{n_1} + R_1t^{-1}, \dots, q_m(t^{-1})I_{n_m} + R_mt^{-1})$$

where $t^r = z$, $q_i(s) \in s^2 \mathbb{C}[s]$ satisfying $q_i \neq q_j$ if $i \neq j$, and $R_i \in M(n_i, \mathbb{C})$. In particular when r = 1, the normal form is said to be *unramified*.

The following is a fundamental fact of the local formal theory of differential equations with irregular singularity.

Theorem 1.2 (Hukuhara-Turrittin-Levelt, see [33] for instance). For any $A \in M(n, \mathbb{C}((z)))$, there exists a field extension $\mathbb{C}((t)) \supset \mathbb{C}((z))$ with $t^r = z$, $r \in \mathbb{Z}_{\geq 1}$ and $X \in \mathrm{GL}(n, \mathbb{C}((t)))$ such that $\overline{X[A]}$ is an HTL normal form in $M(n, \mathbb{C}((t)))$.

We call this $\overline{X[A]}$ the normal form of A with the ramification index r. Let us consider an unramified HTL normal form

$$B = \operatorname{diag} \left(q_1(z^{-1})I_{n_1} + R_1 z^{-1}, \dots, q_m(z^{-1})I_{n_m} + R_m z^{-1} \right)$$

and set

 $k := \max_{i=1,\dots,m} \{ \deg_{\mathbb{C}[z^{-1}]} q_i(z^{-1}) \}.$

We shall consider an orbit of B under the following group action. Let us define $G_k := \operatorname{GL}(n, \mathbb{C}[\![z]\!]/z^k \mathbb{C}[\![z]\!])$ which can be identified with

$$\left\{A_0 + A_1 z + \dots + A_{k-1} z^{k-1} \in \sum_{i=0}^{k-1} M(n, \mathbb{C}) z^i \, \middle| \, A_0 \in \mathrm{GL}(n, \mathbb{C})\right\}.$$

Also define

$$\mathfrak{g}_{k} := M(n, \mathbb{C}[\![z]\!]/z^{k}\mathbb{C}[\![z]\!])$$

$$\cong \left\{ A_{0} + A_{1}z + \dots + A_{k-1}z^{k-1} \, \middle| \, A_{i} \in M(n, \mathbb{C}), \, i = 0, 1, \dots, k-1 \right\}.$$

The group G_k acts on \mathfrak{g}_k by the *adjoint action* $\operatorname{Ad}(g)X := gXg^{-1}$ for $g \in G_k, X \in \mathfrak{g}_k$. The dual vector space \mathfrak{g}_k^* is identified with

$$M(n, z^{-k}\mathbb{C}\llbracket z \rrbracket)/\mathbb{C}\llbracket z \rrbracket) \cong \left\{ \frac{A_k}{z^k} + \dots + \frac{A_1}{z} \middle| A_i \in M(n, \mathbb{C}), i = 1, \dots, k \right\}$$

by the bilinear form

$$\mathfrak{g}_k \times \mathfrak{g}_k^* \ni (A, B) \mapsto \operatorname{Res}(\operatorname{tr}(AB)) \in \mathbb{C}$$

where $\operatorname{Res}(\sum_{i=r}^{\infty} a_i z^i) := a_{-1}$ for $\sum_{i=r}^{\infty} a_i z^i \in \mathbb{C}((z))$. Let us note that the *coadjoint action* of G_k on \mathfrak{g}_k^* is defined by $(\operatorname{Ad}^*(g)f)(X) := f(\operatorname{Ad}(g^{-1})X)$ for $g \in G_k, f \in \mathfrak{g}_k^*, X \in \mathfrak{g}_k$. The coadjoint action induces the action of

 G_k on $M(n, z^{-k}\mathbb{C}[\![z]\!]/\mathbb{C}[\![z]\!])$ defined by $\mathrm{Ad}^*(g)Z := g^{-1}Zg$ for $g \in G_k, Z \in M(n, z^{-k}\mathbb{C}[\![z]\!]/\mathbb{C}[\![z]\!]) \cong \mathfrak{g}_k^*$.

Since we can regard $B \in \mathfrak{g}_k^*$, the orbit of B under the coadjoint action of G_k .

Definition 1.3 (truncated orbit). Let us regard $B \in \mathfrak{g}_k^*$. Then

$$\mathcal{O}_B := \{ \mathrm{Ad}^*(g) B \mid g \in G_k \}$$

is called the *truncated orbit* of B.

Now we are ready to define a generalization of the additive Delinge-Simpson problem for differential equations with unramified irregular singularities. We say that a collection of matrices $(A_1, \ldots, A_s) \in M(n, \mathbb{C})^s$ is *irreducible* if (A_1, \ldots, A_s) has no nontrivial invariant subspace of \mathbb{C}^n , i.e., if a subspace $W \subset \mathbb{C}^n$ satisfies that $A_iW \subset W$ for all $i = 1, \ldots, s$, then $W = \{0\}$ or \mathbb{C}^n . For a differential equation

$$\frac{d}{dz}Y = \left(\sum_{i=1}^{p}\sum_{j=1}^{k_i}\frac{A_{i,j}}{(z-a_i)^j} + \sum_{2\le j\le k_0}A_{0,j}z^{j-2}\right)Y,$$

the *principal term* at the singular point a_i is

$$A_i(z_i) := \sum_{j=1}^{k_i} A_{i,j} z_i^{-j}$$

for each i = 0, ..., p. Here we set $A_{0,1} := -\sum_{i=1}^{p} A_{i,1}$ and $z_i := z - a_i$, $i = 1, ..., p, z_0 := \frac{1}{z}$. This differential equation is said to be *irreducible* if the collection of the matrices $(A_{i,j})_{\substack{0 \le i \le p, \\ 1 \le i \le k_i}}$ is irreducible.

Definition 1.4 (additive Delinge-Simpson problem). Let us take $k_i \in \mathbb{Z}_{\geq 1}$ and unramified HTL normal forms $B_i \in \mathfrak{g}_{k_i}^*$ for $i = 0, 1, \ldots, p$. Then a *solution* of the additive Delinge-Simpson problem for the collection of the unramified HTL normal forms (B_0, B_1, \ldots, B_p) is an irreducible differential equation

$$\frac{d}{dz}Y = \left(\sum_{i=1}^{p}\sum_{j=1}^{k_i}\frac{A_{i,j}}{(z-a_i)^j} + \sum_{2 \le j \le k_0}A_{0,j}z^{j-2}\right)Y$$

such that the principal term at each singular point a_i , i = 0, 1, ..., p satisfies

$$A_i(z) \in \mathcal{O}_{B_i}.$$

Remark 1.5. Let us note that if $k_0 = k_1 = \cdots = k_p = 1$, then $G_{k_i} = \operatorname{GL}(n, \mathbb{C})$ and $\mathfrak{g}_{k_i}^* = M(n, \mathbb{C})$. Thus the truncated orbits \mathcal{O}_{B_i} are just conjugacy classes of $M(n, \mathbb{C})$. Therefore the additive Delinge-Simpson problem in Definition 1.4 contains the original additive Delinge-Simpson problem for Fuchsian differential equations.

1.2. Moduli spaces of meromorphic connections and additive Delinge-Simpson problem. In this section we quickly recall the definition of moduli spaces of meromorphic connections on trivial vector bundles over the Riemann sphere following [3]. For the detailed treatment can be found in the original paper by Boalch [3] and we also refer [16] and their references. The solvability of the additive Delinge-Simpson problems can be seen as the problem determining the necessary and sufficient condition of the nonemptiness of the moduli spaces

Let us recall the notion of meromorphic connections and see their relationship with differential equations. For $f = \sum_{i>-\infty}^{\infty} a_i z^i \in \mathbb{C}((z))$, the order is

$$\operatorname{ord}(f) := \min\{i \mid a_i \neq 0\}.$$

If f = 0, we formally put $\operatorname{ord}(f) = \infty$. For a meromorphic function f locally defined near $a \in \mathbb{P}^1$, we denote the germ of f at a by f_a . We may see $f_a \in \mathbb{C}((z_a))$ by setting $z_a = z - a$ if $a \in \mathbb{C}$ and $z_a = 1/z$ if $a = \infty$ where we take z as the standard coordinate of \mathbb{C} . Then define

$$\operatorname{ord}_a(f) := \operatorname{ord}(f_a).$$

For a meromorphic 1-form ω defined on \mathbb{P}^1 , the order $\operatorname{ord}_a(\omega)$ can be defined as follows. Set $U_1 = \mathbb{P}^1 \setminus \{\infty\}$ and $U_2 = \mathbb{P}^1 \setminus \{0\}$. Let z_i be coordinates of U_i , i = 1, 2, such that $z_1(0) = z_2(\infty) = 0$ and $z_2 = 1/z_1$ in $U_1 \cap U_2$. Then there exist meromorphic functions f_i on U_i such that

$$\omega = f_i \, dz_i$$

on U_i for i = 1, 2. Then define

$$\operatorname{ord}_a(\omega) := \operatorname{ord}_a(f_i)$$

for $a \in U_i, i = 1, 2$.

Let us fix a collection of points $a_0, \ldots, a_p \in \mathbb{P}^1$ and set $S := k_0 a_0 + \cdots + k_p a_p$ as an effective divisor with $k_0, \ldots, k_p > 0$. For $a \in \mathbb{P}^1$ let S(a) be the coefficient of a in S, i.e.,

$$S(a) := \begin{cases} k_i & \text{if } a = a_i \text{ for } i = 0, \dots, p, \\ 0 & \text{otherwise.} \end{cases}$$

For an open set $U \subset \mathbb{P}^1$ we define $\Omega_S(U)$ to be the set of all meromorphic 1-forms ω on U satisfying $\operatorname{ord}_a(\omega) \geq -S(a)$ for any $a \in U$. This correspondence defines the sheaf Ω_S by the natural restriction mappings.

Let \mathcal{E} be a locally free sheaf of rank n on \mathbb{P}^1 , namely a sheaf of modules over the sheaf \mathcal{O} of holomorphic functions on \mathbb{P}^1 satisfying that for any $a \in \mathbb{P}^1$ there exists an open neighbourhood $V \subset \mathbb{P}^1$ such that $\mathcal{E}|_V \cong \mathcal{O}^n|_V$. We may sometimes regard \mathcal{E} as a holomorphic vector bundle over \mathbb{P}^1 .

Definition 1.6 (meromorphic connection). A meromorphic connection is a pair (\mathcal{E}, ∇) of a locally free sheaf \mathcal{E} and a morphism $\nabla \colon \mathcal{E} \to \mathcal{E} \otimes \Omega_S$ of sheaves of \mathbb{C} -vector spaces satisfying

$$\nabla(fs) = df \otimes s + f \otimes \nabla(s)$$

for all $f \in \mathcal{O}(U)$, $s \in \mathcal{E}(U)$ and open subsets $U \subset \mathbb{P}^1$.

Let $U \subset \mathbb{P}^1$ be an open subset which gives a local trivialization of \mathcal{E} and z a local coordinate of U. Then if we fix an identification $\mathcal{E}|_U \cong \mathcal{O}^n|_U$, we can write $\nabla = d - A \, dz$ by $A \in M(n, \mathcal{M}(U))$ on U. Note that if we write $\nabla = d - A' \, dz$ by another identification $\mathcal{E}|_U \cong \mathcal{O}^n|_U$, then A' can be obtained by a holomorphic gauge transformation of A, namely there exists $X \in \mathrm{GL}(n, \mathcal{O}(U))$ such that

$$A' = X[A].$$

Thus we may say that (\mathcal{E}, ∇) defines a holomorphic gauge equivalent class of a local differential equation

$$\frac{d}{dz}Y = AY$$

on $U \subset \mathbb{P}^1$.

In particular, suppose that \mathcal{E} is *trivial*, i.e., $\mathcal{E} \cong \mathcal{O}^n$ and set $U_1 = \mathbb{P}^1 \setminus \{\infty\}$ and $U_2 = \mathbb{P}^1 \setminus \{0\}$ as before. Then if we fix a trivialization $\mathcal{E} \cong \mathcal{O}^n$, we have $\nabla = d - A(z_1)dz_1$ on U_1 with $A(z_1) = (\alpha_{i,j}(z_1))_{i,j=1,\dots,n} \in M(n, \mathbb{C}(z))$ satisfying $\operatorname{ord}_a(\alpha_{i,j}) \geq -S(a)$ for all $a \in U_1$. Similarly on U_2 we have $\nabla = d - B(z_2)dz_2$. Since \mathcal{E} is trivial,

$$A(z_1)dz_1 = B(z_2)dz_2$$
 on $U_1 \cap U_2$.

Namely,

$$B(z_2) = -\frac{A(1/z_2)}{z_2^2}.$$

This is nothing but the coordinate exchange $\zeta = \frac{1}{2}$ for a differential equation

$$\frac{d}{dz}Y = A(z)Y \longmapsto -\zeta^2 \frac{d}{d\zeta}Y = A(1/\zeta)Y.$$

Thus a meromorphic connection (\mathcal{E}, ∇) with a trivial bundle \mathcal{E} on \mathbb{P}^1 corresponds to a meromorphic differential equation $\frac{d}{dz}Y = AY$ with $A = (\alpha_{i,j})_{i,j=1,\dots,n} \in M(n, \mathbb{C}(z))$ satisfying $\operatorname{ord}_a(\alpha_{i,j} dz) \geq -S(a)$ for all $a \in \mathbb{P}^1$, and vice versa. This correspondence is unique up to the choice of $\mathcal{E} \cong \mathcal{O}^n$, i.e., $\operatorname{GL}(n, \mathbb{C})$ -action.

Let $S = k_0 a_0 + \ldots + k_p a_p$ be an effective divisor on \mathbb{P}^1 as before. Define a set of meromorphic connections on \mathbb{P}^1

$$\operatorname{Triv}_{S}^{(n)} := \left\{ (\mathcal{E}, \nabla) \middle| \begin{array}{c} \mathcal{E} : \text{ trivial of rank } n, \\ \nabla \colon \mathcal{E} \to \mathcal{E} \otimes \Omega_{S} \end{array} \right\}.$$

We say $(\mathcal{E}, \nabla) \in \operatorname{Triv}_{S}^{(n)}$ is *stable* if there exists no nontrivial proper subspace $W \subset \mathbb{C}^{n}$ such that the subbundle $\mathcal{W} := W \otimes \mathcal{O} \subset \mathbb{C}^{n} \otimes \mathcal{O} = \mathcal{E}$ is closed under ∇ , i.e.,

$$\nabla(\mathcal{W}) \subset \mathcal{W} \otimes \Omega_S.$$

Let $\mathbf{B} = (B_0, \ldots, B_p) \in M(n, \mathbb{C}((z_i)))^{p+1}$ be a collection of HTL normal forms satisfying $\operatorname{ord}(B_i) = -k_i$ for all $i = 0, \ldots, p$. We write $\nabla|_{a_i} \in \mathcal{O}_{B_i}$ for a connection (\mathcal{E}, ∇) if there exists $A_{a_i} \in M(n, \mathbb{C}((z_{a_i})))$ such that $\nabla = d - A_{a_i} dz_{a_i}$ and $\iota(A_{a_i}) \in \mathcal{O}_{B_i}$ where z_{a_i} is a local coordinate of \mathbb{P}^1 vanishing at a_i and $\iota: M(n, \mathbb{C}((z_{a_i}))) \to M(n, \mathbb{C}((z_{a_i}))/\mathbb{C}[[z_{a_i}]])$ is the natural projection.

Then the moduli space of stable meromorphic connections on trivial bundles is

$$\mathfrak{M}(\mathbf{B}) := \left\{ (\mathcal{E}, \nabla) \in \operatorname{Triv}_{S}^{(n)} \middle| \begin{array}{c} (\mathcal{E}, \nabla) \colon \text{ stable,} \\ \nabla|_{a_{i}} \in \mathcal{O}_{B_{i}} \text{ for all } i = 0, \dots, p \end{array} \right\} / \operatorname{GL}(n, \mathbb{C}).$$

Here $\operatorname{GL}(n,\mathbb{C}) = \operatorname{GL}(n,\mathcal{O}(\mathbb{P}^1))$ acts on $\operatorname{Triv}_S^{(n)}$ as the holomorphic gauge transformation.

Möbius transformation may allow us to suppose $a_0 = \infty \in \mathbb{P}^1$. Then by a trivialization $\mathcal{E} \cong \mathcal{O}^n$ we can identify $(\mathcal{E}, \nabla) \in \operatorname{Triv}^n_S$ with a meromorphic differential equation defined on \mathbb{P}^1 ,

$$\frac{d}{dz}Y = \left(\sum_{i=1}^{p}\sum_{\nu=1}^{k_i}\frac{A_{\nu}^{(i)}}{(z-a_i)^{\nu}} + \sum_{2 \le \nu \le k_0}A_{\nu}^{(0)}z^{\nu-2}\right)Y$$

up to $GL(n, \mathbb{C})$ -action, i.e.,

$$\frac{d}{dz}Y = A(z)Y \longmapsto \frac{d}{dz}Y' = gA(z)g^{-1}Y' \quad (g \in \mathrm{GL}(n,\mathbb{C})).$$

The stability of (\mathcal{E}, ∇) corresponds to the irreducibility of the differential equation. Thus we can regard $\mathfrak{M}(\mathbf{B})$ as a moduli space of meromorphic differential equations on \mathbb{P}^1 ,

$$\mathfrak{M}(\mathbf{B}) = \left\{ \frac{d}{dz} Y = \left(\sum_{i=1}^{p} \sum_{\nu=1}^{k_i} \frac{A_{\nu}^{(i)}}{(z-a_i)^{\nu}} + \sum_{2 \le \nu \le k_0} A_{\nu}^{(0)} z^{\nu-2} \right) Y \\ \left| \begin{array}{c} \text{irreducible,} \\ \sum_{\nu=1}^{k_i} \frac{A_{\nu}^{(i)}}{z^{\nu}} \in \mathcal{O}_{B_i}, i = 0, \dots, p \end{array} \right\} / \operatorname{GL}(n, \mathbb{C}).$$

Thus the solvability of the additive Deligne-Simpson problem is rephrased as the non-emptiness of the moduli space.

Proposition 1.7. There is a solution of the additive Delinge-Simpson problem for $\mathbf{B} = (B_0, \ldots, B_p)$ if and only if $\mathfrak{M}(\mathbf{B}) \neq \emptyset$.

Furthermore, forgetting the location of the singular points, we may regard $\mathfrak{M}(\mathbf{B})$ as a subspace of the orbit space $\prod_{i=0}^{p} \mathcal{O}_{B_i}$,

$$\mathfrak{M}(\mathbf{B}) = \left\{ \mathbf{A} = (A_i(z))_{0 \le i \le p} \in \prod_{i=0}^p \mathcal{O}_{B_i} \middle| \begin{array}{c} \mathbf{A} \text{ is irreducible }, \\ \sum_{i=0}^p \operatorname{Res}\left(A_i(z)\right) = 0 \end{array} \right\} \middle/ \operatorname{GL}(n, \mathbb{C})$$

which is free from locations of a_i in \mathbb{P}^1 . Here

$$\operatorname{Res}\left(\sum_{j=1}^{k} A_j z^{-j}\right) := A_1$$

and we say that $\mathbf{A} = (\sum_{j=1}^{k_i} A_{i,j} z^{-1})_{0 \le i \le p}$ is *irreducible* if $(A_{i,j})_{\substack{0 \le i \le p \\ 1 \le j \le k_i}}$ is irreducible.

2. A REVIEW OF REPRESENTATIONS OF QUIVERS

This section is a review of known results of the representation theory of quivers and theory of quiver varieties. The review will be minimized only for our requirement for the latter sections and we refer original papers for the general theory by Nakajima [29], Crawley-Boevey and Holland [10], Crawley-Boevey [8] and their references.

2.1. Representations of quivers and quiver varieties. Here we recall the definition of representations of quivers and introduce quiver varieties.

Definition 2.1 (quivers). A quiver $Q = (Q_0, Q_1, s, t)$ is the quadruple consisting of Q_0 , the set of *vertices*, and Q_1 , the set of *arrows* connecting vertices in Q_0 , and two maps $s, t : Q_1 \to Q_0$, which associate to each arrow $\rho \in Q_1$ its source $s(\rho) \in \mathbb{Q}_0$ and its target $t(\rho) \in \mathbb{Q}_0$ respectively.

Definition 2.2 (representations of quivers). Let Q be a finite quiver, i.e., Q_0 and Q_1 are finite sets. A representation M of Q is defined by the following data:

- (1) To each vertex a in Q_0 , a finite dimensional \mathbb{C} vector space M_a is attached.
- (2) To each arrow $\rho: a \to b$ in Q_1 , a \mathbb{C} -linear map $\psi_{\rho}: M_a \to M_b$ is attached.

We denote the representation by $M = (M_a, \psi_\alpha)_{a \in Q_0, \alpha \in Q_1}$. The collection of integers defined by $\dim M = (\dim_{\mathbb{C}} M_a)_{a \in Q_0}$ is called the *dimension vector* of M.

For a fixed vector $\alpha \in (\mathbb{Z}_{\geq 0})^{\mathbb{Q}_0}$, the representation space is

$$\operatorname{Rep}(\mathsf{Q}, V, \alpha) = \bigoplus_{\rho \in \mathsf{Q}_1} \operatorname{Hom}_{\mathbb{C}}(V_{s(\rho)}, V_{t(\rho)}),$$

where $V = (V_a)_{a \in Q_0}$ is a collection of finite dimensional \mathbb{C} -vector spaces with $\dim_{\mathbb{C}} V_a = \alpha_a$. If $V_a = \mathbb{C}^{\alpha_a}$ for all $a \in Q_0$, we simply write

$$\operatorname{Rep}\left(\mathsf{Q},\alpha\right) = \bigoplus_{\rho \in \mathsf{Q}_1} \operatorname{Hom}_{\mathbb{C}}(\mathbb{C}^{\alpha_{s(\rho)}}, \mathbb{C}^{\alpha_{t(\rho)}}).$$

The representation space $\operatorname{Rep}(\mathsf{Q}, V, \alpha)$ has an action of $\prod_{a \in \mathsf{Q}_0} \operatorname{GL}(V_a)$. For $(\psi_{\rho})_{\rho \in \mathbf{Q}_1} \in \operatorname{Rep}(\mathbf{Q}, V, \alpha)$ and $g = (g_a) \in \prod_{a \in \mathbf{Q}_0} \operatorname{GL}(V_a)$, then $g \in \mathbf{Q}_0$ $(\psi_{\rho})_{\rho\in\mathbf{Q}_1}\in\operatorname{Rep}\left(\mathbf{Q},V,\alpha\right) \text{ consists of } \psi'_{\rho}=g_{t(\rho)}\psi_{\rho}g_{s(\rho)}^{-1}\in\operatorname{Hom}_{\mathbb{C}}(V_{s(\rho)},V_{t(\rho)}).$

Let $M = (M_a, \psi_{\rho}^M)_{a \in Q_0, \rho \in Q_1}$ and $N = (N_a, \psi_{\rho}^N)_{a \in Q_0, \rho \in Q_1}$ be representa-tions of a quiver Q. Then N is called the *subrepresentation* of M if we have the following:

(1) For each $a \in Q_0$, $N_a \subset M_a$. (2) For each $\rho: a \to b \in Q_1$, $\psi_{\rho}^M|_{N_a} = \psi_{\rho}^N$.

In this case we denote $N \subset M$. Moreover if

- (3) there exists a direct sum decomposition $M_a = N_a \oplus N'_a$ for each $a \in Q_0$,
- (4) for each $\rho: a \to b \in Q_1$, we have $\psi_{\rho}^M|_{N'_a} \subset N'_b$,

then we say M has a direct sum decomposition $M = N \oplus N'$ where $N' = (N'_a, \psi^M_\rho|_{N'_a})_{a \in \mathbb{Q}_0, \rho \in \mathbb{Q}_1}$.

The representation M is said to be *irreducible* if M has no subrepresentations other than M and $\{0\}$. Here $\{0\}$ is the representation of \mathbb{Q} which consists of zero vector spaces and zero linear maps. On the other hand if any direct sum decomposition $M = N \oplus N'$ satisfies either $N = \{0\}$ or $N' = \{0\}$, then M is said to be *indecomposable*.

Let us recall the double of a quiver $\mathsf{Q}.$

Definition 2.3 (double of a quiver). Let $\mathbf{Q} = (\mathbf{Q}_0, \mathbf{Q}_1)$ be a finite quiver. Then the *double quiver* $\overline{\mathbf{Q}}$ of \mathbf{Q} is the quiver obtained by adjoining the reverse arrow $\rho^* \colon b \to a$ to each arrow $\rho \colon a \to b$. Namely $\overline{\mathbf{Q}} = (\overline{\mathbf{Q}}_0 = \mathbf{Q}_0, \overline{\mathbf{Q}}_1 = \mathbf{Q}_1 \cup \mathbf{Q}_1^*)$ where $\mathbf{Q}_1^* = \{\rho^* \colon t(\rho) \to s(\rho) \mid \rho \in \mathbf{Q}_1\}$.

Here we note that the representation space $\operatorname{Rep}(\overline{\mathbb{Q}}, \alpha)$ of the double quiver $\overline{\mathbb{Q}}$ can be regarded as the cotangent bundle of $\operatorname{Rep}(\mathbb{Q}, \alpha)$, namely

$$\operatorname{Rep}(\mathsf{Q},\alpha) \cong \operatorname{Rep}(\mathsf{Q},\alpha) \oplus \operatorname{Rep}(\mathsf{Q},\alpha)^* \cong T^* \operatorname{Rep}(\mathsf{Q},\alpha),$$

since we have the identification

$$\operatorname{Hom}_{\mathbb{C}}(\mathbb{C}^{\alpha_{s(\rho)}},\mathbb{C}^{\alpha_{t(\rho)}})^* \cong \operatorname{Hom}_{\mathbb{C}}(\mathbb{C}^{\alpha_{s(\rho^*)}},\mathbb{C}^{\alpha_{t(\rho^*)}})$$

for each $\rho \in Q_1$. Then we can regard $\operatorname{Rep}(\overline{Q}, \alpha) \cong T^*\operatorname{Rep}(Q, \alpha)$ as a symplectic manifold with the canonical symplectic form

$$\omega(x,y) := \sum_{\rho \in \mathsf{Q}_1} (\operatorname{tr}(x_\rho y_{\rho^*}) - \operatorname{tr}(x_{\rho^*} y_\rho))$$

for $x, y \in T^* \operatorname{Rep}(\mathsf{Q}, \alpha)$, which is invariant under the action of

$$\mathbf{G}(\alpha) := \prod_{a \in \mathsf{Q}_0} \mathrm{GL}(\alpha_a, \mathbb{C}).$$

Then we define a *moment map* of the symplectic manifold $\operatorname{Rep}(\overline{\mathbb{Q}}, \alpha)$ with the $\mathbf{G}(\alpha)$ -action as follows. The map

$$\mu_{\alpha} \colon \operatorname{Rep}(\overline{\mathbb{Q}}, \alpha) \to \operatorname{Lie} \mathbf{G}(\alpha) := \prod_{a \in \mathbb{Q}_0} M(\alpha_a, \mathbb{C})$$

is defined by

$$\mu_{\alpha}(x)_{a} = \sum_{\substack{\rho \in \mathsf{Q}_{1} \\ t(\rho) = a}} x_{\rho} x_{\rho^{*}} - \sum_{\substack{\rho \in \mathsf{Q}_{1} \\ s(\rho) = a}} x_{\rho^{*}} x_{\rho},$$

for $x = (x_{\rho})_{\rho \in \overline{\mathbf{Q}}_1} \in \operatorname{Rep}(\overline{\mathbf{Q}}, \alpha).$

Then the quiver variety is defined as the symplectic reduction of $\operatorname{Rep}(\overline{\mathbb{Q}}, \alpha)$ by the moment map μ .

Definition 2.4 (quiver variety). Let us take a collection of complex numbers $\lambda = (\lambda_a)_{a \in Q_0} \in \mathbb{C}^{Q_0}$ and regard $\lambda = (\lambda_a I_{\alpha_a})_{a \in Q_0} \in \prod_{a \in Q_0} M(\alpha_a, \mathbb{C})$. Then the quiver variety is the symplectic reduction

$$\mathfrak{M}_{\lambda}(\mathsf{Q},\alpha) := \mu_{\alpha}^{-1}(\lambda)/\mathbf{G}(\alpha).$$

Note that the symplectic reduction $\mathfrak{M}_{\lambda}(\mathbf{Q}, \alpha)$ is homeomorphic to the affine quotient scheme $\mu_{\alpha}^{-1}(\lambda)//\mathbf{G}(\alpha)$ by the theory of Kempf-Ness [19] and Kirwan [21].

This variety might have singularities. Thus let us consider the (possibly empty) subspace

$$\mu_{\alpha}^{-1}(\lambda)^{\operatorname{irr}} := \{ x \in \mu_{\alpha}^{-1}(\lambda) \mid x \text{ is irreducible} \}.$$

Then the action of $\mathbf{G}(\alpha)/\mathbb{C}^{\times}$ on this space is proper and moreover free (see King [20]). Thus the homogeneous space

$$\mathfrak{M}_{\lambda}^{\mathrm{reg}}(\mathsf{Q},\alpha) := \mu_{\alpha}^{-1}(\lambda)^{\mathrm{irr}}/\mathbf{G}(\alpha)$$

can be seen as a complex manifold with the symlpectic structure, i.e., a complex symplectic manifold. We call this manifold a quiver variety too.

Remark 2.5. The above quiver varieties are special ones of Nakajima quiver varieties which enjoy rich geometric properties and applications for representation theory and theoretical physics and so on (see [29] for instance).

2.2. Crawley-Boevey's theorems for the geometry of quiver varieties. The regular part $\mathfrak{M}_{\lambda}^{\mathrm{reg}}(\mathsf{Q},\alpha)$ may be empty as we noted above. Thus we recall a necessary and sufficient condition for the non-emptiness of $\mathfrak{M}_{\lambda}^{\mathrm{reg}}(\mathsf{Q},\alpha)$ given by Crawley-Boevey in [8].

First let us introduce the root system of a quiver Q (cf. [18]). Let Q be a finite quiver. From the *Euler form*

$$\langle \alpha, \beta \rangle := \sum_{a \in \mathsf{Q}_0} \alpha_a \beta_a - \sum_{\rho \in \mathsf{Q}_1} \alpha_{s(\rho)} \beta_{t(\rho)}$$

a symmetric bilinear form and quadratic form are defined by

$$\begin{aligned} (\alpha,\beta) &:= \langle \alpha,\beta \rangle + \langle \beta,\alpha \rangle, \\ q(\alpha) &:= \frac{1}{2}(\alpha,\alpha) \end{aligned}$$

and set $p(\alpha) := 1 - q(\alpha)$. Here $\alpha, \beta \in \mathbb{Z}^{\mathbf{Q}_0}$.

For each vertex $a \in Q_0$, define $\epsilon_a \in \mathbb{Z}^{Q_0}$ $(a \in Q_0)$ so that $(\epsilon_a)_a = 1$, $(\epsilon_a)_b = 0$, $(b \in Q_0 \setminus \{a\})$. We call ϵ_a a fundamental root if the vertex a has no edge-loop, i.e., there is no arrow ρ such that $s(\rho) = t(\rho) = a$. Denote by Π the set of fundamental roots. For a fundamental root ϵ_a , define the fundamental reflection s_a by

$$s_a(\alpha) := \alpha - (\alpha, \epsilon_a) \epsilon_a \text{ for } \alpha \in \mathbb{Z}^{\mathbf{Q}_0}.$$

The group $W \subset \operatorname{Aut} \mathbb{Z}^{\mathsf{Q}_0}$ generated by all fundamental reflections is called *Weyl group* of the quiver Q . Note that the bilinear form (,) is *W*-invariant. Similarly we can define the reflection $r_a \colon \mathbb{C}^{\mathsf{Q}_0} \to \mathbb{C}^{\mathsf{Q}_0}$ by

$$r_a(\lambda)_b := \lambda_b - (\epsilon_a, \epsilon_b)\lambda_a$$

for $\lambda \in \mathbb{C}^{Q_0}$ and $a, b \in Q_0$. Define the set of *real roots* by

$$\Delta^{\mathrm{re}} := \bigcup_{w \in W} w(\Pi).$$

For an element $\alpha = (\alpha_a)_{a \in \mathbf{Q}_0} \in \mathbb{Z}^{\mathbf{Q}_0}$ the support of α is the set of ϵ_a such that $\alpha_a \neq 0$, and denoted by supp (α). We say the support of α is connected

if the subquiver consisting of the set of vertices a satisfying $\epsilon_a \in \text{supp}(\alpha)$ and all arrows joining these vertices, is connected. Define the fundamental set $F \subset \mathbb{Z}^{\mathsf{Q}_0}$ by

F := $\left\{ \alpha \in (\mathbb{Z}_{\geq 0})^{\mathbb{Q}_0} \setminus \{0\} \mid (\alpha, \epsilon) \leq 0 \text{ for all } \epsilon \in \Pi, \text{ support of } \alpha \text{ is connected} \right\}.$

Then define the set of *imaginary roots* by

$$\Delta^{\operatorname{im}} := \bigcup_{w \in W} w(F \cup -F).$$

Then the *root system* is

$$\Delta = \Delta^{\rm re} \cup \Delta^{\rm im}.$$

Elements in $\Delta^+ := \Delta \cap (\mathbb{Z}_{\geq 0})^{\mathbf{Q}_0}$ are called *positive roots*.

Now we are ready to see Crawley-Boevey's theorem. For a fixed $\lambda =$ $(\lambda_a) \in \mathbb{C}^{\mathsf{Q}_0}$, the set Σ_{λ} consists of the positive roots satisfying

- (1) $\lambda \cdot \alpha = \sum_{a \in \mathbb{Q}_0} \lambda_a \alpha_a = 0,$ (2) if there exists a decomposition $\alpha = \beta_1 + \beta_2 + \dots + \beta_r$ $(r \ge 2)$, with $\beta_i \in \Delta^+$ and $\lambda \cdot \beta_i = 0$, then $p(\alpha) > p(\beta_1) + p(\beta_2) + \dots + p(\beta_r).$

Theorem 2.6 (Crawley-Boevey. Theorem 1.2 in [8]). Let Q be a finite quiver and $\overline{\mathbf{Q}}$ the double of \mathbf{Q} . Let us fix a dimension vector $\alpha \in (\mathbb{Z}_{\geq 0})^{\mathbf{Q}_0}$ and $\lambda \in \mathbb{C}^{\mathbf{Q}_0}$. Then $\mu_{\alpha}^{-1}(\lambda)^{irr} \neq \emptyset$ if and only if $\alpha \in \Sigma_{\lambda}$. Furthermore, in this case $\mu_{\alpha}^{-1}(\lambda)$ is an irreducible algebraic variety and $\mu_{\alpha}^{-1}(\lambda)^{irr}$ is dense in $\mu_{\alpha}^{-1}(\lambda).$

This provides the following geometric properties of quiver varieties.

Theorem 2.7 (Crawley-Boevey Corollary 1.4 in [8]). If $\alpha \in \sigma_{\lambda}$ then the quiver variety $\mathfrak{M}_{\lambda}(\mathbf{Q},\alpha)$ is reduced and irreducible algebraic variety of dimension $2p(\alpha)$.

Combining these results, we have the following non-emptiness condition of regular parts of quiver varieties.

Corollary 2.8 (Crawley-Boevey [8]). The regular part of quiver variety $\mathfrak{M}_{\lambda}^{reg}(\mathbf{Q},\alpha)$ is non-empty if and only if $\alpha \in \Sigma_{\lambda}$. Furthermore in this case, it is a connected symplectic complex manifold of dimension $2p(\alpha)$.

3. A review of Fuchsian cases

The necessary and sufficient condition for the existence of a solution of the additive Deligne-Simpson problem for Fuchsian differential equations is determined by Crawley-Boevey in [9]. The strategy is as follows. For the additive Deligne-Simpson problem for $\mathbf{C} = (C_0, C_1, \dots, C_p)$, a collection of conjugacy classes in $M(n, \mathbb{C})$, it is shown that there exists a quiver \mathbb{Q} , dimension vector α , and complex parameter λ such that the quiver variety $\mathfrak{M}^{\mathrm{reg}}_{\lambda}(\mathsf{Q},\alpha)$ is isomorphic to the moduli space $\mathfrak{M}(\mathbf{C})$. Thus Theorem 2.6 determines the non-emptiness condition of $\mathfrak{M}(\mathbf{C})$ which is equivalent to the solvability of the additive Deligne-Simpson problem. We shall recall this correspondence between $\mathfrak{M}(\mathbf{C})$ and $\mathfrak{M}_{\lambda}^{reg}(\mathbf{Q}, \alpha)$.

First we construct a representation of a quiver from a conjugacy class C of $M(n, \mathbb{C})$. Let us choose complex numbers ξ_1, \ldots, ξ_d so that

(1)
$$\prod_{i=1}^{d} (A - \xi_i I_n) = 0$$

for all $A \in C$. The minimal polynomial of C is an example of this equation. Set $m_k := \operatorname{rank} \prod_{i=1}^k (A - \xi_i I_n)$ for $k = 1, \ldots, d$. Then let us note that these ξ_1, \ldots, ξ_d and m_1, \ldots, m_d characterize C. Namely $B \in M(n, \mathbb{C})$ is contained in C if and only if B satisfies

$$\operatorname{rank} \prod_{i=1}^{k} (B - \xi_i I_n) = m_k$$

for all k = 1, ..., d. This observation leads us to the following correspondence between the elements in C and some representations of a quiver.

Proposition 3.1 (see Crawley-Boevey [9] and also Lemma A.5 in [16]). Let us fix a conjugacy class C of $M(n, \mathbb{C})$ and choose $\xi_1, \ldots, \xi_d \in \mathbb{C}$ so that the equation (1) holds for all $A \in C$. Set $m_k := \operatorname{rank} \prod_{i=1}^k (A - \xi_i I_n)$ for $k = 1, \ldots, d - 1$ and $A \in C$, also set $m_0 := n$ and $\mathbf{m} := (m_i)_{i=0,\ldots,d-1}$. Define a quiver \mathbb{Q} as below.

$$\underbrace{\bigcirc}_{0}^{\rho_1} \underbrace{\bigcirc}_{1}^{\rho_2} \underbrace{\frown}_{d-1}^{\rho_{d-1}} \underbrace{\bigcirc}_{d-1} \underbrace{\bigcirc}_{d-1}^{\rho_{d-1}} \underbrace{\bigcirc}_{d-1}^{\rho_{d-1}} \underbrace{\bigcirc}_{d-1}^{\rho_{d-1}} \underbrace{\frown}_{d-1}^{\rho_{d-1}} \underbrace{\frown}_{d$$

Also define a subspace of $\operatorname{Rep}(\overline{\mathsf{Q}},\mathbf{m})$ by

$$Z := \left\{ x = (x_{\rho}) \in \operatorname{Rep}\left(\overline{\mathsf{Q}}, \mathbf{m}\right) \middle| \\ \mu_{\mathbf{m}}(x)_{i} = (\xi_{i} - \xi_{i+1})I_{m_{i}} \text{ for all } i = 1, \dots, d-1, \\ x_{\rho} : \text{ injective, } x_{\rho^{*}} : \text{ surjective for all } \rho \in \mathsf{Q}_{1}, \rho^{*} \in \mathsf{Q}_{1}^{*} \right\}$$

Then

$$\Phi_{\xi} \colon \{A \in C\} \longrightarrow Z/\prod_{i=1}^{d-1} \operatorname{GL}(m_i, \mathbb{C})$$

defined below is bijective. For $A \in C$, we define $(M_a, \psi_\rho)_{a \in \mathbb{Q}_0, \rho \in \overline{\mathbb{Q}}_1}$, a representation of $\overline{\mathbb{Q}}$ as follows:

$$M_{0} := \mathbb{C}^{n}, \qquad M_{k} := \operatorname{Im} \prod_{i=1}^{k} (A - \xi_{i} I_{n}) \text{ for all } k = 1, \dots, d - 1,$$

$$\psi_{\rho_{i}} : M_{i} \hookrightarrow M_{i-1} : \text{ inclusion}, \qquad \psi_{\rho_{i}^{*}} = (A - \xi_{i})|_{M_{i-1}}.$$

Then $\Phi_{\xi}(A)$ is the projection of (M_a, ψ_{ρ}) . The inverse map is given by

$$(x_{\rho})_{\rho\in\overline{\mathbf{Q}}_{1}}\mapsto x_{\rho_{1}}x_{\rho_{1}^{*}}+\xi_{1}.$$

Furthermore for any $x = (x_{\rho})_{\rho \in \overline{\mathbb{Q}}_1} \in \mathbb{Z}$ and any subspace $S \subset \mathbb{C}^n$ invariant under $x_{\rho_1}x_{\rho_1^*} + \xi_1$, there exists a subrepresentation y of x such that x = 0 (resp. N = M) if and only if S = 0 (resp. $S = \mathbb{C}^n$).

Let C_0, C_1, \ldots, C_p be a collection of conjugacy classes in $M(n, \mathbb{C})$ and write $\mathbf{C} := (C_0, C_1, \ldots, C_p)$. As we noted before, a conjugacy class C can be seen as a truncated orbit of an HTL normal form of the case k = 1. Thus

$$\mathfrak{M}(\mathbf{C}) := \left\{ (A_i)_{i=0,1,\dots,p} \in \prod_{i=0}^p C_i \middle| \begin{array}{c} (A_i)_{i=0,\dots,p} \text{ is irreducible,} \\ \sum_{i=0}^p A_i = 0 \end{array} \right\} \middle/ \operatorname{GL}(n,\mathbb{C})$$

is a moduli space of Fuchsian differential equations or equivalently that of meromorphic connections defined in Section 1.2. Crawely-Boevey obtained a realization of $\mathfrak{M}(\mathbf{C})$ as a quiver variety.

Theorem 3.2 (Crawley-Boevey [9]). Let C_0, \ldots, C_p be conjugacy classes of $M(n, \mathbb{C})$. For $i = 0, \ldots, p$, choose $\xi_{[i,1]}, \ldots, \xi_{[i,d_i]} \in \mathbb{C}$ so that

$$\prod_{j=1}^{a_i} (A_i - \xi_{[i,j]} I_n) = 0$$

for all $A_i \in C_i$. Let $\xi = (\{\xi_{[i,1]}, \ldots, \xi_{[i,d_i]}\})_{0 \le i \le p}$ be the collection of ordered sets $\{\xi_{[i,1]}, \ldots, \xi_{[i,d_i]}\}$. Set $m_0 := n$ and $m_{[i,j]} := \operatorname{rank} \prod_{k=1}^j (A_i - \xi_{[i,k]}I_n)$ for $j = 1, \ldots, d_i - 1$. Consider the following quiver \mathbb{Q} .



Define $\alpha = (\alpha_a)_{a \in Q_0} \in (\mathbb{Z}_{\geq 0})^{Q_0}$ by $\alpha_0 := m_0$ and $\alpha_{[i,j]} := m_{[i,j]}$ for $i = 0, \ldots, p, j = 1, \ldots, d_i - 1$. Define $\lambda = (\lambda_a)_{a \in Q_0} \in \mathbb{C}^{Q_0}$ by $\lambda_0 := -\sum_{i=0}^p \xi_{[i,1]}$ and $\lambda_{[i,j]} := \xi_{[i,j]} - \xi_{[i,j+1]}$ for $i = 0, \ldots, p, j = 1, \ldots, d_i - 1$.

Then there exists a bijection

$$\Phi_{\xi} \colon \mathfrak{M}(\mathbf{C}) \longrightarrow \mathfrak{M}_{\lambda}^{reg}(\mathbf{Q}, \alpha).$$

Thus Theorem 2.6 solves additive Deligne-Simpson problem.

Theorem 3.3 (Crawley-Boevey [9]). Let C_0, \ldots, C_p be conjugacy classes of $M(n, \mathbb{C})$. Let us choose the quiver \mathbb{Q} , $\alpha \in (\mathbb{Z}_{\geq 0})^{\mathbb{Q}_0}$ and $\lambda \in \mathbb{C}^{\mathbb{Q}_0}$ as Theorem 3.2. Then the additive Deligne-Simpson problem for C_0, \ldots, C_p has a solution if and only if $\alpha \in \Sigma_{\lambda}$.

4. Moduli spaces of meromorphic connections and quiver varieties

In the previous section, we saw that moduli spaces of Fuchisan differential equations are isomorphic to quiver varieties and moreover the solvability of the additive Delinge-Simpson problem for Fuchsian differential equations is determined through these isomorphisms. In this section, we shall give a generalization of this correspondence. Namely we shall consider a collection of HTL normal forms $\mathbf{B} = (B_0, B_1, \dots, B_p)$, and give a correspondence

between the moduli space $\mathfrak{M}(\mathbf{B})$ and a quiver variety. This is first done by Boalch in [4] when the orders of HTL normal forms $k_i = -\operatorname{ord}(B_i)$ for $i = 0, \ldots, p$ satisfy

$$k_0 \leq 3$$
 and $k_1 = \cdots = k_p = 1$.

This result is generalized for arbitrary k_0 by Yamakawa and the author in [16]. Thus when the number of unramified irregular singular points is at most one, it is already known that there exist isomorphisms between moduli spaces of meromorphic connections and quiver varieties.

However in Introduction of [4] Boalch suggested that moduli spaces of meromorphic connections with more than two unramified irregular singular points might not be isomorphic to quiver varieties and gave an example.

Therefore it may not be expected to obtain isomorphisms between arbitrary $\mathfrak{M}(\mathbf{B})$ and quiver varieties. Based on these previous results, for an arbitrary $\mathfrak{M}(\mathbf{B})$, we shall construct an injective map from $\mathfrak{M}(\mathbf{B})$ into a quiver variety which becomes an isomorphism if and only if the number of unramified irregular singular points are less than or equal to one.

4.1. A preliminary example: Differential equations with poles of order 2 and representations of quivers. As we noted above, the correspondence between the moduli spaces of meromorphic connections and quiver varieties is already known if $k_1 = \cdots = k_p = 1$ and k_0 is an arbitrary positive integer. Before going to general cases, we shall see the first nontrivial case $k_0 = k_1 = \cdots = k_p = 2$.

4.1.1. Splitting lemma and truncated orbits. Let $B \in \mathfrak{g}_2^*$ be an HTL normal form,

$$B = \operatorname{diag} \left(c_1 I_{n_1} z^{-2} + R_1 z^{-1}, \dots, c_m I_{n_m} z^{-2} + R_m z^{-1} \right).$$

Here $R_i \in M(n_i, \mathbb{C})$ and $c_i \in \mathbb{C}$, $i = 0, \ldots, p$ satisfying $c_i \neq c_j$ if $i \neq j$.

Let us put $B_{irr} := \text{diag}(c_1 I_{n_1}, \ldots, c_m I_{n_m})$ and denote by $V(c_i) \subset \mathbb{C}^n$ the eigenspace of B_{irr} for each eigenvalue $c_i, i = 1, \ldots, m$. For each $X \in M(n, \mathbb{C}), X_{i,j}$ denotes the $\text{Hom}_{\mathbb{C}}(V(c_j), V(c_i))$ -component of X.

Then for the $G_2 = \operatorname{GL}(n, \mathbb{C}[\![z]\!]/z^2\mathbb{C}[\![z]\!])$ -orbit of B, denoted by \mathcal{O}_B , we have the following lemma which is a direct consequence of the splitting lemma (see the section 3.2 in [2] or section 2.3 in [16] for example).

Lemma 4.1. Let $B \in \mathfrak{g}_2^*$ be the HTL normal form as above. Then \mathcal{O}_B consists of $A(x) = \sum_{i=1}^2 A_i x^{-i} \in \mathfrak{g}_2^*$ satisfying that there exists $G \in \mathrm{GL}(n, \mathbb{C})$ such that

 $G^{-1}A_2G = B_{irr}$ and $(G^{-1}A_1G)_{i,i} \in C_{R_i}$

where C_{R_i} are conjugacy classes of R_i for i = 1, ..., m. Moreover if $G_1, G_2 \in$ GL (n, \mathbb{C}) satisfy $G_i^{-1}A_2G_i = B_{irr}$, i = 1, 2, then $G_2^{-1}G_1 = \text{diag}(h_1, ..., h_m)$ where $h_i \in \text{GL}(n_i, \mathbb{C})$ for i = 1, ..., m.

From this lemma we have the following one to one correspondence.

(2)
$$\mathcal{O}_B \longrightarrow$$

$$\left\{ (G, A_1) \in \operatorname{GL}(n, \mathbb{C}) \times M(n, \mathbb{C}) \middle| \begin{array}{c} (G^{-1}A_1G)_{i,i} \in C_{R_i} \\ \text{for all } i = 1, \dots, m \end{array} \right\} / \prod_{i=1}^m \operatorname{GL}(n_i, \mathbb{C}).$$

Here
$$\prod_{i=1}^{m} \operatorname{GL}(n_i, \mathbb{C})$$
 acts on $\operatorname{GL}(n, \mathbb{C}) \times M(n, \mathbb{C})$ by

$$\prod_{i=1}^{m} \operatorname{GL}(n_i, \mathbb{C}) \times (\operatorname{GL}(n, \mathbb{C}) \times M(n, \mathbb{C})) \longrightarrow \operatorname{GL}(n, \mathbb{C}) \times M(n, \mathbb{C}) \\
((h_1, \dots, h_m), (G, A)) \longmapsto (G \cdot \operatorname{diag}(h_1, \dots, h_m), A).$$

The inverse map is induced by sending (G, A_1) to $GB_{irr}G^{-1}x^{-2} + A_1x^{-1} \in$ \mathcal{O}_B .

Remark 4.2. Let us recall that $T^*\mathrm{GL}(n,\mathbb{C}) \cong \mathrm{GL}(n,\mathbb{C}) \times M(n,\mathbb{C})$. Then the above correspondence can be seen as a special case of the identification of G_k -orbits of HTL normal forms and a symplectic reductions of the extended orbits given by Boalch (see Lemma 2.3 in [3]).

4.1.2. Moduli spaces without irreducibility and quivers. Under the above observation, let us consider the relation between $\mathfrak{M}(\mathbf{B})$ and a quiver variety. Let $B_0, \ldots, B_p \in \mathfrak{g}_2^*$ be HTL normal forms written by

$$B_{i} = \operatorname{diag}\left(c_{[i,1]}I_{n_{[i,1]}}z^{-2} + R_{[i,1]}z^{-1}, \dots, c_{[i,m_{i}]}I_{n_{[i,m_{i}]}}z^{-2} + R_{[i,m_{i}]}z^{-1}\right).$$

Let $V(c_{[i,j]}) \subset \mathbb{C}^n$ be the eigenspace of $(B_i)_{irr}$ for each eigenvalue $c_{[i,j]}$, $i = 0, \ldots, p, \ j = 1, \ldots, m_i$. Let $X_{[i,j],[i'j']}$ be the $\operatorname{Hom}_{\mathbb{C}}(V(c_{[i',j']}), V(c_{[i,j]}))$ component of $X \in M(n, \mathbb{C})$. We may write $X = (X_{[i,j],[i'j']}) \underset{1 \leq j \leq m_i}{1 \leq j' \leq m_{i'}}$.

First let us consider the moduli space $\overline{\mathfrak{M}(\mathbf{B})}$ without the irreducibility, namely,

$$\overline{\mathfrak{M}(\mathbf{B})} := \left\{ (A_i(z))_{i=0,\dots,p} \in \prod_{i=0}^p \mathcal{O}_{B_i} \left| \sum_{i=0}^p \operatorname{Res} A_i(z) = 0 \right\} / \operatorname{GL}(n,\mathbb{C}) \right\}$$

which is isomorphic to

$$\left\{ (G_i, A_i)_{i=0,\dots,p} \in \prod_{i=0}^p \operatorname{GL}(n, \mathbb{C}) \times M(n, \mathbb{C}) \middle| \begin{array}{l} \text{(i)} \ (G_i^{-1}A_iG_i)_{j,j} \in C_{R_{[i,j]}} \\ \text{for all } i = 0, \dots, p \text{ and} \\ j = 1, \dots, m_i \\ \text{(ii)} \ \sum_{i=0}^p A_i = 0 \\ \text{(iii)} \ G_0 = I_n \end{array} \right\} \\ / \prod_{i=0}^p \prod_{j=1}^{m_i} \operatorname{GL}(n_{[i,j]}, \mathbb{C}) \end{array}\right\}$$

from the above identification (2). Here the condition (ii) comes from the condition $\sum_{i=0}^{p} \text{Res}A_i(z) = 0$ and (iii) comes from taking the quotient under the $\operatorname{GL}(n, \mathbb{C})$ -action. We shall give a realization of $\overline{\mathfrak{M}(\mathbf{B})}$ as a representation space of a quiver as follows.

Step 1.

Let us consider the quiver $Q^{(1)}$ defined as follows. The set of vertices is

$$\mathsf{Q}_0^{(1)} := \{0, \dots, p\}.$$

The set of arrows is

$$\mathsf{Q}_{1}^{(1)} := \left\{ \rho_{[i]}^{[0]} \colon 0 \to i \, \Big| \, i = 1, \dots, p \right\}.$$



Fix a dimension vector $\alpha^{(1)} := (\alpha_i)_{i=0,\dots,p}$ so that $\alpha_i := n$ for all $i = 0,\dots,p$. Then we have a bijection,

$$\begin{cases} (G_i, A_i)_{i=0,\dots,p} \in \prod_{i=0}^p \operatorname{GL}(n, \mathbb{C}) \times M(n, \mathbb{C}) \middle| G_0 = I_n, \sum_{i=0}^p A_i = 0 \\ \\ \begin{cases} x = (x_\rho)_{\rho \in \overline{\mathbb{Q}^{(1)}}_1} \in \operatorname{Rep}(\overline{\mathbb{Q}^{(1)}}, \alpha^{(1)}) \middle| & \begin{array}{c} x_{\rho_{[i]}^{[0]}} \in \operatorname{GL}(n, \mathbb{C}) \\ \text{for all } i = 1, \dots, p \end{cases} \end{cases}$$

by setting $x_{\rho_{[i]}^{[0]}} := G_i^{-1}$ and $x_{(\rho_{[i]}^{[0]})^*} := A_i G_i$ for all $i = 1, \ldots, p$. Let us note that from $x = (x_\rho)$ in the target space, setting

$$G_i := x_{\rho_{[i]}^{[0]}}^{-1}, \qquad \qquad A_i := x_{(\rho_{[i]}^{[0]})^*} x_{\rho_{[i]}^{[0]}}$$

for $i = 1, \ldots, p$ and

$$A_0 := -\sum_{i=1^p} x_{(\rho_{[i]}^{[0]})^*} x_{\rho_{[i]}^{[0]}} = \mu_{\alpha^{(1)}}(x)_0,$$

we obtain the inverse map.

Step 2.

In Step 1, we could associate representations of a quiver to $(G_i, A_i)_{i=0,\ldots,p} \in \prod_{i=0}^{p} \operatorname{GL}(n, \mathbb{C}) \times M(n, \mathbb{C})$ satisfying the conditions (ii) $\sum_{i=0}^{p} A_i = 0$ and (iii) $G_0 = I$. However to obtain the one to one correspondence with $\overline{\mathfrak{M}}(\mathbf{B})$, we need one more condition (i) $(G_i^{-1}A_iG_i)_{j,j} \in C_{R_{[i,j]}}$ for $i = 0, \ldots, p$, $j = 1, \ldots, m_i$. Let us recall that

$$G_i^{-1}A_iG_i = x_{\rho_{[i]}^{[0]}}x_{(\rho_{[i]}^{[0]})^*} = \mu_{\alpha^{(1)}}(x)_i$$

for $i = 1, \ldots, p$ and

$$G_0^{-1}A_0G_0 = A_0 = -\sum_{i=0}^p A_i = -\sum_{i=1^p} x_{(\rho_{[i]}^{[0]})^*} x_{\rho_{[i]}^{[0]}} = \mu_{\alpha^{(1)}}(x)_0$$

for $(G_i, A_i)_{i=0,...,p}$ in the domain of the isomorphism in Step 1 and its image $x \in \operatorname{Rep}(\overline{\mathbb{Q}^{(1)}}, \alpha^{(1)})$. To obtain block diagonal components $(G_i^{-1}A_iG_i)_{j,j}$ as images of the moment map, we shall break up the vertex

$$\bigcirc [i, 1] \\ \bigcirc [i, 2] \\ \bigcirc \\ i \\ \bigcirc [i, m_i]$$

for each i = 0, ..., p and define the following quiver $Q^{(2)}$.



Namely, the set of vertices is

$$Q^{(2)}_0 := \{ [i, j] \mid i = 0, \dots, p, j = 1, \dots, m_i \}.$$

The set of arrows is

$$\mathbf{Q}^{(2)}_{1} := \left\{ \rho^{[0,j]}_{[i,j']} \colon [0,j] \to [i,j'] \left| \begin{array}{c} j = 1, \dots, m_{0}, \\ i = 1, \dots, p, \\ j' = 1, \dots, m_{i} \end{array} \right\}.$$

Define $\alpha^{(2)} = (\alpha_a^{(2)})_{a \in \mathbf{Q}_0} \in \mathbb{Z}^{\mathbf{Q}_0}$ by $\alpha_{[i,j]}^{(2)} := \dim_{\mathbb{C}} V(c_{[i,j]}), i = 0, \dots, p,$ $j = 1, \dots, m_i$. Then we have a bijection from $\overline{\mathfrak{M}(\mathbf{B})}$ to an open subset of $\operatorname{Rep}(\overline{\mathbf{Q}^{(2)}}, \alpha^{(2)})/\mathbf{G}(\alpha^{(2)}).$

Proposition 4.3. We use the same notation as above. Then there exists a bijection

$$\Phi: \mathfrak{M}(\mathbf{B}) \longrightarrow \left\{ \begin{array}{l} x = (x_{\rho})_{\rho \in \overline{\mathbf{Q}^{(2)}}_{1}} \\ \in \operatorname{Rep}(\overline{\mathbf{Q}^{(2)}}, \alpha^{(2)}) \end{array} \middle| \begin{array}{l} \det \left(x_{\rho_{[i,j']}^{[0,j]}} \right)_{\substack{1 \le j \le m_{0} \\ 1 \le j' \le m_{i}}} \neq 0 \\ for \ all \ i = 1, \dots, p, \\ \mu_{\alpha^{(2)}}(x)_{[i,j]} \in C_{R_{[i,j]}}, \ [i,j] \in \mathbf{Q}^{(2)}_{0} \end{array} \right\} / \mathbf{G}(\alpha^{(2)}).$$

Proof. It suffices to show that there is a bijection from

$$\left\{ (G_i, A_i)_{i=0,\dots,p} \in \prod_{i=0}^p \operatorname{GL}(n, \mathbb{C}) \times M(n, \mathbb{C}) \middle| \begin{array}{l} \text{(i)} \ (G_i^{-1}A_iG_i)_{j,j} \in C_{R_{[i,j]}} \\ \text{for all } i = 0, \dots, p \text{ and} \\ j = 1, \dots, m_i \\ \text{(ii)} \ \sum_{i=0}^p A_i = 0 \\ \text{(iii)} \ G_0 = I_n \end{array} \right\} \\ / \prod_{i=0}^p \prod_{j=1}^{m_i} \operatorname{GL}(n_{[i,j]}, \mathbb{C})$$

to the target space of the above map. Let $(G_i, A_i)_{i=0,\dots,p}$ be a representative of an element in this space. Then we define $x \in \text{Rep}(\overline{\mathsf{Q}^{(2)}}, \alpha^{(2)})$ as follows.

$$x_{\rho_{[i,j']}^{[0,j]}} = (G_i^{-1})_{[i,j'],[0,j]}, \qquad \qquad x_{\left(\rho_{[i,j']}^{[0,j]}\right)^*} = (A_i G_i)_{[0,j],[i,j']},$$

for $j = 1, ..., m_0, i = 1, ..., p, j' = 1, ..., m_i$. Then

$$\mu_{\alpha^{(2)}}(x)_{[i,j']} = \sum_{j=1}^{m_0} x_{\rho_{[i,j']}^{[0,j]}} x_{\left(\rho_{[i,j']}^{[0,j]}\right)^*} = (G_i^{-1}A_iG_i)_{j',j'} \in C_{R_{[i,j']}}$$

for $i = 1, \ldots, p$ and $j' = 1, \ldots, m_i$. Also

$$\mu_{\alpha^{(2)}}(x)_{[0,j]} = -\sum_{i=1}^{p} \sum_{j'=1}^{m_i} x_{\left(\rho_{[i,j']}^{[0,j]}\right)^*} x_{\rho_{[i,j']}^{[0,j]}} = -\sum_{i=1}^{p} (A_i)_{j,j} = (A_0)_{j,j} \in C_{R_{[0,j]}}$$

for $j = 1, ..., m_0$. Since this correspondence is $\prod_{i=0}^p \prod_{j=1}^{m_i} \operatorname{GL}(n_{[i,j]}, \mathbb{C}) \cong \mathbf{G}(\alpha^{(2)})$ -equivariant, we have the well-defined map. The inverse maps can be defined as we saw in Step 1. Thus it is bijective.

4.1.3. Moduli space and quiver varieties. Now we are ready to consider $\mathfrak{M}(\mathbf{B})$. As we will see later, the irreducibility of differential equations does not coincide with the irreducibility of representations of quiver under the bijection in Proposition 4.3. Thus we shall introduce a weaker condition which is called \mathcal{L} -irreducibility in this paper.

Let us define a sublattice $\tilde{\mathcal{L}}$ of $\mathbb{Z}^{Q^{(2)}_0}$ by

$$\tilde{\mathcal{L}} := \left\{ \beta = (\beta_a)_{a \in \mathbf{Q}^{(2)}_0} \in \mathbb{Z}^{\mathbf{Q}^{(2)}_0} \left| \sum_{j=1}^{m_0} \beta_{[0,j]} = \sum_{j=1}^{m_i} \beta_{[i,j]} \text{ for all } i = 1, \dots, p \right\} \right\}$$

Definition 4.4 ($\tilde{\mathcal{L}}$ -irreducible). An element in

$$\left\{ \begin{array}{l} x = (x_{\rho})_{\rho \in \overline{\mathbf{Q}^{(2)}}_{1}} \\ \in \operatorname{Rep}\left(\overline{\mathbf{Q}^{(2)}}, \alpha^{(2)}\right) \end{array} \middle| \begin{array}{l} \det\left(x_{\rho_{[i,j']}^{[0,j]}}\right)_{\substack{1 \le j \le m_{0} \\ 1 \le j' \le m_{i}}} \neq 0 \\ \text{for all } i = 1, \dots, p, \\ \mu_{\alpha^{(2)}}(x)_{[i,j]} \in C_{R_{[i,j]}}, \ [i,j] \in \mathbf{Q}^{(2)}_{0} \end{array} \right\} / \mathbf{G}(\alpha)$$

is said to be $\tilde{\mathcal{L}}$ -*irreducible*, if it has no proper subrepresentation y with the dimension vector $\operatorname{dim}(y) \in \tilde{\mathcal{L}}$ other than $\{0\}$.

Proposition 4.5. Let $\mathbf{A} = (\sum_{j=1}^{2} A_j^{(i)} z^{-j}) \in \prod_{i=0}^{p} \mathcal{O}_{B_i}$ with $\sum_{i=0}^{p} A_1^{(i)} = 0$ and $x \in \operatorname{Rep}(\overline{\mathbf{Q}^{(2)}}, \alpha^{(2)})$ be the corresponding elements under the map Φ in Proposition 4.3. If \mathbf{A} is irreducible, then x is $\tilde{\mathcal{L}}$ -irreducible and vice versa.

Proof. Suppose that **A** has a nontrivial invariant subspace $W \subseteq \mathbb{C}^n$, i.e., W is invariant under all $A_j^{(i)}$. Set $W^{(i)} := (x_{\rho_{[i,j']}^{[0,j]}})_{\substack{1 \leq j \leq m_0 \\ 1 \leq j' \leq m_i}} W \cong W$ for $i = (x_{\rho_{[i,j']}^{[0,j]}})_{\substack{1 \leq j \leq m_0 \\ 1 \leq j' \leq m_i}} W$

 $1, \ldots, p$ and $W^{(0)} := W$. Also set

$$\begin{split} \tilde{V}_{[i,j]} &:= W^{(i)} \cap V(c_{[i,j]}), \qquad \qquad \tilde{x}_{\rho_{[i,j']}^{[0,j]}} = x_{\rho_{[i,j']}^{[0,j]}}|_{W^{(0)}}, \\ \tilde{x}_{\left(\rho_{[i,j']}^{[0,j]}\right)^*} &= x_{\left(\rho_{[i,j']}^{[0,j]}\right)^*}|_{W^{(i)}}. \end{split}$$

Then $\tilde{x} = (\tilde{V}_a, \tilde{x}_\rho)_{a \in \mathbb{Q}^{(2)}_{0,\rho \in \overline{\mathbb{Q}^{(2)}_{1}}}}$ defines a subrepresentation of x. Since W is $A_2^{(i)}$ -invariant, $W^{(i)}$ is $G_i^{-1} A_2^{(i)} G_i = (B_i)_{irr}$ -invariant. Thus we have $W^{(i)} = \bigoplus_{j=1}^{m_i} \tilde{V}_{[i,j]}$, which shows that $\sum_{j=1}^{m_0} \dim_{\mathbb{C}} \tilde{V}_{[0,j]} = \cdots = \sum_{j=1}^{m_p} \dim_{\mathbb{C}} \tilde{V}_{[p,j]}$. Finally we need to check that $\tilde{x}_{(\rho_{[i,j']}^{[0,j]})^*}(\tilde{V}_{[i,j']}) \subset \tilde{V}_{[0,j]}$. To show this, it suffices to see that $(\tilde{x}_{(\rho_{[i,j']}^{[0,j]})^*})_{1 \leq j \leq m_0}^{1 \leq j \leq m_0} W^{(i)} \subset W$, which follows from the fact that

$$\begin{split} (\tilde{x}_{(\rho_{[i,j']}^{[0,j]})^*})_{\substack{1 \le j \le m_0 \\ 1 \le j' \le m_i}} W^{(i)} &= (x_{(\rho_{[i,j']}^{[0,j]})^*})_{\substack{1 \le j \le m_0 \\ 1 \le j' \le m_i}} W^{(i)} \\ &= (x_{(\rho_{[i,j']}^{[0,j]})^*})_{\substack{1 \le j \le m_0 \\ 1 \le j' \le m_i}} (x_{\rho_{[i,j']}^{[0,j]}})_{\substack{1 \le j \le m_0 \\ 1 \le j' \le m_i}} W \\ &= A_1^{(i)} W \subset W. \end{split}$$

Conversely suppose that x has a nontrivial proper subrepresentation $\tilde{x} = (\tilde{V}_a, \tilde{x}_{\rho})$ satisfying $\sum_{j=1}^{m_0} \dim_{\mathbb{C}} \tilde{V}_{[0,j]} = \cdots = \sum_{j=1}^{m_p} \dim_{\mathbb{C}} \tilde{V}_{[p,j]}$. Then $W = \bigoplus_{j=1}^{m^{(0)}} \tilde{V}_{[0,j]}$ is an **A**-invariant subspace. Indeed W is $(A_1^{(0)}, A_2^{(0)})$ -invariant. Also for $i = 1, \ldots, p$, set $W^{(i)} := (x_{\rho_{[i,j']}})_{\substack{1 \leq j \leq m_0 \\ 1 \leq j' \leq m_i}} W \subset \bigoplus_{j=1}^{m_i} \tilde{V}_{[i,j]}$. Then we

have

$$\sum_{j=1}^{m_i} \dim_{\mathbb{C}} \tilde{V}_{[i,j]} = \sum_{j=1}^{m_0} \dim_{\mathbb{C}} \tilde{V}_{[0,j]} = \dim_{\mathbb{C}} W = \dim_{\mathbb{C}} W^{(i)},$$

which implies that $W^{(i)} = \bigoplus_{j=1}^{m_i} \tilde{V}_{[i,j]}$. Thus since

$$(x_{\rho_{[i,j']}^{[0,j]}})_{\substack{1 \le j \le m_0 \\ 1 \le j' \le m_i}} (x_{(\rho_{[i,j']}^{[0,j]})^*})_{\substack{1 \le j \le m_0 \\ 1 \le j' \le m_i}} = G_i^{-1} A_1^{(i)} G_i,$$

 $W^{(i)}$ is $G_i^{-1}A_1^{(i)}G_i$ -invariant, which shows that $W = G_iW^{(i)}$ is $(A_1^{(i)}, A_2^{(i)})$ -invariant for each $i = 1, \ldots, p$.

Finally let us give a realization of $\mathfrak{M}(\mathbf{B})$ as a subset of a quiver variety $\mathfrak{M}_{\lambda}(\mathbf{Q}, \alpha)$ defined as below.

For each $i = 0, \ldots, p$ and $j = 1, \ldots, m_i$, we choose $\xi_{[i,j,k]} \in \mathbb{C}, k = 1, \ldots, e_{[i,j]}$ so that

$$\prod_{k=1}^{e_{[i,j]}} (R_{[i,j]} - \xi_{[i,j,k]} I_{n_{[i,j]}}) = 0.$$

Then the quiver Q is defined by the set of vertices

$$\mathbf{Q}_{0} := \left\{ \begin{bmatrix} i, j, k \end{bmatrix} \middle| \begin{array}{c} i = 0, \dots, p, \\ j = 1, \dots, m_{i}, \\ k = 1, \dots, e_{[i,j]} - 1 \end{array} \right\}$$

and the set of arrows

$$Q_{1} := \left\{ \rho_{[i,j']}^{[0,j]} \colon [0,j] \to [i,j'] \middle| \begin{array}{c} j = 1, \dots, m_{0}, \\ i = 1, \dots, p, \\ j' = 1, \dots, m_{i} \end{array} \right\}$$
$$\bigsqcup \left\{ \rho_{[i,j,k]} \colon [i,j,k] \to [i,j,k-1] \middle| \begin{array}{c} i = 0, \dots, p, \\ j = 1, \dots, m_{i}, \\ k = 1, \dots, e_{[i,j]} - 1 \end{array} \right\}.$$

Here we set [i, j, 0] := [i, j]. Define a dimension vector $\alpha = (\alpha_a)_{a \in Q_0}$ by

$$\alpha_{[i,j]} := n_{[i,j]} \qquad \qquad \alpha_{[i,j,k]} := \operatorname{rank} \prod_{l=1}^{k} (R_{[i,j]} - \xi_{[i,j,l]} I_{n_{[i,j]}}).$$

Also define $\lambda = (\lambda_a)_{a \in \mathbf{Q}_0} \in \mathbb{C}^{\mathbf{Q}_0}$ by

$$\lambda_{[i,j]} := -\xi_{[i,j,1]}$$
 $\lambda_{[i,j,k]} := \xi_{[i,j,k]} - \xi_{[i,j,k+1]}$

where $\xi_{[i,j,e_{[i,j]}]} := 0$. Let us define a sublattice

$$\mathcal{L} := \left\{ \beta = (\beta_a)_{a \in \mathbf{Q}_0} \in \mathbb{Z}^{\mathbf{Q}_0} \left| \sum_{j=1}^{m_0} \beta_{[0,j]} = \sum_{j=1}^{m_i} \beta_{[i,j]} \text{ for all } i = 1, \dots, p \right\} \subset \mathbb{Z}^{\mathbf{Q}_0} \right\}$$

Then we define a subset of the quiver variety $\mathfrak{M}_{\lambda}(\mathsf{Q},\alpha)$ as follows,

$$\mathfrak{M}_{\lambda}(\mathsf{Q},\alpha)^{\mathrm{dif}} := \mu_{\alpha}^{-1}(\lambda)^{\mathrm{dif}}/\mathbf{G}(\alpha)$$

where

$$\mu_{\alpha}^{-1}(\lambda)^{\operatorname{dif}} := \left\{ x \in \mu_{\alpha}^{-1}(\lambda) \left| \begin{array}{c} x \text{ is } \mathcal{L}\text{-irreducible,} \\ \det \left(x_{\rho_{[i,j']}^{[0,j]}} \right)_{1 \leq j \leq m_0} \neq 0, \, i = 1, \dots, p \end{array} \right\}.$$

Here \mathcal{L} -irreducibility is defined as in Definition 4.4. Then from Proposition 3.1, 4.3, and 4.5, we obtain the following identification.

Theorem 4.6. We have a bijection

$$\mathfrak{M}(\mathbf{B}) \longrightarrow \mathfrak{M}_{\lambda}(\mathbf{Q}, \alpha)^{dif}.$$

Proof. By Proposition 3.1, 4.3, and 4.5, it suffices to see that $x \in \mu^{-1}(\lambda)^{\text{dif}}$ implies that $x_{\rho_{[i,j,k]}}$ are injective and $x_{(\rho_{[i,j,k]})^*}$ are surjective. This can be checked similarly to the proof of Theorem 1 in [9]. Indeed, if there exists $x_{\rho_{[i,j,k]}}$ which is not injective, then there exists a nonzero element $v \in \text{Ker}(x_{\rho_{[i,j,k]}})$. Set $v_k := v$ and $v_{l+1} := \psi_{(\rho_{[i,j,l+1]})^*}(v_l)$ for $l \leq k$. Then the relation

$$x_{\rho_{[i,j,l+1]}} x_{\left(\rho_{[i,j,l+1]}\right)^*} - x_{\left(\rho_{[i,j,l]}\right)^*} x_{\rho_{[i,j,l]}} = \lambda_{[i,j,l]}$$

shows that $x_{\rho_{[i,j,l+1]}}(v_{l+1})$ is a multiple of v_l for $l \ge k$. Thus v_l , $l \le k$, span a subrepresentation of x, which contradicts to the \mathcal{L} -irreducibility of x. A dual argument shows that $x_{(\rho_{[i,j,k]})^*}$ are surjective.

Remark 4.7. In the above theorem, we obtain an isomorphism between the moduli space of meromorphic connections $\mathfrak{M}(\mathbf{B})$ and the subset $\mathfrak{M}_{\lambda}(\mathbf{Q}, \alpha)^{\text{dif}}$ of the quiver variety. However we should notice that $\mathfrak{M}_{\lambda}(\mathbf{Q}, \alpha)^{\text{dif}}$ does not coincide with $\mathfrak{M}_{\lambda}^{\text{reg}}(\mathbf{Q}, \alpha)$ since we imposed the det $(x_{\rho_{[i,j']}})_{\substack{1 \leq j \leq m_i \\ 1 \leq j' \leq m_i}} \neq 0$ and

the \mathcal{L} -irreducibility does not coincide with the irreducibility in general. Thus Crawley-Boevey's theorem (see Theorem 2.6) is not applicable directly to our case.

4.2. Truncated orbits and representations of quivers. Let us recall the description of truncated orbits \mathcal{O}_B of arbitrary orders k as quiver varieties. The description which will be given in this section was obtained by Boalch in [4] when $k \leq 3$ and conjectured for arbitrary k. In this section we shall give a quiver picture of truncated orbits of arbitrary orders following the paper by Yamakawa and the author [16] in which the above Boalch's conjecture was finally settled.

Fix k > 1 and $B = \sum_{i=1}^{k} B_i z^{-i} \in \mathfrak{g}_k^*$, an HTL normal form written by

$$B = \operatorname{diag}\left(q_1(z^{-1})I_{n_1} + R_1 z^{-1}, \dots, q_m(z^{-1})I_{n_m} + R_m z^{-1}\right)$$

where $R_i \in M(n_i, \mathbb{C}), q_i(z^{-1}) \in z^{-2}\mathbb{C}[z^{-1}], i = 1, \dots, m \text{ and } q_i \neq q_j \text{ if } i \neq j.$ To a pair $(j, j'), 1 \leq j \neq j' \leq m$, we attach an integer

(3)
$$d(j,j') := \deg_{\mathbb{C}[x]}(q_j(x) - q_{j'}(x)) - 2.$$

Moreover we set d(j,j) := -1 for the latter use.

Let $\bigoplus_{j=1}^{m(s)} V_{\langle s,j \rangle}$ be the decomposition of \mathbb{C}^n as simultaneous invariant spaces of $\{B_{s+1}, B_{s+2}, \ldots, B_k\}$ for $s = 1, \ldots, k-1$. Especially we write $V_j := V_{\langle 1,j \rangle}$ for $j = 1, \ldots, m = m(1)$.

Let $X_{j,j'}$ be the $\operatorname{Hom}_{\mathbb{C}}(V_{j'}, V_j)$ -component of $X \in M(n, \mathbb{C})$. For $g(z) = \sum_{i=r}^{\infty} g_i z^i \in M(n, \mathbb{C}((z)))$, write $(g(z))_{j,j'} := \sum_{i=r}^{\infty} (g_i)_{j,j'} z^i$, $1 \leq j, j' \leq m$. We denote the $\operatorname{Hom}_{\mathbb{C}}(V_j, \mathbb{C}^n)$ -component of $X \in M(n, \mathbb{C})$ by $X_{*,j}$ for each $i = 1, \ldots, m$. Similarly $X_{j,*}$ denote the $\operatorname{Hom}_{\mathbb{C}}(\mathbb{C}^n, V_j)$ - component. We sometimes use the notation $X = (X_{j,j'})_{1 \leq j,j' \leq m} = (X_{*,j'})_{1 \leq j \leq m}$.

Let $\pi_s: J_s = \{1, \ldots, m(s)\} \to J_{s+1} = \{1, \ldots, m(s+1)\}$ be the natural surjection such that $V_{\langle s,j \rangle} \subset V_{\langle s+1,\pi_s(j) \rangle}$. Define the total ordering $\{1 < 2 < \cdots < m\}$ on J_1 and also define total orderings on $J_s, s = 2, \ldots, k-1$, so that

if
$$j_1 < j_2$$
, then $\pi_s(j_1) \le \pi_s(j_2)$, $j_1, j_2 \in J_s$.

Let us define the subgroup of G_k by

$$G_k^o := \left\{ \sum_{i=0}^{k-1} A_i z^i \in G_k \, \middle| \, A_0 = I_n \right\}.$$

Similarly define the subspace $\mathfrak{g}_k^o := z\mathfrak{g}_k = M\left(n, z\mathbb{C}[\![z]\!]/z^k\mathbb{C}[\![z]\!]\right)$ of \mathfrak{g}_k , which can be identified with

$$\left\{\sum_{i=0}^{k-1} A_i z^i \in \mathfrak{g}_k \,\middle|\, A_0 = 0\right\}.$$

Then the dual space $(\mathfrak{g}_k^o)^*$ can be identified with

$$\left\{ \sum_{i=0}^{k-1} A_i z^{-i-1} \in \mathfrak{g}_k^* \, \middle| \, A_0 = 0 \right\}.$$

For $A = \sum_{i=1}^{k} A_i z^{-i} \in \mathfrak{g}_k^*$, set $A_{irr} := \sum_{i=2}^{k} A_i z^{-i}$ and $A_{res} := A_1$. Then we define the following two orbits

$$\mathcal{O}_B^o := \{ g B g^{-1} \mid g \in G_k^o \} \subset \mathfrak{g}_k^*,$$
$$\mathcal{O}_{B_{\operatorname{irr}}} := \{ g B_{\operatorname{irr}} g^{-1} \mid g \in G_k^o \} \subset (\mathfrak{g}_k^o)^*.$$

Let us define the subgroup $H \subset \operatorname{GL}(n, \mathbb{C})$ by

$$H = \{h = \operatorname{diag}(h_1, \dots, h_m) \mid h_i \in \operatorname{GL}(n_i, \mathbb{C}), i = 1, \dots, m\}.$$

The following proposition links \mathcal{O}_B^o with \mathcal{O}_B .

Proposition 4.8 (cf. Lemmas 2.2 and 2.4 in [3]). Set

$$\operatorname{Ad}_{H}(\mathcal{O}_{B}^{o}) := \left\{ hAh^{-1} \in \mathfrak{g}_{k}^{*} \mid h \in H, A \in \mathcal{O}_{B}^{o} \right\}.$$

Then we have a bijection

$$\operatorname{GL}(n,\mathbb{C})\times_H\operatorname{Ad}_H(\mathcal{O}_B^o)\xrightarrow{\sim}\mathcal{O}_B.$$

Here $\operatorname{GL}(n, \mathbb{C}) \times_H \operatorname{Ad}_H(\mathcal{O}_B^o) = (\operatorname{GL}(n, \mathbb{C}) \times \operatorname{Ad}_H(\mathcal{O}_B^o)) / \sim$, the equivalence relation \sim is defined by $(g, A) \sim (gh^{-1}, hAh^{-1})$ for $h \in H$.

Proof. Let us send $(g, A) \in \operatorname{GL}(n, \mathbb{C}) \times_H \operatorname{Ad}_H(\mathcal{O}_B^o)$ to $gAg^{-1} \in \mathcal{O}_B$ and show that this map is well-defined and bijective. If $(g, A) = (g', A') \in$ $\operatorname{GL}(n, \mathbb{C}) \times_H \operatorname{Ad}_H(\mathcal{O}_B^o)$, then there exists $h \in H$ such that $g' = gh^{-1}$ and $A' = hAh^{-1}$. Thus $g'A'(g')^{-1} = gh^{-1}hAh^{-1}hg^{-1} = gAg^{-1}$. Let us see the surjectivity. For any $A' \in \mathcal{O}_B$, there exists $g = g_0 + g_1x + \cdots \in G_k$ such that $g^{-1}A'g = B$. Then $(g_0^{-1}, g_0A'g_0^{-1}) \in \operatorname{GL}(n, \mathbb{C}) \times \operatorname{Ad}_H(\mathcal{O}_B^o)$ and $g_0^{-1}(g_0A'g_0^{-1})g_0 = A'$, which shows the surjectivity. Next we shall see the injectivity. Suppose that $(g, A), (g', A') \in \operatorname{GL}(n, \mathbb{C}) \times_H \operatorname{Ad}_H(\mathcal{O}_B^o)$ are sent to the same element, i.e., $gAg^{-1} = g'A'(g')^{-1}$. Then putting $h = g^{-1}g'$, we have $A' = h^{-1}Ah$. On the other hand, we may assume $A, A' \in \mathcal{O}_B^o$. Thus there exist $b, b' \in G_k^o$ such that $b^{-1}Ab = (b')^{-1}A'b' = B$ which implies $B = b^{-1}Ab = (hb')^{-1}A(hb')$. Lemma 2.9 in [16] shows that the constant term of $b'hb^{-1} \in G_k$ is contained in $\operatorname{Stab}_H(B_1)$, the stabilizer of B_1 in H, i.e., $h \in \operatorname{Stab}_H(B_1) \subset H$. □

According to the ordering on each J_s , $s = 1, \ldots, k - 1$, let us define parabolic subalgebras of $M(n, \mathbb{C})$ as below,

$$\mathfrak{p}(s)^{+} := \bigoplus_{\substack{j_{1}, j_{2} \in J_{i}, \\ j_{1} \ge j_{2}}} \operatorname{Hom}_{\mathbb{C}}(V_{\langle s, j_{1} \rangle}, V_{\langle s, j_{2} \rangle}),$$
$$\mathfrak{p}(s)^{-} := \bigoplus_{\substack{j_{1}, j_{2} \in J_{i}, \\ j_{1} \le j_{2}}} \operatorname{Hom}_{\mathbb{C}}(V_{\langle s, j_{1} \rangle}, V_{\langle s, j_{2} \rangle}),$$

and similarly nilpotent subalgebras

$$\mathfrak{u}(s)^{+} := \bigoplus_{\substack{j_{1}, j_{2} \in J_{i}, \\ j_{1} > j_{2}}} \operatorname{Hom}_{\mathbb{C}}(V_{\langle s, j_{1} \rangle}, V_{\langle s, j_{2} \rangle}),$$
$$\mathfrak{u}(s)^{-} := \bigoplus_{\substack{j_{1}, j_{2} \in J_{i}, \\ j_{1} < j_{2}}} \operatorname{Hom}_{\mathbb{C}}(V_{\langle s, j_{1} \rangle}, V_{\langle s, j_{2} \rangle}),$$

for $s = 1, \ldots, k - 1$. Note that $\mathfrak{p}(s)^{\pm} = \mathfrak{h}(s) \oplus \mathfrak{u}(s)^{\pm}$ where

$$\mathfrak{h}(s) := \bigoplus_{j \in J_i} \operatorname{End}_{\mathbb{C}}(V_{\langle s, j \rangle}).$$

Lemma 4.9. (cf. Lemma 3.4 in [16]) An element $g \in G_k^o$ preserves $B + (\mathfrak{u}_1^+ \oplus \mathfrak{u}_1^-)x^{-1}$, i.e., $g^{-1}bg \in B + (\mathfrak{u}_1^+ \oplus \mathfrak{u}_1^-)x^{-1}$ for all $b \in B + (\mathfrak{u}_1^+ \oplus \mathfrak{u}_1^-)x^{-1}$ if and only if

$$g \in \mathcal{H}_k = \left\{ h := \sum_{i=1}^{k-1} h_i z^i \in G_k^o \, \middle| \, h_i \in \mathfrak{h}_{i+1} \text{ for } i = 1, \dots, k-1 \right\}.$$

Here we put $\mathfrak{h}_k := M(n, \mathbb{C}).$

Proof. We know that \mathcal{H}_k is the stabilizer of B_{irr} in $(\mathfrak{g}_k^o)^*$, see Lemma 3.4 in [16] for example. Thus if $g \in G_k^o$ preserves $B + (\mathfrak{u}_1^+ \oplus \mathfrak{u}_1^-)z^{-1}$, then $g \in \mathcal{H}_k$. Conversely take $g \in \mathcal{H}_k$ and put $u_1 z^{-1} = gBg^{-1} - B \in M(n, \mathbb{C})z^{-1}$. Then $u_1 = B_1 + \sum_{t=2}^k \sum_{s=1}^{t-1} g_s B_t g'_{t-s-1} - B_1$. Here we put $g = I_n + g_1 z + g_2 z^2 + \cdots$ and $g^{-1} = I_n + g'_1 z + g'_2 z^2 + \cdots$. Then we have

$$u_{1} = \sum_{t=2}^{k} B_{t} \sum_{s=1}^{t-1} g_{s} g'_{t-s-1} + \sum_{t=2}^{k} (g_{t-1}B_{t} - B_{t}g_{t-1})$$
$$= \sum_{t=2}^{k} B_{t} \cdot 0 + \sum_{t=2}^{k} (g_{t-1}B_{t} - B_{t}g_{t-1})$$
$$= \sum_{t=2}^{k} (g_{t-1}B_{t} - B_{t}g_{t-1}) \in \mathfrak{u}_{1}^{+} \oplus \mathfrak{u}_{1}^{-}.$$

Here we put $g_0 = g'_0 = I_n$. Thus we are done.

Let us define subsets of G_k^o ,

$$\mathcal{P}_{k}^{\pm} := \left\{ \sum_{i=0}^{k-1} P_{i} z^{i} \in G_{k}^{o} \, \middle| \, P_{i} \in \mathfrak{p}_{i+1}^{\pm}, i = 1, \dots, k-1 \right\},\$$
$$\mathcal{U}_{k}^{\pm} := \left\{ \sum_{i=0}^{k-1} U_{i} z^{i} \in G_{k}^{o} \, \middle| \, U_{i} \in \mathfrak{u}_{i+1}^{\pm}, i = 1, \dots, k-1 \right\},\$$

and subspaces of \mathfrak{g}_k^o and $(\mathfrak{g}_k^o)^*$,

$$\mathfrak{U}_{k}^{\pm} := \left\{ \sum_{i=1}^{k-1} U_{i} z^{i} \left| U_{i} \in \mathfrak{u}_{i+1}^{\pm}, i = 1, \dots, k-1 \right\}, \\ (\mathfrak{U}_{k}^{\pm})^{*} := \left\{ \sum_{i=1}^{k-1} U_{i} z^{-i-1} \left| U_{i} \in \mathfrak{u}_{i+1}^{\pm}, i = 1, \dots, k-1 \right\}.$$

Here we put $\mathfrak{p}_k^{\pm} := M(n, \mathbb{C})$ and $\mathfrak{u}_k^{\pm} := \{0\}$. Let us note that \mathcal{P}_k^{\pm} are subgroups of G_k^o but \mathcal{U}_k^{\pm} are not closed under the multiplication.

Lemma 4.10 (cf. Lemma 3.8 in [16]). The \mathcal{P}_k^+ -orbit through B in \mathfrak{g}_k^* is $B + (\mathfrak{U}_k^-)^* \oplus (\mathfrak{u}_1^+ \oplus \mathfrak{u}_1^-) z^{-1}$.

Proof. First we see that \mathcal{P}_k^+ -orbit through B is included in $B + (\mathfrak{U}_k^-)^* \oplus (\mathfrak{u}_1^+ \oplus \mathfrak{u}_1^-)x^{-1}$. From Lemma 3.8 in [16], the \mathcal{P}_k^+ -orbit through B_{irr} in $(\mathfrak{g}_k^o)^*$ is $B_{\mathrm{irr}} + (\mathfrak{U}_k^-)^*$. Thus it suffices to show that $(pBp^{-1})_{\mathrm{res}} \in (B_1 + (\mathfrak{u}_1^+ \oplus \mathfrak{u}_1^-)) z^{-1}$ for all $p \in \mathcal{P}_k^+$. Let us set $u := (pBp^{-1})_{\mathrm{res}} - B_1$. Then $u = \sum_{t=2}^k \sum_{s=1}^{t-1} p_s B_t p'_{t-s-1}$

where $p = p_0 + p_1 x + p_2 x_2 + \cdots$, $p^{-1} = p'_0 + p'_1 x + p'_2 x^2 + \cdots$. Let h_s (resp. h'_s) be the \mathfrak{h}_s -component of p_s (resp. $p'_s) \in \mathfrak{p}_s^+ = \mathfrak{h}_s \oplus \mathfrak{u}_s^+$. Then

$$u_{i,i} = \left(\sum_{t=2}^{k} \sum_{s=1}^{t-1} p_s B_t p'_{t-s-1}\right)_{i,i} = \left(\sum_{t=2}^{k} \sum_{s=1}^{t-1} h_s B_t h'_{t-s-1}\right)_{i,i}$$
$$= \left(\sum_{t=2}^{k} B_t \sum_{s=1}^{t-1} h_s h'_{t-s-1}\right)_{i,i} + \sum_{t=2}^{k} (h_{t-1} B_t - B_t h_{t-1})_{i,i} = 0$$

for i = 1, ..., m. Here $X_{i,i}$ denote the $\operatorname{End}_{\mathbb{C}}(V_i)$ -component of $X \in M(n, \mathbb{C})$. Thus $u \in \mathfrak{u}_1^+ \oplus \mathfrak{u}_1^-$ as required.

Conversely take an arbitrary element $D \in B + (\mathfrak{U}_k^-)^* \oplus (\mathfrak{u}_1^+ \oplus \mathfrak{u}_1^-)z^{-1}$. Lemma 3.8 in [16] shows that there exists $p \in \mathcal{P}_k^+$ such that $(pDp^{-1})_{irr} = B_{irr}$. Namely $pDp^{-1} = B_{irr} + (h+u)x^{-1}$ where $h \in \mathfrak{h}_1, u \in \mathfrak{u}_1^+ \oplus \mathfrak{u}_1^-$. Thus there exists $p' \in \mathcal{P}_k^+$ such that $p'D(p')^{-1} = D' = B_{irr} + hx^{-1}$, an HTL normal form. Moreover recall that if $D' \in \mathcal{O}_B^o$ is of HTL normal form, then D' = B, cf. Lemma 2.9 in [16]. Thus D is in \mathcal{P}_k^+ -orbit through B.

Lemma 4.11 (Lemma 3.5 in [16]). For any $g \in G_k^o$, there uniquely exist $u_- \in \mathcal{U}_k^-$ and $p_+ \in \mathcal{P}_k^+$ such that $g = u_- p_+$.

For $A \in \mathcal{O}_{B_{\mathrm{irr}}}$, take $g \in G_k^o$ so that $g^{-1}Ag = B_{\mathrm{irr}}$ and decompose $g = u_-p_+$ as above. Note that u_- does not depend on the choice of g because the stabilizer of B_{irr} is contained in \mathcal{P}_k^+ . Thus u_- is uniquely determined by $A \in \mathcal{O}_{B_{\mathrm{irr}}}$. Then let us put $Q = u_- - I_n$, $A' = u_-^{-1}A$ and $P = A'|_{(\mathfrak{U}_k^-)^*}$.

Proposition 4.12 (Theorem 3.6 in [16]). The map

$$\begin{array}{cccc} \Phi \colon & \mathcal{O}_{B_{irr}} & \longrightarrow & \mathfrak{U}_k^- \times (\mathfrak{U}_k^-)^* \\ & A & \longmapsto & (Q, P) \end{array}$$

is bijective.

Proposition 4.13. Let us take $A \in \mathcal{O}_B^o$ and set $(Q, P) = \Phi(A_{irr})$. Then the following equations in \mathfrak{g}_k^* hold for $l = 1, \ldots, m$.

$$B_{l,l} - A_{l,l} = -Q_{l,1}P_{1,l} - Q_{l,2}P_{2,l} - \dots - Q_{l,l-1}P_{l-1,l} + P_{l,l+1}Q_{l+1,l} + P_{l,l+2}Q_{l+2,l} + \dots + P_{l,m}Q_{m,l}$$

Proof. First we note that putting $u = I_n + Q$, we have $u^{-1}Au \in B + (\mathfrak{U}_k^-)^* \oplus (\mathfrak{u}_1^+ \oplus \mathfrak{u}_1^-)z^{-1}$ by Lemma 4.10. Denote the $(\mathfrak{U}_k^-)^*$ and $(\mathfrak{u}_1^+ \oplus \mathfrak{u}_1^-)z^{-1}$ components of $u^{-1}Au$ by R and U respectively. Since Au = u(B + R + U), we have

(4)
$$B_{l,l} + \sum_{s=1}^{l-1} Q_{l,s}(R+U)_{s,l} = A_{l,l} + \sum_{s=l+1}^{m} A_{l,s}Q_{s,l}$$

Recalling that $Q \in M(n, z\mathbb{C}[z])$, we have $Q_{l,s}(R+U)_{s,l} = Q_{l,s}R_{s,l}$ in the above equation.

Let us set $\overline{A} = A_{irr}$. Note that

$$\bar{A}_{s,t} = \sum_{j=1}^{s-1} Q_{s,j} P_{j,t} + P_{s,t}$$

for $1 \leq s < t \leq m$. Then right hand side of the proposition is written as follows,

$$\begin{split} \bar{A}_{l,l+1}Q_{l+1,l} + \cdots + \bar{A}_{l,m}Q_{m,l} \\ &- Q_{l,1}(P_{1,l} + P_{1,l+1}Q_{l+1,l} + \cdots + P_{1,m}Q_{m,l}) \\ &- \cdots \\ &- Q_{l,l-1}(P_{l-1,l} + P_{l-1,l+1}Q_{l+1,l} + \cdots + P_{l-1,m}Q_{m,l}) \\ = &A_{l,l+1}Q_{l+1,l} + \cdots + A_{l,m}Q_{m,l} - Q_{l,1}R_{1,l} - \cdots - Q_{l,l-1}R_{l-1,l} \end{split}$$

Here we use fact that each component of $Q_{l,l'}$ (l > l') is in $z\mathbb{C}[z]$, which deduces $A_{l',l}Q_{l,l'} = \overline{A}_{l',l}Q_{l,l'}$. Then the above equation and the equation (4) induce the required equations.

Proposition 4.14. The map

$$\begin{split} \Psi_B \colon & \mathcal{O}_B^o \longrightarrow (\mathfrak{U}_k^- \times (\mathfrak{U}_k^-)^*) \times (\mathfrak{u}_1^+ \oplus \mathfrak{u}_1^-) \\ & A \longmapsto (\Phi(A_{irr}), A_{res}|_{\mathfrak{u}_1^+ \oplus \mathfrak{u}_1^-}) \end{split}$$

is bijective.

Proof. For $A \in \mathcal{O}_B^o$, let us put $(Q, P) = \Phi(A_{irr})$. Then Proposition 4.13 shows that $A_{res}|_{\mathfrak{h}_1}$ is uniquely determined by (Q, P) and B, which implies that Ψ_B is injective.

Let us take an arbitrary $((Q, P), U) \in (\mathfrak{U}_k^- \times (\mathfrak{U}_k^-)^*) \times (\mathfrak{u}_1^+ \oplus \mathfrak{u}_1^-)$. Define $h = \operatorname{diag}(h_1, \ldots, h_m) \in \mathfrak{h}_1$

by

$$h_i = R_i + \left(\sum_{j=1}^{i-1} Q_{i,j} P_{j,i} - \sum_{j=i+1}^m P_{i,j} Q_{j,i}\right)_{\text{res}}$$

Here recall that $R_i = (B_{\text{res}})_{i,i}$ for i = 1, ..., m. Then there exists an HTL normal form $B' \in \mathfrak{g}_k^*$ such that $B'_{\text{irr}} = B_{\text{irr}}$ and $\Phi^{-1}((Q, P)) + (h + U)x^{-1}$ is contained in some $\mathcal{O}_{B'}^o$. However B = B' by Proposition 4.13 and the construction of h. Thus $\Phi^{-1}((Q, P)) + (h + U)x^{-1}$ is the inverse image of ((Q, P), U) by Ψ_B . Hence Ψ_B is surjective. \Box

Let C_{R_i} be the conjugacy class of each R_i , block diagonal component of the residue of B. Recalling that $h^{-1}Bh = B_{irr} + h^{-1}B_{res}h$ for $h \in H$, we have one to one correspondence,

$$\operatorname{Ad}_{H}(B) := \{ hBh^{-1} \mid h \in H \} \longrightarrow \prod_{i=1}^{m} C_{R_{i}}$$
$$\longrightarrow ((B'_{\operatorname{res}})_{i,i})_{i=1,\dots,m}$$

Also recall that $\operatorname{Ad}_H(\mathcal{O}_B^o)$ is the disjoint union of $\mathcal{O}_{B'}^o$ for $B' \in \operatorname{Ad}_H(B)$, i.e., $\operatorname{Ad}_H(\mathcal{O}_B^o) = \bigsqcup_{B' \in \operatorname{Ad}_H(B)} \mathcal{O}_{B'}^o$, see Lemma 2.9 in [16].

For any $B' \in \operatorname{Ad}_H(B)$ we can define $\Psi_{B'}$ as in the above proposition. Thus we can define the map

$$\Psi \colon \mathrm{Ad}_{H}(\mathcal{O}_{B}^{o}) = \bigsqcup_{B' \in \mathrm{Ad}_{H}(B)} \mathcal{O}_{B'}^{o} \longrightarrow \left(\prod_{i=1}^{m} C_{R_{i}}\right) \times \left(\left(\mathfrak{U}_{k}^{-} \times (\mathfrak{U}_{k}^{-})^{*}\right) \times (\mathfrak{u}_{1}^{+} \oplus \mathfrak{u}_{1}^{-})\right)$$

by $\Psi(A) = (((B'_{res})_{i,i})_{1 \le i \le m}, \Psi_{B'}(A))$ for $A \in \mathcal{O}_{B'}^o$ with $B' = \sum_{i=1}^{k-1} B'_i x^{-i} \in S_B$. Then it is bijective.

Under these preparations, now we can define a quiver ${\sf Q}$ as follows. The set of vertices is

$$Q_0 := \{0\} \cup \{1, \ldots, m\}.$$

The set of arrows is

$$Q_1 := \left\{ \rho_{i,i'}^{[j]} \colon i \to i' \middle| \begin{array}{l} 1 \le i < i' \le m, \\ 1 \le j \le d(i,i') \end{array} \right\} \cup \left\{ \rho_i \colon 0 \to i \mid i = 1, \dots, m \right\}.$$

Fix the dimension vector $\alpha = (\alpha_a)_{a \in Q_0}$ defined by $\alpha_0 =: n$ and $\alpha_i := \dim_{\mathbb{C}} V_i$, $i = 1, \ldots, m$.

Let us construct a map from $\operatorname{Ad}_H(\mathcal{O}_B^o)$ to the representation space of $\overline{\mathbb{Q}}$. For $A \in \mathcal{O}_{B'}^o$, $B' \in \operatorname{Ad}_H(B)$, we set $(Q, P) = \Phi(A_{\operatorname{irr}})$ and define the representation $x_A \in \operatorname{Rep}(\overline{\mathbb{Q}}, \alpha)$ as follows:

$$(x_A)_{\rho_{i,i'}^{[j]}} := P_{i,i'}^{[j]}, \qquad (x_A)_{(\rho_{i,i'}^{[j]})^*} := Q_{i',i}^{[j]}, (x_A)_{\rho_i} := (I_n)_{i,*}, \qquad (x_A)_{\rho_i^*} := (A_{\text{res}})_{*,i},$$

for i, i' = 1, ..., m. Here we set $P = \sum_{i=1}^{k-2} P^{[i]} z^{-i-1}$ and $Q = \sum_{i=1}^{k-2} Q^{[i]} z^{i}$. Then Proposition 4.13 tells us that $\mu_{\alpha}(x_A)_i = (B'_{\text{res}})_{i,i} \in C_{R_i}$ for i = 1, ..., m.

Proposition 4.15. The following map is bijective,

$$\tilde{\Psi} \colon \mathrm{Ad}_{H}(\mathcal{O}_{B}^{o}) \longrightarrow \left\{ x \in \mathrm{Rep}\left(\overline{\mathbb{Q}}, \alpha\right) \middle| \begin{array}{c} (\psi_{\rho_{i}})_{1 \leq i \leq m} = I_{n}, \ \mu_{\alpha}(x)_{i} \in C_{R_{i}} \\ for \ i = 1, \dots, m \end{array} \right\},$$

which is defined by $\tilde{\Psi}(A) := x_A$ for $A \in \operatorname{Ad}_H(\mathcal{O}_B^o)$ as above. Moreover $\tilde{\Psi}$ preserves *H*-actions, i.e., $\tilde{\Psi}(hAh^{-1}) = h \cdot x_A$ for all $h \in H$.

Proof. Proposition 4.14 shows that $\tilde{\Psi}$ is bijective. The last assertion can be directly checked.

Finally we can obtain a correspondence between \mathcal{O}_B and representations of $\overline{\mathbb{Q}}$.

Proposition 4.16. There exists a bijection

$$\mathcal{O}_B \cong \operatorname{GL}(n, \mathbb{C}) \times_H \operatorname{Ad}_H(\mathcal{O}_B^o) \longrightarrow \left\{ x \in \operatorname{Rep}\left(\overline{\mathbb{Q}}, \alpha\right) \middle| \begin{array}{c} \det(x_{\rho_i})_{1 \le i \le m} \neq 0, \mu_\alpha(x)_i \in C_{R_i} \\ for \ i = 1, \dots, m \end{array} \right\} / \prod_{i=1}^m \operatorname{GL}(\alpha_i, \mathbb{C}).$$

Proof. Let us define a map $\overline{\Psi}$ from $\operatorname{GL}(n, \mathbb{C}) \times \operatorname{Ad}_H(\mathcal{O}_B^o)$ to

$$\left\{ x \in \operatorname{Rep}\left(\overline{\mathsf{Q}}, \alpha\right) \middle| \begin{array}{c} \det\left(x_{\rho_i}\right)_{1 \le i \le m} \ne 0, \ \mu_{\alpha}(x)_i \in C_{R_i} \\ \text{for } i = 1, \dots, m \end{array} \right\}$$

For $(g, A) \in \operatorname{GL}(n, \mathbb{C}) \times \operatorname{Ad}_H(\mathcal{O}_B^o), x = \overline{\Psi}((g, A)) =$ is defined as below:

$$\begin{aligned} x_{\rho_{i,i'}^{[j]}} &= P_{i,i'}^{[j]}, & x_{(\rho_{i,i'}^{[j]})^*} = Q_{i',i}^{[j]}, \\ x_{\rho_i} &= (g^{-1})_{i,*}, & x_{\rho_i^*} = ((gA)_{\text{res}})_{*,i}, \end{aligned}$$

where $(Q, P) = \Phi(A_{irr})$ and write $P = \sum_{i=1}^{k-2} P^{[i]} z^{-i-1}$, $Q = \sum_{i=1}^{k-2} Q^{[i]} z^i$. Proposition 4.15 shows that this map is bijective. Moreover we can directly check that this map preserves *H*-actions. Thus we are done.

For example, let us consider an HTL normal form $B = \sum_{i=1}^{4} B_i z^{-i}$ such that

$B_4 = \operatorname{diag} \left(a_1^{(4)}, a_2^{(4)}, a_2^{(4)}, a_2^{(4)} \right),$	$B_3 = \operatorname{diag}(*, a_1^{(3)}, a_2^{(3)}, a_2^{(3)}),$
$B_2 = \operatorname{diag}(*, *, a_1^{(2)}, a_2^{(2)}),$	$B_1 = \text{diag}(*, *, *, *),$

where $a_1^{(i)} \neq a_2^{(i)}$. Then the corresponding quiver is as follows.



Proposition 4.17 (cf. Lemma 3.1 in [16]). Let $(g, A) \in GL(n, \mathbb{C}) \times_H$ $Ad_H(\mathcal{O}_B^o) \cong \mathcal{O}_B$ and $x \in \operatorname{Rep}(\overline{\mathbb{Q}}, \alpha)$ the corresponding elements under the isomorphism in Proposition 4.16.

- (i) Let us suppose that x has a subrepresentation $N = (N_a, \psi_\rho)_{a \in Q_0, \rho \in \overline{Q}_1}$ satisfying dim_C $\bigoplus_{i=1}^m N_i = \dim_{\mathbb{C}} N_0$, then $W = \bigoplus_{i=1}^m N_i$ is invariant under all A_i where $A = \sum_{i=1}^k A_i z^{-i}$ and similarly $gW = N_0$ is invariant under all $gA_i g^{-1}$.
- (ii) Any subspace $S \subset \mathbb{C}^n$ invariant under all A_i is homogeneous with respect to the decomposition $\mathbb{C}^n = \bigoplus_{i=1}^m V_i$. Moreover $N_i = S \cap V_i$, $i=1,\ldots,m$, and $N_0 = gS$ define the subrepresentation $N = (N_a, \psi_\rho)$ where ψ_ρ are restrictions of x_ρ on N_i .

Proof. If x has a subrepresentation N as above, then Lemma 3.10 in [16] shows that W is invariant under A_i for $i \ge 2$ and obviously invariant under A_1 since $A_1 = (x_{\rho_i^*})_{1 \le i \le m} (x_{\rho_i})_{1 \le i \le m}$.

Conversely let $S \subset \mathbb{C}^n$ be a subspace invariant under all A_i . We need to check that all ψ_{ρ} are well-defined, namely $x_{\rho}(N_{s(\rho)}) \subset N_{t(\rho)}$ which are already checked in Lemma 3.10 in [16] for $\rho_{i,i'}^{[j]}$ and $(\rho_{i,i'}^{[j]})^*, i, i' = 1, \ldots, m, j = 1, \ldots, d(i, i')$. For ρ_i , we have $x_{\rho_i}(N_i) = gN_i \subset gS = N_0$. For ρ_i^* , we have $x_{\rho_i^*}(N_0) = V_i \cap A_1 g^{-1} N_0 = V_i \cap A_1 S \subset N_i$.

4.3. Quivers associated with differential equations. Now we are ready to consider a correspondence between moduli spaces $\mathfrak{M}(\mathbf{B})$ of arbitrary k_i and subsets of quiver varieties $\mathfrak{M}_{\lambda}(\mathbf{Q}, \alpha)$ as we saw in Section 4.1 under the restriction $k_0 = \cdots = k_p = 2$.

For i = 0, ..., p, let us fix a collection of nonzero positive integers k_i and HTL normal forms $B_i = \sum_{j=1}^{k_i} B_j^{(i)} z^{-j} \in \mathfrak{g}_{k_i}^*$. Then write

$$B_{i} = \operatorname{diag}\left(q_{[i,1]}(z^{-1})I_{n_{[i,1]}} + R_{[i,1]}z^{-1}, \dots, q_{[i,m_{i}]}(z^{-1})I_{n_{[i,m_{i}]}} + R_{[i,m_{i}]}z^{-1}\right)$$

for i = 0, ..., p where $q_{[i,j]}(z^{-1}) \in z^{-2}\mathbb{C}[z^{-1}]$ satisfying $q_{[i,j]} \neq q_{[i,j']}$ if $j \neq j'$ and $R_{[i,j]} \in M(n_{[i,j]}, \mathbb{C})$.

For each i = 0, ..., p, decompose $\mathbb{C}^n = \bigoplus_{j=1}^{m_i(s)} V_{\langle s,j \rangle}^{(i)}$ as simultaneous $(B_{s+1}^{(i)}, \ldots, B_{k_i}^{(i)})$ -invariant subspaces. In particular we write $V_{[i,j]} := V_{\langle 1,j \rangle}^{(i)}$ for i = 0, ..., p and $j = 1, ..., m_i$. Here we note $m_i(1) = m_i$.

For each pair $j, j' \in \{1, \ldots, m_i\}$, attach the integer $d_i(j, j')$ defined by

$$d_i(j,j') := \deg_{\mathbb{C}[z^{-1}]}(q_{[i,j]}(z^{-1}) - q_{[i,j']}(z^{-1})) - 2$$

if $j \neq j'$ or $d_i(j, j') := -1$ if j = j'. Set $I_{irr} := \{i \in \{0, \dots, p\} \mid m_i > 1\} \cup \{0\}$ and $I_{reg} := \{0, \dots, p\} \setminus I_{irr}$.

Remark 4.18. Suppose that $m_i = 1$ for some $i \in \{0, \ldots, p\}$. Then the truncated orbit of the normal form B_i is trivial, namely $\mathcal{O}_{B_i} \cong C_{R_{[i,1]}}$. Thus I_{irr} can be seen as the set of singular points at which truncated orbits are nontrivial and we add the point 0 as a "base point" to I_{irr} .

Now consider the following quiver $Q^{\rm irr}$. The set of vertices is

$$\mathsf{Q}_0^{_{
m Irr}} := \{ [i,j] \mid i \in I_{
m irr}, \, j = 1, \dots, m_i \}$$
 .

Also define

$$\begin{aligned} \mathsf{Q}_{1}^{0 \to i_{\mathrm{irr}}} &:= \left\{ \rho_{[i,j']}^{[0,j]} \colon [0,j] \to [i,j'] \left| \begin{array}{c} j = 1, \dots, m_{0}, \\ i \in I_{\mathrm{irr}} \setminus \{0\}, \\ j = 1, \dots, m_{i} \end{array} \right\}, \\ \mathsf{Q}_{1}^{B_{i}} &:= \left\{ \rho_{[i,j],[i,j']}^{[k]} \colon [i,j] \to [i,j'] \left| \begin{array}{c} i \in I_{\mathrm{irr}}, \\ 1 \leq j < j' \leq m_{i}, \\ 1 \leq k \leq d_{i}(j,j') \end{array} \right\}. \end{aligned} \right. \end{aligned}$$

Then the set of vertices is

$$\mathsf{Q}_1^{\mathrm{irr}} := \mathsf{Q}_1^{0 \to i_{\mathrm{irr}}} \sqcup \bigsqcup_{i \in I_{\mathrm{irr}}} \mathsf{Q}_1^{B_i}.$$

Let us define the dimension vector $\alpha^{\text{irr}} = (\alpha_a)_{a \in \mathbf{Q}_0^{\text{irr}}}$ by $\alpha_{[i,j]} := n_{[i,j]}$.

Proposition 4.19. There exists a bijection

$$\begin{split} \Phi \colon \left\{ \left(\sum_{j=1}^{k_i} A_j^{(i)} z^{-j} \right)_{0 \leq i \leq p} \in \prod_{i=0}^p \mathcal{O}_{B_i} \left| \begin{array}{c} \sum_{i=0}^p A_1^{(i)} = 0, \\ irreducible \end{array} \right\} / \operatorname{GL}(n, \mathbb{C}) \longrightarrow \\ \left\{ \begin{array}{c} \left(\sum_{j=1}^{k_i} A_j^{(i)} z^{-j} \right) \in \mathcal{O}_{B_i} \\ \operatorname{Rep}(\overline{\mathbb{Q}^{irr}}, \alpha^{irr}) \times \prod_{i \in I_{reg}} \mathcal{O}_{B_i} \end{array} \right| \begin{array}{c} \mathcal{L}^{irr} \text{-} irreducible, \\ \operatorname{det} \left(x_{\rho_{[i,j']}^{[0,j]}} \right)_{\substack{1 \leq j \leq m_0 \\ 1 \leq j' \leq m_i}} \neq 0, \ i \in I_{irr} \setminus \{0\}, \\ \mu_{\alpha^{irr}}(x)_{[i,j]} \in C_{R_{[i,j]}}, \ i \in I_{irr} \\ / \prod_{a \in \mathbb{Q}_0^{irr}} \operatorname{GL}(\alpha_a, \mathbb{C}). \end{split} \end{split}$$

Here

$$\mathcal{L}^{irr} := \left\{ \beta \in \mathbb{Z}^{\mathsf{Q}_0^{irr}} \left| \sum_{j=1}^{m_0} \beta_{[0,j]} = \sum_{j=1}^{m_i} \beta_{[i,j]} \text{ for all } i \in I_{irr} \setminus \{0\} \right\}.$$

Proof. Setting $H_0 := \{ \operatorname{diag}(h_1, \ldots, h_{m_0}) \mid h_i \in \operatorname{GL}(n_{[0,i]}, \mathbb{C}) \}$, we can identify $\mathcal{O}_{B_0}/\operatorname{GL}(n, \mathbb{C})$ with $\operatorname{Ad}_{H_0}(\mathcal{O}_{B_0}^o)/H_0$, see Proposition 4.8. Let us set

$$V := \left\{ \begin{array}{c} \left(\sum_{j=1}^{k_i} A_j^{(i)} z^{-j}\right)_{0 \le i \le p} \in \\ \operatorname{Ad}_{H_0}(\mathcal{O}_{B_0}^o) \times \prod_{i \in I_{\operatorname{irr}} \setminus \{0\}} \mathcal{O}_{B_i} \times \prod_{i \in I_{\operatorname{reg}}} \mathcal{O}_{B_i} \end{array} \right| \sum_{i=0}^p A_1^{(i)} = 0 \right\}.$$

Then we can identify

$$\left\{ \left(\sum_{j=1}^{k_i} A_j^{(i)} z^{-j} \right)_{0 \le i \le p} \in \prod_{i=0}^p \left| \sum_{i \in I_{\operatorname{irr}}} A_1^{(i)} = 0 \right\} / \operatorname{GL}(n, \mathbb{C}) \right\}$$

with V/H_0 .

Comparing with Proposition 4.3, we see that Propositions 4.15 and 4.16 give a bijection from V to

$$\begin{cases} x \in \operatorname{Rep}(\overline{\mathbf{Q}^{\operatorname{irr}}}, \alpha^{\operatorname{irr}}) \middle| & \det\left(x_{\rho_{[i,j']}^{[0,j]}}\right)_{\substack{1 \le j \le m_0 \\ 1 \le j' \le m_i}} \neq 0, \ i \in I_{\operatorname{irr}} \setminus \{0\}, \\ & \mu_{\alpha^{\operatorname{irr}}}(x)_{[i,j]} \in C_{R_{[i,j]}}, i \in I_{\operatorname{irr}} \\ & \swarrow \\ & \bigwedge \\ & A_{a \in \mathbf{Q}_0^{\operatorname{irr}} \setminus \{[0,j]|j=1,\ldots,m_0\}} \operatorname{GL}(\alpha_a, \mathbb{C}) \end{cases}$$

and the bijection preserves H_0 -actions. Thus we have the required bijection. The correspondence between the irreducibility and \mathcal{L}^{irr} -irreducibility follows from Proposition 4.17 and the arguments in the proof of Proposition 4.3. \Box

Finally as we saw in Section 3, we shall associate conjugacy classes of residue matrices of HTL normal forms to representations of quivers. For each $R_{[i,j]}$, $i = 0, \ldots, p$ and $j = 1, \ldots, m_i$, let us choose $\xi_1^{[i,j]}, \ldots, \xi_{e_{[i,j]}}^{[i,j]} \in \mathbb{C}$ so that

$$\prod_{k=1}^{\mathbb{P}[i,j]} (R_{[i,j]} - \xi_k^{[i,j]}) = 0.$$

Let $\xi = \left(\left\{\xi_1^{[i,j]}, \dots, \xi_{e_{[i,j]}}^{[i,j]}\right\}\right)_{\substack{0 \le i \le p, \\ 1 \le j \le m_i}}$ be the collection of the ordered sets $\left\{\xi_1^{[i,j]}, \dots, \xi_{e_{[i,j]}}^{[i,j]}\right\}.$

Now consider the following quiver Q. Set

$$\mathbf{Q}_{0}^{\text{leg}} := \left\{ \begin{bmatrix} i, j, k \end{bmatrix} \middle| \begin{array}{c} i = 0, \dots, p, \\ j = 1, \dots, m^{(i)}, \\ k = 1, \dots, e_{[i,j]} - 1 \end{array} \right\}$$

Then the set of vertices is

$$\mathsf{Q}_0 := \mathsf{Q}_0^{\operatorname{irr}} \sqcup \mathsf{Q}_0^{\operatorname{leg}}.$$

Also set

$$\begin{aligned} \mathsf{Q}_{1}^{\mathrm{leg}_{i}} &:= \left\{ \rho_{[i,j,k]} \colon [i,j,k] \to [i,j,k-1] \left| \begin{array}{c} j = 1, \dots, m_{i}, \\ k = 2, \dots, e_{[i,j]} - 1 \end{array} \right\}, \\ \mathsf{Q}_{1}^{\mathrm{leg}_{i} \to B_{i}} &:= \left\{ \rho_{[i,j,1]} \colon [i,j,1] \to [i,j] \mid j = 1, \dots, m_{i} \right\}, \\ \mathsf{Q}_{1}^{\mathrm{leg}_{i} \to 0} &:= \left\{ \rho_{[0,j]}^{[i,1,1]} \colon [i,1,1] \to [0,j] \mid i \in I_{\mathrm{reg}}, \, j = 1, \dots, m_{0} \right\}. \end{aligned}$$

The set of arrows is

$$\begin{split} \mathsf{Q}_1 &:= \mathsf{Q}_1^{0 \to I_{\mathrm{irr}}} \sqcup \bigsqcup_{i \in I_{\mathrm{irr}}} \left(\mathsf{Q}_1^{B_i} \sqcup \mathsf{Q}_1^{\mathrm{leg}_i \to B_i} \sqcup \mathsf{Q}_1^{\mathrm{leg}_i} \right) \\ & \sqcup \bigsqcup_{i \in I_{\mathrm{reg}}} \left(\mathsf{Q}_1^{\mathrm{leg}_i \to 0} \sqcup \mathsf{Q}_1^{\mathrm{leg}_i} \right). \end{split}$$

For example, let us consider the following $\mathbf{B} = (B_0, B_1, B_2)$.

$$B^{(0)} = \begin{pmatrix} a_4^{(0)} & & & \\ & a_4^{(0)} & & \\ & & & a_4^{(0)} & \\ & & & & b_4^{(0)} \end{pmatrix} z^{-4} + \begin{pmatrix} a_3^{(0)} & & & \\ & & a_3^{(0)} & & \\ & & & b_3^{(0)} & \\ & & & & c_3^{(0)} \end{pmatrix} z^{-3} + \begin{pmatrix} a_2^{(0)} & & & \\ & & & b_2^{(0)} & & \\ & & & & c_2^{(0)} & \\ & & & & & c_2^{(0)} & \\ & & & & & & c_2^{(0)} \end{pmatrix} z^{-2} + \begin{pmatrix} \xi_1^{[0,1]} & & & \\ & & \xi_1^{[0,2]} & & \\ & & & & & & \xi_1^{[0,4]} \end{pmatrix} z^{-1},$$

$$B_{1} = \begin{pmatrix} a_{2}^{(1)} & & \\ & a_{2}^{(1)} & \\ & & a_{2}^{(1)} \\ & & & b_{2}^{(1)} \end{pmatrix} z^{-2} + \begin{pmatrix} \xi_{1}^{[1,1]} & & \\ & & \xi_{2}^{[1,1]} & \\ & & & \xi_{3}^{[1,1]} \\ & & & & \xi_{1}^{[1,2]} \end{pmatrix} z^{-1},$$

$$B_2 = \begin{pmatrix} \xi_1^{[2,1]} & & \\ & \xi_2^{[2,1]} & \\ & & \xi_3^{[2,1]} \\ & & & \xi_4^{[2,1]} \end{pmatrix} z^{-1}.$$

Here any distinct two of $\{a_j^{(i)}, b_j^{(i)}, c_j^{(i)}, d_j^{(i)}\}$ stand for distinct complex numbers and $\xi_k^{[i,j]} \neq \xi_{k'}^{[i,j]}$ if $k \neq k'$.

Then we can associate the following quiver to this **B**.



Define the dimension vector $\alpha = (\alpha_a)_{a \in Q_0}$ by

$$\alpha_{[i,j]} := n_{[i,j]}, \qquad \qquad \alpha_{[i,j,k]} := \operatorname{rank} \prod_{l=1}^{k} (R_{[i,j]} - \xi_l^{[i,j]}).$$

Also define $\lambda = (\lambda_a)_{a \in Q_0}$ by

$$\begin{split} \lambda_{[i,j]} &:= -\xi_1^{[i,j]}, & \text{for } i \in I_{\text{irr}} \setminus \{0\}, \ j = 1, \dots, m_i, \\ \lambda_{[0,j]} &:= -\xi_1^{[0,j]} - \sum_{i \in I_{\text{reg}}} \xi_1^{[i,1]} & \text{for } j = 1, \dots, m_0, \\ \lambda_{[i,j,k]} &:= \xi_k^{[i,j]} - \xi_{k+1}^{[i,j]} & \text{for } \substack{i = 0, \dots, p, \ j = 1, \dots, m_i, \\ k = 1, \dots, e_{[i,j]} - 1. \end{split}$$

Then Propositions 3.1 and 4.19 show the following.

Proposition 4.20. Let B_0, \ldots, B_p be HTL normal forms chosen as above. Then there exists a bijection

$$\begin{split} \Phi_{\xi} \colon \left\{ \left(\sum_{j=1}^{k_i} A_j^{(i)} x^{-j} \right)_{0 \leq i \leq p} \in \prod_{i=0}^p \mathcal{O}_{B_i} \middle| \sum_{i=0}^p A_1^{(i)} = 0 \right\} / \operatorname{GL}(n, \mathbb{C}) \\ \to \left\{ x \in \mu^{-1}(\lambda) \subset \operatorname{Rep}(\overline{\mathbb{Q}}, \alpha) \middle| \\ \det \left(x_{\rho_{[i,j']}^{[0,j]}} \right)_{1 \leq j \leq m_0} \neq 0, \, i \in I_{irr} \setminus \{0\}, \\ \left(x_{\rho_{[0,j]}^{[i,1,1]}} \right)_{1 \leq j \leq m_0} : \mathbb{C}^{\alpha_{[i,1,1]}} \to \bigoplus_{j=1}^{m_0} \mathbb{C}^{\alpha_{[0,j]}}, \text{ injective, } i \in I_{reg}, \\ \left(x_{\left(\rho_{[0,j]}^{[i,1,1]}\right)^*} \right)_{1 \leq j \leq m_0} : \bigoplus_{j=1}^{m_0} \mathbb{C}^{\alpha_{[0,j]}} \to \mathbb{C}^{\alpha_{[i,1,1]}}, \text{ surjective, } i \in I_{reg}, \\ x_{\rho_{[i,j,k]}}, \text{ injective, } x_{(\rho_{[i,j,k]})^*}, \text{ surjective} \right\} / \prod_{a \in \mathbb{Q}_0} \operatorname{GL}(\alpha_a, \mathbb{C}). \end{split}$$

Finally we shall close this section giving the isomorphism from the moduli space of meromorphic connections $\mathfrak{M}(\mathbf{B})$ as follows.

Define a sublattice of $\mathbb{Z}^{\mathsf{Q}_0}$,

$$\mathcal{L} := \left\{ \beta \in \mathbb{Z}^{\mathbf{Q}_0} \left| \sum_{j=1}^{m^{(0)}} \beta_{[0,j]} = \sum_{j=1}^{m^{(i)}} \beta_{[i,j]} \text{ for all } i \in I_{\mathrm{irr}} \setminus \{0\} \right\}.$$

Set $\mathcal{L}^+ := \mathcal{L} \cap (\mathbb{Z}_{\geq 0})^{\mathsf{Q}_0}$.

Definition 4.21 (\mathcal{L} -irreducible). If $x \in \operatorname{Rep}(\overline{\mathbb{Q}}, \alpha)$ has no nontrivial proper subrepresentation $\{0\} \neq y \subsetneq x$ with $\dim y \in \mathcal{L}$, then x is said to be \mathcal{L} -*irreducible*.

Then we define a subset of the quiver variety $\mathfrak{M}_{\lambda}(\mathsf{Q},\alpha)$ as follows,

$$\mathfrak{M}_{\lambda}(\mathsf{Q},\alpha)^{\mathrm{dif}} := \mu_{\alpha}^{-1}(\lambda)^{\mathrm{dif}}/\mathbf{G}(\alpha)$$

where

$$\mu_{\alpha}^{-1}(\lambda)^{\operatorname{dif}} := \left\{ x \in \mu_{\alpha}^{-1}(\lambda) \left| \begin{array}{c} x \text{ is } \mathcal{L}\text{-irreducible,} \\ \det\left(x_{\rho_{[i,j']}^{[0,j]}}\right)_{\substack{1 \leq j \leq m_0 \\ 1 \leq j' \leq m_i}} \neq 0, \ i \in I_{\operatorname{irr}} \setminus \{0\} \end{array} \right\}.$$

Also define

$$\mu_{\alpha}^{-1}(\lambda)^{\det} := \left\{ x \in \mu_{\alpha}^{-1}(\lambda) \, \middle| \, \det \left(x_{\rho_{[i,j']}^{[0,j]}} \right)_{\substack{1 \le j \le m_0 \\ 1 \le j' \le m_i}} \neq 0, \, i \in I_{\mathrm{irr}} \setminus \{0\} \right\}$$

for the latter use.

Theorem 4.22. The restriction of the map in Proposition 4.20,

$$\Phi_{\xi} \colon \mathfrak{M}(\mathbf{B}) \longrightarrow \mathfrak{M}_{\lambda}(\mathsf{Q}, \alpha)^{dif}$$

is well-defined and bijective.

Proof. The map Φ_{ξ} in Proposition 4.20 sends irreducible $\mathbf{A} \in \prod_{i=0}^{p} \mathcal{O}_{B_{i}}$ to the \mathcal{L} -irreducible representation $\Phi_{\xi}(\mathbf{A})$ and vice versa by Propositions 3.1 and 4.19. Thus we only need to check that the \mathcal{L} -irreducibility of a representation $x, \rho \in \mu^{-1}(\lambda)^{\text{dif}}$ implies that $\left(x_{\rho_{0,j}^{[i,1,1]}}\right)_{1 \leq j \leq m_{0}}$ and $x_{\rho_{[i,j,k]}}$ are injective, and $\left(x_{\left(\rho_{0,j}^{[i,1,1]}\right)^{*}}\right)_{1 \leq j \leq m_{0}}$ and $x_{\left(\rho_{[i,j,k]}\right)^{*}}$ are surjective. This can be shown as in the proof of Theorem 4.6.

5. MIDDLE CONVOLUTIONS AND REFLECTIONS

In the previous section we gave an isomorphism $\mathfrak{M}(\mathbf{B}) \cong \mathfrak{M}_{\lambda}(\mathbf{Q}, \alpha)^{\text{dif}}$. However $\mathfrak{M}_{\lambda}(\mathbf{Q}, \alpha)^{\text{dif}}$ does not coincide with $\mathfrak{M}_{\lambda}^{\text{reg}}(\mathbf{Q}, \alpha)$ in general. Namely, we can not refer to Theorem 1.2 in [8] by Crawley-Boevey (see Corollary 2.8) for the non-emptiness of $\mathfrak{M}(\mathbf{B})$ in our general setting. Therefore in the remaining sections, we shall determine the condition for the non-emptiness of $\mathfrak{M}_{\lambda}(\mathbf{Q}, \alpha)^{\text{dif}}$ tracing the case of $\mathfrak{M}_{\lambda}^{\text{reg}}(\mathbf{Q}, \alpha)$ done by Crawley-Boevey in [8]. In the case of $\mathfrak{M}_{\lambda}^{\text{reg}}(\mathbf{Q}, \alpha)$, reflection functors (see [10] and [30]) plays very impotent role as we can see in Kac's theorem for the existence of indecomposable representations of quivers in [18]. Unfortunately it, however, is easy

to see that all reflection functors on $\mathfrak{M}(\mathbf{Q}, \alpha)$ do not preserve $\mathfrak{M}_{\lambda}(\mathbf{Q}, \alpha)^{\text{dif}}$ because of the open condition $\det(x_{\rho_{[i,j']}^{[0,j]}})_{\substack{1 \leq j \leq m_0 \\ 1 \leq j' \leq m_i}} \neq 0$. Thus in this section we shall introduce some operations for $\mathfrak{M}_{\lambda}(\mathbf{Q}, \alpha)^{\text{dif}}$ which are induced from operations originally defined for $\mathfrak{M}(\mathbf{B})$, so-called middle convolutions, ad-

operations originally defined for $\mathfrak{M}(\mathbf{B})$, so-called middle convolutions, additions, etc. The map in Proposition 4.20 enable us to define analogous operations on $\mathfrak{M}_{\lambda}(\mathbf{Q}, \alpha)^{\text{dif}}$.

In this section we retain the notation in the previous section.

5.1. A review of middle convolutions. Let us give a review of middle convolutions on differential equations with irregular singular points. The middle convolution is originally defined by N. Katz in [24] and reformulated as an operation on Fuchsian systems by Dettweiler-Reiter [13], see also [12] and Völklein's paper [34]. There are several studies to generalize the middle convolution to non-Fuchsian differential equations, see [1],[25],[32],[37] for example. Among them we shall give a review of middle convolutions following [37].

From $\mathbf{A} = (\sum_{j=1}^{k_i} A_j^{(i)} z^{-j})_{0 \le i \le p} \in \prod_{i=0}^p \mathcal{O}_{B_i}$, let us construct a 5-tuple (V, W, T, Q, P) consisting of \mathbb{C} -vector spaces V, W and $T \in \operatorname{End}_{\mathbb{C}}(W), Q \in \operatorname{Hom}_{\mathbb{C}}(W, V), P \in \operatorname{Hom}_{\mathbb{C}}(V, W)$. Set $V := \mathbb{C}^n$ and $\widehat{W}_i := V^{\oplus k_i}$ for $i = 0, \ldots, p$. Then define

$$\widehat{Q}_{i} := (A_{k_{i}}^{(i)}, A_{k_{i}-1}^{(i)}, \dots, A_{1}^{(i)}) \in \operatorname{Hom}_{\mathbb{C}}(\widehat{W}_{i}, V),
\widehat{P}_{i} := \begin{pmatrix} 0 \\ \vdots \\ 0 \\ \operatorname{Id}_{V} \end{pmatrix} \in \operatorname{Hom}_{\mathbb{C}}(V, \widehat{W}_{i}), \ \widehat{N}_{i} := \begin{pmatrix} 0 & \operatorname{Id}_{V} & 0 \\ 0 & \ddots & \\ & \ddots & \operatorname{Id}_{V} \\ 0 & & 0 \end{pmatrix} \in \operatorname{End}_{\mathbb{C}}(\widehat{W}_{i}).$$

Setting

$$\begin{split} \widehat{W} &:= \bigoplus_{i=0}^{p} \widehat{W}_{i}, \\ \widehat{T} &:= (\widehat{N}_{i})_{0 \leq i \leq p} \in \bigoplus_{i=0}^{p} \operatorname{End}_{\mathbb{C}}(\widehat{W}_{i}) \subset \operatorname{End}_{\mathbb{C}}(\widehat{W}), \\ \widehat{Q} &:= (\widehat{Q}_{i})_{0 \leq i \leq p} \in \bigoplus_{i=0}^{p} \operatorname{Hom}_{\mathbb{C}}(\widehat{W}_{i}, V) = \operatorname{Hom}_{\mathbb{C}}(\widehat{W}, V), \\ \widehat{P} &:= (\widehat{P}_{i})_{0 \leq i \leq p} \in \bigoplus_{i=0}^{p} \operatorname{Hom}_{\mathbb{C}}(V, \widehat{W}_{i}) = \operatorname{Hom}_{\mathbb{C}}(V, \widehat{W}), \end{split}$$

we have a 5-tuple $(V, \widehat{W}, \widehat{T}, \widehat{Q}, \widehat{P})$. Further setting

$$\widehat{A}_{i} := \begin{pmatrix} A_{k_{i}}^{(i)} & A_{k_{i}-1}^{(i)} & \cdots & A_{1}^{(i)} \\ & A_{k_{i}}^{(i)} & \ddots & \vdots \\ & & \ddots & A_{k_{i}-1}^{(i)} \\ 0 & & & A_{k_{i}}^{(i)} \end{pmatrix} \in \operatorname{End}_{\mathbb{C}}(\widehat{W}_{i}),$$

we define $W_i := \widehat{W}_i / \operatorname{Ker} \widehat{A}_i$ and $W := \bigoplus_{i=0}^p W_i$. Then T, Q, P are the maps induced from $\widehat{T}, \widehat{Q}, \widehat{P}$ respectively.

Definition 5.1 (Yamakawa [37]). The 5-tuple (V, W, T, Q, P) given above is called the *canonical datum* for $\mathbf{A} \in \prod_{i=0}^{p} \mathcal{O}_{B_i}$.

Lemma 5.2. Let (V, W, T, Q, P) is the canonical datum of $\mathbf{A} \in \prod_{i=0}^{p} \mathcal{O}_{B_i}$. Then

$$\dim_{\mathbb{C}} W = \sum_{i=0}^{p} \sum_{j=0}^{k_i-1} \left(n - \dim_{\mathbb{C}} \bigcap_{l=0}^{j} \operatorname{Ker} B_{k_i-l}^{(i)} \right)$$

Proof. For $\mathbf{A} = (\sum_{j=1}^{k_i} A_j^{(i)} z^{-j})_{0 \le i \le p} \in \prod_{i=0}^p \mathcal{O}_{B_i}$, there exists $g^{(i)} = g_0^{(i)} + g_1^{(i)} z + \dots + g_{k_i-1}^{(i)} z^{k_i-1} \in G_{k_i}$ such that $(g^{(i)})^{-1} (\sum_{j=1}^{k_i} A_j^{(i)} z^{-j}) g^{(i)} = B_i$. Thus

$$\begin{pmatrix} g_0^{(i)} & g_1^{(i)} & \cdots & g_{k_i-1}^{(i)} \\ g_0^{(i)} & \ddots & \vdots \\ & \ddots & g_1^{(i)} \\ 0 & & g_0^{(i)} \end{pmatrix}^{-1} \begin{pmatrix} A_{k_i}^{(i)} & A_{k_i-1}^{(i)} & \cdots & A_1^{(i)} \\ & A_{k_i}^{(i)} & \ddots & \vdots \\ & & \ddots & A_{k_i-1}^{(i)} \\ 0 & & & A_{k_i}^{(i)} \end{pmatrix} \begin{pmatrix} g_0^{(i)} & g_1^{(i)} & \cdots & g_{k_i-1}^{(i)} \\ & g_0^{(i)} & \ddots & \vdots \\ & & \ddots & g_1^{(i)} \\ 0 & & & & g_0^{(i)} \end{pmatrix}$$
$$= \begin{pmatrix} B_{k_i}^{(i)} & B_{k_i-1}^{(i)} & \cdots & B_1^{(i)} \\ & B_{k_i}^{(i)} & \ddots & \vdots \\ & & \ddots & B_{k_i-1}^{(i)} \\ 0 & & & & B_{k_i}^{(i)} \end{pmatrix} = \hat{B}_i.$$

This implies that $\dim_{\mathbb{C}} \operatorname{Ker} \widehat{A}_i = \dim_{\mathbb{C}} \operatorname{Ker} \widehat{B}_i$ as required.

Fix $t \in \{0, \ldots, p\}$, take a polynomial $p_t(z^{-1}) = \sum_{j=1}^{k_t} p_j^{(t)} z^{-j} \in z^{-1} \mathbb{C}[z^{-1}]$ and define an operation, called *addition*, as follows. For an element $\mathbf{A} = (A_i(x^{-1}))_{0 \leq i \leq p} \in \prod_{i=0}^p \mathcal{O}_{B_i}$, we define $\operatorname{Add}_{p_t(z^{-1})}^{(t)}(\mathbf{A}) := (A'_i(z^{-1}))_{0 \leq i \leq p}$ by

$$A'_i(z^{-1}) := \begin{cases} A_i(z^{-1}) & \text{if } i \neq t, \\ A_t(z^{-1}) - p_t(x^{-1}) & \text{if } i = t. \end{cases}$$

Then $\operatorname{Add}_{p_t(x^{-1})}^{(t)}(\mathbf{A}) \in \prod_{i=0}^p \mathcal{O}_{B'_i}$ where

$$B'_i := \begin{cases} B_i & \text{if } i \neq t, \\ B_t - p_t(z^{-1}) & \text{if } i = t. \end{cases}$$

Set

$$\mathcal{J}_i := \{[i,j] \mid j = 1, \dots, m_i\}$$

for $i = 0, \ldots, p$ and

$$\mathcal{J} := \prod_{i=0}^p \mathcal{J}_i.$$

Then let us define

$$Add_{\mathbf{i}} := \prod_{i=0}^{p} Add_{q_{[i,j_i]}(z^{-1}) + \xi_1^{[i,j_i]} z^{-1}},$$

for $\mathbf{i} = ([i, j_i])_{0 \le i \le p} \in \mathcal{J}$. Here we use the notation $\prod_{i \in \{a, b, \dots, \}} f_i = f_a \circ$ $f_b \circ \cdots$ and note that the operators $\operatorname{Ad}_{q_{[i,j_i]}(z^{-1})+\xi_1^{[i,j_i]}z^{-1}}^{(i)}$ for $i \in [0, \dots, p]$ are commutative.

Take $\mathbf{A} = (A_i(z^{-1}))_{0 \le i \le p} \in \prod_{i=0}^p \mathcal{O}_{B_i}$ satisfying $\sum_{i=0}^p \operatorname{Res} A_i(z^{-1}) = 0.$ Suppose that we can choose $\mathbf{i} \in \mathcal{J}$ so that

$$\xi_{\mathbf{i}} := \sum_{i=0}^{p} \xi_{1}^{[i,j_{i}]} \neq 0.$$

Let (V, W, T, Q, P) be the canonical datum of Add_i(A). Following Example 3 in [37], we construct a new 5-tuple (V', W, T, Q', P') as follows. Note that $QP = -\xi_i Id_V$. Thus Q and P are surjective and injective respectively. Let us set $V' := \operatorname{Coker} P$ and $Q' \colon W \to V'$, the natural projection. Then we have the split exact sequence

$$0 \longrightarrow V \xrightarrow{P} W \xrightarrow{Q'} V' \longrightarrow 0$$

with the left splitting $(-\xi_{\mathbf{i}}^{-1}Q)P = \mathrm{Id}_V$. Then from the splitting, we can define $P': V' \to W$ be the injection such that $Q'(\xi_i^{-1}P') = \mathrm{Id}_{V'}$. Then we have a 5-tuple (V', W, T, Q', P').

Next we set Q'_i (resp. P'_i) to be the $\operatorname{Hom}_{\mathbb{C}}(W_i, V)$ (resp. $\operatorname{Hom}_{\mathbb{C}}(V, W_i)$) component of Q' (resp. P'). Also set N_i to be the $\operatorname{End}_{\mathbb{C}}(W_i)$ -component of T. Define

$$(A')_{j}^{(i)} := Q'_{i} N_{i}^{j-1} P'_{i}$$

and $\mathbf{A}' := (A'_i(z^{-1}))_{0 \le i \le p}$ where $A'_i(z^{-1}) := \sum_{i=1}^{k_i} (A')_i^{(i)} z^{-j}$. We note that $\sum_{i=0}^{p} (A')_{1}^{(i)} = Q'P' = \xi_{\mathbf{i}} \mathrm{Id}_{V'}.$

Finally let us set

$$\mathbf{A}'' := \mathrm{Add}_{\mathbf{i}}^{-1} \circ \mathrm{Add}_{2\xi_{\mathbf{i}}z^{-1}}^{(0)}(\mathbf{A}').$$

Then $\mathbf{A}'' = (A''_i(x^{-1}))_{0 \le i \le p}$ satisfies that $\sum_{i=0}^{p} \operatorname{Res} A''_i(x^{-1}) = 0$. Let us denote \mathbf{A}'' by $\mathrm{mc}_{\mathbf{i}}(\mathbf{A})$ and call the operator $\mathrm{mc}_{\mathbf{i}}$ the *middle convolution* at \mathbf{i} . Let us recall basic properties of middle convolutions.

Lemma 5.3. Let (V, W, T, Q, P) be the canonical datum of the above $mc_i(\mathbf{A})$. Then $\dim_{\mathbb{C}} W = \sum_{i=0}^{p} w_i$ where

$$w_{i} := \sum_{j=1}^{m_{i}} \left((d_{i}(j, j_{i}) + 1) \dim_{\mathbb{C}} V_{[i,j]} \right) \\ + (n - \dim_{\mathbb{C}} V_{[i,j_{i}]}) + \operatorname{rank}(R_{[i,j_{i}]} - \xi_{[i,j_{i},1]}).$$

Proof. From Lemma 5.2, we have

$$w_i = \sum_{t=0}^{k_i-1} \left(n - \dim_{\mathbb{C}} \bigcap_{l=0}^t \operatorname{Ker} \tilde{B}_{k_i-l}^{(i)} \right)$$

where $\tilde{B}_i := B_i - \left(q_{[i,j_i](z^{-1}) + \xi_1^{[i,j_i]}z^{-1}} \right) I_n$ for i = 0, ..., p. Suppose that $0 \le 1$ $t \leq k_i - 2$. Then we have $V_{[i,j]} \subset \bigcap_{l=0}^t \operatorname{Ker} \tilde{B}_{k_i-l}^{(i)}$ if and only if $\deg_{\mathbb{C}[z^{-1}]}(q_{[i,j]} - 1)$

 $q_{[i,j_i]}) < k_i - t$. Equivalently, we have

$$n - \dim_{\mathbb{C}} \bigcap_{l=0}^{t} \operatorname{Ker} \tilde{B}_{k_{i}-l}^{(i)} = \sum_{j \in \{j' \mid d_{i}(j',j_{i}) \ge k_{i}-t-2\}} \dim_{\mathbb{C}} V_{[i,j]}$$

for $0 \le t \le k_i - 2$. Thus

$$\sum_{t=0}^{k_i-2} \left(n - \dim_{\mathbb{C}} \bigcap_{l=0}^t \operatorname{Ker} \tilde{B}_{k_i-l}^{(i)} \right) = \sum_{j=0}^{m_i} (d_i(j,j_i) + 1) \dim_{\mathbb{C}} V_{[i,j]}.$$

Combining this computation and the fact

$$\dim_{\mathbb{C}} \operatorname{Ker} \bigcap_{l=0}^{k_i-1} \tilde{B}_{k_i-l}^{(i)} = \dim_{\mathbb{C}} \operatorname{Ker}(R_{[i,j_i]} - \xi_1^{[i,j_i]}),$$

we obtain the required result.

Proposition 5.4 (Yamakawa [37]). Let us take $\mathbf{A} = (A_i(z^{-1}))_{0 \le i \le p} \in \prod_{i=0}^p \mathcal{O}_{B_i}$ satisfying $\sum_{i=0}^p \operatorname{Res} A_i(z^{-1}) = 0$. Suppose we can choose $\mathbf{i} \in \mathcal{J}$ so that $\xi_{\mathbf{i}} \ne 0$.

- (1) If \mathbf{A} is irreducible, then $mc_i(\mathbf{A})$ is irreducible.
- (2) If \mathbf{A} is irreducible,

$$mc_i \circ mc_i(\mathbf{A}) \sim \mathbf{A}$$

i.e., there exists $g \in GL(n, \mathbb{C})$ such that

$$\mathrm{mc}_{\mathbf{i}} \circ \mathrm{mc}_{\mathbf{i}}(\mathbf{A}) = g\mathbf{A}g^{-1} := (gA_i(z^{-1})g^{-1})_{0 \le i \le p}$$

(3) Let us define elements in $M(n'_{[i,j]}, \mathbb{C})$ by

$$R'_{[i,j]} := \begin{cases} R_{[i,j]} + (d_i(j,j_i) + 2)\xi_{\mathbf{i}}I_{n_{[i,j]}} & \text{if } i \neq 0, \\ R_{[0,j]} + d_0(j,j_0)\xi_{\mathbf{i}}I_{n_{[0,j]}} & \text{if } i = 0 \end{cases}$$

for all $i \in \{0, \ldots, p\}$ and $j \in \{1, \ldots, m_i\} \setminus \{j_i\}$. Here we put $n'_{[i,j]} := n_{[i,j]}$.

Further define $R'_{[i,j_i]} \in M(n'_{[i,j_i]}, \mathbb{C})$ for i = 1, ..., p so that equations hold,

$$\operatorname{rank} (R'_{[i,j_i]} - \xi_1^{[i,j_i]}) = \operatorname{rank} (R_{[i,j_i]} - \xi_1^{[i,j_i]}),$$

$$\operatorname{rank} (R'_{[i,j_i]} - \xi_1^{[i,j_i]}) \prod_{k=2}^{l} (R'_{[i,j_i]} - \xi_k^{[i,j_i]} - \xi_i)$$

$$= \operatorname{rank} \prod_{k=1}^{l} (R_{[i,j_i]} - \xi_k^{[i,j_i]}), \quad l = 2, \dots, e_{[i,j_i]}.$$

Similarly define $R'_{[0,j_0]} \in M(n'_{[0,j_0]}, \mathbb{C})$ so that the following equations hold.

$$\operatorname{rank} \left(R_{[0,j_0]}' - \xi_1^{[0,j_0]} + 2\xi_{\mathbf{i}} \right) = \operatorname{rank} \left(R_{[0,j_0]} - \xi_1^{[0,j_0]} \right),$$

$$\operatorname{rank} \left(R_{[0,j_0]}' - \xi_1^{[0,j_0]} + 2\xi_{\mathbf{i}} \right) \prod_{k=2}^l \left(R_{[0,j_0]}' - \xi_k^{[0,j_0]} + \xi_{\mathbf{i}} \right)$$

$$= \operatorname{rank} \prod_{k=1}^l \left(R_{[0,j_0]} - \xi_k^{[0,j_0]} \right), \quad l = 2, \dots, e_{[0,j_0]}$$

Here we put

$$n'_{[i,j_i]} := n_{[i,j_i]} + \dim_{\mathbb{C}} W - 2n.$$

Finally define

$$B'_{i} := \operatorname{diag} \left(q_{[i,1]}(z^{-1})I_{n'_{[i,1]}} + R'_{[i,1]}z^{-1}, \dots, q_{[i,m_{i}]}(z^{-1})I_{n'_{[i,m_{i}]}} + R'_{[i,m_{i}]}z^{-1} \right) for \ i = 0, \dots, p. Then \ \operatorname{mc}_{\mathbf{i}}(\mathbf{A}) \in \prod_{i=0}^{p} \mathcal{O}_{B'_{i}}.$$

Proof. Corollary 4 in [37] shows (1). Lemma 11, Remark 14 and Proposition 16 in [37] show (2).

By this proposition, we obtain the bijection

 $\mathrm{mc}_{\mathbf{i}} \colon \mathfrak{M}(\mathbf{B}) \longrightarrow \mathfrak{M}(\mathbf{B}')$

where $\mathbf{B}' := (B'_i)_{0 \le i \le p}$ defined in the proposition.

Proposition 5.4 shows the following.

Proposition 5.5. Let ξ and $\mathfrak{M}_{\lambda}(\mathbf{Q}, \alpha)^{dif}$ be same as in Theorem 4.22. Suppose that we can choose $\mathbf{i} = ([i, j_i])_{0 \leq i \leq p} \in \mathcal{J}$ so that

$$\lambda_{\mathbf{i}} := \sum_{i \in I_{irr}} \lambda_{[i,j_i]} = -\xi_{\mathbf{i}} \neq 0$$

Define $\mathrm{mc}_{\mathbf{i}}(\alpha) := (\alpha'_a)_{a \in \mathsf{Q}_0} \in \mathbb{Z}^{\mathsf{Q}_0}$ and $\mathrm{mc}_{\mathbf{i}}(\lambda) := (\lambda'_a)_{a \in \mathsf{Q}_0} \in \mathbb{C}^{\mathsf{Q}_0}$ by

$$\begin{split} \alpha'_{[i,j]} &:= \begin{cases} \alpha_{[i,j]} & \text{if } j \neq j_i, \\ \alpha_{[i,j_i]} + n_{\mathbf{i}} & \text{if } j = j_i, \end{cases} \\ \alpha'_{[i,j,k]} &:= \alpha_{[i,j,k]}, \end{cases} \\ \lambda'_{[i,j]} &:= \begin{cases} \lambda_{[i,j_i]} & \text{if } [i,j] = [i,j_i] \text{ and } i \neq 0, \\ \lambda_{[0,j_0]} - 2\lambda_{\mathbf{i}} & \text{if } [i,j] = [0,j_0], \\ \lambda_{[i,j]} + (d_i(j,j_i) + 2)\lambda_{\mathbf{i}} & \text{if } i \neq 0 \text{ and } j \neq m_i, \\ \lambda_{[0,j]} + d_0(j,j_0)\lambda_{\mathbf{i}} & \text{if } i = 0 \text{ and } j \neq m_0, \end{cases} \\ \lambda'_{[i,j,k]} &:= \begin{cases} \lambda_{[i,j,k]} & \text{if } [i,j,k] \neq [i,j_i,1], \\ \lambda_{[i,j,1]} + \lambda_{\mathbf{i}}. \end{cases} \end{split}$$

Here

$$n_{\mathbf{i}} := \sum_{i \in I_{irr}} \sum_{j=1}^{m^{(i)}} (d_i(j, j_i) + 1) \alpha_{[i,j]} + \sum_{i \in I_{irr}} ((n - \alpha_{[i,j_i]}) + \alpha_{[i,j_i,1]}) + \sum_{i \in I_{reg}} \alpha_{[i,1,1]} - 2n.$$

Then there exists a bijection

$$\mathrm{mc}_{\mathbf{i}} \colon \mathfrak{M}_{\lambda}(\mathsf{Q}, \alpha)^{dif} \longrightarrow \mathfrak{M}_{\mathrm{mc}_{\mathbf{i}}(\lambda)}(\mathsf{Q}, \mathrm{mc}_{\mathbf{i}}(\alpha))^{dif}.$$

Proof. We already have the bijection

$$\mathrm{mc}_i\colon\mathfrak{M}(\mathbf{B})\longrightarrow\mathfrak{M}(\mathbf{B}')$$

for **B'** defined as in Proposition 5.4. If we choose $\xi' = \left(\left\{(\xi')_1^{[i,j]}, \dots, (\xi')_{e_{[i,j]}}^{[i,j]}\right\}\right)_{\substack{0 \le i \le p\\1 \le j \le m^{(i)}}}$ so that

$$(\xi')_{k}^{[i,j]} = \begin{cases} \xi_{k}^{[i,j]} + (d_{i}(j,j_{i}) + 2)\xi_{\mathbf{i}} & \text{if } i \neq 0 \text{ and } j \neq j_{i}, \\ \xi_{k}^{[0,j]} + d_{0}(j,j_{0})\xi_{\mathbf{i}} & \text{if } i = 0 \text{ and } j \neq j_{0}, \\ \xi_{1}^{[i,j_{i}]} & \text{if } i \neq 0, \ j = j_{i} \text{ and } k = 1, \\ \xi_{k}^{[i,j_{i}]} + \xi_{\mathbf{i}} & \text{if } i \neq 0, \ j = j_{i} \text{ and } k \neq 1, \\ \xi_{1}^{[0,j_{0}]} - 2\xi_{\mathbf{i}} & \text{if } i = 0, \ j = j_{0} \text{ and } k = 1, \\ \xi_{k}^{[0,j_{0}]} - \xi_{\mathbf{i}} & \text{if } i = 0, \ j = j_{0} \text{ and } k = 1, \end{cases}$$

then we have the bijection

$$\Phi_{\xi'} \colon \mathfrak{M}(\mathbf{B}') \longrightarrow \mathfrak{M}_{\mathrm{mc}_{\mathbf{i}}(\lambda)}(\mathsf{Q}, \mathrm{mc}_{\mathbf{i}}(\alpha))^{\mathrm{dif}}$$

by Theorem 4.22. Thus we have the bijection $\Phi_{\xi'} \circ \mathrm{mc}_{\mathbf{i}} \circ \Phi_{\xi}^{-1} \colon \mathfrak{M}_{\lambda}(\mathsf{Q}, \alpha)^{\mathrm{dif}} \longrightarrow$ $\mathfrak{M}_{\mathrm{mc}_{\mathbf{i}}(\lambda)}(\mathsf{Q},\mathrm{mc}_{\mathbf{i}}(\alpha))^{\mathrm{dif}}.$

The last equation is obtained by Lemma 5.3 as follows. Since $n_{\mathbf{i}} = n'_{[i,j_i]} - n'_{[i,j_i]}$ $n_{[i,j_i]}$, we have

$$\begin{split} n_{\mathbf{i}} &= n'_{[i,j_i]} - n_{[i,j_i]} = \dim_{\mathbb{C}} W - 2n \\ &= \sum_{i \in I_{\mathrm{irr}}} \sum_{\substack{1 \le j \le m_i \\ j \ne j_i}} d_i(j,j_i) \alpha_{[i,j]} + \sum_{i \in I_{\mathrm{irr}}} (2(n - \alpha_{[i,j_i]}) + \alpha_{[i,j_i,1]}) \\ &+ \sum_{i \in I_{\mathrm{reg}}} \alpha_{[i,1,1]} - 2n \\ &= \sum_{i \in I_{\mathrm{irr}}} \sum_{j=1}^{m_i} (d_i(j,j_i) + 1) \alpha_{[i,j]} + \sum_{i \in I_{\mathrm{irr}}} ((n - \alpha_{[i,j_i]}) + \alpha_{[i,j_i,1]}) \\ &+ \sum_{i \in I_{\mathrm{reg}}} \alpha_{[i,1,1]} - 2n. \end{split}$$

5.2. More operations on $\mathfrak{M}_{\lambda}(\mathsf{Q},\alpha)^{\mathrm{dif}}$. Proposition 5.5 enable us to define mc_i as an operation for $\mathfrak{M}_{\lambda}(\mathsf{Q},\alpha)^{\mathrm{dif}}$. We shall introduce some other operations.

Let us note that the map Φ_{ξ} in Proposition 4.20 depends on the order of $\{\xi_1^{[i,j]}, \ldots, \xi_{e_{[i,j]}}^{[i,j]}\}$ for $i = 0, \ldots, p$ and $j = 1, \ldots, m_i$. Thus we shall see what happens when we change the order. Let us define

$$\sigma_s^{[i_0,j_0]}(\xi) := (\{\zeta_1^{[i,j]}, \dots, \zeta_{e_{[i,j]}}^{[i,j]}\})_{\substack{0 \le i \le p\\ 1 \le j \le m_i}}$$

by

$$\zeta_l^{[i,j]} := \begin{cases} \xi_l^{[i,j]} & \text{if } [i,j] \neq [i_0,j_0], \\ \xi_{\sigma_s(l)}^{[i_0,j_0]} & \text{if } [i,j] = [i_0,j_0] \end{cases}$$

where σ_s is the permutation (s, s + 1). Then we can extend these permutations to operations on representations of $\overline{\mathsf{Q}}$,

$$\sigma_s^{[i_0,j_0]} \colon \mathfrak{M}_{\lambda}(\mathsf{Q},\alpha)^{\mathrm{dif}} \longrightarrow \mathfrak{M}_{\lambda'''}(\mathsf{Q},\alpha''')^{\mathrm{dif}}$$

defined by $\Phi_{\sigma_s^{[i_0,j_0]}(\xi)} \circ \Phi_{\xi}^{-1}$. Here α''' and λ''' can be computed as below.

Proposition 5.6. The above α''' and λ''' are defined as follows,

$$\begin{aligned} \alpha''' &= \begin{cases} \alpha & \text{if } \xi_s^{[i_0,j_0]} = \xi_{s+1}^{[i_0,j_0]}, \\ s_{[i_0,j_0,s]}(\alpha) & \text{otherwise}, \end{cases} \\ \lambda''' &= \begin{cases} \lambda & \text{if } \xi_s^{[i_0,j_0]} = \xi_{s+1}^{[i_0,j_0]}, \\ r_{[i_0,j_0,s]}(\lambda) & \text{otherwise}. \end{cases} \end{aligned}$$

Proof. If $\xi_s^{[i_0,j_0]} = \xi_{s+1}^{[i_0,j_0]}$, then $\Phi_{\sigma_s^{[i_0,j_0]}(\xi)} \circ \Phi_{\xi}^{-1} = \Phi_{\xi} \circ \Phi_{\xi}^{-1} = \text{id.}$ Thus we obtain the result. Next we suppose $\xi_s^{[i_0,j_0]} \neq \xi_{s+1}^{[i_0,j_0]}$. To see that $\alpha' = s_{[i_0,j_0,s]}(\alpha)$, it suffices to show that

$$\operatorname{rank} \prod_{l=1}^{s-1} (R_{[i_0,j_0]} - \xi_l^{[i_0,j_0]}) - \operatorname{rank} \prod_{l=1}^s (R_{[i_0,j_0]} - \xi_l^{[i_0,j_0]})$$
$$= \operatorname{rank} (R_{[i_0,j_0]} - \xi_{s+1}^{[i_0,j_0]}) \prod_{l=1}^{s-1} (R_{[i_0,j_0]} - \xi_l^{[i_0,j_0]}) - \operatorname{rank} \prod_{l=1}^{s+1} (R_{[i_0,j_0]} - \xi_l^{[i_0,j_0]}),$$

and

$$\begin{aligned} & \operatorname{rank} \prod_{l=1}^{s} (R_{[i_{0},j_{0}]} - \xi_{l}^{[i_{0},j_{0}]}) - \operatorname{rank} \prod_{l=1}^{s+1} (R_{[i_{0},j_{0}]} - \xi_{l}^{[i_{0},j_{0}]}) \\ & = \operatorname{rank} \prod_{l=1}^{s-1} (R_{[i_{0},j_{0}]} - \xi_{l}^{[i_{0},j_{0}]}) - \operatorname{rank} (R_{[i_{0},j_{0}]} - \xi_{s+1}^{[i_{0},j_{0}]}) \prod_{l=1}^{s-1} (R_{[i_{0},j_{0}]} - \xi_{l}^{[i_{0},j_{0}]}). \end{aligned}$$

These equations follow from the following fact. Let us suppose that $A, A' \in M(n, \mathbb{C})$ satisfy AA' = A'A and $\operatorname{Ker} A \cap \operatorname{Ker} A' = \{0\}$. Let V be an A'-invariant subspace of \mathbb{C}^n . Then we have

(5)
$$\dim_{\mathbb{C}} V - \dim_{\mathbb{C}} A' V = \dim_{\mathbb{C}} A V - \dim_{\mathbb{C}} A A' V.$$

Indeed, setting $\widetilde{V} = V \cap \operatorname{Ker} A$ and $\widetilde{W} = A'V \cap \operatorname{Ker} A$, we have that A' gives an injection from $\widetilde{V} \to \widetilde{W}$ since $\operatorname{Ker} A \cap \operatorname{Ker} A' = \{0\}$. Since $\widetilde{W} \subset \widetilde{V}$, this implies that $\widetilde{W} = \widetilde{V}$, which shows the equation (5).

Since $\xi_s^{[i_0,j_0]} \neq \xi_{s+1}^{[i_0,j_0]}$, $(R_{[i_0,j_0]} - \xi_s^{[i_0,j_0]})$ and $(R_{[i_0,j_0]} - \xi_{s+1}^{[i_0,j_0]})$ satisfy the above assumption. Then equation (5) gives the required equations.

The remaining assertion follows from a direct computation.

Furthermore let us introduce an operation which is trivial on $\mathfrak{M}(\mathbf{B})$. Let $t_{(i,0)}$ be the permutation (i,0) for $i \in I_{irr} \setminus \{0\}$. Define

$$t_{(i,0)}(\mathbf{B}) := (B_{t_{(i,0)}(j)})_{j=0,\dots,p}$$

Then we can define the bijection

$$\begin{array}{rccc} T_{(i,0)} \colon & \mathfrak{M}(\mathbf{B}) & \longrightarrow & \mathfrak{M}(t_{(i,0)}(\mathbf{B})) \\ & (A_j(z^{-1}))_{0 \le j \le p} & \longmapsto & (A_{t_{(i,0)}(j)}(z^{-1}))_{0 \le j \le p} \end{array}$$

Then $t_{(i,0)}(\xi)$, $t_{(i,0)}(\mathbb{Q})$, $t_{(i,0)}(\alpha)$ and $t_{(i,0)}(\lambda)$ can be defined by replacing i with 0 and vice versa. Thus we can define the bijection

$$T_{(i,0)} \colon \mathfrak{M}_{\lambda}(\mathsf{Q},\alpha)^{\mathrm{dif}} \longrightarrow \mathfrak{M}_{t_{(i,0)}(\lambda)}(t_{(i,0)}(\mathsf{Q}), t_{(i,0)}(\alpha))^{\mathrm{dif}}$$

by $\Phi_{t_{(i,0)}(\xi)} \circ T_{(i,0)} \circ \Phi_{\xi}^{-1}$.

5.3. Middle convolution and reflection. As we saw in the above proposition which shows that permutations on ξ can be obtained by reflection, middle convolutions mc_i can also be obtained by reflections as follows.

For $\mathbf{i} = ([i, j_i])_{0 \le i \le p} \in \mathcal{J}$, let us define $\epsilon_{\mathbf{i}} \in \mathbb{Z}^{\mathsf{Q}_0}$ by

$$(\epsilon_{\mathbf{i}})_a := \begin{cases} 1 & \text{if } a = [i, j_i], i \in I_{\text{irr}}, \\ 0 & \text{otherwise.} \end{cases}$$

We note that ϵ_i for $i \in \mathcal{J}$ are positive real roots of Q. Let us define

$$s_{\mathbf{i}}(\beta) := \beta - (\beta, \epsilon_{\mathbf{i}})\epsilon_{\mathbf{i}}$$

for $\mathbf{i} \in \mathcal{J}$ and $\beta \in \mathbb{Z}^{Q_0}$.

Let us see this reflection s_i can be obtained by a product of simple reflections.

Lemma 5.7. Let us take $\mathbf{i} = ([i, j_i])_{0 \le i \le p} \in \mathcal{J}$. Then we have

$$\left(\prod_{i\in I_{irr}\setminus\{0\}}s_{[i,j_i]}\right)\circ s_{[0,j_0]}\circ\left(\prod_{i\in I_{irr}\setminus\{0\}}s_{[i,j_i]}\right)(\beta)=s_{\mathbf{i}}(\beta)$$

for any $\beta \in \mathbb{Z}^{\mathsf{Q}_0}$.

Proof. Set $r := \prod_{i \in I_{irr} \setminus \{0\}} s_{[i,j_i]}$ for short. Note that r is an involution and $\epsilon_{\mathbf{i}} = r(\epsilon_{[0,j_0]})$. Then

$$\begin{aligned} r \circ s_{[0,j_0]} \circ r(\beta) &= r(r(\beta) - (r(\beta), \epsilon_{[0,j_0]}) \epsilon_{[0,j_0]}) \\ &= r^2(\beta) - (\beta, r^{-1}(\epsilon_{[0,j_0]})) r(\epsilon_{[0,j_0]}) \\ &= \beta - (\beta, r(\epsilon_{[0,j_0]})) r(\epsilon_{[0,j_0]}) \\ &= \beta - (\beta, \epsilon_{\mathbf{i}}) \epsilon_{\mathbf{i}} \\ &= s_{\mathbf{i}}(\beta). \end{aligned}$$

This lemma tells us that mc_i can be regarded as a reflection and a product of simple reflections as below.

Proposition 5.8. Retain the notation in Proposition 5.5. Then we have

$$\mathrm{mc}_{\mathbf{i}}(\alpha) = s_{\mathbf{i}}(\alpha)$$
$$= \left(\prod_{i \in I_{irr} \setminus \{0\}} s_{[i,j_i]}\right) \circ s_{[0,j_0]} \circ \left(\prod_{i \in I_{irr} \setminus \{0\}} s_{[i,j_i]}\right) (\alpha).$$

Proof. From the definition of $mc_i(\alpha)$ given in Proposition 5.5, it suffices to show

$$n_{\mathbf{i}} = -(\alpha, \epsilon_{\mathbf{i}}).$$

Indeed

$$\begin{aligned} (\alpha, \epsilon_{\mathbf{i}}) &= \sum_{i \in I_{\mathrm{irr}}} (\alpha, \epsilon_{[i,j_i]}) \\ &= \sum_{i \in I_{\mathrm{irr}} \setminus \{0\}} \left(2\alpha_{[i,j_i]} - \sum_{\substack{1 \le j \le m_i \\ j \ne j_i}} d_i(j,j_i) \alpha_{[i,j]} - \alpha_{[i,j_i,1]} - \sum_{j=1}^{m_0} \alpha_{[0,j]} \right) \\ &+ 2\alpha_{[0,j_0]} - \sum_{\substack{1 \le j \le m_0 \\ j \ne j_0}} d_0(j,j_0) \alpha_{[0,j]} - \alpha_{[0,j_0,1]} - \sum_{i \in I_{\mathrm{irr} \setminus \{0\}}} \sum_{j=1}^{m_i} \alpha_{[i,j]} \\ &- \sum_{i \in I_{\mathrm{reg}} \alpha_{[i,1,1]}} \alpha_{[i,j]}. \end{aligned}$$

Recalling that $d_i(j,j) = -1$ and $\sum_{j=1}^{m_i} \alpha_{[i,j]} = n$, we can continue the above computation,

$$\begin{split} (\alpha, \epsilon_{\mathbf{i}}) &= -\sum_{i \in I_{\mathrm{irr}}} \left(\sum_{j=1}^{m_{i}} (d_{i}(j, j_{i}) \alpha_{[i,j]}) + (n - \alpha_{[i,j_{i}]}) + \alpha_{[i,j_{i},1]} \right) \\ &- \sum_{i \in I_{\mathrm{reg}}} \alpha_{[i,1,1]} - (\#I_{\mathrm{irr}} - 2)n \\ &= -\sum_{i \in I_{\mathrm{irr}}} \left(\sum_{j=1}^{m_{i}} (d_{i}(j, j_{i} + 1) \alpha_{[i,j]}) + (n - \alpha_{[i,j_{i}]}) + \alpha_{[i,j_{i},1]} \right) \\ &- \sum_{i \in I_{\mathrm{reg}}} \alpha_{[i,1,1]} + 2n \\ &= -n_{\mathbf{i}}. \end{split}$$

These observations lead us to define transformations on $\mathbb{Z}^{\mathsf{Q}_0} \times \mathbb{C}^{\mathsf{Q}_0}$ as an analogy of middle convolutions and other operations. For $(\beta, \nu) \in \mathbb{Z}^{\mathsf{Q}_0} \times \mathbb{C}^{\mathsf{Q}_0}$ with $\nu_{\mathbf{i}} = \sum_{i \in I_{\mathrm{irr}}} \nu_{[i,j_i]} \neq 0$ define

$$s_{\mathbf{i}}((\beta,\nu)) := (\mathrm{mc}_{\mathbf{i}}(\beta'), \mathrm{mc}_{\mathbf{i}}(\nu')).$$

Also for $(\beta, \nu) \in \mathbb{Z}^{\mathsf{Q}_0} \times \mathbb{C}^{\mathsf{Q}_0}$ with $\nu_{[i,j,k]} \neq 0$, define

$$s_{[i,j,k]}((\beta,\nu)) := (s_{[i,j,k]}(\beta), r_{[i,j,k]}(\nu)).$$

Let us define

$$\mathcal{S} := \left\{ \left(\beta, \nu\right) \in \mathcal{L} \times \mathbb{C}^{\mathsf{Q}_0} \, \middle| \, \beta \cdot \nu = \sum_{a \in \mathsf{Q}_0} \beta_a \nu_a = 0 \right\}.$$

`

Then we can see that s_i and $s_{[i,j,k]}$ preserve \mathcal{S} , see Propositions 5.5 and 5.6.

5.4. Irreducibility and \mathcal{L} -irreducibility. The \mathcal{L} -irreducibility is a weaker condition than the usual irreducibility. We shall show that if we shift the parameter λ by using the operation Add, then these two irreducibility can be identical.

Fix $i_0 \in I_{irr} \setminus \{0\}$ and define an operation on S as an analogue of $\operatorname{Add}_{z^{-1}}^{(i_0)} \circ \operatorname{Add}_{-z^{-1}}^{(0)}$ as follows. Let us define $z^{(i_0)} = (z_a^{(i_0)})_{a \in \mathsf{Q}_0} \in \mathbb{C}^{\mathsf{Q}_0}$ by

$$z_{[i,j]}^{(i_0)} = \begin{cases} 1 & \text{if } i = i_0, \\ -1 & \text{if } i = 0, \\ 0 & \text{otherwise}, \end{cases}$$
$$z_{[i,j,k]}^{(i_0)} = 0.$$

Then let us define

$$\operatorname{add}_{\gamma}^{(i_0)} \colon \begin{array}{ccc} \mathcal{S} & \longrightarrow & \mathcal{S} \\ (\beta, \nu) & \longmapsto & (\beta, \nu + \gamma z^{(i_0)}) \end{array}$$

for $i_0 \in I_{irr} \setminus \{0\}$ and $\gamma \in \mathbb{C}$.

For $\nu \in \mathbb{C}^{Q_0}$, let R^+_{ν} be the set of positive roots β of \mathbb{Q} satisfying $\beta \cdot \nu = 0$. Denote $\mathcal{L} \cap R^+_{\nu}$ by \widetilde{R}^+_{ν} . The subset Σ_{ν} of R^+_{ν} consists of β satisfying that $p(\beta) > \sum_t p(\beta_t)$ for any decomposition $\beta = \beta_1 + \cdots + \beta_r$ with $r \ge 2$ and $\beta_t \in R^+_{\nu}$. Similarly define $\widetilde{\Sigma}_{\nu}$ consisting of $\beta \in \widetilde{R}^+_{\nu}$ satisfying that $\beta \cdot \nu = 0$ and $p(\beta) > \sum_t p(\beta_t)$ for any decomposition $\beta = \beta_1 + \cdots + \beta_r$ with $r \ge 2$ and $\beta_t \in \widetilde{R}^+_{\nu}$.

If $\beta, \beta' \in (\mathbb{Z}_{\geq 0})^{\mathbb{Q}_0}$ satisfy that $\beta'_a \leq \beta_a$ for all $a \in \mathbb{Q}_0$, then we write $\beta' \leq \beta$.

Lemma 5.9. Fix $(\beta, \nu) \in S$. There exist $\gamma_i \in \mathbb{C}$ for $i \in I_{irr} \setminus \{0\}$ such that $\nu' = \nu + \sum_{i \in I_{irr} \setminus \{0\}} \gamma_i z^{(i)}$ satisfies the following. If $\beta' \in (\mathbb{Z}_{\geq 0})^{\mathbb{Q}_0}$ satisfies that $\beta' \leq \beta$ and $\beta' \cdot \nu' = 0$, then $\beta' \in \mathcal{L}$.

Proof. Let F_{β} be the set of all elements β' in $(\mathbb{Z}_{\geq 0})^{\mathbb{Q}_0}$ satisfying $\beta' \leq \beta$ and $\beta' \notin \mathcal{L}$. Note that F_{β} is a finite set. Define a closed subset of $\mathbb{C}^{\mathbb{Q}_0}$ by

$$V_{\beta} := \bigcup_{\beta' \in F_{\beta}} \{ \eta \in \mathbb{C}^{\mathsf{Q}_0} \mid \beta' \cdot \eta = 0 \}.$$

Namely, if $\nu \notin V_{\beta}$, then $\beta' \cdot \nu = 0$ and $\beta' \leq \beta$ imply $\beta' \in \mathcal{L}$. Thus let us suppose $\nu \in V_{\beta}$. Consider the affine space $W_{\nu} := \{\nu + \sum_{i \in I_{\mathrm{irr}} \setminus \{0\}} t_i z^{(i)} \mid t_i \in \mathbb{C}\}$. Then $W_{\nu} \cap \{\eta \in \mathbb{C}^{\mathsf{Q}_0} \mid \beta' \cdot \eta = 0\}$ is a proper closed subset of W_{μ} for any $\beta' \in F_{\beta}$. Indeed, since $\beta' \notin \mathcal{L}$ there exists $i_0 \in I_{\mathrm{irr}} \setminus \{0\}$ such

that $\sum_{j=1}^{m^{(0)}} \beta'_{[0,j]} \neq \sum_{j=1}^{m^{(i_0)}} \beta'_{[i_0,j]}$. Then the line $\{\nu + tz^{(i_0)} \mid t \in \mathbb{C}\} \subset W_{\nu}$ is not contained in the hyperplane $\{\eta \in \mathbb{C}^{\mathsf{Q}_0} \mid \beta' \cdot \eta = 0\}$. Thus dim $W_{\nu} >$ dim $(W_{\nu} \cap \{\eta \in \mathbb{C}^{\mathsf{Q}_0} \mid \beta' \cdot \eta = 0\})$ for any $\beta' \in F_{\beta}$ since W_{μ} is an irreducible algebraic set. This shows the inequality,

 $\dim W_{\nu} \cap V_{\beta} = \max_{\beta' \in F_{\beta}} \left\{ \dim \left(W_{\nu} \cap \{ \eta \in \mathbb{C}^{\mathsf{Q}_0} \mid \beta' \cdot \eta = 0 \} \right) \right\} < \dim W_{\nu}.$

Hence there exists $\nu' \in W_{\nu}$ which is not contained in V_{β} as required. \Box

Lemma 5.10. We have $\widetilde{\Sigma}_{\nu} = \widetilde{\Sigma}_{\nu+\gamma z^{(i)}}$ for any $i \in I_{irr} \setminus \{0\}$ and $\gamma \in \mathbb{C}$.

Proof. Recall that $(\beta, \nu) \in S$ if and only if $(\beta, \nu + \gamma z^{(i)}) \in S$.

Obviously $\mathcal{L} \cap \Sigma_{\nu} \subset \widetilde{\Sigma}_{\nu}$. The above lemmas show that for any $\beta \in \widetilde{\Sigma}_{\nu}$, there exists $\nu' \in \mathbb{C}^{Q_0}$ such that $\beta \in \Sigma_{\nu'}$.

Proposition 5.11. For any $\beta \in \widetilde{\Sigma}_{\nu}$, there exist $\gamma_i \in \mathbb{C}$ for $i \in I_{irr} \setminus \{0\}$ such that $\beta \in \Sigma_{\nu + \sum_{i \in I_{irr} \setminus \{0\}} \gamma_i z^{(i)}}$.

Proof. For $\beta \in \widetilde{\Sigma}_{\nu}$, let us choose γ_i as in Lemma 5.9 and set $\nu' = \nu + \sum_{i \in I_{irr} \setminus \{0\}} \gamma_i z^{(i)}$. Then Lemma 5.9 shows that $\beta \in \Sigma_{\nu'}$.

Lemma 5.12. Suppose that $\mu^{-1}(\lambda)^{dif} \neq \emptyset$. Fix $i_0 \in I_{irr}$ and $\gamma \in \mathbb{C}$. Then there exists a $\mathbf{G}(\alpha)$ -equivariant analytic bijection

$$\operatorname{add}_{\gamma}^{(i_0)} \colon \mu_{\alpha}^{-1}(\lambda)^{\operatorname{dif}} \longrightarrow \mu_{\alpha}^{-1}(\lambda + \gamma z^{(i_0)})^{\operatorname{dif}}.$$

Proof. The required map is obtained by $\Phi_{\xi'} \circ \operatorname{Add}_{-\gamma z^{-1}}^{(i_0)} \circ \operatorname{Add}_{\gamma z^{-1}}^{(0)} \circ \Phi_{\xi}^{-1}$ with suitable ξ and ξ' . Thus it follows that the map preserves the \mathcal{L} -irreducibility since Add preserves the irreducibility of differential equations.

We can directly check that for $x \in \mu_{\alpha}^{-1}(\lambda)^{\text{dif}}$, its image $x' := \operatorname{add}_{\gamma}^{(i_0)}(x) \in \mu_{\alpha}^{-1}(\lambda + \gamma z^{(i_0)})$ is written as follows. Set

$$x_{\rho_{i_0}} := \left(x_{\rho_{[i_0,j']}^{[0,j]}}\right)_{\substack{1 \le j \le m_0 \\ 1 \le j' \le m_{i_0}}}, \qquad x_{\rho_{i_0}^*} := \left(x_{(\rho_{[i_0,j']}^{[0,j]})^*}\right)_{\substack{1 \le j \le m_0 \\ 1 \le j' \le m_{i_0}}}.$$

Then

$$x'_{(\rho_{[i_0,j']}^{[0,j]})^*} = \left(x_{\rho_{i_0}^*} + \gamma \cdot x_{\rho_{i_0}}^{-1}\right)_{[0,j],[i_0,j']}$$

for $1 \leq j \leq m_0$ and $1 \leq j' \leq m_{i_0}$ and

$$x'_{\rho} = x_{\rho}$$

for the remaining $\rho \in \overline{\mathbb{Q}}_1$, which tells us that the map is analytic. \Box **Theorem 5.13.** If $\mu_{\alpha}^{-1}(\lambda)^{dif} \neq \emptyset$, then $\alpha \in \widetilde{\Sigma}_{\lambda}$.

Proof. Let us suppose that there exists an \mathcal{L} -irreducible representation $x \in \mu_{\alpha}^{-1}(\lambda)^{\text{det}}$. Choose $\gamma_i \in \mathbb{C}$ for $i \in I_{\text{irr}} \setminus \{0\}$ as in Lemma 5.9 and put $\lambda' = \lambda + \sum_{i \in I_{\text{irr}} \setminus \{0\}} \gamma_i z^{(i)}$. Then the operation $\prod_{i \in I_{\text{irr}} \setminus \{0\}} \operatorname{add}_{\gamma_i}^{(i)}$ sends x to the \mathcal{L} -irreducible element $x' \in \mu_{\alpha}^{-1}(\lambda')^{\text{det}}$. However Lemma 5.9 shows that if an element in $\mu_{\alpha}^{-1}(\lambda')$ is \mathcal{L} -irreducible, then it is irreducible. Thus x' is irreducible, which shows $\alpha \in \mathcal{L} \cap \Sigma_{\lambda'}$ by Crawley-Boevey's result (see Theorem 2.6). Hence $\alpha \in \mathcal{L} \cap \Sigma_{\lambda'} \subset \widetilde{\Sigma}_{\lambda'} = \widetilde{\Sigma}_{\lambda}$ by Lemma 5.10.

We close this section by seeing that $\mathfrak{M}_{\lambda}(\mathsf{Q},\alpha)^{\mathrm{dif}}$ is a connected manifold if it is non-empty. Thus the moduli space of connections $\mathfrak{M}(\mathbf{B})$ is also to be connected complex manifold.

Theorem 5.14. If $\mathfrak{M}_{\lambda}(\mathbb{Q}, \alpha)^{dif}$ is non-empty, then it is a connected complex manifold. Thus $\mathfrak{M}(\mathbf{B})$ is a connected complex manifold if it is non-empty, under the identification by the isomorphism Φ_{ξ} given in Theorem 4.22.

Proof. Let us take $\gamma_i \in \mathbb{C}$ for $i \in I_{irr} \setminus \{0\}$ as in Lemma 5.9 and set $\lambda' := \lambda + \sum_{i \in I_{irr}\{0\}} \gamma_i z^{(i)}$. Then it suffices to consider $\mathfrak{M}_{\lambda'}(\mathbb{Q}, \alpha)^{dif}$ by Lemma 5.12. Note that Lemma 5.9 shows that the \mathcal{L} -irreducibility coincides with the irreducibility in $\mu_{\alpha}^{-1}(\lambda')$. Thus $\mathfrak{M}_{\lambda'}(\mathbb{Q}, \alpha)^{dif}$ becomes an open subset of the complex manifold $\mathfrak{M}_{\lambda'}^{\operatorname{reg}}(\mathsf{Q},\alpha)$.

To show the connectedness, let us recall that $\mu_{\alpha}^{-1}(\lambda')$ is irreducible topological space from Theorem 1.2 in [8] and $\mu_{\alpha}^{-1}(\lambda')^{\text{irr}}$ is open in $\mu_{\alpha}^{-1}(\lambda')$. Thus

$$\mu_{\alpha}^{-1}(\lambda')^{\text{dif}} = \left\{ x \in \mu_{\alpha} - 1(\lambda')^{\text{irr}} \left| \det \left(x_{\rho_{[i,j']}^{[0,j]}} \right)_{\substack{1 \le j \le m_0 \\ 1 \le j' \le m_i}} \neq 0, \ i \in I_{\text{irr}} \setminus \{0\} \right\} \right.$$
connected since it is open in the irreducible space $\mu_{\alpha}^{-1}(\lambda')$.

is connected since it is open in the irreducible space $\mu_{\alpha}^{-1}(\lambda')$.

Theorem 5.15. If $\mathfrak{M}(\mathbf{B}) \neq \emptyset$, there exist $\lambda' \in \mathbb{C}^{\mathsf{Q}_0}$ and the injection

$$\Phi\colon\mathfrak{M}(\mathbf{B})\hookrightarrow\mathfrak{M}^{reg}_{\lambda'}(\mathsf{Q},\alpha)$$

whose image is $\left(\mu_{\alpha}^{-1}(\lambda')^{det} \cap \mu_{\alpha}^{-1}(\lambda')^{irr}\right) / \mathbf{G}(\alpha).$

Proof. Let us choose λ' as in the proof of Theorem 5.14. Then $\mathfrak{M}(\mathbf{B}) \cong \mathfrak{M}_{\lambda}(\mathbf{Q},\alpha)^{\text{dif}} \cong \mathfrak{M}_{\lambda'}(\mathbf{Q},\alpha)^{\text{dif}} = \left(\mu_{\alpha}^{-1}(\lambda')^{\text{det}} \cap \mu_{\alpha}^{-1}(\lambda')^{\text{irr}}\right) / \mathbf{G}(\alpha) \subset \mathfrak{M}_{\lambda'}^{\text{reg}}(\mathbf{Q},\alpha).$

6. \mathcal{L} -fundamental set

This section and the next one are dedicated to show the converse of Theorem 5.13. Namely we shall show that if $\alpha \in \widetilde{\Sigma}_{\lambda}$, then $\mu_{\alpha}^{-1}(\lambda)^{\text{dif}} \neq \emptyset$. For this purpose, first let us introduce an analogue of the fundamental set F.

Definition 6.1 (\mathcal{L} -fundamental set). Let us define the subset of \mathcal{L} by

$$\tilde{F} := \left\{ \beta \in \mathcal{L}^+ \setminus \{0\} \left| \begin{array}{c} (\beta, \epsilon_a) \le 0 \text{ for all } a \in \mathcal{J} \cup \mathsf{Q}_0^{\text{leg}}, \\ \text{support of } \beta \text{ is connected} \end{array} \right\} \right\}$$

and call \mathcal{L} -fundamental set.

The aim of this section is to show that \tilde{F} consists of positive imaginary roots of \mathbb{Q} and $\mu_{\alpha}^{-1}(\lambda)^{\text{dif}} \neq \emptyset$ if $\alpha \in \widetilde{\Sigma}_{\lambda} \cap \tilde{F}$.

6.1. A symmetric Kac-Moody root lattice associated with \mathcal{L} . As we saw in the previous section, the sublattice $\mathcal{L} \subset \mathbb{Z}^{Q_0}$ has the action of $\langle s_a \mid a \in \mathcal{J} \cup Q_0^{\text{leg}} \rangle$ which is a subgroup of the Weyl group of Q. We would like to define an analogy of root system on \mathcal{L} and regard \tilde{F} as the fundamental set of positive imaginary roots of \mathcal{L} . For this purpose, we shall define a Kac-Moody root lattice \mathcal{M} and regard \mathcal{L} as a quotient lattice of \mathcal{M} . And then \tilde{F} will be the image of the fundamental set of positive imaginary roots of \mathcal{M} .

6.1.1. Lift of \mathcal{L} to a Kac-Moody root lattice. Let us note that \mathcal{L} is generated by

$$\left\{\epsilon_a \mid a \in \mathcal{J} \cup \mathsf{Q}_0^{\mathrm{leg}}\right\}$$

see Theorem 3.6 in [14]. It is easy to verify that

(6)
$$(\epsilon_{\mathbf{i}}, \epsilon_{\mathbf{i}'}) = 2 - \sum_{\substack{0 \le i \le p \\ j_i \ne j'_i}} (d_i(j_i, j'_i) + 2),$$

(7)
$$(\epsilon_{\mathbf{i}}, \epsilon_{[i,j,k]}) = \begin{cases} -1 & \text{if } j = j_i \text{ and } k = 1, \\ 0 & \text{otherwise,} \end{cases}$$

(8)
$$(\epsilon_{[i,j,k]}, \epsilon_{[i',j',k']}) = \begin{cases} 2 & \text{if } [i,j,k] = [i',j',k'], \\ -1 & \text{if } (i,j) = (i',j') \text{ and } |k-k'| = 1, \\ 0 & \text{otherwise}, \end{cases}$$

cf. section 3.2 in [14]. Here $\mathbf{i} = ([i, j_i])_{0 \le i \le p}$, $\mathbf{i}' = ([i, j'_i])_{0 \le i \le p} \in \mathcal{J}$. Thus we consider a new lattice \mathcal{M} generated by the set of indeterminate

$$\mathcal{C} := \left\{ c_a \mid a \in \mathcal{J} \cup \mathsf{Q}_0^{\mathrm{leg}} \right\},\,$$

namely all $c_a \in C$ have no relations, and define a symmetric bilinear form (,) on \mathcal{M} in accordance with equations (6), (7) and (8).

We can attach \mathcal{M} to a diagram, called *Dynkin diagram*, regarding elements in \mathcal{C} as vertices and connecting $c, c' \in \mathcal{C}$ by |(c, c')| edges if $c \neq c'$. We say $c, c' \in \mathcal{C}$ are *connected* if there exists a sequence $c_0 = c, c_1, \ldots, c_r = c'$ in \mathcal{C} such that $(c_{i-1}, c_i) \neq 0$ for all $i = 1, \ldots, r$. Then we may define *Dynkin diagram* of $\gamma \in \mathcal{M}$ which is a subdiagram obtained by connecting the vertices in $\operatorname{supp}(\beta)$ in the same manner.

Also we can define reflections s_a on \mathcal{M} by

$$s_a(\gamma) := \gamma - (\gamma, c_a)c_a$$

for $a \in \mathcal{J} \cup \mathsf{Q}_0^{\text{leg}}$ and $\gamma \in \mathcal{M}$. Let us denote the set of all positive elements in \mathcal{M} by \mathcal{M}^+ .

Then the inclusion $\mathcal{L} \hookrightarrow \mathbb{Z}^{\mathsf{Q}_0}$ induces

$$\Xi\colon \mathcal{M}\longrightarrow \mathbb{Z}^{\mathsf{Q}_0}$$

where for $\gamma = \sum_{c \in \mathcal{C}} \gamma_c c \in \mathcal{M}$, the image $\Xi(\gamma) = (\beta_a)_{a \in \mathbf{Q}_0}$ is given by

$$\beta_{[i,j]} := \sum_{\{\mathbf{i} = ([i,j_i]) \in \mathcal{J} | j_i = j\}} \gamma_{c_{\mathbf{i}}}$$
$$\beta_{[i,j,k]} := \gamma_{c_{[i,j,k]}}.$$

Proposition 6.2 (Theorem 3.6 in [14]). We have the following.

- (1) We have $(\gamma, \gamma') = (\Xi(\gamma), \Xi(\gamma'))$ for any $\gamma, \gamma' \in \mathcal{M}$.
- (2) The image of Ξ is \mathcal{L} .
- (3) The map Ξ is injective if and only if

$$\#\{i \in \{0, \dots, p\} \mid m_i > 1, i = 0, \dots, p\} \le 1.$$

(4) For $\gamma \in \mathcal{M}$ and $a \in \mathcal{J} \cup \mathsf{Q}_0^{leg}$, we have

$$\Xi(s_a(\gamma)) = s_a(\Xi(\gamma))$$

From this proposition \mathcal{M} can be seen as a "lift" of \mathcal{L} to a Kac-Moody root lattice in which s_i for $i \in \mathcal{J}$ are simple reflections.

6.1.2. Some special examples. The Dynkin diagram of \mathbb{Z}^{Q_0} can be defined as well as that of \mathcal{M} and the Dynkin diagram of $\beta \in \mathbb{Z}^{Q_0}$ is also defined. Here we note that these diagrams coincide with diagrams obtained by forgetting the orientation of the quiver Q and the subquiver associated with $\operatorname{supp}(\beta)$ respectively. Let us compare Dynkin diagrams of $\beta \in \mathcal{L}$ and the inverse image $\Xi^{-1}(\beta) \in \mathcal{M}$ in the following special cases for the latter use.

The kernel of Ξ is a big space in general. Thus if we consider the inverse image of an element $\beta \in \mathcal{L}$, it is convenient to restrict Ξ to some smaller space as follows. Fix $\beta \in \mathcal{L}$. Define $\mathcal{J}_{\beta} := \{([i, j_i])_{i=0,...,p} \in \mathcal{J} \mid \beta_{[i,j_i]} \neq 0 \text{ for all } i \in I_{\text{irr}}\}, \ \mathsf{Q}_0^{\text{leg}}(\beta) := \text{supp}(\beta) \cap \mathsf{Q}_0^{\text{leg}}, \text{ and a sublattice } \mathcal{M}_{\beta} := \sum_{\{a \in \mathcal{J}_{\beta} \cup \mathsf{Q}_0^{\text{leg}}(\beta)\}} \mathbb{Z}c_a$. Denote the set of all positive elements in \mathcal{M}_{β} by \mathcal{M}_{β}^+ . We write the restriction of Ξ on \mathcal{M}_{β} by Ξ_{β} .

First consider $\beta \in \mathcal{L}$ satisfying that

$$\{[i, j] \mid \beta_{[i, j]} \neq 0\} = \{[i, 1]\} \text{ for all } i \in I_{\text{irr}} \setminus \{0\},$$
$$\{[0, j] \mid \beta_{[0, j]} \neq 0\} = \{[0, 1], \dots, [0, m'_0]\}$$

for some $m'_0 \leq m_0$, and $\operatorname{supp}(\beta)$ is connected. Then Proposition 6.2 implies that $\Xi_{\beta} \colon \mathcal{M}_{\beta} \to \mathcal{L}$ is injective and we can show that this bijection preserves Dynkin diagrams of elements in \mathcal{M} and \mathcal{L} in the following way.

Proposition 6.3. For the above $\beta \in \mathcal{L}$, let us define

$$\beta^{red} := \left(\prod_{i \in I_{irr} \setminus \{0\}} s_{[i,1,e_{[i,1]}-1]} \circ \dots \circ s_{[i,1,2]} \circ s_{[i,1,1]} \circ s_{[i,1]}\right) (\beta).$$

Then Dynkin diagram of β^{red} coincides with that of $\Xi_{\beta}^{-1}(\beta) \in \mathcal{M}_{\beta}$.

Proof. From Proposition 6.2, the map Ξ_{β} is injective. And note the identification

$$\mathcal{J}_{\beta} = \{1, \dots, m'_0\} \times (\prod_{i=1}^p \{1\}) \longrightarrow \{1, \dots, m'_0\}$$
$$(j_0, 1_1, \dots, 1_p) \longmapsto j_0$$

Thus for $\beta \in \mathcal{L}$ of the form

$$\beta = \sum_{j=1}^{m'_0} \left(\beta_{[0,j]} \epsilon_{[0,j]} + \sum_{k=1}^{e_{[0,j]}-1} \beta_{[0,j,k]} \epsilon_{[0,j,k]} \right) + \sum_{i \in I_{irr} \setminus \{0\}} \left(\beta_{[i,1]} \epsilon_{[i,1]} + \sum_{k=1}^{e_{[i,1]}-1} \beta_{[i,1,k]} \epsilon_{[i,1,k]} \right) + \sum_{i \in I_{reg}} \sum_{k=1}^{e_{[i,1]}-1} \beta_{[i,1,k]} \epsilon_{[i,1,k]},$$

the inverse image $\tilde{\gamma} := \Xi_{\beta}^{-1}(\beta)$ is written by

$$\begin{split} \tilde{\beta} &= \sum_{j \in \mathcal{J}_{\beta} = \{1, \dots, m'_{0}\}} \left(\beta_{[0,j]} c_{j} + \sum_{k=1}^{e_{[0,j]}-1} \beta_{[0,j,k]} c_{[0,j,k]} \right) \\ &+ \sum_{i \in I_{irr} \setminus \{0\}} \sum_{k=1}^{e_{[i,1]}-1} \beta_{[i,1,k]} c_{[i,1,k]} \\ &+ \sum_{i \in I_{reg}} \sum_{k=1}^{e_{[i,1]}-1} \beta_{[i,1,k]} c_{[i,1,k]}. \end{split}$$

Noting that $\beta_{[i,1]} = \sum_{j=1}^{m_0} \beta_{[0,j]}$ for $i = 1, \ldots, p$ since $\beta \in \mathcal{L}$, we can obtain that

$$\beta^{\text{red}} = \sum_{j=1}^{m'_0} \left(\beta_{[0,j]} \epsilon_{[0,j]} + \sum_{k=1}^{e_{[0,j]}-1} \beta_{[0,j,k]} \epsilon_{[0,j,k]} \right) \\ + \sum_{i \in I_{\text{irr}} \setminus \{0\}} \left(\beta_{[i,1,1]} \epsilon_{[i,1]} + \sum_{k=1}^{e_{[i,1]}-2} \beta_{[i,1,k+1]} \epsilon_{[i,1,k]} \right) \\ + \sum_{i \in I_{\text{reg}}} \sum_{k=1}^{e_{[i,1]}-1} \beta_{[i,1,k]} \epsilon_{[i,1,k]},$$

from direct computation. Let us recall that $(\epsilon_{[i,j]}, \epsilon_{[i,j,1]}) = 1$ and

$$(\epsilon_{[i,j,k-1]}, \epsilon_{[i,j,k]}) = (c_{[i,j,k-1]}, c_{[i,j,k]}) = 1$$

for $i = 0, ..., p, j = 1, ..., m_i$ and $k = 2, ..., e_{[i,1]} - 1$. Also

for $j, j' = 1,, m_0$,
for $j = 1, \ldots, m_0, i \in I_{irr} \setminus \{0\},\$
for $j = 1,, m_0, j \in I_{reg}$,
for $j = 1,, m_0$.

All the other pairs in $\operatorname{supp}(\beta)$ (resp. $\operatorname{supp}(\tilde{\beta})$) are zero. Thus we are done.

Next let us consider the following special $\beta \in \mathcal{L}$, $\{[0, j] \mid \beta_{[0, j]} \neq 0\} = \{[0, 1], [0, 2]\}, \quad \{[1, j] \mid \beta_{[1, j]} \neq 0\} = \{[1, 1], [1, 2]\},$ $\{[i, j] \mid \beta_{[i, j]} \neq 0\} = \{[i, 1]\} \text{ for all } i \in I_{irr} \setminus \{0, 1\},$ $d_0(1, 2) = d_1(1, 2) = 1,$

and

$$\mathsf{Q}_0^{\mathrm{leg}} \cap \mathrm{supp}(\beta) = \emptyset.$$

Then the Dynkin diagram of \mathcal{M}_{β} is

 $c_{\mathbf{i}_2}$ $c_{\mathbf{i}_4}$ $c_{\mathbf{i}_1}$ where $\mathbf{i}_1, \dots, \mathbf{i}_4 \in \mathcal{J} =$

 $\{1_0, 2_0\} \times \{1_1, 2_1\} \times \{1_2\} \times \dots \times \{1_p\}$ are defined by

$$\mathbf{i}_1 := (1_0, 1_1, 1_2, \dots, 1_p), \qquad \mathbf{i}_2 := (2_0, 1_1, 1_2, \dots, 1_p), \\ \mathbf{i}_3 := (1_0, 2_1, 1_2, \dots, 1_p), \qquad \mathbf{i}_4 := (2_0, 2_1, 1_2, \dots, 1_p).$$

In this case corresponding subquiver Q_{β} generated by supp (β) is



Then the image of $\tilde{\beta} = \beta_1 c_{\mathbf{i}_1} + \dots + \beta_4 c_{\mathbf{i}_4} \in \mathcal{M}_\beta$ by $\Xi_\beta \colon \mathcal{M}_\beta \to \mathcal{L}$ is $\beta := \Xi(\tilde{\beta}) = (\beta_1 + \beta_3)\epsilon_{[0,1]} + (\beta_2 + \beta_4)\epsilon_{[0,2]} + (\beta_1 + \beta_2)\epsilon_{[1,1]} + (\beta_3 + \beta_4)\epsilon_{[1,2]} + (\beta_1 + \dots + \beta_4)\epsilon_{[r,1]} + (\beta_1 + \dots + \beta_4)\epsilon_{[r,1]}.$

Let us define

$$\beta^{\text{red}} := \left(\prod_{i=2}^{r} s_{[i,1]}\right) (\beta).$$

Then the Dynkin diagram of β^{red} is of type $A_3^{(1)}$,



if $\beta_s + \beta_t \neq 0$ for all (s, t) = (1, 2), (1, 3), (2, 4), (3, 4).

Remark 6.4. Recalling the orientation of Q, we can check that reflections defining β^{reg} , $\prod_{i \in I_{\text{irr}} \setminus \{0\}} s_{[i,1,e_{[i,1]}-1]} \circ \cdots \circ s_{[i,1,2]} \circ s_{[i,1,1]} \circ s_{[i,1]}$ in Proposition 6.3 and $\prod_{i=2}^{r} s_{[i,1]}$ in the latter example, are products of *admissible reflections*, i.e., reflection functors of Bernstein-Gel'fand-Ponomarev on representations of Q associated to these reflections are well-defined. The detail of the reflection functor can be found in [18] for example.

6.1.3. Inverse image of a positive imaginary root is a positive imaginary root. The following lemma shows that if $\beta \in \mathcal{L}^+$, then $\Xi_{\beta}^{-1}(\beta) \cap \mathcal{M}_{\beta}^+ \neq \emptyset$. Namely there exist at least one positive element in the inverse image of a positive element in \mathcal{L} .

Lemma 6.5 (Lemma 3 in [15]). Take $\beta \in \mathcal{L}^+ \setminus \{0\}$ and set

$$\overline{m}_i := \max\{j \in \{1, \dots, m_i\} \mid \beta_{[i,j]} \neq 0\},\\ \underline{m}_i := \min\{j \in \{1, \dots, m_i\} \mid \beta_{[i,j]} \neq 0\},$$

for $i \in I_{irr}$. Further set $\overline{\mathbf{i}} := ([i, \overline{m}_i])_{0 \le i \le p}$ and $\underline{\mathbf{i}} := ([i, \underline{m}_i])_{0 \le i \le p}$ where we put $\overline{m}_i = \underline{m}_i := 1$ for $i \in I_{reg}$.

Then there exists $\tilde{\beta} \in \mathcal{M}^+_{\beta}$ such that $\Xi(\tilde{\beta}) = \beta$ and $\tilde{\beta}_{c_{\overline{i}}} \cdot \tilde{\beta}_{c_{\overline{i}}} \neq 0$.

From this lemma and the above proposition, we can see the following.

Corollary 6.6. For $\beta \in \tilde{F}$ we have $q(\beta) \leq 0$.

Proof. Let us take $\tilde{\beta} \in \mathcal{M}^+$ as Lemma 6.5. Then Proposition 6.2 says that $(\tilde{\beta}, c_a) \leq 0$ for all $a \in \mathcal{J} \cup \mathsf{Q}_0^{\text{leg}}$. Thus $q(\beta) = q(\tilde{\beta}) \leq 0$, see the proof of Proposition 5.2 in [23] for example.

For $\beta \in \mathcal{L}$, let us define

$$\operatorname{Irr}(\beta) = \left\{ i \in I_{\operatorname{irr}} \middle| \begin{array}{c} \text{there exist at least two distinct } j, j' \text{ such that} \\ \beta_{[i,j]} \neq 0 \text{ and } \beta_{[i,j']} \neq 0 \end{array} \right\}$$

Then we can identify

$$\mathcal{J}_{\beta} \cong \prod_{i \in \operatorname{Irr}(\beta)} \{ [i, j] \mid \beta_{[i, j]} \neq 0 \}.$$

Proposition 6.2 shows that if $\#\operatorname{Irr}(\beta) \leq 1$ for $\beta \in \mathcal{L}^+$, then Ξ_{β} is injective, which implies that $\Xi_{\beta}^{-1}(\beta) = \{\tilde{\beta}\}$. Here we take $\tilde{\beta}$ as in Lemma 6.5.

Next we shall see that for $\beta \in \tilde{F}$, the support of $\tilde{\beta} \in \mathcal{M}^+_{\beta}$ in Lemma 6.5 is connected with one exception.

Lemma 6.7. For $\mathbf{i} = ([i, j_i])_{0 \le i \le p}$, $\mathbf{i}' = ([i, j'_i])_{0 \le i \le p} \in \mathcal{J}$ and $\mathbf{i} \ne \mathbf{i}'$, we have $(c_i, c_{i'}) = 0$ if and only if the following are satisfied;

- (1) $\#\{i \mid j_i \neq j'_i\} = 1,$ (2) setting $\{i_0\} := \{i \mid j_i \neq j'_i\}$, we have $d_{i_0}(j_{i_0}, j'_{i_0}) = 0.$

Proof. Since $d_i(j, j') \ge 0$ for $j \ne j'$, $(c_{\mathbf{i}}, c_{\mathbf{i}'}) = 2 - \sum_{\substack{0 \le i \le p \\ j_i \ne j'_i}} (d_i(j_i, j'_i) + 2) < 0$ if $\#\{i \mid j_i \ne j'_i\} \ge 2$. Also if $\{i_0\} = \{i \mid j_i \ne j'_i\}$, then $(c_{\mathbf{i}}, c_{\mathbf{i}'}) = 2 - (d_i(j_i, j'_i) + 2) < 0$. 2). Thus we obtain the result.

Lemma 6.8. For $\beta \in \tilde{F}$, take $\tilde{\beta} \in \mathcal{M}^+$ as in Lemma 6.5.

- (1) If $c_{[i,j,k]} \in \operatorname{supp}(\tilde{\beta})$, then $c_{[i,j,k']} \in \operatorname{supp}(\tilde{\beta})$ for $k' \leq k$ and there exists $\mathbf{i} = ([i', j_{i'}])_{i'=0,\dots,p} \text{ satisfying } j_i = j \text{ such that } c_{\mathbf{i}} \in \operatorname{supp}(\tilde{\beta}).$
- (2) If one of the following is satisfied,
 - (a) #Irr $(\beta) \ge 3$,
 - (b) there exist $[i, j], [i, j'] \in \operatorname{supp}(\beta)$ such that $d_i(j, j') \ge 1$,

then any distinct $c_{\mathbf{i}}, c_{\mathbf{i}'} \in \operatorname{supp}(\tilde{\beta})$ for $\mathbf{i}, \mathbf{i}' \in \mathcal{J}_{\beta}$ are connected.

Proof. Let us note that $\sum_{\{\mathbf{i}\in\mathcal{J}\mid j_i=j\}}\tilde{\beta}_{c_{\mathbf{i}}}\geq\tilde{\beta}_{c_{[i,j,1]}}\geq\tilde{\beta}_{c_{[i,j,2]}}\geq\tilde{\beta}_{c_{[i,j,e_{[i,j]}-1]}}$ since $\beta \in F$. This shows (1).

Let us suppose that $\#\operatorname{Irr}(\beta) \geq 3$. If $(c_i, c_{i'}) \neq 0$ for a pair of distinct $c_{\mathbf{i}}, c_{\mathbf{i}'} \in \operatorname{supp}(\beta)$, then they are obviously connected. Thus we consider the case $(c_i, c_{i'}) = 0$. Then $\#\{i \mid j_i \neq j'_i\} = 1$ by Lemma 6.7. The hypothesis $\#\operatorname{Irr}(\beta) \geq 3$ assures that the existence of $c_{\mathbf{i}''} \in \operatorname{supp}(\beta)$ such that $\#\{i \mid j_i \neq j_i\}$ $j''_i \neq 1$ and $\#\{i \mid j'_i \neq j''_i\} \neq 1$ are satisfied. Namely $(c_i, c_{i''}) \leq -1$ and $(c_{\mathbf{i}'}, c_{\mathbf{i}''}) \leq -1$. Thus $c_{\mathbf{i}}$ and $c_{\mathbf{i}'}$ are connected.

Next suppose that there exist $[\underline{i}, \underline{j}], [\underline{i}, \underline{j'}] \in \operatorname{supp}(\beta)$ such that $d_{\underline{i}}(\underline{j}, \underline{j'}) \ge 1$.

We note that for any $i \in \{0, ..., p\}$ and $j, j', j'' \in \{1, ..., m_i\}$, the triangle inequality

$$d_i(j, j') \le \max \{ d_i(j, j''), d_i(j'', j') \}$$

holds (see equations (26) in [15]). Thus $d_{\underline{i}}(j,\underline{j}) \ge 1$ or $d_{\underline{i}}(j,\underline{j}') \ge 1$ holds for $j \in \{1, \ldots, m_{\underline{i}}\} \setminus \{\underline{j}, \underline{j}'\}$ since the condition $d_{\underline{i}}(j,\underline{j}) = d_{\underline{i}}(j,\underline{j}') = 0$ breaks the triangle inequality and $d_{\underline{i}}(\underline{j},\underline{j}') \ge 1$.

Let us fix $\mathbf{i} = ([i, j_i])_{0 \le i \le p} \in \mathcal{J}$ such that $c_{\mathbf{i}} \in \operatorname{supp}(\tilde{\beta})$ and define $\mathcal{J}_{\mathbf{i},\underline{i}} := \{([i, j'_i])_{0 \le i \le p} \in \mathcal{J} \mid j'_i = j_i \text{ for all } i \ne \underline{i}\}$. Then for all $\mathbf{i}', \mathbf{i}'' \in \mathcal{J}_{\mathbf{i},\underline{i}}, c_{\mathbf{i}'}$ and $c_{\mathbf{i}''}$ are connected. Indeed define $\mathbf{i}_{\underline{j}} = ([i, j^{(1)}_i])_{0 \le i \le p} \in \mathcal{J}_{\mathbf{i},\underline{i}}$ and $\mathbf{i}_{\underline{j}'} = ([i, j^{(2)}_i])_{0 \le i \le p} \in \mathcal{J}_{\mathbf{i},\underline{i}}$ by

$$j_i^{(1)} = \begin{cases} \underline{j} & \text{if } i = \underline{i}, \\ \overline{j_i} & \text{otherwise} \end{cases}$$

and

$$j_i^{(2)} = \begin{cases} \underline{j'} & \text{if } i = \underline{i}, \\ \overline{j_i} & \text{otherwise} \end{cases}$$

respectively. Then $c_{\mathbf{i}'}$ and $c_{\mathbf{i}''}$ are connected with $c_{\mathbf{i}_{\underline{j}}}$ or $c_{\mathbf{i}_{\underline{j}'}}$ since $d_{\underline{i}}(j,\underline{j}) \geq 1$ or $d_{\underline{i}}(j,\underline{j}') \geq 1$ holds for $j \in \{1, \ldots, m_{\underline{i}}\} \setminus \{\underline{j}, \underline{j}'\}$ as we saw above. Moreover $c_{\mathbf{i}_{\underline{j}}}$ and $c_{\mathbf{i}_{j'}}$ are connected. Thus $c_{\mathbf{i}'}$ and $c_{\mathbf{i}''}$ are connected.

Moreover $c_{\mathbf{i}_{\underline{j}}}$ and $c_{\mathbf{i}_{\underline{j}'}}$ are connected. Thus $c_{\mathbf{i}'}$ and $c_{\mathbf{i}''}$ are connected. Finally consider $c_{\mathbf{i}'''} \in \operatorname{supp}(\tilde{\beta})$ with $\mathbf{i}''' = ([i, j_i'''])_{0 \le i \le p} \notin \mathcal{J}_{\mathbf{i},\underline{i}}$, namely there exists $0 \le i_0 (\ne \underline{i}) \le p$ such that $j_{i_0}''' \ne j_i$. Then we can choose $\mathbf{i}''' = (i, j_i''')_{0 \le i \le p} \in \mathcal{J}_{\mathbf{i},\underline{i}}$ so that $j_{\underline{i}'}'' \ne j_{\underline{i}''}''$ since $\#\{1, \ldots, m_{\underline{i}}\} \ge 2$. Thus

$$(c_{\mathbf{i}'''}, c_{\mathbf{i}''''}) \le 2 - (d_{i_0}(j_{i_0}''', j_{i_0}) + 2) + (d_{\underline{i}}(j_{\underline{i}''}'', j_{\underline{i}}) + 2) < 0.$$

Summing up these results, we can conclude that any distinct $c_{\mathbf{i}^{(1)}}, c_{\mathbf{i}^{(2)}} \in \operatorname{supp}(\tilde{\beta})$ is connected with some $c_{\mathbf{i}^{(3)}}, c_{\mathbf{i}^{(4)}} \in \operatorname{supp}(\tilde{\beta})$, $\mathbf{i}^{(3)}, \mathbf{i}^{(4)} \in \mathcal{J}_{\mathbf{i},\underline{i}}$ respectively. Since $c_{\mathbf{i}^{(3)}}$ and $c_{\mathbf{i}^{(4)}}$ are connected, $c_{\mathbf{i}^{(1)}}$ and $c_{\mathbf{i}^{(2)}}$ are connected as well.

Proposition 6.9. For $\beta \in \tilde{F}$ take $\tilde{\beta} \in \mathcal{M}^+$ as in Lemma 6.5. Then the support of $\tilde{\beta}$ is connected or the following holds. There exist $i_1, i_2 \in I_{irr}$ and $j_{s,t} \in \{1, \ldots, m_{i_s}\}$ for s, t = 1, 2 such that $Irr(\beta) = \{i_1, i_2\}$ and $supp(\tilde{\beta}) = \{c_{\mathbf{i}_{u,v}} \mid u, v = 1, 2\}$ where

$$\mathbf{i}_{u,v} := ([i_1, j_{1,u}], [i_2, j_{2,v}]) \in \mathcal{J}_{\beta} \cong \prod_{s=1}^2 \{ [i_s, j_{s,t}] \mid t = 1, 2 \}$$

and

$$(c_{\mathbf{i}_{u,v}}, c_{\mathbf{i}_{u'v'}}) = \begin{cases} 0 & \text{if } u = u' \text{ or } v = v', \\ -2 & \text{otherwise}, \end{cases}$$

namely the Dynkin diagram of $\operatorname{supp}(\tilde{\beta})$ is case we have $q(\beta) = 0$. $c_{\mathbf{i}_{1,1}} \\ c_{\mathbf{i}_{1,2}} \\ c_{\mathbf{i}_{1,2}}$. In the latter

Proof. From (1) and (2)-(a) in Lemma 6.8, it suffices to consider the case $\#\operatorname{Irr}(\beta) \leq 2$. First suppose that $\#\operatorname{Irr}(\beta) \leq 1$. Then there exists $i_0 \in \{0,\ldots,p\}$ and we may identify $\mathcal{J}_{\beta} \cong \{[i_0,1],\ldots,[i_0,m'_{i_0}]\}$ with some $m'_{i_0} \leq m_{i_0}$. Suppose that $\operatorname{supp}(\tilde{\beta})$ is not connected. Then $c_{[i,j,k]} \notin \operatorname{supp}(\tilde{\beta})$ for any $i \neq i_0, j = 1,\ldots,m_i$ and $k = 1,\ldots,e_{[i,j]} - 1$. Indeed, all $c_{[i,j,k]} \in \operatorname{supp}(\tilde{\beta})$ ($i \neq i_0$) are connected with all c_i ($\mathbf{i} \in \mathcal{J}_{\beta}$) from (1) in Lemma 6.8. And any $c_{[i_0,j,k]} \in \operatorname{supp}(\tilde{\beta})$ is connected with one of c_i for $\mathbf{i} \in \mathcal{J}_{\beta}$. Thus $\operatorname{supp}(\tilde{\beta})$ is connected. Moreover $(c_{\mathbf{i}}, c_{\mathbf{i'}}) = 0$ for all $\mathbf{i} \neq \mathbf{i'} \in \mathcal{J}_{\beta}$ from (2)-(b) in Lemma 6.8. Thus connected components of the Dynkin diagram of $\operatorname{supp}(\tilde{\beta})$ are of type A_n , $\underbrace{\bigcirc}_{c_{\mathbf{i}}} \underbrace{\frown}_{c_{[i_0,j,2]}} \underbrace{c_{[i_0,j,3]}} c_{[i_0,j,3]}$

 $\{[i_0, 1], \ldots, [i_0, m'_{i_0}]\}$. However the root system of type A_n has no imaginary root which contradict to $\beta \in \widetilde{F}$.

The remaining case is $\#Irr(\beta) = 2$. We may suppose that $Irr(\tilde{\beta}) = \{0, 1\}$ without loss of generality.

Recall that for $\overline{\mathbf{i}} = ([0, \overline{m_0}], [1, \overline{m_1}])$ and $\mathbf{i} = ([0, \underline{m_0}], [1, \underline{m_1}]), c_{\overline{\mathbf{i}}}, c_{\overline{\mathbf{i}}} \in \operatorname{supp}(\tilde{\beta})$ and $(c_{\overline{\mathbf{i}}}, c_{\overline{\mathbf{i}}}) \leq 2 - (d_0(\overline{m_0}, \underline{m_0}) + 2) - (d_1(\overline{m_1}, \underline{m_1}) + 2) < 0$. Let us suppose that $\operatorname{supp}(\tilde{\beta})$ is not connected. Then $c_{[i,j,k]} \notin \operatorname{supp}(\tilde{\beta})$ for $i \neq 0, 1$ as above. And there exists $c_{\mathbf{i}} \in \operatorname{supp}(\tilde{\beta})$ such that neither $c_{\overline{\mathbf{i}}}$ nor $c_{\overline{\mathbf{i}}}$ is connected with $c_{\mathbf{i}}$. Here \mathbf{i} must be $\mathbf{i}_1 = ([0, \overline{m_0}], [1, \underline{m_1}])$ or $\mathbf{i}_2 = ([0, \underline{m_0}], [1, \overline{m_1}])$ because of the condition 1 in Lemma 6.7. And $d_i(\overline{m_i}, \underline{m_i}) = 0$ for i = 0, 1. We may suppose $\mathbf{i} = \mathbf{i}_1$. Then $\operatorname{supp}(\tilde{\beta}) \cap \{c_{\mathbf{j}} \mid \mathbf{j} \in \mathcal{J}\} \subset \{c_{\mathbf{i}}, c_{\overline{\mathbf{i}}}, c_{\mathbf{i}_1}, c_{\mathbf{i}_2}\}$. Indeed, if there exists $\mathbf{i}' \in \mathcal{J}$ such that $c_{\mathbf{i}'} \in \operatorname{supp}(\tilde{\beta}) \setminus \{c_{\mathbf{i}}, c_{\overline{\mathbf{i}}}, c_{\mathbf{i}_1}, c_{\mathbf{i}_2}\}$, then the condition 1 in Lemma 6.7 shows that $c_{\mathbf{i}'}$ must be connected with all $c_{\mathbf{i}}, c_{\overline{\mathbf{i}}}, c_{\mathbf{i}_1}, c_{\mathbf{i}_2}$. For



Then $c_{\mathbf{i}}$ is connected with $c_{\underline{\mathbf{i}}}$ and $c_{\overline{\mathbf{i}}}$, which is a

contradiction.

example,

Let us note that $c_{\mathbf{i}_1}$ can be connected only with $c_{\mathbf{i}_2}$ in $\operatorname{supp}(\beta)$ because if there exists $c_{[i,j,1]}$ which is connected with $c_{\mathbf{i}_1}$, then $c_{[i,j,1]}$ must be connected with $c_{\underline{\mathbf{i}}}$ or $c_{\overline{\mathbf{i}}}$ which implies that $c_{\mathbf{i}_1}$ is connected with them. The same argument shows that $c_{\mathbf{i}_2}$ can be connected only with $c_{\mathbf{i}_1}$ in $\operatorname{supp}(\tilde{\beta})$. Thus if $c_{\mathbf{i}_2} \notin \operatorname{supp}(\tilde{\beta})$, then $c_{\mathbf{i}_1}$ is isolated in $\operatorname{supp}(\tilde{\beta})$, which contradicts to that $\beta \in \tilde{F}$.

Hence the remaining possibility is supp $(\tilde{\beta}) = \{c_{\mathbf{i}}, c_{\mathbf{i}}, c_{\mathbf{i}_1}, c_{\mathbf{i}_2}\}$ and

$(c_{\underline{\mathbf{i}}}, c_{\overline{\mathbf{i}}}) = -2,$	$(c_{\underline{\mathbf{i}}},c_{\mathbf{i}_i})=0, i=1,2,$
$(c_{\mathbf{i}_1}, c_{\mathbf{i}_2}) = -2,$	$(c_{\overline{\mathbf{i}}},c_{\mathbf{i}_i})=0, i=1,2.$

6.2. Wild case. Let us separate our argument into the two cases, namely the wild case, $q(\beta) < 0$, and the tame case, $q(\beta) = 0$, for $\beta \in \tilde{F}$. We consider the wild case first.

Proposition 6.10. Let $\beta = \gamma_1 + \cdots + \gamma_r \in \tilde{F}$ with $q(\beta) < 0, r \ge 2$ and $\gamma_1, \ldots, \gamma_r \in \mathcal{L}^+ \setminus \{0\}$ then $q(\beta) < q(\gamma_1) + \cdots + q(\gamma_r)$.

Proof. For the above $\gamma_1, \ldots, \gamma_r$, take $\tilde{\gamma_1}, \ldots, \tilde{\gamma_r} \in \mathcal{M}^+ \setminus \{0\}$ as in Lemma 6.5 and define $\tilde{\beta} = \tilde{\gamma_1} + \cdots + \tilde{\gamma_r}$. Then $\tilde{\beta}$ satisfies conditions in Lemma 6.5. Thus the support of $\tilde{\beta}$ is connected from Proposition 6.9. Recall that $(\tilde{\beta}, c_a) \leq 0$ for all $a \in \mathcal{J} \cup \mathsf{Q}_0^{\text{leg}}$ and the assumption $q(\beta) < 0$. Then the standard argument (see Lemma 2 in [28] for example) shows that $q(\beta) = (\tilde{\beta}, \tilde{\gamma}) < \sum_{i=1}^r (\tilde{\gamma_i}, \tilde{\gamma_i}) = \sum_{i=1}^r q(\gamma_i)$.

Let us fix $\beta \in \widetilde{F}$ with $q(\beta) < 0$. Define a non-empty open subset of $\operatorname{Rep}(\mathsf{Q},\beta)$ by

$$\operatorname{Rep}(\mathsf{Q},\beta)^{\operatorname{det}} :=$$

$$\left\{ x \in \operatorname{Rep}(\mathsf{Q},\beta) \, \middle| \, \det \left(x_{\rho_{[i,j']}^{[0,j]}} \right)_{\substack{1 \leq j \leq m_0 \\ 1 \leq j' \leq m_i}} \neq 0, \, i \in I_{\operatorname{irr}} \backslash \{0\} \right\}.$$

Lemma 6.11. If $x \in \text{Rep}(\mathsf{Q},\beta)^{det}$ is decomposed as $x = x_1 \oplus \cdots \oplus x_r$ in $\text{Rep}(\mathsf{Q},\beta)$, then $\dim x_i \in \mathcal{L}^+$ for all $i = 1, \ldots, r$.

Proof. Since $x \in \operatorname{Rep}(\mathbb{Q}, \beta)^{\operatorname{det}}$, subrepresentations x_t with $\gamma_t = \dim x_t$ for $t = 1, \ldots, r$ satisfy $\sum_{j=1}^{m_0} (\gamma_t)_{[0,j]} \leq \sum_{j=1}^{m_i} (\gamma_t)_{[i,j]}$ for all $i \in I_{\operatorname{irr}} \setminus \{0\}$. If there exists "<" among these inequalities, then $\beta = \gamma_1 + \cdots + \gamma_r \notin \mathcal{L}^+$ which contradicts to the assumption $\beta \in \widetilde{F} \subset \mathcal{L}^+$. Thus $\gamma_t \in \mathcal{L}^+$ for all $t = 1, \ldots, r$.

Let us recall the notion of generic decomposition. A decomposition $\beta = \gamma_1 + \cdots + \gamma_r, \ \gamma_t \in (\mathbb{Z}_{\geq 0})^{\mathbb{Q}_0} \setminus \{0\}$, is called the generic decomposition if

$$\operatorname{Ind}(\mathsf{Q};\gamma_1,\ldots,\gamma_r) = \left\{ x_1 \oplus \cdots \oplus x_r \in \operatorname{Rep}(\mathsf{Q},\beta) \middle| \begin{array}{c} \dim x_t = \gamma_t \text{ and } x_t \text{ are} \\ \text{indecomposable for } t = 1,\ldots,r \end{array} \right\}$$

contains a non-empty open dense subset of $\operatorname{Rep}(\mathbb{Q},\beta)$. It is known that the generic decomposition uniquely exists for any $\beta' \in (\mathbb{Z}_{\geq 0})^{\mathbb{Q}_0} \setminus \{0\}$, see Proposition 2.7 in [28] for example.

Proposition 6.12. Let us take $\beta \in \widetilde{F}$ with $q(\beta) < 0$. If $\beta = \gamma_1 + \cdots + \gamma_r$ is the generic decomposition of $\beta \in \widetilde{F}$, then r = 1.

Proof. If $\beta = \gamma_1 + \cdots + \gamma_r$ is the generic decomposition, then

$$\operatorname{Ind}(\mathsf{Q};\gamma_1,\ldots,\gamma_r)\cap\operatorname{Rep}(\mathsf{Q},\beta)^{\det}\neq\emptyset.$$

Thus $\gamma_i \in \mathcal{L}^+$ for $i = 1, \ldots, r$ by Lemma 6.11. Then Proposition 6.10 shows that $q(\beta) < q(\gamma_1) + \cdots + q(\gamma_r)$ if $r \ge 2$. This contradicts to that $\beta = \gamma_1 + \cdots + \gamma_r$ is the generic decomposition by the standard argument, see Theorem 3.3 in [28] for example. Thus r = 1.

Corollary 6.13. If $\beta \in \widetilde{F}$ and $q(\beta) < 0$, then β is a positive root of Q.

Proof. Proposition 6.12 shows that $\operatorname{Rep}(\mathbb{Q},\beta)$ contains a indecomposable representation. Then Kac's theorem (Theorem 1.10 in [18]) tells us that β is a positive root of \mathbb{Q} .

Corollary 6.14. Let us take $\beta \in \widetilde{\Sigma}_{\nu}$ and suppose that $\beta \in \widetilde{F}$ and $q(\beta) < 0$. Then $\mu_{\beta}^{-1}(\nu)^{dif} \neq \emptyset$.

Proof. Let us take ν' as in Lemma 5.9 and show that $\mu_{\beta}^{-1}(\nu')^{\text{dif}} \neq \emptyset$. Let us note that $\sum_{j=1}^{m_i} \beta_{[i,j]} \geq 1$ for all $i \in I_{\text{irr}}$ since $\operatorname{supp}(\beta)$ is connected. Then Proposition 6.12 shows that the subset Z of $\operatorname{Rep}(\mathbf{Q},\beta)$ consisting of all indecomposable representations is a dense subset. Thus $Z \cap \operatorname{Rep}(\mathbf{Q},\beta)^{\det} \neq \emptyset$. Then Theorem 3.3 in [8] shows that $\mu_{\beta}^{-1}(\nu')^{\det} \neq \emptyset$. Moreover Theorem 1.2 in [8] says that the set of all irreducible representations in the irreducible topological set $\mu_{\beta}^{-1}(\nu')$ is a dense subset. Here we note that $\beta \in \Sigma_{\nu'}$. Thus the non-empty open subset $\nu_{\beta}^{-1}(\nu')^{\det}$ contains a irreducible representation x. Thus $\mu_{\beta}^{-1}(\nu')^{\dim} \neq \emptyset$ which shows that $\mu_{\beta}^{-1}(\nu)^{\dim} \neq \emptyset$ by Lemma 5.12. \Box

6.3. Tame case. Let us consider the remaining case $q(\beta) = 0$ for $\beta \in \tilde{F}$. The proof depends on a classification of Dynkin diagrams of $\operatorname{supp}(\beta)$. To recall the classification, first let us introduce the shape of $\beta \in \mathcal{L}$.

Definition 6.15 (shape). Fix a Kac-Moody root lattice $L = \bigoplus_{i \in I} \mathbb{Z}\alpha_i$ and $\alpha = \sum_{i \in I} m_i \alpha_i \in L$. For the Dynkin diagram of the support of α , we attach each coefficient m_i of α to the vertex corresponding to α_i , then we obtain the diagram with the coefficients, which we call the *shape* of α .

By using this we define shapes of elements in \mathcal{L} as follows.

Definition 6.16. For $\beta \in \mathcal{L}$, the *shape* of β is the set of shapes of elements in $\Xi_{\beta}^{-1}(\beta) \subset \mathcal{M}_{\beta}$.

Example 6.17. For example, suppose p = 1, $m_0 = m_1 = 2$, $e_{[i,j]} = 1$ (i = 0, 1 and j = 1, 2), $d_0(1, 2) = d_1(1, 2) = 0$. Consider $\beta = \epsilon_{[0,1]} + \epsilon_{[0,2]} + \epsilon_{[1,1]} + \epsilon_{[0,2]} + \epsilon_{[1,2]} + \epsilon_{[1,2$

 $\epsilon_{[1,2]}$. Then the shape of β is 1-adenote $\{x_a \mid a \in \mathbb{Z}\}$ by $x_a \ (a \in \mathbb{Z})$. $(a \in \mathbb{Z})$, where we simply

Suppose p = 0, $m_0 = 4$, $d_0(i, j) = 1$ for $1 \le i < j \le 4$ and $e_{0,\nu} = 1$ for $1 \le \nu \le 4$. Then the shape of $\beta = \sum_{\nu=1}^4 \epsilon_{[0,\nu]}$ is $\begin{array}{c} 1 \\ 1 \\ 1 \end{array} \begin{array}{c} 0 \\ 1 \end{array} \begin{array}{c} 0 \\ 1 \end{array}$.

Then a classification of shapes of $\alpha \in \tilde{F}$ is know as follows.

Theorem 6.18 (Theorem 9 in [15]). For $\beta \in \tilde{F}$ with $q(\beta) = 0$, there exists a positive integer m and the shape of β is one of the following.



Here the last shape corresponds to the latter case in Proposition 6.9. For the other shapes, the condition $\#\operatorname{Irr}(\beta) \leq 1$ holds, namely $\Xi^{-1}(\beta) = \{\tilde{\beta}\}.$

By using this classification, we can show that $\mu_{\beta}^{-1}(\nu) \neq \emptyset$ for $\beta \in \widetilde{\Sigma}_{\nu} \cap \tilde{F}$ and $q(\beta) = 0$.

Theorem 6.19. For $\beta \in \widetilde{\Sigma}_{\nu} \cap \widetilde{F}$ and $q(\beta) = 0$, we have $\mu_{\beta}^{-1}(\nu)^{dif} \neq \emptyset$.

Proof. Applying $T_{(i,0)}: \mathfrak{M}_{\nu}(\mathbb{Q},\beta)^{\operatorname{dif}} \to \mathfrak{M}_{t_{(i,0)}(\nu)}(t_{(i,0)}(\mathbb{Q}), t_{(i,0)}(\beta))^{\operatorname{dif}}$ defined in Section 5.2 if necessary, we can assume $0 \in \operatorname{Irr}(\beta)$ if $\operatorname{Irr}(\beta) \neq \emptyset$. Since $\beta \in \widetilde{\Sigma}_{\nu} \cap \widetilde{F}$ and $q(\beta) = 0$, the coefficients of β is indivisible, i.e., have no common divisor other than 1, namely the integer m = 1 in Theorem 6.18 for $\Xi^{-1}(\beta)$. Then we obtain that $\beta^{\operatorname{reg}}$ defined in Section 6.1.2 is the indivisible null root of a Euclidean root system. Indeed, if the shape of β is one of the first 7 cases in Theorem 6.18 which are indivisible null roots of Euclidean root systems. Proposition 6.3 shows that $\beta^{\operatorname{reg}}$ is an indivisible null root. The last case in Theorem 6.18 corresponds to the second example in Section 6.1.2. Thus $\beta^{\operatorname{reg}}$ is the indivisible null root of the root system of type $A_3^{(1)}$.

Since it is known that indivisible null roots of Euclidean root systems are Schur roots of quivers, thus we obtain that β^{reg} is a Schur root. As we noted in Remark 6.4, β^{reg} is obtained by the product of admissible reflections from β . This implies that β is a Schur root as well. Therefore general elements of Rep(Q, β) are indecomposable. Then we have the result by the same argument in Corollary 6.14.

Combining Corollary 6.13 and the proof in Theorem 6.19, we have the following theorem.

Theorem 6.20. The \mathcal{L} -fundamental set is a subset of the set of positive imaginary roots of Q.

Also Corollary 6.14 and Theorem 6.19 show the following theorem.

Theorem 6.21. For $\beta \in \widetilde{\Sigma}_{\nu} \cap \widetilde{F}$, we have $\mu_{\beta}^{-1}(\nu)^{dif} \neq \emptyset$.

7. EXISTENCE OF \mathcal{L} -IRREDUCIBLE REPRESENTATIONS

In this section we shall give a proof of our main theorem. First we recall some basic properties of real roots of Q in \mathcal{L}^+ . The following lemma shows that $\epsilon_{\mathbf{i}}$ behaves as a simple root in $\widetilde{R}^+ := \Delta^+ \cap \mathcal{L}^+$ for each $\mathbf{i} \in \mathcal{J}$.

Lemma 7.1. If we have $s_{\mathbf{i}}(\alpha) \notin \widetilde{R}^+$ for some $\mathbf{i} \in \mathcal{J}$ and $\alpha \in \widetilde{R}^+$, then $\alpha = \epsilon_{\mathbf{i}}$.

Proof. Let us take $\alpha \in \widetilde{R}^+$ and $\mathbf{i} = ([i, j_i]) \in \mathcal{J}$ as above. By Proposition 5.7, $s_{\mathbf{i}}(\alpha) = \left(\prod_{i \in I_{\mathrm{irr}} \setminus \{0\}} s_{[i, j_i]}\right) \circ s_{[0, j_0]} \circ \left(\prod_{i \in I_{\mathrm{irr}} \setminus \{0\}} s_{[i, j_i]}\right) (\alpha)$.

Suppose that $\alpha_1 = \left(\prod_{i \in I_{\mathrm{irr}} \setminus \{0\}} s_{[i,j_i]}\right)(\alpha) \notin \Delta^+$. Then there exists $i_0 \in I_{\mathrm{irr}} \setminus \{0\}$ such that $\alpha_1 = -\epsilon_{[i_0,j_{i_0}]}$ since $s_{[i,j_i]}$ for $i \in I_{\mathrm{irr}} \setminus \{0\}$ are commutative. This implies that $\alpha = \epsilon_{[i_0,j_{i_0}]} \notin \mathcal{L}$ which contradicts to $\alpha \in \widetilde{R}^+$.

Next suppose that $\alpha_2 = s_{[0,j_0]} \circ \left(\prod_{i \in I_{irr} \setminus \{0\}} s_{[i,j_i]}\right)(\alpha) \notin \Delta^+$. Then $\alpha_2 = -\epsilon_{[0,j_0]}$ which shows that $\alpha = \epsilon_i$ and $s_i(\alpha) = -\epsilon_i$.

Finally suppose that $\alpha_2 \in \Delta^+$. Then $\alpha_3 = s_i(\alpha) = -\epsilon_{[i_0, j_{i_0}]}$ for some $i_0 \in I_{irr} \setminus \{0\}$, which shows that $\alpha \notin \widetilde{R}^+$ as above. This is a contradiction. \Box

Lemma 7.2. Take $\beta \in \widetilde{R}^+$ satisfying $\sum_{j=1}^{m_i} \beta_{[i,j]} > 0$ for all $i \in I_{irr}$. Let us suppose that there exists $a \in \mathcal{J} \cup Q_0^{leg}$ such that $\beta' = s_a(\beta)$ satisfies that $\beta' \in \widetilde{R}^+$ and $\beta'_{[i,j]} = 0$ for all $i \in I_{irr}$ and $j = 1, \ldots, m_i$. Then there exist $\mathbf{i} = ([i, j_i])_{0 \le i \le p} \in \mathcal{J}, i_0 \in \{0, \ldots, p\}$ and $l \in \{1, \ldots, e_{[i_0, j_{i_0}]} - 1\}$ such that $a = \mathbf{i}$ and $\beta = \epsilon_{\mathbf{i}} + \epsilon_{[i_0, j_{i_0}, 1]} + \cdots + \epsilon_{[i_0, j_{i_0}, l]}$. In this case β is a real root.

Proof. From the assumption, $\operatorname{supp}(\beta') \subset \mathsf{Q}_0^{\operatorname{leg}}$. Since $\beta' \in \widetilde{R}^+$, there exist $i_0 \in \{0, \ldots, p\}, \ j^{i_0} \in \{1, \ldots, m_{i_0}\}$ and $1 \leq k < l \leq m_{i_0}$ such that $\beta' = \epsilon_{[i_0, j^{i_0}, k]} + \cdots + \epsilon_{[i_0, j^{i_0}, l]}$. From the assumption, there exists $a \in \mathcal{J} \cup \mathsf{Q}_0^{\operatorname{leg}}$ such that $s_a(\beta') = \beta$ and $\sum_{j=1}^{m^{(i)}} \beta_{[i,j]} > 0$. Thus there exists $\mathbf{i} = ([i, j_i]) \in \mathcal{J}$ such that $j_{i_0} = j^{i_0}$ and $a = \mathbf{i}$. Also it follows that k = 1.

Lemma 7.3. Let us take $\beta = \epsilon_i$ or $\beta = \epsilon_i + \epsilon_{i_0, j_{i_0}, 1} + \dots + \epsilon_{i_0, j_{i_0}, l}$ for some $\mathbf{i} = ([i, j_i])_{0 \leq i \leq p} \in \mathcal{J}, i_0 \in \{0, \dots, p\}$ and $l \in \{1, \dots, e_{[i_0, j_{i_0}]} - 1\}$. Then we have $\mu_{\beta}^{-1}(\nu)^{dif} \neq \emptyset$ for any $\nu \in \mathbb{C}^{\mathsf{Q}_0}$ satisfying $\beta \in \widetilde{\Sigma}_{\nu}$.

Proof. Applying the operator $T_{(i_0,0)}$ defined in Section 5.2 if necessary, we may assume $i_0 = 0$. Let us choose $\gamma_i \in \mathbb{C}$ for all $i \in I_{irr} \setminus \{0\}$ such that $\beta \in \Sigma_{\nu'}$ where $\nu' := \nu + \sum_{i \in I_{irr} \setminus \{0\}} \gamma_i z^{(i)}$ as in Proposition 5.11. Note that that $\nu'_{[i,j_i]} \neq 0$ for all $i \in I_{irr} \setminus \{0\}$. Indeed if there exists $i_1 \in I_{irr} \setminus \{0\}$ such that $\nu'_{[i,j_i_1]} = 0$, then we have the decomposition $\beta = \epsilon_{[i_1,j_{i_1}]} + (\beta - \epsilon_{[i_1,j_{i_1}]})$ such that $\nu' \cdot \epsilon_{[i_1,j_{i_1}]} = 0$, $\nu' \cdot (\beta - \epsilon_{[i_1,j_{i_1}]}) = 0$, $\epsilon_{[i_1,j_{i_1}]}, (\beta - \epsilon_{[i_1,j_{i_1}]}) \in \Delta^+$, and $p(\epsilon_{[i_1,j_{i_1}]}) = p(\beta - \epsilon_{[i_1,j_{i_1}]}) = 0$. This contradicts to the assumption $\beta \in \Sigma_{\nu'}$. Thus we obtain that any $x \in \mu_{\beta}^{-1}(\nu)$ satisfies that $\det(x_{\rho_{[i,j]}^{[0,j]}})_{\substack{1 \leq j \leq m_0 \\ 1 \leq j' \leq m_i}} = x_{\rho_{[i_j_i]}^{[0,j_0]}} \neq 0$ for all $i \in I_{irr} \setminus \{0\}$ because $x_{\rho_{[i,j_i]}^{[0,j_0]}} x_{(\rho_{[i,j_i]}^{[0,j_0]})^*} = \nu_{[i,j_i]} \neq 0$. Therefore

 $\mu_{\beta}^{-1}(\nu')^{\text{dif}} = \mu_{\beta}^{-1}(\nu')^{\text{irr}} \neq \emptyset \text{ since } \beta \in \Sigma_{\nu'}. \text{ Applying } \left(\prod_{i \in I_{\text{irr}} \setminus \{0\}} \operatorname{add}_{\gamma_i z}^{(i)}\right)^{-1},$ we obtain $\mu_{\beta}^{-1}(\nu)^{\text{dif}} \neq \emptyset$.

Now let us study the structure of $\tilde{\Sigma}_{\mu}$. Let ~ be the smallest equivalence relation on \mathcal{S} with $(\alpha, \lambda) \sim s_a(\alpha, \lambda)$ for $a \in \mathcal{J} \cup \mathsf{Q}_0^{\mathrm{leg}}$ whenever s_a is defined. We write $\mathbb{N}\widetilde{R}^+_{\lambda}$ for the set of sums of elements of $\widetilde{R}^+_{\lambda}$ including 0. Then the following statements are obtained by the same arguments in [8].

Lemma 7.4 (cf. Lemma 5.1 in [8]). Given any pair (α, λ) with $\alpha \in \mathbb{N}\widetilde{R}^+_{\lambda}$, if $a \in \mathcal{J} \cup \mathsf{Q}_0^{leg}$ with $\lambda_a = 0$ and $(\alpha, \epsilon_a) > 0$, then $\alpha - \epsilon_a \in \mathbb{N}\widetilde{R}^+_{\lambda}$.

Proof. The proof is just an analogy of the proof of Lemma 5.1 in [8]. We can write $\alpha = \sum_{t=1}^{r} \gamma^{(t)}$ as a sum of positive roots. If any $\gamma^{(t)}$ is equal to ϵ_a , then we are done. Otherwise all $s_a(\gamma^{(t)})$ are positive roots by Lemma 7.1, so in $\widetilde{R}^+_{\lambda}$. Thus $s_a(\alpha) = \alpha - (\alpha, \epsilon_a)\epsilon_a \in \mathbb{N}\widetilde{R}^+_{\lambda}$. Then adding on a suitable number of copies of $\epsilon_a \in \widetilde{R}^+_{\lambda}$, it follows that $\alpha - \epsilon_a \in \mathbb{N}\widetilde{R}^+_{\lambda}$.

Lemma 7.5 (cf. Lemma 5.2 in [8]). If $(\alpha, \lambda) \sim (\alpha', \lambda')$ then

- (1) $\alpha \in \widetilde{R}^+_{\lambda}$ if and only if $\alpha' \in \widetilde{R}^+_{\lambda'}$, (2) $\alpha \in \mathbb{N}\widetilde{R}^+_{\lambda}$ if and only if $\alpha' \in \mathbb{N}\widetilde{R}^+_{\lambda'}$,
- (3) $\alpha \in \widetilde{\Sigma}_{\lambda}$ if and only if $\alpha' \in \widetilde{\Sigma}_{\lambda'}$.

Proof. Lemma 7.1 enable us to apply the same argument as in Lemma 5.2 in [8] to this lemma. \square

Lemma 7.6 (cf. Lemma 5.3 in [8]). Given any pair (α, λ) with $\alpha \in \mathbb{N}\widetilde{R}^+_{\lambda}$, there is an equivalent pair (α', λ') with the property that $(\alpha', \epsilon_a) \leq 0$ whenever $\lambda'_a \neq 0 \text{ for } a \in \mathcal{J} \cup \mathsf{Q}_0^{leg}.$

Proof. This follows form Lemma 7.5 as well as the proof of Lemma 5.3 in [8].

The following Lemmas 7.7, 7.8 and Theorem 7.9 can be shown by the arguments in the proofs of the corresponding statements in [8] without any change.

Lemma 7.7 (cf. Lemma 5.4 in [8]). Suppose that $0 \neq \alpha \in \mathbb{N}\widetilde{R}^+_{\lambda}$ and $(\alpha, \epsilon_a) \leq 0$ for all $a \in \mathcal{J} \cup \mathsf{Q}_0^{leg}$ with $\lambda_a \neq 0$. If $(\beta, \alpha - \beta) \leq -2$ whenever $\beta, \alpha - \beta$ are nonzero and in $\mathbb{N}\widetilde{R}^+_{\lambda}$, then α is either ϵ_a where $a \in \mathcal{J} \cup \mathsf{Q}_0^{leg}$ or in the \mathcal{L} -fundamental set.

Lemma 7.8 (cf. Lemma 5.5 in [8]). If $0 \neq \alpha \in \mathbb{N}\widetilde{R}^+_{\lambda}$ and $(\beta, \alpha - \beta) \leq -2$ whenever $\beta, \alpha - \beta$ are nonzero and in $\widetilde{R}^+_{\lambda}$, then $\alpha \in \widetilde{R}^+_{\lambda}$.

Theorem 7.9 (cf. Theorem 5.6 in [8]). If $\alpha \in \mathcal{L}^+$ then $\alpha \in \widetilde{\Sigma}_{\lambda}$ if and only if $0 \neq \alpha \in \mathbb{N}\widetilde{R}^+_{\lambda}$ and $(\beta, \alpha - \beta) \leq -2$ whenever $\beta, \alpha - \beta$ are nonzero and in $\mathbb{N}\widetilde{R}^+_{\lambda}$.

Combining these results, we can obtain the following theorem as in [8].

Theorem 7.10 (cf. Theorem 5.8 in [8]). If $\alpha \in \widetilde{\Sigma}_{\lambda}$ then there is an equivalent pair (α', λ') with α' which is either ϵ_a where $a \in \mathcal{J} \cup \mathsf{Q}_0^{leg}$ or in the \mathcal{L} -fundamental set.

Theorem 7.11. For $\beta \in \widetilde{\Sigma}_{\nu}$, we have $\mu_{\beta}^{-1}(\nu)^{dif} \neq \emptyset$.

Proof. By Theorems 6.21, 7.10 and Lemma 7.2, we may assume $(\beta, \nu) \sim (\epsilon_a, \nu')$, $a \in \mathcal{J} \cup \mathsf{Q}_0^{\text{leg}}$. If $a \in \mathcal{J}$, then Lemma 7.3 shows that $\mu_{\beta}^{-1}(\nu) \neq \emptyset$. Suppose $a \in \mathsf{Q}_0^{\text{leg}}$. If $\sum_{j=1}^{m_i} \beta_{[i,j]} > 0$ for all $i \in I_{\text{irr}}$, then Lemma 7.2 and Theorem 7.10 imply that there exists a sequence $a_1, \ldots, a_r \in \mathcal{J} \cup \mathsf{Q}_0^{\text{leg}}$ such that $(\beta^{(k)}, \nu^{(k)}) = s_{a_k} s_{a_{k-1}} \cdots s_{a_1}(\beta, \nu)$ are well defined and $\sum_{j=1}^{m_i} \beta_{[i,j]}^{(k)} > 0$ $(i \in I_{\text{irr}})$ for all $k = 1, \ldots, r$ and moreover $\beta^{(r)} = \epsilon_i + \epsilon_{[i_0,j_0,1]} + \cdots$ as in Lemma 7.2. Then we have $\mu_{\beta}^{-1}(\nu)^{\text{dif}} \neq \emptyset$ by Lemma 7.3. If $\sum_{j=1}^{m_i} \beta_{[i,j]} = 0$ for all $i \in I_{\text{irr}}$, then $\mu_{\beta}^{-1}(\nu)^{\text{dif}} = \mu_{\beta}^{-1}(\nu)^{\text{irr}} \neq \emptyset$.

Then Theorems 5.13 and 7.11 show our main theorem.

Theorem 7.12. We use the same notation as in Section 5. Let us consider the additive Deligne-Simpson problem for k_0, \ldots, k_p and the HTL normal forms $B_i \in \mathfrak{g}_{k_i}^*$ for $i = 0, \ldots, p$. Then the problem has a solution if and only if $\alpha \in \widetilde{\Sigma}_{\lambda}$.

Corollary 7.13. Let $\mathbf{B} = (B_i)_{0 \le i \le p} \in \prod_{i=0}^{p} \mathfrak{g}_{k_i}$ be the collection of HTL normal forms as above theorem. Then $\mathfrak{M}(\mathbf{B}) \neq \emptyset$ if and only if $\alpha \in \widetilde{\Sigma}_{\lambda}$.

References

- D. Arinkin, Rigid irregular connections on P¹, Compos. Math., 146, no. 5 (2010), 1323–1338.
- [2] W. Balser, Formal power series and linear systems of meromorphic ordinary differential equations. Universitext, Springer-Verlag, New York, 2000.
- [3] P. Boalch, Symplectic manifolds and isomonodromic deformations, Adv. Math. 163 (2) (2001), 137–205.
- [4] P. Boalch, Irregular connections and Kac-Moody root systems, 2008, arXiv:0806.1050.
- [5] P. Boalch, Simply-laced isomonodromy systems, Publ. Math. IHES, 116, no. 1 (2012), 1–68.
- [6] P. Boalch, Geometry and braing of Stokes data; Fission and wild character varieties, Ann. of Math. 179, (2014), 301–365.
- [7] C. Bremer, D. Sage, Moduli spaces of irregular singular connections, Int. Math. Res. Not. IMRN 2013, no.8, 1800–1872.
- [8] W. Crawley-Boevey, Geometry of the moment map for representations of quivers, Compos. Math. 126, no.3 (2001), 257–293.
- [9] W. Crawley-Boevey, On matrices in prescribed conjugacy classes with no common invariant subspace and sum zero, Duke Math. J. 118, no. 2 (2003), 339–352.
- [10] W. Crawley-Boevey and P. Holland, Noncommutative deformations of Kleinian sigularities, Duke Math. J. 92 (1998), 605–635.
- [11] W. Crawley-Boevey and P. Shaw, Multiplicative preprojective algebras, middle covolution and Deligne-Simpson problem, Adv. Math., 201, (2006), 180–208.
- [12] M. Dettweiler, S. Reiter, An algorithm of Katz and its application to the inverse Galois problem, J. Symbolic Comput., 30, no. 6 (2000), 761–798.
- [13] M. Dettweiler, S. Reiter, Middle convolution of Fuchsian systems and the construction of rigid differential systems, J. Algebra, 318, no. 1 (2007), 1–24.
- [14] K. Hiroe, Linear differential equations on P¹ and root systems, J. Algebra, 382, (2013), 1–38.

- [15] K. Hiroe, T. Oshima, A classification of roots of symmetric Kac-Moody root systems and its application, in Symmetries, integrable systems and representations, Springer-Verlag, 2013, 195-241.
- [16] K. Hiroe, D. Yamakawa, Moduli spaces of meromorphic connection and quiver varieties, Adv. Math., 266 (2014), 120–151.
- [17] M. Inaba, M.-H. Saito. Moduli of unramified irregular singular parabolic connections on a smooth projective curve, Kyoto J. Math. 53 (2013), no.2, 433–482.
- [18] V. Kac, Root systems, representations of quivers and invariant theory, Lecture Notes in Mathematics Vol.996, Springer- Verlag, Berlin, 1983, pp. 74–108.
- [19] G. Kempf and L. Ness, The length of vectors in representation spaces in Algebraic Geometry, LNM 732 (1979), 233–244.
- [20] A. King, Moduli of representations of finite dimensional algebras, Quart. J. of Math. 45 (1994), 515–530.
- [21] F. Kirwan, Cohomology of quotients in symplectic and algebraic geometry, Math. Notes 31, Princeton Univ. Press, Princeton, 1984.
- [22] M. Jimbo, T. Miwa, K. Ueno, Monodromy preserving deformations of linear ordinary differential equations with rational coefficients I, Physica D 2, no.2 (1981), 306–352.
- [23] V. Kac, Infinite-dimensional Lie algebras. Third edition. Cambridge University Press, Cambridge, 1990.
- [24] N. Katz, Rigid local systems. Annals of Mathematics Studies, vol. 139. Princeton University Press, 1996.
- [25] H. Kawakami, Generalized Okubo systems and the middle convolution, Int. Math. Res. Not. IMRN, 2010, no. 17, 3394–3421.
- [26] V. Kostov, The Deligne-Simpson problem-a survey, J. Algebra 281, no.1 (2004), 83– 108.
- [27] V. Kostov, Additive Deligne-Simpson problem for non-Fuchsian systems, Funkcialaj Ekvacioj 53 (2010), 395–410.
- [28] H. Kraft, Ch. Riedtmann, Geometry of representations of quivers, in Representations of algebras (Durham, 1985), London Math. Soc. Lecture Note Ser., 116, 109–145, Cambridge Univ. Press, Cambridge, 1986.
- [29] H. Nakajima, Instanton on ALE spaces, quiver varieties, and Kac-Moody algebras, Duke Math. J. 76 (2) (1994), 365–416.
- [30] H. Nakajima, Reflection functors for quiver varieties and Weyl group actions, Math. Ann. 327 (2003), no. 4, 671–721.
- [31] C. Simpson, Products of Matrices, in Differential geometry, global analysis, and topology (Halifax, NS, 1990), Canadian Math. Soc. Conf. Proc. 12, Amer. Math. Soc. Providence, RI, 1991, 157–185.
- [32] K. Takemura, Introduction to middle convolution for differential equations with irregular singularities, in New trends in quantum integrable systems, 393–420, World Sci. Publ., Hackensack, NJ, 2011.
- [33] W. Wasow, Asymptotic expansions for ordinary differential equations. Pure and Applied Mathematics, Vol. XIV Interscience Publishers John Wiley & Sons, Inc 1965.
- [34] H. Völklein, The braid group and linear rigidity, Geom. Dedicata, 84, no. 1–3 (2001), 135–150.
- [35] D. Yamakawa, Geometry of Multiplicative Preprojective Algebra, Int. Math. Res. Pap. IMRP 2008 (2008), 77pp.
- [36] D. Yamakawa, Quiver varieties with multiplicities, Weyl groups of non-symmetric Kac-Moody algebras, and Painlevé equations, SIGMA Symmetry Integrability Geom. Methods Appl. 6 (2010), 215–262.
- [37] D. Yamakawa, Middle convolution and Harnad duality, Math. Ann., 349, no. 1 (2011), 215–262.

E-mail address: kazuki@josai.ac.jp

DEPARTMENT OF MATHEMATICS, JOSAI UNIVERSITY,, 1-1 KEYAKIDAI SAKADO-SHI SAITAMA 350-0295 JAPAN.