# NON-REMOVABLE TERM ERGODIC ACTION SEMIGROUPS/GROUPS

ALI SARIZADEH

ABSTRACT. In this work, we introduce the concept of term ergodicity for action semigroups and construct semigroups on two dimensional manifolds which are  $C^{1+\alpha}$ -robustly term ergodic. Moreover, we illustrate the term ergodicity by some exciting examples.

At last, we study a problem in the context of circle packing which is concerned to term ergodic.

# INTRODUCTION

As is well known, in ordinary dynamical system (M, f), where M is a compact metric space and f is a map from M to itself, an invariant probability measure  $\mu$  is said to be ergodic if every invariant measurable set is either of zero or full  $\mu$ -measure. The definition of ergodicity, for action semigroups/groups generated by  $\mathcal{G} = \{g_1, \ldots, g_s\}$  on M, is naturally extended to quasi-invariant measure<sup>1</sup>. Recall that a quasi-invariant measure is ergodic if every measurable invariant set with respect to  $\mathcal{G}$  is either of zero or full measure. In view of topological theory, counterpart to ergodicity is minimality. More precisely, an action semigroup/group is said to be minimal if each closed invariant subset is either empty or coincides with the whole of space. Minimal systems have been studied extensively by many authors, see for instance [2, 4, 7, 5, 9, 1]. Authors in [5], provided an example of an action semigroup/group generated by two circle diffeomorphisms, that is robustly minimal in the  $C^1$ -topology. In [4], this one-dimensional example is generalized to an action semigroup/group on higher-dimensional manifolds which is also  $C^1$ -robustly minimal. Recently, in [7], the authors provided an action semigroup generated by two diffeomorphisms on any compact manifold that is  $C^1$ -robustly minimal. In typical papers, finding a local invariant set with non-empty interior plays a key role.

Now, we are going to concentrate on some relations between ergodicity and minimality in action semigroups/groups. Notice, there are some examples of ergodic action semigroups/groups having global fixed points. So, in

<sup>2000</sup> Mathematics Subject Classification. 26A18, 37A99, 28A20.

*Key words and phrases.* action semigroup, action group, minimality, ergodicity, robust property, circle packing.

<sup>&</sup>lt;sup>1</sup>A measure  $\mu$  is said to be quasi-invariant if  $(g_i)_*\mu$  is absolutely continuous with respect to  $\mu$  for every element  $g_i$  in  $\mathcal{G}$ .

#### A. SARIZADEH

general, ergodicity does not imply minimality. Thus, in the opposite direction, a natural question arises: which system having the minimal property is ergodic? To answer this question one can refer to an earlier result contained in [2] which allows to solve the following conjecture concerning the one-dimensional case in the affirmative under some additional assumptions, although the conjecture and the question are apparently far from each other.

**Conjecture 1.** Every minimal smooth action of a finitely generated group on the circle is ergodic with respect to the Lebesgue measure.

In this note, we prove the above mentioned conjecture under some assumption which is not unusual. Actually, by this assumption, we insure that our result does not have any contradiction with results by Furstenberg [3].

On the other hand, if there exists a relation between minimality and ergodicity, it is natural to get, as a corollary, robustness of ergodic systems. By these results, we provide an example of some action semigroup/group which is both of robustly minimal and robustly term ergodic.

Finally, as an application of this note, we study the branch of mathematics generally known as *circle packing*. The packing problem is concerned with how to pack a number of objects, each with given shape and size, into bounded region without overlap (see more details on this context [6], also see http://hydra.nat.uni-magdeburg.de/packing/cci/#Overview). Here, we consider a problem of packing of circles with unequal radii and some additional property into a given circle. In fact, by benefit the concept of term ergodicity, we this problem in the circle packing.

### 1. Main results

We begin to introduce some definitions and notations and then formulate our main results. Throughout this paper, M stands for a smooth compact Riemannian manifold and Vol is normalized volume. Also, consider the space  $\text{Diff}^1(M)$  of  $C^1$  diffeomorphisms of M, endowed with the  $C^1$ -topology. A point  $x \in M$  is a Lebesgue density point of measurable set  $A \subseteq M$  if

$$\lim_{r \to 0} \operatorname{Vol}(A : B(x, r)) = \frac{\operatorname{Vol}(A \cap B(x, r))}{\operatorname{Vol}(B(x, r))} = 1,$$

where B(x, r) is the geodesic ball of radius r centered about x. Denote by DP(A) the set of Lebesgue density points of a measurable set A. By Lebesgue density point theorem, for every measureable set A,

$$\operatorname{Vol}(A \bigtriangleup DP(A)) = 0.$$

Now, consider a collection of diffeomorphisms  $\mathcal{G} = \{g_1, \ldots, g_s\}$  on M. Write  $\mathcal{G}^{-1} = \{g_1^{-1}, \ldots, g_s^{-1}\}$ . The action semigroup  $\langle \mathcal{G} \rangle^+$  generated by  $\mathcal{G}$  is given by

$$\langle \mathcal{G} \rangle^+ = \{h: M \to M: h = g_{i_n} \circ \cdots \circ g_{i_1}, i_j \in \{1, \dots, s\}\} \cup \{\mathrm{id}: M \to M\}$$

Furthermore, the action group  $\langle \mathcal{G} \rangle$  generated by  $\mathcal{G}$  is defined by  $\langle$  $\mathcal{G} \bigcup \mathcal{G}^{-1} >^+$ . So, every action group is not more than an action semigroup. Notice that  $\lim_{r\to\infty} f_{\omega}^r \notin \mathcal{G} >^+$  where  $f_{\omega_1,\ldots,\omega_r}^r = f_{\omega_n} \circ f_{\omega_1,\ldots,\omega_{r-1}}^{r-1}$ . Also, we denote the reverse iteration by  $\hat{f}_{\omega_1,\dots,\omega_r}^r = f_{\omega_1} \circ f_{\omega_2,\dots,\omega_r}^{r-1}$ . Let us recall that the subset  $\mathcal{K}$  of M is invariant with respect to an action

semigroup  $\Gamma$  generated by  $\mathcal{G} = \{g_1, \ldots, g_s\}$  if

$$\mathcal{K} = \bigcup_{i=1}^{s} g_i(\mathcal{K}).$$

Also,  $\Gamma$  is said to be *minimal* if each closed invariant subset A of M with respect to  $\Gamma$  is empty or coincides with M.

Observe that for ordinary dynamical system (M, f), the minimality of f is equivalent to that of  $f^{-1}$ . This is not the case for dynamical systems with several maps: there exists a minimal action semigroup  $\langle f_1, \ldots, f_s \rangle^+$  on the circle such that  $\langle f_1^{-1}, \ldots, f_s^{-1} \rangle^+$  is not minimal [9].

**Definition 1.** Given an action semigroup  $\Gamma$  generated by  $\mathcal{G} = \{g_1, \ldots, g_s\}$ and a probability measure space  $(M, \mathcal{M}, \mu)$  which  $\mu$  is quasi-invariant with respect to  $\Gamma$ . The measure  $\mu$  is called ergodic if for every measurable set  $B \in \mathcal{M}$  with

$$g_i^{-1}(B) = B; \quad \forall \ i = 1, \dots, s$$

we have that either  $\mu(B) = 0$  or  $\mu(B) = 1$ .

Here, to obtain term ergodic results for an action semigroup/group, we need just  $C^{1+\alpha}$ -regularity. In this regard, we begin by stating term ergodic result for an action semigroup.

**Theorem A.** Every boundaryless compact two dimensional manifold M admits a finite set of  $C^{1+\alpha}$ -diffeomorphisms that generates a  $C^{1+\alpha}$ -robustly term ergodic action semigroup with respect to volume measure.

We recall that a property  $\mathcal{P}$  holds  $C^r$ -robustly for action semigroup  $\Gamma$ generated by  $\mathcal{G} = \{g_1, \ldots, g_s\}$  if it holds for action semigroup  $\Gamma$  generated by  $\mathcal{F} = \{f_1, \ldots, f_s\}$  whose elements are  $C^r$ -perturbations of elements of  $\mathcal{G}$ .

Since every action group is action semigroup the following corollary is an immediately consequence of Theorem A.

**Corollary 1.1.** Theorem A is valid for action groups.

### 2. Some new results about minimality

First of all, we state some results about minimality on compact manifolds in any dimension.

**Lemma 2.1.** Let M be a compact manifold and action semigroup generated by homeomorphisms  $\mathcal{G} = \{g_1, \ldots, g_s\}$  be a minimal. Then every invariance set respect to each  $g_i$  for i = 1, ..., s and its complement are dens in the whole space.

*Proof.* Suppose that

$$g_i(\mathcal{B}) = \mathcal{B}; \quad \forall \ i = 1, \dots, s.$$

So we have

$$\mu_i(\overline{\mathcal{B}}) = \overline{\mathcal{B}}; \quad \forall \ i = 1, \dots, s.$$

When  $\overline{\mathcal{B}}$  is closed and  $\langle \mathcal{G} \rangle^+$  is minimal on M then  $\overline{\mathcal{B}} = M$ . On the other hand  $g_i(\mathcal{B}^c) = \mathcal{B}^c$  for i = 1, ..., s, where  $\mathcal{B}^c$  is complement of  $\mathcal{B}$ . By minimality of  $\langle \mathcal{G} \rangle^+$  and invariance of the subset  $\mathcal{B}^c$  with respect to each generator, one can have density of this subset in M; that is  $\overline{\mathcal{B}^c} = M$ .  $\Box$ 

It follows that

$$\mathcal{B} \cap B(x,r) \neq \emptyset$$
 and  $\mathcal{B}^c \cap B(x,r) \neq \emptyset$ 

for each real number r > 0 and  $x \in M$ .

**Lemma 2.2.** Under the assumption of lemma 2.1 on a compact Riemannian manifold M, density points of every invariance set respect to each  $g_i$  for i = 1, ..., s, with positive volume is dens in the whole space.

*Proof.* Suppose that  $0 < \operatorname{Vol}(\mathcal{B})$  and

$$q_i(\mathcal{B}) = \mathcal{B}; \quad \forall \ i = 1, \dots, s.$$

If  $DP(\mathcal{B})$  is not dense then there exists a neighborhood  $B(x_0, r_0)$  for some point  $x_0$  of M so that  $B(x_0, r_0) \cap DP(\mathcal{B}) = \emptyset$ . By Lebesgue density point theorem, one can have  $\operatorname{Vol}(B(x_0, r_0) \cap \mathcal{B}) = 0$ , equivalently  $\operatorname{Vol}(B(x_0, r_0) \cap \mathcal{B}^c) =$  $\operatorname{Vol}(B(x_0, r_0))$ . We remark that the volume measure is a quasi-invariant for any  $C^1$ -diffeomorphism. So, the union of iterates  $B(x_0, r_0) \cap \mathcal{B}$  under  $\mathcal{G}$ has zero volume measure. Now, the assumption of minimality  $\langle \mathcal{G} \rangle^+$  and invariance of  $\mathcal{B}^c$  under  $\mathcal{G}$  yields a contradiction with  $0 < \operatorname{Vol}(\mathcal{B})$ . Hence  $\overline{DP(\mathcal{B})} = M$ .

Notice that, similarly, when  $0 < Vol(\mathcal{B}) < 1$  both the subsets  $DP(\mathcal{B})$  and  $DP(\mathcal{B}^c)$  are dense.

**Lemma 2.3.** Let M be a compact connected two dimensional manifold. Then there exist a set  $\mathcal{H} = \{h_1, \ldots, h_k\}$  of  $C^1$ -diffeomorphisms on M and invariant set  $\Delta$  with respect to  $\mathcal{H}$  and nonempty interior so that  $\langle \mathcal{H} \rangle^+$ is minimal on  $\Delta$  and for every  $x \in \Delta$  and  $i = 1, \ldots, k$ ,  $Dh_i$  at x have two complex eigenvalues.

*Proof.* Let A be rotation matrix by angle  $\theta = 179^{\circ}$ . Define a linear map T as the follows,

$$T(x,y) = \kappa \cdot A(x,y);$$
 for  $ever(x,y) \in \mathbb{R}^2$ ,

where  $3/4 < \kappa < 1$ . The choice of  $\theta$  insure that each eigenvalues of T at each point is complex. Put  $V = B(0, \delta)$ . Clearly,  $T(\overline{V}) \subset V$  and for every  $x \in \overline{V}$ , DT at x have two complex eigenvalues  $\lambda, \overline{\lambda}$  with  $|\lambda| = |\overline{\lambda}| > 3/4$ . For every  $y \in W$ , define

$$S_u(x) = T(x-y) + y: \quad \forall \ x \in V$$

where  $W = \{x \in V : |x| = \frac{3}{4}\delta\}$ . Observe that, by construction of T and  $S_u$ , the following set

$$T(V) \bigcup \left[\bigcup_{y \in W} S_y(V)\right]$$

is a cover for  $\overline{V}$ . Therefor, it has a finite subcover as  $\overline{V} \subset T(V) \bigcup [\bigcup_{i=1}^{k-1} S_i(V)]$ Take  $\mathcal{H}_0 = \{T, S_1, \ldots, S_{k-1}\}$ . Since every element of  $\mathcal{H}_0$  is contractions, there is a ball U that is mapped into itself by T and  $S_1, \ldots, S_{k-1}$ , i.e.,  $\mathcal{H}_0(U) = T(U) \bigcup [\bigcup_{i=1}^{k-1} S_i(U)] \subset U$  (when  $\kappa$  is very close to 3/4 one can take  $U = B(0, 16\delta)$  to have this property). Thus

$$\lim_{n \to \infty} \mathcal{H}_0^n(U) = \lim_{n \to \infty} \mathcal{H}_0^{n-1}[\mathcal{H}_0(U)] = \Delta$$

is the unique non-empty invariant set with respect to  $\mathcal{H}_0$  which the interior of it is non-empty. Notice that the semigroup generated by  $\mathcal{H}_0$  on  $\Delta$  is minimal.

Take a gradient Morse-Smale vector fields  $\dot{x} = \nabla F_i(x)$  on M with a unique hyperbolic attracting equilibrium  $p_i$ , for  $i = 1, \ldots, k$  (see e.g. Theorem 3.35 of [10], for the existence of Morse functions F with unique extrema) and let  $h_i$  be its time-1 map. We may assume that each  $p_i$  belong to an open neighborhood  $\hat{V}$  and the each eigenvalue of  $Dh_i(p_i)$  are complex.

Now, working in a coordinate chart on a small open neighborhood  $\hat{V} \subset \mathbb{R}^2$ and  $U \subset \hat{V}$ . One can assume that  $h_1 = T$  and  $h_i = S_i$  on U for  $i = 2, \ldots, k$ . The action semigroup generated by  $\mathcal{H} = \{h_1, \ldots, h_k\}$  is minimal on the set  $\Delta$ .

Since the construction used in Lemma 2.3 is  $C^1$ -robust, by the similar argument used in [4] and [7], we can have a finite extension  $\mathcal{G}$  of  $\mathcal{H}$  so that  $\langle \mathcal{G} \rangle^+$  is  $C^1$ -robust minimal on M.

**Corollary 2.4.** There is finite extension of  $\mathcal{H}$  to finite set  $\mathcal{G}$  which  $\langle \mathcal{G} \rangle^+$  is  $C^1$ - robustly minimal.

## 3. Robustly term ergodic: Proof of Theorem A

We used the following lemma to prove of main theorem.

**Lemma 3.1** (Bounded distortion in the Hutchinson attractor). Consider a finite family  $\mathcal{H} = \{h_1, \ldots, h_k\} \subset \operatorname{Hom}(M)$  where each  $h_i$  is a contracting  $C^{1+\alpha}$ -map of the closure of an open set  $D \subset M$ . Then, there exists  $L_{\mathcal{H}} > 0$  such that for every  $n \in \mathbb{N}$  and  $\omega \in \Sigma_k^+$ ,

$$L_{\mathcal{H}}^{-1} < \left| \frac{\det(D\hat{h}_{\omega}^{n}(x))}{\det(D\hat{h}_{\omega}^{n}(y))} \right| < L_{\mathcal{H}} \quad for \ all \ x, y \in \Delta_{\mathcal{H}}$$

where  $\triangle_{\mathcal{H}}$  is the Hutchinson attractor of IFS( $\mathcal{H}$ ) in  $\overline{D}$ .

*Proof.* Define  $\Phi$  : GL(dim(M),  $\mathbb{R}$ )  $\to \mathbb{R}$  by  $\Phi(A) = \log |\det(A)|$  and  $\digamma_i(x) = \Phi(Dh_i(x))$ , for any  $i = 1, \ldots, k$ . Note that by assumption,  $\log |\det Dh_i|$  is  $\alpha$ -Hölder and thus for every  $x, y \in \Delta_{\mathcal{H}}$  and  $1 \le i \le k$ 

$$|\mathcal{F}_i(x) - \mathcal{F}_i(y)| \le C ||x - y||^{\alpha},$$

for some constants C > 0. On the other hand, for every  $\omega = \omega_1 \omega_2 \cdots \in \Sigma_k^+$ ,  $\|\hat{h}^n_{\omega}(x) - \hat{h}^n_{\omega}(y)\| \le \|Dh_{\omega_1}\| \|\hat{h}^{n-1}_{\omega_2,\dots,\omega_n}(x) - \hat{h}^{n-1}_{\omega_2,\dots,\omega_n}(y)\| \le \xi^n \|x-y\| \le \xi^n \operatorname{diam}(\Delta_{\mathcal{H}}),$ where

$$\xi = \sup_{x \in \Delta_{\mathcal{H}}, \ 1 \le i \le k} \|Dh_i(x)\| < 1.$$

Hence,

$$\log \frac{|\det\left(D\hat{h}_{\omega}^{n}(x)\right)|}{|\det\left(D\hat{h}_{\omega}^{n}(y)\right)|} = \sum_{i=0}^{n-1} |\mathcal{F}_{\omega_{i}}(\hat{h}_{\omega}^{i}(x)) - \mathcal{F}_{\omega_{i}}(\hat{h}_{\omega}^{i}(y))| \le C \sum_{i=0}^{n-1} \|\hat{h}_{\omega}^{i}(x) - \hat{h}_{\omega}^{i}(y)\|^{\alpha}$$
$$\le C \sum_{i=0}^{n-1} \{\xi^{i} \operatorname{diam}(\triangle_{\mathcal{H}})\}^{\alpha} \le C M(\operatorname{diam}(\triangle_{\mathcal{H}}))^{\alpha} \sum_{i=0}^{\infty} (\xi^{\alpha})^{i}.$$

Taking  $L_{\mathcal{H}} = \exp\{C\xi^{\alpha}(\operatorname{diam}(\Delta_{\mathcal{H}}))^{\alpha}/(1-\xi^{\alpha})\}$  the desired inequality holds.

Now, we will prove Theorem A.

Proof of Theorem A. Suppose that M is two dimensional manifold. Consider Lemma 2.3 for  $C^{1+\alpha}$ -diffeomorphisms  $\mathcal{H} = \{h_1, \ldots, h_k\}$ , which will provide an invariant set  $\Delta$  with nonempty interior respect to action semigroup  $\langle \mathcal{H} \rangle^+$  so that the  $Dh_i(x)$  have complex eigenvalue for each  $i = 1, \ldots, k$  and  $x \in \Delta^\circ$ . Take the subset U of M so that is mapped into itself by  $h_i$ , i.e.,  $\mathcal{H}(U) = \bigcup_{i=1}^k h_i(U) \subset U$  and also take  $\langle \mathcal{G} \rangle^+$  is a finite  $C^{1+\alpha}$ -extension of  $\mathcal{H}$  which is  $C^{1+\alpha}$ -robustly minimal on M (see Corollary 2.4).

We claim that  $< \mathcal{G}^{-1} >^+$  is term ergodic. To this end, suppose that  $0 < \operatorname{Vol}(\mathcal{B}) < 1$  and

$$g_i(\mathcal{B}) = \mathcal{B}; \quad \forall \ i = 1, \dots, s.$$

Now, let  $p \in \Delta^{\circ}$  and  $J = B(p, \hat{\delta}) \subset \Delta^{\circ}$ . Since  $Dh_i(x)$  have two complex eigenvalue  $\lambda_i, \overline{\lambda_i}$  for every  $x \in U$  and  $i = 1, \ldots, k$ , the image of an open ball under the apply of  $h_i$  is a ball, too.

On the other hand, there is  $\omega \in \Sigma_k^+$  so that  $\bigcap_r \hat{h}^r_{\omega}(U) = \{p\}$  and  $\lim_{r\to\infty} \operatorname{diam}(\hat{h}^r_{\omega}(U)) \to 0$ . So,  $\hat{h}^r_{\omega}(U)$  is a ball for each r and  $\operatorname{diam}(\hat{h}^r_{\omega}(U) \to 0$  as  $r \to \infty$ . Define

 $r_0 = \min\{r : \operatorname{diam}(\hat{h}^r_{\omega}(U)) < \delta\}.$ 

Clearly,  $p \in \hat{h}^{r_0}_{\omega}(U) \subset B(p, \delta)$ . Since  $\hat{h}^{r_0}_{\omega}(U)$  is a ball with diam $(\hat{h}^{r_0}_{\omega}(U)) < \delta$  and diam $(\hat{h}^{r_0-1}_{\omega}(U)) \ge \delta$ , then

$$\operatorname{diam}(h_i(B(p,\delta/2))) \le \operatorname{diam}(h_{\omega}^{r_0}(U)) < \delta$$

and

$$\frac{\operatorname{Vol}(h_i(B(p,\delta/2)))}{\operatorname{Vol}(J)} \le \frac{\operatorname{Vol}(h_{\omega}^{r_0}(U))}{\operatorname{Vol}(J)}.$$

Notice that  $h^{r_0}_{\omega}(U) \subset J$  and  $h_i(\mathcal{B}^c) = \mathcal{B}^c$  for every  $i = 1, \ldots, k$  when  $\mathcal{H}$  is a subset of  $\mathcal{G}$ . So, one can have  $h^{r_0}_{\omega}(\mathcal{B}^c \cap U) \subseteq \mathcal{B}^c \cap h^{r_0}_{\omega}(U)$ .

Hence, we have

$$\frac{\operatorname{Vol}(\mathcal{B}^{c} \cap J)}{\operatorname{Vol}(J)} \geq \frac{\operatorname{Vol}(\mathcal{B}^{c} \cap \hat{h}_{\omega}^{r_{0}}(U))}{\operatorname{Vol}(J)} \\
\geq \frac{\operatorname{Vol}(\mathcal{B}^{c} \cap \hat{h}_{\omega}^{r_{0}}(U))}{\operatorname{Vol}(\hat{h}_{\omega}^{r_{0}}(U))} \cdot \frac{\operatorname{Vol}(\hat{h}_{\omega}^{r_{0}}(U))}{\operatorname{Vol}(J)} \\
\geq \frac{\operatorname{Vol}(\hat{h}_{\omega}^{r_{0}}(\mathcal{B}^{c} \cap U))}{\operatorname{Vol}(\hat{h}_{\omega}^{r_{0}}(U))} \cdot \frac{\operatorname{Vol}(h_{i}(B(p, \delta/2)))}{\operatorname{Vol}(J)}.$$

On the other hand  $\mathcal{B}^c$  is forward  $\mathcal{H}$ -invariant,

$$\frac{\operatorname{Vol}(\hat{h}^{r_0}_{\omega}(\mathcal{B}^c \cap U))}{\operatorname{Vol}(\hat{h}^{r_0}_{\omega}(U))} \ge L_{\mathcal{H}}^{-1} \frac{\operatorname{Vol}(\mathcal{B}^c \cap U)}{\operatorname{Vol}(U)}$$

Indeed, the last inequality is implied by the bounded distortion result, Lemma 3.1. Indeed, since

$$\operatorname{Vol}(\hat{h}^{r_0}_{\omega}(\mathcal{B}^c \cap U)) = \int_{\mathcal{B}^c \cap U} |\det D\hat{h}^{r_0}_{\omega}| \, d\operatorname{Vol} \ge \operatorname{Vol}(\mathcal{B}^c \cap U) \inf_{x \in \Delta} |\det D\hat{h}^{r_0}_{\omega}(x)|,$$
$$\operatorname{Vol}(\hat{h}^{r_0}_{\omega}(U)) = \int_U |\det \hat{h}^{r_0}_{\omega}| \, d\operatorname{Vol} \le \operatorname{Vol}(U) \sup_{x \in \Delta} |\det D\hat{h}^{r_0}_{\omega}(x)|$$

This means that  $\frac{\operatorname{Vol}(\mathcal{B}^c \cap J)}{\operatorname{Vol}(J)}$  is bounded from below for every neighborhood J of p. So,  $p \notin DP(\mathcal{B})$ . Similarly, One can prove that  $p \notin DP(\mathcal{B}^c)$ . Hence

$$\Delta^{\circ} \bigcap (DP(\mathcal{B}) \bigcup DP(\mathcal{B}^{c})) = \emptyset$$

which is a contradiction with Lemma 2.2 and the proof is completed.  $\Box$ 

Observe that ergodicity may be removed from some ordinary dynamical systems under a perturbation which an irrational translation is such a system. But the following example, containing irrational translation, is robust term ergodic. Moreover, it is shown that sufficiently close to such system in the  $C^{1+\alpha}$ -topology, there is a term ergodic action semigroup which each of generators are not ergodic.

**Example 3.2.** Suppose  $f_1$  is a north-south pole  $C^2$ -diffeomorphism of the circle  $\mathbb{S}^1$  possessing an attracting fixed point p as a north pole and a repelling fixed point q as south pole with multipliers

$$1/2 < f'_1(p) < 1$$
 and  $1/2 < (f_1^{-1})'(q) < 1$ .

Consider the map  $f_2 = R_{\lambda}$  where  $R_{\lambda}$  is the rotation by irrational angle  $\lambda$  on the circle. Observe that, both systems  $\langle f_1, f_2 \rangle^+$  and  $\langle f_1^{-1}, f_2^{-1} \rangle^+$  are  $C^{1+\alpha}$ - robust minimal [5]. Thus the systems  $\langle f_1, f_2 \rangle^+$  and  $\langle f_1^{-1}, f_2^{-1} \rangle^+$ 

#### A. SARIZADEH

are  $C^{1+\alpha}$ -robust term ergodic. Moreover, take a rational number  $\gamma$  sufficiently close to  $\lambda$  so that the system  $\langle f_1, R_{\gamma} \rangle^+$  is minimal, too. Clearly, none of  $f_1, f_1^{-1}, R_\gamma, R_\gamma^{-1}$  is neither ergodic nor minimal but,  $\langle f_1, R_\gamma \rangle^+$  is both of minimal and term ergodic.

# 4. A PROBLEM ON CIRCLE PACKING CONCERNING TO ERGODICITY

Let M be a compact 2-dimensional manifold and  $\Gamma = \langle \mathcal{G} | \mathcal{G}^{-1} \rangle^+$  be a minimal action group generated by homeomorphisms  $\mathcal{G} = \{g_1, \ldots, g_s\}$ . Suppose that  $\mathcal{B}$  is an invariant set with respect to each generator of  $\Gamma$ , that is

$$g_i(\mathcal{B}) = \mathcal{B}; \quad \forall \ i = 1, \dots, s.$$

Assume that  $0 < Vol(\mathcal{B}) < 1$ . By Lemmas 2.1 and 2.2, we insure that  $\mathcal{B}, \mathcal{B}^c, DP(\mathcal{B}), DP(\mathcal{B}^c)$  are dense in M.

Now, let y be an arbitrary point of  $DP(\mathcal{B})$ . By definition of density point, one can find  $\delta > 0$  so that

$$\operatorname{Vol}(\mathcal{B} \bigcap B(y, \delta)) > \frac{3}{4} \operatorname{Vol}(B(y, \delta)).$$

One kind of circle packing problem, with respect to dynamical system, may be as follows.

**Problem 4.1.** Under the above assumptions, is there a family  $\{B(p, \delta_p)\}_{p \in \mathcal{P}}$ with  $\mathcal{P} \subset DP(\mathcal{B}^c) \cap B(y, \delta)$  so that

- $\begin{array}{ll} (\mathrm{i}) & B(p,\delta_p) \subseteq B(y,\delta), \\ (\mathrm{ii}) & \forall \; p,q \in \mathcal{P} \; \mathrm{with} \; p \neq q; \quad B(p,\delta_p) \cap B(q,\delta_q) = \emptyset, \\ (\mathrm{iii}) & \frac{2}{3} \operatorname{Vol}(B(y,\delta)) < \operatorname{Vol}(\bigcup_{p \in \mathcal{P}} B(p,\delta_p)) \; \mathrm{and} \\ (\mathrm{iv}) \; \operatorname{Vol}(\mathcal{B}^c \cap B(p,\delta_p)) > \frac{1}{2} \operatorname{Vol}(B(p,\delta_p)). \end{array}$

In general and without any additional assumption, the answer of the problem is negative. Actually, one can prove that the action group  $\Gamma$  is term ergodic with respect to volume measure if there is a family  $\{B(p, \delta_p)\}_{p \in \mathcal{P}}$ which satisfies in the properties (i)-(iv). Indeed, one can have

$$\operatorname{Vol}(\mathcal{B}^{c} \cap B(y, \delta)) = \operatorname{Vol}[\bigcup_{p \in \mathcal{P}} (\mathcal{B}^{c} \cap B(p, \delta_{p}))]$$
$$= \sum_{p \in \mathcal{P}} \operatorname{Vol}(\mathcal{B}^{c} \cap B(p, \delta_{p}))$$
$$> \sum_{p \in \mathcal{P}} \frac{1}{2} \operatorname{Vol}(B(p, \delta_{p}))$$
$$= \frac{1}{2} \operatorname{Vol}(\cup_{p \in \mathcal{P}} B(p, \delta_{p}))$$
$$= \frac{1}{3} \operatorname{Vol}(B(y, \delta)),$$

which is a contradiction and so  $\operatorname{Vol}(\mathcal{B}) \in \{0, 1\}$ . Therefore, by Theorem A, minimality of action group  $\langle \mathcal{G} \bigcup \mathcal{G}^{-1} \rangle^+$  implies term ergodicity of it which is lead to the following counterexample.

**Example 4.2.** In [3], Furstenberg constructed an analytic diffeomorphism T of tours which preserve Haar measure and is minimal but not ergodic. So,  $\Gamma = \langle T, T^{-1} \rangle^+$  is minimal action group which is not a term ergodic. This is a contradiction.

#### Acknowledgments

We thank A. Fakhari, A. J. Homburg, M. Nassiri, A. Navas and F. Khosh-Ahang for useful discussions and suggestions.

## References

- 1. P. G. Barrientos and A. Raibekas Dynamics of iterated function systems on the circle close to rotations. ArXiv e-prints, Mar. 2013.
- B. Deroin, V. Kleptsyn, A. Navas. On the question of ergodicity for minimal group actions on the circle. Moscow Math. Journal 9:263–303, 2009.
- H. Furstenberg Strict Ergodicity and Transformation of the Torus. American Journal of Mathematics Vol. 83, No. 4, pp. 573-601, 1961.
- F. H. Ghane, A. Homburg, A. Sarizadeh. C<sup>1</sup>-robustly minimal iterated function systems. Sto. Day. 10:155-160, 2010.
- A. S. Gorodetskiĭ and Yu. S. Il'yashenko. Certain properties of skew products over a horseshoe and a solenoid. Grigorchuk, R. I. (ed.), Dynamical systems, automata, and infinite groups. Proc. Steklov Inst. Math. 231, 90-112, 2000.
- 6. A. Grosso, A.R.M.J.U. Jamali, M. Locatelli, F. Schoen, Solving the problem of packing equal and unequal circles in a circular container, Optimization Online, March 2008.
- 7. A. Homburg, M. Nassiri. Robust minimality of iterated function systems with two generators. To appear in Ergodic Theory and Daynamical Systems.
- 8. J. E. Hutchinson. Fractals and self-similarity. Indiana Univ. Math. J., 30:713747, 1981.
- V. A. Kleptsyn and M. B. Nalskii. Contraction of orbits in random dynamical systems on the circle. *Funct. Anal. Appl.*, 38(4):267–282, 2004.
- 10. Y. Matsumoto. An introduction to Morse theory, volume 208 of Translations of Mathematical Monographs. American Mathematical Society, Providence, RI, 2002.

DEPARTMENT OF MATHEMATICS, ILAM UNIVERSITY, P.O.Box 69315-516, ILAM, IRAN. *E-mail address:* ali.sarizadeh@gmail.com