A note on $U_q(D_4^{(3)})$ $\binom{3}{4}$ - Demazure crystals

Alyssa M. Armstrong and Kailash C. Misra

ABSTRACT. We show that there exists a suitable sequence $\{w^{(k)}\}_{k\geq 0}$ of Weyl group elements for the perfect crystal $B = B^{1,3l}$ such that the path realizations of the Demazure crystals $B_{w^{(k)}}(l\Lambda_2)$ for the quantum affine algebra $U_q(D_4^{(3)})$ have tensor product like structure with mixing index $\kappa = 1$.

1. Introduction

Consider the quantum affine algebra $U_q(\mathfrak{g})$ (cf. [[13](#page-9-0)]) associated with the affine Lie algebra $\mathfrak g$ (cf. [[2](#page-9-1)]). For a dominant integral weight λ of level $l > 0$, let $V(\lambda)$ denote the integrable highest weight $U_q(\mathfrak{g})$ -module with highest weight λ and the pair $(L(\lambda), B(\lambda))$ denote its crystal base ([[6](#page-9-2)], [[8](#page-9-3)], [[12](#page-9-4)]). The crystal $B(\lambda)$ has many interesting combinatorial properties and can be realized as elements (called paths) in the semi-infinite tensor product $\cdots \otimes B \otimes B \otimes B$ where B is a perfect crystal of level l for $U_q(\mathfrak{g})$ [[3](#page-9-5)]. A perfect crystal B of level l for the quantum affine algebra $U_q(\mathfrak{g})$ can be thought of as a crystal for a level zero representation of the derived subalgebra $U_q'(\mathfrak{g})$ with certain properties.

Let W denote the Weyl group for $\mathfrak g$ and $U_q^+(\mathfrak g)$ denote the positive part of $U_q(\mathfrak g)$. For each $w \in W$, the $U_q^+(\mathfrak{g})$ -submodule $V_w(\lambda)$ generated by the one-dimensional extremal weight space $V(\lambda)_{w\lambda}$ of $V(\lambda)$ is called a Demazure module. In [[9](#page-9-6)], Kashiwara showed that the Demazure crystal $B_w(\lambda)$ for the Demazure module $V_w(\lambda)$ is a subset of the crystal $B(\lambda)$ for $V(\lambda)$ satisfying a certain recursive property. In [[10](#page-9-7)], it has been shown that under certain conditions, the Demazure crystal $B_w(\Lambda)$ as a subset of $\cdots \otimes B \otimes B \otimes B$ has tensor product-like structure. In this realization a certain parameter κ , called the *mixing index* enters the picture. It is conjectured that for all quantum affine algebras $\kappa \leq 2$. It is known that for $\lambda = l\Lambda$, where Λ is a dominant weight of level 1 the mixing index $\kappa = 1$ for $\mathfrak{g} = A_n^{(1)}, B_n^{(1)}, C_n^{(1)}, D_n^{(1)}, A_{2n-1}^{(2)}, D_{n+1}^{(2)}, A_{2n}^{(2)}$ $\mathfrak{g} = A_n^{(1)}, B_n^{(1)}, C_n^{(1)}, D_n^{(1)}, A_{2n-1}^{(2)}, D_{n+1}^{(2)}, A_{2n}^{(2)}$ $\mathfrak{g} = A_n^{(1)}, B_n^{(1)}, C_n^{(1)}, D_n^{(1)}, A_{2n-1}^{(2)}, D_{n+1}^{(2)}, A_{2n}^{(2)}$ [[11](#page-9-8)], $D_4^{(3)}$ [[14](#page-9-9)] and $G_2^{(1)}$ [1].

In this paper using the perfect crystal $B = B^{1,3l}$ of level 3l constructed in [[7](#page-9-11)] for $\mathfrak{g} = D_4^{(3)}$ we show that there exists a suitable sequence $\{w^{(k)}\}_{k \geq 0}$ of Weyl group elements in W such that the Demazure crystals $B_{w^{(k)}}(l\Lambda_2)$ has tensor product like structures with $\kappa = 1$. We observe that the dominant weight Λ_2 has level 3.

²⁰¹⁰ Mathematics Subject Classification. Primary 17B37,17B10; Secondary 17B67. Partially supported by NSA grant, H98230-12-1-0248.

2. Quantum affine algebras and the perfect crystals

In this section we recall necessary facts in crystal base theory for quantum affine algebras. Our basic references for this section are $\mathbf{[2]}, \mathbf{[13]}, \mathbf{[8]}, \mathbf{[3]},$ and $\mathbf{[5]}$ $\mathbf{[5]}$ $\mathbf{[5]}$.

Let $I = \{0, 1, ..., n\}$ be the index set and let $A = (a_{ij})_{i,j \in I}$ be an affine Cartan matrix and $D = diag(s_0, s_1, \ldots, s_n)$ be a diagonal matrix with all $s_i \in \mathbb{Z}_{>0}$ such that DA is symmetric. The *dual weight lattice* P^{\vee} is defined to be the free abelian group $P^{\vee} = \mathbb{Z}h_0 \oplus \mathbb{Z}h_1 \oplus \cdots \oplus \mathbb{Z}h_n \oplus \mathbb{Z}d$ of rank $n+2$, whose complexification $\mathfrak{h} = \mathbb{C} \otimes P^{\vee}$ is called the *Cartan subalgebra*. We define the linear functionals α_i and $Λ_i$ $(i \in I)$ on $η$ by

 $\alpha_i(h_i) = a_{ii}, \alpha_i(d) = \delta_{i0}, \Lambda_i(h_i) = \delta_{ii}, \Lambda_i(d) = 0 \ (i, j \in I).$

The α_i 's are called the *simple roots* and the Λ_i 's are called the *fundamental weights*. We denote by $\Pi = {\alpha_i \mid i \in I}$ the set of simple roots. We also define the *affine* weight lattice to be $P = {\lambda \in \mathfrak{h}^* \mid \lambda(P^{\vee}) \subset \mathbb{Z}}$. The quadruple (A, P^{\vee}, Π, P) is called an affine Cartan datum. We denote by g the affine Kac-Moody algebra corresponding to the affine Cartan datum (A, P^{\vee}, Π, P) (see [[2](#page-9-1)]). Let δ denote the null root and c denote the canonical central element for $\mathfrak g$ (see [[2](#page-9-1), Ch. 4]). Now the affine weight lattice can be written as $P = \mathbb{Z}\Lambda_0 \oplus \mathbb{Z}\Lambda_1 \oplus \cdots \oplus \mathbb{Z}\Lambda_n \oplus \mathbb{Z}\delta$. Let $P^+ = \{\lambda \in P \mid \lambda(h_i) \geq 0 \text{ for all } i \in I\}.$ The elements of P are called the affine weights and the elements of P^+ are called the *affine dominant integral weights*.

Let $\overline{P}^{\vee} = \mathbb{Z}h_0 \oplus \cdots \oplus \mathbb{Z}h_n$, $\overline{\mathfrak{h}} = \mathbb{C} \otimes_{\mathbb{Z}} \overline{P}^{\vee}$, $\overline{P} = \mathbb{Z}\Lambda_0 \oplus \mathbb{Z}\Lambda_1 \oplus \cdots \oplus \mathbb{Z}\Lambda_n$ and $\bar{P}^+=\{\lambda\in\bar{P}\mid \lambda(h_i)\geq 0\,\,\text{for all}\,\,i\in I\}.$ The elements of \bar{P} are called the *classical* weights and the elements of \bar{P}^+ are called the *classical dominant integral weights*. The level of a (classical) dominant integral weight λ is defined to be $l = \lambda(c)$. We call the quadruple $(A, \overline{P}^{\vee}, \Pi, \overline{P})$ the *classical Cartan datum*.

For the convenience of notation, we define $[k]_x = \frac{x^k - x^{-k}}{x - x^{-1}}$ $\frac{x}{x-x^{-1}}$, where k is an integer and x is a symbol. We also define $\begin{bmatrix} m \\ u \end{bmatrix}$ k 1 \boldsymbol{x} $=\frac{[m]_x!}{[1]_x!}$ $\frac{[k]_x!}{[k]_x! [m-k]_x!}$, where m and k are nonnegative integers, $m \ge k \ge 0$, $[k]_x! = [k]_x [k-1]_x \cdots [1]_x$ and $[0]_x! = 1$.

The quantum affine algebra $U_q(\mathfrak{g})$ is the quantum group associated with the affine Cartan datum (A, P^{\vee}, Π, P) . That is, it is the associative algebra over $\mathbb{C}(q)$ with unity generated by $e_i, f_i (i \in I)$ and $q^h(h \in P^{\vee})$ satisfying the following defining relations:

(i)
$$
q^0 = 1, q^h q^{h'} = q^{h+h'}
$$
 for all $h, h' \in P^{\vee}$,
\n(ii) $q^h e_i q^{-h} = q^{\alpha_i(h)} e_i, q^h f_i q^{-h} = q^{-\alpha_i(h)} f_i$ for $h \in P^{\vee}$,
\n(iii) $e_i f_j - f_j e_i = \delta_{ij} \frac{K_i - K_i^{-1}}{q_i - q_i^{-1}}$ for $i, j \in I$, where $q_i = q^{s_i}$ and $K_i = q^{s_i h_i}$,
\n(iv)
$$
\sum_{k=0}^{1-a_{ij}} (-1)^k e_i^{(1-a_{ij}-k)} e_j e_i^{(k)} = 0
$$
 for $i \neq j$,
\n(v)
$$
\sum_{k=0}^{1-a_{ij}} (-1)^k f_i^{(1-a_{ij}-k)} f_j f_i^{(k)} = 0
$$
 for $i \neq j$,

where $e_i^{(k)} = \frac{e_i^k}{[k]_{q_i}!}$, and $f_i^{(k)} = \frac{f_i^k}{[k]_{q_i}!}$. We denote by $U_q'(\mathfrak{g})$ the subalgebra of $U_q(\mathfrak{g})$ generated by $e_i, f_i, K_i^{\pm 1}$ ($i \in I$). The algebra $U'_q(\mathfrak{g})$ can be regarded as the quantum group associated with the classical Cartan datum $(A, \overline{P}^{\vee}, \Pi, \overline{P})$.

DEFINITION 2.1. An affine crystal (respectively, a classical crystal) is a set B together with the maps wt : $B \to P$ (respectively, wt : $B \to \overline{P}$), \tilde{e}_i , $\tilde{f}_i : B \to B \cup \{0\}$ and $\varepsilon_i, \varphi_i : B \to \mathbb{Z} \cup \{-\infty\}$ $(i \in I)$ satisfying the following conditions:

- (i) $\varphi_i(b) = \varepsilon_i(b) + \langle h_i, \text{wt}(b) \rangle$ for all $i \in I$,
- (ii) $\operatorname{wt}(\tilde{e}_i b) = \operatorname{wt}(b) + \alpha_i \text{ if } \tilde{e}_i b \in B,$
- (iii) $\operatorname{wt}(\tilde{f}_i b) = \operatorname{wt}(b) \alpha_i$ if $\tilde{f}_i b \in B$,
- (iv) $\varepsilon_i(\tilde{e}_i b) = \varepsilon_i(b) 1$, $\varphi_i(\tilde{e}_i b) = \varphi_i(b) + 1$ if $\tilde{e}_i b \in B$,
- (v) $\varepsilon_i(\tilde{f}_i b) = \varepsilon_i(b) + 1$, $\varphi_i(\tilde{f}_i b) = \varphi_i(b) 1$ if $\tilde{f}_i b \in B$,
- (vi) $\tilde{f}_i b = b'$ if and only if $b = \tilde{e}_i b'$ for $b, b' \in B, i \in I$,
- (vii) If $\varphi_i(b) = -\infty$ for $b \in B$, then $\tilde{e}_i b = \tilde{f}_i b = 0$.

DEFINITION 2.2. Let B_1 and B_2 be affine or classical crystals. A *crystal mor*phism (or morphism of crystals) $\Psi : B_1 \to B_2$ is a map $\Psi : B_1 \cup \{0\} \to B_2 \cup \{0\}$ such that

- (i) $\Psi(0) = 0$,
- (ii) if $b \in B_1$ and $\Psi(b) \in B_2$, then $wt(\Psi(b)) = wt(b)$, $\varepsilon_i(\Psi(b)) = \varepsilon_i(b)$, and $\varphi_i(\Psi(b)) = \varphi_i(b)$ for all $i \in I$,
- (iii) if $b, b' \in B_1$, $\Psi(b), \Psi(b') \in B_2$ and $\tilde{f}_i b = b'$, then $\tilde{f}_i \Psi(b) = \Psi(b')$ and $\Psi(b) = \tilde{e}_i \Psi(b')$ for all $i \in I$.

A crystal morphism $\Psi : B_1 \to B_2$ is called an *isomorphism* if it is a bijection from $B_1 \cup \{0\}$ to $B_2 \cup \{0\}$.

For crystals B_1 and B_2 , we define the *tensor product* $B_1 \otimes B_2$ to be the set $B_1 \times B_2$ whose crystal structure is given as follows:

$$
\tilde{e}_i(b_1 \otimes b_2) = \begin{cases}\n\tilde{e}_i b_1 \otimes b_2 & \text{if } \varphi_i(b_1) \ge \varepsilon_i(b_2), \\
b_1 \otimes \tilde{e}_i b_2 & \text{if } \varphi_i(b_1) < \varepsilon_i(b_2),\n\end{cases}
$$
\n
$$
\tilde{f}_i(b_1 \otimes b_2) = \begin{cases}\n\tilde{f}_i b_1 \otimes b_2 & \text{if } \varphi_i(b_1) > \varepsilon_i(b_2), \\
b_1 \otimes \tilde{f}_i b_2 & \text{if } \varphi_i(b_1) \le \varepsilon_i(b_2),\n\end{cases}
$$
\n
$$
\text{wt}(b_1 \otimes b_2) = \text{wt}(b_1) + \text{wt}(b_2),
$$
\n
$$
\varepsilon_i(b_1 \otimes b_2) = \max(\varepsilon_i(b_1), \varepsilon_i(b_2) - \langle h_i, \text{wt}(b_1) \rangle),
$$
\n
$$
\varphi_i(b_1 \otimes b_2) = \max(\varphi_i(b_2), \varphi_i(b_1) + \langle h_i, \text{wt}(b_2) \rangle).
$$

Let B be a classical crystal. For an element $b \in B$, we define

$$
\varepsilon(b) = \sum_{i \in I} \varepsilon_i(b) \Lambda_i, \quad \varphi(b) = \sum_{i \in I} \varphi_i(b) \Lambda_i.
$$

DEFINITION 2.3. Let l be a positive integer. A classical crystal B is called a perfect crystal of level l if

- (1) there exists a finite dimensional $U_q'(\mathfrak{g})$ -module with a crystal basis whose crystal graph is isomorphic to B,
- (2) $B \otimes B$ is connected,
- (3) there exists a classical weight $\lambda_0 \in \bar{P}$ such that $\text{wt}(B) \subset \lambda_0 + \sum$ $i\neq 0$ $\mathbb{Z}_{\leq 0} \alpha_i,$
	- $#(B_{\lambda_0})=1$, where $B_{\lambda_0}=\{b\in B\mid \text{ wt}(b)=\lambda_0\},$
- (4) for any $b \in B$, $\langle c, \varepsilon(b) \rangle \geq l$,

(5) for any $\lambda \in \bar{P}^+$ with $\lambda(c) = l$, there exist unique $b^{\lambda}, b_{\lambda} \in B$ such that $\varepsilon(b^{\lambda}) = \lambda = \varphi(b_{\lambda}).$

The following crystal isomorphism theorem plays a fundamental role in the theory of perfect crystals.

THEOREM 2.4. [[3](#page-9-5)] Let B be a perfect crystal of level l $(l \in \mathbb{Z}_{\geq 0})$. For any $\lambda \in \bar{P}^+$ with $\lambda(c) = l$, there exists a unique classical crystal isomorphism

 $\Psi: B(\lambda) \stackrel{\sim}{\longrightarrow} B(\varepsilon(b_{\lambda})) \otimes B$ given by $u_{\lambda} \longmapsto u_{\varepsilon(b_{\lambda})} \otimes b_{\lambda},$

where u_{λ} is the highest weight vector in $B(\lambda)$ and b_{λ} is the unique vector in B such that $\varphi(b_\lambda) = \lambda$.

Set $\lambda_0 = \lambda$, $\lambda_{k+1} = \varepsilon(b_{\lambda_k})$, $b_0 = b_{\lambda_0}$, $b_{k+1} = b_{\lambda_{k+1}}$. Applying the above crystal isomorphism repeatedly, we get a sequence of crystal isomorphisms

$$
B(\lambda) \longrightarrow B(\lambda_1) \otimes B \longrightarrow B(\lambda_2) \otimes B \otimes B \longrightarrow \cdots
$$

\n $u_{\lambda} \longmapsto u_{\lambda_1} \otimes b_0 \longmapsto u_{\lambda_2} \otimes b_1 \otimes b_0 \longmapsto \cdots$

In this process, we get an infinite sequence $\mathbf{p}_{\lambda} = (b_k)_{k=0}^{\infty} \in B^{\otimes \infty}$, which is called the ground-state path of weight λ . Let $\mathcal{P}(\lambda) := \{ \mathbf{p} = (p(k))_{k=0}^{\infty} \in B^{\otimes \infty} \mid p(k) \in$ $B, p(k) = b_k$ for all $k \gg 0$. The elements of $\mathcal{P}(\lambda)$ are called the λ -paths. The following result gives the *path realization* of $B(\lambda)$.

PROPOSITION 2.5. [[3](#page-9-5)] There exists an isomorphism of classical crystals

$$
\Psi_{\lambda}: B(\lambda) \xrightarrow{\sim} \mathcal{P}(\lambda)
$$
 given by $u_{\lambda} \mapsto \mathbf{p}_{\lambda}$,

where u_{λ} is the highest weight vector in $\mathcal{B}(\lambda)$.

3. $U_q(D_4^{(3)})$ - Perfect crystals

In this section we recall the perfect crystal $B^{1,l}$ for the quantum affine algebra $U_q(D_4^{(3)})$ of level $l > 0$ constructed in [[7](#page-9-11)].

First we fix the data for $D_4^{(3)}$. Let $\{\alpha_0, \alpha_1, \alpha_2\}$, $\{h_0, h_1, h_2\}$ and $\{\Lambda_0, \Lambda_1, \Lambda_2\}$ be the set of simple roots, simple coroots and fundamental weights, respectively. The Cartan matrix $A = (a_{ij})_{i,j=0,1,2}$ is given by

$$
A = \left(\begin{array}{rrr} 2 & -1 & 0 \\ -1 & 2 & -3 \\ 0 & -1 & 2 \end{array} \right),
$$

and its Dynkin diagram is given by:

$$
\circ_0 - \circ_1 \Leftarrow \circ_2
$$

The standard null root δ and the canonical central element c are given by

 $\delta = \alpha_0 + 2\alpha_1 + \alpha_2$ and $c = h_0 + 2h_1 + 3h_2$,

where $\alpha_0 = 2\Lambda_0 - \Lambda_1 + \delta$, $\alpha_1 = -\Lambda_0 + 2\Lambda_1 - \Lambda_2$, $\alpha_2 = -3\Lambda_1 + 2\Lambda_2$. For positive integer l define the set

$$
B = B^{1,l} = \left\{ b = (x_1, x_2, x_3, \bar{x}_3, \bar{x}_2, \bar{x}_1) \in (\mathbb{Z}_{\geq 0})^6 \middle| \begin{array}{l} x_3 \equiv \bar{x}_3 \; (\text{mod } 2), \\ \sum_{i=1,2} (x_i + \bar{x}_i) + \frac{x_3 + \bar{x}_3}{2} \leq l \end{array} \right\}.
$$

For $b = (x_1, x_2, x_3, \bar{x}_3, \bar{x}_2, \bar{x}_1) \in B$ we denote

(3.1)
$$
s(b) = x_1 + x_2 + \frac{x_3 + \bar{x}_3}{2} + \bar{x}_2 + \bar{x}_1, t(b) = x_2 + \frac{x_3 + \bar{x}_3}{2},
$$

(3.2)
$$
z_1 = \bar{x}_1 - x_1
$$
, $z_2 = \bar{x}_2 - \bar{x}_3$, $z_3 = x_3 - x_2$, $z_4 = \frac{\bar{x}_3 - x_3}{2}$,

and

$$
(3.3) \quad \mathcal{A} = (0, z_1, z_1 + z_2, z_1 + z_2 + 3z_4, z_1 + z_2 + z_3 + 3z_4, 2z_1 + z_2 + z_3 + 3z_4).
$$

Now we define conditions (E_1) - (E_6) and (F_1) - (F_6) as follows.

$$
(3.4)
$$
\n
$$
\begin{cases}\n(F_1) & z_1 + z_2 + z_3 + 3z_4 \le 0, z_1 + z_2 + 3z_4 \le 0, z_1 + z_2 \le 0, z_1 \le 0, \\
(F_2) & z_1 + z_2 + z_3 + 3z_4 \le 0, z_2 + 3z_4 \le 0, z_2 \le 0, z_1 > 0, \\
(F_3) & z_1 + z_3 + 3z_4 \le 0, z_3 + 3z_4 \le 0, z_4 \le 0, z_2 > 0, z_1 + z_2 > 0, \\
(F_4) & z_1 + z_2 + 3z_4 > 0, z_2 + 3z_4 > 0, z_4 > 0, z_3 \le 0, z_1 + z_3 \le 0, \\
(F_5) & z_1 + z_2 + z_3 + 3z_4 > 0, z_3 + 3z_4 > 0, z_3 > 0, z_1 \le 0, \\
(F_6) & z_1 + z_2 + z_3 + 3z_4 > 0, z_1 + z_3 + 3z_4 > 0, z_1 + z_3 > 0, z_1 > 0.\n\end{cases}
$$

 (E_i) $(1 \le i \le 6)$ is defined from (F_i) by replacing > (resp. \le) with \ge (resp. \lt). Then for $b = (x_1, x_2, x_3, \bar{x}_3, \bar{x}_2, \bar{x}_1) \in B$, we define $\tilde{e}_i(b), \tilde{f}_i(b), \varepsilon_i(b), \varphi_i(b)$, $i = 0, 1, 2$ as follows. We use the convention: $(a)_+ = \max(a, 0)$.

$$
\tilde{e}_0(b) = \begin{cases}\n(x_1 - 1, \ldots) & \text{if } (E_1), \\
(\ldots, x_3 - 1, \bar{x}_3 - 1, \ldots, \bar{x}_1 + 1) & \text{if } (E_2), \\
(\ldots, x_3 - 2, \ldots, \bar{x}_2 + 1, \ldots) & \text{if } (E_3), \\
(\ldots, x_2 - 1, \ldots, \bar{x}_3 + 2, \ldots) & \text{if } (E_4), \\
(x_1 - 1, \ldots, x_3 + 1, \bar{x}_3 + 1, \ldots) & \text{if } (E_5), \\
(\ldots, \bar{x}_1 + 1) & \text{if } (E_6), \\
(\ldots, x_3 + 1, \bar{x}_3 + 1, \ldots, \bar{x}_1 - 1) & \text{if } (F_1), \\
(\ldots, x_3 + 1, \bar{x}_3 + 1, \ldots, \bar{x}_1 - 1) & \text{if } (F_2), \\
(\ldots, x_3 + 2, \ldots, \bar{x}_2 - 1, \ldots) & \text{if } (F_3), \\
(\ldots, x_2 + 1, \ldots, \bar{x}_3 - 2, \ldots) & \text{if } (F_4), \\
(x_1 + 1, \ldots, x_3 - 1, \bar{x}_3 - 1, \ldots) & \text{if } (F_5), \\
(\ldots, \bar{x}_1 - 1) & \text{if } (F_6),\n\end{cases}
$$

$$
\tilde{e}_1(b) = \begin{cases}\n(\ldots, \bar{x}_2 + 1, \bar{x}_1 - 1) & \text{if } z_2 \ge (-z_3)_+, \\
(\ldots, x_3 + 1, \bar{x}_3 - 1, \ldots) & \text{if } z_2 < 0 \le z_3, \\
(x_1 + 1, x_2 - 1, \ldots) & \text{if } (z_2)_+ < -z_3, \\
(x_1 - 1, x_2 + 1, \ldots) & \text{if } (z_2)_+ \le -z_3, \\
(\ldots, x_3 - 1, \bar{x}_3 + 1, \ldots) & \text{if } z_2 \le 0 < z_3, \\
(\ldots, \bar{x}_2 - 1, \bar{x}_1 + 1) & \text{if } z_2 > (-z_3)_+, \\
\tilde{e}_2(b) = \begin{cases}\n(\ldots, \bar{x}_3 + 2, \bar{x}_2 - 1, \ldots) & \text{if } z_4 \ge 0, \\
(\ldots, x_2 + 1, x_3 - 2, \ldots) & \text{if } z_4 < 0, \\
(\ldots, x_2 - 1, x_3 + 2, \ldots) & \text{if } z_4 \le 0, \\
(\ldots, \bar{x}_3 - 2, \bar{x}_2 + 1, \ldots) & \text{if } z_4 > 0.\n\end{cases}
$$

$$
\varepsilon_1(b) = \bar{x}_1 + (\bar{x}_3 - \bar{x}_2 + (x_2 - x_3)_{+})_+, \quad \varphi_1(b) = x_1 + (x_3 - x_2 + (\bar{x}_2 - \bar{x}_3)_{+})_+,
$$

\n
$$
\varepsilon_2(b) = \bar{x}_2 + \frac{1}{2}(x_3 - \bar{x}_3)_{+}, \quad \varphi_2(b) = x_2 + \frac{1}{2}(\bar{x}_3 - x_3)_{+},
$$

\n
$$
\varepsilon_0(b) = l - s(b) + \max \mathcal{A} - (2z_1 + z_2 + z_3 + 3z_4), \quad \varphi_0(b) = l - s(b) + \max \mathcal{A}.
$$

For $b \in B$ if $\tilde{e}_i(b)$ or $\tilde{f}_i(b)$ does not belong to B, namely, if x_j or \bar{x}_j for some j becomes negative or $s(b)$ exceeds l, we understand it to be 0.

The following is one of the main results in [[7](#page-9-11)]:

THEOREM 3.1. [[7](#page-9-11)] For the quantum affine algebra $U_q(D_4^{(3)})$ the set $B = B^{1,l}$ equipped with the maps \tilde{e}_i , \tilde{f}_i , ϵ_i , φ_i , $i = 0, 1, 2$ is a perfect crystal of level l.

As was shown in [[7](#page-9-11)], the minimal elements are given by:

 $(B)_{\min} = \{(\alpha, \beta, \beta, \beta, \beta, \alpha) | \alpha, \beta \in \mathbb{Z}_{\geq 0}, 2\alpha + 3\beta \leq l\}.$

4. Demazure Modules and Demazure crystals

Let W denote the Weyl group for the affine Lie algebra $\mathfrak g$ generated by the simple reflections $\{r_i | i \in I\}$, where $r_i(\mu) = \mu - \mu(h_i)\alpha_i$ for all $\mu \in \mathfrak{h}^*$. For $w \in W$ let $l(w)$ denote the length of w and \prec denote the Bruhat order on W. Let $U_q^+(\mathfrak{g})$ be the subalgebra of $U_q(\mathfrak{g})$ generated by the e_i 's. For $\lambda \in P^+$ with $\lambda(d) = 0$, consider the irreducible integrable highest weight $U_q(\mathfrak{g})$ -module $V(\lambda)$ with highest weight λ and highest weight vector u_λ . Let $(L(\lambda), B(\lambda))$ denote the crystal basis for $V(\lambda)$ ([[6](#page-9-2)],[[8](#page-9-3)]). It is known that for $w \in W$, the extremal weight space $V(\lambda)_{w\lambda}$ is one dimensional. Let $V_w(\lambda)$ denote the $U_q^+(\mathfrak{g})$ -module generated by $V(\lambda)_{w\lambda}$. These modules $V_w(\lambda)$ ($w \in W$) are called the *Demazure modules*. They are finite dimensional subspaces of $V(\lambda)$ and satisfy the properties: $V(\lambda) = \bigcup_{w \in W} V_w(\lambda)$ and for $w, w' \in W$ with $w \preceq w'$ we have $V_w(\lambda) \subseteq V_{w'}(\lambda)$.

In 1[9](#page-9-6)93, Kashiwara [9] showed that for each $w \in W$, there exists a subset $B_w(\lambda)$ of $B(\lambda)$ such that

$$
\frac{V_w(\lambda) \cap L(\lambda)}{V_w(\lambda) \cap qL(\lambda)} = \bigoplus_{b \in B_w(\lambda)} \mathbb{Q}b.
$$

The set $B_w(\lambda)$ is the crystal for the Demazure module $V_w(\lambda)$. The Demazure crystal $B_w(\lambda)$ has the following recursive property:

(4.1) If
$$
r_i w \succ w
$$
, then $B_{r_i w}(\lambda) = \bigcup_{n \geq 0} \tilde{f}_i^n B_w(\lambda) \setminus \{0\}.$

Suppose $\lambda(c) = l \geq 1$, and suppose B be a perfect crystal of level l for $U_q(\mathfrak{g})$. Then the crystal $B(\lambda)$ is isomorphic to the set of paths $\mathcal{P}(\lambda) = \mathcal{P}(\lambda, B)$ (see Propo-sition [2.5\)](#page-3-0). Under this isomorphism the highest weight element $u_{\lambda} \in B(\lambda)$ is identified with the ground state path $\mathbf{p}_{\lambda} = (\cdots \otimes b_3 \otimes b_2 \otimes b_1)$. We recall the path realizations of the Demazure crystals $B_w(\lambda)$ from [[10](#page-9-7)]. Fix positive integers d and κ. For a sequence of integers $\left\{ i_a^{(j)} \middle| j \geq 1, 1 \leq a \leq d \right\} \subset \{0, 1, \ldots, n\}$, we define the subsets $\left\{ B_a^{(j)} \, \middle| \, j \ge 1, 0 \le a \le d \right\}$ of B by

$$
B_0^{(j)} = \{b_j\}, B_a^{(j)} = \bigcup_{k \geq 0} \tilde{f}_{i_a^{(j)}}^k B_{a-1}^{(j)} \setminus \{0\}.
$$

Next we define $B_a^{(j+1,j)}, j \geq 1, 0 \leq a \leq d$ by

$$
B_0^{(j+1,j)}=B_0^{(j+1)}\otimes B_d^{(j)}, B_a^{(j+1,j)}=\bigcup_{k\geq 0} \tilde{f}_{i_a^{(j+1)}}^k B_{a-1}^{(j+1,j)}\backslash\left\{0\right\},
$$

and continue until we define

$$
B_0^{(j+\kappa-1,...,j)} = B_0^{(j+\kappa-1)} \otimes B_d^{(j+\kappa-2,...,j)}
$$

\n
$$
B_a^{(j+\kappa-1,...,j)} = \bigcup_{k \ge 0} \tilde{f}_{i_a^{(j+\kappa-1)}}^k B_{a-1}^{(j+\kappa-1,...,j)} \setminus \{0\}.
$$

Furthermore, we define a sequence $\{w^{(k)}\}$ of elements in the Weyl group W by

$$
w^{(0)} = 1, w^{(k)} = r_{i_a^{(j)}} w^{(k-1)}, \text{ for } k > 0.
$$

Here, j and a are uniquely determined from k by the relation $k = (j-1)d + a, j \geq$ $1, 1 \le a \le d$. Now for $k \ge 0$, we define subsets $\mathcal{P}^{(k)}(\lambda, B)$ of $\mathcal{P}(\lambda, B)$ as follows:

$$
\mathcal{P}^{(0)}(\lambda,B) = {\mathbf{p}}_{\lambda} \},
$$

and for $k > 0$,

$$
\mathcal{P}^{(k)}(\lambda, B) = \begin{cases} \cdots \otimes B_0^{(j+2)} \otimes B_0^{(j+1)} \otimes B_a^{(j,\ldots,1)} & \text{if } j < \kappa \\ \cdots \otimes B_0^{(j+2)} \otimes B_0^{(j+1)} \otimes B_a^{(j,\ldots,j-\kappa+1)} \otimes B^{\otimes (j-\kappa)} & \text{if } j \ge \kappa \end{cases}
$$

The following theorem shows that under certain conditions, the path realizations of the Demazure crystals $B_{w^{(k)}}(\lambda)$ is isomorphic to $\mathcal{P}^{(k)}(\lambda, B)$ and hence have tensor product-like structures.

THEOREM 4.1. [[10](#page-9-7)] Let $\lambda \in \bar{P}^+$ with $\lambda(c) = l$ and B be a perfect crystal of level l for the quantum affine Lie algebra $U_q(\mathfrak{g})$. For fixed positive integers d and κ , suppose we have a sequence of integers $\left\{ i_a^{(j)} | j \geq 1, 1 \leq a \leq d \right\} \subset \{0, 1, 2, \ldots, n\}$ satisfying the conditions:

- (1) for any $j \ge 1$, $B_d^{(j+\kappa-1,...,j)} = B_d^{(j+\kappa-1,...,j+1)} \otimes B$,
- (2) for any $j \ge 1, 1 \le a \le d$, $\langle \lambda_j, h_{i_a(j)} \rangle \le \varepsilon_{i_a(j)}(b), b \in B_{a-1}^{(j)}$, and
- (3) the sequence of elements $\left\{w^{(k)}\right\}_{k\geq 0}$ is an increasing sequence with respect to the Bruhat order.

.

Then we have $B_{w^{(k)}}(\lambda) \cong \mathcal{P}^{(k)}(\lambda, B)$.

The positive integer κ in Theorem [4.1](#page-6-0) is called the *mixing index*. It is conjectured that for any affine Lie algebra \mathfrak{g} , the mixing index $\kappa \leq 2$. It is known that the mixing index κ is dependent on the choice of the perfect crystal (see [[15](#page-9-13)]). For $\lambda = l\Lambda$, (where Λ is a dominant weight of level one) and the perfect crystal $B = B^{1,l}$ it is known that there exists a suitable sequence of Weyl group elements $\{w^{(k)}\}\$ which satisfy the conditions in Theorem [4.1](#page-6-0) with $\kappa = 1$ for g any classical quantum affine Lie algebra [[11](#page-9-8)], $U_q(D_4^{(3)})$ [[14](#page-9-9)], and $U_q(G_2^{(1)})$ $U_q(G_2^{(1)})$ $U_q(G_2^{(1)})$ [1].

For $b \in B$, let $\tilde{f}_i^{\max}(b)$ denote $\tilde{f}_i^{\varphi_i(b)}(b)$. For $j \geq 1$, we set

$$
b_0^{(j)} = b_j
$$
, $b_a^{(j)} = \tilde{f}_{i_a^{(j)}}^{\max}(b_{a-1}^{(j)})$ $(a = 1, 2, \cdots, d)$.

The following Proposition (10) (10) (10) , Proposition 2) will be useful to check the validity of condition (3) in Theorem [4.1.](#page-6-0)

PROPOSITION 4.2. [[10](#page-9-7)] For $w \in W$, if $\langle w\mu, h_i \rangle > 0$ for some $\mu \in \bar{P}^+$, then $r_jw \succ w$.

5. $U_q(D_4^{(3)})\text{-}\mathbf{Denazure}$ crystals

In this section we show that for the perfect crystal of level $3l$ for the quantum affine algebra $U_q(D_4^{(3)})$ given in Section 3, there is a suitable sequence of Weyl group elements $\{w^{(k)}\}$ which satisfy the conditions $(1), (2)$ and (3) for $\lambda = l\Lambda_2$ and hence Theorem [4.1](#page-6-0) holds in this case with $\kappa = 1$. Thus we have path realizations of the corresponding Demazure crystals with tensor product-like structures.

For $\lambda = l\Lambda_2, l \geq 1$, the $l\Lambda_2$ -minimal element in the perfect crystal $B = B^{1,3l}$ is $\bar{b} = (0, l, l, l, l, 0)$ and in this case $\lambda_j = \lambda = l\Lambda_2$ for $j \geq 1$. Hence $b_j = \bar{b}$ for all $j \geq 1$. Set $d = 6$ and choose the sequence $\{i_a^{(j)} | j \geq 1, 1 \leq a \leq 6\}$ defined by

(5.1)
$$
i_1^{(j)} = i_3^{(j)} = 2, i_2^{(j)} = i_4^{(j)} = i_6^{(j)} = 1, i_5^{(j)} = 0.
$$

Hence, by the action of \tilde{f}_i on B we have, for $j \geq 1$

(5.2)
\n
$$
b_0^{(j)} = (0, l, l, l, l, 0),
$$
\n
$$
b_1^{(j)} = \tilde{f}_2^{\max}(b_0^{(j)}) = (0, 0, 3l, l, l, 0),
$$
\n
$$
b_2^{(j)} = \tilde{f}_1^{\max}(b_1^{(j)}) = (0, 0, 0, 4l, l, 0),
$$
\n
$$
b_3^{(j)} = \tilde{f}_2^{\max}(b_2^{(j)}) = (0, 0, 0, 0, 3l, 0),
$$
\n
$$
b_4^{(j)} = \tilde{f}_1^{\max}(b_3^{(j)}) = (0, 0, 0, 0, 0, 3l),
$$
\n
$$
b_5^{(j)} = \tilde{f}_0^{\max}(b_4^{(j)}) = (3l, 0, 0, 0, 0, 0),
$$
\n
$$
b_6^{(j)} = \tilde{f}_1^{\max}(b_5^{(j)}) = (0, 3l, 0, 0, 0, 0),
$$

We define conditions P and $Q_j, 1 \leq j \leq 6$ for $b \in B$ as follows:

$$
\begin{cases}\n(P): z_3 \geq 0; z_3 + 3z_4 \geq (-2z_2)_+; z_1 + z_2 + z_3 + 3z_4 \geq 0; t(b) < 2l; s(b) < 3l \\
(Q_1): z_3 < 0; z_4 \geq 0; z_1 + z_2 + 3z_4 \geq 0; z_1 + 2z_2 + z_3 + 3z_4 \geq 0; t(b) \leq 2l; s(b) \leq 3l \\
(Q_2): z_2 \geq 0; z_4 < 0; z_3 + 3z_4 < 0; z_1 + z_2 \geq 0; z_1 + 2z_2 + z_3 \geq 0; s(b) \leq 3l \\
(Q_3): z_2 \geq 0; z_4 < 0; z_3 + 3z_4 < 0; z_1 + z_2 < 0; z_2 + z_3 \geq 0; s(b) \leq 3l \\
(Q_4): z_2 \geq 0; z_3 \geq 0; z_1 + z_2 + z_3 + 3z_4 < 0; z_3 + 3z_4 \geq 0; t(b) < 2l; s(b) \leq 3l \\
(Q_5): x_1 > 0; z_3 \geq 0; z_3 + 3z_4 \geq 0; z_1 + z_2 + z_3 + 3z_4 \geq 0; z_1 + 2z_2 + z_3 + 3z_4 \geq 0; \\
t(b) < 2l; s(b) \leq 3l \\
(Q_6): x_1 > 0; z_3 < 0; z_4 \geq 0; z_1 + z_2 + 3z_4 < 0; z_2 + 3z_4 > 0; z_2 + z_3 \geq 0; t(b) < 2l; s(b) \leq 3l\n\end{cases}
$$

By direct calculations it can be seen that the subsets $\left\{ B_a^{(j)} \middle| j \ge 1, 0 \le a \le 6 \right\}$ of B are given as follows.

$$
B_0^{(j)} = \{(0, l, l, l, l, 0)\}
$$

\n
$$
B_1^{(j)} = B_0^{(j)} \cup \{(0, x_2, x_3, l, l, 0) \mid z_3 > 0, s(b) = 3l\}
$$

\n
$$
B_2^{(j)} = B_1^{(j)} \cup \{(0, x_2, x_3, \bar{x}_3, l, 0) \mid z_2 < 0, z_3 \ge 0, s(b) = 3l\}
$$

\n
$$
B_3^{(j)} = B_2^{(j)} \cup \{(0, x_2, x_3, \bar{x}_3, \bar{x}_2, 0) \mid z_3 \ge 0, z_3 + 3z_4 \ge 0, t(b) < 2l, s(b) = 3l\}
$$

\n
$$
B_4^{(j)} = B_3^{(j)} \cup \{(0, x_2, x_3, \bar{x}_3, \bar{x}_2, \bar{x}_1) \mid \bar{x}_1 > 0, z_3 \ge 0, z_3 + 3z_4 \ge (-2z_2)_+, t(b) < 2l, s(b) = 3l\}
$$

\n
$$
B_5^{(j)} = B_4^{(j)} \cup C \cup D_1 \cup D_2 \cup \dots D_6
$$

\n
$$
B_6^{(j)} = B,
$$

where $C = \{(0, x_2, x_3, \bar{x}_3, \bar{x}_2, \bar{x}_1) | (P) \text{ holds}\}\$ and for $1 \leq j \leq 6$, $D_j = \{(x_1, x_2, x_3, \bar{x}_3, \bar{x}_2, \bar{x}_1) \mid (Q_j) \text{ holds}\}.$

The following lemma is useful and follows by easy calculations.

LEMMA 5.1. Let
$$
k \in \mathbb{Z}_{>0}
$$
 and $k = 6(j - 1) + a, j \ge 1, 1 \le a \le 6$. Then we have
\n
$$
w^{(k)} \Lambda_2 = \Lambda_2 - m_0 \alpha_0 - m_1 \alpha_1 - m_2 \alpha_2
$$
 where
\n
$$
m_0 = \begin{cases} 3j^2 + 3j & \text{if } a = 1, 2, 3, 4 \\ 3j^2 + 9j + 6 & \text{if } a = 5, 6 \end{cases}
$$
\n
$$
m_1 = \begin{cases} 6j^2 + 3j, & \text{if } a = 1 \\ 6j^2 + 9j + 3, & \text{if } a = 2, 3 \\ 6j^2 + 12j + 6, & \text{if } a = 4, 5 \\ 3j^2 + 15j + 9, & \text{if } a = 6 \end{cases}
$$
\n
$$
m_2 = \begin{cases} 3j^2 + 3j + 1, & \text{if } a = 1, 2 \\ 3j^2 + 6j + 3, & \text{if } a = 3, 4, 5, 6 \end{cases}
$$

THEOREM 5.2. For $\lambda = l\Lambda_2$, $l \geq 1$ and the given perfect crystal $B = B^{1,3l}$ for the quantum affine algebra $U_q(D_4^{(3)})$ with $d=6$ and the sequence $\{i_a^{(j)}\}$ given in [\(5.1\)](#page-7-0), conditions (1), (2) and (3) in Theorem [4.1](#page-6-1) hold with $\kappa = 1$. Hence we have path realizations of the corresponding Demazure crystals $B_{w^{(k)}}(l\Lambda_2)$ for $U_q(D_4^{(3)})$ with tensor product-like structures.

PROOF. We have already shown above by explicit descriptions of the subsets $B_a^{(j)}$ that $B_6^{(j)} = B$ which implies that condition (1) in Theorem [4.1](#page-6-1) holds for $\kappa = 1$.

Observe that $\langle l\Lambda_2, h_{i_a^{(j)}} \rangle = 0 \leq \varepsilon_{i_a^{(j)}}(b)$ for all $b \in B_{a-1}^{(j)}, a = 2, 4, 5, 6$. Also $\langle l\Lambda_2, h_{i_a^{(j)}}\rangle = l$ for $a = 1,3$. Observe that for all $b = (x_1, x_2, x_3, \bar{x}_3, \bar{x}_2, \bar{x}_1) \in$ $B_0^{(j)}$ or $B_2^{(j)}$ we have $\bar{x}_2 = l$. Hence $\varepsilon_{i_a^{(j)}}(b) = \bar{x}_2 + \frac{1}{2}(x_3 - \bar{x}_3)_+ = l + \frac{1}{2}(x_3 - \bar{x}_3)_+ \ge l$ for $a = 1, 3$ and condition (2) holds.

To prove condition (3), we use Lemma [5.1](#page-8-0) to obtain:

$$
\Big\langle w^{(k)} \Lambda_2, h_{i_{a+1}^{(j)}} \Big\rangle \ = \begin{cases} 3j+2 & \text{if } a=2 \\ 3j+3 & \text{if } a=3,5 \\ 3j+4 & \text{if } a=6 \\ 6j+3 & \text{if } a=1, \\ 6j+6 & \text{if } a=4, \end{cases}
$$

where $k = 6(j-1) + a, j \ge 1$. Hence $\langle w^{(k)} \Lambda_2, h_{i_{a+1}^{(j)}} \rangle > 0$ for all $j \ge 1$. Therefore, by Proposition [4.2,](#page-7-1) $w^{(k+1)} = r_{i_{a+1}^{(j)}} w^{(k)} \succ w^{(k)}$, which implies that condition (3) holds.

 \Box

References

- [1] R. L. Jayne, K. C. Misra, On Demazure crystals for $U_q(G_2^{(1)})$, Proc. Amer. Math. Soc. 139(7) (2011), 2343 - 2356.
- [2] V. G. Kac, Infinite-dimensional Lie algebras, Third edition, Cambridge, 1990.
- [3] S-J. Kang, M. Kashiwara, K. C. Misra, T. Miwa, T. Nakashima, A. Nakayashiki, Affine crystals and vertex models, Int. J. Mod. Phys. 7 (suppl. 1A) (1992), 449-484.
- [4] S-J. Kang, M. Kashiwara, K. C. Misra, T. Miwa, T. Nakashima, A. Nakayashiki, Perfect crystals of quantum affine Lie algebras, Duke Math. J. 68 (1992), 499-607.
- [5] S-J. Kang, M. Kashiwara, K. C. Misra, Crystal bases of Verma modules for quantum affine Lie Algebras, Compositio Math. 92 (1994), 299-325.
- [6] M. Kashiwara, Crystalizing the q-analogue of universal enveloping algebras, Commun. Math. Phys. 133 (1990), 249-260.
- [7] M. Kashiwara, K. C. Misra, M. Okado and D. Yamada, Perfect crystals for $U_q(D_4^{(3)})$, J. Algebra, 317, no.1, (2007), 392-423.
- [8] M. Kashiwara, On crystal bases of the q-analogue of universal enveloping algebras, Duke Math. J. 63 (1991), 465-516.
- [9] M. Kashiwara, The crystal base and Littelmann's refined Demazure character formula, Duke Math. J. **71** (1993), 839-858.
- [10] A. Kuniba, K. C. Misra, M. Okado and J. Uchiyama, Demazure modules and perfect crystals, Commun. Math. Phys. 192 (1998), 555-567.
- [11] A. Kuniba, K. C. Misra, M. Okado, T. Takagi and J. Uchiyama, Crystals for Demazure Modules of Classical Affine Lie Algebras, J. Algebra 208 (1998), 185-215.
- [12] G. Lusztig, Canonical bases arising from quantized enveloping algebras, J. Amer. Math. Soc. 3 (1990), 447-498.
- [13] G. Lusztig, Introduction to quantum groups, Birkhäuser Boston Inc., 1993.
- [14] K. C. Misra, On Demazure crystals for $U_q(D_4^{(3)})$, Contemporary Math. **392** (2005), 83-93.
- [15] K. C. Misra, V. Williams, Combinatorics of quantum affine Lie algebra representations, Contemporary Math. 343 (2004), 167-189.

Department of Mathematics, North Carolina State University, Raleigh, NC 27695- 8205

E-mail address: amarmstr@ncsu.edu

Department of Mathematics, North Carolina State University, Raleigh, NC 27695- 8205

E-mail address: misra@ncsu.edu