Unraveling quantum mechanical effects in water using isotopic fractionation

Thomas E. Markland * and B. J. Berne †

*Department of Chemistry, Stanford University, Stanford, California 94305-5080, and †Department of Chemistry, Columbia University, New York, New York 10027 Submitted to Proceedings of the National Academy of Sciences of the United States of America

When two phases of water are at equilibrium, the ratio of hydrogen isotopes in each is slightly altered due to their different phase affinities. This isotopic fractionation process can be utilized to analyze water's movement in the world's climate. Here we show that equilibrium fractionation ratios, an entirely quantum mechanical property, also provide a sensitive probe to assess the magnitude of nuclear quantum fluctuations in water. By comparing the predictions of a series of water models, we show that those describing the OH chemical bond as rigid or harmonic greatly over-predict the magnitude of isotope fractionation. Models that account for anharmonicity in this coordinate are shown to provide much more accurate results due to their ability to give partial cancellation between inter and intra-molecular quantum effects. These results give evidence of the existence of competing quantum effects in water and allow us to identify how this cancellation varies across a wide range of temperatures. In addition, this work demonstrates that simulation can provide accurate predictions and insights into hydrogen fractionation.

water | quantum effect | isotope | fractionation

Water within earth's atmosphere is naturally composed of the stable hydrogen isotopes Hydrogen (H) and Deuterium (D). During cycles of evaporation, condensation and precipitation, these isotopes naturally undergo partial separation due to their differing masses thereby leading to different H/D ratios in the two phases. This process of fractionation has a number of fortuitous consequences which are utilized in hydrology and geology. For instance, by comparing the ratio of H to D, one can estimate the origins of a water sample, the temperature at which it was formed, and the altitude at which precipitation occurred [1, 2]. Equilibrium fractionation, where the two phases are allowed to equilibrate their H/D ratio, is entirely a consequence of the effects of quantum mechanical fluctuations on water's hydrogen bond network. Quantum mechanical effects such as zero-point energy and tunneling are larger for H due to its lower

Despite numerous studies, the extent to which quantum fluctuations affect water's structure and dynamics remains a subject of considerable debate. It has long been appreciated that one effect of quantum fluctuations in water is the disruption of hydrogen bonding, leading to de-structuring of the liquid and faster dynamics [3, 4, 5, 6]. However, more recent work has suggested that a competing quantum effect may exist in water [7, 8], namely that the quantum kinetic energy in the OH covalent bond allows it to stretch and form shorter and stronger hydrogen bonds, which partially cancels the disruptive effect. This hydrogen bond strengthening has only been recently appreciated, as many original studies drew their conclusions based on models with rigid or harmonic bonds which are unable to describe this behavior. The degree of quantum effect cancellation depends sensitively on the anharmonicity of the OH stretch and the temperature. These parameters tune the balance between the lower frequency hydrogen bonding disruption, which will dominate at lower temperatures, and the higher frequency hydrogen bond strengthening effect, which will dominate at higher temperatures when rotations become essentially classical.

If such a large degree of cancellation existed at ambient temperature, it would be highly fortuitous both in terms of the biological effects of heavy water, which is only mildy toxic to humans [9], as

well as the ability to use heavy solvents in 2D-IR and NMR spectroscopies, where deuteration is assumed not to dramatically alter the structure or dynamics observed. However, the size of this cancellation remains elusive since empirical quantum models of water are typically fit to reproduce its properties when used in path integral simulations and the two *ab initio* path integral studies performed have not produced a consistent picture [7, 10]. In addition, many of these simulation studies compare the properties of water to those of its classical counterpart, but "classical" water is physically unrealizable even at relatively high temperatures, since water still has significant quantum effects present in its vibrations.

In this paper, we use equilibrium fractionation ratios as a sensitive probe to assess the magnitude of quantum mechanical effects in water. Fractionation ratios can be directly related to quantum kinetic energy differences between H and D in liquid water and its vapor and can be calculated exactly for a given water potential energy model using path integral simulations. The large number of accurate experimental measurements of these ratios allows for sensitive comparisons of theory and experiment over a wide range of temperatures [11]. In the present work, we show what features are needed in a water model to accurately predict these ratios by decomposing the contributions to the free energy difference leading to fractionation. This in turn leads to a simple explanation of the inversion of the fractionation ratios seen experimentally at high temperatures, where D is favored over H in the vapor phase [11].

Calculating Fractionation Ratios

The liquid-vapor fractionation ratio, α_{l-v} , is defined as

$$\alpha_{l-v} = \frac{(x_{D,l}/x_{H,l})}{(x_{D,v}/x_{H,v})} = e^{-\Delta A/k_B T},$$
 [1]

where x_Z is the mole fraction of isotope Z, l denotes the liquid phase, and v denotes the vapor phase. In the second equality, ΔA is the Helmholtz free energy corresponding to the process

$$H_2O_{(l)} + HOD_{(v)} \rightleftharpoons H_2O_{(v)} + HOD_{(l)}.$$
 [2]

In this work we consider the dilute deuterium (D) limit which reflects the situation found in the Earth's atmosphere where it is 6000 times less common than H. In this limit, we consider the free energy of exchanging a single D atom in a vapor water molecule with an H atom in a liquid water molecule, with all other molecules being H₂O. The

Reserved for Publication Footnotes

free energy difference can be calculated from the thermodynamic integration expression [12]

$$\Delta A = \int_{m_H}^{m_D} dm_Z \left(\frac{\langle K_l(m_Z) \rangle - \langle K_v(m_Z) \rangle}{m_Z} \right), \quad [3]$$

where $\langle K_l(m_Z) \rangle$ and $\langle K_v(m_Z) \rangle$ are the kinetic energy expectation values for a hydrogen isotope of mass m_Z in a water molecule HOZ in the liquid and vapor phases respectively. The kinetic energy can be calculated exactly for a given potential energy model of water using a path integral molecular dynamics (PIMD) simulation. These simulations exploit the exact isomorphism between a system of quantum mechanical particles and that of a set of classical ring polymers in which the spread of a polymer is directly related to that quantum particle's position uncertainty[13]. The kinetic energy for the particle Zin the molecule HOZ can be calculated from these simulations using the centroid virial estimator [14, 15]

$$K = \frac{3k_B T}{2} - \frac{1}{2n} \sum_{k=1}^{n} \Delta \mathbf{r}_Z^{(k)} \cdot \mathbf{f}_Z^{(k)},$$
 [4]

where T is the temperature and k_B is the Boltzmann constant. Here, $\Delta \mathbf{r}_Z^{(k)}$ is vector from the kth bead representing particle Z to the center of the ring polymer and $\mathbf{f}_Z^{(k)}$ is the force on that bead as shown in Fig. 1. The first term in this expression is the classical kinetic energy and is independent of the surrounding environment, thus it is identical in the vapor and liquid phase. The second term is the kinetic energy associated with confinement of a quantum particle. This confinement depends on the forces exerted on the particle by the surrounding molecules. Equilibrium fractionation is thus an entirely quantum mechanical phenomenon.

PIMD simulations were performed using 1000 water molecules with n=32 ring polymer beads used for the imaginary time discretization. Previously described evolution and thermostatting procedures were used [16]. The computational cost of these calculations was reduced by using the ring polymer contraction technique with a cutoff of 5A, which for this system size leads to more than an order of magnitude speed-up compared to a standard PIMD implementation [17, 18]. The integral in Eq. 3 was performed using the midpoint rule with 11 masses m_Z evenly spaced between m_H and m_D . Calculations were performed at the experimental coexistence densities [19].

To model the interactions within and between water molecules, we used the q-SPCFw [5] and q-TIP4P/F [8] models which have previously been shown to accurately reproduce many of water's properties in PIMD simulations of liquid water. Both models are flexible, use point charges, and have a harmonic description of the bending mode. However, while q-SPCFw uses a purely harmonic description of the OH stretch

$$V_{\rm OH}(r) = \frac{1}{2}k_r(r - r_{\rm eq})^2,$$
 [5]

q-TIP4P/F contains anharmonicity by modeling the stretch as a Morse expansion truncated at fourth order

$$V_{\rm OH}(r) = D_r \left[\alpha_r^2 (r - r_{\rm eq})^2 - \alpha_r^3 (r - r_{\rm eq})^3 + \frac{7}{12} \alpha_r^4 (r - r_{\rm eq})^4 \right].$$
 [6]

Here r is the distance between the oxygen and hydrogen atom and the parameters are given in Refs. [5] and [8]. The anharmonicity in the q-TIP4P/F model makes the observed quantum mechanical effects much smaller than previously predicted from harmonic or rigid models and gave rise to the idea of "competing quantum effects" in water[8].

Both models have previously been shown to accurately reproduce many of water's properties in PIMD simulations of liquid water. Due to their simple potential form such models are generally less transferable to other phases than more sophisticated polarizable or ab initio descriptions. However partially adiabatic centroid molecular dynamics simulations [20] have shown that the anharmonic stretch (Eq. 6) allows reasonable agreement to be obtained in the observed frequency shifts in the infrared spectrum in going from liquid to gaseous water as well as from pure light to pure heavy water [8, 21, 22]. These models were chosen to assess the importance of anharmonicity in the OH stretch using a "zeroth order" description of liquid water. Hence, as we show below, they offer a straightforward way to assess competing quantum effects in water.

Results and Discussion

Figure 2 shows the fractionation factors calculated from our PIMD simulations compared to the experimental data of Ref. [11]. For consistency with the experimental data, we plot $10^3 \ln \alpha_{l-v}$ which is simply $-10^3\Delta A/k_BT$. Since $\Delta A/k_BT$ is generally small, $e^{-\Delta A/k_BT}\simeq 1-\Delta A/k_BT$ thus an experimental value at 280 K of 100 corresponds to 10% more D residing in the liquid than the vapor. Above 500 K, the experimental data shows a well characterized region of inverse fractionation where D becomes more favored in the vapor than in the liquid, as shown in the inset of Fig. 2.

Turning to the simulated data, we observe that the harmonic q-SPC/Fw model over-predicts the magnitude of fractionation at 300 K by a factor of 3, and does not fall to the value observed experimentally at 300 K until the temperture is raised to 450 K. In contrast, the q-TIP4P/F model is in error by only 25% at the lowest temperature and approaches the experimental values more closely at higher temperatures. It also correctly shows inverse fractionation above 540 K. A previous study using the rigid SPC/E model and \hbar^2 perturbation theory, found a H/D fractionation of 450 at 300 K which is \sim 5 times higher than that seen experimentally [23]. However, it is not clear whether this was purely due to the use of a rigid model or whether the approximate \hbar^2 perturbation technique used to obtain the fractionation ratios was also at fault.

While Fig.2 demonstrates that the q-TIP4P/F model provides much better agreement with the experimental data than q-SPCFw, it is not immediately clear what aspect of the parameterization causes this. To better understand the origins of this effect, we constructed two models which we denote Aq-SPC/Fw and Hq-TIP4P/F. In the former, the q-SPC/Fw water model has its harmonic OH stretch replaced by a fourth order Morse expansion using the parameters of q-TIP4P/F; in the latter, the Morse potential of q-TIP4P/F is truncated at the harmonic term (see Eqs. 5 and 6). The anharmonic variant of q-SPC/Fw gives results as good as q-TIP4P/F, while the harmonic version of q-TIP4P/F fares as poorly as q-SPCFw. In other words, the accurate prediction of the fractionation ratios in liquid water is tied to the anharmonicity in the OH direction and is rather insensitive to the other parameters. This sheds light on a previous study where a sophisticated rigid polarizable model gave identical predictions for H/D fractionation to the simple fixed-charge rigid SPC/E model, i.e. varying the intermolecular potential alone does not give the flexibility required to accurately reproduce the experimental fractionation ratios

To determine the reason for the inversion observed in fractionation above 500 K in both experiment and the q-TIP4P/F model, we decompose the contributions to the fractionation ratio by noting that the quantum contribution to the kinetic energy in Eq. 4 is the dot product of two vectors. The overall kinetic energy is invariant to the coordinate system used to evaluate it, thus when the kinetic energy is calculated in the standard Cartesian basis all three components will average to the same number due to the isotropy of the liquid. To gain further insight, we instead use the internal coordinates of the water molecule and determine the contribution to $10^3 \ln \alpha_{l-v}$ arising from the OH bond vector, a vector in the plane of the molecule, and the vector perpendicular to the molecular plane as shown in Fig. 1. This is similar to the approach taken by Lin et. al. in the different context of investigating the proton momentum disitribution in ice [24]. The

results of this decomposition are shown in Table 1 for 300 K where D is experimentally seen to favor the liquid, and Table 2 for 620 K where D is experimentally seen to favor the vapor.

From Table 1, we see that all of the models have largely similar contributions from the two directions orthogonal to the OH bond and that both are positive and therefore favor the D excess in the liquid. The contribution perpendicular to the plane is noticeably larger than the contribution in the plane. As shown in Eq. 3, the values depend on the change in the quantum kinetic energy between H and D in the liquid and the vapor, which in turn is determined by how much the H or D atom's position uncertainty is restricted by interacting with the other water molecules in the liquid or vapor. Since in the vapor there is little confinement in plane orthogonal to the water, a larger contribution from that direction is expected; in the liquid, other molecules are present which restrict expansion in that coordinate.

In all cases the OH contribution is negative, indicating that there is less confinement in the position of the H atom in that direction in the liquid than in the vapor because the hydrogen atom participates in hydrogen bonding allowing the OH chemical bond to stretch more easily. However, comparing the anharmonic models (Aq-SPCFw and q-TIP4P/F) with their harmonic counterparts (q-SPCFw and Hq-TIP4P/F), we observe a 10 fold increase in the values arising from the OH contribution which gives rise to a larger cancellation of the positive contributions from the two orthogonal vectors. It is this cancellation that leads to the much better agreement with the experimental data at 300 K.

We now turn now to Table 2, which shows the contributions to the fractionation ratio at 620 K, a regime where experimentally D is preferred in the lighter phase. This is closer to the classical limit where the fractionation would be zero, thus, as expected, each component is reduced in magnitude compared to the lower temperature data in Table 1. However, the relative decrease in each component varies. The contributions arising from the in-plane and out-of-plane contributions orthogonal to OH decrease by a factor of 7-8 whereas the OH contribution falls by a factor of only \sim 4. For the anharmonic models, the negative OH contribution outweighs the positive components in the other two directions leading to an inversion of the fractionation compared to that seen at 300 K in agreement with the experimental observation of -2. The reason the OH component falls off more slowly is that this direction is dominated by stretching of the OH chemical bond, which is a high frequency coordinate, so even at high temperatures, quantum mechanics plays a noticeable role. In contrast, the two directions orthogonal to the OH direction are lower frequency and so approach the classical limit more rapidly as the temperature is increased. Thus, the reason the H/D fractionation is low around 600 K is not due to the fact that all contributions are individually low but rather that they nearly exactly cancel at this point due to the different rates at which the components approach the classical limit

Finally, we computed the fractionation ratio for the TTM3-F water model[25], which is known to have a very large cancellation of its quantum mechanical effects at 300 K [8, 26]. This model was fit to ab initio calculations using a potential form incorporating anharmonic flexibility, geometry dependent charges, and polarizability on the oxygen site. As such, it represents the current "gold standard" of parameterized water models and has been used extensively in recent studies probing the effects of quantum mechanical fluctuations on water [21, 27, 28]. To see if the large cancellation of quantum effects predicted by this model is consistent with experimental fractionation ratios, we calculated the value at 300 K which yielded a value of $10^3 \ln \alpha_{l-v} = -57$. This is in qualitative and quantitative disagreement with the experimental results, since it predicts D to be favored in the vapor at all temperatures. This model therefore overpredicts competing quantum effects in water and hence care should be taken concerning its predictions on the effects of quantum fluctuations on water's structure and dynamics. However, based on our discussion above it is likely that reparamaterization of the OH bond anharmonicity could correct this discrepancy.

Conclusion

In conclusion, we have shown that including anharmonicity in the OH bond when modeling water is essential to obtain agreement with the experimentally observed H/D fractionation ratios and that these ratios provide an excellent method to assess the accuracy of the quantum effects predicted by models of water. Since it has recently been shown that the competition between quantum mechanical effects applies to other hydrogen bonded systems [29], it is likely that many of our conclusions will be relevant to understanding isotopic fractionation in these systems. Additionally, while we only considered equilibrium fractionation in this work, which can be calculated exactly for a given potential energy model using PIMD simulations, many water processes occurring in the world's atmosphere are nonequilibrium ones. While including the effects of quantum fluctuations on dynamics is a much more challenging feat, the recent development of efficient condensed phase quantum dynamics approaches [30, 31, 32] should allow insights to be gained into kinetic fractionation processes. These directions will form the basis of future work.

ACKNOWLEDGMENTS. The authors gratefully thank Joseph Morrone and David Selassie for helpful comments and a critical reading of this manuscript. This research was supported by a grant to B.J.B. from the National Science Foundation (NSF-CHE-0910943).

- 1. Hoefs J (1997) Stable Isotope Geochemistry (Springer-Verlag, Berlin).
- Worden J, Noone D, Bowman K (2007) Importance of rain evaporation and continental convection in the tropical water cycle. Nature 445:528.
- Kuharski RA, Rossky PJ (1984) Quantum mechanical contributions to the structure of liquid water. Chem. Phys. Lett. 103:357–362.
- Wallqvist A, Berne BJ (1985) Path-integral simulation of pure water. Chem. Phys. Lett. 117:214–219.
- Paesani F, et al. (2006) An accurate and simple quantum model for liquid water. J. Chem. Phys. 125:184507–11.
- Miller TF, Manolopoulos DE (2005) Quantum diffusion in liquid water from ring polymer molecular dynamics. J. Chem. Phys. 123:154504.
- Chen B, Ivanov I, Klein ML, Parrinello M (2003) Hydrogen bonding in water. Phys. Rev. Lett. 91:215503.
- Habershon S, Markland TE, Manolopoulos DE (2009) Competing quantum effects in the dynamics of a flexible water model. J. Chem. Phys. 131:024501–11.
- Kushner D, Baker A, Dunstall T (1999) Pharmacological uses and perspectives of heavy water and deuterated compounds. Can. J. Physiol. Pharmacol. 77:79

 –88.
- Morrone JA, Car R (2008) Nuclear quantum effects in water. Phys. Rev. Lett. 101:017801.
- Horita J, Wesolowski DJ (1994) Liquid-vapor fractionation of oxygen and hydrogen isotopes of water from the freezing to the critical temperature. Geochimica et Cosmochimica Acta 58:3425–3427.

- Vanicek J, Miller WH (2007) Efficient estimators for quantum instanton evaluation
 of the kinetic isotope effects: Application to the intramolecular hydrogen transfer
 in pentadiene. J. Chem. Phys. 127:114309.
- Parrinello M, Rahman A (1984) Study of an f center in molten kcl. J. Chem. Phys. 80:860.
- Herman MF, Bruskin EJ, Berne BJ (1982) On path integral monte carlo simulations. J. Chem. Phys. 76:5150–5155.
- Cao J, Berne BJ (1989) On energy estimators in path integral Monte Carlo simulations: Dependency of accuracy on algorithm. J. Chem. Phys. 91:6359.
- Ceriotti M, Parrinello M, Markland TE, Manolopoulos DE (2010) Efficient stochastic thermostatting of path integral molecular dynamics. J. Chem. Phys. 133:124104.
- Markland TE, Manolopoulos DE (2008) An efficient ring polymer contraction scheme for imaginary time path integral simulations. J. Chem. Phys. 129:024105.
- Markland TE, Manolopoulos DE (2008) A refined ring polymer contraction scheme for systems with electrostatic interactions. Chem. Phys. Lett. 464:256 – 261.
- Wagner W, Pruβ A (2002) The lapws formulation 1995 for the thermodynamic properties of ordinary water substance for general and scientific use. J. Phys. Chem. Ref. Data 31:387.
- Hone TD, Rossky PJ, Voth GA (2006) A comparative study of imaginary time path integral based methods for quantum dynamics. The Journal of Chemical Physics 124:154103.

- Habershon S, Fanourgakis GS, Manolopoulos DE (2008) Comparison of path integral molecular dynamics methods for the infrared absorption spectrum of liquid water. J. Chem. Phys. 129:074501.
- Paesani F, Voth GA (2010) A quantitative assessment of the accuracy of centroid molecular dynamics for the calculation of the infrared spectrum of liquid water. The Journal of Chemical Physics 132:014105.
- Chialvo AA, Horita J (2009) Liquid-vapor equilibrium and isotopic fractionation of water: How well can classical water models predict it? J. Chem. Phys. 130:094509.
- Lin L, Morrone JA, Car R, Parrinello M (2011) Momentum distribution, vibrational dynamics, and the potential of mean force in ice. Phys. Rev. B 83:220302—.
- Fanourgakis GS, Xantheas SS (2008) Development of transferable interaction potentials for water. v. extension of the flexible, polarizable, thole-type model potential (ttm3-f, v. 3.0) to describe the vibrational spectra of water clusters and liquid water. J. Chem. Phys. 128:074506.
- Zeidler A, et al. (2011) Oxygen as a site specific probe of the structure of water and oxide materials. Phys. Rev. Lett. 107:145501.

- Paesani F (2011) Hydrogen bond dynamics in heavy water studied with quantum dynamical simulations. Phys. Chem. Chem. Phys. 13:19865.
- Liu J, et al. (2011) Insights in quantum dynamical effects in the infrared spectroscopy of liquid water from a semiclassical study with an ab initio-based flexible and polarizable force field. J. Chem. Phys. 135:244503.
- Li XZ, Walker B, Michaelides A (2011) Quantum nature of the hydrogen bond. Proc. Nat. Acad. Sci. USA 108:6369.
- Cao J, Voth (1994) The formulation of quantum statistical mechanics based on the feynman path centroid density. i. equilibrium properties. J, Chem. Phys. 100:5093.
- Wang H, Sun X, Miller WH (1998) Semiclassical approximations for the calculation of thermal rate constants for chemical reactions in complex molecular systems. *J. Chem. Phys.* 108:9726.
- Craig IR, Manolopoulos DE (2004) Quantum statistics and classical mechanics: Real time correlation functions from ring polymer molecular dynamics. J. Chem. Phys. 121:3368.

4 | www.pnas.org — — Footline Author




Fig. 1. Graphical representation of the vectors introduced in the text for a water molecule HOZ where Z is a hydrogen isotope of mass, m_Z . For clarity the H and O atoms are shown as single atoms whereas in the PIMD formalism they are also mapped onto ring polymers.




Fig. 2. Fractionation factors as a function of temperature for the q-SPCFw and q-TIP4P/F water models which describe the OH bond stretch using harmonic and anharmonic functions respectively. The experimental results are taken from Ref. [11]. The inset shows the high temperature region where both experiment and the q-TIP4P/F model show an inversion in the fractionation where D is preferentially found in the vapor phase. The curves through the calculated points are simply guides for the eye.

Table 1. Decomposition of the components leading to the fractionation ratio, $10^3 \ln \alpha_{l-v}$ at 300 K. OH is the contribution to $10^3 \ln \alpha_{l-v}$ arising from quantum kinetic energy down the OH direction, plane is that arising from the vector in the molecular plane orthogonal to the OH bond and Orth is the contribution down the vector orthogonal to the molecular plane. The type column denotes whether the water model for which the data was calculated uses a harmonic or anharmonic description of the OH bond.

Model	Stretch	ОН	Plane	Orth	Total
q-SPCFw	Harmonic	-17	92	141	216
Hq-TIP4P/F	Harmonic	-15	84	146	215
q-TIP4P/F	Anharmonic	-152	92	164	104
Aq-SPCFw	Anharmonic	-149	98	151	100

Table 2. Decomposition of the components leading to the fractionation ratio, $10^3 \ln \alpha_{l-v}$ at 620 K. The labeling is identical to that in Table 1.

Mod	lel	Stretch	ОН	Plane	Orth	Total
q-SP0	CFw	Harmonic	-3	13	20	30
Hq-TIP	4P/F	Harmonic	-3	11	20	28
q-TIP4	₽/F	Anharmonic	-40	13	22	-5
Aq-SP	CFw	Anharmonic	-40	13	21	-6

6 | www.pnas.org — Footline Author