Sharp constants in Riemannian L^p -Gagliardo-Nirenberg inequalities *[†]

Jurandir Ceccon[‡]; Carlos E. Durán[§]

Departamento de Matemática, Universidade Federal do Paraná,

Caixa Postal 019081, 81531-990, Curitiba, PR, Brazil

August 23, 2018

Abstract

Let (M, g) be a smooth compact Riemannian manifold of dimension $n \ge 2$, $1 and <math>1 \le q < r < p^* = \frac{np}{n-p}$ be real parameters. This paper concerns to the validity of the optimal Gagliardo-Nirenberg inequality

$$\left(\int_{M} |u|^{r} \, dv_{g}\right)^{\frac{\tau}{r\theta}} \leq \left(A_{opt}\left(\int_{M} |\nabla_{g}u|^{p} \, dv_{g}\right)^{\frac{\tau}{p}} + B_{opt}\left(\int_{M} |u|^{p} \, dv_{g}\right)^{\frac{\tau}{p}}\right) \left(\int_{M} |u|^{q} \, dv_{g}\right)^{\frac{\tau(1-\theta)}{\theta q}}$$

This kind of inequality is studied in Chen and Sun (Nonlinear Analysis 72 (2010), pp. 3159-3172) where the authors established its validity when $2 and (implicitly) <math>\tau = 1$. Here we solve the case $p \ge r$ and introduce one more parameter $1 \le \tau \le \min\{p, 2\}$. Moreover, we prove the existence of extremal function for the optimal inequality above.

1 Introduction

Inequalities of Gagliardo-Nirenberg type ([19] and [25]) contain by varying the parameters some classical inequalities, such as the Moser [23] and Nash [24] inequalities. Moreover, by taking limits of the parameters, we get logarithmic [15] and Sobolev [27] inequalities.

Optimal inequalities of Gagliardo-Nirenberg type have been extensively studied, both in the Euclidean and Riemannian contexts; e.g, [1], [5], [7], [13], [14] for the Euclidean case and [2], [3], [4], [6], [8], [9], [12], [17], [21] for Riemannian manifolds.

^{*2010} Mathematics Subject Classification: 58J05, 53C21, 35J20

[†]Key words: sharp Gagliardo-Nirenberg inequalities, extremal maps, best constant

[‡]E-mail addresses: ceccon@ufpr.br (J. Ceccon)

[§]E-mail addresses: cduran@ufpr.br (C. Durán)

The optimal cases of Euclidean Gagliardo-Nirenberg inequalities are used, for example, for finding sharp criteria for the global existence for nonlinear Schrödinger equations (see [11] or [26]), and optimal decay rate of the intermediate asymptotics of solutions to nonlinear diffusion equations (see [14]). Recently, the Riemannian Gagliardo-Nirenberg optimal constants studied in [8] where applied by [22] to obtain global existence theorems for Zakharov system in \mathbb{T}^2 . A particularly important family of applications of optimal Gagliardo-Nirenberg inequalities is the transition to optimal *Entropy inequalities*, in the spirit of [10, 15].

Denote by $D^{p,q}(\mathbb{R}^n)$ the completion of $C_0^{\infty}(\mathbb{R}^n)$ under the norm

$$||u||_{D^{p,q}(\mathbb{R}^n)} = \left(\int_{\mathbb{R}^n} |\nabla u|^p \, dx\right)^{\frac{1}{p}} + \left(\int_{\mathbb{R}^n} |u|^q \, dx\right)^{\frac{1}{q}}.$$

The Euclidean Gagliardo-Nirenberg inequality states that there exists A > 0, such that for any function $u \in D^{p,q}(\mathbb{R}^n)$,

$$\left(\int_{\mathbb{R}^n} |u|^r \, dx\right)^{\frac{p}{r\theta}} \le A\left(\int_{\mathbb{R}^n} |\nabla u|^p \, dx\right) \left(\int_{\mathbb{R}^n} |u|^q \, dx\right)^{\frac{p(1-\theta)}{\theta q}},\qquad (GN_E(A))$$

where $1 , <math>1 \le q < r < p^* = \frac{np}{n-p}$ and $\theta = \frac{np(r-q)}{r(q(p-n)+np)} \in (0,1)$ is the interpolation parameter. Define

$$A(p,q,r,n)^{-1} = \inf_{u \in D^{p,q}(\mathbb{R}^n)} \{ ||\nabla u||_{L^p(\mathbb{R}^n)}^p ||u||_{L^q(\mathbb{R}^n)}^{\frac{p(1-\theta)}{\theta}}; ||u||_{L^r(\mathbb{R}^n)} = 1 \}$$

Note that this constant is well defined since the right hand side infimum is a positive number by $GN_E(A)$. Inequality $GN_E(A(p,q,r,n))$ is called *optimal Euclidean Gagliardo-Nirenberg inequality* and the constant A(p,q,r,n) is the *best constant* in this inequality. A non-zero function realizing equality in $GN_E(A(p,q,r,n))$ is said to be an *extremal function*. The existence of such function is established in this case by using standard classical methods of Calculus of Variations.

We now consider the Riemannian case. Let (M, g) be a smooth, compact Riemannian manifold without boundary of dimension $n \ge 2$. Using standard arguments, we obtain a Riemannian version of the Euclidean inequality $GN_E(A)$: there exists positive constants C, D such that for all u in the Sobolev space $H^{1,p}(M)$, we have

$$\left(\int_{M} |u|^{r} dv_{g}\right)^{\frac{p}{r\theta}} \leq \left(C \int_{M} |\nabla_{g}u|^{p} dv_{g} + D \int_{M} |u|^{p} dv_{g}\right) \left(\int_{M} |u|^{q} dv_{g}\right)^{\frac{p(1-\theta)}{\theta_{q}}},\tag{1}$$

where $1 , <math>1 \le q < r < p^*$ and $\theta = \frac{np(r-q)}{r(q(p-n)+np)} \in (0,1)$ is the interpolation parameter.

It is a simple matter to add an additional parameter $\tau \ge 1$ in this inequality. We will work with

$$\left(\int_{M} |u|^{r} \, dv_{g}\right)^{\frac{\tau}{r\theta}} \leq \left(A\left(\int_{M} |\nabla_{g}u|^{p} \, dv_{g}\right)^{\frac{\tau}{p}} + B\left(\int_{M} |u|^{p} \, dv_{g}\right)^{\frac{\tau}{p}}\right) \left(\int_{M} |u|^{q} \, dv_{g}\right)^{\frac{\tau(1-\theta)}{\theta_{q}}}, \qquad (GN_{R}(A, B))$$

with p, q, r, θ as above. Note that when $\tau = p$ we recover (1).

The presence of this parameter represents the study of the Gagliardo-Niremberg inequalities for the family of equivalent norms in $W^{1,p}$ given by

$$\left(\left(\int_{M} |\nabla_{g} u|^{p} dv_{g}\right)^{\frac{\tau}{p}} + \left(\int_{M} |u|^{p} dv_{g}\right)^{\frac{\tau}{p}}\right)^{1/\tau}$$

A similar parameter was considered by Druet ([17]) in the context of Sobolev inequalities (his θ corresponds to our τ), in the process of solving a conjecture of Aubin (conjecture 2 of [2]).

Observe that the non-sharp inequality $GN_R(A, B)$, implies that

$$A \ge A(p,q,r,n)^{\frac{\tau}{p}} , \qquad (2)$$

for any $1 and <math>1 \le q < r < p^*$. This is shown by taking an appropriate localized test function, with support contained in a small enough normal neighbourhood so that the metric is almost Euclidean, compare [18].

We now study the optimal inequality. Having two constants, the optimality can be defined in two ways. We follow the more interesting one from the PDE viewpoint (see chapters 4 and 5 [20]): define the *first Riemannian* L^p -Gagliardo-Nirenberg best constant by

 $A_{opt} = \inf \{ A \in \mathbb{R} : \text{ there exists } B \in \mathbb{R} \text{ such that } GN_R(A, B) \text{ is valid} \}.$

This optimal constant is positive by (2). Moreover,

$$A_{opt}^{\frac{p}{\tau}} \ge A(p,q,r,n) , \qquad (3)$$

for any $1 and <math>1 \le q < r < p^*$. Then the first optimal Riemannain L^p -Gagliardo-Nirenberg inequality means that there exists a constant $B \in \mathbb{R}$ such that, for any $u \in H^{1,p}(M)$,

$$\left(\int_{M} |u|^{r} dv_{g}\right)^{\frac{\tau}{r\theta}} \leq \left(A_{opt}\left(\int_{M} |\nabla_{g}u|^{p} dv_{g}\right)^{\frac{\tau}{p}} + B\left(\int_{M} |u|^{p} dv_{g}\right)^{\frac{\tau}{p}}\right) \left(\int_{M} |u|^{q} dv_{g}\right)^{\frac{\tau(1-\theta)}{\theta q}},$$

is valid. In contrast with the Euclidean case, the validity of the optimal inequality is delicate since as $A \to A_{opt}$ the corresponding B might in principle go to infinity. In fact, when $\tau = p > 2$ there exists cases where the optimal inequality is not valid, depending on the geometry of (M, g), (see [8] and [16]).

Assuming that $GN_R(A_{opt}, B)$ holds, we can define the second Riemannian L^p -Gagliardo-Nirenberg best constant by

$$B_{opt} = \inf\{B \in \mathbb{R}; GN_R(A_{opt}, B) \ is \ valid\}.$$

Since constant non zero functions belong to $H^{1,p}(M)$, the constant B_{opt} satisfies

$$B_{opt} \ge |M|^{-\frac{\tau}{n}} , \tag{4}$$

where |M| denotes the volume of (M, g).

Then, by the optimal Riemannain L^p -Gagliardo-Nirenberg inequality we mean that for all $u \in H^{1,p}(M)$,

$$\left(\int_{M} |u|^{r} \, dv_{g}\right)^{\frac{\tau}{r\theta}} \leq \left(A_{opt}\left(\int_{M} |\nabla_{g}u|^{p} \, dv_{g}\right)^{\frac{\tau}{p}} + B_{opt}\left(\int_{M} |u|^{p} \, dv_{g}\right)^{\frac{\tau}{p}}\right) \left(\int_{M} |u|^{q} \, dv_{g}\right)^{\frac{\tau(1-\theta)}{\theta q}}$$

is valid. A non-zero function satisfying equality in $GN_R(A_{opt}, B_{opt})$ is called an *extremal function*.

We now state the main results of this paper:

Theorem 1. Let (M, g) be a smooth, compact Riemannian manifold without boundary of dimension $n \ge 2, 1 \le \tau \le p$ and $1 \le q < r \le p < n$. If $1 \le \tau \le \min\{p, 2\}$ then $GN_R(A_{opt}, B)$ is always valid for some B.

Theorem 1 complements the results of Chen and Sun [12]; for $\tau = 1$, they deal with the case p < r whereas we study the case $p \ge r$. The case p = r, Nash inequalities, is of particular interest since starting form Nash inequalities one can obtain entropy inequalities in the same spirit as in [10]; the classical case $\tau = p = r = 2$, q = 1 is treated by Humbert [21]; the conjecture of Aubin (conjeture 2 in [2]) is set up with $\tau = \frac{p}{p-1}$. The condition $1 \le \tau \le \min\{p, 2\}$ extends a similar condition for Sobolev inequalities present in Druet's solution [17] of Aubin's conjecture.

We remark that the arguments used in the proof are of *non-local* nature, in contrast with the local techniques used in [12]; the local arguments being inadequate when r < p. These non-local ideas allow the proof of an L^r concentration result (section 2.2) and a more refined pointwise estimate of certain maximizers (section 2.3). This refinement is essential for the case r < p.

Having theorem 1 allows the consideration of the second optimal constant; now by definition the optimal inequality $GN_R(A_{opt}, B_{opt})$ holds. We have

Theorem 2. Let (M, g) be a smooth, compact Riemannian manifold without boundary of dimension $n \ge 2, 1 \le \tau \le p$ and $1 \le q < r \le p < n$. If $1 \le \tau < 2$ then $GN_R(A_{opt}, B_{opt})$ admits an extremal function.

2 Proof of Theorem 1

We proceed by contradiction: assume inequality $GN_R(A_{opt}, B)$ is false for all B; this means that for any $\alpha > 0$ there exists $u \in H^{1,p}(M)$ such that

$$\left(\int_{M} |u|^{r} dv_{g}\right)^{\frac{\tau}{r\theta}} \left(\int_{M} |u|^{q} dv_{g}\right)^{-\frac{\tau(1-\theta)}{\theta q}} - \alpha \left(\int_{M} |u|^{p} dv_{g}\right)^{\frac{\tau}{p}} > A_{opt} \left(\int_{M} |\nabla_{g} u|^{p} dv_{g}\right)^{\frac{\tau}{p}}.$$
(5)

Consider the space $E = \{u \in H^{1,p}(M) : ||\nabla_g u||_{L^p(M)} = 1\}$. By suitably normalizing, we can assume the an u satisfying the previous inequality belongs to E, that is

$$\left(\int_{M} |u|^{r} \, dv_{g}\right)^{\frac{\tau}{r\theta}} \left(\int_{M} |u|^{q} \, dv_{g}\right)^{-\frac{\tau(1-\theta)}{\theta_{q}}} - \alpha \left(\int_{M} |u|^{p} \, dv_{g}\right)^{\frac{\tau}{p}} > A_{opt} \,. \tag{6}$$

Consider now the functional $J_{\alpha}: E \to \mathbb{R}$ given by the left-hand side of the preceeding inequality. We will show:

- 1. J_{α} always admit a maximizer \tilde{u}_{α} , that is, $J_{\alpha}(\tilde{u}_{\alpha}) = \sup_{u \in E} J_{\alpha} =: \nu_{\alpha} > A_{opt}$, which satisfies an elliptic PDE as Euler-Lagrange equation. It will be simpler to work with the normalization $u_{\alpha} = \tilde{u}_{\alpha}/||\tilde{u}_{\alpha}||_{L^{r}(M)}$, which, by homogeneity, satisfies inequality (5). This step is studied in section 2.1
- 2. Aided by the Euler-Lagrange equation, satisfied by u_{α} , we show that it is concentrated around its maximum x_0 in a sense to be made precise in section 2.2, and it also satisfies a pointwise estimate that quantifies the rate of decay of u_{α} in terms of the distance to x_0 ; this is done in section 2.3.
- 3. The previous item allows to localize the integrations in a small normal coordinate chart, and control the non-Euclidean terms in the Cartan expansion of the metric-volume-gradient etc. Then using the Euclidean Gagliardo-Nirenberg inequality will furnish the desired contradiction.

2.1 Maximizers and their Euler-Lagrange equations

For each $\alpha > 0$, consider the functional J_{α}

$$J_{\alpha}(u) = \left(\int_{M} |u|^{r} dv_{g}\right)^{\frac{\tau}{r\theta}} \left(\int_{M} |u|^{q} dv_{g}\right)^{-\frac{\tau(1-\theta)}{\theta q}} - \alpha \left(\int_{M} |u|^{p} dv_{g}\right)^{\frac{\tau}{p}}$$

defined on the space E, and

$$\nu_{\alpha} = \sup_{u \in E} J_{\alpha}(u) > A_{opt} \,. \tag{7}$$

Note that ν_{α} is well-defined and finite since there are constants A, B such that $GN_R(A, B)$ holds.

Assume first q > 1. Since J_{α} is of class C^1 , by using standard variational arguments, we find a maximizer $\tilde{u}_{\alpha} \in E$ of J_{α} , i.e.

$$J_{\alpha}(\tilde{u}_{\alpha}) = \nu_{\alpha} = \sup_{u \in E} J_{\alpha}(u) .$$
(8)

When q = 1, the functional J_{α} is not C^1 ; however in this case, following Humbert [21] we obtain the existence of extremal satisfying the corresponding Euler-Lagrange equation. From now on, the arguments are similar in the two cases q > 1 and q = 1. Thereby, we will focus our attention only on the case q > 1.

By (8), \tilde{u}_{α} satisfies the Euler-Lagrange equation

$$\frac{1}{\theta} ||\tilde{u}_{\alpha}||_{L^{r}(M)}^{\frac{\tau-r\theta}{\theta}} ||\tilde{u}_{\alpha}||_{L^{q}(M)}^{-\frac{\tau(1-\theta)}{\theta}} \tilde{u}_{\alpha}^{r-1} - \frac{(1-\theta)}{\theta} ||\tilde{u}_{\alpha}||_{L^{r}(M)}^{\frac{\tau}{\theta}} ||\tilde{u}_{\alpha}||_{L^{q}(M)}^{-\frac{\tau(1-\theta)+\theta q}{\theta}} \tilde{u}_{\alpha}^{q-1} - \alpha ||\tilde{u}_{\alpha}||_{L^{p}(M)}^{\tau-p} \tilde{u}_{\alpha}^{p-1} = \nu_{\alpha} \Delta_{p,g} \tilde{u}_{\alpha} , \quad (9)$$

where $\Delta_{p,g} = -\text{div}_g(|\nabla_g|^{p-2}\nabla_g)$ is the *p*-Laplace operator of *g*. Because $\nabla_g |\tilde{u}_{\alpha}| = \pm \nabla_g \tilde{u}_{\alpha}$, we can assume $\tilde{u}_{\alpha} \ge 0$. We now set $u_{\alpha} = \frac{\tilde{u}_{\alpha}}{||\tilde{u}_{\alpha}||_{L^r(M)}}$. Writing the Euler-Lagrange equation in terms of u_{α} , we have

$$\lambda_{\alpha}^{-1} A_{\alpha} \Delta_{p,g} u_{\alpha} + \alpha A_{\alpha}^{\frac{\tau}{p}} ||u_{\alpha}||_{L^{p}(M)}^{\tau-p} u_{\alpha}^{p-1} + \frac{1-\theta}{\theta} ||u_{\alpha}||_{L^{q}(M)}^{-q} u_{\alpha}^{q-1} = \frac{1}{\theta} u_{\alpha}^{r-1} \quad \text{on} \quad M ,$$
(10)

where $||u_{\alpha}||_{L^{r}(M)} = 1$,

$$A_{\alpha} = \left(\int_{M} u_{\alpha}^{q} \, dv_{g}\right)^{\frac{p(1-\theta)}{\theta_{q}}}$$

and

$$\lambda_{\alpha} = \nu_{\alpha}^{-1} ||u_{\alpha}||_{L^{q}(M)}^{\frac{(p-\tau)(1-\theta)}{\theta}} ||\tilde{u}_{\alpha}||_{L^{r}(M)}^{\tau-p}.$$

By Tolksdorf's regularity theory (see [28]), it follows that
$$u_{\alpha}$$
 is of class C^{1} .

We now highlight two important consequences of the Euler-Lagrange equation.

$$\lambda_{\alpha}^{-1} \ge A_{opt}^{\frac{p}{\tau}} , \qquad (11)$$

and

$$\lim_{\alpha \to \infty} A_{\alpha} = 0.$$
 (12)

In order to show (11), note that taking \tilde{u}_{α} as test function in (9), we have

$$\nu_{\alpha} \leq ||\tilde{u}_{\alpha}||_{L^{r}(M)}^{\frac{\tau}{\theta}} ||\tilde{u}_{\alpha}||_{L^{q}(M)}^{-\frac{\tau(1-\theta)}{\theta}}$$

Putting together the previous inequality, (7) and noting that $\tau \leq p$, we get

$$\lambda_{\alpha} = \nu_{\alpha}^{-1} ||\tilde{u}_{\alpha}||_{L^{r}(M)}^{\frac{(\tau-p)(1-\theta)}{\theta}} ||\tilde{u}_{\alpha}||_{L^{q}(M)}^{-\frac{(\tau-p)(1-\theta)}{\theta}} ||\tilde{u}_{\alpha}||_{L^{r}(M)}^{\tau-p} = \nu_{\alpha}^{-1} \left(||\tilde{u}_{\alpha}||_{L^{r}(M)}^{\frac{\tau}{\theta}} ||\tilde{u}_{\alpha}||_{L^{q}(M)}^{-\frac{\tau-p}{\tau}} \right)^{\frac{\tau-p}{\tau}} \leq \nu_{\alpha}^{-\frac{p}{\tau}} \leq A_{opt}^{-\frac{p}{\tau}} .$$

The limit (12) is shown as follows: first,

$$1 = \int_{M} u_{\alpha}^{r} dv_{g} \le |M|^{1 - \frac{r}{p}} \left(\int_{M} u_{\alpha}^{p} dv_{g} \right)^{\frac{r}{p}}$$

and thus we have $||u_{\alpha}||_{L^{p}(M)} > c > 0$ for all α . Using now (10), we obtain

$$\alpha A_{\alpha}^{\frac{\tau}{p}} \left(\int_{M} u_{\alpha}^{p} dv_{g} \right)^{\frac{\tau}{p}} \le c , \qquad (13)$$

which proves claim (12).

2.2 L^r -concentration

Let $x_{\alpha} \in M$ be a maximum point of u_{α} , that is,

$$u_{\alpha}(x_{\alpha}) = ||u_{\alpha}||_{L^{\infty}(M)}.$$

$$(14)$$

Throughout this section, we use the notation $\lim_{\sigma,\alpha\to\infty}$ to mean $\lim_{\sigma\to\infty}\lim_{\alpha\to\infty}$. Our aim here is to establish that

$$\lim_{\sigma, \alpha \to \infty} \int_{B(x_{\alpha}, \sigma a_{\alpha})} u_{\alpha}^{r} \, dv_{g} = 1 \,, \tag{15}$$

where

$$a_{\alpha} = A_{\alpha}^{\frac{r}{np-nr+pr}}.$$
(16)

Let us proceed. Since np - nr + pr > 0 and by (12), $a_{\alpha} \to 0$ as $\alpha \to \infty$.

In order to work in local coordinates, we pullback the metric and the function u_{α} to a ball $B(0,\sigma)$ in $T_{x_{\alpha}}M$. The pullbacks will be by exponential map precomposed by a dilation: if $\delta_L : TM \to TM$ denotes the dilation by L, $\delta_L(x) = Lx$ and $E_{\alpha} = \exp_{x_{\alpha}} \circ \delta_{a_{\alpha}}$, define

$$h_{\alpha} = E_{\alpha}^{*}g,$$

$$\varphi_{\alpha} = a_{\alpha}^{\frac{n}{r}}u_{\alpha} \circ E_{\alpha}.$$
(17)

Note that since $a_{\alpha} \to 0$ as $\alpha \to \infty$, these are well defined: for α large enough, $B(0, a_{\alpha}\sigma)$ will be contained inside of a set where $\exp_{x_{\alpha}}$ is a diffeomorphism.

By (10), one easily deduces that

$$\lambda_{\alpha}^{-1}\Delta_{p,h_{\alpha}}\varphi_{\alpha} + \alpha A_{\alpha}^{\frac{\tau}{p}-1} ||u_{\alpha}||_{L^{p}(M)}^{\tau-p} a_{\alpha}^{p}\varphi_{\alpha}^{p-1} + \frac{1-\theta}{\theta}\varphi_{\alpha}^{q-1} = \frac{1}{\theta}\varphi_{\alpha}^{r-1} \quad \text{on} \quad B(0,\sigma) \,.$$

$$\tag{18}$$

By (11) and applying the Moser's iterative scheme (see [23]) to this last equation, we see that, for α large enough,

$$a_{\alpha}^{n}||u_{\alpha}||_{L^{\infty}(M)}^{r} = \sup_{B(0,\frac{\sigma}{2})}\varphi_{\alpha}^{r} \le c \int_{B(0,\sigma)}\varphi_{\alpha}^{r} dh_{\alpha} = c \int_{B(x_{\alpha},\sigma a_{\alpha})}u_{\alpha}^{r} dv_{g} \le c$$

This estimate together with

$$1 = \int_{M} u_{\alpha}^{r} \, dv_{g} \le ||u_{\alpha}||_{L^{\infty}(M)}^{r-q} \int_{M} u_{\alpha}^{q} dv_{g} = \left(||u_{\alpha}||_{L^{\infty}(M)} \, a_{\alpha}^{\frac{n}{r}}\right)^{r-q}$$

gives

$$1 \le ||u_{\alpha}||_{L^{\infty}(M)} a_{\alpha}^{\frac{n}{r}} \le c.$$

$$\tag{19}$$

We will use in the sequel that (19) means that the limiting behaviour of $||u_{\alpha}||_{L^{\infty}(M)}$ and $a_{\alpha}^{-\frac{n}{r}}$ have the same order.

In particular, there exists a constant c > 0 such that

$$\int_{B(0,\sigma)} \varphi_{\alpha}^{r} \, dh_{\alpha} \ge c > 0 \tag{20}$$

for α large enough.

Now using Cartan expansion in normal coordinates and (19), we have for each $\sigma > 0$, that

$$\int_{B(0,\sigma)} \varphi_{\alpha}^{p} dx \leq c \int_{B(0,\sigma)} \varphi_{\alpha}^{p} dh_{\alpha} = c \ a_{\alpha}^{\frac{n}{r}(p-r)} \int_{B(x_{\alpha},\sigma a_{\alpha})} u_{\alpha}^{p} dv_{g} \leq c a_{\alpha}^{\frac{n}{r}(p-r)} ||u_{\alpha}||_{L^{\infty}(M)}^{p} \sigma^{n} a_{\alpha}^{n} \leq c(\sigma) ,$$

where $c(\sigma) \to \infty$ when $\sigma \to \infty$, but $c(\sigma)$ being independent of α . We have also

$$\int_{B(0,\sigma)} |\nabla \varphi_{\alpha}|^p \, dx \le c \int_{B(0,\sigma)} |\nabla_{h_{\alpha}} \varphi_{\alpha}|^p \, dh_{\alpha} = cA_{\alpha} \int_{B(x_{\alpha},\sigma a_{\alpha})} |\nabla_g u_{\alpha}|^p \, dv_g \le c A_{opt}^{-\frac{p}{\tau}}.$$

Therefore there exists $\varphi \in W^{1,p}(\mathbb{R}^n)$ such that, for some subsequence, $\varphi_{\alpha} \rightharpoonup \varphi$ in $W^{1,p}_{loc}(\mathbb{R}^n)$. For each $\sigma > 0$, we have

$$\int_{B(0,\sigma)} \varphi^q \, dx = \lim_{\alpha \to \infty} \int_{B(0,\sigma)} \varphi^q_\alpha \, dh_\alpha = \lim_{\alpha \to \infty} \frac{\int_{B(x_\alpha,\sigma a_\alpha)} u^q_\alpha \, dv_g}{\int_M u^q_\alpha \, dv_g} \le 1$$

and

$$\int_{B(0,\sigma)} \varphi^r \, dx = \lim_{\alpha \to \infty} \int_{B(0,\sigma)} \varphi^r_\alpha \, dh_\alpha = \lim_{\alpha \to \infty} \int_{B(x_\alpha,\sigma a_\alpha)} u^r_\alpha \, dv_g \le 1 \; .$$

In particular,

$$\varphi \in L^q(\mathbb{R}^n) \cap L^r(\mathbb{R}^n) .$$
(21)

Let $\eta \in C_0^1(\mathbb{R})$ be a cutoff function such that $\eta = 1$ on $[0, \frac{1}{2}]$, $\eta = 0$ on $[1, \infty)$ and $0 \le \eta \le 1$. Define $\eta_{\alpha,\sigma}(x) = \eta((\sigma a_\alpha)^{-1}d_g(x, x_\alpha))$. Choosing $u_\alpha \eta_{\alpha,\sigma}^p$ as a test function in (10), one gets

$$\lambda_{\alpha}^{-1} A_{\alpha} \int_{M} |\nabla_{g} u_{\alpha}|^{p} \eta_{\alpha,\sigma}^{p} dv_{g} + \lambda_{\alpha}^{-1} A_{\alpha} \int_{M} |\nabla_{g} u_{\alpha}|^{p-2} \nabla_{g} u_{\alpha} \cdot \nabla_{g} (\eta_{\alpha,\sigma}^{p}) u_{\alpha} dv_{g} + \frac{1-\theta}{\theta} \frac{\int_{M} u_{\alpha}^{q} \eta_{\alpha,\sigma}^{p} dv_{g}}{\int_{M} u_{\alpha}^{q} dv_{g}} \leq \frac{1}{\theta} \int_{M} u_{\alpha}^{r} \eta_{\alpha,\sigma}^{p} dv_{g}.$$

$$(22)$$

We now show that

$$\lim_{\sigma,\alpha\to\infty} A_{\alpha} \int_{M} |\nabla_{g} u_{\alpha}|^{p-2} \nabla_{g} u_{\alpha} \cdot \nabla_{g} (\eta^{p}_{\alpha,\sigma}) u_{\alpha} \, dv_{g} = 0.$$
⁽²³⁾

Taking u_{α} as test function, by (10) we have

$$A_{\alpha} \int_{M} |\nabla_{g} u_{\alpha}|^{p} dv_{g} \leq \lambda_{\alpha} \leq A_{opt}^{-\frac{p}{\tau}}$$

Therefore, it suffices to establish that

$$A_{\alpha} \int_{M} u_{\alpha}^{p} |\nabla_{g} \eta_{\alpha,\sigma}|^{p} \, dv_{g} \le \frac{c}{\sigma^{p}} \,. \tag{24}$$

Using (16) and (19), we derive

$$A_{\alpha} \int_{M} u_{\alpha}^{p} |\nabla_{g} \eta_{\alpha,\sigma}|^{p} dv_{g} \leq c \frac{A_{\alpha} ||u_{\alpha}||_{L^{\infty}(M)}^{p-r}}{\sigma^{p} a_{\alpha}^{p}} \int_{M} u_{\alpha}^{r} dv_{g} \leq c \frac{A_{\alpha} a_{\alpha}^{-\frac{n}{r}(p-r)}}{\sigma^{p} a_{\alpha}^{p}} \leq \frac{c}{\sigma^{p}}.$$

Therefore (24) holds and (23) is valid.

Replacing (11) and (23) in (22) and because λ_{α}^{-1} is limited, one arrives at

$$\theta \lim_{\sigma,\alpha\to\infty} \left(A_{opt}^{\frac{p}{\tau}} A_{\alpha} \int_{M} |\nabla_{g} u_{\alpha}|^{p} \eta_{\alpha,\sigma}^{p} dv_{g} \right) + (1-\theta) \lim_{\sigma,\alpha\to\infty} \frac{\int_{M} u_{\alpha}^{q} \eta_{\alpha,\sigma}^{p} dv_{g}}{\int_{M} u_{\alpha}^{q} dv_{g}} \leq \lim_{\sigma,\alpha\to\infty} \int_{M} u_{\alpha}^{r} \eta_{\alpha,\sigma}^{p} dv_{g}$$

To rewrite this inequality in a suitable format, we first remark that

$$\left|\frac{\int_{M} u_{\alpha}^{q} \eta_{\alpha,\sigma}^{p} \, dv_{g}}{\int_{M} u_{\alpha}^{q} \, dv_{g}} - \frac{\int_{M} u_{\alpha}^{q} \eta_{\alpha,\sigma}^{q} \, dv_{g}}{\int_{M} u_{\alpha}^{q} \, dv_{g}}\right| \leq \frac{\int_{B(x_{\alpha},\sigma a_{\alpha}) \setminus B(x_{\alpha},\sigma a_{\alpha}/2)} u_{\alpha}^{q} \, dv_{g}}{\int_{M} u_{\alpha}^{q} \, dv_{g}} = \int_{B(0,\sigma) \setminus B(0,\sigma/2)} \varphi_{\alpha}^{q} \, dh_{\alpha}$$

and the above right-hand side converges to 0 as $\sigma \to \infty$, by (21). So,

$$\lim_{\sigma,\alpha\to\infty} \frac{\int_M u^q_\alpha \eta^p_{\alpha,\sigma} \, dv_g}{\int_M u^q_\alpha \, dv_g} = \lim_{\sigma,\alpha\to\infty} \frac{\int_M u^q_\alpha \eta^q_{\alpha,\sigma} \, dv_g}{\int_M u^q_\alpha \, dv_g}$$

In a similar way,

$$\left| \int_{M} u_{\alpha}^{r} \eta_{\alpha,\sigma}^{p} \, dv_{g} - \int_{M} u_{\alpha}^{r} \eta_{\alpha,\sigma}^{r} \, dv_{g} \right| \leq \int_{B(x_{\alpha},\sigma a_{\alpha}) \setminus B(x_{\alpha},(\sigma a_{\alpha})/2)} u_{\alpha}^{r} \, dv_{g} = \int_{B(0,\sigma) \setminus B(0,\sigma/2)} \varphi_{\alpha}^{r} dh_{\alpha} ,$$

using (21), so that

$$\lim_{\sigma,\alpha\to\infty}\int_M u^r_\alpha\eta^r_{\alpha,\sigma}\ dv_g = \lim_{\sigma,\alpha\to\infty}\int_M u^r_\alpha\eta^p_{\alpha,\sigma}\ dv_g$$

Consequently, we can write

$$\theta \lim_{\sigma,\alpha\to\infty} \left(A_{opt}^{\frac{p}{\tau}} A_{\alpha} \int_{M} |\nabla_{g} u_{\alpha}|^{p} \eta_{\alpha,\sigma}^{p} dv_{g} \right) + (1-\theta) \lim_{\sigma,\alpha\to\infty} \frac{\int_{M} u_{\alpha}^{q} \eta_{\alpha,\sigma}^{q} dv_{g}}{\int_{M} u_{\alpha}^{q} dv_{g}} \le \lim_{\sigma,\alpha\to\infty} \int_{M} u_{\alpha}^{r} \eta_{\alpha,\sigma}^{r} dv_{g} .$$
(25)

On the other hand, for $\varepsilon > 0$ let the constant $B_{\varepsilon} > 0$, independent of α , such that

$$\left(\int_{M} u_{\alpha}^{r} \eta_{\alpha,\sigma}^{r} \, dv_{g}\right)^{\frac{\tau}{r\theta}} \leq \left((A_{opt} + \varepsilon) \left(\int_{M} |\nabla_{g}(u_{\alpha}\eta_{\alpha,\sigma})|^{p} \, dv_{g} \right)^{\frac{\tau}{p}} + B_{\varepsilon} \left(\int_{M} u_{\alpha}^{p} \eta_{\alpha,\sigma}^{p} \, dv_{g} \right)^{\frac{\tau}{p}} \right) \left(\int_{M} u_{\alpha}^{q} \eta_{\alpha,\sigma}^{q} \, dv_{g} \right)^{\frac{\tau(1-\theta)}{\theta_{q}}}$$

From the definition of A_{α} , Young inequality and $(x+y)^p \leq x^p + cx^{p-1}y + cy^p$ for $x, y \geq 0$, one has

$$\left(\int_{M} u_{\alpha}^{r} \eta_{\alpha,\sigma}^{r} dv_{g} \right)^{\frac{\tau}{r\theta}} \leq c \left(A_{\alpha} \int_{M} u_{\alpha}^{p} dv_{g} \right)^{\frac{\tau}{p}}$$

$$+ (A_{opt} + \varepsilon) \left(\left((1+\varepsilon) \int_{M} |\nabla_{g} u_{\alpha}|^{p} \eta_{\alpha,\sigma}^{p} dv_{g} + c(\varepsilon) \int_{M} u_{\alpha}^{p} |\nabla_{g} \eta_{\alpha,\sigma}|^{p} dv_{g} \right)^{\frac{\tau}{p}} \right) \left(\int_{M} u_{\alpha}^{q} \eta_{\alpha,\sigma}^{q} dv_{g} \right)^{\frac{\tau(1-\theta)}{\theta_{q}}} .$$

$$= inr (12) \quad (24) \quad \text{lattime a graph of a set of a generate.}$$

Then, using (13), (24), letting $\alpha, \sigma \to \infty$ and $\varepsilon \to 0$, one gets

$$\lim_{\sigma,\alpha\to\infty} \left(\int_M u^r_\alpha \eta^r_{\alpha,\sigma} \, dv_g \right)^{\frac{p}{r\theta}} \leq \lim_{\sigma,\alpha\to\infty} \left(A^{\frac{p}{r}}_{opt} A_\alpha \int_M |\nabla_g u_\alpha|^p \eta^p_{\alpha,\sigma} \, dv_g \right) \lim_{\sigma,\alpha\to\infty} \left(\frac{\int_M u^q_\alpha \eta^q_{\alpha,\sigma} \, dv_g}{\int_M u^q_\alpha \, dv_g} \right)^{\frac{p(1-\theta)}{\theta_q}} \,. \tag{26}$$

Let

$$X = \lim_{\sigma, \alpha \to \infty} \left(A_{opt}^{\frac{p}{\tau}} A_{\alpha} \int_{M} |\nabla_{g} u_{\alpha}|^{p} \eta_{\alpha, \sigma}^{p} dv_{g} \right), \quad Y = \lim_{\sigma, \alpha \to \infty} \frac{\int_{M} u_{\alpha}^{q} \eta_{\alpha, \sigma}^{q} dv_{g}}{\int_{M} u_{\alpha}^{q} dv_{g}}, \quad Z = \lim_{\sigma, \alpha \to \infty} \int_{M} u_{\alpha}^{r} \eta_{\alpha, \sigma}^{r} dv_{g}.$$

Clearly, $X, Y, Z \leq 1$ and (25) and (26) may be rewritten as

$$\begin{cases} \theta X + (1-\theta)Y \le Z\\ Z \le X^{\frac{r\theta}{p}}Y^{\frac{r(1-\theta)}{q}}. \end{cases}$$
(27)

By (20), one also has Z > 0, so that X, Y > 0.

Assertion (15) follows readily by proving that Z = 1. For this, we will consider the behavior of u_{α} outside $B(x_{\alpha}, \sigma a_{\alpha})$. Indeed, let $\zeta_{\alpha,\sigma} = 1 - \eta_{\alpha,\sigma}$ on *M*. Inequalities (25) and (26) remain valid for $\zeta_{\alpha,\sigma}$ in place of $\eta_{\alpha,\sigma}$. In other words,

$$\theta \lim_{\sigma,\alpha\to\infty} \left(A_{opt}^{\frac{p}{\tau}} A_{\alpha} \int_{M} |\nabla_{g} u_{\alpha}|^{p} \zeta_{\alpha,\sigma}^{p} dv_{g} \right) + (1-\theta) \lim_{\sigma,\alpha\to\infty} \frac{\int_{M} u_{\alpha}^{q} \zeta_{\alpha,\sigma}^{q} dv_{g}}{\int_{M} u_{\alpha}^{q} dv_{g}} \leq \lim_{\sigma,\alpha\to\infty} \int_{M} u_{\alpha}^{r} \zeta_{\alpha,\sigma}^{r} dv_{g}$$

and

$$\lim_{\sigma,\alpha\to\infty} \left(\int_M u_\alpha^r \zeta_{\alpha,\sigma}^r \, dv_g \right)^{\frac{p}{r\theta}} \le \lim_{\sigma,\alpha\to\infty} \left(A_{opt}^{\frac{p}{r}} A_\alpha \int_M |\nabla_g u_\alpha|^p \zeta_{\alpha,\sigma}^p \, dv_g \right) \lim_{\sigma,\alpha\to\infty} \left(\frac{\int_M u_\alpha^q \zeta_{\alpha,\sigma}^q \, dv_g}{\int_M u_\alpha^q \, dv_g} \right)^{\frac{p(1-\theta)}{\theta_q}}$$

ilar way, we denote

In a simi ay,

$$\tilde{X} = \lim_{\sigma, \alpha \to \infty} \left(A_{opt}^{\frac{p}{\tau}} A_{\alpha} \int_{M} |\nabla_{g} u_{\alpha}|^{p} \zeta_{\alpha, \sigma}^{p} dv_{g} \right), \quad \tilde{Y} = \lim_{\sigma, \alpha \to \infty} \frac{\int_{M} u_{\alpha}^{q} \zeta_{\alpha, \sigma}^{q} dv_{g}}{\int_{M} u_{\alpha}^{q} dv_{g}}, \quad \tilde{Z} = \lim_{\sigma, \alpha \to \infty} \int_{M} \zeta_{\alpha, \sigma}^{r} u_{\alpha}^{r} dv_{g},$$

so that

$$\begin{cases} \theta \tilde{X} + (1-\theta)\tilde{Y} \leq \tilde{Z} \\ \tilde{Z} \leq \tilde{X}^{\frac{r\theta}{p}} \tilde{Y}^{\frac{r(1-\theta)}{q}}. \end{cases}$$
(28)

.

We next assert that

$$Y + \tilde{Y} = 1 \quad \text{and} \quad Z + \tilde{Z} = 1.$$
⁽²⁹⁾

,

To justify the first equality, let us write

$$1 = \frac{\int_M u^q_\alpha \eta^q_{\alpha,\sigma} \, dv_g}{\int_M u^q_\alpha \, dv_g} + \frac{\int_M u^q_\alpha (1 - \eta^q_{\alpha,\sigma}) \, dv_g}{\int_M u^q_\alpha \, dv_g} \, .$$

Then,

$$\left|\frac{\int_{M} u_{\alpha}^{q}(1-\eta_{\alpha,\sigma}^{q}) \, dv_{g} - \int_{M} u_{\alpha}^{q} \zeta_{\alpha,\sigma}^{q} \, dv_{g}}{\int_{M} u_{\alpha}^{q} \, dv_{g}}\right| \leq \left|\frac{\int_{B(x_{\alpha},a_{\alpha}\sigma) \setminus B(x_{\alpha},a_{\alpha}\sigma/2)} u_{\alpha}^{q} dv_{g}}{\int_{M} u_{\alpha}^{q} dv_{g}}\right| = \int_{B(0,\sigma) \setminus B(0,\sigma/2)} \varphi_{\alpha}^{q} \, dh_{\alpha} \, .$$

Since $\varphi \in L^q(\mathbb{R}^n)$, we get

$$\lim_{\sigma,\alpha\to\infty}\frac{\int_M u^q_{\alpha}\zeta^q_{\alpha,\sigma}dv_g}{\int_M u^q_{\alpha}dv_g} = \lim_{\sigma,\alpha\to\infty}\frac{\int_M u^q_{\alpha}(1-\eta^q_{\alpha,\sigma})dv_g}{\int_M u^q_{\alpha}dv_g}$$

which yields $Y + \tilde{Y} = 1$. Analogously, since $\varphi \in L^r(\mathbb{R}^n)$,

$$1 = \int_M u_\alpha^r \eta_{\alpha,\sigma}^r \, dv_g + \int_M u_\alpha^r (1 - \eta_{\alpha,\sigma}^r) \, dv_g$$

and

$$\left| \int_{M} u_{\alpha}^{r} (1 - \eta_{\alpha,\sigma}^{r}) \, dv_{g} - \int_{M} u_{\alpha}^{r} \zeta_{\alpha,\sigma}^{r} \, dv_{g} \right| \leq \int_{B(x_{\alpha},a_{\alpha}\sigma) \setminus B(x_{\alpha},a_{\alpha}\sigma/2)} u_{\alpha}^{r} \, dv_{g} = \int_{B(0,\sigma) \setminus B(0,\sigma/2)} \varphi_{\alpha}^{r} \, dh_{\alpha}$$

which gives $Z + \tilde{Z} = 1$.

We are now ready to prove that Z = 1. The first inequality in (27) lead us to three possible alternatives:

(a) $X \leq Z$ and $Y \leq Z$, (b) $Y \leq Z \leq X$, (c) $X \leq Z \leq Y$.

If (a) holds, the second inequality in (27) implies

$$1 \le Z^{\frac{r\theta}{p} + \frac{r(1-\theta)}{q} - 1}.$$

But the definition of θ furnishes $\frac{r\theta}{p} + \frac{r(1-\theta)}{q} - 1 > 0$, so that Z = 1. Suppose then that the item (b) holds. Again, by (27),

$$Z \ge \theta X + (1-\theta)Y = \frac{r\theta}{p}X + (1-\frac{r\theta}{p})Y + \theta(1-\frac{r}{p})X + \theta(-1+\frac{r}{p})Y.$$

Using the assumption $p \ge r, Y \le X$ and Young's inequality, we derive

$$Z \ge X^{\frac{r\theta}{p}}Y^{\frac{p-r\theta}{p}} + \theta \frac{p-r}{p}(X-Y) \ge X^{\frac{r\theta}{p}}Y^{\frac{p-r\theta}{p}},$$

so that the second inequality in (27) immediately yields

$$1 \le Y^{\frac{r(1-\theta)}{q} - \frac{p-r\theta}{p}}$$

Since $\frac{r(1-\theta)}{q} - \frac{p-r\theta}{p} > 0$, one has Y = 1. Thus, evoking (28) and (29), one easily deduces that Z = 1. Finally, we come to the alternative (c). We first show that Y = 1. Otherwise, by (29), one has $\tilde{Y} > 0$ and this implies, by (28) and $\theta < 1$, that $\tilde{X} > 0$ and $\tilde{Z} > 0$. Since, by hypothesis, $Z \leq Y$, again thanks to (29), one concludes that $\tilde{Y} \leq \tilde{Z}$. Therefore, applying the previously discussed cases (a) and (b) with \tilde{X} , \tilde{Y} and \tilde{Z} in the place of X, Y and Z, one arrives at $\tilde{Z} = 1$. But this contradicts the fact that Z > 0, so that Y = 1. As before, (28) and (29) produce Z = 1 and this finishes the proof of the L^r -concentration.

2.3 Pointwise estimates

The aim of this section is to prove the following pointwise estimate of the decay of u_{α} in terms of the distance to its maximum:

For any constant $\lambda > 0$ there exists a constant $c_{\lambda} > 0$, independent of α , such that

$$d_g(x, x_\alpha)^\lambda u_\alpha(x) \le c_\lambda \ a_\alpha^{\lambda - \frac{n}{r}}$$

for all $x \in M$ and α large enough.

The proof proceeds by contradiction. Suppose that the assertion above is false. Then, there exist $\lambda_0 > 0$ and $y_\alpha \in M$ such that $f_\alpha(y_\alpha) \to \infty$ as $\alpha \to \infty$, where

$$f_{\alpha}(x) = d_g(x, x_{\alpha})^{\lambda_0} u_{\alpha}(x) \ a_{\alpha}^{-\lambda_0 + \frac{n}{r}} .$$

Assume, without loss of generality, that $f_{\alpha}(y_{\alpha}) = ||f_{\alpha}||_{L^{\infty}(M)}$. From (19), we have

$$f_{\alpha}(y_{\alpha}) \leq c \, \frac{u_{\alpha}(y_{\alpha})}{||u_{\alpha}||_{L^{\infty}(M)}} d_g(x_{\alpha}, y_{\alpha})^{\lambda_0} a_{\alpha}^{-\lambda_0} \leq c d_g(x_{\alpha}, y_{\alpha})^{\lambda_0} a_{\alpha}^{-\lambda_0} \,,$$

so that

$$d_g(x_\alpha, y_\alpha)a_\alpha^{-1} \to \infty.$$
(30)

For any fixed $\sigma > 0$ and $\varepsilon \in (0, 1)$, we next show that

$$B(y_{\alpha}, \varepsilon d_g(x_{\alpha}, y_{\alpha})) \cap B(x_{\alpha}, \sigma a_{\alpha}) = \emptyset$$
(31)

for α large enough. Clearly, this assertion follows from

$$d_g(x_\alpha, y_\alpha) \ge \sigma a_\alpha + \varepsilon d(x_\alpha, y_\alpha) \,.$$

But the above inequality is equivalent to

$$d_g(x_\alpha, y_\alpha)(1-\varepsilon)a_\alpha^{-1} \ge \sigma$$
,

which is clearly satisfied, since $d_g(x_\alpha, y_\alpha)a_\alpha^{-1} \to \infty$ and $1 - \varepsilon > 0$. We claim that exists a constant c > 0 such that

$$u_{\alpha}(x) \le c u_{\alpha}(y_{\alpha}) \tag{32}$$

for all $x \in B(y_{\alpha}, \varepsilon d_g(x_{\alpha}, y_{\alpha}))$ and α large enough. In fact, for each $x \in B(y_{\alpha}, \varepsilon d_g(x_{\alpha}, y_{\alpha}))$, we have

$$d_g(x, x_\alpha) \ge d_g(x_\alpha, y_\alpha) - d_g(x, y_\alpha) \ge (1 - \varepsilon) d_g(x_\alpha, y_\alpha).$$

Thus,

$$d_g(y_\alpha, x_\alpha)^{\lambda_0} u_\alpha(y_\alpha) a_\alpha^{-\lambda_0 + \frac{n}{r}} = f_\alpha(y_\alpha) \ge f_\alpha(x) = d_g(x, x_\alpha)^{\lambda_0} u_\alpha(x) a_\alpha^{-\lambda_0 + \frac{n}{r}}$$
$$\ge (1 - \varepsilon)^{\lambda_0} d_g(y_\alpha, x_\alpha)^{\lambda_0} u_\alpha(x) a_\alpha^{-\lambda_0 + \frac{n}{r}},$$

so that

$$u_{\alpha}(x) \le \left(\frac{1}{1-\varepsilon}\right)^{\lambda_0} u_{\alpha}(y_{\alpha})$$

for all $x \in B(y_{\alpha}, \varepsilon d_g(x_{\alpha}, y_{\alpha}))$ and α large enough. This proves our claim. Since $p \ge r$, by (16) and (19), one has

$$A_{\alpha}^{\frac{1}{p}}u_{\alpha}(y_{\alpha})^{\frac{p-r}{p}} \leq A_{\alpha}^{\frac{1}{p}}||u_{\alpha}||_{L^{\infty}(M)}^{\frac{p-r}{p}} \leq c \ a_{\alpha} \to 0.$$

So, in the same way as in (17), we can now define, inside of B(0,2) and for α large enough, $E_{\alpha} = \exp_{y_{\alpha}} \circ \delta_{A_{\alpha}^{\frac{1}{p}} u_{\alpha}(y_{\alpha})^{\frac{p-r}{p}}}$, and pullback

$$h_{\alpha} = E_{\alpha}^{\star} g$$

$$\psi_{\alpha} = u_{\alpha} (y_{\alpha})^{-1} u_{\alpha} \circ E_{\alpha}$$

From (10), it readily follows that

$$\lambda_{\alpha}^{-1} \Delta_{p,h_{\alpha}} \psi_{\alpha} + \alpha A_{\alpha}^{\frac{\tau}{p}} ||u_{\alpha}||_{L^{p}(M)}^{\tau-p} u_{\alpha}(y_{\alpha})^{p-r} \psi_{\alpha}^{p-1} + \frac{1-\theta}{\theta} ||u_{\alpha}||_{L^{q}(M)}^{-q} u_{\alpha}(y_{\alpha})^{q-r} \psi_{\alpha}^{q-1} = \frac{1}{\theta} \psi_{\alpha}^{r-1} \quad \text{on} \quad B(0,2) \,. \tag{33}$$

In particular,

$$\int_{B(0,2)} |\nabla_{h_{\alpha}}\psi_{\alpha}|^{p-2} \nabla_{h_{\alpha}}\psi_{\alpha} \cdot \nabla_{h_{\alpha}}\phi \, dv_{h_{\alpha}} \le c \int_{B(0,2)} \psi_{\alpha}^{r-1}\phi \, dv_{h_{\alpha}}$$

for all positive test function $\phi \in C_0^1(B(0,2))$. So, by the Moser's iterative scheme and (19), one deduces that

$$1 \leq \sup_{B(0,\frac{1}{4})} \psi_{\alpha}^{r} \leq c \int_{B(0,\frac{1}{2})} \psi_{\alpha}^{r} \, dv_{h_{\alpha}} = c \left(A_{\alpha}^{\frac{\theta_{q}}{p(1-\theta)}} u_{\alpha}(y_{\alpha})^{r-q} \right)^{-\frac{n(1-\theta)}{\theta_{q}}} \int_{B(y_{\alpha},\frac{1}{2}A_{\alpha}^{\frac{1}{p}} u_{\alpha}(y_{\alpha})^{\frac{p-r}{p}})} u_{\alpha}^{r} \, dv_{g}$$
$$\leq c \left(\frac{||u_{\alpha}||_{L^{\infty}(M)}}{u_{\alpha}(y_{\alpha})} \right)^{\frac{np-rn+pr}{p}} \int_{B(y_{\alpha},\frac{1}{2}A_{\alpha}^{\frac{1}{p}} u_{\alpha}(y_{\alpha})^{\frac{p-r}{p}})} u_{\alpha}^{r} \, dv_{g} \, .$$

For simplicity, we rewrite this last inequality as

$$0 < c \le m_{\alpha}^{\varrho} \int_{B(y_{\alpha}, \frac{1}{2}A_{\alpha}^{\frac{1}{p}} u_{\alpha}(y_{\alpha})^{\frac{p-r}{p}})} u_{\alpha}^{r} dv_{g}, \qquad (34)$$

where $m_{\alpha} = \frac{||u_{\alpha}||_{L^{\infty}(M)}}{|u_{\alpha}(y_{\alpha})}$ and $\varrho = \frac{np-rn+pr}{p} > 0$. By (16), (19) and (30), note that $B(y_{\alpha}, \frac{1}{2}A_{\alpha}^{\frac{1}{p}}u_{\alpha}(y_{\alpha})^{\frac{p-r}{p}}) \subset B(y_{\alpha}, \varepsilon d(x_{\alpha}, y_{\alpha}))$ for α large enough. So, the L^{r} concentration property (15) combined with (31) provide

$$\int_{B(y_{\alpha},\frac{1}{2}A_{\alpha}^{\frac{1}{p}}u_{\alpha}(y_{\alpha})^{\frac{p-r}{p}})} u_{\alpha}^{r} dv_{g} \to 0$$

when $\alpha \to \infty$.

So, we have that

 $\lim_{\alpha \to \infty} m_\alpha = \infty \; .$

Our goal now is to establish a contradiction to (34). Initially, from (19) and (32), we have

$$m_{\alpha}^{\varrho} \int_{D_{\alpha}} u_{\alpha}^{r} \, dv_{g} \le m_{\alpha}^{\varrho} ||u_{\alpha}||_{L^{\infty}(D_{\alpha})}^{r} (A_{\alpha}^{\frac{1}{p}} u_{\alpha}(y_{\alpha})^{\frac{p-r}{p}})^{n} \le c m_{\alpha}^{\varrho} u_{\alpha}(y_{\alpha})^{r} (A_{\alpha}^{\frac{1}{p}} u_{\alpha}(y_{\alpha})^{\frac{p-r}{p}})^{n} \le c \,, \tag{35}$$

where $D_{\alpha} = B(y_{\alpha}, A_{\alpha}^{\frac{1}{p}} u_{\alpha}(y_{\alpha})^{\frac{p-r}{p}})$. Consider the function $\eta_{\alpha}(x) = \eta(A_{\alpha}^{-\frac{1}{p}}d_g(x,y_{\alpha})u_{\alpha}(y_{\alpha})^{\frac{r-p}{p}})$, where $\eta \in C_0^1(\mathbb{R})$ is a cutoff function such that $\eta = 1$ on $[0,\frac{1}{2}], \eta = 0$ on $[1,\infty)$ and $0 \le \eta \le 1$. Taking $u_{\alpha}\eta^p_{\alpha}$ as a test function in (10), one has

$$\begin{split} \lambda_{\alpha}^{-1}A_{\alpha}\int_{M}|\nabla_{g}u_{\alpha}|^{p}\eta_{\alpha}^{p} \, dv_{g} + \lambda_{\alpha}^{-1}pA_{\alpha}\int_{M}|\nabla_{g}u_{\alpha}|^{p-2}u_{\alpha}\eta_{\alpha}^{p-1}\nabla_{g}u_{\alpha}\cdot\nabla_{g}\eta_{\alpha} \, dv_{g} + \alpha A_{\alpha}^{\frac{\tau}{p}}||u_{\alpha}||_{L^{p}(M)}^{\tau-p}\int_{M}u_{\alpha}^{p}\eta_{\alpha}^{p} \, dv_{g} \\ + \frac{1-\theta}{\theta}\frac{\int_{M}u_{\alpha}^{q}\eta_{\alpha}^{p}}{\int_{M}u_{\alpha}^{q}dv_{g}} \, dv_{g} = \frac{1}{\theta}\int_{M}u_{\alpha}^{r}\eta_{\alpha}^{p} \, dv_{g} \, . \end{split}$$

By Hölder and Young's inequalities,

$$\left|\int_{M} |\nabla_{g} u_{\alpha}|^{p-2} u_{\alpha} \eta_{\alpha}^{p-1} \nabla_{g} u_{\alpha} \cdot \nabla_{g} \eta_{\alpha} \, dv_{g}\right| \leq \varepsilon \int_{M} |\nabla_{g} u_{\alpha}|^{p} \eta_{\alpha}^{p} \, dv_{g} + c_{\varepsilon} \int_{M} |\nabla_{g} \eta_{\alpha}|^{p} u_{\alpha}^{p} \, dv_{g}$$

Also, by (19) and (32), it follows that

$$A_{\alpha} \int_{M} |\nabla_{g} \eta_{\alpha}|^{p} u_{\alpha}^{p} \, dv_{g} \leq A_{\alpha} (A_{\alpha}^{-\frac{1}{p}} u_{\alpha}(y_{\alpha})^{\frac{r-p}{p}})^{p} \int_{D_{\alpha}} u_{\alpha}^{p} \, dv_{g} \leq c u_{\alpha}(y_{\alpha})^{r} (A_{\alpha}^{\frac{1}{p}} u_{\alpha}(y_{\alpha})^{\frac{p-r}{p}})^{n} \leq c m_{\alpha}^{-\varrho} \,. \tag{36}$$

Consequently, combining these inequalities with (35), one arrives at

$$A_{\alpha} \int_{M} |\nabla_{g} u_{\alpha}|^{p} \eta_{\alpha}^{p} dv_{g} + c\alpha A_{\alpha}^{\frac{\tau}{p}} ||u_{\alpha}||_{L^{p}(M)}^{\tau-p} \int_{M} u_{\alpha}^{p} \eta_{\alpha}^{p} dv_{g} + c \frac{\int_{M} u_{\alpha}^{q} \eta_{\alpha}^{p} dv_{g}}{\int_{M} u_{\alpha}^{q} dv_{g}} \le cm_{\alpha}^{-\varrho}.$$

$$(37)$$

Now, since p > 1, the non-sharp Riemannian Gagliardo-Nirenberg inequality produces

$$\left(\int_{B(y_{\alpha},\frac{1}{2}A_{\alpha}^{\frac{1}{p}}u_{\alpha}(y_{\alpha})^{\frac{p-r}{p}})} u_{\alpha}^{r} dv_{g}\right)^{\frac{p}{r\theta}} \leq \left(\int_{M} (u_{\alpha}\eta_{\alpha}^{p})^{r} dv_{g}\right)^{\frac{p}{r\theta}} \leq c \left(\int_{M} |\nabla_{g}u_{\alpha}|^{p}\eta_{\alpha}^{p} dv_{g}\right) \left(\int_{M} (u_{\alpha}\eta_{\alpha}^{p})^{q} dv_{g}\right)^{\frac{p(1-\theta)}{\theta q}} + c \left(\int_{M} |\nabla_{g}u_{\alpha}|^{p}\eta_{\alpha}^{p} dv_{g}\right) \left(\int_{M} (u_{\alpha}\eta_{\alpha}^{p})^{q} dv_{g}\right)^{\frac{p(1-\theta)}{\theta q}} + c \left(\int_{M} (u_{\alpha}\eta_{\alpha}^{p})^{p} dv_{g}\right) \left(\int_{M} (u_{\alpha}\eta_{\alpha}^{p})^{q} dv_{g}\right)^{\frac{p(1-\theta)}{\theta q}}.$$
(38)

Thanks to (36), (37) and since $q \ge 1$, we then can estimate each term of the right-hand side of (38). Indeed, we have

$$\int_{M} |\nabla_{g} u_{\alpha}|^{p} \eta_{\alpha}^{p} \, dv_{g} \left(\int_{M} (u_{\alpha} \eta_{\alpha}^{p})^{q} \, dv_{g} \right)^{\frac{p(1-\theta)}{\theta_{q}}} \leq A_{\alpha} \int_{M} |\nabla_{g} u_{\alpha}|^{p} \eta_{\alpha}^{p} \, dv_{g} \left(\frac{\int_{M} u_{\alpha}^{q} \eta_{\alpha}^{p} \, dv_{g}}{\int_{M} u_{\alpha}^{q} dv_{g}} \right)^{\frac{p(1-\theta)}{\theta_{q}}} \leq cm_{\alpha}^{-\varrho(1+\frac{p(1-\theta)}{\theta_{q}})},$$

$$\int_{M} |\nabla_{g} \eta_{\alpha}|^{p} u_{\alpha}^{p} \, dv_{g} \left(\int_{M} (u_{\alpha} \eta_{\alpha}^{p})^{q} \, dv_{g} \right)^{\frac{p(1-\theta)}{\theta_{q}}} \leq A_{\alpha} \int_{M} |\nabla_{g} \eta_{\alpha}|^{p} u_{\alpha}^{p} \, dv_{g} \left(\frac{\int_{M} u_{\alpha}^{q} \eta_{\alpha}^{p} \, dv_{g}}{\int_{M} u_{\alpha}^{q} dv_{g}} \right)^{\frac{p(1-\theta)}{\theta_{q}}} \leq cm_{\alpha}^{-\varrho(1+\frac{p(1-\theta)}{\theta_{q}})}.$$
(13) and because $p \geq \tau$, we have

$$\alpha A_{\alpha}^{\frac{\tau-p}{p}} ||u_{\alpha}||_{L^{p}(M)}^{\tau-p} > c\alpha^{\frac{p}{\tau}} > c > 0 ,$$

 \mathbf{SO}

By

$$\int_{M} (u_{\alpha}\eta_{\alpha}^{p})^{p} \, dv_{g} \left(\int_{M} (u_{\alpha}\eta_{\alpha}^{p})^{q} \, dv_{g} \right)^{\frac{p(1-\theta)}{\theta_{q}}} \leq A_{\alpha} \int_{M} u_{\alpha}^{p}\eta_{\alpha}^{p} \, dv_{g} \left(\frac{\int_{M} u_{\alpha}^{q}\eta_{\alpha}^{p} \, dv_{g}}{\int_{M} u_{\alpha}^{q} dv_{g}} \right)^{\frac{p(1-\theta)}{\theta_{q}}} \leq cm_{\alpha}^{-\varrho(1+\frac{p(1-\theta)}{\theta_{q}})}$$

Replacing these three estimates in (38), one gets

$$\left(\int_{B(y_{\alpha},\frac{1}{2}A_{\alpha}^{\frac{1}{p}}u_{\alpha}(y_{\alpha})^{\frac{p-r}{p}})} u_{\alpha}^{r} dv_{g}\right)^{\frac{p}{r\theta}} \leq cm_{\alpha}^{-\varrho(1+\frac{p(1-\theta)}{\theta q})},$$

so that

$$m_{\alpha}^{\varrho} \int_{B(y_{\alpha}, \frac{1}{2}A_{\alpha}^{\frac{p}{p}} u_{\alpha}(y_{\alpha})^{\frac{p-r}{p}})} u_{\alpha}^{r} dv_{g} \leq c m_{\alpha}^{\varrho(1-\frac{r\theta}{p}-\frac{r(1-\theta)}{q})}.$$

Since $m_{\alpha} \to \infty$ and

$$1 - \frac{r\theta}{p} - \frac{r(1-\theta)}{q} < 0$$

we derive

$$m_{\alpha}^{\varrho} \int_{B(y_{\alpha}, \frac{1}{2}A_{\alpha}^{\frac{1}{p}} u_{\alpha}(y_{\alpha})^{\frac{p-r}{p}})} u_{\alpha}^{r} dv_{g} \to 0,$$

when $\alpha \to \infty$. But this contradicts (34).

2.4 The final argument in the proof of Theorem 1

In the sequel, we will perform several estimates by using the L^r -concentration and the pointwise estimation. By the scale invariance of the problem, we can assume that the radius of injectivity of M grater than one.

Let $\eta \in C_0^1(\mathbb{R})$ be a cutoff function as in the previous section and define $\eta_\alpha(x) = \eta(d_g(x, x_\alpha))$. From the inequality $GN_E(A(p, q, r, n))$ and by (3), we have

$$\left(\int_{B(0,1)} u_{\alpha}^{r} \eta_{\alpha}^{r} dx\right)^{\frac{p}{r\theta}} \leq A(p,q,r,n) \left(\int_{B(0,1)} |\nabla(u_{\alpha}\eta_{\alpha})|^{p} dx\right) \left(\int_{B(0,1)} u_{\alpha}^{q} \eta_{\alpha}^{q} dx\right)^{\frac{p(1-\theta)}{\theta q}}$$
$$\leq A_{opt}^{\frac{p}{\tau}} \left(\int_{B(0,1)} |\nabla(u_{\alpha}\eta_{\alpha})|^{p} dx\right) \left(\int_{B(0,1)} u_{\alpha}^{q} \eta_{\alpha}^{q} dx\right)^{\frac{p(1-\theta)}{\theta q}}.$$

Expanding the metric g in normal coordinates around x_{α} , one locally gets

$$(1 - cd_g(x, x_\alpha)^2) \, dv_g \le dx \le (1 + cd_g(x, x_\alpha)^2) \, dv_g \tag{39}$$

and

$$|\nabla(u_{\alpha}\eta_{\alpha})|^{p} \leq |\nabla_{g}(u_{\alpha}\eta_{\alpha})|^{p} (1 + cd_{g}(x, x_{\alpha})^{2}).$$

$$\tag{40}$$

Thus,

$$\left(\int_{B(0,1)} u_{\alpha}^{r} \eta_{\alpha}^{r} \, dx\right)^{\frac{p}{r\theta}}$$

$$\leq \left(A_{\alpha}A_{opt}^{\frac{p}{\tau}}\int_{M}|\nabla_{g}(u_{\alpha}\eta_{\alpha})|^{p} dv_{g} + cA_{\alpha}\int_{M}|\nabla_{g}(u_{\alpha}\eta_{\alpha})|^{p}d_{g}(x,x_{\alpha})^{2} dv_{g}\right)\left(\frac{\int_{B(0,1)}u_{\alpha}^{q}\eta_{\alpha}^{q} dx}{\int_{M}u_{\alpha}^{q} dv_{g}}\right)^{\frac{p(1-\theta)}{\theta_{q}}}$$

Applying then the inequalities

$$|\nabla_g(u_\alpha\eta_\alpha)|^p \le |\nabla_g u_\alpha|^p \eta_\alpha^p + c |\eta_\alpha \nabla_g u_\alpha|^{p-1} |u_\alpha \nabla_g \eta_\alpha| + c |u_\alpha \nabla_g \eta_\alpha|^p ,$$

(10) and (11), we have

$$A_{opt}^{\frac{p}{\tau}} A_{\alpha} \int_{M} |\nabla_{g} u_{\alpha}|^{p} dv_{g} \leq 1 - \alpha \left(A_{\alpha} \int_{M} u_{\alpha}^{p} dv_{g}\right)^{\frac{\tau}{p}}$$

one easily checks that

$$\left(\int_{B(0,1)} u_{\alpha}^{r} \eta_{\alpha}^{r} dx\right)^{\frac{p}{r\theta}}$$

$$\leq \left(1 - \alpha \left(A_{\alpha} \int_{M} u_{\alpha}^{p} dv_{g}\right)^{\frac{\tau}{p}} + cF_{\alpha} + cG_{\alpha} + cA_{\alpha} \int_{B(x_{\alpha},1)\setminus B(x_{\alpha},\frac{1}{2})} u_{\alpha}^{p} dv_{g}\right) \left(\frac{\int_{B(0,1)} u_{\alpha}^{q} \eta_{\alpha}^{q} dx}{\int_{M} u_{\alpha}^{q} dv_{g}}\right)^{\frac{p(1-\theta)}{\theta q}}, \quad (41)$$

where

$$F_{\alpha} = A_{\alpha} \int_{M} |\nabla_{g} u_{\alpha}|^{p} \eta_{\alpha}^{p} d_{g}(x, x_{\alpha})^{2} dv_{g}$$

and

$$G_{\alpha} = A_{\alpha} \int_{M} |\nabla_{g} u_{\alpha}|^{p-1} \eta_{\alpha}^{p-1} u_{\alpha} |\nabla_{g} \eta_{\alpha}| \, dv_{g} \,.$$

We now estimate F_{α} and G_{α} . Note that by (9), taking u_{α} as test function, we have

$$A_{\alpha} \int_{M} |\nabla_{g} u_{\alpha}|^{p} dv_{g} \leq \lambda_{\alpha} \leq A_{opt}^{-\frac{p}{\tau}} .$$

Then applying Hölder inequality, (16), the definition of φ_{α} , the inequality above and the pointwise estimate, we obtain

$$G_{\alpha} \leq \left(A_{\alpha} \int_{M} |\nabla_{g} u_{\alpha}|^{p} dv_{g}\right)^{\frac{p-1}{p}} \left(A_{\alpha} \int_{B(x_{\alpha},1) \setminus B(x_{\alpha},\frac{1}{2})} u_{\alpha}^{p} dv_{g}\right)^{\frac{1}{p}} \leq c \left(A_{\alpha} \int_{B(x_{\alpha},1) \setminus B(x_{\alpha},\frac{1}{2})} u_{\alpha}^{p} dg(x,x_{\alpha})^{p} dv_{g}\right)^{\frac{1}{p}}$$

$$\leq c \left(a_{\alpha}^{\frac{np-nr+pr}{r}+p-\frac{np}{r}+n} \int_{B(0,a_{\alpha}^{-1}) \setminus B(0,\frac{a_{\alpha}^{-1}}{2})} \varphi_{\alpha}^{p} |x|^{p} dh_{\alpha}\right)^{\frac{1}{p}} \leq c_{\lambda} a_{\alpha}^{2} \left(\int_{\mathbb{R}^{n} \setminus B(0,1)} |x|^{p(1-\lambda)} dx\right)^{\frac{1}{p}} \leq c a_{\alpha}^{2}, \quad (42)$$

with λ being large enough. Similarly,

$$A_{\alpha} \int_{M} |\nabla_{g} u_{\alpha}|^{p-1} \eta_{\alpha}^{p} u_{\alpha} d_{g}(x, x_{\alpha}) dv_{g} \leq \left(A_{\alpha} \int_{M} |\nabla_{g} u_{\alpha}|^{p} dv_{g}\right)^{\frac{p-1}{p}} \left(A_{\alpha} \int_{B(x_{\alpha}, 1)} u_{\alpha}^{p} d_{g}(x, x_{\alpha})^{p} dv_{g}\right)^{\frac{1}{p}} \\ \leq c a_{\alpha}^{2} \left(\int_{B(0, a_{\alpha}^{-1})} \varphi_{\alpha}^{p} |x|^{p} dh_{\alpha}\right)^{\frac{1}{p}} \leq c a_{\alpha}^{2} \left(1 + \int_{\mathbb{R}^{n} \setminus B(0, 1)} |x|^{p(1-\lambda)} dx\right)^{\frac{1}{p}} \leq c a_{\alpha}^{2} .$$

$$(43)$$

Now taking $u_{\alpha} d_g^2 \eta_{\alpha}^p$ as a test function in (10), one easily checks that

$$F_{\alpha} = A_{\alpha} \int_{M} |\nabla_g u_{\alpha}|^p \eta_{\alpha}^p d_g(x, x_{\alpha})^2 \, dv_g \le c \int_{B(x_{\alpha}, 1)} u_{\alpha}^r d_g(x, x_{\alpha})^2 \, dv_g + cA_{\alpha} \int_{M} |\nabla_g u_{\alpha}|^{p-1} \eta_{\alpha}^p u_{\alpha} d_g(x, x_{\alpha}) \, dv_g + cG_{\alpha} \, dv_g + cG$$

Therefore, by (42) and (43),

$$F_{\alpha} \le c \int_{B(x_{\alpha},1)} u_{\alpha}^{r} d_{g}(x,x_{\alpha})^{2} dv_{g} + ca_{\alpha}^{2}.$$

Again using the pointwise estimate, we get

$$\int_{B(x_{\alpha},1)} u_{\alpha}^{r} d_{g}(x,x_{\alpha})^{2} dv_{g} \leq ca_{\alpha}^{2} \left(1 + \int_{B(0,a_{\alpha}^{-1})\setminus B(0,1)} \varphi_{\alpha}^{r} |x|^{2} dx \right) \leq ca_{\alpha}^{2} \left(1 + \int_{\mathbb{R}^{n}\setminus B(0,1)} |x|^{2-\lambda r} dx \right) \leq ca_{\alpha}^{2}$$
(44)

for λ big enough. Consequently,

$$F_{\alpha} \le ca_{\alpha}^2 \quad \text{and} \quad G_{\alpha} \le ca_{\alpha}^2 \,.$$

$$\tag{45}$$

Proceeding as in (42) and since p > 1, we get

$$A_{\alpha} \int_{B(x_{\alpha},1)\setminus B(x_{\alpha},\frac{1}{2})} u_{\alpha}^{p} dv_{g} \leq cA_{\alpha} \int_{B(x_{\alpha},1)\setminus B(x_{\alpha},\frac{1}{2})} u_{\alpha}^{p} d_{g}(x,x_{\alpha})^{p} dv_{g} \leq ca_{\alpha}^{2p} \leq ca_{\alpha}^{2} .$$

Inserting this estimate and (45) in (41), one arrives at

$$\left(\int_{B(x_{\alpha},1)} u_{\alpha}^{r} \eta_{\alpha}^{r} dx\right)^{\frac{p}{r\theta}} \leq \left(1 - \alpha \left(A_{\alpha} \int_{M} u_{\alpha}^{p} dv_{g}\right)^{\frac{\tau}{p}} + ca_{\alpha}^{2}\right) \left(\frac{\int_{B(x_{\alpha},1)} u_{\alpha}^{q} \eta_{\alpha}^{q} dx}{\int_{M} u_{\alpha}^{q} dv_{g}}\right)^{\frac{p(1-\theta)}{\theta q}}.$$
(46)

Now by (39) and the mean value theorem, we obtain

$$\left(\int_{M} u_{\alpha}^{r} \eta_{\alpha}^{r} dx\right)^{\frac{p}{r\theta}} \ge \left(\int_{M} u_{\alpha}^{r} \eta_{\alpha}^{r} dv_{g} - c \int_{M} u_{\alpha}^{r} \eta_{\alpha}^{r} d_{g}(x, x_{\alpha})^{2} dv_{g}\right)^{\frac{p}{r\theta}}$$
$$\ge 1 - c \int_{M \setminus B(x_{\alpha}, 1)} u_{\alpha}^{r} dv_{g} - c \int_{M} u_{\alpha}^{r} \eta_{\alpha}^{r} d_{g}(x, x_{\alpha})^{2} dv_{g}$$

and

$$\left(\frac{\int_{B(x_{\alpha},1)} u_{\alpha}^{q} \eta_{\alpha}^{q} dx}{\int_{M} u_{\alpha}^{q} dv_{g}}\right)^{\frac{p(1-\theta)}{\theta_{q}}} \leq \left(\frac{\int_{M} u_{\alpha}^{q} \eta_{\alpha}^{q} dv_{g} + c \int_{M} u_{\alpha}^{q} \eta_{\alpha}^{q} dg(x, x_{\alpha})^{2} dv_{g}}{\int_{M} u_{\alpha}^{q} dv_{g}}\right)^{\frac{p(1-\theta)}{\theta_{q}}} \\ \leq \left(\frac{\int_{M} u_{\alpha}^{q} \eta_{\alpha}^{q} dv_{g}}{\int_{M} u_{\alpha}^{q} dv_{g}}\right)^{\frac{p(1-\theta)}{\theta_{q}}} + c \frac{\int_{M} u_{\alpha}^{q} \eta_{\alpha}^{q} dg(x, x_{\alpha})^{2} dv_{g}}{\int_{M} u_{\alpha}^{q} dv_{g}} \leq 1 + c \frac{\int_{M} u_{\alpha}^{q} \eta_{\alpha}^{q} dg(x, x_{\alpha})^{2} dv_{g}}{\int_{M} u_{\alpha}^{q} dv_{g}}$$

Replacing these two estimates in (46), one gets

$$\alpha \left(A_{\alpha} \int_{M} u_{\alpha}^{p} \, dv_{g}\right)^{\frac{\tau}{p}} \leq ca_{\alpha}^{2} + c \frac{\int_{M} u_{\alpha}^{q} \eta_{\alpha}^{q} d_{g}(x, x_{\alpha})^{2} \, dv_{g}}{\int_{M} u_{\alpha}^{q} \, dv_{g}} + c \int_{M} u_{\alpha}^{r} \eta_{\alpha}^{r} d_{g}(x, x_{\alpha})^{2} \, dv_{g} + c \int_{M \setminus B(x_{\alpha}, 1)} u_{\alpha}^{r} \, dv_{g} \, dv_{g} + c \int_{M \setminus B(x_{\alpha}, 1)} u_{\alpha}^{r} \, dv_{g} \, dv_{g} \, dv_{g} + c \int_{M \setminus B(x_{\alpha}, 1)} u_{\alpha}^{r} \, dv_{g} \, dv_{g$$

Using the pointwise estimate with $\lambda r - n = 2$, we have

$$\int_{M \setminus B(x_{\alpha},1)} u_{\alpha}^{r} \, dv_{g} \leq c \int_{M \setminus B(x_{\alpha},1)} u_{\alpha}^{r} d(x,x_{\alpha})^{\lambda r} \, dv_{g} \leq c_{\lambda} a_{\alpha}^{\lambda r-n} = c \, a_{\alpha}^{2}$$

So, by this estimate and (44), one concludes that

$$\alpha \left(A_{\alpha} \int_{M} u^{p}_{\alpha} \, dv_{g} \right)^{\frac{\tau}{p}} \leq ca^{2}_{\alpha} + c \frac{\int_{M} u^{q}_{\alpha} \eta^{q}_{\alpha} d_{g}(x, x_{\alpha})^{2} \, dv_{g}}{\int_{M} u^{q}_{\alpha} \, dv_{g}} \,. \tag{47}$$

By the pointwise estimate, for λ big enough

$$\frac{\int_{M} u_{\alpha}^{q} \eta_{\alpha}^{q} d_{g}(x, x_{\alpha})^{2} dv_{g}}{\int_{M} u_{\alpha}^{q} dv_{g}} \leq \frac{\int_{B(x_{\alpha}, 1)} u_{\alpha}^{q} d_{g}(x, x_{\alpha})^{2} dv_{g}}{\int_{M} u_{\alpha}^{q} dv_{g}}$$
$$= a_{\alpha}^{2} \int_{B(0, a_{\alpha}^{-1})} \varphi_{\alpha}^{q} |x|^{2} dh_{\alpha} \leq ca_{\alpha}^{2} \left(1 + c_{\lambda} \int_{\mathbb{R}^{n} \setminus B(0, 1)} |x|^{2 - \lambda q} dx\right) \leq c a_{\alpha}^{2} .$$

Introducing this inequality in (47), we readily deduce that

$$\alpha \left(A_{\alpha} \int_{M} u_{\alpha}^{p} \, dv_{g} \right)^{\frac{\tau}{p}} \le c \, a_{\alpha}^{2}$$

Then, since

$$\int_{M} u_{\alpha}^{p} \, dv_{g} \ge \int_{B(x_{\alpha}, a_{\alpha})} u_{\alpha}^{p} \, dv_{g} = a_{\alpha}^{\frac{n(r-p)}{r}} \int_{B(0,1)} \varphi_{\alpha}^{p} \, dh_{\alpha} \ge ca_{\alpha}^{\frac{n(r-p)}{r}}$$

for α large enough and since $a_{\alpha}^{\frac{n(r-p)}{r}}A_{\alpha} = a_{\alpha}^{p}$, one arrives at

$$\alpha a_{\alpha}^{\tau} \le c \; a_{\alpha}^2 \, .$$

Finally, because $\tau \leq 2$, we arrive at the contradiction

3 Existence of extremals for the optimal Gagliardo-Nirenberg inequality

Let us now prove Theorem 2

By Theorem 1 we have that the inequality

$$\left(\int_{M} |u|^{r} \, dv_{g}\right)^{\frac{\tau}{r\theta}} \leq \left(A_{opt}\left(\int_{M} |\nabla_{g}u|^{p} \, dv_{g}\right)^{\frac{\tau}{p}} + B_{opt}\left(\int_{M} |u|^{p} \, dv_{g}\right)^{\frac{\tau}{p}}\right) \left(\int_{M} |u|^{q} \, dv_{g}\right)^{\frac{\tau(1-\theta)}{\theta q}}$$

is valid for all $u \in H^{1,p}(M)$.

Let $\alpha > 0$ and $c_{\alpha} = B_{opt} - \alpha^{-1}$. Define

$$J_{\alpha}(u) = \left(\int_{M} |u|^{r} dv_{g}\right)^{\frac{\tau}{r\theta}} \left(\int_{M} |u|^{q} dv_{g}\right)^{-\frac{\tau(1-\theta)}{\theta q}} - c_{\alpha} \left(\int_{M} |u|^{p} dv_{g}\right)^{\frac{\tau}{p}}.$$

By the definition of B_{opt} we have

$$\nu_{\alpha} = \sup_{u \in E} J_{\alpha}(u) > A_{opt} \,,$$

where $E = \{ u \in H^{1,p}(M) : ||\nabla_g u||_{L^p(M)} = 1 \}$. Clearly, this supremum is well-defined.

In the same fashion as in section 2.1, we will give the proof for the C^1 -case q > 1, the case q = 1 being taken care or by the methods contained in [21]. Since then J_{α} is of class C^1 , by using standard variational arguments, we find a maximizer $\tilde{u}_{\alpha} \in E$ of J_{α} , and then

$$J_{\alpha}(\tilde{u}_{\alpha}) = \nu_{\alpha} = \sup_{u \in E} J_{\alpha}(u)$$

The function \tilde{u}_{α} satisfies the Euler-Lagrange equation

$$\frac{1}{\theta}||\tilde{u}_{\alpha}||_{L^{r}(M)}^{\frac{\tau-r\theta}{\theta}}||\tilde{u}_{\alpha}||_{L^{q}(M)}^{-\frac{\tau(1-\theta)}{\theta}}\tilde{u}_{\alpha}^{r-1}-\frac{(1-\theta)}{\theta}||\tilde{u}_{\alpha}||_{L^{r}(M)}^{\frac{\tau}{\theta}}||\tilde{u}_{\alpha}||_{L^{q}(M)}^{-\frac{\tau(1+\theta)+\theta q}{\theta}}\tilde{u}_{\alpha}^{q-1}-c_{\alpha}||\tilde{u}_{\alpha}||_{L^{p}(M)}^{\tau-p}\tilde{u}_{\alpha}^{p-1}=\nu_{\alpha}\Delta_{p,g}\tilde{u}_{\alpha},$$

where $\Delta_{p,g} = -\text{div}_g(|\nabla_g|^{p-2}\nabla_g)$ is the *p*-Laplace operator of *g*. Because $\nabla_g|\tilde{u}_{\alpha}| = \pm \nabla_g \tilde{u}_{\alpha}$, we can assume $\tilde{u}_{\alpha} \ge 0$ and again be Tolksdorf regularity [28], we have that $\tilde{u}_{\alpha} \in C^1(M)$. Setting $u_{\alpha} = \frac{\tilde{u}_{\alpha}}{||\tilde{u}_{\alpha}||_{L^r(M)}}$, we find

$$\lambda_{\alpha}^{-1} A_{\alpha} \Delta_{p,g} u_{\alpha} + c_{\alpha} A_{\alpha}^{\frac{\tau}{p}} ||u_{\alpha}||_{L^{p}(M)}^{\tau-p} u_{\alpha}^{p-1} + \frac{1-\theta}{\theta} ||u_{\alpha}||_{L^{q}(M)}^{-q} u_{\alpha}^{q-1} = \frac{1}{\theta} u_{\alpha}^{r-1} \quad \text{on} \quad M \,, \tag{48}$$

where $||u_{\alpha}||_{L^{r}(M)} = 1$,

$$A_{\alpha} = \left(\int_{M} u_{\alpha}^{q} \ dv_{g}\right)^{\frac{p(1-\theta)}{\theta_{q}}}$$

and

$$\lambda_{\alpha} = \nu_{\alpha}^{-1} ||u_{\alpha}||_{L^{q}(M)}^{\frac{(p-\tau)(1-\theta)}{\theta}} ||\tilde{u}_{\alpha}||_{L^{r}(M)}^{\tau-p} .$$

Up to taking a subsequence, we can assume that there exists $A \in \mathbb{R}$ such that

$$\lim_{\alpha \to \infty} A_{\alpha} = A \; .$$

Then there are two possibilities: either A = 0 or A > 0. We presently show that A = 0 cannot happen. Indeed, if A = 0 then (12) of Theorem 1 holds. Then we can follow the proof of theorem 1 and we get

$$c_{\alpha} \leq c \ a_{\alpha}^{2-\tau}$$
,

as in the end of section 2.4. Since $\tau < 2$, $c_{\alpha} = B_{opt} - \alpha^{-1}$ and $\lim_{\alpha \to \infty} a_{\alpha} = 0$, we have that above inequality contradicts (4). Therefore A > 0.

Using this fact in (48), we see that there exists c > 0 such that

$$\int_{M} |\nabla_{g} u_{\alpha}|^{p} dv_{g} + \left(\int_{M} u_{\alpha}^{p} dv_{g}\right)^{\frac{\tau}{p}} \leq c$$

for all α . Then, up to a subsequence, $u_{\alpha} \rightharpoonup u_0$ in $H^{1,p}(M)$. Since $||u_{\alpha}||_{L^r(M)} = 1$ for all α , we have that $||u_0||_{L^r(M)} = 1$. From (48) we have

$$\int_{M} |\nabla_{g} u_{\alpha}|^{p-2} \nabla_{g} u_{\alpha} \nabla_{g} h dv_{g} \leq c \int_{M} u_{\alpha}^{r-1} h dv_{g} ,$$

for an arbitrary test function $h \ge 0$. Then by Moser's iterative scheme,

$$\sup_{x \in M} u_{\alpha} \le c \left(\int_{M} u_{\alpha}^{r} dv_{g} \right)^{\frac{1}{r}} \le c ,$$

for all α .

Using Tolksdorf regularity in (48), it follows that $u_{\alpha} \to u_0$ in $C^1(M)$.

The function \tilde{u}_{α} satisfies

$$\left(\int_{M} \tilde{u}_{\alpha}^{r} dv_{g}\right)^{\frac{\tau}{r\theta}} \geq \left(A_{opt}^{\frac{p}{\tau}} \left(\int_{M} |\nabla_{g} \tilde{u}_{\alpha}|^{p} dv_{g}\right)^{\frac{\tau}{p}} + (B_{opt} - \frac{1}{\alpha}) \left(\int_{M} \tilde{u}_{\alpha}^{p} dv_{g}\right)^{\frac{\tau}{p}}\right) \left(\int_{M} \tilde{u}_{\alpha}^{q} dv_{g}\right)^{\frac{\tau(1-\theta)}{\theta_{q}}}$$

and since $u_{\alpha} = \frac{\tilde{u}_{\alpha}}{||\tilde{u}_{\alpha}||_{L^{r}(M)}}$, we get

$$1 \ge \left(A_{opt}^{\frac{p}{\tau}} \left(\int_{M} |\nabla_{g} u_{\alpha}|^{p} dv_{g}\right)^{\frac{\tau}{p}} + (B_{opt} - \frac{1}{\alpha}) \left(\int_{M} u_{\alpha}^{p} dv_{g}\right)^{\frac{\tau}{p}}\right) \left(\int_{M} u_{\alpha}^{q} dv_{g}\right)^{\frac{\tau(1-\theta)}{\theta q}}$$

Taking the limit in α in this inequality, we find

$$1 \ge \left(A_{opt}^{\frac{p}{\tau}} \left(\int_{M} |\nabla_g u_0|^p dv_g\right)^{\frac{\tau}{p}} + B_{opt} \left(\int_{M} u_0^p dv_g\right)^{\frac{\tau}{p}}\right) \left(\int_{M} u_0^q dv_g\right)^{\frac{\tau(1-\theta)}{\theta_q}}$$

And then u_0 is an extremal function for $GN_R(A_{opt}, B_{opt})$.

References

- M. Agueh Sharp Gagliardo-Nirenberg Inequalities via p-Laplacian type equations, Nonlinear Diff. Eq. Appl. 15 (2008) 457-472.
- [2] T. Aubin Problèmes isopérimétriques et espaces de Sobolev, J. Differential Geom. 11 (1976) 573-598.
- [3] T. Aubin, Y.Y Li On the best Sobolev inequality, J. Math. Pures Appl. 78 (1999) 353-387.
- [4] D. Bakry, T. Coulhon, M. Ledoux, L. Sallof-Coste Sobolev inequalities in disguise, Indiana J. Math. 44 (1995) 1033-1074.
- [5] W. Beckner Estimates on Moser Embedding, Potential Analysis 20 (2004) 345-359.
- [6] C. Brouttelande The best-constant problem for a family of Gagliardo-Nirenberg inequalities on a compact Riemannian manifold, Proc. R. Soc. Edinb. 46 (2003) 147-157.
- [7] E. Carlen, M. Loss Sharp constants in Nash's inequality, Int. Math. Res. Not. 7 (1993) 213-215.
- [8] J. Ceccon, M. Montenegro Optimal L^p-Riemannian Gagliardo-Nirenberg inequalities, Math. Z. 258 (2008) 851-873.
- [9] J. Ceccon, M. Montenegro Optimal Riemannian L^p-Gagliardo-Nirenberg inequalities revisited, J. Differential Equations 254, 6 (2013) 2532-2555.
- [10] J. Ceccon, M. Montenegro Sharp L^p-entropy inequalities on manifolds, arXiv:1307.7115 [math.AP].
- [11] J. Chen, B. Guo Sharp constant of an improved Gagliardo-Nirenberg inequality and its application, Ann. Mat. Pura Appl. 4, n. 2, 190 (2011) 341-354.
- [12] W. Chen, X. Sun Optimal improved L_p-Riemannian Gagliardo-Nirenberg inequalities, Nonlinear Analysis 72 (2010) 3159-3172.

- [13] D. Cordero-Erausquin, B. Nazaret, C. Villani A mass-transportation approach to sharp Sobolev and Gagliardo-Nirenberg inequalities, Adv. Math. 182 (2004) 307-332.
- [14] M. Del Pino, J. Dolbeault Best constants for Gagliardo-Nirenberg inequalities and applications to nonlinear diffusions, J. Math. Pures Appl. 81 (2002) 847-875.
- [15] M. Del Pino, J. Dolbeault The optimal Euclidean L^p-Sobolev logarithmic inequality, J. Funct. Anal. 197 (2003) 151-161.
- [16] O. Druet Optimal Sobolev inequalities of arbitrary order on compact Riemannian manifolds, J. Funct. Anal. 159 (1998) 217-242.
- [17] O. Druet The best constants problem in Sobolev inequalities, Math. Ann. 314 (1999) 327-346.
- [18] O. Druet, E. Hebey, M. Vaugon Optimal Nash's inequalities on Riemannian manifolds: the influence of geometry, Int. Math. Res. Not. 14 (1999) 735-779.
- [19] E. Gagliardo Proprietà di alcune classi di funzioni in piu variabili, Ricerche Mat. 7 (1958) 102-137.
- [20] E. Hebey Nonlinear Analysis on Manifolds: Sobolev Spaces and Inequalities, Courant Institute of Mathematical Sciences, Lecture Notes in Mathematics, 5 (1999).
- [21] E. Humbert Best constants in the L^2 -Nash inequality, Proc. R. Soc. Edinb. 131 (2001) 621-646.
- [22] N. Kishimoto, M. Maeda Construction of blow-up solutions for Zakharov system on T², Ann. I. H. Poincaré 30 (2013), n. 5, 791-824.
- [23] J. Moser On Harnack's theorem for elliptic differential equations, Commun. Pure Appl. Math. 14 (1961) 577-591.
- [24] J. Nash Continuity of solutions of parabolic and elliptic equations, Am. J. Math. 80 (1958) 931-954.
- [25] L. Nirenberg On elliptic partial differential equations, Ann. Sc. Norm. Super. Pisa Cl. Sci. 13 (1959) 115-162.
- [26] J. Shu, J. Zhang Global Existence for a System of Weakly Coupled Nonlinear Schrödinger Equations, Math. Notes 86, 5-6 (2009) 650-654.
- [27] S. Sobolev Sur un théorème d'analyse fonctionnelle, Rec. Math. [Mat. Sbornik] N.S. 46 (1938) 471-497.
- [28] P. Tolksdorf Regularity for a more general class of quasilinear elliptic equations, J. Differential Equations, 51 (1) (1984) 126-150.