BOULIGAND-SEVERI *k*-TANGENTS AND STRONGLY SEMISIMPLE MV-ALGEBRAS

LEONARDO MANUEL CABRER

ABSTRACT. An algebra A is said to be strongly semisimple if every principal congruence of A is an intersection of maximal congruences. We give a geometrical characterisation of strongly semisimple MV-algebras in terms of Bouligand-Severi k-tangents. The latter are a k-dimensional generalisation of the classical Bouligand-Severi tangents.

Each semisimple MV-algebra A is isomorphic to a separating MV-algebra of continuous [0, 1]-valued maps defined on a compact Hausdorff space X, which turns out to be homeomorphic to the maximal spectral space of A. In the particular case when A is *n*-generated, it is no loss of generality to assume that X is a compact subset of $[0, 1]^n$ and A is isomorphic to the MV-algebra $\mathcal{M}(X)$ of McNaughton maps of $[0, 1]^n$ restricted to X (see [5, Thm. 3.6.7]).

Following Dubuc and Poveda [6], we say that an MV-algebra A is strongly semisimple if all its principal quotients are semisimple. In [4], Busaniche and Mundici characterise those sets $X \subseteq [0,1]^2$ having the property that the MValgebra $\mathcal{M}(X)$ to be strongly semisimple. Their result (Theorem 0.2 below) is formulated in terms of the following classical notion (see [2, 11, 12]; also see [1, p.16], [8, pp.14 and 133] for modern reformulations):

Definition 0.1. Let $\emptyset \neq X \subseteq \mathbb{R}^n$ and $x \in \mathbb{R}^n$. A Bouligand-Severi tangent of X at x is a unit vector $u \in \mathbb{R}^n$ such that X contains a sequence x_1, x_2, \ldots with the following properties:

- (i) $x_i \neq x$ for all i,
- (ii) $\lim_{i\to\infty} x_i = x$, and
- (iii) $\lim_{i \to \infty} (x_i x) / ||x_i x|| = u.$

Theorem 0.2. [4, Thm. 2.4] Let $X \subseteq [0,1]^2$ be a closed set. The MV-algebra $\mathcal{M}(X)$ is not strongly semisimple iff there exist a point $x \in X$, a unit vector $u \in \mathbb{R}^2$, and a real number $\lambda > 0$ such that

- (i) *u* is a Bouligand-Severi tangent of *X* at *x*,
- (ii) $\operatorname{conv}(x, x + \lambda u) \cap X = \{x\}, and$
- (iii) the coordinates of x and $x + \lambda u$ are rational.

In Theorem 2.4 this result will be generalised to all finitely generated MValgebras, using the higher-order Bouligand-Severi tangents defined in 2.1-2.2.

Key words and phrases. MV-algebra, strongly semisimple, ideal, Bouligand-Severi tangent, rational polyhedron.

AMS2010. Primary: 06D35. Seondary: 54C40, 49J52, 52B05, 03B50.

This research was supported by a Marie Curie Intra European Fellowship within the 7th European Community Framework Program (ref. 299401-FP7-PEOPLE-2011-IEF).

L.M. CABRER

1. Preliminaries

1.1. Semisimple MV-algebras. We refer the reader to [5] for background on MV-algebras. We let $\mathcal{M}([0,1]^n)$ denote the MV-algebra of piecewise (affine) linear continuous functions $f: [0,1]^n \to [0,1]$, such that each linear piece of f has integer coefficients, with the pointwise operations of the standard MV-algebra [0,1]. $\mathcal{M}([0,1]^n)$ is the free *n*-generator MV-algebra. More generally, for any nonempty subset $X \subseteq [0,1]^n$ we denote by $\mathcal{M}(X)$ the MV-algebra of restrictions to X of the functions in $\mathcal{M}([0,1]^n)$. For every $f \in \mathcal{M}(X)$ we let $Zf = f^{-1}(0)$.

By an *ideal* of an MV-algebra A we mean the kernel of an (MV-)homomorphism. An ideal is *principal* if it is singly generated. For each $a \in A$, the principal ideal $\langle a \rangle$ generated by a is the set $\{b \in A \mid \text{for some } m \in \mathbb{Z}_{>0}, b \leq ma\}$. An ideal I is *maximal* if $I \neq A$, and whenever J is an ideal such that $I \subseteq J$, then J = I or J = A. For each closed set $X \subseteq [0,1]^n$ and $x \in X$, the set $I_x = \{f \in \mathcal{M}(X) \mid f(x) = 0\}$ is a maximal ideal of $\mathcal{M}(X)$. Moreover, for each maximal ideal I of $\mathcal{M}(X)$, there exists a uniquely determined $x \in X$ such that $I = I_x$

An MV-algebra A is said to be *semisimple* if the intersection of its maximal ideals is $\{0\}$. Each semisimple MV-algebra is isomorphic to a separating MV-algebra of continuous maps from a compact Hausdorff space into [0, 1]. In particular, if A is an *n*-generated semisimple MV-algebra then there is a closed set $X \subseteq [0, 1]^n$ such that $A \cong \mathcal{M}(X)$.

An MV-algebra A is strongly semisimple if every principal ideal of A is an intersection of maximal ideals of A. Equivalently, A is strongly semisimple if for each $a \in A$, the quotient algebra $A/\langle a \rangle$ is semisimple.

1.2. Simplicial Geometry. We refer to [7], [10] and [13] for background in elementary polyhedral topology.

For any set $\{v_0, \ldots, v_m\} \subseteq \mathbb{R}^n$, $\operatorname{conv}(v_0, \ldots, v_m)$ denotes its *convex hull*. If $\{v_0, \ldots, v_m\}$ are affinely independent then $S = \operatorname{conv}(v_0, \ldots, v_m)$ is an *m*-simplex. For any $V \subseteq \{v_0, \ldots, v_m\}$, the convex hull $\operatorname{conv}(V)$ is called a *face* of *S*. If |V| = m - 1 then $\operatorname{conv}(V)$ is called a *facet* of *S*.

For any *m*-simplex $S = \operatorname{conv}(v_0, \ldots, v_m) \subseteq \mathbb{R}^n$, we let aff S denote the affine hull of S, i.e.,

aff
$$S = \{\sum_{i=0}^{m} \lambda_i v_i \mid \text{for some } \lambda_i \in \mathbb{R}, \text{ with } \sum_{i=0}^{m} \lambda_i = 1\}$$

= $v_0 + \mathbb{R}(v_1 - v_0) + \dots + \mathbb{R}(v_m - v_{m-1}).$

Further, we write relint S for the relative interior of S, that is, the topological interior of S in the relative topology of aff(S). For each $v \in \mathbb{R}^n$, ||v|| denotes the euclidean norm of v in \mathbb{R}^n . For each $0 < \delta \in \mathbb{R}$ and $v \in \mathbb{R}^n$ we use the notation $B(\delta, v) = \{w \in \mathbb{R}^n \mid ||v - w|| < \delta\}$ for the open ball of radius δ centred at v. Then

relint $S = \{ v \in \mathbb{R}^n \mid \text{for some } \delta > 0, \ B(\delta, v) \cap \text{aff } S \subseteq S \}.$

For later use we record here some elementary properties of simplexes.

Lemma 1.1. Let $T \subseteq \mathbb{R}^n$ be a simplex and F a face of T. If $x \in T$ then

 $x \in \operatorname{relint} F$ iff F is the smallest face of T such that $x \in F$.

Moreover, for any simplex S contained in T we have

 $S \subseteq F$ iff $F \cap \operatorname{relint} S \neq \emptyset$.

Notation and Terminology. Given $x \in \mathbb{R}^n$, a k-tuple $u = (u_1, \ldots, u_k)$ of pairwise orthogonal unit vectors in \mathbb{R}^n and a k-tuple $\lambda = (\lambda_1, \ldots, \lambda_k) \in \mathbb{R}^k_{>0}$, we write

$$C_{x,u,\lambda} = \operatorname{conv}(x, x + \lambda_1 u_1, \dots, x + \lambda_1 u_1 + \dots + \lambda_k u_k),$$

For any k-tuple $a = (a_1, \ldots, a_k)$ and $l = 1, \ldots, k$ we let a(l) be an abbreviation of the initial segment (a_1, \ldots, a_l) . Then the simplex $C_{x,u(l),\lambda(l)}$ is a face of $C_{x,u,\lambda}$.

Lemma 1.2. [3, Prop. 2.2] For each $x \in \mathbb{R}^n$ and k-tuple $u = (u_1, \ldots, u_k)$ of pairwise orthogonal unit vectors in \mathbb{R}^n , the family of (x, u)-simplexes ordered by inclusion is down-directed. That is, if C_1 and C_1 are (x, u)-simplexes, then $C_1 \cap C_2$ contains an (x, u)-simplex.

1.3. Rational Polyhedra and \mathbb{Z} -maps. An *m*-simplex $S = \operatorname{conv}(v_0, \ldots, v_m)$ is said to be *rational* if the coordinates of each vertex of S are rational numbers. A subset P of \mathbb{R}^n is said to be a *rational polyhedron* if there are rational simplexes T_1, \ldots, T_l such that $P = T_1 \cup \cdots \cup T_l$.

Given a rational polyhedron P, a triangulation of P is a simplicial complex Δ such that $P = \bigcup \Delta$ and each simplex $S \in \Delta$ is rational. Given triangulations Δ and Σ of P, we say that Δ is a subdivision of Σ if every simplex of Δ is contained in a simplex of Σ .

For v a rational point in \mathbb{R}^n we let $\operatorname{den}(v)$ denote the least common denominator of the coordinates of v. A rational *m*-simplex $S = \operatorname{conv}(v_0, \ldots, v_m) \subseteq \mathbb{R}^n$ is called *regular* if the set of vectors $\{\operatorname{den}(v_0)(v_0, 1), \ldots, \operatorname{den}(v_m)(v_m, 1)\}$ is part of a basis of the free abelian group \mathbb{Z}^{n+1} . By a *regular triangulation* of a rational polyhedron Pwe understand a triangulation of P consisting of regular simplexes.

Given polyhedra $P \subseteq \mathbb{R}^n$ and $Q \in \mathbb{R}^m$, a map $\eta: P \to Q$ is called a \mathbb{Z} -map if there is a triangulation Δ of P such that on every simplex T of Δ , η coincides with an affine linear map $\eta_T: \mathbb{R}^n \to \mathbb{R}^m$ with integer coefficients. In particular, $f \in \mathcal{M}([0,1]^n)$ iff it is a \mathbb{Z} -map form $[0,1]^n$ to [0,1].

For later use we recall here some properties of regular triangulations and \mathbb{Z} -maps. (See [9, Chapters 2,3] for the proofs.)

Lemma 1.3. Let P and Q be rational polyhedra and Δ a rational triangulation of P. If $Q \subseteq P$, there exists a regular triangulation Δ' of P which is a subdivision of Δ and also satisfies $Q = \bigcup \{S \in \Delta' \mid S \subseteq Q\}$.

Lemma 1.4. Let P and Q be rational polyhedra, and $\eta: P \to Q$ a \mathbb{Z} -map. If R is a rational polyhedron contained in Q, then $\eta^{-1}(R)$ is a rational polyhedron.

Lemma 1.5. Let P and Q be rational polyhedra, $\eta: P \to Q$ a \mathbb{Z} -map and Δ a triangulation of P. Then there is a regular triangulation ∇ of P which is a subdivision of Δ and has the property that the restriction $\eta \upharpoonright_S$ of η to S is (affine) linear for each $S \in \nabla$.

Lemma 1.6. Let Δ be a regular triangulation of a polyhedron $P \subseteq \mathbb{R}^m$ and V the set of vertices of the simplexes of Δ . Suppose the map $f: V \to \mathbb{R}^n$ has the property that f(v) is a rational vector of \mathbb{R}^n and $\operatorname{den}(f(v))$ divides $\operatorname{den}(v)$ for each $v \in V$. Then there exists a unique \mathbb{Z} -map $\mu: P \to \mathbb{R}^n$ satisfying:

- (i) μ is linear on each simplex of Δ ,
- (ii) $\mu \upharpoonright V = f$.

2. Strong semisimplicity and Bouligand-Severi tangents

Here we introduce k-dimensional Bouligand-Severi tangents, replacing the unit vector u of Definition 0.1 by a k-tuple $u = (u_1, \ldots, u_k)$ of pairwise orthogonal unit vectors in \mathbb{R}^n . For each $l \leq k$, let

$$\mathbf{p}_l: \mathbb{R}^n \to \mathbb{R}u_1 + \cdots + \mathbb{R}u_l$$

denote the orthogonal projection map onto the linear subspace of \mathbb{R}^n generated by u_1, \ldots, u_l .

Definition 2.1. A k-tuple $u = (u_1, \ldots, u_k)$ of pairwise orthogonal unit vectors in \mathbb{R}^n is said to be a *Bouligand-Severi tangent of* X at x of degree k (for short, u is a k-tangent of X at x) if X contains a sequence of points x_1, x_2, \ldots converging to x, such that no vector $x_i - x$ lies in $\mathbb{R}u_1 + \cdots + \mathbb{R}u_k$ and upon defining $x_i^1 = (x_i - x)/||x_i - x||$ and inductively,

$$x_{i}^{l} = \frac{x_{i} - x - \mathsf{p}_{l-1}(x_{i} - x)}{||x_{i} - x - \mathsf{p}_{l-1}(x_{i} - x)||} \quad (l \le k),$$

it follows that $\lim_{i\to\infty} x_i^s = u_s$, for each $s \in \{1, \ldots, k\}$. The sequence x_1, x_2, \ldots is said to *determine u*.

Conditions (ii) and (iii) in Theorem 2.3 have the following generalisation:

Definition 2.2. A k-tangent $u = (u_1, \ldots, u_k)$ of $X \subseteq \mathbb{R}^n$ at x is rationally outgoing if there is a rational simplex S, together with a face $F \subseteq S$ and a k-tuple $\lambda = (\lambda_1, \ldots, \lambda_k) \in \mathbb{R}_{>0}$ such that $S \supseteq C_{x,u,\lambda}$, $F \not\supseteq C_{x,u,\lambda}$ and $F \cap X = S \cap X$.

Remarks. When k = 1, Definition 2.1 amounts to the classical Definition 0.1 of a Bouligand-Severi tangent of a closed set in \mathbb{R}^n . Any subsequence of x_0, x_1, \ldots also determines the tangent u. Further, if $u = (u_1, \ldots, u_k)$ is a k-tangent of X at x then $u(l) = (u_1, \ldots, u_l)$ is an l-tangent of X at x for each $l \in \{1, \ldots, k\}$.

If u is a rationally outgoing k-tangent of $X \subseteq \mathbb{R}^n$ then, trivially, k < n. In particular if n = 2 then necessarily k is equal to 1, and there is $\epsilon > 0$ such that $\operatorname{conv}(x, x + \epsilon u_1)$ is a rational polyhedron and $X \cap \operatorname{conv}(x, x + \epsilon u_1) = \{x\}$. The main result of [4] (Theorem 0.2 above) can now be restated as follows:

Theorem 2.3. Let $X \subseteq [0,1]^2$ be a closed set. The MV-algebra $\mathcal{M}(X)$ is strongly semisimple iff X does not have a rationally outgoing 1-tangent.

The main result of our paper is the following generalisation of Theorem 2.3:

Theorem 2.4. For any closed $X \subseteq [0,1]^n$ the following conditions are equivalent:

- (i) The MV-algebra $\mathcal{M}(X)$ is strongly semisimple.
- (ii) For no k = 1, ..., n 1, X has a rationally outgoing k-tangent.

Each direction of the equivalence in Theorem 2.4 depends on a key property of rationally outgoing k-tangents. Accordingly, the proof is divided in two parts, each of them proved in a separate section.

3. PROOF OF THEOREM 2.4: (I) \Rightarrow (II)

Lemma 3.1. Let $P \subseteq \mathbb{R}^n$ be a polyhedron, $X \subseteq P$ a closed set, and $u = (u_1, \ldots, u_k)$ a k-tangent of X at x. Then P contains an (x, u)-simplex.

Proof. Let y_1, y_2, \ldots be a sequence of elements in X determining the tangent u. Let Δ be a triangulation of P, and S a simplex of P such that $\{i \mid y_i \in S\}$ is infinite. Since S is closed, $x \in S$. Let x_1, x_2, \ldots be the subsequence of y_1, y_2, \ldots whose elements lie in S. Then x_1, x_2, \ldots determines the k-tangent $u = (u_1, \ldots, u_k)$ of $X \cap S$ at x.

We will first prove

(1)
$$x + \mathbb{R}u_1 + \dots + \mathbb{R}u_k \subseteq \text{aff } S.$$

For all i we have

$$x + \mathbb{R}x_i^1 = x + \mathbb{R}\frac{x - x_i}{||x - x_i||} = x + \mathbb{R}(x - x_i) \subseteq \text{aff } S.$$

Since aff S is closed then $x + \mathbb{R}u_1 \subseteq$ aff S. Suppose we have proved

 $x + \mathbb{R}u_1 + \dots + \mathbb{R}u_{l-1} \subseteq \text{aff } S.$

Let $y \in x + \mathbb{R}u_1 + \cdots + \mathbb{R}u_{l-1}$. Then

$$y + \mathbb{R}x_i^l = y + \mathbb{R}\frac{x - x_i - \mathsf{p}_{l-1}(x - x_i)}{||x - x_i - \mathsf{p}_{l-1}(x - x_i)||}$$
$$= y + \mathbb{R}(x - x_i - \mathsf{p}_{l-1}(x - x_i))$$
$$\subseteq y + \mathbb{R}(x - x_i) + \mathbb{R}u_1 + \dots + \mathbb{R}u_{l-1} \subseteq \operatorname{aff} S$$

Again, $y + \mathbb{R}u_l \subseteq \operatorname{aff} S$ and $x + \mathbb{R}u_1 + \cdots + \mathbb{R}u_l \subseteq \operatorname{aff} S$. This concludes the proof of (1).

We shall now find $\lambda = (\lambda_1, \ldots, \lambda_k) \in \mathbb{R}^k_{>0}$ such that $C_{x,u,\lambda} \subseteq S$. To this purpose we will prove the following stronger statement:

Claim: For each $l \leq k$, there exists $\lambda(l) = (\lambda_1, \ldots, \lambda_l) \in \mathbb{R}^l_{>0}$ such that

- (i) $C_{x,u(l),\lambda(l)} \subseteq S$, and (ii) if F is a face of S such that $z_l = x + \lambda_1 u_1 + \cdots + \lambda_l u_l \in F$, then $C_{x,u(l),\lambda(l)} \subseteq F.$

The proof is by induction on $l = 1, \ldots, k - 1$.

Basis Step (l = 1):

In case $x \in \operatorname{relint} S$ there exists $\epsilon > 0$ such that $B(\epsilon, x) \cap \operatorname{aff} S \subseteq \operatorname{relint} S$. Then setting $\lambda_1 = \epsilon/2$, by (1) we obtain $C_{x,u(1),\lambda_1} \subseteq B(\epsilon, x) \cap \text{aff } S \subseteq \text{relint } S$, from which both (i) and (ii) immediately follow.

In case x does not lie in relint S, let H be an arbitrary facet of S containing x as an element. Let aff H^+ be the half-space of aff S with boundary aff H and containing S. For each $\rho > 0$ we have the inclusion $x + \rho(x_i - x) \in \text{aff } H^+$. Since aff H^+ is closed, then $x + \mathbb{R}_{>} u_1 \subseteq$ aff H^+ . As a consequence,

(2)
$$x + \mathbb{R}_{\geq 0} u_1 \subseteq \bigcap \{ \text{ aff } G^+ \mid G \text{ a facet of } S \text{ containing } x \}.$$

For some $\epsilon_1 > 0$ the points $x + \epsilon_1 u_1$ must lie in S. (For otherwise some facet K of S has the property that for each $\epsilon > 0$, $x + \epsilon u_1 \in \text{aff } S \setminus \text{aff } K^+$, where aff K^+ is the half-space of aff S with boundary aff K and containing S. From $x \in S \subseteq$ aff H^+ we get $x \in S \cap \text{aff } K = K$ contradicting (2).)

Now let $\lambda_1 = \epsilon_1/2$. Then, $C_{x,\lambda_1,u_1} \subseteq \operatorname{conv}(x, x + \epsilon_1 u_1) \subseteq S$, and (i) is settled. Let F be a face of S containing $x + \lambda_1 u_1$. Since $x + \lambda_1 u_1$ lies in the relative interior of $\operatorname{conv}(x, x + \epsilon_1 u_1) \subseteq S$, by Lemma 1.1 $C_{x,u_1,\lambda_1} \subseteq \operatorname{conv}(x, x + \epsilon_1 u_1) \subseteq F$. This proves (ii), and concludes the proof of the basis step.

Inductive Step:

Assume that our claim holds for l < k. Then there exists $\lambda(l) = (\lambda_1, \ldots, \lambda_l)$ in $\mathbb{R}^l_{>0}$ such that $C_{x,u(l),\lambda(l)} \subseteq S$ and if F is a face of S containing $x + \lambda_1 u_1 + \cdots + \lambda_l u_l$, then $C_{x,u(l),\lambda(l)} \subseteq F$.

For the rest of the proof let $z_l = x + \lambda_1 u_1 + \cdots + \lambda_l u_l$ and F_l be the face of S such that $z_l \in \operatorname{relint} F_l$.

As in the basis step, in case $F_l = S$, there exists $\epsilon \in \mathbb{R}_{>0}$ such that

$$B(\epsilon, z_l) \cap \operatorname{aff} S \subseteq \operatorname{relint} S.$$

Setting $\lambda_{l+1} = \epsilon/2$ and $\lambda(l+1) = (\lambda(l), \lambda_{l+1})$, by (1) we get

 $z_l + \lambda_{l+1} u_{l+1} \in \operatorname{aff} S \cap B(\epsilon, z_l) \subseteq \operatorname{relint} S.$

Since S is a simplex and $C_{x,u(l),\lambda(l)} \subseteq S$, then

$$C_{x,u(l+1),\lambda(l+1)} = \operatorname{conv}(C_{x,u(l),\lambda(l)} \cup \{z_l + \lambda_{l+1}u_{l+1}\}) \subseteq S,$$

which proves (i) and (ii).

In case F_l is a proper face of S, let H be an arbitrary facet of S containing F_l . Let aff H^+ be the closed half-space of aff S with boundary aff H containing S. From (1), we obtain

(3)
$$z_l + \mathbb{R}_{>0} u_{l+1} \subseteq \bigcap \{ \text{ aff } G^+ \mid G \text{ a facet of } S \text{ containing } F_l \}.$$

Therefore, $z_l + \mathbb{R}_{\geq 0} u_{l+1} \cap S \neq \{z_l\}$. For otherwise, arguing by way of contradiction, there is a facet K of S such that $z_l + \epsilon u_{l+1} \in \operatorname{aff} S \setminus \operatorname{aff} K^+$ for each $\epsilon > 0$. Since $z_l \in \operatorname{aff} K^+$ then $z_l \in K$. Since F_l is the smallest face of S containing z_l , it follows that $F_l \subseteq K$, contradicting (3).

We have just proved that there exists $\epsilon_{l+1} > 0$ such that $z_l + \epsilon_{l+1}u_{l+1} \in S$. Letting $\lambda_{l+1} = \epsilon/2$ we have

$$C_{x,u(l+1),\lambda(l+1)} = \operatorname{conv}(C_{x,u(l),\lambda(l)} \cup \{z_l + \lambda_{l+1}u_{l+1}\}) \subseteq S,$$

which settles (i).

For any face F of S such that $z_l + \lambda_{l+1}u_1 \in F$, by Lemma 1.1 we have $z_l \in \operatorname{conv}(z_l, z_l + \epsilon_{l+1}u_{l+1}) \subseteq F$. Therefore, $F_l \subseteq F$, and by inductive hypothesis $C_{x,u(l),\lambda(l)} \subseteq F$, whence

$$C_{x,u(l+1),\lambda(l+1)} = \operatorname{conv}(C_{x,u(l),\lambda(l)} \cup \{z_l + \lambda_{l+1}u_{l+1}\}) \subseteq F.$$

Having thus proved (ii), the claim is settled and the lemma is proved.

Proof of Theorem 2.4: (i) \Rightarrow (ii). Let $u = (u_1, \ldots, u_k)$ be a rationally outgoing ktangent of X at x, with the intent of proving that $\mathcal{M}(X)$ is not strongly semisimple. With reference to Definition 2.1, let S be a rational k-simplex together with a proper face $F \subseteq S$ and reals $\lambda = (\lambda_1, \ldots, \lambda_k) \in \mathbb{R}^k_{>0}$ such that $C_{x,u,\lambda} \subseteq S, S \cap X = F \cap X$, and $C_{x,u,\lambda} \not\subseteq F$.

By Lemma 1.3 there exists a regular triangulation Δ of $[0,1]^n$ such that $S = \bigcup \{T \in \Delta \mid T \subseteq S\}$ and $F = \bigcup \{R \in \Delta \mid R \subseteq F\}$. Let $f, g \in \mathcal{M}([0,1]^n)$ be the uniquely determined maps which are (affine) linear over each simplex of Δ , and for each vertex v (of a simplex) in Δ satisfy the conditions

$$f(v) = \begin{cases} 0 & \text{if } v \in F; \\ 1 & \text{otherwise;} \end{cases} \qquad g(v) = \begin{cases} 0 & \text{if } v \in S; \\ 1 & \text{otherwise.} \end{cases}$$

The existence of f and g follows from Lemma 1.6. Observe that Zg = S and Zf = F. Then:

(4)
$$Zf \upharpoonright_X = X \cap Zf = X \cap F = X \cap S = X \cap Zg = Zg \upharpoonright_X.$$

This proves that $f \upharpoonright_X$ belongs to a maximal ideal of $\mathcal{M}(X)$ iff $g \upharpoonright_X$ does. To complete the proof of (i) \Rightarrow (ii) it suffices to settle the following

Claim. $f \upharpoonright_X$ does not belong to the ideal $\langle g \upharpoonright_X \rangle$ generated by $g \upharpoonright_X$.

As a matter of fact, arguing by way of contradiction and letting the integer m > 0satisfy $f \upharpoonright_X \leq mg \upharpoonright_X$, it follows that X is contained in the rational polyhedron $P = \{y \in [0,1]^n \mid f(y) \leq mg(y)\}$. An application of Lemma 3.1 yields $\lambda' = (\lambda'_1, \ldots, \lambda'_k) \in \mathbb{R}^k_{>0}$ such that the simplex $C_{x,u,\lambda'}$ is contained in P. By Lemma 1.2 there exists $\epsilon = (\epsilon_1, \ldots, \epsilon_k) \in \mathbb{R}^k_{<0}$ such that $C_{x,u,\epsilon} \subseteq C_{x,u,\lambda} \cap C_{x,u,\lambda'}$.

Since g, as well as mg, vanish over S, then f vanishes over $C_{x,u,\epsilon}$. Therefore, $C_{x,u,\epsilon} \subseteq Zf = F$. From this we obtain $\emptyset \neq \operatorname{relint} C_{x,u,\epsilon} \cap F \subseteq \operatorname{relint} C_{x,u,\lambda} \cap F$. Since F is a face of S, by Lemma 1.1 $C_{x,u,\lambda} \subseteq F$, which contradicts our assumption $C_{x,u,\lambda} \not\subseteq F$.

This completes the proof of the claim, as well as of the (i) \Rightarrow (ii) direction of Theorem 2.4.

4. Proof of Theorem 2.4 (II) \Rightarrow (I)

Lemma 4.1. Let $P \subseteq \mathbb{R}^n$ be a rational polyhedron and $X \subseteq P$ a closed set. If $\eta: P \to \mathbb{R}^2$ is a \mathbb{Z} -map such that $\eta(X)$ has a rationally outgoing 1-tangent, then for some $k \in \{1, ..., n-1\}$, X has a rationally outgoing k-tangent.

Proof. Let $u \in \mathbb{R}^2$ be a rationally outgoing 1-tangent of $\eta(X)$ at x. Since u is outgoing there exists $\epsilon > 0$ such that both vertices of the segment $\operatorname{conv}(x, x + \epsilon u)$ are rational, and

(5)
$$\operatorname{conv}(x, x + \epsilon u) \cap \eta(X) = \{x\}.$$

By Lemma 1.4, both $\eta^{-1}(\{x\})$ and $\eta^{-1}(\operatorname{conv}(x, x + \epsilon u))$ are rational polyhedra contained in *P*. By Lemmas 1.3 and 1.5, there exists a regular triangulation Δ of *P* such that η is (affinely) linear on each simplex of Δ and

(6)
$$\eta^{-1}(\{x\}) = \bigcup \{R \in \Delta \mid R \subseteq \eta^{-1}(\{x\})\},\$$

(7)
$$\eta^{-1}(\operatorname{conv}(x, x + \epsilon u)) = \bigcup \{ U \in \Delta \mid U \subseteq \eta^{-1}(\operatorname{conv}(x, x + \epsilon u)) \}$$

The rest of the proof is framed in three steps.

Step 1: Let x_1, x_2, \ldots be a sequence of elements of $\eta(X)$ determining the rationally outgoing 1-tangent u of $\eta(X)$ at x. There exists $T \in \Delta$ such that the set $\{i \mid x_i \in \eta(T \cap X)\}$ is infinite. The compactness of T yields a sequence z_1, z_2, \ldots in T such that $\eta(z_1), \eta(z_2), \ldots$ is a subsequence of x_1, x_2, \ldots and $z = \lim_{i \to \infty} z_i$ exists. Since η is continuous and $X \cap T$ is closed, we have $\eta(z) = x$ and $z \in X \cap T$. Since η is (affine) linear on T, there is a $2 \times n$ integer matrix A and a vector $b \in \mathbb{Z}^2$ such that

$$\eta(y) = Ay + b$$
 for each $y \in T$.

Step 2: We claim that there exists $k \in \{1, ..., n\}$ together with k orthogonal unital vectors $w_1, ..., w_k \in \mathbb{R}^n$ such that

- (i) $Aw_k \neq 0$, and
- (ii) $Aw_j = 0$ for each j < k,

(iii) $w = (w_1, \ldots, w_k)$ is a k-tangent of X at z determined by a subsequence of z_0, z_1, \ldots

The vectors w_1, \ldots, w_k are constructed by the following inductive procedure: Basis Step:

From $\eta(z_i) - \eta(z) \notin \mathbb{R}u$ it follows that $z_i \neq z$ for all *i*, whence each vector $z_i^1 = (z_i - z)/||z_i - z||$ is well defined. Without loss of generality we can assume that z_0^1, z_1^1, \ldots tends to some unit vector w_1 . (If not, using the compactness of (n-1)-dimensional sphere of radius 1 we can take a converging subsequence of z_0^1, z_1^1, \ldots , and call w_1 its limit.) Observe that w_1 is a 1-tangent of X at z. If $Aw_1 \neq 0$ then $w = w_1$ proves the claim. Otherwise we proceed inductively.

Inductive Step:

Suppose we have obtained for some l, an l-tangent $w(l) = (w_1, \ldots, w_l)$ of X at z, and $Aw_i = 0$ for each $i \in \{1, \ldots, l\}$. Observe that l < n. (For otherwise, since for each $1 \leq i < j \leq l$, the vectors w_i and w_j are pairwise orthogonal, then A is the zero matrix, which is contradicts $Az + b = \eta(z) \neq \eta(z_i) = Az_i + b$.) Since $Aw_i = 0$ for each $i \in \{1, \ldots, l\}$, then

$$A(z+\delta_1w_1+\cdots+\delta_lw_l)+b=A(z)+b=\eta(z)\neq\eta(z_i) \text{ for each } \delta_1,\ldots,\delta_l\in\mathbb{R}.$$

It follows that $z_i - z \notin \mathbb{R}w_1 + \cdots + \mathbb{R}w_l$, and the vectors

$$z_i^{l+1} = \frac{z_i - z - p_l(z_i - z)}{||z_i - z - p_l(z_i - z)||}$$

are well defined. Taking, if necessary, a subsequence of the z_i and denoting by z_j its *j*th element, we can further assume that $\lim_{j\to\infty} z_j^{l+1} = w_{l+1}$ for some vector w_{l+1} .

By construction, the unit vector w_{l+1} is orthogonal to each w_j with $j \leq l$, and $w(l+1) = (w_1, \ldots, w_l, w_{l+1})$ is an (l+1)-tangent of X at z. If $Aw_{l+1} \neq 0$ we fix k = l+1 and w = w(l+1) is a k-tangent satisfying the properties of the claim. If not, we proceed inductively. This proves the claim and completes Step 2.

Step 3: Let $w = (w_1, \ldots, w_k)$ be the k-tangent of X at z obtained in Step 2. We will prove that w is rationally outgoing.

Since w is also a k-tangent of $X \cap T$ at z, by Lemma 3.1, there exists a k-tuple $\gamma = (\gamma_1, \ldots, \gamma_k) \in \mathbb{R}^k_{>0}$ such that $C_{z,w,\gamma} \subseteq T$. Since $Aw_j = 0$ for each j < k then $\eta(y) = \eta(z) = x$ for each $y \in C_{z,w(k-1),\gamma(k-1)}$. We can write

$$0 \neq Aw_{k} = \lim_{i \to \infty} Az_{i}^{k} = \lim_{i \to \infty} A\left(\frac{z_{i} - z - \mathbf{p}_{k-1}(z_{i} - z)}{||z_{i} - z - \mathbf{p}_{k-1}(z_{i} - z)||}\right)$$
$$= \lim_{i \to \infty} \frac{A(z_{i}) - A(z)}{||z_{i} - z - \mathbf{p}_{k-1}(z_{i} - z)||}$$
$$= \lim_{i \to \infty} \frac{\eta(z_{i}) - \eta(z)}{||z_{i} - z - \mathbf{p}_{k-1}(z_{i} - z)||} \cdot \frac{||\eta(z_{i}) - \eta(z)||}{||\eta(z_{i}) - \eta(z)||}$$
$$= \lim_{i \to \infty} \frac{\eta(z_{i}) - \eta(z)}{||\eta(z_{i}) - \eta(z)||} \cdot \frac{||\eta(z_{i}) - \eta(z)||}{||z_{i} - z - \mathbf{p}_{k-1}(z_{i} - z)||} \cdot$$

Since $0 \neq u = \lim_{i \to \infty} (\eta(z_i) - \eta(z))/||\eta(z_i) - \eta(z)||$, then for some c > 0 we can write

$$c = \lim_{i \to \infty} \frac{||\eta(z_i) - \eta(z)||}{||z_i - z - \mathbf{p}_{k-1}(z_i - z)||}$$
 and $Aw_k = cu$

8

Now, let us set $\lambda_j = \gamma_j$ for j < k, and $\lambda_k = \min\{\gamma_k, \epsilon/c\}$. Then $C_{z,w,\lambda} \subseteq C_{z,w,\gamma} \subseteq T$. For any $y \in C_{z,w,\lambda}$ there is $0 \le \delta \le \lambda_k \le \epsilon/c$ with

(8)
$$\eta(y) = A(z) + \delta A(w_k) + b = \eta(z) + \delta cu = x + \delta cu.$$

It follows that

(9)
$$\eta(C_{z,w,\lambda}) \subseteq \operatorname{conv}(x, x + \epsilon u).$$

To conclude the proof, let S be the smallest face of T such that $C_{z,w,\lambda}$ is contained in S. By (7) and (9), $S \subseteq \eta^{-1}(\operatorname{conv}(x, x + \epsilon u))$. By (6), $S \cap \eta^{-1}(\{x\})$ is a union of faces of S. The linearity of η on S ensures that $S \cap \eta^{-1}(\{x\})$ is convex, whence a face of S.

Letting $F = S \cap \eta^{-1}(\{x\})$, from (5), it follows that $S \cap X = F \cap X$. Moreover by (8), $\eta(z + \lambda_1 w_1 + \dots + \lambda_k w_k) = \eta(z) + \lambda_k cu \neq x$. Then $C_{z,w,\lambda} \not\subseteq F$. We have shown that the k-tangent $w = (w_1, \dots, w_k)$ is rationally outgoing. This concludes Step 3 and completes the proof of the lemma.

Proof of Theorem 2.4: (ii) \Rightarrow (i). By way of contradiction, let $f, g \in \mathcal{M}([0,1]^n)$ be such that $f \upharpoonright_X$ does not belong to the ideal generated by $g \upharpoonright_X$ and that $f \upharpoonright_X$ belongs to a maximal ideal of $\mathcal{M}(X)$ iff $g \upharpoonright_X$ does. Let A be the subalgebra of $\mathcal{M}(X)$ generated by $f \upharpoonright_X$ and $g \upharpoonright_X$. By [4, 4.1], A is not strongly semisimple. Let the map $\eta: X \to [0,1]^2$ be defined by $\eta = (f \upharpoonright_X, g \upharpoonright_X)$. By [9, 3.6], $A \cong \mathcal{M}(\eta(X))$, whence $\mathcal{M}(\eta(X))$ is not strongly semisimple. By Theorem 2.3, $\eta(X)$ has a rationally outgoing 1-tangent. By Lemma 4.1, for some $k \in \{1, \ldots, n-1\} X$ has a rationally outgoing k-tangent.

Acknowledgment: I would like to thank the valuable comments and suggestions made by Daniele Mundici on previous drafts of this paper.

References

- R.I.Bot, S.M. Grad, G.Wanka: Duality in vector optimization Springer-Verlag, New York, 2009.
- [2] H. Bouligand: Sur les surfaces dépourvues de points hyperlimites Ann. Soc. Polon. Math. vol 9 (1930), 32–41.
- [3] M. Busaniche, D. Mundici: Geometry of Robinson consistency in Łukasiewicz logic Ann. Pure Appl. Log. vol 147 (2007), 1–22.
- [4] M. Busaniche, D. Mundici: Bouligand-Severi tangents in MV-algebras Rev. Mat. Iberoamericana (To appear). arXiv:1204.2147
- [5] R. Cignoli, I.M.L. D'Ottaviano, D. Mundici: Algebraic Foundations of many-valued Reasoning Kluwer Academic Publishers, Dordrecht, Trends in Logic vol. 7, 2000.
- [6] E. Dubuc, Y. Poveda: Representation theory of MV-algebras Ann. Pure Appl. Log. vol 161 (2010), 1024–1046.
- [7] G. Ewald: Combinatorial convexity and algebraic geometry Springer-Verlag, New York, Grad. Texts in Math. vol. 168, 2000.
- [8] B. S. Mordukhovich: Variational Analysis and Generalized Differentiation I: Basic Theory Springer-Heidelberg, 2006.
- [9] D. Mundici: Advanced Łukasiewicz calculus and MV-algebras Springer-Verlag, Berlin, New York, Trends in Logic vol. 35, 2011.
- [10] C.P. Rourke and B.J. Sanderson: Introduction to piecewise-linear topology Springer-Verlag, New York, 1972.
- [11] F. Severi: Conferenze di geometria algebrica (Raccolte da B. Segre), Stabilimento tipolitografico del Genio Civile, Roma, 1927, and Zanichelli, Bologna, 1927–1930.
- [12] F. Severi: Su alcune questioni di topologia infinitesimale Ann. Soc. Polon. Math. vol 9 (1931), 97–108.

L.M. CABRER

[13] J.R. Stallings: Lectures on Polyhedral Topology Tata Institute of Fundamental Research, Mumbay, 1967.

(l.cabrer@disia.unifi.it) LEONARDO MANUEL CABRER: DEPARTMENT OF COMPUTER SCIENCE, STATISTICS AND APPLICATIONS "GIUSEPPE PARENTI", UNIVERSITY OF FLORENCE, VIALE MORGAGNI 59 – I-50134, FLORENCE - ITALY

10