Upper Bound for Critical Probability of Site Percolation on Triangular Lattice is $1/2^{\frac{3}{2}}$

Marko Puljic*

Center for Computational Science at Tulane University

(Dated: August 24, 2018)

In triangular lattice, the upper bound for critical probability of site percolation is $\frac{1}{2^{\frac{3}{2}}} \approx 0.3535$.

I. INTRODUCTION AND DESCRIPTION OF \mathbb{Z}^2

In site percolation, vertices (sites) of a graph are open with probability p, and there is the smallest $p=p_H$, critical p, for which open vertices form an open path the long way across a graph, so a vertex at the origin is a part of an infinite connected open vertex set, [3]. Smirnov found that for triangular lattice $p_H=1/2$, [1], but there is the traversal, from the origin upwards, so that an infinite connected open vertex set exists for $p_H=1/2^{\frac{3}{2}}\approx 0.3535$.

In finite graph \mathbb{Z}_k^2 , 2 pairs of opposite arcs are k edges away from the vertex at origin, Fig. 1. Basis $\mathcal{B}(\mathbb{Z}^2)$ of edges and the integers a_i assign the place to vertex $\mathbf{v} \in \mathbb{Z}_k^2$:

$$\mathbf{v}_0 = 0 = \text{vertex at origin of } \mathbb{Z}_k^2 \subset \mathbb{Z}^2$$

$$\mathcal{B}\left(\mathbb{Z}^d\right) = \{\uparrow_1, \uparrow_2, \uparrow_3, ..., \uparrow_d\}, \quad -\uparrow_1 = \downarrow_1 \& -\uparrow_2 = \downarrow_2$$

$$\mathbf{v} = a_1 \uparrow_1 + a_2 \uparrow_2, \ \mathbf{v}_0 \to \mathbf{v} = \text{path from } \mathbf{v}_0 \text{ to } \mathbf{v}$$

$$|\mathbf{v}_0 \to \mathbf{v}| = ||\mathbf{v}|| = |a_1(\mathbf{v})| + |a_2(\mathbf{v})|$$

Places of neighbors of vertex $\mathbf{v} \in \mathbb{Z}^2$ can be partition into 2 up-step neighbors traversed via \uparrow_1 and \uparrow_2 and 2 downstep neighbors traversed via \downarrow_1 and \downarrow_2 . For \uparrow_1 , there is matching \downarrow_1 , and for \uparrow_2 , there is matching \downarrow_2 :

$$\mathcal{N}(\mathbf{v}, \mathbb{Z}^2)$$
 = neighbors of \mathbf{v}
 $\mathcal{N}_u(\mathbf{v}, \mathbb{Z}^2)$ = up-step neighbors of $\mathbf{v} = {\mathbf{v} + \uparrow_1, \mathbf{v} + \uparrow_2}$




FIG. 1: In arcs, $\sum_{i} |a_{i}(\mathbf{v})| = 4$ (outside dotted curve). In the convenient pair of opposite arcs $\{\mathcal{A}_{4}(\mathbb{Z}^{2}), \mathcal{A}_{-4}(\mathbb{Z}^{2})\}_{1}$, $\mathcal{A}_{4}(\mathbb{Z}^{2}) = \{4\uparrow_{1}, 3\uparrow_{1}+1\uparrow_{2}, 2\uparrow_{1}+2\uparrow_{2}, 1\uparrow_{1}+3\uparrow_{2}, 4\uparrow_{2}\}.$

 \mathbf{v}_0 is k edges away from the arcs and it can be any vertex in \mathbb{Z}^2 . Two pairs of opposite arcs in \mathbb{Z}^2_k look the same and any pair, by rotation of \mathbb{Z}^2 , can be a convenient pair $\{\mathcal{A}_k(\mathbb{Z}^2), \mathcal{A}_{-k}(\mathbb{Z}^2)\}_1$:

$$\left\{ \mathcal{A}_k(\mathbb{Z}^2), \mathcal{A}_{-k}(\mathbb{Z}^2) \right\}_1 = \text{one of 2 pairs of opposite arcs}$$

 $\mathcal{A}_k(\mathbb{Z}^2) \cap \mathcal{A}_{-k}(\mathbb{Z}^2) = \emptyset$

$$\mathcal{A}_k(\mathbb{Z}^2) = \left\{ \begin{array}{c} \mathbf{v} : \|\mathbf{v}\| = k \ \& \ a_1(\mathbf{v}), a_2(\mathbf{v}) \ge 0 \\ \text{or} \\ \mathbf{v} \text{ built with } \uparrow_1 \ \& \ \uparrow_2 \text{ edges only} \end{array} \right\}$$

The shortest traversal from \mathbf{v}_0 to $\mathcal{A}_k(\mathbb{Z}^2)$ is a traversal via vertices in $\mathcal{N}_u(\mathbf{v}, \mathbb{Z}^2)$, which are one edge closer to $\mathcal{A}_k(\mathbb{Z}^2)$:

$$1^{st} : \mathcal{A}_{1} \left(\mathbb{Z}^{2} \right) = \{ \uparrow_{1}, \uparrow_{2} \}$$

$$2^{nd} : \mathcal{A}_{2} \left(\mathbb{Z}^{2} \right) = \bigcup_{\mathbf{v} \in \mathcal{A}_{1}(\mathbb{Z}^{2})} \mathcal{N}_{u}(\mathbf{v}, \mathbb{Z}^{2})$$

$$\vdots$$

$$k^{th} : \mathcal{A}_{k} \left(\mathbb{Z}^{2} \right) = \bigcup_{\mathbf{v} \in \mathcal{A}_{k-1}(\mathbb{Z}^{2})} \mathcal{N}_{u}(\mathbf{v}, \mathbb{Z}^{2})$$

II. TRIANGULAR LATTICE PERCOLATION

Lattice \mathbb{Z}^2 is percolating when the open vertices form an open path connecting \mathbf{v}_0 with the vertices in the opposite sides of \mathbb{Z}_k^2 and $k \to \infty$, so we need to know the number of paths connecting \mathbf{v}_0 and vertices in $\mathcal{A}_k(\mathbb{Z}^2)$:

$$p = \text{probability of vertex being open}$$

$$\psi\left(\mathbb{Z}^2, p\right) = \text{number of percolating paths in } \mathbb{Z}^2$$

 \mathbb{Z}^2 is embedded in \mathbf{T} , which has two pairs of opposite sides and two definitions of $\mathcal{B}(\mathbf{T})$, Fig. 2:

 $\mathbf{T} = \text{triangular lattice with embedded } \mathbb{Z}^2$

$$\mathcal{B}(\mathbf{T}) = \begin{cases} \uparrow_{1}, \uparrow_{2}, \uparrow_{1} + \uparrow_{2} = \uparrow_{1,2} = -\downarrow_{1,2} \\ \text{or} \\ \uparrow_{1}, \uparrow_{2}, \uparrow_{1} + \downarrow_{2} = \rightarrow_{1,2} = -\leftarrow_{1,2} \end{cases}$$
$$\|\uparrow_{1,2}\| = \|\rightarrow_{1,2}\| = 1$$
$$\mathcal{N}_{u}(\mathbf{v}, \mathbf{T}) = \begin{cases} \mathbf{v} + \uparrow_{1}, \mathbf{v} + \uparrow_{2}, \mathbf{v} + \uparrow_{1,2} \\ \text{or} \\ \mathbf{v} + \uparrow_{1}, \mathbf{v} + \uparrow_{2} \end{cases}$$

^{*}Marko Puljic: mpuljic@tulane.edu




FIG. 2: Triangular lattice is built by adding $\uparrow_1+\uparrow_2$ and $\downarrow_1+\downarrow_2$ (left) or by adding $\uparrow_1+\downarrow_2$ and $\downarrow_1+\uparrow_2$ (right) to each $\mathbf{v}\in\mathbb{Z}^2$.

Arc $\mathcal{A}_k(\mathbf{T})$, k edges away from $\mathbf{v}_0 \in \mathbf{T}$, contains the vertices in $\mathcal{A}_{k+i}(\mathbb{Z}^d)$ traversed via $\mathcal{N}_u(\mathbf{v}, \mathbf{T})$:

$$\mathcal{A}_{k}(\mathbf{T}) = \bigcup_{\mathbf{v} \in \mathcal{A}_{k-1}(\mathbf{T})} \mathcal{N}_{u}(\mathbf{v}, \mathbf{T}) : \mathcal{A}_{1}(\mathbf{T}) = \{\uparrow_{1}, \uparrow_{2}, \uparrow_{1,2}\}$$

$$n_{k}(\mathcal{A}_{k+i}(\mathbb{Z}^{2}), \mathbf{T}) = \text{number of paths to } \mathcal{A}_{k+i}(\mathbb{Z}^{2})$$

$$n_k(\mathcal{A}_k(\mathbf{T})) = \sum_i n_k \left(\mathcal{A}_{k+i}\left(\mathbb{Z}^2\right), \mathbf{T}\right)$$

For vertices in arc $\mathcal{A}_1(\mathbf{T})$, 2 paths end in $\mathcal{A}_1(\mathbb{Z}^2)$ and 1 path ends in $\mathcal{A}_2(\mathbb{Z}^2)$:

$$\mathcal{A}_1(\mathbf{T}) = \{\mathbf{v}_0 + \uparrow_1, \mathbf{v}_0 + \uparrow_2, \mathbf{v}_0 + \uparrow_{1,2}\} \Rightarrow \begin{matrix} 2 & \rightarrow_1 \\ 1 & \rightarrow_2 \end{matrix} \Rightarrow \begin{matrix} |\rightarrow_1| = 2 \\ |\rightarrow_2| = 1 \end{matrix}$$

 \rightarrow_1 = path from \mathbf{v}_0 to a vertex in embedded $\mathcal{A}_1(\mathbb{Z}^2)$ $|\rightarrow_1|$ = number of paths from \mathbf{v}_0 to $\mathcal{A}_1(\mathbb{Z}^2)$

From $A_1(\mathbf{T})$, the number of paths doubles for up-steps from $\mathcal{A}_1(\mathbb{Z}^2)$ to $\mathcal{A}_2(\mathbb{Z}^2)$ and from $\mathcal{A}_2(\mathbb{Z}^2)$ to $\mathcal{A}_3(\mathbb{Z}^2)$. The number of paths does not change for up-steps from $\mathcal{A}_1(\mathbb{Z}^2)$ to $\mathcal{A}_3(\mathbb{Z}^2)$ and from $\mathcal{A}_2(\mathbb{Z}^2)$ to $\mathcal{A}_4(\mathbb{Z}^2)$:

$$\mathcal{A}_{2}(\mathbf{T}) = \bigcup_{\mathbf{v} \in \mathcal{A}_{1}(\mathbf{T})} \mathcal{N}_{u}(\mathbf{v}, \mathbf{T}) \Rightarrow \begin{vmatrix} 2 | \rightarrow_{1} | & \cdots \rightarrow_{2} \\ 1 | \rightarrow_{1} | & \cdots \rightarrow_{3} \\ 2 | \rightarrow_{2} | & \cdots \rightarrow_{3} \\ 1 | \rightarrow_{2} | & \cdots \rightarrow_{4} \end{vmatrix} \begin{vmatrix} \cdots \rightarrow_{2} | = 1 \cdot 2^{2} \\ \cdots \rightarrow_{3} | = 2 \cdot 2^{1} \\ \cdots \rightarrow_{4} | = 1 \cdot 2^{0} \end{vmatrix}$$

 $|.\rightarrow_2|$ =number of paths ending in $\mathcal{A}_2(\mathbb{Z}^2)$

When counting the paths from \mathbf{v}_0 to the arcs $\mathcal{A}_k(\mathbb{Z}^2)$, $\mathcal{A}_{k+1}(\mathbb{Z}^2), ..., \mathcal{A}_{2k}(\mathbb{Z}^2)$, the coefficients c_{i_k} follow the rule generated by Pascal's triangle, [4]:

$$c_{i_k} \cdot 2^{k-i} = \text{number of paths to } \mathcal{A}_{k+i} \left(\mathbb{Z}^2 \right) \text{ after } k^{th} \text{ step } 1^{st} : 1, 1 \Rightarrow 1 \cdot 2^1 + 1 \cdot 2^0 \text{ paths to } \mathcal{A}_1(\mathbf{T})$$

$$2^{nd}$$
: 1, 2, 1 \Rightarrow 1 · 2² + 2 · 2¹ + 1 · 2⁰ paths to $\mathcal{A}_2(\mathbf{T})$

$$3^{rd}$$
: $1, 3, 3, 1 \Rightarrow 1 \cdot 2^3 + 3 \cdot 2^2 + 3 \cdot 2^1 + 1 \cdot 2^0$ paths

$$4^{th}$$
: $1 = c_{0_4}, 4 = c_{1_4}, 6 = c_{2_4}, 4 = c_{3_4}, 1 = c_{4_4}$

$$\sum_{i=0}^{i=k} \binom{k}{i} \cdot 2^{k-i} = \text{number of paths to } \mathcal{A}_k(\mathbf{T})$$

$$\binom{k}{i} \cdot 2^{k-i} = \text{number of paths to } \mathcal{A}_{k+i} \left(\mathbb{Z}^2 \right)$$

When odd or even $k \to \infty$,

$$p_H(\mathbf{T}) \leq \min p \text{ for which } \binom{k}{i} \cdot 2^{k-i} \cdot p^k \geq 1$$

$$\binom{k}{\frac{k-1}{2}} \cdot 2^{k-\left(\frac{k-1}{2}\right)} = \text{number of paths to } \mathcal{A}_{k+\frac{k-1}{2}}(\mathbb{Z}^2)$$
or
$$\binom{k}{\frac{k}{2}-1} \cdot 2^{k-\left(\frac{k}{2}-1\right)} = \text{number of paths to } \mathcal{A}_{k+\frac{k}{2}-1}(\mathbb{Z}^2)$$

$$\psi(\mathbf{T}, p) = 2^{\frac{1}{2}} \lim_{k \to \infty} \left(2^{\frac{1}{2}} \cdot {k \choose \frac{k-1}{2}}^{\frac{1}{k}} \cdot p \right)^k$$
or

From
$$\mathcal{A}_1(\mathbf{T})$$
, the number of paths doubles for up-steps from $\mathcal{A}_1(\mathbb{Z}^2)$ to $\mathcal{A}_2(\mathbb{Z}^2)$ and from $\mathcal{A}_2(\mathbb{Z}^2)$ to $\mathcal{A}_3(\mathbb{Z}^2)$. The number of paths does not change for up-steps from $\mathcal{A}_1(\mathbb{Z}^2)$ to $\mathcal{A}_3(\mathbb{Z}^2)$ and from $\mathcal{A}_2(\mathbb{Z}^2)$ to $\mathcal{A}_4(\mathbb{Z}^2)$:

$$\mathcal{A}_2(\mathbf{T}) = \bigcup_{\mathbf{v} \in \mathcal{A}_1(\mathbf{T})} \mathcal{N}_u(\mathbf{v}, \mathbf{T}) \Rightarrow \begin{cases} 2|\rightarrow_1| & ... \rightarrow_2 \\ 1|\rightarrow_1| & ... \rightarrow_3 \\ 2|\rightarrow_2| & ... \rightarrow_4| = 1 \cdot 2^0 \end{cases} \Rightarrow p_H(\mathbf{T}) \leq \begin{cases} \lim_{k \to \infty} \frac{1}{2^{\frac{1}{2}} \cdot \left(\frac{k}{k-1}\right)^{\frac{1}{k}}} \\ \text{or} \\ \lim_{k \to \infty} \frac{1}{2^{\frac{1}{2}} \cdot \left(\frac{k}{k-1}\right)^{\frac{1}{k}}} \\ \text{or} \\ \lim_{k \to \infty} \frac{1}{2^{\frac{1}{2}} \cdot \left(\frac{k}{k-1}\right)^{\frac{1}{k}}} \\ \lim_{k \to \infty} \frac{1}{2^{\frac{1}{2}} \cdot \left(\frac{k}{k-1}\right)^{\frac{1}{k}}} \end{cases} \Rightarrow 0.3535$$

$$... \rightarrow_2 = \text{path ending in } \mathcal{A}_2(\mathbb{Z}^2)$$

$$... \rightarrow_2 = \text{path ending in } \mathcal{A}_2(\mathbb{Z}^2)$$

$$\lim_{k\to\infty} \left(\frac{k}{\frac{k-1}{2}}\right)^{\frac{1}{k}} \approx 2$$
, [2][5].

- [1] Stanislav Smirnov. Critical percolation in the plane: Conformal invariance, cardy's formula, scaling limits. Comptes Rendus de l'Académie des Sciences-Series I-Mathematics, 333(3):239–244, 2001.
- [2] Herbert Robbins. A remark on stirling's formula. The American Mathematical Monthly, 62(1):26–29, 1955.
- [3] Path is a walk via edges visiting each vertex only once.
- [4] From The On-Line Encyclopedia of Integer Sequences at http://oeis.org.
- [5] Limits at http://www.wolframalpha.com.