A CANONICAL PARTITION OF THE PRIMES OF LOGIC FUNCTIONS

SIDNIE FEIT

Copyright 2012, 2013, 2014 Sidnie Feit

1. Introduction

This paper deals with Boolean functions, called *logic functions* in circuit and VLSI theory. A Boolean function in n binary variables,

$$f(x_1,\ldots,x_n)\to\{0,1\},\ x_i\in\{0,1\},\ i=1,\ldots,n$$

assigns a value of 1 (TRUE) or 0 (FALSE) to each of the 2^n n-tuples that make up its domain. A Boolean function can be represented by a formula built using AND, OR and NOT operations. In this paper, '*' represents AND, '+' represents OR, and x' represents the negation NOT x (also called the complement of x).

An AND of variables (some of which may be negated) is called a *product*, and an OR of products is called a *sum-of-products*. Every Boolean function can in fact be represented by a sum-of-products formula, such as

(1)
$$f = x_2 * x_4' * x_8 * x_9 + x_1 * x_5 * x_8' * x_{10} * x_{11} + x_1 * x_6' * x_{10}' * x_{11}$$

A sum-of-products representation of a Boolean function is far from unique. Any Boolean function can be represented by an enormous number of distinct, logically equivalent sums-of-products.

Definition 1.1. A product X is an *implicant* for Boolean function f if the assignment of variable values that make product X = 1 also makes f = 1.

For example, if

$$(2) X = x_2 * x_4' * x_8 * x_9 * x_{12}$$

X = 1 if and only if $x_4 = 0$ and the rest of the variables equal 1. Substituting these values into the sum-of-products in (1) results in f = 1. Hence X is an implicant for f.

Definition 1.2. A prime is an implicant that no longer is an implicant if any factor is removed.

¹The book [6] by Hachtel and Somenzi provides a clear presentation of the role of Logic (i.e., Boolean) functions in VLSI theory.

X in (2) is not prime, because if we remove factor x_{12} , then $Y = x_2 * x'_4 * x_8 * x_9$ is an implicant. Note that the set of factors of any implicant must include the set of factors of a prime implicant. In this example, Y is a prime.

The set consisting of all of the primes for function f is called its *complete set of primes*. The sum of all of the primes in the complete set for f is a canonical (but usually extremely inefficient) representation of f.

There is a classic problem associated with the primes of a function.

```
Given a Boolean function f(x_1, ..., x_n) and a cost function C such that C assigns a positive cost to each Boolean product, C is additive, that is, given a set of Boolean products X_1, ..., X_n, Cost(X_1, ..., X_n) = Cost(X_1) + \cdots + Cost(X_n), find a minimum-cost set of primes whose sum is equal to f.
```

This problem originated in the world of logic circuits, and initially was aimed at finding a minimum-cost sum of products equal to a given logic (Boolean) function. Quine[9] proved that a minimum-cost sum-of-products for a Boolean function f must consist of a sum of primes if any definition of cost is used in which the addition of a single factor to a formula increases the cost.

Here, we look for minimum-cost sums-of-primes for somewhat more general cost functions. All of the cost functions in this paper are assumed to be positive and additive. A set of primes whose sum is equivalent to f will be called a *basis* for f. The cost of a basis is equal to the sum of the costs of the members of the basis. A solution to the classic problem is given by a basis whose cost is minimal with respect to the given cost function.

This paper presents algorithms that partition the complete set of primes for f into N+2 canonical disjoint subsets such that for any positive additive cost function C, solving the minimization problem consists of solving N separate minimization problems. The algorithms calculate the following:

- (1) Essential Primes, which must be part of any basis for f,
- (2) $Unnecessary\ Primes$ that cannot be part of a minimum-cost basis for f for any positive additive cost function,
- (3) N unique disjoint sets of primes, PS_1, \ldots, PS_N associated with 'covering' tables TS_1, \ldots, TS_N such that a minimum-cost basis for a positive, additive cost function C consists of

```
\{Essential\ Primes\} \cup QS_1(C) \cup \cdots \cup QS_N(C)
where QS_i(C) \subset PS_i and QS_i is a minimum-cost 'cover' for PS_i,
as determined by C and TS_i.
```

Covering relationships are determined by an operation, $Cascade(RS, TS) \longrightarrow T$, where RS is a set of products and TS is the table of covering relationships associated with a set of primes PS. RS covers PS if and only if Cascade(RS, TS) is an empty table. Thus, $QS_i \subset PS_i$ is a minimum-cost set such that $Cascade(QS_i, TS_i)$ is empty.

Tables and covering relationships in this paper are somewhat unusual due to the iterative nature of the algorithms. QS_{i+1} covers PS_{i+1} relative to the fact that all products already

covered by the primsets $PS_1, \dots PS_i$ have been removed from table TS_{i+1} (and from the problem).

The paper also contains results that solve or simplify the problem of finding QS_i when TS_i satisfies certain conditions. If a subproblem satisfies the hypotheses of Theorem 6.1, the Span Basis Theorem, a solution consists of one easily identified prime. If a subproblem satisfies the hypothesis of Theorem 6.2, the Independent Prime Decomposition Theorem, the subproblem can be decomposed into smaller subproblems, and at least one of these is solved by an easily identified prime.

The input data for the partitioning algorithms consists of the primes for f, the products generated by consensus combinations of the primes, and a table $triples(f) = \{(r, i, j)\}$ that expresses the covering relationships between products.

A key new concept in this paper is that of Ancestor Sets, whose elements are easily derived from the triples(f) table. The Ancestor Theorem that follows is an indispensable result in this paper.

Theorem 1.1 (Ancestor Theorem). If A is an Ancestor Set for f and C is a positive additive cost function, then every minimum-cost basis consists of the union of the Essential Primes, a minimum-cost cover QS(C) for the primes PS that belong to ancestor set A, and a minimum-cost cover for the set of primes that are not in A (and are not covered by the union of the Essential Primes and PS).

An Independent Ancestor Set is an Ancestor Set that does not contain any other Ancestor Set. The Independent Ancestor Sets are unique and disjoint, and each PS_i in the partition is equal to the set of primes in an Independent Ancestor Set A_i .

A straightforward algorithm applied to the *triples* table identifies a unique initial group of disjoint Independent Ancestor Sets, say, A_1, \ldots, A_k , with primesets PS_1, \ldots, PS_k . Because of the Ancestor Theorem, every minimum-cost basis for f must include minimum-cost covers for each primeset PS_i , $i = 1, \ldots, k$.

But then all products covered by the union of the Essential Primes and the PS_i can be removed from the triples table and from the partitioning problem. The remaining Independent Ancestor Sets are found by iterating the algorithm on the reduced triples table until the triples table is empty. A minimum-cost cover for each new Independent Ancestor Set A_j will be part of a minimum-cost basis for f.

If there are K products and the triples table has M rows, the partitioning algorithms execute with worst-case complexity $M * log_2(M) * K$. When there are some Essential Primes, the number of products and the number of rows in the table are reduced in size significantly.

The algorithms in this paper calculate the primes, the relevant consensus combinations of primes, the Independent Ancestor Sets A_i , the PS_i , and the tables TS_i . Although the algorithmic proofs are valid, the worst-case size of the data can be enormous. First of all, to get a sense of the number of primes that could be associated with a Boolean function, McMullen and Shearer [8] found the following upper bound and constructed a function that attained the upper bound:

4

If m equals the minimum number of products in a sum-of products that represents f, then the number of primes for f is less than or equal to $2^m - 1$.

Furthermore, the task of finding a minimum-cost $QS_i(C)$ can be framed as an integer programming problem, and a general integer programming problem can be NP-hard.

However, it is possible that new algorithms may be discovered that split up an oversized problem by extracting some Ancestor Sets from the problem. Also, the tables for some Ancestor Sets satisfy hypotheses that make it possible to find minimum-cost covers in linear or polynomial time.

But it is not surprising that much effort has been devoted to finding algorithms that lead to approximate solutions. The classic problem has been studied extensively using a geometric model that represents products as "cubes" and merges smaller cubes into larger ones. This model underlies various versions of the popular heuristic ESPRESSO algorithm [2], which originally was developed by Robert Brayton at IBM. ESPRESSO frequently is used to compute approximations to a minimum-cost basis. Sometimes an exact solution is sought, and in this case, the first step in ESPRESSO is an attempt to calculate the complete set of primes for f.²

2. The Consensus Identity and Consensus Operations

This section and the sections that follow include statements of the type

$$formula E = formula F.$$

In Boolean algebra, such a statement represents 'formula E is logically equivalent to formula F' or 'formula E represents the same Boolean function as formula F.' This paper follows the usual practice of using an equals sign to denote logical equivalence.

In the following, small letters such as x, y, z represent binary (0 or 1) logic variables. Capital letters, including P, Q, R, U, V, W, X, Y, Z will represent products whose factors are binary variables or their negations.

The standard consensus identity and the consensus operation of Boolean algebra play a major role in this paper. The consensus identity for binary variables x, y, and z, is simple:

$$x * y + x' * z = x * y + x' * z + y*z$$

Note that the products x * y and x' * z have one reversal variable — that is, one variable, x, that is not negated in one product and is negated, x', in the other. The identity also holds when we replace y and z with products Y and Z such that the pair Y and Z has no reversal variable.

$$x * Y + x' * Z = x * Y + x' * Z + Y*Z$$

Product Y * Z is called the *consensus* of x * Y with x' * Z, written

$$Y * Z = (x * Y \circ x' * Z).$$

²Another approach uses data structures called Binary Decision Diagrams (see [3]) represent a Boolean function as a rooted, directed, acyclic graph. These diagrams have led to advances in formal verification of circuits and logic minimization.

The consensus operation $(U \circ V)$ for products U and V is defined and valid only for products U and V that have exactly one reversal variable. The result consists of the product of all of the factors in U and V (with duplicates removed) except for the reversal factors. If $(U \circ V)$ and $(Y \circ W)$ are valid, then

$$U + V + Y + W = U + V + (UoV) + Y + W + (YoW)$$

If $((U \circ V) \circ (Y \circ W))$ is valid, then

$$U + V + Y + W = U + V + (U \circ V) + Y + W + (Y \circ W) + ((\mathbf{U} \circ \mathbf{V}) \circ (\mathbf{Y} \circ \mathbf{W}))$$

Definition 2.1. Suppose that $\{X_1, \dots X_k\}$ is a set of distinct products. Then

$$Consensus(X_1, \ldots, X_k)$$

equals $\{X_1,\ldots,X_k\}$ and all products generated by their valid consensus combinations.

The following procedure, here called *Loose Iterated Consensus*, is an orderly way to generate $Consensus(X_1, \ldots, X_k), k > 1$. Initially, start with the list of the distinct products, X_1, \ldots, X_k and set i = 1.

- (1) Increase i by 1. Compute the consensus of X_i with each prior product in the list. If a result X is new, append it to the list.
- (2) If X_i is not the last product, repeat starting with (1). If X_i is the last product, stop.

Then $Consensus(X_1, \ldots, X_k)$ equals the set of products in the final list,

$$X_1,\ldots,X_k,X_{k+1},\ldots,X_n.$$

The following Lemma follows from the consensus identity.

Lemma 2.1 (Consensus Lemma). Given a Boolean function f expressed as sum-of-products $X_1 + \cdots + X_k$, then the sum of the products in $Consensus(X_1, ..., X_k)$ is logically equivalent to f, and every product in $Consensus(X_1, ..., X_k)$ is an implicant for f.

Definition 2.2. Product Y is covered by $\{X_1, \ldots, X_k\}$ if $Y \in Consensus(X_1, \ldots, X_k)$.

2.1. Computing the Complete Set of Primes. The classic method of finding the complete set of primes associated with a Boolean function that is defined by a sum-of-products formula is based on a theorem proved by Brown [4] and presented in more modern language in many textbooks. The wording here differs slightly from the usual language.

One of the most basic identities in Boolean Algebra is used when computing the complete set of primes. If U, V, and W are products and V = U * W, then U + V = U + U * W = U. A special case is U + U = U.

Theorem 2.1 (Brown's Complete Set of Primes Theorem). A sum-of-products formula for f is the sum of the complete set of primes for the function if and only if

- (1) No product's factors are a subset of any other product's factors.
- (2) If the consensus of two products Y and Z in the sum exists, $X = (Y \circ Z)$, then either X = some product in the sum, or there is a product R in the sum such that the factors in R are a proper subset of the factors in X.

6

The classic *Iterated Consensus* algorithm for calculating the complete set of primes for a sum-of-products, is based on Brown's Theorem. Note that the Iterated Consensus of X_1, \ldots, X_m , here called **IT** (X_1, \ldots, X_m) , differs from $Consensus(X_1, \ldots, X_m)$. When computing **IT**, if the factors of X are a proper subset in any other product Z, product Z is removed.

Definition 2.3. Suppose that $f = X_1 + \cdots + X_m$ and there is no X_i whose factors are a subset of another product's factors. The *Iterated Consensus*, $\mathbf{IT}(X_1, \dots, X_m)$ is defined as follows. Starting with X_2 , compute the consensus of each product with each prior product. For each valid consensus result X,

- (1) If the factors in X are a proper subset of the factors in any other product(s) on the list, remove the product(s).
- (2) If the factors of any X_k on the list are a subset of the factors of X, discard X.
- (3) If no X_k is a subset of X, append X to the list.

The resulting sum satisfies the hypotheses of Brown's Theorem and hence the products are the complete set of primes for the function.³

At every step in performing the Iterated Consensus, the sum of the products on the list remains equivalent to the Boolean function f defined by the original sum-of-products. Thus the sum of the complete set of primes is equal to the original function f. The same set of primes will be generated from any sum-of-products that is equal to f. In particular, if the set of primes Q_1, \ldots, Q_m is a basis for f, then

$$\mathbf{IT}(Q_1,...,Q_m)$$
 = the complete set of primes for f .

2.2. ALL, the Consensus of the Complete Set of Primes. Once the complete set of primes of f is known, the relationships between the primes are hidden within the relationships between all of the products generated by computing consensus combinations of the primes. For example, we may have primes P, $\{Q_i\}$ with

$$P = (\ (Q_1 \circ Q_2) \circ (Q_8 \circ Q_{10})\)$$

where $X = (Q_1 \circ Q_2)$ and $Y = (Q_8 \circ Q_{10})$ are not primes. Note that if Q_1, Q_2, Q_8 , and Q_{10} belonged to a minimum-cost basis, then P could not belong to the minimum-cost basis since it already is covered.

To capture all relationships, after finding the primes, let

$$ALL = Consensus$$
 (complete set of primes).

That is, *All* is generated by performing a Loose Iterated Consensus, where only duplicate products are discarded.⁴ The next lemma follows immediately.

³Another method, which often is more efficient, is to calculate the primes by recursive use of Boole's Expansion Theorem. See [6], page 139.

⁴More efficient methods for calculating ALL can be devised, but this simple method suffices for the purpose of defining the set.

Lemma 2.2. Primes P_1, \ldots, P_k are a basis for f if and only if

$$Consensus(P_1, \ldots, P_k) = ALL.$$

As mentioned earlier, the number of products that need to be analyzed and the number of relationships between products are diminished if some of the products are *Essential*. Recall that product Y is covered by $\{X_1, \ldots, X_k\}$ if $Y \in Consensus(X_1, \ldots, X_k)$.

Definition 2.4. A prime P is Essential if P is not covered by the other primes.

All Essential Primes have to be part of any minimum-cost basis. Since their fixed positive cost is irrelevant to computing a minimum-cost basis, in this paper they are called *free*.

The Essential Primes can be calculated directly from any sum-of-primes formula for f using the following Theorem of Sasao [10].

Theorem 2.2. Suppose that function f is expressed as a sum of primes,

$$P + V_1 + \dots V_n$$
.

Then P is an essential prime if and only if P is not covered by

$$Consensus(P * V_1, \dots, P * V_n, (P \circ V_1), \dots, (P \circ V_n)).$$

2.3. Free Products, Non-Free Products, and Spans. The products generated by computing $Consensus(Essential\ Primes)$ are covered by the Essentials, and hence also can be viewed as free products. The set of all free products is denoted by $\langle Free \rangle$. Thus,

$$\langle Free \rangle = Consensus(Essential\ Primes).$$

For any positive additive cost function C, the Essential Primes are a minimum-cost cover for $\langle Free \rangle$. The set of free primes that are not Essential cannot belong to any minimum-cost basis and hence belong to the set of $Unnecessary\ Primes$.

Definition 2.5. Every prime that is not Essential is called a non-Essential prime. A non-Essential prime that is not free is called a non-free prime.

If all primes are free, the problem is solved because the unique minimum-cost basis for any positive additive cost function C consists of the Essential Primes. From now on, we assume that there are some non-free primes. The set of products of interest is

$$NONFREE$$
 = the non-free products in ALL

We will order and number NONFREE with the non-free primes first. It is convenient to number the non-free primes $\{P_k\}$ in order of increasing cost. (This will make it trivial to select a prime with the least cost from a set of equivalent primes.)

Definition 2.6. The set of non-free primes that turn out to be *Unnecessary* will be called *Surplus*.

Definition 2.7. The Span of a set of non-free products S is defined by

$$Span(S) =$$
the non-free products in $Consensus(< Free >, S)$.

Definition 2.8. Given a set of non-free products S, non-free product X is *covered* by Span(S) if $X \in Span(S)$.

Definition 2.9. A set of non-free primes R_1, \ldots, R_g for f is a *cover* for the non-free primes of f if

$$Span(R_1, ..., R_q) = NONFREE$$

Note that if $Span(R_1, \ldots, R_q)$ covers the non-free primes of f, then

Essential Primes,
$$R_1, \ldots, R_q$$

is a basis for f.

Clearly, any minimum-cost basis must consist of the Essential Primes and a minimum-cost cover, Q_1, \ldots, Q_k , for the non-free primes.

Lemma 2.3. Let P_1, \ldots, P_n be the non-free primes. Then each non-free P_k is covered by the span of the other non-free primes.

Proof. If P_k was not covered by the span of the other non-free primes it would have to be an Essential or other free prime, contradicting the assumption that it is non-free.

3. The triples(f) Table

Consensus relationships are used to build triples(f), a table that includes all useful covering relationships and is the starting point of all calculations.

The Essential Primes must be in any basis, so we assume that they have been chosen as the initial basis members and hence all free products are already covered.

A table entry, (r, i, j) corresponds to a consensus relationship, $X_r = (X_i \circ X_j)$. This relationship states that if X_i and X_j are covered, then X_r is covered. Note that:

- (1) When X_r is free, X_r already is covered by the Essentials so the relationship is irrelevant and is not included.
- (2) If both X_i and X_j are free, then X_r is free, so the relationship is not included.
- (3) If X_r and X_i are non-free and X_j is free, then, since X_j already is covered, X_r is covered whenever X_i is covered. This is tabulated as (r, i, 0).⁵ Thus, all free X_j products will be represented by 0s in the table, and all product indices in a table represent non-free products.
- (4) If X_r, X_i , and X_j all are non-free, then the entry is tabulated as (r, i, j).

The (r, i, 0) entries are called triples0 entries and the (r, i, j) entries are called triples3 entries.

In addition to the triples0 entries generated by consensus operations, there is a second set of valuable triples0 cover relationships. By definition, non-free prime P_k covers every product in $Span(P_k)$. When $Span(P_k)$ is maximal (see Definition 6.1) and closed (see Definition 6.2, then for each $X_r \in Span(P_k)$, we append (r, k, 0) to triples0.

⁵When taking the consensus of a non-free with a free, by convention, the free product will always be assumed to occupy the X_j position.

The calculations will produce many duplicate (r, i, 0) results. Efficient code can discard duplicates as they arise (for example, by using a sorted list and hashed list insertion).

It is convenient to refer to products by their indices. Thus, we can say that an (r, i, j) entry states that if i and j are covered then r is covered.

As a rule of thumb, a greater number of (r, i, j), j > 0) entries lead to slower execution, while more (r, i, 0) entries lead to faster execution. Fortunately, when there are free products, some (r, i, j), j > 0 entries are useless and can be discarded.

Specifically, if there is an (r, i, 0) entry, i covered implies r covered; thus, stating 'i AND j covered imply r covered' is redundant and any such (r, i, j) entry can be discarded. The same argument holds for (r, i, j) if there is an (r, j, 0) entry. An (r, i, j) entry that is not useless is called useful.

The triples(f) table consists of the union of the remaining entries in triples0 and triples3. The algorithm in Appendix A arranges the products in NONFREE in a favorable order and the algorithm in Appendix B can be used to generate triples. These algorithms are just examples, and can be used for problems of small size. More efficient algorithms are required for big problems.

Note that by the definition of Span, the set of non-free primes R_1, \ldots, R_g covers the non-free primes if and only if $Span(R_1, \ldots, R_g)$ covers every index in triples(f).

3.1. Inactive Products. An r-index that does not appear anywhere as an i or j does not cover anything, and is called *inactive*. After removing the entries for inactive r-indices, others may become inactive. An iteration clears out these entries, (including inactive primes) leaving only active indices. An inactive prime cannot be part of any minimum-cost basis, and belongs to Surplus.

From now on, we assume that useless and inactive entries have been removed from triples(f). Let triples denote the resulting table of useful, active products. Once the initial triples table has been defined, no further calculations of consensus products are needed. All needed information is in triples.

3.2. Cascades Relative to a Table. Given a set of products S and a *triples* table T, the operation Cascade(S,T) plays a key role in the algorithms used to discover the PS_i . The purpose of Cascade(S,T) is to

generate the list $\{S, \text{ all other products in } T \text{ covered by } S\}$, remove all indices in S or covered by S from T.

When the identity of the table T is clear from the current context, we can write Cascade(S) instead of Cascade(S,T).

The cascade process is iterative. If S covers additional indices S1 and $S \cup S1$ covers additional indices S2, then S covers $S \cup S1 \cup S2$. Thus, an initial cascade list grows through iteration until no more new indices are found. To cascade set S through the given table:

- (1) Set the initial cascade list to S.
- (2) For each s in the cascade list, remove all (s, x, y) entries from the table.
- (3) For the remaining table entries, for each index in an i or j position that belongs to the cascade list, replace the index with 0.

```
24,67,0
25, 32, 0
                 \implies (25, 0, 0)
26, 32, 0)
                 \implies (26, 0, 0)
27, 32, 0)
                 \implies (27, 0, 0)
30, 32, 0)
                 \implies (30, 0, 0)
30, 35, 37
                                     \implies deleted!
31, 25, 37
                                     \implies (31, 0, 37) \implies (31, 0, 0)
32, 27, 31
                 \implies deleted!
33, 24, 39)
35, 27, 0
                                     \Longrightarrow (35, 0, 0)
35, 30, 0)
                                     \implies (35, 0, 0)
                                     \implies (37, 0, 0)
37, 30, 0)
38, 35, 40)
                                                          \implies (38, 0, 40) = (38, 40, 0)
```

FIGURE 1. Computing Cascade(32). (r, 0, 0) indicates that r is covered.

```
( 24, 67, 0)
( 33, 24, 39)
( 38, 40, 0)
```

FIGURE 2. The result after cascading 32.

- (4) A (t,0,0) entry indicates that t is covered. Append t to the cascade list.
- (5) Iterate for the set of new indices in the cascade list, until no new (t, 0, 0) index is found.

Figure 1 illustrates substitutions for $S = \{32\}$ into the table in the figure. After replacing 32 with 0, indices 25, 26, 27, and 30 are added to the cascade list.

- . Then these new values need to be replaced with 0's in the table. Iterations continue until no further changes to the list occur. In this example, the final cascade list consists of 32 and 25, 26, 27, 30, 31, 35, and 37, and the final table (see Figure 2) has only three entries.
- 3.3. Spans Relative to a Table. Earlier, Span(S) was defined to be the set of all non-free products in Consensus(< Free >, S), in other words, the non-free products that are covered by S. From now on, spans will be calculated relative to whatever local triples table T is under study.
- **Definition 3.1.** Span(S,T), the span of a set S relative to triples table T, is equal to the union of S with the set of products in T that are covered by S. That is, the span is the cascade list that results from cascading S through a copy of T. When the relevant table T is clear from the context, the span can be written Span(S).

The span of an individual prime often provides useful information when trying to calculate a minimum-cost cover $QS_i \subset PS_i$.

4. Ancestor Sets

In this section and the ones that follow, all products are active and non-free. P, Q, and R will represent active non-free primes. X, Y, or Z can represent any active product in NONFREE, including primes.

At this point, we are ready to define and compute the Ancestor Sets associated with a triples table T.

Definition 4.1. Given triples table T, for any triple (r, i, 0) i is a parent of r. For (r, i, j), j > 0, i and j are parents of r.

Definition 4.2. The Ancestor Set of non-free product X_r , $Anc(X_r)$, (also written Anc(r)) consists of the union of all of the parents of r with the parents of parents, parents of parents of parents, ... iterated until there no new ancestors are added.

A 0/1 bitmap can be used to represent the parents of each non-free product. A simple (but inefficient) way to find Anc(r) is to set the initial A(r)=parents(r) and then let

$$A(r) = A(r)$$
 OR parents(each parent).

Then, for each new member of A(r), set

$$A(r) = A(r)$$
 OR $parents(each new member)$

until no new ancestors are added.

Definition 4.3. The Ancestor Set of a non-free prime is called a *Prime Ancestor Set*.

Later, it will be shown that only Prime Ancestor Sets need to be calculated, and in fact, there are shortcuts that identify a small subset of the primes whose ancestor sets actually need to be calculated. The algorithms for computing ancestor sets make use of the following obvious lemma.

Lemma 4.1. If $Y \in Anc(X)$ then $Anc(Y) \subset Anc(X)$.

Definition 4.4. An Ancestor Set is *Independent* if it does not properly contain any other Ancestor Set.

The Independent Ancestor Sets are the critical ones. Lemma 4.5 will show that every Independent Ancestor Set is a Prime Ancestor Set.

Definition 4.5. For Ancestor Set A, triples(A) is defined to be the set of all triples (r, i, j) in T such that r, i, and j, if j > 0, belong to A.

Lemma 4.2. Every non-free product has some ancestors that are non-free primes.

Proof. For the non-free primes, this follows directly from Lemma 2.3. For other non-free products, it follows from the fact that all of the non-free products were generated by $Consensus(\langle Free \rangle, non-free\ primes)$.

Definition 4.6. Given an Ancestor Set A, the *primeset* for A is the set of all primes that belong to A.

Definition 4.7. Primes Q_1, \ldots, Q_t cover A if A is equal to the span of Q_1, \ldots, Q_t in triples(A).

Lemma 4.3. Suppose that A is an Ancestor Set and $PS = \{P_1, \ldots, P_m\}$ is its primeset. Then PS covers A with respect to triples(A), that is,

$$A = Span(P_1, \dots, P_m)$$
 in $triples(A)$.

Proof. Every non-free product X is generated by consensus combinations that include non-free primes, (as well as free primes that are represented by 0s in the table). P_1, \ldots, P_m are the only primes in A and hence are the only non-free prime ancestors of any $X \in A$. Hence A must be contained in their span.

Lemma 4.4. Primes $QS = Q_1, \dots, Q_t$ cover A if and only if they cover the primeset PS of A.

Proof. Since by definition, the primeset PS for A is contained in A, if QS is a cover for A it must cover PS. Conversely, if QS covers PS, the span of QS must contain A.

Definition 4.8. An Ancestor Set A(Y) is closed with respect to Y if $Y \in A(Y)$.

4.1. Statement and Proof of the Ancestor Theorem. The Ancestor Theorem is a crucial result for this paper. The statement is true for any Ancestor Set Anc(X), although it will only need to be applied to Independent Ancestor Sets.

The key fact is that every minimum-cost cover for an Ancestor Set A has to be chosen from the primeset for A, since all other primes are irrelevant to covering A.

Theorem 4.1 (Ancestor Theorem). Let A be an Ancestor Set, PS be its primeset, and RS be the set of primes that are not covered by the union of the Essential Primes with PS. Every minimum-cost basis for the primes consists of the Essential Primes, a minimum-cost subset of PS that covers PS, and a minimum-cost subset of RS that covers RS.

Proof. By the definition of Ancestor Sets, every non-free prime ancestor of any $P \in PS$ belongs to PS. Let Q_1, \ldots, Q_k be a minimum-cost cover for all of the non-free primes. That is,

 $Span(Q_1, \ldots, Q_k)$ with respect to T covers all of the non-free primes.

Some minimum-cost subset of $\{Q_i\}$ covers the primeset of A. Suppose that this subset consists of Q_1, \ldots, Q_t . If some Q_i , $i \leq t$ does not belong to PS (and hence does not belong to A), it is not an ancestor of any product in PS, or in A. The span of the remaining Q_j must cover PS, which contradicts Q_1, \ldots, Q_t being a minimum-cost set. Hence Q_1, \ldots, Q_t must all belong to PS and must cover PS.

Suppose that Q_1, \ldots, Q_t covers PS, but is not a minimum-cost cover for PS. If the set of primes $\{R_1, \ldots, R_m\} \subset PS$, is a minimum-cost cover for PS, then R_1, \ldots, R_m covers every product in PS including the primes Q_1, \ldots, Q_t , so $R_1, \ldots, R_m, Q_{t+1}, \ldots, Q_k$ is a better cover for the non-free primes, contradicting the assumption that Q_1, \ldots, Q_k is a minimum-cost cover. Finally, no prime that is already covered by the union of the Essential Primes with PS can be part of a minimum-cost basis.

At this point, the Ancestor Sets may not be disjoint. An Ancestor Set may contain or be contained in other Ancestor Sets.

Corollary 4.1. If Ancestor Set A is contained in Ancestor Set B, then a minimum-cost cover for B contains a minimum-cost cover for A.

4.2. Properties of Independent Ancestor Sets. Recall that Anc(X) is Independent if it does not properly contain Anc(Y) for any $Y \in Anc(X)$. Note that this implies that if Anc(X) is Independent, Anc(X) = Anc(Y) for every $Y \in Anc(X)$.

Lemma 4.5. Every Independent Ancestor Set is a Prime Ancestor Set.

Proof. For each prime Q in the primeset of Independent Ancestor Set A, A = Anc(Q).

There always is at least one Independent Ancestor Set, namely an Ancestor Set of smallest size. The following *Independence Lemma* characterizes Independent Ancestor Sets.

Lemma 4.6 (Independence Lemma.). Suppose that A is an Ancestor Set and Q_1, \ldots, Q_k is its primeset. Then A is an Independent Ancestor Set if and only if

$$Anc(Q_1) = Anc(Q_2) = \cdots = Anc(Q_k) = A.$$

Proof. Suppose that A is Independent. Since each $Q_i \in A$, $Anc(Q_i) \subset A$. But $Anc(Q_i)$ cannot be properly contained in A so $Anc(Q_i) = A$.

For the converse, suppose that $Anc(Q_i) = A$ for every prime Q_i in the primeset of A. Suppose that B is an Ancestor Set properly contained in A. Every Ancestor Set must contain primes. Suppose that prime $Q \in B$. Then $Anc(Q) \subset B$. Since $B \subset A$, $Q \in A$. But by hypothesis, Anc(Q) = A, which contradicts B properly contained in A.

Lemma 4.7. If A_m and A_k are distinct Independent Ancestor Sets, then A_m and A_k are disjoint, that is, $A_m \cap A_k$ is empty.

Proof. If
$$X \in A_m$$
 and $X \in A_k$, then $A_m = Anc(X) = A_k$.

Note that the primesets of distinct independent ancestor sets A_m and A_k are disjoint, and $triples(A_m)$ is disjoint from $triples(A_k)$.

4.3. Finding the Initial Independent Ancestor Sets. Suppose that the *triples* table is known and has been stripped of all inactive or useless entries. The procedure that follows uncovers the independent ancestor sets iteratively, one batch at a time. The steps are easy to follow, and prove that the iteration will succeed.⁶

After the first batch of Independent Ancestor Sets has been generated, *triples* will be replaced by a smaller table and a fresh batch of Independent Ancestor Sets will be calculated from the smaller table.

Iteration continues until the triples table is empty. We assume that at each step in the discussion that follows, triples identifies the table that is current. The steps that follow

⁶However, the algorithm in Appendix C bypasses the calculation of most of the ancestor sets. It computes batches of independent ancestor sets directly using a method that converges quite quickly.

generate the first batch of Independent Ancestor Sets. Note that initially, many Ancestor Sets will have the same entries.

- (1) Since all Independent Ancestor Sets are Prime Ancestor Sets, build the Prime Ancestor Sets for *triples*.⁷
- (2) Order the list of Prime Ancestor Sets by increasing size.
- (3) Discard all Ancestor Sets Anc(P) such that $P \notin Anc(P)$. (P will belong to Surplus.)
- (4) The smallest Ancestor Set, Anc(Q), must be Independent. Move it to the list of Independents.
- (5) Every Ancestor Set B such that $Q \in B$ equals or contains Anc(Q). Discard these sets.
- (6) If any Ancestor Sets remain, iterate from (4).

In the two sections that follow, we assign table $triples(A_i)$ to each Independent Ancestor Set A_i and then reduce the size of triples in preparation for finding the next batch of Independent Ancestor Sets. Every Independent Ancestor Set that has been discovered causes a piece of the current triples table to be removed and additional covered indices in triples to be set to 0.

4.4. Assigning $triples(A_k)$ to Independent A_k . Suppose that A_k is Independent. Recall that $triples(A_k)$ is the set of all triples (r, i, j) such that r, i, and j, if j > 0, belong to A_k . Table $triples(A_k)$ contains all covering relationships for each $r \in A_k$. Given a positive additive cost function, this information determines a minimum-cost set of primes that covers PS_k (and A_k) and is part of a minimum-cost basis for f.

Since the Independent Ancestor Sets $\{A_k\}$ are disjoint, the arrays $triples(A_k)$ also are disjoint. The $triples(A_k)$ tables are called the *Independent triples tables*.

All of the entries in these tables now can be removed from the main *triples* table.

4.5. Cascading the Union of the Independent Ancestor Sets. As was noted in Section 4.1, the Ancestor Theorem shows that the minimum-cost covers of the current set of Independent Ancestor Sets are disjoint pieces of the overall minimum-cost basis.

A minimum-cost cover of PS depends only on triples(A) and the given cost function C. From the point of view of the other Independent Ancestor Sets, this separate piece of the problem can be viewed as solved. All of the indices covered by PS can be removed from the problem by cascading A through triples.

We already have removed each triples(A) from the table but other downstream indices may be covered by the indices in A. Note that the individual Independent Ancestor Sets do not have to be cascaded separately. If A_1, \ldots, A_m form the current batch of Independent Ancestor Sets, let AU denote their union. The covered indices are set to 0 by performing a removal cascade of AU through the current triples. That is, set the initial cascade list to AU and,

(1) Replace each index belonging to the cascade list with 0 in the current triples.

⁷The algorithm in Appendix C builds the first batch of independent sets directly and quickly. However, the steps listed here are easy to follow, and prove that the iteration will succeed.

- (2) An (s,0,0) entry indicates that s has been covered by AU. Remove all (s,x,y) entries and append index s to the newIndex cascade list.
- (3) Iterate, setting all new s indices to 0 in the triples table, until no new covered indices are found.

The final cascade list equals $AU + extra\ indices$. The extra indices that correspond to primes belong to Surplus.

As a result of the cascade, all indices that were in AU and any extra indices that were covered by AU have been removed from the current triples and hence have been removed from the problem. Specifically, if any X_t in an Independent A belongs to non-Independent B, then, since $Anc(X_t) = A$, all of A is contained in B. A cascade of A through triples(B) would remove all indices in A from B and also would remove any extra indices in B that are covered by A.

4.6. The Iterative Ancestor Set Algorithm. At this point, the initial Independent Ancestor Sets $\{A_k\}$ and tables $triples(A_k)$ have been set aside and the current triples has been reduced and simplified. If the current table is not empty, iterating the preceding steps for the current table will reveal new disjoint Independent Ancestor Sets with their own associated tables. Since the size of the current triples table decreases at each iteration, the process will terminate.

The overall process is summarized below. The following operations build the list of Independent Ancestor Sets and their disjoint primesets.

- (1) Remove duplicate entries and useless entries (i.e. (r, i, j) when there is an entry (r, i, 0) or (r, j, 0)) from the current triples. Remove inactive indices.
- (2) Find the Prime Ancestor Sets for the current triples.
- (3) Identify a new batch of Independent Ancestor Sets.
- (4) For each Independent Ancestor Set A_k , let PS_k be its primeset and find $triples(A_k)$. Associate $triples(A_k)$ with PS_k and remove the $triples(A_k)$ entries from the current triples table.
- (5) Cascade the union of the new Independent Ancestor Sets through the current *triples* table in order to remove all other products covered by the current batch of Independent Ancestor Sets from the problem.
- (6) If the resulting table is not empty, iterate from step 1.

Since the smallest Ancestor Set in a batch always is Independent, at each iteration, the total number of Independent Ancestor Sets will increase and the number of entries in the current *triples* will decrease and eventually become zero.

At that point, the set of Independent Ancestor Sets $\{A_i\}$ is complete and all of the disjoint primesets PS_i have been identified.

5. Summary of the Theorems

We summarize the results of the preceding sections in the following theorems.

Theorem 5.1 (Cascade List). The cascade list resulting from cascading the union of the canonical Independent primesets PS_i of Boolean function f through triples(f) contains all of the products in triples(f).

Proof. Each PS_i covers A_i . The reduction process cascades the union of the Independent Ancestor Sets A_i through the table, covering all of the active, non-free products. Since the active products cover all of the inactive non-free products all non-free products will be covered.

Theorem 5.2. Given a Boolean function f with Independent Ancestor Sets A_i and primesets $PS_i \subset A_i$, let Surplus be the set of non-free primes that do not belong to $\bigcup PS_i$. Then no prime in Surplus can be part of any minimum-cost basis.

Proof. By definition, $P \in Surplus$ does not belong to any PS_i and hence is not an ancestor of any prime that belongs to any PS_i . The products in A_i can only be covered by a subset QS_i of PS_i that covers PS_i , and the union of the QS_i covers all non-free products, including the products in Surplus. Therefore, any prime $P \in Surplus$ that was part of a basis would be covered by other primes that are in the basis, so the basis would not have minimum cost.

Theorem 5.3 (Canonical Partition). Let f be a Boolean function. There is a canonical partition of the complete set of primes of f into disjoint sets of primes,

$$(Essential\ Primes) \cup (PS_1) \cup \cdots \cup (PS_N) \cup (Unnecessary\ Primes)$$

where

- (1) The Essential Primes must be part of any basis for f.
- (2) PS_1, \ldots, PS_N are the primesets of the Independent Ancestor Sets A_1, \ldots, A_N for f.
- (3) The Independent Ancestor Sets for f are disjoint.
- (4) The Unnecessary Primes consist of the union of the free non-essential primes with Surplus, and an Unnecessary Prime cannot be part of a minimum-cost basis for any positive, additive cost function.
- (5) For each PS_i , there is a canonical triples table TS_i that contains all covering relationships between the products that belong to A_i .
- (6) Given a positive additive cost function C, any minimum-cost basis for f with respect to C must consist of the union of the Essential Primes with subsets $QS_i(C) \subset PS_i$ such that $QS_i(C)$ is a minimum-cost cover of PS_i . (That is, $Cascade(QS_i(C), TS_i)$ is empty.)

The sections that follow deal with the spans of primes, which are the keys to the solutions QS_i of some uncomplicated Independent Ancestor Sets A_i .

6. Spans of Individual Primes and Simple Solutions

Suppose that A is an Independent Ancestor Set for Boolean function f and PS is the primeset for A. The set of spans of the individual primes $P \in PS$ are of special interest. Recall that the span of a prime P in A equals the products in A that are covered by P.

Lemma 6.1. Prime P is an ancestor of every product in Span(P).

Proof. Clear from the definition of the span.

Definition 6.1. The span of prime $P_r \in A$ is maximal if the span includes at least one prime other than P_r and $Span(P_r)$ is not properly contained in the span of any other prime in A.

Definition 6.2. The span of prime $P_r \in A$ is closed if triples(A) includes a triple (r, i, j) such that $X_i \in Span(P_r)$ and, if j > 0, $X_j \in Span(P_r)$.

Lemma 6.2. If $Q \in Span(P)$ then $Span(Q) \subset Span(P)$.

Proof. Clear from the definition of a span.

Definition 6.3. Primes P and Q are equivalent if and only if P is in the span of Q and Q is in the span of P.

Lemma 6.3 (Equivalence Lemma). If P and Q are equivalent, then Span(P) = Span(Q) and the span is closed with respect to both P and Q.

Proof. By Lemma (6.2),

$$Span(Q) \subset Span(P)$$

$$Span(P) \subset Span(Q)$$
.

For equivalent primes P and Q, whenever P is covered, Q also is covered and vice-versa. A set of primes that are equivalent can be represented by the one with the lowest cost.

Lemma 6.4. If P is equivalent to Q, then

$$Anc(P) = Anc(Q).$$

Anc(P) and Anc(Q) are closed with respect to P and Q.

Proof. Anc(P) = Anc(Q) since P is an ancestor of Q and Q is an ancestor of P, so $Anc(P) \subset Anc(Q)$ and $Anc(Q) \subset Anc(P)$. $P \in Anc(P)$ since $P \in Anc(Q) = Anc(P)$. Similarly, $Q \in Anc(Q)$.

6.1. **The Span Basis Theorem.** It is not unusual for an Independent Ancestor Set to be equal to the span of one of its primes. The Span Basis Lemma states that in this case a minimum-cost cover for A equals a minimum-cost cover for P.

Lemma 6.5 (Span Basis Lemma). Suppose that A=Anc(P) is an Independent Ancestor Set and $P \in A$. If Span(P) in triples(A) equals A, then Span(P) is maximal and closed, and $Q_1, \ldots Q_t$ is a minimum-cost cover for A if and only if it is a minimum-cost cover for P.

Proof. It is clear that Span(P) is maximal and closed. Any cover for A covers P, and any cover for P covers A. Hence, both have the same minimum-cost covers.

The most desirable result of the Span Basis Lemma would be that a single prime P is the minimum-cost cover for A. The Span Basis Theorem that follows shows that this is the case for the most common cost functions when P is equal to the minimum-cost prime in its equivalence class. In this case, P must belong to any minimum-cost basis for f.

Definition 6.4. Cost function C is a *constant additive* cost function if C(X) equals a constant c for each product X, and C is additive.

Theorem 6.1 (Span Basis Theorem). Suppose that $f(x_1, ..., x_n)$ is a Boolean function, and

- (1) A=Anc(P) is an Independent Ancestor Set and $P \in A$,
- (2) Span(P) in triples(A) = A,
- (3) for every prime Q equivalent to P, $C(P) \leq C(Q)$,
- (4) the cost function is either constant additive or satisfies Quine's hypothesis (namely that the addition of a single factor to a formula increases the cost of the formula).

Then P is a minimum-cost cover for A, and P (or an equivalent prime with the same cost) must be part of any minimum-cost basis.

Proof. Condition (3) can be guaranteed when the non-free primes are ordered by cost, and the span of the least-cost prime in a set of equivalent ones is always selected to represent the set of equivalent spans. In this case, any minimum cost cover of P other than P would consist of two or more primes. If C is constant then the cost of any cover of P with two or more primes is > C(P). If C satisfies Quine's hypothesis, then for any cover consisting of two or more primes, the union of the factors of the primes in such a cover of P would have to properly contain every factor of P along with other factors. Thus if the cost function satisfies Quine's hypothesis, such a cover could not be minimum-cost.

6.2. Splitting an Independent Ancestor Set. It may appear that it is not possible to further collapse an Independent Ancestor Set, but in fact, an Independent Ancestor Set can be decomposed, (simplifying its solution) if it contains an *Independent prime*.

Definition 6.5. Suppose that prime P belongs to Independent Ancestor Set A. We say that prime P is Independent if Span(P) in triples(A) is maximal and closed, $Span(P) \neq A$, and P does not belong to the span of its complement in A.

In this case, any minimum-cost cover for A consists of a minimum-cost cover for P (and hence, for Span(P)) plus a minimum-cost cover of the primes in A and outside of Span(P).

The independence of P can be checked by computing the span in triples(A) of the set of products in A that do not belong to Span(P). If P is not covered, then P is Independent.

A good candidate for independence is the prime in A with the largest span, such that Span(P) is maximal and closed, and such that for every entry corresponding to (P, X_i, X_j) in the triples table, Xi, and Xj, $(if X_j > 0)$, belong to Span(P). The following Theorem describes the partition in more detail.

Theorem 6.2 (Independent Prime Decomposition Theorem). If $P \in A$ is Independent, then A can be split into Span(P) and one or more smaller Ancestor Sets.

Proof. The splitting algorithm is:

- (1) Extract from triples(A): $triples(Span(P)) = \{(r, i, j)\}$ such that r, i, and j (if j > 0)) belong to Span(P).
- (2) Cascade Span(P) through the remaining entries.
- (3) Calculate the Independent Ancestor Sets for the remaining triples.

Thus, the solution to A consists of a minimum-cost cover for Span(P), as in Section(6.1) plus the solutions to the second-generation Independent Ancestor Sets associated with the reduced table.

If C is constant or satisfies Quine's hypothesis and $C(P) \le C(Q)$ for every prime Q equivalent to Independent prime P, then $P \in Anc(P)$ is a minimum-cost cover for Span(P) and P (or an equivalent prime with the same cost) must be part of any minimum-cost basis.

Note that one or more of the second-generation Independent Ancestor Sets might contain Independent primes, and the process could iterate several times. However, the size of the new Ancestor Sets and of the their tables decreases at each iteration and the number of steps is linear in the data.

APPENDIX A. AN ALGORITHM FOR REORDERING THE NON-FREE PRODUCTS IN ALL

The following algorithm is convenient for problems of moderate size. The algorithm reorders ALL while also computing the span of each non-free prime P_k , k = 1...M, as specified in definition 2.7, that is,

$$Span(P_k) = Consensus(\langle Free \rangle, P_k).$$

The reordering places all non-free products that are in the span of a non-free prime before any products that are not covered by an individual prime. Recall that $Span(P_k)$ is closed if there are non-free products $X_i, X_j \in Span(P_k)$ with $P_k = (X_i \circ X_j)$.

The algorithm also generates some special entries for the triples0 table. Since by definition, P_k covers every X_s in its span, each triple (s, k, 0) is a valid entry for triples0. However, the algorithm below only saves entries that arise from maximal closed spans.

- (1) Initialize newALL = Essentials, non-Essential frees, non-free primes.
- (2) For each P_k ,
 - (a) compute $Span(P_k)$.
 - (b) for each non-free $Y \in Span(P_k)$ that is not in newAll, append Y to newALL.
- (3) For each $Z \in ALL$ that is not in newALL, append Z to newAll.
- (4) Sort the spans by decreasing size and identify the maximal closed spans,

$$Span(Q_1), \dots Span(Q_h).$$

(5) For each $X_r \in Span(Q_i)$, append (r, i, 0) to triples0.

APPENDIX B. AN ALGORITHM FOR GENERATING THE triples TABLE

The first set of entries for triples0 were generated in Appendix A, and consist of (r, i, 0) such that X_r belongs to a maximal closed $Span(P_i)$.

The algorithm that follows generates ordinary triples0 entries by taking the loose Iterated Consensus of non-free, non-prime products X_i with free products.

Then the algorithm calculates triples3 entries by taking the consensus of pairs of non-free products X_i and X_j . For each non-free result X_r , (r, i, j) is tested to see if it is useful. Recall that (r, i, j) is useful if there is no (r, i, 0) or (r, j, 0) in triples0. Only useful entries are appended to triples3.

It is not difficult to organize the data in ways that speed up the processing.

- (1) Set productList equal to the non-free products in newALL.
- (2) Part 1. Calculate ordinary *triples*0 entries:
 - (a) For each non-prime X_i on the productList, compute the consensus of X_i with each free product. If the result $X_r = (X_i \ o \ free)$ is non-free and (r, i, 0) is not a duplicate, append it to triples0.
- (3) Part 2. Calculate the *triples*3 entries:
 - (a) Perform Loose Iterated Consensus on the entries in productList.
 - (b) Test each non-free result $X_r = (X_i \circ X_j)$. If it is useful, append (r, i, j) to triples3.

Note that there has to be at least one table entry (r, i, j) for each non-free product X_r . After removing all inactive entries, there are entries for each active non-free product. It is convenient to sort the table by r - values.

APPENDIX C. A DIRECT ALGORITHM FOR INDEPENDENT ANCESTOR SETS

We assume that the non-free primes $P_1, \ldots P_n$ have been arranged in order of increasing cost. We compute the ancestor sets of the primes starting from the most costly.

If P_i , i < n, is a parent of P_n , then $Anc(P_i) \subseteq Anc(P_n)$. If the sets are equal, the least costly prime P_i will be preferred to P_n as the representative prime for the set. If $Anc(P_i)$ is a proper subset, $Anc(P_n)$ cannot be independent. In either case, P_n can be discarded.

If P_i , i < n, is a grandparent of P_n , then again $Anc(P_i) \subseteq Anc(P_n)$ and P_n can be discarded. More generally, checking each generation of the ancestors of the remaining primes (processed in order of decreasing cost) leads to more discards. The number of primes whose ancestor sets require full expansion shrinks quickly.

One last check of the remaining set of primes is needed. If $P_j \in Anc(P_i)$ for j > i, then $Anc(P_j)$ must be a proper subset of $Anc(P_i)$; otherwise, $P_i \in Anc(P_j)$ and P_j would have been discarded. Thus, P_i can be discarded. When this test has been completed, the remaining primes and their ancestor sets form the first batch of Independent ancestor sets.

After extracting each set $triples(Anc(P_k))$ and cascading the union of the ancestor sets in this batch through the triples table, the process is iterated until triples is empty.

Once all of the independent sets and their associated triples have been found, each Independent set can be checked to find the prime in the set whose span (within its triples set) is largest. If $Q \in Anc(P)$ and Span(Q) = Anc(Q) (and hence equals Anc(P)) and the cost function is constant additive or satisfies Quine's hypothesis, then Anc(Q) is solved. If Span(Q) is not equal to Anc(Q) but Span(Q) is maximal and closed, then Q can be tested to see whether it is independent, which would enable the ancestor set to be split.

References

- [1] S. B. Akers. Binary Decision Diagrams. *IEEE Transactions on Computers*, C-27(6) (June 1978), 509516.
- [2] R. K. Brayton, G. D. Hachtel, C. T. McMullen, and A. L. Sangiovanni-Vincentelli. Logic Minimization Algorithms for VLSI Synthesis. Kluwer Academic Publishers, (1990).
- [3] R. T. Boute. The Binary Decision Machine as a programmable controller. *EUROMICRO Newsletter*, Vol. 1(2) (January 1976), 1622.
- [4] F. M. Brown. Boolean Reasoning: The Logic of Boolean Equations, Second Edition. *Dover Publications*, Appendix A, (2003), 259-265.
- [5] C. Y. Lee. Representation of Switching Circuits by Binary-Decision Programs. *Bell Systems Technical Journal*, 38 (1959), 985999.
- [6] G. Hachtel and F. Somenzi. Logic Synthesis and Verification Algorithms, Springer, (2006).
- [7] E. J. McCluskey, Junior. Minimization of Boolean Functions. Bell Syst. Technical Journal, 35 (Nov. 1956), 1417-1444.
- [8] C. McMullen and J. Shearer. Prime implicants, minimum covers, and the complexity of logic simplification. *IEEE Trans. Comput.* C-35(8), (1986), 761-762.
- [9] W. Quine. The problem of simplifying truth functions. Amer. Math. Monthly 59, (1952), 521-531.
- [10] T. Sasao. Input variable assignment and output phase optimization of PLA's. IEEE Transactions on Computers C-33 (Oct 1984), 879-894.