Approximate constraint satisfaction requires large LP relaxations

Siu On Chan* James R. Lee[†] Prasad Raghavendra[‡] David Steurer[§]

Abstract

We prove super-polynomial lower bounds on the size of linear programming relaxations for approximation versions of constraint satisfaction problems. We show that for these problems, polynomial-sized linear programs are no more powerful than programs arising from a constant number of rounds of the Sherali–Adams hierarchy.

In particular, any polynomial-sized linear program for Max Cut has an integrality gap of $\frac{1}{2}$ and any such linear program for Max 3-Sat has an integrality gap of $\frac{7}{8}$.

1 Introduction

Linear programming is one of the most powerful tools known for finding approximately optimal solutions to NP-hard problems. We refer to the books [Vaz01, WS11] which each contain a wealth of examples. If $P \neq NP$, then for many such problems we do not expect polynomial-sized linear programs (LPs) to compute arbitrarily good approximations to the optimal solution. (More formally, if $NP \nsubseteq P/poly$, then such LPs cannot exist [Yan91].)

Thus a line of research has sought to prove lower bounds on the efficacy of small linear programs. The construction of integrality gaps for specific LPs has long been a topic of interest in approximation algorithms. Arora, Bollobás, and Lovász [ABL02] initiated a more systematic study; they explored the limitations of LPs arising from lift-and-project hierarchies like those of Lovász and Schrijver [LS91] and Sherali and Adams [SA90]. There has now been an extensive amount of progress made in this area; one can see a sampling in the section on previous work.

Arguably, the ultimate goal of this study is to prove unconditional lower bounds for *every* sufficiently small LP. Since linear programming is *P*-complete

^{*}Microsoft Research New England

[†]University of Washington

[‡]U. C. Berkelev

[§]Cornell University

under various notions of reduction, this would require proving that NP does not have polynomial-size circuits (see, e.g., the discussion in [Yan91]). But one could still hope to complete this program for LPs that use the natural encoding of the underlying combinatorial problem.

We make progress toward this goal for the class of constraint satisfaction problems (CSPs). For instance, we prove that every polynomial-sized LP for Max Cut has an integrality gap of $\frac{1}{2}$, answering a question from [BFPS12]. As another example, every such LP for Max 3-Sat has an integrality gap of $\frac{7}{8}$, and every such LP for Max 2-Sat has an integrality gap of $\frac{3}{4}$. In fact, in both cases these integrality gaps hold for families of LPs of size up to $n^{o(\frac{\log n}{\log \log n})}$.

Corresponding upper bounds for all three problems can be achieved by simple polynomial-sized LPs. For Max 3-Sat, a $\frac{7}{8}$ -approximation is best-possible assuming P \neq NP [Hås01]. For Max Cut, the seminal SDP-based algorithm of Goemans and Williamson [GW95] achieves a 0.878-approximation. In this case, our result yields a strict separation between the power of polynomial-sized LPs and SDPs for a natural optimization problem. Interestingly, even a simple spectral algorithm can do strictly better than 1/2 for Max Cut [Tre12].

To establish these lower bounds, we show that for approximating CSPs, polynomial-sized LPs are exactly as powerful as those programs arising from O(1) rounds of the Sherali–Adams hierarchy. We are then able to employ the powerful Sherali–Adams gaps that appear in prior work. This offers a potential framework for understanding the power of linear programs for many problems by relating their expressive power to that of the very explicit Sherali–Adams hierarchy.

In Section 1.2, we discuss our approach for the specific example of Max Cut, including the class of LPs to which our results apply. Section 2 is devoted to a review of CSPs and their linear relaxations. There we explain our basic approach to proving lower bounds by exhibiting an appropriate separating hyperplane. We also review the Sherali–Adams hierarchy for CSPs. In Section 3, we present the technical components of our approach, as well as the proof of our main theorem.

Finally, Section 4 contains an illustrative discussion of how Sherali–Adams gap examples can be used to construct corresponding gaps for *symmetric* LPs. This connection is quantitatively stronger than our result for general LPs. We refer to Section 5 for a discussion of future directions.

Recent work. Since initial publication of this manuscript, there has been substantial followup work building on the ideas presented here. The papers [LRST14, FSP13] establish a connection between *symmetric* semidefinite programs and the Sum-of-Squares hierarchy by analogy with our work in Section 4. In [LRS15], a connection between *general* semidefinite extended formulations and the Sum-of-Squares hierarchy is established; in particular, the authors prove

exponential lower bounds on the semidefinite extension complexity of explicit polytopes (like the TSP polytopes). Finally, our models for approximation via linear programs are extended and refined in the work [BPZ15]; the authors show that a suitable notion of reduction within the model allows one to derive lower bounds for additional problems (other than CSPs).

1.1 History and context

Extended formulations. In a seminal paper, Yannakakis [Yan91] proved that every symmetric LP (i.e., one whose formulation is invariant under permutations of the variables) for TSP has exponential size. Only recently was a similar lower bound given for general LPs. More precisely, Fiorini, et al. [FMP+12] show that the extension complexity of the TSP polytope is at least $2^{\Omega(\sqrt{n})}$ for n-vertex graphs.

Braun, et al. [BFPS12] expand the notion of extension complexity to include approximation problems and show that approximating Max Clique within $O(n^{1/2-\varepsilon})$ requires LPs of size $2^{\Omega(n^{\varepsilon})}$. Building on that work, Braverman and Moitra [BM13] show that approximating Max Clique within $O(n^{1-\varepsilon})$ requires LPs of size $2^{\Omega(n^{\varepsilon})}$. We remark that the encoding of Max Clique used in the later two works is somewhat lacking. Specifically, these lower bounds do not encompass, for instance, standard relaxations for Max Clique, including those given by the Sherali-Adams hierarchy.

These three latter papers all use Yannakakis' connection between extension complexity and non-negative rank (see [FMP+12] for a detailed discussion). They are based on increasingly more sophisticated analyses of a single family of slack matrices first defined in [FMP+12] (and extended to the approximation setting by [BFPS12]). Closely related slack matrices are employed in a recent paper of Rothvoss [Rot14] to show exponential lower bounds on the extension complexity of the matching polytope. A significant contribution of the present work is that the connection between general LPs and the Sherali–Adams hierarchy allows one to employ a much richer family of hard instances.

LP and **SDP** hierarchies. As mentioned previously, starting with the works [ABL02, ABLT06], the efficacy of LP and SDP hierarchies for approximation problems has been extensively studied. We refer to the survey of Laurent [Lau03] for a discussion of the various hierarchies and their relationships.

We mention a few results that will be quite useful for us. Fernández de la Vega and Mathieu [FdlVKM07] showed that for any fixed $\varepsilon>0$ and k, Max Cut has an integrality gap of $\frac{1}{2}+\varepsilon$ even after k rounds of the Sherali–Adams hierarchy. In a paper of Charikar, Makarychev, and Makarychev [CMM09], it is shown that Max Cut and Vertex Cover have integrality gaps of $\frac{1}{2}+\varepsilon$ and $2-\varepsilon$, respectively, for $n^{\Omega(\varepsilon)}$ rounds of the Sherali–Adams hierarchy.

In work of Schoenebeck [Sch08], tight bounds are given on the number of rounds needed to approximate k-CSPs in the Lasserre hierarchy (which, in particular, is stronger than the Sherali–Adams hierarchy). For instance, he shows that for every $\varepsilon > 0$, Max 3-Sat has a $\frac{7}{8} + \varepsilon$ integrality gap even after $\Omega(n)$ rounds. One should consult also the much earlier work of Grigoriev [Gri01] which achieves an equivalent family of lower bounds stated in the dual setting of Positivstellensatz proof systems. There are also Sherali–Adams integrality gaps for CSPs with a pairwise independent predicate, due to Benabbas et al. [BGMT12].

Strong separation between nonnegative rank and smooth nonnegative rank. We remark that all previous lower bounds for nonnegative rank (at least in the context of extended formulations) are robust with respect to small multiplicative perturbations [Rot14, FMP⁺12, BFPS12, BM13]. Concretely, if we define the ε -smooth nonnegative rank of a matrix A as

$$\operatorname{rank}_{+,\varepsilon}(A) := \min \left\{ \operatorname{rank}_{+}(A') \mid (1-\varepsilon)A_{ij} \leq A'_{ij} \leq (1+\varepsilon)A_{ij} \right\},\,$$

then all previous lower bounds for nonnegative rank also lower bound the ε -smooth version for some absolute constant $\varepsilon > 0$. A related generalization of nonnegative rank is approximate nonnegative rank that allows additive instead of multiplicative error. ¹ This version of nonnegative rank is equivalent to the smooth rectangle bound [KMSY14].

In contrast, the matrices studied in this work turn out to have only polynomial approximate and smooth nonnegative rank. In this sense, our superpolynomial lower bounds on the nonnegative rank of these matrices give the first separation between nonnegative rank and smooth nonnegative rank. See Section 3.4 for a discussion.

1.2 Outline: Max Cut

We now present the basic details of our approach applied to the Max Cut problem. To this end, consider a graph G = (V, E) with |V| = n. For any $S \subseteq V$, we use

$$G(S) \stackrel{\text{def}}{=} \frac{|E(S, \bar{S})|}{|E|}$$

to denote the fraction of edges of G crossing the cut (S, \bar{S}) . The maximum cut value of G is opt $(G) = \max_{S \subseteq V} G(S)$.

¹All previous lower bounds also hold for this generalization of nonnegative rank. However, some of the lower bound arguments do not apply to the additive-error setting. (For example, arguments that rely on the zero / non-zero pattern of the matrix.)

The standard LP. To construct an LP for computing (or approximating) opt(G), it is natural to introduce variables $x = (x_1, x_2, ..., x_n) \in \{-1, 1\}^n$ corresponding to the vertices of G. One can then write, for instance,

opt(G) =
$$\max_{x \in \{-1,1\}^n} \frac{1}{|E|} \sum_{\{i,j\} \in E} \frac{1 - x_i x_j}{2}$$
.

To convert this computation into a linear program, we need to *linearize* the objective function.

The usual way is to introduce new LP variables $y = (y_{i,j}) \in \mathbb{R}^{\binom{n}{2}}$ meant to represent the quantities $(1 - x_i x_j)/2$. Now consider the vector $v_G \in \{0, 1\}^{\binom{n}{2}}$ such that $(v_G)_{\{i,j\}} = 1$ precisely when $\{i, j\} \in E$. Given that we have linearized both the graph G and the cut variable x, we can consider the LP relaxation

$$\mathcal{L}(G) = \max_{y \in P} \langle v_G, y \rangle,$$

where P is any polytope containing all the vectors y such that $y_{i,j} = (1-x_ix_j/2)$ for some $x \in \{-1,1\}^n$. The standard relaxation corresponds to a polytope P defined by the constraints $\{0 \le y_{i,j} \le 1 : i, j \in V\}$ and

$$\{y_{i,j} \leq y_{i,k} + y_{k,j}, y_{i,j} + y_{i,k} + y_{k,j} \leq 2 : i, j, k \in V\}$$
.

Clearly P is characterized by $O(n^3)$ inequalities.

Arbitrary linearizations. But it is important to point out that, for our purposes, any linearization of the natural formulation of Max Cut suffices. We only require that there is a number $D \in \mathbb{N}$ such that:

- 1. For every graph G, we have a vector $v_G \in \mathbb{R}^D$.
- 2. For every cut $S \subseteq V$, we have a vector $y_S \in \mathbb{R}^D$.
- 3. For all graphs *G* and vectors y_S , the condition $G(S) = \langle v_G, y_S \rangle$ holds.

Now any polytope $P \subseteq \mathbb{R}^D$, such that $y_S \in P$ for every $S \subseteq V$, yields a viable LP relaxation: $\mathcal{L}(G) = \max_{y \in P} \langle v_G, y \rangle$. The *size* of this relaxation is simply the number of facets of P, i.e. the number of linear inequalities needed to specify P.

Remark 1.1. We stress that the polytope *P* depends only on the input size. This is akin to lower bounds in non-uniform models of computation like circuits wherein there is a single circuit for all inputs of a certain size. The input graph *G* is used only to define the objective function being maximized. In other words, the variables and constraints of the linear program are fixed for each input size while the objective function is defined by the input. To the best of our knowledge, all linear and semidefinite programs designed for approximating max-CSP problems are subsumed by relaxations of this nature.

In Section 3, we prove that every such relaxation of polynomial size has an integrality gap of $\frac{1}{2}$ for Max Cut. We now give an informal outline of the proof.

Proving a lower bound. In Theorem 2.3, we recall that if there is an LP relaxation \mathcal{L} of size R, then a simple application of Farkas' Lemma shows that there are non-negative functions $q_1, \ldots, q_R : \{-1, 1\}^n \to \mathbb{R}_{\geqslant 0}$ such that for every graph G, there are coefficients $\lambda_0, \lambda_1, \ldots, \lambda_R \geqslant 0$ satisfying

$$\mathcal{L}(G) - G(x) = \lambda_0 + \lambda_1 q_1(x) + \dots + \lambda_R q_R(x). \tag{1.1}$$

for all $x \in \{-1, 1\}^n$. (Note that we have earlier viewed G as a function on cuts and we now view it as a function on $\{-1, 1\}^n$ by associating these vectors with cuts.)

One should think of (1.1) as saying that $\mathcal{L}(G) - G \in \text{cone}(\mathbb{1}, q_1, q_2, \dots, q_R)$, where the latter object is the cone generated by $\{\mathbb{1}, q_1, q_2, \dots, q_R\}$ inside the Hilbert space $L^2(\{-1, 1\}^n)$ of real-valued functions, and $\mathbb{1}$ denotes the function that is identically 1. These functions $q_i : \{-1, 1\}^n \to \mathbb{R}_{\geqslant 0}$ encode the slack of each constraint of the LP. Thus if the *i*th LP constraint is of the form $\langle A_i, z \rangle \leqslant b_i$, then $q_i(x) = b_i - \langle A_i, y_{S_x} \rangle$ where y_{S_x} is the cut vector corresponding to $x \in \{-1, 1\}^n$.

Consider some $m \ll n$. The d-round Sherali–Adams relaxation for an m-vertex graph G_0 has value $SA_d(G_0) \leqslant c$ if and only if there exist a family of non-negative d-juntas $\{f_i : \{-1,1\}^m \to \mathbb{R}_{\geqslant 0}\}$ such that

$$c - G_0 = \sum_i \lambda_i f_i \,, \tag{1.2}$$

where $\lambda_i \ge 0$ for each i. We recall that a d-junta is a function whose value depends on at most d of its inputs. See Section 2.1 for an explanation of (1.2).

In particular, if G_0 is such that $SA_d(G_0) > c$, then no such representation (1.2) with d-juntas can exist. Our goal is to use (1.1) to find a graph G on n vertices such that $opt(G) = opt(G_0)$, and such that G_0 has a representation of the form (1.2) with $c = \mathcal{L}(G)$. This will show that $\mathcal{L}(G) \ge SA_d(G_0)$, completing our proof. (Recall that since we are dealing with maximization problems and $opt(G) = opt(G_0)$, this means that our LP is not doing better than Sherali–Adams.)

This proceeds in three steps: First, we argue that, by a truncation argument, it suffices to consider functions $\{q_i\}$ that are sufficiently smooth. Then in Section 3.1, we show that any sufficiently smooth q_i can be approximated (in a certain weak sense) by a K-junta q'_i for K which may be quite large (e.g., $K = n^{0.2}$).

In Section 3.2, we employ a random restriction argument: By planting the m-vertex instance G_0 at random inside a larger graph G (on n vertices), we can ensure that for every q_i' , the set of significant coordinates when restricted to G_0 is much smaller; in fact, we show that with high probability over the random planting, every such q_i' has only d significant coordinates in the support of G_0 . Here we use crucially the fact that we have only R functions $\{q_i\}$, where $R \leq n^{\alpha d}$ for some small constant $\alpha > 0$.

In particular, applying (1.1) to G and then restricting our attention to the vertices in $V(G_0)$, this yields a representation of the form

$$\mathcal{L}(G) - G_0 = \lambda_0 + \sum_{i=1}^{R} \lambda_i q_i |_{V(G_0)}, \qquad (1.3)$$

and, when restricted to G_0 , every q_i is weakly approximated by a d-junta q_i' . More specifically, all the low-degree Fourier coefficients of $q_i - q_i'$ are small. Now, the fact that (1.3) holds and each q_i is approximately a d-junta will yield that $\mathcal{L}(G) \geqslant SA_d(G_0)$, taking (1.2) into consideration. Here we remain vague, but the reader should note that this implication would follow immediately if each $q_i|_{V(G_0)}$ were actually a d-junta.

This will hold true as long as the "approximation" does not hurt us too much. One might think that our approximation is too weak: We only know that q_i' approximates q_i on $V(G_0)$ in the low-degree part. Now we use the fact that the d-round Sherali–Adams relaxation is only capable of perceiving low-degree functions (more technically, the d-round Sherali–Adams functional introduced in Section 2.1 is a degree-d multilinear polynomial). In particular, it suffices that the low-degree parts of q_i and q_i' are close.

The ingredients are all put together in Section 3.3, where one can find the proof of our main theorem for general CSPs.

2 Background

We now review the maximization versions of boolean CSPs, their linear programming relaxations, and related issues.

Throughout the paper, for a function $f: \{-1,1\}^n \to \mathbb{R}$, we write $\mathbb{E} f = 2^{-n} \sum_{x \in \{-1,1\}^n} f(x)$. If $g: \{-1,1\}^n \to \mathbb{R}$, we denote the inner product $\langle f,g \rangle = \mathbb{E}[fg]$ on the Hilbert space $L^2(\{-1,1\}^n)$. Recall that any $f: \{-1,1\}^n \to \mathbb{R}$ can be written uniquely in the Fourier basis as $f = \sum_{\alpha \subseteq [n]} \hat{f}(\alpha) \chi_{\alpha}$, where $\chi_{\alpha}(x) = \prod_{i \in \alpha} x_i$ and $\hat{f}(\alpha) = \langle f, \chi_{\alpha} \rangle$. A function f is called a d-junta for $d \in [n]$ if f depends only on a subset $S \subseteq [n]$ of coordinates with $|S| \leq d$. In other words, f can be written as $f = \sum_{\alpha \subseteq S} \hat{f}(\alpha) \chi_{\alpha}$.

We say that f is a *density* if it is non-negative and satisfies $\mathbb{E} f = 1$. For such an f, we let μ_f denote the corresponding probability measure on $\{-1,1\}^n$. Observe that for any $g: \{-1,1\}^n \to \mathbb{R}$, we have $\mathbb{E}_{x \sim \mu_f}[g(x)] = \langle f,g \rangle$.

Constraint Satisfaction Problems. Constraint satisfaction problems form a broad class of discrete optimization problems that include, for example, Max Cut and Max 3-Sat. For simplicity of presentation, we will focus on constraint satisfaction problems with a boolean alphabet, though similar ideas extend to larger domains (of constant size). One can consult [LRS15, §7].

For a finite collection $\Pi = \{P\}$ of k-ary predicates $P \colon \{-1,1\}^k \to \{0,1\}$, we let Max- Π denote the following optimization problem: An instance $\mathfrak I$ consists of boolean variables X_1, \ldots, X_n and a collection of Π -predicates $P_1(X), \ldots, P_m(X)$ over these variables. A Π -predicate is a predicate $P_0 \colon \{-1,1\}^n \to \{0,1\}$ such that $P_0(X) = P(X_{i_1}, \ldots, X_{i_k})$ for some $P \in \Pi$ and distinct indices $i_1, \ldots, i_k \in [n]$. The objective is to find an assignment $x \in \{-1,1\}^n$ that satisfies as many of the predicates as possible, that is, which maximizes

$$\mathfrak{I}(x) \stackrel{\text{def}}{=} \frac{1}{m} \sum_{i=1}^{m} P_i(x).$$

We denote the optimal value of an assignment for \mathfrak{I} as opt(\mathfrak{I}) = $\max_{x \in \{-1,1\}^n} \mathfrak{I}(x)$.

Examples: Max Cut corresponds to the case where Π consists of the binary inequality predicate. For Max 3-Sat, Π contains all eight 3-literal disjunctions, e.g., $X_1 \vee \bar{X}_2 \vee \bar{X}_3$.

Linear Programming Relaxations for CSPs. In order to write an LP relaxation for such a problem, we need to linearize the objective function. For $n \in \mathbb{N}$, let Max- Π_n be the set of Max- Π instances on n variables. An *LP-relaxation of size R* for Max- Π_n consists of the following.

Linearization: Let D be a natural number. For every $\mathfrak{I} \in \operatorname{Max-}\Pi_n$, we associate a vector $\tilde{\mathfrak{I}} \in \mathbb{R}^D$ and for every assignment $x \in \{-1,1\}^n$, we associate a point $\tilde{x} \in \mathbb{R}^D$, such that $\mathfrak{I}(x) = \langle \tilde{\mathfrak{I}}, \tilde{x} \rangle$ for all $\mathfrak{I} \in \operatorname{Max-}\Pi_n$ and all $x \in \{-1,1\}^n$.

Feasible region: A closed, convex (possibly unbounded) polyhedron $P \subseteq \mathbb{R}^D$ described by R linear inequalities, such that $\tilde{x} \in P$ for all assignments $x \in \{-1,1\}^n$. Note that the polytope P is independent of the instance \mathfrak{I} of Max-Π_n.

Given an instance $\mathfrak{I} \in Max-\Pi_n$, the LP relaxation \mathcal{L} has value

$$\mathcal{L}(\mathfrak{I}) \stackrel{\text{def}}{=} \max_{y \in P} \langle \tilde{\mathfrak{I}}, y \rangle.$$

Since $\tilde{x} \in P$ for all assignments $x \in \{-1,1\}^n$ and $\langle \tilde{\mathfrak{I}}, \tilde{x} \rangle = \mathfrak{I}(x)$, we have $\mathcal{L}(\mathfrak{I}) \geqslant \operatorname{opt}(\mathfrak{I})$ for all instances $\mathfrak{I} \in \operatorname{Max-}\Pi_n$.

Remark 2.1. For concreteness, there is a "universal linearization" for CSPs that one can always use (this is sometimes referred to as the "vertex extended formulation"). One views $x \mapsto \mathfrak{I}(x)$ as a multilinear polynomial over $\{-1,1\}^n$. In the Fourier basis $\{\chi_\alpha : \alpha \subseteq [n]\}$, one would have $\tilde{\mathfrak{I}} = \sum_\alpha \hat{\mathfrak{I}}(\alpha)\chi_\alpha$ and $\tilde{x} = \sum_\alpha \chi_\alpha(x)\chi_\alpha$. Note that if the Π_n contains k-ary predicates, then $\tilde{\mathfrak{I}}$ and \tilde{x} are multilinear polynomials of degree at most k.

Remark 2.2. Of course, in the preceding linearization, the number of variables is now 2^n . But if the number of defining inequalities small, one can reduce the number of variables via an appropriate linear transformation; see [FMP⁺12].

Symmetric Linear Programs. A symmetric LP is one for which the linearization is symmetric under any permutation of the input variables. More precisely, let us suppose \mathcal{L} is a linear program for Max- Π_n that associates to each instance \mathfrak{I} , a linearization $\tilde{\mathfrak{I}} \in \mathbb{R}^D$ and to every assignment $x \in \{-1,1\}^n$ a point $\tilde{x} \in \mathbb{R}^D$.

Let $\operatorname{Sym}(n)$ denote the symmetric group on $\{1, 2, ..., n\}$. Note that $\operatorname{Sym}(n)$ acts naturally on elements $x \in \mathbb{R}^n$ by permutation of the coordinates. Specifically, for a permutation $\sigma \in \operatorname{Sym}(n)$ and $x \in \{-1, 1\}^n$, let $\sigma x = \left(x_{\sigma(1)}, x_{\sigma(2)}, ..., x_{\sigma(n)}\right)$. This action extends to an action of $\operatorname{Sym}(n)$ on functions $f : \{-1, 1\}^n \to \mathbb{R}$ by defining $\sigma f(x) = f(\sigma x)$ for $\sigma \in \operatorname{Sym}(n)$.

We say that the linear program \mathcal{L} is *symmetric* if the following holds: For every permutation $\sigma \in \operatorname{Sym}(n)$, there exists a corresponding permutation $\tilde{\sigma} \in \operatorname{Sym}(D)$ such that for every assignment $x \in \{-1, 1\}^n$,

$$\widetilde{\sigma x} = \widetilde{\sigma}\widetilde{x}$$
.

and the feasible region $P \subset \mathbb{R}^D$ remains invariant under the permutation $\tilde{\sigma}$ of coordinates, i.e.,

$$\tilde{\sigma}P = P$$
.

To the best of our knowledge, all linear and semidefinite programming relaxations designed for approximating max-CSP problems have been symmetric relaxations. In general, assymetric relaxations could be much more powerful as demonstrated by Kaibel et al. [KPT10] who show that asymmetric LPs can be superpolynomially smaller than symmetric LPs for optimizing over log *n*-sized partial matchings.

(c,s)-approximation. For $c > s \ge 0$, we say that a linear programming relaxation \mathcal{L} for Max- Π_n is a (c,s)-approximation if $\mathcal{L}(\mathfrak{I}) \le c$ for all instances $\mathfrak{I} \in \text{Max-}\Pi_n$ with $\text{opt}(\mathfrak{I}) \le s$. We also say that \mathcal{L} achieves an α -factor approximation if $\mathcal{L}(\mathfrak{I}) \le \alpha \text{ opt}(\mathfrak{I})$ for all $\mathfrak{I} \in \text{Max-}\Pi_n$.

The following theorem is inspired by Yannakakis's characterization of exact linear programming relaxations. It appears in similar form in previous works [Pas12] and [BFPS12, Thm. 1]. For simplicity, we specialize it here for constraint satisfaction problems.

Theorem 2.3. For every $c, s \in [0, 1]$, there exists an LP relaxation of size at most R that achieves a (c, s)-approximation for Max- Π_n if and only if there exist non-negative functions $q_1, \ldots, q_R : \{-1, 1\}^n \to \mathbb{R}_{\geqslant 0}$ such that for every instance $\mathfrak{I} \in \operatorname{Max-}\Pi_n$ with $\operatorname{opt}(\mathfrak{I}) \leqslant s$, the function $c - \mathfrak{I}$ is a nonnegative combination of the functions q_1, \ldots, q_R and \mathfrak{I} , i.e.

$$c - \mathfrak{I} \in \left\{ \lambda_0 + \sum_{i=1}^R \lambda_i q_i \mid \lambda_0, \lambda_1, \dots, \lambda_R \ge 0 \right\}. \tag{2.1}$$

Moreover, if the LP relaxation is symmetric then there exist nonnegative functions $q_1, \ldots, q_R : \{-1, 1\}^n \to \mathbb{R}_{\geqslant 0}$ witnessing (2.1) and such that $\{q_1, \ldots, q_R\}$ is closed under the action of Sym(n).

Proof. First, we prove that the existence of an LP relaxation of size R yields a representation of the form (2.1). Consider a natural number D and linearizations $\tilde{\mathfrak{I}}, \tilde{x} \in \mathbb{R}^D$ for every $\mathfrak{I} \in \operatorname{Max-}\Pi_n$ and $x \in \{-1,1\}^n$. Let $P \subseteq \mathbb{R}^D$ be specified by R linear inequalities $\langle A_i, y \rangle \leq b_i$, and such that $\tilde{x} \in P$ for every $x \in \{-1,1\}^n$. We define the function $q_i : \{-1,1\}^n \to \mathbb{R}_+$ by $q_i(x) = b_i - \langle A_i, \tilde{x} \rangle$.

Consider now any instance \mathfrak{I} with opt(\mathfrak{I}) $\leq s$. By assumption, we have $\mathcal{L}(\mathfrak{I}) \leq c$, meaning that $c \geq \langle y, \tilde{\mathfrak{I}} \rangle$ holds for all $y \in P$. Now Farkas' Lemma [Sch03, Corollary 5.3c] tells us that every valid linear inequality over P can be written as a non-negative combination of the inequalities $\{b_i - \langle A_i, y \rangle \geq 0 : i = 1, 2, \dots, R\}$, and the inequality $1 \geq 0$. This yields the existence of non-negative numbers $\{\lambda_i\}$ such that $c - \langle y, \tilde{\mathfrak{I}} \rangle = \lambda_0 + \sum_{i=1}^R \lambda_i (b_i - \langle A_i, y \rangle)$ holds for all $y \in P$.

In particular, this holds for every \tilde{x} , where $x \in \{-1,1\}^n$. Now, a defining property of the linearization is that $\langle \tilde{x}, \tilde{\mathfrak{I}} \rangle = \mathfrak{I}(x)$ for every $x \in \{-1,1\}^n$. Thus we have arrived at a representation of the form (2.1).

We now show the reverse implication. Consider functions $\{q_i\}$ satisfying (2.1). We will let $D=2^n$ and the D-dimensional Hilbert space for our linearization will be $L^2(\{-1,1\}^n)$, which we identify with the linear span of the Fourier characters $\{\chi_\alpha:\alpha\subseteq[n]\}$. We use the linearization appearing in Remark 2.1. We may think of each q_i as lying in $L^2(\{-1,1\}^n)$. Define a polyhedron $P\subseteq L^2(\{-1,1\}^n)$ by $P=\{y\in L^2(\{-1,1\}^n):\langle y,q_i\rangle\geqslant 0,i=1,\ldots,R\}$. This yields an LP of size at most R since $\langle \tilde{x},q_i\rangle=q_i(x)\geqslant 0$ for every i and $x\in\{-1,1\}^n$. Now (2.1) tells us that whenever opt(\mathfrak{I}) $\leqslant s$, the inequality $\langle y,\tilde{\mathfrak{I}}\rangle\leqslant c$ is valid over P, implying that $\mathcal{L}(\mathfrak{I})\leqslant c$. Thus our LP is a (c,s)-approximation.

Finally, suppose the LP relaxation is symmetric. By definition, for every $\sigma \in \operatorname{Sym}(n)$, there exists a $\tilde{\sigma} \in \operatorname{Sym}(D)$ such that $\widetilde{\sigma x} = \tilde{\sigma} \tilde{x}$ for all $x \in \{-1, 1\}^n$ and the polytope P is invariant under the action of $\tilde{\sigma}$. We may assume that P is full-dimensional, and moreover that the facet-defining inequalities $\langle A_i, y \rangle \leq b_i$ are normalized so that $||A_i||_2 = 1$.

Consider an inequality of $\langle A_i, y \rangle \leq b_i$ of the polyhedron P and the corresponding function $q_i: \{-1,1\}^n \to \mathbb{R}$ defined by $q_i(x) = b_i - \langle A_i, \tilde{x} \rangle$. Since P is invariant under the action of $\tilde{\sigma}$, the faces of P are mapped to each other by the permutation $\tilde{\sigma}$. Now by our choice of normalization $||A_i||_2 = 1$ for i = 1, 2, ..., R, the facet-defining inequality $\langle A_i, \tilde{\sigma}y \rangle \leq b_i$ is the same as $\langle A_j, y \rangle \leq b_j$ for some $j \in \{1, 2, ..., R\}$. Hence for all $x \in \{-1, 1\}^n, q_i(\sigma x) = b_i - \langle A_i, \tilde{\sigma}\tilde{x} \rangle = b_j - \langle A_j, \tilde{x} \rangle = q_j(x)$. This implies that one can choose the family $Q = \{q_1, ..., q_R\}$ of functions to be invariant under the action of Sym(n).

A communication model. The characterization in Theorem 2.3 has an illustrative interpretation as a two-party, one-way communication complexity problem: Alice's input is a Max- Π instance $\mathfrak I$ with opt($\mathfrak I$) $\leq s$. Bob's input is an assignment $x \in \{-1,1\}^n$. Their goal is to compute the value $\mathfrak I(x)$ in expectation. To this end, Alice sends Bob a randomized message containing at most L bits. Given the

message Bob outputs deterministically a number v such that $v \le c$. The protocol is correct if for every input pair (\mathfrak{I}, x) , the expected output satisfies $\mathbb{E} v = \mathfrak{I}(x)$ (the expectation is over Alice's randomness).

An *L*-bit protocol for this communication problem yields an LP relaxation of size 2^L : If Bob outputs a value v(x, i) based on message i from Alice, then define $q_i(x) = c - v(x, i)$. This yields 2^L non-negative functions satisfying the conditions of Theorem 2.3.

On the other hand, if there exist $R = 2^L$ functions $\{q_1, q_2, ..., q_R\}$ as in Theorem 2.3, then by adding the constant function q_0 and an appropriate $\lambda_0 \ge 0$, we may assume that $\sum_{i=0}^R \lambda_i = 1$, i.e. that we have a convex combination instead of a non-negative combination. This yields a strategy for Alice and Bob: Alice sends an index $i \in \{0, 1, ..., R\}$, drawn from a distribution depending on \mathfrak{I} (specified by the coefficients $\{\lambda_i\}$), and then Bob outputs $c - q_i(x) \le c$.

Example: Suppose the optimization problem is Max Cut. In this case, Alice receives a graph G = (V, E) and Bob a cut $S \subseteq V$. If Alice sends Bob a uniformly random edge $\{u, v\} \in E$ and Bob outputs the value $|\mathbb{1}_S(u) - \mathbb{1}_S(v)|$, the result is a communication (in expectation) protocol using at most $\log_2\binom{n}{2}$ bits of communication. In this communication protocol, the value output by Bob is always at most 1. Therefore, this corresponds to a trivial (1, s)-approximation for Max Cut for every s < 1. In any protocol achieving a less trivial approximation, Bob would have to always output numbers strictly less than 1.

A similar communication in expectation model is considered in [FFGT11], where they show that the communication complexity is equal to the logarithm of the non-negative rank (up to an additive constant) of the associated slack matrix. There is an important distinction, however; their model involves communicating a slack matrix in expectation (the value $c - \Im(x)$), while the model here deals directly with the underlying combinatorial problem (the value $\Im(x)$).

2.1 Sherali-Adams LP relaxations for CSPs

A primary component of our approach involves leveraging known integrality gaps for the Sherali–Adams (SA) hierarchy. To that end, we now give a brief overview of Sherali–Adams LP relaxations. For a more detailed account, we refer the reader to [Lau03].

A d-round Sherali–Adams LP relaxation for a Max- Π_n instance will consist of variables $\{X_S : S \subseteq [n], |S| \le d\}$ for all products of up to degree-d on the n variables. These variables $\{X_S : |S| \le d\}$ are to be thought of as the moments up to degree-d of the variables, under a purported distribution.

An important property of an SA solution $\{X_S : |S| \le d\}$ is that these moments agree with a set of local marginal distributions. In particular, for every set $S \subseteq [n]$ with $|S| \le d$ there exists a distribution μ_S over $\{-1,1\}^S$ such that,

$$\mathbb{E}_{x \sim \mu_S} \chi_A(x) = X_A \qquad \forall A \subseteq S.$$

In an alternate but equivalent terminology, a d-round SA instance can be thought of as d-local expectation functional (d- ℓ .e.f.). Fix $n \ge 1$. We define a d-local expectation functional to be a linear functional $\tilde{\mathbb{E}}$ on degree-d n-variate multilinear polynomials such that $\tilde{\mathbb{E}}\mathbb{1} = 1$ and $\tilde{\mathbb{E}}P \ge 0$ for every degree-d multilinear polynomial P that is nonnegative over $\{-1,1\}^n$ and depends only on d variables. In terms of the local marginal distributions, $\tilde{\mathbb{E}}: \{-1,1\}^n \to \mathbb{R}$ is the unique linear functional satisfying

$$\tilde{\mathbb{E}} \chi_S = \mathbb{E}_{\substack{x \sim \mu_S \\ x \sim \mu_S}} \chi_S(x) \qquad \forall |S| \leq d, S \subseteq [n], \tag{2.2}$$

and $\tilde{\mathbb{E}} \chi_S = 0$ for |S| > d, $S \subseteq [n]$.

The *d-round Sherali–Adams value* of a Max- Π_n instance $\mathfrak I$ is defined as

$$SA_{d}(\mathfrak{I}) \stackrel{\text{def}}{=} \max_{d-\ell.\text{e.f. } \tilde{\mathbb{E}}} \tilde{\mathbb{E}} \mathfrak{I}. \qquad (2.3)$$

This optimization problem can be implemented by an $n^{O(d)}$ -sized linear programming relaxation for Max- Π_n . (Notice that $\tilde{\mathbb{E}}$ is a $\left(\sum_{i=0}^d \binom{n}{i}\right)$ -dimensional object.) In particular, if d-rounds of Sherali–Adams achieve a (c,s)-approximation for Max- Π_n , then so do general $n^{O(d)}$ -sized LP relaxations.

We remark that a d- ℓ .e.f. $\tilde{\mathbb{E}}$ is a linear functional, but using self-duality of $L^2(\{-1,1\}^n)$, we may also think of $\tilde{\mathbb{E}} \in L^2(\{-1,1\}^n)$. It has the Fourier representation

$$\tilde{\mathbb{E}} = \sum_{|\alpha| \leq d} \tilde{\mathbb{E}}(\chi_{\alpha}) \chi_{\alpha} .$$

We will use this representation freely.

Lemma 2.4. If $\tilde{\mathbb{E}}$ is a d- ℓ .e.f. on $L^2(\{-1,1\}^n)$, then the following properties hold:

i For any non-negative d-junta $f: \{-1,1\}^n \to \mathbb{R}_+$, we have $\tilde{\mathbb{E}} f \ge 0$.

ii For any $\alpha \subseteq [n]$, we have $|\tilde{\mathbb{E}} \chi_{\alpha}| \leq 1$.

$$iii \parallel \tilde{\mathbb{E}} \parallel_{\infty} \leq \sum_{i=0}^{d} \binom{n}{i}$$
.

Proof. Property (i) follows directly from the definition of a d- ℓ .e.f.. Property (ii) follows from (2.2). Property (iii) follows from (ii) using the fact that $\tilde{\mathbb{E}}$ has at most $\sum_{i=0}^{d} \binom{n}{i}$ non-zero Fourier coefficients.

It might help the reader, at this point to recall Theorem 2.3 and the representation (2.1). Suppose that we had such a representation where the family of functions $\{q_i\}$ were all d-juntas. Fix an instance $\mathfrak I$ and let $\tilde{\mathbb E}$ denote an optimal solution to (2.3). Applying $\tilde{\mathbb E}$ to the right-hand side of (2.1) yields

$$\widetilde{\mathbb{E}}\left[\lambda_0 + \lambda_1 q_1 + \dots + \lambda_R q_R\right] = \lambda_0 + \sum_{i=1}^R \lambda_i \,\widetilde{\mathbb{E}}\left[q_i\right] \geqslant 0,$$

using Lemma 2.4(i). On the other hand, applying it to the left-hand-side yields $\tilde{\mathbb{E}}(c-\mathfrak{I})=c-\mathsf{SA}_d(\mathfrak{I})$. Altogether, we conclude that $\mathsf{SA}_d(\mathfrak{I})\leqslant c$. In particular, this holds for any $c\geqslant \mathcal{L}(\mathfrak{I})$, hence $\mathsf{SA}_d(\mathfrak{I})\leqslant \mathcal{L}(\mathfrak{I})$, implying that in this special case (when all the q_i functions are d-juntas), the Sherali–Adams relaxation is at least as good as the given LP. In general, our approach will be to *approximate* the $\{q_i\}$ functions by juntas, and then apply a variant of this reasoning.

Remark 2.5. Some work on Sherali–Adams relaxations for Max Cut focus on edge variables instead of vertex variables. This includes [FdlVKM07, CMM09]. In those papers, the d-round Sherali–Adams relaxation consists of variables $\{X_S: S \subseteq {[n] \choose 2}, |S| \le d\}$ for every subset of d edges in the complete graph. Since their base polytope also includes triangle inequalities, any ${d \choose 2}$ -round Sherali–Adams solution with edge variables can be converted to a d-round solution for vertex variables. One should observe that the d-round vertex relaxation is at least as strong as the d-round edge relaxation.

Moreover, both papers [FdlVKM07, CMM09] actually prove a lower bound against the *d*-round vertex version and then argue that this yields a lower bound for the weaker edge relaxation. For general max-CSPs, the vertex version is arguably the canonical relaxation, and it is perhaps misguided to consider the edge version even for Max Cut. In [Sch08] (which studies general CSPs), the more natural vertex version is considered.

A major benefit of the "extended formulation" model to which our results apply is that the edge/vertex relaxation distinctions are not relevant; in fact no specific meaning is ascribed to the variables of the LP. All that matters is the number of defining inequalities.

3 Sherali-Adams and general LPs

Our main theorem is that general LP relaxations are no more powerful than Sherali–Adams relaxations (in the polynomial-size regime).

Theorem 3.1 (Main). Fix a positive number $d \in \mathbb{N}$, and a sequence of k-ary CSPs {Max- Π_n }, with $k \leq d$. Suppose that the d-round Sherali–Adams relaxation cannot achieve a (c,s)-approximation for Max- Π_n for every n. Then no sequence of LP relaxations of size at most $n^{d/2}$ can achieve a (c,s)-approximation for Max- Π_n for every n.

We prove the following result for super-polynomial sized linear programs in Section 3.3.

Theorem 3.2. Consider a function $f: \mathbb{N} \to \mathbb{N}$. Suppose that the f(n)-round Sherali–Adams relaxation cannot achieve a (c,s)-approximation for Max- Π_n . Then for all sufficiently large n, no LP relaxation of size at most $n^{f(n)^2}$ can achieve a (c,s)-approximation for Max- Π_N , where $N \leq n^{10f(n)}$.

In particular, by choosing $f(n) \times n^{\varepsilon}$ for $\varepsilon > 0$, and $n \times ((\log N)/(\log \log N))^{1/\varepsilon}$, known Sherali–Adams gaps for Max Cut [CMM09] and Max 2-Sat, Max 3-Sat [Sch08] imply the same integrality gaps for LPs of size $n^{o(\frac{\log n}{\log \log n})}$.

3.1 High-entropy distributions vs. juntas

Our first goal is to observe the following consequence of Chang's Lemma [Cha02] (and, specifically, the proof in [IMR14]).

Lemma 3.3. Let $q: \{-1,1\}^n \to \mathbb{R}_{\geqslant 0}$ be a density and let μ_q denote the corresponding measure on $\{-1,1\}^n$. If μ_q has entropy at least n-t for some $t \leqslant n$, then for every $1 \leqslant d \leqslant n$ and $\gamma > 0$, there exists a set $J \subseteq [n]$ with

$$|J| \le \frac{2td}{\gamma^2} \tag{3.1}$$

such that for all subsets $\alpha \not\subseteq J$ with $|\alpha| \leqslant d$, we have $|\hat{q}(\alpha)| \leqslant \gamma$.

Proof. Consider some $\gamma > 0$ and let $S = \{ |\alpha| \le d : |\hat{q}(\alpha)| > \gamma \}$. Let $S' \subseteq S$ denote a maximal set of linearly independent elements over \mathbb{F}_2^n . In [IMR14], it is proved that $|S'| \le 2\gamma^{-2}t$. Let $J = \bigcup_{\alpha \in S'} \alpha$ so that $|J| \le 2d\gamma^{-2}t$.

Remark 3.4. The claim in [IMR14] (namely Lemma 2 in [IMR14]) is only stated for a $q = (2^n/|A|)\mathbf{1}_A$ that is the (scaled) characteristic function of a subset $A \subseteq \{-1,1\}^n$, but the proof only uses the entropy of q. A formal statement with a somewhat different proof can be found in [LRS15, §7].

Discussion of Lemma 3.3. It is interesting to note examples for which Lemma 3.3 cannot be improved much. First, suppose that n is odd, and consider the density coming from majority on n bits:

$$q(x) = 2 \cdot \mathbf{1}_{\{x_1 + \dots + x_n > 0\}}. \tag{3.2}$$

The corresponding measure μ_q has entropy n-1. In this case, we have $|\hat{q}(\alpha)| \approx n^{-d/2}$ for $|\alpha| = d$, d odd, and $d \ll \sqrt{n}$. Thus (3.1) is essentially tight for t = d = 1.

Consider the task of obtaining $|J| = n^{1-\delta}$ and $\gamma = n^{-\omega(1)}$, for some $\delta > 0$. This is the interesting range of parameters in the next section. For the majority density (3.2), this is clearly impossible in light of our discussion. On the other hand, if one could obtain a rate of decay of the form $n^{-c(d)}$, with $c(d) \to \infty$ as $d \to \infty$ on the non-junta low-degree Fourier coefficients, then one could improve our main theorem (see (3.7)).

Unfortunately, the next example shows that this is impossible. Let $k, n \in \mathbb{N}$ be such that k divides n, and partition $\{1, 2, ..., n\} = B_1 \cup B_2 \cup \cdots \cup B_{n/k}$ into n/k disjoint blocks, each of size k. Consider the density

$$q(x) = \frac{k2^k}{n} \sum_{i=1}^{n/k} \mathbf{1}_{\{x_j = 1 \forall j \in B_i\}}.$$

This function has a transitive symmetry, and thus for k = o(n), does not admit an interesting junta set of size o(n). On the other hand for any $\alpha \subseteq B_i$, we have

$$|\hat{q}(\alpha)| = |\langle q, \chi_{\alpha} \rangle| = \frac{k}{n} 2^{k-|\alpha|}.$$

If we put $k = \sqrt{n}$, then we do not have an appreciable decay of the form $n^{-c(d)}$ with $c(d) \to \infty$ as $d \to \infty$.

But not all hope is lost: It is plainly clear that *q* can be approximated by a *non-negative combination of non-negative k-juntas*. Furthermore, an approximation of this form would be just as good for us in the arguments that follow. Thus another possible direction for improving our lower bounds significantly would be to prove a variant of Lemma 3.3 using an approximation by convex combinations of non-negative juntas, such that one achieves a strong form of decay on the Fourier coefficients.

Some improvement is possible in this case: In the setting of Lemma 3.3, one can achieve a non-negative combination of k-juntas with $k = O(td/\gamma)$ (as opposed to γ^2); see [LRS15, §7]. But this approach too reaches a bottleneck: Suppose that m divides n and partition $[n] = S_1 \cup S_2 \cup \cdots \cup S_{n/m}$ where $|S_i| = m$. Consider functions of the form $q(x) = f\left(\chi_{S_1}(x), \chi_{S_2}(x), \ldots, \chi_{S_{n/m}}(x)\right)$ where $\chi_{S}(x) = \prod_{i \in S} x_i$ is the corresponding Fourier character and $f: \{-1,1\}^{n/m} \to \mathbb{R}_{\geqslant 0}$ is a function. The effect of this operation is to lift the low-degree Fourier coefficients of f to higher-degree coefficients of f, cutting off the hope for a strong form of decay. For instance, if f and f is the majority density on f bits (as in (3.2)).

3.2 Random restrictions

We first recall the following standard estimates (see, e.g., [McD98]). Suppose X_1, \ldots, X_n are i.i.d {0, 1} random variables with $\mathbb{E}[X_i] = p$. Then,

$$\mathbb{P}\left(\sum_{i=1}^{n} X_i \geqslant \frac{pn}{2}\right) \geqslant 1 - e^{-pn/8} \,. \tag{3.3}$$

Furthermore,

$$\mathbb{P}\left(\sum_{i=1}^{n} X_{i} \geqslant t\right) \leqslant \sum_{S \in \binom{n}{t}} \mathbb{P}\left(\sum_{i \in S} X_{i} = t\right) \leqslant \binom{n}{t} \cdot p^{t} \leqslant (pn)^{t}$$
(3.4)

Lemma 3.5. For any $d \in \mathbb{N}$, the following holds. Let Q be a collection of densities $q: \{-1,1\}^n \to \mathbb{R}_{\geqslant 0}$ such that the corresponding measures μ_q have entropy at least n-t. If $|Q| \le n^{d/2}$, then for all integers m with $3 \le m \le n/4$, there exists a set $S \subseteq [n]$ such that

$$- |S| = m$$

– For each $q \in Q$, there exists a set of at most d coordinates $J(q) \subseteq S$ such that under the distribution μ_q , all d-wise correlations in S - J(q) are small. Quantitatively, we have

$$|\hat{q}(\alpha)| \le \left(\frac{16mtd}{\sqrt{n}}\right)^{1/2} \quad \forall \alpha \subseteq S, \alpha \not\subseteq J(q), |\alpha| \le d$$

Proof. We will sample the set $S \subseteq [n]$ by including each element independently with probability 2m/n, then argue that with non-zero probability, both the conditions on S hold.

First, by (3.3), we have $|S| \ge m$ with probability at least $1 - e^{-m/4} > 1/2$.

Fix $\gamma = \left(\frac{16mtd}{\sqrt{n}}\right)^{1/2}$. By Lemma 3.3, for each $q \in Q$ there exists a set J'(q) of at most $\frac{2td}{\gamma^2} \le \frac{\sqrt{n}}{8m}$ coordinates such that for all subsets $\alpha \nsubseteq J'(q)$ with $|\alpha| \le d$, we have $|\hat{q}(\alpha)| \le \gamma$.

The set J(q) for a distribution q is given by $J(q) = J'(q) \cap S$. By (3.4), we can write

$$\mathbb{P}\left[|J'(q)\cap S|\geqslant d\right]\leqslant \left(\frac{2m}{n}\cdot|J'(q)|\right)^d\leqslant \left(\frac{2m}{n}\cdot\frac{\sqrt{n}}{8m}\right)^d\leqslant \frac{1}{4^dn^{d/2}}\,.$$

The existence of the set *S* follows by taking a union bound over all the $|Q| \le n^{d/2}$ densities in the family *Q*. Note that we have concluded with $|S| \ge m$, but we can remove some elements from *S* to achieve |S| = m.

3.3 Proof of Main Theorem

In this subsection, we will prove Theorem 3.1 and Theorem 3.2. Let $m \le n$ be parameters $m, n \in \mathbb{N}$ to be chosen later. Consider an instance \mathfrak{I}_0 of Max- Π_m . Recalling (2.3), let $\tilde{\mathbb{E}}$ be a corresponding optimal d- ℓ .e.f., i.e. such that $\tilde{\mathbb{E}}[\mathfrak{I}_0] = SA_d(\mathfrak{I}_0)$.

Suppose that \mathcal{L} is an LP relaxation of size at most $R \leq n^{d/2}$ for Max- Π_n . Our goal is to show that there exists an instance \mathfrak{I} that is a "shift" of \mathfrak{I}_0 , and a value $\varepsilon_n > 0$ such that $\mathcal{L}(\mathfrak{I}) \geq \mathrm{SA}_d(\mathfrak{I}_0) - \varepsilon_n$, with $\varepsilon_n \to 0$ as $n \to \infty$. By "shift," we mean a planting of the instance \mathfrak{I}_0 on some subset of the variables $\{1, 2, \ldots, n\}$. Since $\mathrm{opt}(\mathfrak{I}) = \mathrm{opt}(\mathfrak{I}_0)$, we will conclude our proof by taking $\varepsilon_n \to 0$.

By Theorem 2.3, there are densities $q_1, q_2, \dots, q_R : \{-1, 1\}^n \to \mathbb{R}_{\geqslant 0}$ such that for every Max- Π_n instance \mathfrak{I} , we have

$$\mathcal{L}(\mathfrak{I}) - \mathfrak{I} = \lambda_0(\mathfrak{I}) + \sum_{i=1}^{R} \lambda_i(\mathfrak{I}) q_i, \qquad (3.5)$$

for some non-negative numbers $\lambda_i(\mathfrak{I})$ depending on \mathfrak{I} . For some $t \ge 0$ to be chosen later, let

$$Q_t = \{1 \le i \le R : ||q_i||_{\infty} \le 2^t\}.$$

Observing that the left-hand side of (3.5) is pointwise at most 1, for any $i \notin Q_t$, we must have $\lambda_i(\mathfrak{I}) \leq 2^{-t}$ for every instance \mathfrak{I} . At this point, one should also observe that $\sum_{i=0}^R \lambda_i(\mathfrak{I}) \leq 1$ by taking expectations over both sides of (3.5).

If $i \in Q_t$, then since $||q_i||_{\infty} \le 2^t$, we can lower bound the entropy of μ_{q_i} as follows,

$$H(\mu_{q_i}) = \sum_{x \in \{-1,1\}^n} \frac{q_i(x)}{2^n} \log \frac{2^n}{q_i(x)} \geqslant \left(\sum_{x \in \{-1,1\}^n} \frac{q_i(x)}{2^n}\right) \cdot \log \frac{2^n}{\|q_i\|_{\infty}} = \log \frac{2^n}{\|q_i\|_{\infty}} \geqslant n - t.$$

Apply Lemma 3.5 to the set of densities with index in Q_t , and let $S \subseteq [n]$ with |S| = m be the subset whose existence is guaranteed. Let \mathfrak{I}_S denote the instance \mathfrak{I}_0 planted on the subset S, and similarly let $\tilde{\mathbb{E}}_S$ be the Sherali–Adams functional $\tilde{\mathbb{E}}$ planted on S. Equation (3.5) gives us a representation of the form

$$\mathcal{L}(\mathfrak{I}_S) - \mathfrak{I}_S = \lambda_0(\mathfrak{I}_S) + \sum_{i=1}^R \lambda_i(\mathfrak{I}_S) q_i, \qquad (3.6)$$

For each $i \in Q_t$, let $q_i^S = \sum_{\alpha \subseteq S} \hat{q}_i(\alpha) \chi_\alpha$. Observe that q_i^S is the conditional density on the variables in S (equivalently, we obtain q_i^S by averaging over all variables outside S). By our application of Lemma 3.5, we can write $q_i^S = \tilde{q}_i^S + e_i$ where $\tilde{q}_i^S = \sum_{\alpha \subseteq J(q_i) \cap S} \hat{q}_i(\alpha) \chi_\alpha$ is a non-negative d-junta and $|\hat{e}_i(\alpha)| \le \left(\frac{16mtd}{\sqrt{n}}\right)^{1/2}$ for all $|\alpha| \le d$.

Using the fact that $\tilde{\mathbb{E}}_S$ only depends on variables in S, we have $\tilde{\mathbb{E}}_S(q_i) = \tilde{\mathbb{E}}_S(q_i^S)$ for all $i \in Q_t$. Also observe that for $i \notin Q_t$, we have $|\tilde{\mathbb{E}}_S(q_i)| \le |\tilde{\mathbb{E}}_S||_{\infty} \le {m \choose d}$, first using $\mathbb{E} q_i = 1$ and then using property (iii) of Lemma 2.4. Now we apply $\tilde{\mathbb{E}}_S$ to both sides of (3.6) to obtain

$$\mathcal{L}(\mathfrak{I}_{S}) - \mathsf{SA}_{d}(\mathfrak{I}_{0}) = \lambda_{0}(\mathfrak{I}_{S}) + \sum_{i \in Q_{t}} \lambda_{i}(\mathfrak{I}_{S}) \tilde{\mathbb{E}}_{S}(q_{i}^{S}) + \sum_{i \notin Q_{t}} \lambda_{i}(\mathfrak{I}_{S}) \tilde{\mathbb{E}}_{S}(q_{i})$$

$$\geqslant \sum_{i \in Q_{t}} \lambda_{i}(\mathfrak{I}_{S}) \left(\tilde{\mathbb{E}}_{S}(\tilde{q}_{i}^{S}) + \tilde{\mathbb{E}}_{S}(e_{i})\right) - ||\tilde{\mathbb{E}}_{S}||_{\infty} \sum_{i \notin Q_{t}} \lambda_{i}(\mathfrak{I}_{S})$$

$$\geqslant \sum_{i \in Q_{t}} \lambda_{i}(\mathfrak{I}_{S}) \tilde{\mathbb{E}}_{S}(e_{i}) - \binom{m}{d} n^{d/2} 2^{-t},$$

where in the final line we have used the fact that \tilde{q}_i^S is a non-negative d-junta (along with property (i) of Lemma 2.4), the fact that $\lambda_i(\mathfrak{I}_S) \leq 2^{-t}$ for $i \notin Q_t$, and our assumption that the total number of indices $i \in \{1, 2, ..., R\}$ is at most $n^{d/2}$.

Finally, it remains to observe that

$$|\tilde{\mathbb{E}}_{S}(e_{i})| \leq \sum_{\alpha \subseteq S} |\tilde{\mathbb{E}}_{S}[\chi_{\alpha}]| \cdot |\hat{e}_{i}(\alpha)| \leq {m \choose d} \left(\frac{16mtd}{\sqrt{n}}\right)^{1/2},$$
 (3.7)

where we have employed property (ii) of Lemma 2.4. Plugging this estimate into the preceding inequality yields

$$\mathcal{L}(\mathfrak{I}_S) - \mathsf{SA}_d(\mathfrak{I}_0) \geqslant -\binom{m}{d} \left(\frac{16mtd}{\sqrt{n}}\right)^{1/2} - \binom{m}{d} n^{d/2} 2^{-t}.$$

If we set $t = d \log_2 n$, then $\mathcal{L}(\mathfrak{I}_S) \ge SA_d(\mathfrak{I}_0) - \varepsilon_n$, where

$$\varepsilon_n = O\left(\frac{m^d \sqrt{md \log n}}{n^{1/4}}\right). \tag{3.8}$$

Clearly for m,d fixed, we have $\varepsilon_n \to 0$ as $n \to \infty$, completing the proof of Theorem 3.1.

Proof of Theorem 3.2. Fix an instance size m and put d = f(m). In the preceding argument, require that n grows like $m^{10d} = m^{10f(m)}$ so that $\varepsilon_n = o(1)$ (see (3.8)). The lower bound achieved is $n^{d/2} \ge m^{5f(m)^2}$.

3.4 Nonnegative rank

The lower bounds of Theorem 3.1 can be stated equivalently in terms of nonnegative rank. We recall that the nonnegative rank of a nonnegative matrix $A \in \mathbb{R}_+^{m \times n}$ is defined by

$$\operatorname{rank}_+(A) = \min \left\{ r : A_{ij} = \langle u_i, v_j \rangle \text{ for some } \{u_i, v_j\} \subseteq \mathbb{R}_+^r \right\}.$$

Fix $n \ge 1$. Let $M = (M_{G,x})$ be the matrix indexed by n-vertex Max Cut instances with Max Cut value at most s (e.g., $s = 1/2 + \gamma$) and bipartitions $x \in \{\pm 1\}^n$ such that

$$M_{G,x}=c-G(x)\,,$$

where G(x) denotes the fraction of edges crossing the bipartition corresponding to x. A corollary of Theorem 3.1 is that $\operatorname{rank}_+(M) \ge n^{\Omega\left(\frac{\log n}{\log \log n}\right)}$.

Define for $\varepsilon > 0$, the ε -smooth nonnegative rank of a matrix A as

$$\operatorname{rank}_{+,\varepsilon}(A) := \min \left\{ \operatorname{rank}_{+}(A') \mid (1-\varepsilon)A_{ij} \leqslant A'_{ij} \leqslant (1+\varepsilon)A_{ij} \ \forall i,j \right\},\,$$

Our main result shows that M has superpolynomial nonnegative rank. We claim that M has only polynomial nonnegative approximate rank for every fixed $\varepsilon > 0$. (Since the entries of M are bounded from above and bounded away from 0, the notions of approximate and smooth nonnegative rank coincide.) In order to demonstrate that M has small approximate nonnegative rank, for each $t \in \mathbb{N}$, we will exhibit a matrix M' that approximates M well, $\left|M'_{G,x} - M_{G,x}\right| \leqslant 2^{-\Omega(t)}$ for all G and x, but has a small nonnegative rank, i.e., $\operatorname{rank}_+(M') \leqslant n^{O(t)}$. To this end, we will use the reformulation of nonnegative rank as a communication model discussed in Section 2. Consider the following communication protocol between Alice and Bob:

- Alice receives as input an *n*-vertex graph *G* with Max Cut value at most *s*.
- Bob receives as input a bipartition x ∈ {±1} n .
- Alice chooses t edges e_1, \ldots, e_t of G independently at random and sends the endpoints of the sampled edges to Bob.
- Bob computes what fraction θ of the edges e_1, \dots, e_t cross the bipartition x. If $\theta > c$, then Bob outputs 0 else Bob outputs $c \theta$.

Let M' be the matrix computed by the above protocol, i.e., $M'_{G,x}$ is the expected value of Bob's output when Alice receives the graph G and Bob receives x as input. This protocol yields a rank- $2^{t\cdot 2\log n}$ nonnegative factorization of the matrix M', as follows.

Let $\theta_{G,x}$ be the random variable given by Bob's output when the inputs for Alice and Bob are G and x, respectively. Then, $M'_{G,x} = c - \mathbb{E} \theta_{G,x}$ for all G and x.

At the same time, we have

$$M_{G,x} = c - \mathbb{E}\left[\theta_{G,x}\right] - \delta_{G,x},$$

where

$$\delta_{G,x} \leq 2 \mathbb{P}[\theta_{G,x} > c]$$
.

In words, the discrepancy between the computed matrix M' and the target matrix M is accounted for by the probability of the events $\{\theta_{G,x} > c\}$. However, since $G(x) \le s < c$ is bounded away from c by some constant, a standard Chernoff bound yields

$$\mathbb{P}[\theta_{G,x}>c] \leq 2^{-\Omega(t)}\,,$$

for all G and x. It follows that the matrix M' satisfies $\operatorname{rank}_+(M') \leq n^{O(t)}$ and $\max_{G,x}|M'_{G,x}-M_{G,x}| \leq 2^{-\Omega(t)}$.

4 Symmetric linear programs

We will now prove the following theorem relating Sherali–Adams gaps to those for symmetric LPs for Max Cut. While this connection holds more generally for max-CSP problems, we will focus on Max Cut for clarity. Recent work has extended these ideas to problems like TSP [LRST14], and also to a connection between symmetric SDPs and the Sum-of-Squares hierarchy [LRST14, FSP13].

Theorem 4.1. Fix a k-ary CSP Max- Π over the boolean domain. Suppose that, for some numbers m, d > 0, the d-round Sherali–Adams relaxation for Max- Π_m cannot achieve a (c,s)-approximation. Then no symmetric LP of size $\leq \binom{n}{d}$ can achieve a (c,s)-approximation on Max- Π_n where n = 2m.

We note here that the Sherali-Adams hierarchy produces symmetric linear programs. Hence, the above result can be viewed as asserting that Sherali-Adams hierarchy is *complete* for the class of symmetric linear programs.

By appealing to the known Sherali–Adams gaps for Max Cut [CMM09] and Max 2-Sat, Max 3-Sat [Sch08], we get the same integrality gaps for arbitrary symmetric LPs. For example, in the case of Max Cut, we obtain the following lower bound.

Corollary 4.2. For every $\varepsilon > 0$, there exists $\delta > 0$ such that no symmetric linear program of size $2^{n^{\delta}}$ yields a $(1 - \varepsilon, \frac{1}{2} + \varepsilon)$ -approximation for Max Cut.

In order to prove Theorem 4.1, we will need the following characterization of symmetric function families.

Lemma 4.3. Suppose a family of functions $Q = \{q_i : \{-1,1\}^n \to \mathbb{R} : i = 1,2,...,R\}$ is closed under the action of Sym(n). If $R < \binom{n}{d}$ for d < n/4, then each function q_i depends only on a subset $J_i \subseteq [n]$ of at most d coordinates and possibly the value of the sum $\sum_{i=1}^n x_i$.

Proof. Here we will need a few basic notions about group actions. A group \mathcal{G} *acts* on a universe \mathcal{X} , if each element $g \in \mathcal{G}$ *permutes* the elements of the universe \mathcal{X} , and this action commutes with the group operation. Formally, a group action is defined by a map $\iota : \mathcal{G} \times \mathcal{X} \to \mathcal{X}$ such that $\iota(g, \iota(h, x)) = \iota(gh, x)$ for all $g, h \in \mathcal{G}$ and $x \in \mathcal{X}$. For convenience, we will denote $g \cdot x \stackrel{\text{def}}{=} \iota(g, x)$.

For an element $x \in X$, its orbit $\operatorname{Orb}(x)$ is given by $\operatorname{Orb}(x) = \{g \cdot x \mid g \in G\}$ and its stabilizer is given by $\operatorname{Stab}(x) = \{g \in G \mid g \cdot x = x\}$. A basic fact from group theory is that for every action of a finite group G and every G and every G and G it holds that $|\operatorname{Stab}(x)| \cdot |\operatorname{Orb}(x)| = |G|$.

The group $\operatorname{Sym}(n)$ of all permutations on n elements acts on the space of functions over $\{-1,1\}^n$ by permutation of the coordinates. Let $\operatorname{Orb}(f)$ denote the orbit of a function f under the action of $\operatorname{Sym}(n)$, and let $\operatorname{Stab}(f)$ denote the stabilizer of f. Since Q is closed under this action, it contains the orbits of each of the functions g_1, \ldots, g_R .

This implies that for each $i \in [R]$, we have $|\operatorname{Orb}(q_i)| < \binom{n}{d}$. Since $|\operatorname{Orb}(q_i)| \cdot |\operatorname{Stab}(q_i)| = |\operatorname{Sym}(n)| = n!$, we conclude that for each $i \in [R]$, $|\operatorname{Stab}(q_i)| \ge d!(n-d)!$. At this point, we appeal to the following group theoretic fact that we borrow from the work of Yannakakis [Yan91].

Lemma 4.4 ([Yan91, Claim 2]). Let H be a group of permutations whose index in $\operatorname{Sym}(n)$ is at most $\binom{n}{d}$ for some d < n/4. Then there exists a set $J \subseteq \{1, 2, ..., n\}$ of size at most d such that H contains all even permutations that fix the elements of J.

By Lemma 4.4, the stabilizer subgroup $\operatorname{Stab}(q_i)$ contains all even permutations that fix a subset of coordinates J_i with $|J_i| \leq d$. We claim that $\operatorname{Stab}(q_i)$ contains all permutations that fix the coordinates in J_i . We know that for every $x \in \{-1,1\}^n$, and every even permutation $\sigma \in \operatorname{Sym}(\bar{J}_i)$ we have, $q_i(x) = q_i(\sigma x)$. Here, we use $\operatorname{Sym}(\bar{J}_i)$ to denote the subgroup of $\operatorname{Sym}(n)$ fixing elements in J_i .

For every $x \in \{-1,1\}^n$, there will be two coordinates $a,b \in \overline{J}_i$ such that $x_a = x_b$. Let π_{ab} denote the transposition that swaps a and b. Since $\pi_{ab}(x) = x$, we have $q_i(\pi_{ab}(x)) = q_i(x)$. So for even permutations $\sigma \in \text{Sym}(\overline{J}_i)$,

$$q_i(\sigma \pi_{ab} x) = q_i(\pi_{ab} x) = q_i(x).$$

As σ varies over all even permutations in $\text{Sym}(\bar{J}_i)$, $\sigma \pi_{ab}$ varies over all odd permutations in $\text{Sym}(\bar{J}_i)$, leading to the conclusion that $\text{Sym}(\bar{J}_i) \subseteq \text{Stab}(q_i)$.

This symmetry of the function $q_i(x)$ implies that it depends only on the assignment to coordinates in J_i and the hamming weight of the assignment to coordinates in \bar{J}_i , i.e. the value $\sum_{j=1}^n x_j - \sum_{j \in J_i} x_j$. This shows that q_i is a function depending only on the coordinates in J_i and the value $\sum_{i \in [n]} x_i$.

We are now in position to prove the main theorem of this section.

Proof of Theorem 4.1. Let \mathcal{L} be a symmetric LP relaxation for Max- Π_n of size $R \leq \binom{n}{d}$. Supposing that this relaxation achieves a (c,s)-approximation, we will derive a contradiction.

By applying Theorem 2.3, there exists a family of functions $Q = \{q_1, \ldots, q_R : \{-1, 1\}^n \to \mathbb{R}_{\geqslant 0}\}$ such that for every instance \mathfrak{I} of Max- Π_n with opt(\mathfrak{I}) $\leqslant s$, we have

$$c - \mathfrak{I} = \lambda_0 + \sum_{i=1}^R \lambda_i q_i.$$

for some non-negative weights $\{\lambda_i\}_{i=0}^R$. Moreover, the family of functions Q is invariant under the action of $\operatorname{Sym}(n)$. Therefore, by Lemma 4.3, each of the functions $q_i \in Q$ depends on a set J_i of at most d coordinates and possibly the value $\sum_{i=1}^n x_i$.

Fix an instance \mathfrak{I} of Max- Π_m on which the d-round Sherali–Adams relaxation fails to achieve a (c, s)-approximation, i.e.,

$$\operatorname{opt}(\mathfrak{I}) \leq s$$
 and $\operatorname{SA}_d(\mathfrak{I}) > c$.

For n = 2m, construct an instance \mathfrak{I}' of Max- Π_n by including m additional dummy variables in \mathfrak{I} with no constraints among them. Concretely, if X_1, \ldots, X_n are variables in \mathfrak{I}' , then restricted to the variables X_1, \ldots, X_m , the constraints are identical to \mathfrak{I} while there are no constraints among X_{m+1}, \ldots, X_n .

For an assignment $x \in \{-1,1\}^n$, we will denote $x_A = (x_1, ..., x_m)$ and $x_B = (x_{m+1}, ..., x_n)$. In this notation, it is easy to see that for every assignment x,

$$\mathfrak{I}'(x)=\mathfrak{I}(x_A)\,.$$

By construction, we have $\operatorname{opt}(\mathfrak{I}') = \operatorname{opt}(\mathfrak{I}) \leqslant s$. Since the symmetric LP relaxation \mathcal{L} yields a (c,s)-approximation to $\operatorname{Max-}\Pi_n$, there exist $\{\lambda_i \geqslant 0\}_{i=0}^R$ such that

$$c - \mathfrak{I}' = \lambda_0 + \sum_{i=1}^R \lambda_i q_i.$$

Using $\Im(x) = \Im(x_A)$, we can rewrite the above identity as,

$$c - \mathfrak{I}(x_A) = \lambda_0 + \sum_{i=1}^R \lambda_i q_i(x_A, x_B).$$

Define $h_i: \{-1,1\}^m \to \mathbb{R}_{\geq 0}$ as $h_i(x) = q_i(x,-x)$. Setting $x_B = -x_A$ in the above identity, we arrive at

$$c - \mathfrak{I}(x_A) = \lambda_0 + \sum_i \lambda_i h_i(x_A). \tag{4.1}$$

Recall that each of the functions q_i depends on a subset J_i of at most d coordinates and possibly the value of $\sum_{i=1}^{n} x_i$. This implies that $h_i(x) = q_i(x, -x)$ is a d-junta, since the sum of all the coordinates of (x, -x) is always equal to 0. In particular, the identity in (4.1) expresses the function $c - \mathfrak{I}$ as a non-negative combination of d-juntas.

Let $\tilde{\mathbb{E}}$ denote the d-round Sherali–Adams functional for the instance \mathfrak{I} . Apply the $\tilde{\mathbb{E}}$ functional to both sides of (4.1) to obtain a contradiction. By Lemma 2.4 and the fact that each h_i is a non-negative d-junta, we have $\tilde{\mathbb{E}}[h_i] \geqslant 0$. On the other hand, the left hand side $\tilde{\mathbb{E}}(c-\mathfrak{I}) = c - \mathsf{SA}_d(\mathfrak{I}) < 0$.

5 Conclusion

We have shown that for constraint satisfaction problems, there is an intimate relationship between general polynomial-sized linear programs and those arising from O(1) rounds of the Sherali–Adams hierarchy. There are a few natural questions that readily suggest themselves.

Firstly, our quantitative bounds are far from optimal. For instance, it is known that the integrality gap of $1/2 + \varepsilon$ for Max Cut persists for $n^{c_{\varepsilon}}$ rounds of Sherali-Adams hierarchy, where c_{ε} is some constant depending on ε [CMM09], while we are only able to prove an integrality gap for LPs of size $n^{o(\frac{\log n}{\log \log n})}$. This is due to the factor of m^d appearing in our Fourier estimate (3.7).

Question 5.1. Is it the case that for approximating (boolean) max-CSP problems on n variables, linear programs of size R(n) are only as powerful as those arising from poly($\frac{\log R(n)}{\log n}$) rounds of the Sherali–Adams hierarchy?

Secondly, given the connection for linear programs, it is natural to suspect that a similar phenomenon holds for SDPs.

Question 5.2. For max-CSP problems, is there a connection between the efficacy of general SDPs and those from the Sum-of-Squares SDP hierarchy [Las01, Par00]?

As mentioned in the introduction, recent work [LRS15] yields a positive solution to this question, although the approach has similar limitations to those highlighted in Question 5.1.

Finally, our techniques have made very strong use of the product structure on the space of feasible assignments for CSPs. One might hope to extend these connections to other types of problems like TSP and finding maximum-weight perfect matchings in general graphs [Rot14, Yan91] or approximations for vertex cover. See [BPZ15] for progress on the latter problem.

Acknowledgements

We thank the anonymous referees for many useful suggestions and observations. S. O. Chan was supported by NSF grants CCF-1118083 and CCF-1017403. P. Raghavendra was supported by NSF Career Award CCF-1343104 and an Alfred P. Sloan Fellowship. J. R. Lee was supported by NSF grants CCF-1217256 and CCF-0905626. D. Steurer was supported by NSF grants, an Alfred P. Sloan Fellowship, and a Microsoft Research Faculty Fellowship.

References

- [ABL02] Sanjeev Arora, Béla Bollobás, and László Lovász, *Proving integrality gaps without knowing the linear program*, FOCS, 2002, pp. 313–322. 1, 3
- [ABLT06] Sanjeev Arora, Béla Bollobás, László Lovász, and Iannis Tourlakis, Proving integrality gaps without knowing the linear program, Theory of Computing 2 (2006), 19–51. 3
- [BFPS12] Gábor Braun, Samuel Fiorini, Sebastian Pokutta, and David Steurer, *Approximation limits of linear programs (beyond hierarchies)*, FOCS, 2012, pp. 480–489. 2, 3, 4, 9
- [BGMT12] Siavosh Benabbas, Konstantinos Georgiou, Avner Magen, and Madhur Tulsiani, *SDP gaps from pairwise independence*, Theory of Computing 8 (2012), no. 12, 269–289. 4
- [BM13] Mark Braverman and Ankur Moitra, *An information complexity approach to extended formulations*, STOC, ACM, 2013, pp. 161–170. 3,
- [BPZ15] Gábor Braun, Sebastian Pokutta, and Daniel Zink, *Inapproximability* of combinatorial problems via small LPs and SDPs, STOC (New York, NY, USA), ACM, 2015, pp. 107–116. 3, 23

- [Cha02] Mei-Chu Chang, *A polynomial bound in Freiman's theorem*, Duke Math. J. **113** (2002), no. 3, 399–419. MR 1909605 (2003d:11151) **14**
- [CMM09] Moses Charikar, Konstantin Makarychev, and Yury Makarychev, Integrality gaps for Sherali-Adams relaxations, STOC, ACM, 2009, pp. 283–292. 3, 13, 14, 20, 22, 27, 28
- [FdIVKM07] Wenceslas Fernández de la Vega and Claire Kenyon-Mathieu, *Linear programming relaxations of Maxcut*, SODA, 2007, pp. 53–61. 3, 13, 27, 28
- [FFGT11] Yuri Faenza, Samuel Fiorini, Roland Grappe, and Hans Raj Tiwary, Extended formulations, non-negative factorizations and randomized communication protocols, arXiv:1105.4127, 2011. 11
- [FMP+12] Samuel Fiorini, Serge Massar, Sebastian Pokutta, Hans Raj Tiwary, and Ronald de Wolf, *Linear vs. semidefinite extended formulations: exponential separation and strong lower bounds*, STOC, 2012, pp. 95–106. 3, 4, 8
- [FSP13] Hamza Fawzi, James Saunderson, and Pablo A. Parrilo, *Equivariant* semidefinite lifts and sum-of-squares hierarchies, arXiv:1312.6662, 2013. 2, 19
- [Gri01] Dima Grigoriev, *Linear lower bound on degrees of Positivstellensatz calculus proofs for the parity*, Theoret. Comput. Sci. **259** (2001), no. 1-2, 613–622. MR 1832812 (2002e:03093) 4
- [GW95] Michel X. Goemans and David P. Williamson, *Improved approximation algorithms for maximum cut and satisfiability problems using semidefinite programming*, Journal of the ACM **42** (1995), 1115–1145.
- [Hås01] Johan Håstad, *Some optimal inapproximability results*, Journal of the ACM **48** (2001), no. 4, 798–859. MR 2144931 (2006c:68066) 2
- [IMR14] Russell Impagliazzo, Cristopher Moore, and Alexander Russell, An entropic proof of Chang's inequality, SIAM J. Discrete Math. 28 (2014), no. 1, 173–176. MR 3162401 14
- [KMSY14] Gillat Kol, Shay Moran, Amir Shpilka, and Amir Yehudayoff, *Approximate nonnegative rank is equivalent to the smooth rectangle bound.*, Electronic Colloquium on Computational Complexity (ECCC), vol. 21, 2014, p. 46. 4
- [KPT10] Volker Kaibel, Kanstantsin Pashkovich, and Dirk Oliver Theis, Symmetry matters for the sizes of extended formulations, IPCO, 2010, pp. 135–148. 9

- [Las01] Jean B. Lasserre, Global optimization with polynomials and the problem of moments, SIAM J. Optim. 11 (2000/01), no. 3, 796–817. MR 1814045 (2002b:90054) 22
- [Lau03] Monique Laurent, A comparison of the Sherali-Adams, Lovász-Schrijver, and Lasserre relaxations for 0-1 programming, Math. Oper. Res. (2003), 470–496. 3, 11
- [LRS15] James R. Lee, Prasad Raghavendra, and David Steurer, *Lower bounds on the size of semidefinite programming relaxations*, STOC (New York, NY, USA), ACM, 2015, pp. 567–576. 2, 7, 14, 15, 23
- [LRST14] James R. Lee, Prasad Raghavendra, David Steurer, and Ning Tan, On the power of symmetric LP and SDP relaxations, CCC, IEEE, 2014. 2, 19
- [LS91] László Lovász and Alexander Schrijver, Cones of matrices and setfunctions and 0-1 optimization, SIAM J. Optim. 1 (1991), 166–190.
- [McD98] Colin McDiarmid, Concentration, Probabilistic methods for algorithmic discrete mathematics, Algorithms Combin., vol. 16, Springer, Berlin, 1998, pp. 195–248. MR 1678578 (2000d:60032) 15
- [Par00] Pablo Parrilo, Structured semidefinite programs and semialgebraic geometry methods in robustness and optimization, Ph.D. thesis, California Institute of Technology, 2000. 22
- [Pas12] Kanstantsin Pashkovich, Extended formulations for combinatorial polytopes, Ph.D. thesis, Magdeburg Universität, 2012. 9
- [Rot14] Thomas Rothvoß, *The matching polytope has exponential extension complexity*, STOC (New York, NY, USA), ACM, 2014, pp. 263–272. 3, 4, 23
- [SA90] Hanif D. Sherali and Warren P. Adams, *A hierarchy of relaxations between the continuous and convex hull representations for zero-one programming problems*, SIAM J. Discrete Math. **3** (1990), 411–430. **1**, 27, 28
- [Sch03] Alexander Schrijver, Combinatorial optimization. Polyhedra and efficiency. Vol. A, Algorithms and Combinatorics, vol. 24, Springer-Verlag, Berlin, 2003, Paths, flows, matchings, Chapters 1–38. MR 1956924 (2004b:90004a) 10
- [Sch08] Grant Schoenebeck, *Linear level Lasserre lower bounds for certain k-CSPs*, FOCS, IEEE, 2008, pp. 593–602. 4, 13, 14, 20

- [Tre12] Luca Trevisan, *Max cut and the smallest eigenvalue*, SIAM J. Comput. **41** (2012), no. 6, 1769–1786. MR 3029271 2
- [Vaz01] Vijay V. Vazirani, *Approximation algorithms*, Springer-Verlag, Berlin, 2001. MR 1851303 (2002h:68001) 1
- [WS11] David P. Williamson and David B. Shmoys, *The design of approximation algorithms*, Cambridge University Press, Cambridge, 2011.
- [Yan91] Mihalis Yannakakis, *Expressing combinatorial optimization problems by linear programs*, J. Comput. System Sci. **43** (1991), no. 3, 441–466. 1, 2, 3, 20, 23

A What is Sherali-Adams?

Our definition of Sherali–Adams relaxation differs from the definition in prior works (in particular, the works that proved lower bounds on the size of Sherali-Adams relaxations for approximating CSPs) [SA90, FdlVKM07, CMM09]. This discrepancy stems from the fact that traditionally LP hierarchies like Sherali-Adams are applied to integer linear programming formulations of a problem, whereas our relaxations can be viewed as applying the analogous reasoning to a more direct formulation of the problem. It turns out that the latter approach typically leads to relaxations that are easier to describe and a-priori more powerful.

We will argue that the two versions of Sherali–Adams are equivalent for the problems we consider, in the sense that each relaxation in one hierarchy is captured by a relaxation in the other hierarchy of comparable size (at most a polynomial factor more constraints).

We remark that our relaxations are equivalent to the viewpoint of Sherali–Adams as a collection of mutually-consistent "local distributions over assignments." This viewpoint was used in previous works for proving lower bounds. These previous works show that this viewpoint captures the power of Sherali–Adams. We will argue that this viewpoint is indeed equivalent to the Sherali–Adams hierarchy.

A.1 Edge-based Sherali–Adams relaxations for Max Cut

The cut polytope CUT_n is the convex hull of all vectors $y \in \{0,1\}^{\binom{n}{2}}$ such that there exists a bipartition $x \in \{\pm 1\}^n$ with $y_{i,j} = \mathbb{1}_{\{x_i \neq x_i\}}$ for all $i \neq j \in [n]$. We can formulate Max Cut as the problem of optimizing a linear function of the form $\sum_{ij \in E(G)} y_{i,j}$ over CUT_n for a graph G. The standard LP relaxation of CUT_n is the *metric polytope* METRIC_n, which consists of all vectors $y \in [0, 1]^{\binom{n}{2}}$ that satisfy the inequalities $y_{i,j} \le y_{i,k} + y_{k,j}$ and $y_{i,j} + y_{i,k} + y_{k,j} \le 2$ for all $i, j, k \in [n]$. This $O(n^3)$ size LP relaxation corresponds to an exact integer linear programming (ILP) formulation in the sense that the convex hull of the integer vectors METRIC_n \cap $\{0,1\}_{2}^{\binom{n}{2}}$ is precisely the cut polytope CUT_n. In our notation, the level-r Sherali-Adams relaxation of this ILP formulation consists of all linear functionals $\tilde{\mathbb{E}}$ on $L_2(\{0,1\}^{\binom{n}{2}})$ such that $\tilde{\mathbb{E}}1=1$ and $\tilde{\mathbb{E}}f\cdot\ell\geqslant 0$ for every non-negative *r*-junta f on $\{0,1\}^{\binom{n}{2}}$ and every linear function ℓ on $\{0,1\}^{\binom{n}{2}}$ corresponding to one of the defining linear inequalities of METRIC_n, i.e., ℓ is of the form $y_{i,j}$, $1 - y_{i,j}$, $y_{i,k} + y_{k,j} - y_{i,j}$, or $2 - y_{i,j} - y_{i,k} - y_{k,j}$ for some $i, j, k \in [n]$. The value of the level-rSherali-Adams relaxation for a Max Cut instance G is the maximum value of $\tilde{\mathbb{E}} \sum_{i \in E(G)} y_{i,j}$ over all linear functionals $\tilde{\mathbb{E}}$ that satisfy the previous conditions. (From our description it is not immediately clear that this optimization problem has a small linear programming formulation. However note that $n^{O(r)}$ linear

inequalities are enough to define the set of all admissible linear functionals $\tilde{\mathbb{E}}$. Hence, we can reduce this problem to a linear program of size $n^{O(r)}$. It's also possible, but somewhat cumbersome, to describe this small linear program explicitly [SA90, FdIVKM07, CMM09].)

A.1.1 Why is this hierarchy of relaxations equivalent to the previously described hierarchy?

Let G be any graph. First, consider any k-local pseudo-expectation $\tilde{\mathbb{E}}_x$ as defined before. We will construct an equivalent linear functional $\tilde{\mathbb{E}}_y$ for the level-r Sherali–Adams relaxation with r = k/2 - 3. Recall that $\tilde{\mathbb{E}}_x$ is a linear functional on $L_2(\{\pm 1\}^n)$ such that $\tilde{\mathbb{E}}_x$ 1 = 1 and $\tilde{\mathbb{E}}_x$ $f \ge 0$ for every nonnegative k-junta f. We define a linear functional $\tilde{\mathbb{E}}_y$ on $L_2(\{0,1\}^{\binom{n}{2}})$ as follows,

$$\tilde{\mathbb{E}}_{y} f = \tilde{\mathbb{E}}_{x}(f \circ \varphi),$$

Next, consider any linear functional $\tilde{\mathbb{E}}_y$ for the level-r Sherali–Adams relaxation. We will construct an equivalent k-local pseudo-expectation $\tilde{\mathbb{E}}_x$ for k=r. We define $\tilde{\mathbb{E}}_x$ as follows,

$$\tilde{\mathbb{E}}_x f = \tilde{\mathbb{E}}_y (f \circ \psi) \,,$$

where $\psi(y)_i = 1$ if $y_{i,1} = 0$ and $\psi(y)_i = -1$ if $y_{i,1} = 1$. In words, we assign 1 to all vertices i on the same side of the bipartition as vertex 1 and -1 to all vertices on the other side. (Algebraically, ψ is defined by $\psi(y)_i = 1 - 2y_{i,1}$.) This linear functional satisfies $\tilde{\mathbb{E}}_x 1 = \tilde{\mathbb{E}}_y 1 = 1$ and $\tilde{\mathbb{E}}_x f \ge 0$ for every nonnegative k-junta f (because $f \circ \psi$ is also a nonnegative k-junta). It remains to show that $\tilde{\mathbb{E}}_x \sum_{ij \in E(G)} (1 - x_i x_j)/2 = \tilde{\mathbb{E}}_y \sum_{ij \in E(G)} y_{i,j}$. By our construction of the functional $\tilde{\mathbb{E}}_x$, we have $\tilde{\mathbb{E}}_x (1 - x_i x_j)/2 = \tilde{\mathbb{E}}_y (y_{i,1} - y_{j,1})^2$ and thus it's enough to establish $\tilde{\mathbb{E}}_y (y_{i,1} - y_{j,1})^2 - y_{i,j} = 0$. To simplify notation let us assume i = 2 and j = 3. Let $\mathbb{1}_{000}, \ldots, \mathbb{1}_{111}$ be the indicators for the eight possible assignments for the variables $y_{1,2}, y_{1,3}, y_{2,3}$. Since $1 = \mathbb{1}_{000} + \cdots + \mathbb{1}_{111}$ as functions over $\{0,1\}^{\binom{n}{2}}$, it is enough

to verify that $\tilde{\mathbb{E}}_y \mathbb{1}_{abc} \cdot \left((y_{1,2} - y_{1,3})^2 - y_{1,2} \right) = 0$ for all $a,b,c \in \{0,1\}$. Note that the identity $(y_{1,2} - y_{1,3})^2 = y_{2,3}$ holds if y is one of the vertices of CUT_n. (In words, vertices 2 and 3 are on different sides of the bipartition if and only if exactly one of them is on the same side as vertex 1.) We claim that either $\tilde{\mathbb{E}}_y \mathbb{1}_{abc} = 0$ or $(a,b,c) \in \text{CUT}_3 \cap \{0,1\}^3$. This claim implies the desired identity,

$$\begin{split} \tilde{\mathbb{E}}(y_{1,2} - y_{1,3})^2 - y_{2,3} &= \sum_{a,b,c \in \{0,1\}} \tilde{\mathbb{E}} \, \mathbb{1}_{abc} \cdot \left((y_{1,2} - y_{1,3})^2 - y_{2,3} \right) \\ &= \sum_{a,b,c \in \{0,1\}} \tilde{\mathbb{E}} \, \mathbb{1}_{abc} \cdot \left((a - b)^2 - c \right) \\ &= \sum_{(a,b,c) \in \text{CUT}_3 \cap \{0,1\}^3} \tilde{\mathbb{E}} \, \mathbb{1}_{abc} \cdot \left((a - b)^2 - c \right) \quad \text{(by claim)} \\ &= 0 \end{split}$$

(The second step uses that $\mathbb{1}_{abc} \cdot \left((y_{1,2} - y_{1,3})^2 - y_{2,3} \right) = \mathbb{1}_{abc} \cdot \left((a - b)^2 - c \right)$ for all y.) It remains to prove the claim. Since $\text{CUT}_n \cap \{0,1\}^n = \text{METRIC}_n \cap \{0,1\}^n$, it is enough to show that for every $a,b,c \in \{0,1\}$ and every valid linear inequality $\{\ell \geq 0\}$ for METRIC₃ either $\tilde{\mathbb{E}}_y \mathbb{1}_{abc} = 0$ or $\ell(a,b,c) \geq 0$. Indeed, since $\mathbb{1}_{abc}$ is a nonnegative 3-junta and $\{\ell(y_{1,2},y_{1,3},y_{2,3})\}$ is a valid linear inequality for METRIC_n,

$$0 \leq \tilde{\mathbb{E}}_y \mathbb{1}_{abc} \cdot \ell(y_{1,2}, y_{1,3}, y_{2,3}) = \tilde{\mathbb{E}}_y \mathbb{1}_{abc} \cdot \ell(a, b, c),$$

which means that either $\tilde{\mathbb{E}}_y \mathbb{1}_{abc} = 0$ or $\ell(a,b,c) \ge 0$. (The second step uses that $\mathbb{1}_{abc} \cdot \ell(y_{1,2},y_{1,3},y_{2,3}) = \mathbb{1}_{abc} \cdot \ell(a,b,c)$ for all y.)