Goal oriented adaptivity in the IRGNM for parameter identification in PDEs I: reduced formulation

B. Kaltenbacher

A. Kirchner

er S. Veljović

June 3, 2021

Abstract

In this paper we study adaptive discretization of the iteratively regularized Gauss-Newton method IRGNM with an a posteriori (discrepancy principle) choice of the regularization parameter in each Newton step and of the stopping index. We first of all prove convergence and convergence rates under some accuracy requirements formulated in terms of four quantities of interest. Then computation of error estimators for these quantities based on a weighted dual residual method is discussed, which results in an algorithm for adaptive refinement. Finally we extend the results from the Hilbert space setting with quadratic penalty to Banach spaces and general Tikhonov functionals for the regularization of each Newton step.

1 Introduction

Parameter identification problems in partial differential equations (PDEs) can often be written as nonlinear ill-posed operator equations

$$F(q) = g,\tag{1}$$

where F is a nonlinear operator between Hilbert spaces Q and G and where the given data g^{δ} is noisy with the noise level δ :

$$||g - g^{\delta}|| \le \delta. \tag{2}$$

Throughout this paper we will assume that a solution q^{\dagger} to (1) exists.

In case of inverse problems for PDEs, F is the composition of a parameter-to-solution map

$$\begin{array}{rcccc} S \colon Q & \to & V \\ q & \mapsto & u \end{array}$$

with some measurement operator

$$\begin{array}{rccc} C \colon V & \to & G \\ & u & \mapsto & g \,, \end{array}$$

where V is an appropriate Hilbert space. Here, we will write the underlying (possibly nonlinear) PDE in its weak form:

For
$$q \in Q$$
 find $u \in V$: $A(q, u)(v) = (f, v) \quad \forall v \in W$, (3)

where u denotes the PDE solution, q some searched for coefficient or boundary function, and $f \in W^*$ is some given right hand side in the dual of some Hilbert space W. We will assume that the PDE (3) and especially also its linearization at (q, u) is uniquely and stably solvable.

For the stable solution of (1) with noisy data, we consider the iteratively regularized Gauss-Newton method (IRGNM) first of all (section 2) in the reduced form

$$q^{k,\delta} = q^{k-1,\delta} - (F'(q^{k-1,\delta})^*F'(q^{k-1,\delta}) + \alpha_k I)^{-1}(F'(q^{k-1,\delta})^*(F(q^{k-1,\delta}) - g^{\delta}) + \alpha_k(q^{k-1,\delta} - q_0))$$
(4)

or equivalently

$$q^{k,\delta} \in \arg\min_{q}(\mathcal{T}_{\alpha_{k}}(q)) = \arg\min_{q} \|F'(q^{k-1,\delta})(q-q^{k-1,\delta}) + F(q^{k-1,\delta}) - g^{\delta}\|_{G}^{2} + \alpha_{k}\|q-q_{0}\|_{Q}^{2},$$
(5)

(see, e.g., [1, 17] and the references therein). For some all-at-once formulations of the IRGNM we refer to [16] (part II of this paper).

The regularization parameter α_k and the overall stopping index k_* have to be chosen in an appropriate way in order to guarantee convergence. We will here use an inexact Newton / discrepancy principle type strategy, as it has been shown to yield convergence of the IRGNM even in a Banach space setting in [18], see also [13] for a convergence analysis in a still more general setup but with different parameter choice strategies for α_k and k_* .

Our aim is to consider adaptively discretized versions of the formulations (4) defined by replacing the spaces Q, V, W with finite dimensional counterparts Q_h, V_h, W_h (using possibly different discretizations of V, W in (9) and (8)). These should be sufficiently precise so that the convergence results from the continuous setting can be carried over, but save computational effort by using degrees of freedom only where really necessary. For this purpose we will make use of goal oriented error estimators ([2, 3]), that control the error in some quantities of interest I, which are functionals of the variables q, u, w (see (7)-(9) below). We follow the concept proposed in [10], where an inexact Newton method for the computation of a regularization parameter according to the discrepancy principle is combined with adaptive refinement using goal oriented error estimators. While [10] is limited to linear inverse problems, in [15] the idea has been extended to the nonlinear case. Different from [15], we do not treat the nonlinear problem directly here, but use an iterative solution algorithm, the iteratively regularized Gauss-Newton method (4), (5) and treat a sequence of linearized problems instead.

The remainder of this paper is organized as follows. In Section 2 we formulate the Newton step equation as linear quadratic optimal control problem and derive its discretization together with certain quantities of interest, whose precision will be crucial for obtaining convergence results for the overall regularized Newton iteration. This will be substantiated in the convergence and convergence rates results provided in the subsections 2.1 and 2.2. Subsection 2.3 describes computation of the required error estimators by a goal oriented approach and Subsection 2.4 provides the full algorithm. The method and its analysis is extended to a setting with general data misfit and regularization terms in Subsection 2.5. We conclude with a few remarks in Section 3.

2 Reduced form of the discretized IRGNM

We consider the iteration rule (4) for solving the optimization problem

$$\min_{q \in Q} \|F'(q^{\delta,k-1})(q-q^{\delta,k-1}) + F(q^{\delta,k-1}) - g^{\delta}\|_{G}^{2} + \frac{1}{\beta_{k}} \|q-q_{0}\|_{Q}^{2},$$
(6)

where the regularization parameter β_k is updated in each Gauss Newton iteration according to an inexact Newton method guaranteeing a relaxed version of the discrepancy principle (see Step 15 in Algorithm 3, [10, 15]). Note that although the domain $\mathcal{D}(F)$ might be a strict subset of Q, we need not explicitly restrict q to $\mathcal{D}(F)$ in this minimization, since we will assume that $\mathcal{D}(F)$ contains a ball of radius ρ around q_0 and prove that all iterates remain in this ball, cf. (33). So minimizers over $\mathcal{D}(F)$ will automatically be minimizers over Q.

We start with a detailed description of a single iteration step (4) for fixed (discretized) previous iterate $q^{\delta,k-1} = q_{\text{old}} \in Q$ in a continuous and later in the discretized setting actually used in computations, along with the quantities of interest required in error estimation and adaptive refinement.

We formulate the optimization problem (6) as optimal control problem

$$\min_{(q,u_{\text{old}},w)\in Q\times V\times V} \|C'(u_{\text{old}})(w) + C(u_{\text{old}}) - g^{\delta}\|_{G}^{2} + \frac{1}{\beta_{k}}\|q - q_{0}\|_{Q}^{2}$$
(7)

s.t.
$$A(q_{\text{old}}, u_{\text{old}})(v) = f(v)$$
 $\forall v \in W,$ (8)

$$A'_u(q_{\text{old}}, u_{\text{old}})(w, v) = -A'_q(q_{\text{old}}, u_{\text{old}})(q - q_{\text{old}}, v) \qquad \forall v \in W,$$
(9)

since for a solution $q^{\delta,k}$ of (6) $(q^{\delta,k}, S(q_{\text{old}}), S'(q_{\text{old}})(q^{\delta,k} - q_{\text{old}}))$ solves (7)-(9).

In most of this section we omit the superscript δ (denoting dependence on the noisy data) in order to be able to better indicate the difference between continuous and discretized quantities.

We consider the following quantities of interest

$$\tilde{I}_{1}: Q \times V \times V \times \mathbb{R} \to \mathbb{R}, \quad (q, u_{\text{old}}, w, \beta) \quad \mapsto \|C'(u_{\text{old}})(w) + C(u_{\text{old}}) - g^{\delta}\|_{G}^{2} + \frac{1}{\beta}\|q - q_{0}\|_{Q}^{2} \\
\tilde{I}_{2}: V \times V \to \mathbb{R}, \qquad (u_{\text{old}}, w) \qquad \mapsto \|C'(u_{\text{old}})(w) + C(u_{\text{old}}) - g^{\delta}\|_{G}^{2} \\
\tilde{I}_{3}: V \to \mathbb{R}, \qquad u_{\text{old}} \qquad \mapsto \|C(u_{\text{old}}) - g^{\delta}\|_{G}^{2},$$
(10)

i.e. we assume the knowledge about error estimates

$$\begin{split} \eta_1 &\geq \left| \tilde{I}_1(q, u_{\text{old}}, w, \beta) - \tilde{I}_1(q_h, u_{\text{old},h}, w_h, \beta) \right| \\ \eta_2 &\geq \left| \tilde{I}_2(u_{\text{old}}, w) - \tilde{I}_2(u_{\text{old}h}, w_h) \right| \\ \eta_3 &\geq \left| \tilde{I}_3(u_{\text{old}}) - \tilde{I}_3(u_{\text{old}h}) \right|, \end{split}$$

where $q_h, u_{\text{old},h}, w_h$ is a discrete approximate solution to (7), which will be concretised in the following. The error bounds η_1 , η_2 and η_3 will be estimated using goal oriented error estimators cf. Section 2.3.

Additionally we define the functionals

$$I_{1}: Q \times Q \times \mathbb{R} \to \mathbb{R}, \quad (q_{\text{old}}, q, \beta) \quad \mapsto \|F'(q_{\text{old}})(q - q_{\text{old}}) + F(q_{\text{old}}) - g^{\delta}\|_{G}^{2} + \frac{1}{\beta} \|q - q_{0}\|_{Q}^{2}$$

$$I_{2}: Q \times Q \to \mathbb{R}, \qquad (q_{\text{old}}, q) \qquad \mapsto \|F'(q_{\text{old}})(q - q_{\text{old}}) + F(q_{\text{old}}) - g^{\delta}\|_{G}^{2}$$

$$I_{3}: Q \to \mathbb{R}, \qquad q_{\text{old}} \qquad \mapsto \|F(q_{\text{old}}) - g^{\delta}\|_{G}^{2}$$

$$I_{4}: Q \to \mathbb{R}, \qquad q \qquad \mapsto \|F(q) - g^{\delta}\|_{G}^{2}, \qquad (11)$$

which can be seen as reduced versions of (10), since for a solution (q, u_{old}, w) of (7)-(9) and $u \in V$ fulfilling

$$A(q, u)(v) = f(v) \qquad \forall v \in W$$
(12)

there holds

$$\tilde{I}_{1}(q, u_{\text{old}}, w, \beta) = I_{1}(q_{\text{old}}, q, \beta), \qquad \tilde{I}_{2}(u_{\text{old}}, w) = I_{2}(q_{\text{old}}, q),
\tilde{I}_{3}(u_{\text{old}}) = I_{3}(q_{\text{old}}), \qquad \tilde{I}_{3}(u) = I_{4}(q).$$
(13)

The (continuous) quantities of interest in the k-th iteration step are then defined as follows: For a solution $(q^k, u_{\text{old}}^k, w^k)$ of (7) for given $q_{\text{old}} = q_{\text{old}}^k$ and $\beta = \beta_k$ and u^k fulfilling

$$A(q^k, u^k)(v) = f(v) \qquad \forall v \in W$$
(14)

in the k-th iteration let

$$I_{1}^{k} \coloneqq \tilde{I}_{1}(q^{k}, u_{\text{old}}^{k}, w^{k}, \beta_{k}) = I_{1}(q^{k}, q_{\text{old}}^{k}, \beta_{k})$$

$$= \|F'(q_{\text{old}}^{k})(q^{k} - q_{\text{old}}^{k}) + F(q_{\text{old}}^{k}) - g^{\delta}\|_{G}^{2} + \frac{1}{\beta_{k}}\|q^{k} - q_{0}\|_{Q}^{2}$$

$$I_{2}^{k} \coloneqq \tilde{I}_{2}(q_{\text{old}}^{k}, w^{k}) = I_{2}(q_{\text{old}}^{k}, q^{k}) = \|F'(q_{\text{old}}^{k})(q^{k} - q_{\text{old}}^{k}) + F(q_{\text{old}}^{k}) - g^{\delta}\|_{G}^{2}$$

$$I_{3}^{k} \coloneqq \tilde{I}_{3}(u_{\text{old}}^{k}) = I_{3}(q_{\text{old}}^{k}) = \|F(q_{\text{old}}^{k}) - g^{\delta}\|_{G}^{2}$$

$$I_{4}^{k} \coloneqq \tilde{I}_{3}(u^{k}) = I_{4}(q^{k}) = \|F(q^{k}) - g^{\delta}\|_{G}^{2}.$$
(15)

To formulate the quantities of interest (10) for a discrete setting, we consider finite element spaces Q_h, V_h, W_h to Q, V, W, and S_h denotes the discrete solution operator of the state equation. The discretized version of the optimal control problem (7) for given $q_{\text{old}} \in Q_h$ can then be formulated as

$$\min_{(q,u_{\text{old}},w)\in Q_h\times V_h\times V_h} \|C'(u_{\text{old}})(w) + C(u_{\text{old}}) - g^{\delta}\|_G^2 + \frac{1}{\beta}\|q - q_0\|_Q^2$$
(16)

subject to

$$A(q_{\text{old}}, u_{\text{old}})(v) = f(v) \qquad \forall v \in W_h \tag{17}$$

$$A'_u(q_{\text{old}}, u_{\text{old}})(w, v) + A'_q(q_{\text{old}}, u_{\text{old}})(q - q_{\text{old}}, v) = 0 \qquad \forall v \in W_h .$$
(18)

Equation (17) is equivalent to $u_{\text{old}} = S_h(q_{\text{old}})$ and (18) is equivalent to $w = S'_h(q_{\text{old}})(q - q_{\text{old}})$, such that the reduced form of (16) reads

$$\min_{q \in Q_h} \|F'_h(q_{\text{old}})(q - q_{\text{old}}) + F_h(q_{\text{old}}) - g^{\delta}\|_G^2 + \frac{1}{\beta} \|q - q_0\|_Q^2$$
(19)

with $F_h = C \circ S_h$.

Remark 1. One can think of using different discretizations (V_h, W_h) for (17) and $(\tilde{V}_h, \tilde{W}_h)$ for (18) (see Algorithm 1), which we do not indicate here in order to avoid a too complicated setup that would probably not lead to much gain in computational efficiency.

Then the discrete quantities of interest in the reduced form (i.e. the discrete counterparts to (11)) are defined by

$$I_{1,h}: Q \times Q \times \mathbb{R} \to \mathbb{R}, \quad (q_{\text{old}}, q, \beta) \quad \mapsto \|F'_h(q_{\text{old}})(q - q_{\text{old}}) + F_h(q_{\text{old}}) - g^{\delta}\|_G^2 + \frac{1}{\beta} \|q - q_0\|_Q^2$$

$$I_{2,h}: Q \times Q \to \mathbb{R}, \quad (q_{\text{old}}, q) \quad \mapsto \|F'_h(q_{\text{old}})(q - q_{\text{old}}) + F_h(q_{\text{old}}) - g^{\delta}\|_G^2$$

$$I_{3,h}: Q \to \mathbb{R}, \quad q_{\text{old}} \quad \mapsto \|F_h(q_{\text{old}}) - g^{\delta}\|_G^2$$

$$I_{4,h}: Q \to \mathbb{R}, \quad q \quad \mapsto \|F_h(q) - g^{\delta}\|_G^2, \quad (20)$$

such that consistent with (13) for a solution $(q_h, u_{\text{old},h}, w_h)$ of the discretized problem (16)-(18) there holds

$$\tilde{I}_{1}(q_{h}, u_{\text{old},h}, w_{h}, \beta_{h}) = I_{1,h}(q_{\text{old}}, q_{h}, \beta), \qquad \tilde{I}_{2}(u_{\text{old},h}, w_{h}) = I_{2,h}(q_{\text{old}}, q_{h}),
\tilde{I}_{3}(u_{\text{old},h}) = I_{3,h}(q_{\text{old}}), \qquad \tilde{I}_{3}(u_{h}) = I_{4,h}(q_{h}).$$

Correspondingly, the discrete quantities of interest in the k-th iteration step (i.e. the discrete counterparts to (15)) for a solution $(q_h^k, u_{\text{old},h}^k, w_h^k)$ of (16) for given $q_{\text{old}} = q_{\text{old}}^k \in Q_h$ can be formulated as

$$I_{1,h}^{k} \coloneqq I_{1}(q_{h_{k}}^{k}, u_{\text{old},h_{k}}^{k}, w_{h_{k}}^{k}, \beta_{h_{k}}^{k}) = I_{1,h_{k}}^{k}(q_{h_{k}}^{k}, q_{\text{old}}^{k}, \beta_{k}) \\
= \|F_{h_{k}}^{\prime}(q_{\text{old}}^{k})(q_{h_{k}}^{k} - q_{\text{old}}^{k}) + F_{h_{k}}(q_{\text{old}}^{k}) - g^{\delta}\|_{G}^{2} + \frac{1}{\beta_{k}}\|q_{h_{k}}^{k} - q_{0}\|_{Q}^{2} \\
I_{2,h}^{k} \coloneqq \tilde{I}_{2}(q_{\text{old}}^{k}, w_{h_{k}}^{k}) = I_{2,h_{k}}^{k}(q_{\text{old}}^{k}, q_{h_{k}}^{k}) = \|F_{h_{k}}^{\prime}(q_{\text{old}}^{k})(q_{h_{k}}^{k} - q_{\text{old}}^{k}) + F_{h_{k}}(q_{\text{old}}^{k}) - g^{\delta}\|_{G}^{2} \\
I_{3,h}^{k} \coloneqq \tilde{I}_{3}(u_{\text{old},h_{k}}^{k}) = I_{3,h_{k}}^{k}(q_{\text{old}}^{k}) = \|F_{h_{k}}(q_{\text{old}}^{k}) - g^{\delta}\|_{G}^{2} \\
I_{4,h}^{k} \coloneqq \tilde{I}_{3}(u_{h_{k}}^{k}) = I_{4,h_{k}}^{k}(q_{h_{k}}^{k}) = \|F_{h_{k}}(q_{h_{k}}^{k}) - g^{\delta}\|_{G}^{2},$$
(21)

where we introduced the notation h_k (replacing h), denoting the discretization in step k, in order to distiguish between the possibly different discretizations during the iterative process in the following.

Note that the norms in G and in Q (and later on also the one in V) as well as the operator C and the semilinear form $a: Q \times V \times W \to \mathbb{R}$ defined by the relation $a(q, u)(v) = \langle A(q, u), v \rangle_{W^*, W}$ (where $\langle ., . \rangle_{W^*, W}$ denotes the duality pairing between W^* and W) are assumed to be evaluated exactly.

At the end of each iteration step we set

$$q_{\text{old}}^{k+1} \coloneqq q_h^k \,. \tag{22}$$

Remark 2. The sequence of iterates we actually consider is the discrete one $(q_{h_k}^k)_{k \in \mathbb{N}}$, which we also update according to (22). Besides that, for theoretical purposes we keep a sequence of continuous iterates $(q^k)_{k \in \mathbb{N}}$, where each member q^k of this sequence emerges from a member $q_{\text{old}}^k = q_{k-1}^{h_{k-1}}$ of the sequence of discretized iterates $(q_{h_k}^k)_{k \in \mathbb{N}}$, but not from q^{k-1} , see Figure 1. One of the reasons for the necessity of considering this auxiliary continuous iterates is the key inequality (34) in the proof of the convergence theorem below, which makes use of minimality



Figure 1: Sequence of discretized iterates and auxiliary sequence of continuous iterates

of the iterate q_k in all of Q (and not only in the finite dimensional subspace Q_h) thus allowing for comparison to the infinite dimensional exact solution q^{\dagger} .

We stress once more that the discretization may be different in each iteration, as indicated by the superscripts h_k , h_{k-1} here. In order to keep the notation readable we will suppress the iteration index k in the superscript h_k whenever this is possible without causing confusion.

Remark 3. In view of (22) and the last two identities in (13) one might think that $I_{3,h}^{k+1} = \|F(q_{\text{old}}^{k+1}) - g^{\delta}\|_{G}^{2}$ and $I_{4,h}^{k} = \|F(q_{h}^{k}) - g^{\delta}\|_{G}^{2}$ are the same, but this is not the case, since there holds indeed

$$u_h^k = S_{h_k}(q_h^k) = S_{h_k}(q_{\text{old}}^{k+1})$$

for $h = h_k$, i.e. with respect to the discretization from step k, but

$$u_{\text{old},h}^{k+1} = S_{h_{k+1}}(q_{\text{old}}^{k+1})$$

for $h = h_{k+1}$, i.e. with respect to the discretization from step k + 1. Due to the possibly different discretizations, in general there holds

$$u_h^k \neq u_{\text{old},h}^{k+1}$$
.

Also $I_3^{k+1} = \|F(q_{\text{old}}^{k+1}) - g^{\delta}\|_G^2$ and $I_4^k = \|F(q^k) - g^{\delta}\|_G^2$ are not the same, because

$$q_{\text{old}}^{k+1} \coloneqq q_h^k \neq q^k \,,$$

see Figure 1.

Remark 4. Note that even the discretizations h_k for fixed k can differ in the different quantities of interest during one Gauss Newton iteration cf. Algorithm 1. Tracking the proof of the main convergence result Theorem 1 the reader can verify that only $I_{1,h}^k$ and $I_{2,h}^k$ have to be evaluated on the same mesh, since in the proof we will need the identity

$$I_{1,h}^{k} = I_{2,h}^{k} + \frac{1}{\beta_{k}} \|q_{h_{k}}^{k} - q_{0}\|^{2}, \qquad (23)$$

which is guaranteed by assuming exact evaluation of the Q-norm $\|q_h^k - q_0\|_Q$.

In order to assess and – by adaptive refinement – to control the differences

$$|I_{i,h}^k - I_i^k| \le \eta_i^k, \quad i \in \{1, 2, 3, 4\}$$
(24)

between the exact quantities of interest and their counterparts resulting from discretization, we will make use of goal oriented error estimators, which will be explained in more detail in Section 2.3.

We select β_k according to an inexact Newton condition (cf. [11, 20]) which can be interpreted as a discrepancy principle with "noise level" $\tilde{\theta}I_{3,h}^k$

$$\tilde{\underline{\theta}}I_{3,h}^k \le I_{2,h}^k \le \overline{\theta}I_{3,h}^k \,, \tag{25}$$

i.e.,

$$\frac{\tilde{\theta}}{\tilde{\theta}} \|F_h(q_{\text{old}}^k) - g^{\delta}\|_G^2 \le \|F_h'(q_{\text{old}}^k)(q_h^k - q_{\text{old}}^k) + F_h(q_{\text{old}}^k) - g^{\delta}\|_G^2 \le \tilde{\tilde{\theta}} \|F_h(q_{\text{old}}^k) - g^{\delta}\|_G^2,$$

for some $0 < \tilde{\underline{\theta}} \leq \tilde{\overline{\theta}} < \frac{1}{2}$. Note that this regularization parameter can be computed in an efficient manner according to [10], see Theorem 1 there. We mention in passing that the latter would as well allow us to use the continuous version I_2^k in (25), but we prefer to formulate the condition with the discretized actually computed quantities anyway.

The overall Newton iteration is stopped according to a generalized discrepancy principle

$$k_* = \min\{k \in \mathbb{N} : I_{3,h}^k \le \tau^2 \delta^2\}.$$
(26)

In our convergence analysis we will use the following weak sequential closedness assumption on F:

$$(q_n \rightharpoonup q \land F(q_n) \to g) \Rightarrow (q \in \mathcal{D}(F) \land F(q) = g)$$
(27)

for all $\{q_n\}_{n\in\mathbb{N}}\subseteq Q$ together with the tangential cone condition (also often called Scherzer condition)

$$\|F(q) - F(\bar{q}) - F'(q)(q - \bar{q})\|_G \le c_{tc} \|F(q) - F(\bar{q})\|_G \qquad \forall q, \bar{q} \in \mathcal{B}_\rho(q_0) \subseteq \mathcal{D}(F) \subseteq Q \quad (28)$$

for some $\rho > 0$, $0 < c_{tc} < 1$, which are both typical conditions in the analysis of regularization methods for nonlinear ill-posed problems cf., e.g., [9, 17] and the references therein.

2.1 Convergence

Theorem 1. Let F satisfy the weak sequential closedness condition (27) and the tangential cone condition (28) with $c_{tc} < \frac{1}{4}$ sufficiently small. Let, further, $\tau > 0$ be chosen sufficiently large and $0 < \frac{\tilde{\theta}}{\tilde{\theta}} < \tilde{\overline{\theta}}$ sufficiently small, such that

$$2\left(c_{tc}^{2} + \frac{(1+c_{tc})^{2}}{\tau^{2}}\right) < \underline{\tilde{\theta}} \quad and \quad \frac{2\overline{\tilde{\theta}} + 4c_{tc}^{2}}{1 - 4c_{tc}^{2}} < 1.$$

$$\tag{29}$$

Finally, let for the discretization error with respect to the quantities of interest (24) hold, where η_1^k , η_2^k , η_3^k , η_4^k are selected such that

$$\eta_1^k + 2c_{tc}^2 \eta_3^k \le \left(\frac{\tilde{\theta}}{\ell} - 2\left(c_{tc}^2 + \frac{(1+c_{tc})^2}{\tau^2}\right)\right) I_{3,h}^k \tag{30}$$

$$I_{3,h}^{k} \le (1+c_3)I_{4,h}^{k-1} + r^{k} \quad and \quad (1+c_3)\frac{2\overline{\theta} + 4c_{tc}^2}{1-4c_{tc}^2} \le c_2 < 1 \tag{32}$$

for some constants $c_1, c_2, c_3 > 0$, and a sequence $r^k \to 0$ as $k \to \infty$ (where the second condition in (32) is possible due to the right inequality in (29)).

Then with β_k and $h = h_k$ fulfilling (25) and k_* selected according to (26) there holds

(i) For any solution $q^{\dagger} \in \mathcal{B}_{\rho}(q_0)$ of (1)

$$\|q_h^k - q_0\|_Q^2 \le \|q^{\dagger} - q_0\|_Q^2 \quad \forall k < k_* ,$$
(33)

- (ii) k_* is finite,
- (iii) $q_{\text{old}}^{k_*} = q_{h_{k_*}-1}^{k_*-1}$ converges (weakly) subsequentially to a solution of (1) as $\delta \to 0$ in the sense that it has a weakly convergent subsequence and each weakly convergent subsequence converges strongly to a solution of (1). If the solution q^{\dagger} to (1) is unique, then $q_{\text{old}}^{k_*}$ converges strongly to q^{\dagger} as $\delta \to 0$.
- *Proof.* (i): For k = 0, (33) trivially holds. For all $1 \le k < k_*$ and any solution q^{\dagger} of (1) we have by (24) and minimality of q^k

$$I_{1,h}^{k} \leq I_{1}^{k} + \eta_{1}^{k} = \|F'(q_{\text{old}}^{k})(q^{\dagger} - q_{\text{old}}^{k}) + F(q_{\text{old}}^{k}) - g^{\delta}\|_{G}^{2} + \frac{1}{\beta_{k}}\|q^{\dagger} - q_{0}\|_{Q}^{2} + \eta_{1}^{k}.$$
 (34)

In here, according to (2), (26) and (28), as well as the inequality $(a + b)^2 \leq 2a^2 + 2b^2$ for arbitrary $a, b \in \mathbb{R}$ we can estimate as follows

$$\begin{aligned} \|F'(q_{\text{old}}^{k})(q^{\dagger} - q_{\text{old}}^{k}) + F(q_{\text{old}}^{k}) - g^{\delta}\|_{G}^{2} \\ &\leq \left(\|F'(q_{\text{old}}^{k})(q^{\dagger} - q_{\text{old}}^{k}) + F(q_{\text{old}}^{k}) - F(q^{\dagger})\|_{G} + \delta\right)^{2} \\ &\leq \left(c_{tc}\|F(q^{\dagger}) - F(q_{\text{old}}^{k})\|_{G} + \delta\right)^{2} \\ &\leq \left(c_{tc}\left(\|g^{\delta} - F(q_{\text{old}}^{k})\|_{G} + \delta\right) + \delta\right)^{2} \\ &= \left(c_{tc}\sqrt{I_{3}^{k}} + (1 + c_{tc})\delta\right)^{2} \\ &\leq 2c_{tc}^{2}I_{3}^{k} + 2(1 + c_{tc})^{2}\delta^{2} \\ &\leq 2c_{tc}^{2}(I_{3,h}^{k} + \eta_{3}^{k}) + 2(1 + c_{tc})^{2}\frac{I_{3,h}^{k}}{\tau^{2}} \\ &= 2\left(c_{tc}^{2} + \frac{(1 + c_{tc})^{2}}{\tau^{2}}\right)I_{3,h}^{k} + 2c_{tc}^{2}\eta_{3}^{k}. \end{aligned}$$
(35)

On the other hand, from (23), (25) it follows that

$$I_{1,h}^{k} = I_{2,h}^{k} + \frac{1}{\beta_{k}} \|q_{h}^{k} - q_{0}\|_{Q}^{2} \ge \underline{\tilde{\theta}} I_{3,h}^{k} + \frac{1}{\beta_{k}} \|q_{h}^{k} - q_{0}\|_{Q}^{2}$$
(36)

which together with the previous inequality and (30) gives

$$\begin{split} \underline{\tilde{\theta}} I_{3,h}^k + \frac{1}{\beta_k} \| q_h^k - q_0 \|_Q^2 &\leq I_1^k + \eta_1^k \\ &= \| F'(q_{\text{old}}^k)(q^{\dagger} - q_{\text{old}}^k) + F(q_{\text{old}}^k) - g^{\delta} \|_G^2 + \frac{1}{\beta_k} \| q^{\dagger} - q_0 \|_Q^2 + \eta_1^k \\ &\leq 2 \left(c_{tc}^2 + \frac{(1 + c_{tc})^2}{\tau^2} \right) I_{3,h}^k + \frac{1}{\beta_k} \| q^{\dagger} - q_0 \|_Q^2 + \eta_1^k + 2c_{tc}^2 \eta_3^k \\ &\leq \underline{\tilde{\theta}} I_{3,h}^k + \frac{1}{\beta_k} \| q^{\dagger} - q_0 \|_Q^2 \,, \end{split}$$

which implies (33).

(ii): Furthermore, for all $1 \le k < k_*$ we have by the triangle inequality as well as (28) and (25)

$$\begin{aligned}
\sqrt{I_4^k} &= \|F(q^k) - g^{\delta}\|_G \\
&\leq \|F'(q_{\text{old}}^k)(q^k - q_{\text{old}}^k) + F(q_{\text{old}}^k) - g^{\delta}\|_G + \|F'(q_{\text{old}}^k)(q^k - q_{\text{old}}^k) - F(q^k) + F(q_{\text{old}}^k)\|_G \\
&\leq \sqrt{I_2^k} + c_{tc}\|F(q^k) - F(q_{\text{old}}^k)\|_G \\
&\leq \sqrt{\tilde{\theta}}I_{3,h}^k + \eta_2^k + c_{tc}(\sqrt{I_4^k} + \sqrt{I_3^k}),
\end{aligned}$$
(37)

hence by $(a+b)^2 \leq 2a^2 + 2b^2$ for all $a, b \in \mathbb{R}$

$$I_4^k \le 2(\overline{\theta}I_{3,h}^k + \eta_2^k) + 2c_{tc}^2(2I_4^k + 2I_3^k),$$

which implies

$$I_4^k \le \frac{1}{1 - 4c_{tc}^2} \left(2\tilde{\overline{\theta}} I_{3,h}^k + 2\eta_2^k + 4c_{tc}^2 I_3^k \right)$$

With (24) and (32) we can further deduce

$$\begin{split} I_{4,h}^{k} &\leq \frac{1}{1 - 4c_{tc}^{2}} \left((2\tilde{\overline{\theta}} + 4c_{tc}^{2})I_{3,h}^{k} + 2\eta_{2}^{k} + 4c_{tc}^{2}\eta_{3}^{k} \right) + \eta_{4}^{k} \\ &\leq \frac{2\tilde{\overline{\theta}} + 4c_{tc}^{2}}{1 - 4c_{tc}^{2}} (1 + c_{3})I_{4,h}^{k-1} + \frac{1}{1 - 4c_{tc}^{2}} \left((2\tilde{\overline{\theta}} + 4c_{tc}^{2})r^{k} + 2\eta_{2}^{k} + 4c_{tc}^{2}\eta_{3}^{k} \right) + \eta_{4}^{k} \\ &\leq c_{2}I_{4,h}^{k-1} + \frac{1}{1 - 4c_{tc}^{2}} \left((2\tilde{\overline{\theta}} + 4c_{tc}^{2})r^{k} + 2\eta_{2}^{k} + 4c_{tc}^{2}\eta_{3}^{k} \right) + \eta_{4}^{k} \,. \end{split}$$

With the notation

$$a^{i} \coloneqq \frac{1}{1 - 4c_{tc}^{2}} \left((2\tilde{\bar{\theta}} + 4c_{tc}^{2})r^{i} + 2\eta_{2}^{i} + 4c_{tc}^{2}\eta_{3}^{i} \right) + \eta_{4}^{i} \quad \forall i \in \{1, 2, \dots, k\}$$
(38)

there follows recursively

$$I_{4,h}^{k} \le c_{2}^{k} I_{4,h}^{0} + \sum_{j=0}^{k-1} c_{2}^{j} a^{k-j} .$$
(39)

Note that by the second part of (31), the second part of (32) and the fact that $r^k \to 0$ as $k \to \infty$ (by definition of r^k), we have $c_2^k I_{4,h}^0 + \sum_{j=0}^{k-1} c_j^j a^{k-j} \to 0$ as $k \to \infty$. So, if the discrepancy principle never got active (i.e., $k_* = \infty$), the sequence $(I_{4,h}^k)_{k \in \mathbb{N}}$ and therewith by assumption (31) also $(I_{3,h}^k)_{k \in \mathbb{N}}$ would be bounded by a sequence tending to zero as $k \to \infty$, which implies that $I_{3,h}^k$ would fall below $\tau^2 \delta^2$ for k sufficiently large, thus yielding a contradiction. Hence the stopping index $k_* < \infty$ is well-defined and finite.

(iii): With (2), (24), (31) and definition of k_* , we have

$$\|F(q_{\text{old}}^{k_*}) - g\|_G \le \sqrt{I_3^{k_*}} + \delta \le \sqrt{I_{3,h}^{k_*} + \eta_3^{k_*}} + \delta \le \sqrt{(1+c_1)I_{3,h}^{k_*}} + \delta \le (\sqrt{1+c_1}\tau + 1)\delta \to 0$$
(40)

as $\delta \to 0$. Thus, due to (ii) (33) $q_{\text{old}}^{k_*} = q_{h_{k_*-1}}^{k_*-1}$ has a weakly convergent subsequence $(q_{\text{old}}^{k_*(\delta_l)})_{l \in \mathbb{N}}$ and due to the weak sequential closedness of F and (40) the limit $q^* \in \mathcal{B}_{\rho}(q_0)$ of every weakly convergent subsequence is a solution to F(q) = g.

Strong convergence of $(q_{\text{old}}^{k_*(\delta_l)})_{l\in\mathbb{N}}$ to q^* follows from the standard argument

$$\|q_{\text{old}}^{k_{*}(\delta_{l})} - q^{*}\|_{Q}^{2} = \underbrace{\|q_{\text{old}}^{k_{*}(\delta_{l})} - q_{0}\|_{Q}^{2}}_{\leq \|q^{*} - q_{0}\|_{Q}^{2}} + \|q^{*} - q_{0}\|_{Q}^{2} - 2 \underbrace{\langle q_{\text{old}}^{k_{*}(\delta_{l})} - q_{0}, q^{*} - q_{0}\rangle_{Q}}_{\rightarrow \|q^{*} - q_{0}\|_{Q}^{2}} \text{by weak convergence}$$

(with $\langle ., . \rangle_Q$ denoting the scalar product in Q), where we have used the fact that in (33) we can replace q^{\dagger} by q^* since the latter solves (1).

Remark 5. Note that estimate (34) can alternatively be obtained by using stationarity instead of minimality of q^k (which is equivalent by convexity): For all $dq \in Q$

$$0 = \langle F'(q_{\text{old}}^k)(q^k - q_{\text{old}}^k) + F(q_{\text{old}}^k) - g^{\delta}, F'(q_{\text{old}}^k)(dq) \rangle_G + \frac{1}{\beta_k} \langle q^k - q_0, dq \rangle_Q$$

(with $\langle ., . \rangle_G$ denoting the scalar product in G). With $dq = q^k - q^{\dagger}$ this yields

$$\begin{aligned} 0 &= \|F'(q_{\text{old}}^k)(q^k - q_{\text{old}}^k) + F(q_{\text{old}}^k) - g^{\delta}\|_G^2 \\ &- \langle F'(q_{\text{old}}^k)(q^k - q_{\text{old}}^k) + F(q_{\text{old}}^k) - g^{\delta}, F'(q_{\text{old}}^k)(q^{\dagger} - q_{\text{old}}^k) + F(q_{\text{old}}^k) - g^{\delta} \rangle_G \\ &+ \frac{1}{\beta_k} \|q^k - q_0\|_Q^2 - \frac{1}{\beta_k} \langle q^k - q_0, q^{\dagger} - q_0 \rangle_Q \,, \end{aligned}$$

hence by Cauchy-Schwarz and $ab \leq \frac{1}{2}a^2 + \frac{1}{2}b^2 \ \forall a, b \in \mathbb{R}$

$$\begin{aligned} \|F'(q_{\text{old}}^k)(q^k - q_{\text{old}}^k) + F(q_{\text{old}}^k) - g^{\delta}\|_G^2 + \frac{1}{\beta_k} \|q^k - q_0\|_Q^2 \\ &\leq \|F'(q_{\text{old}}^k)(q^{\dagger} - q_{\text{old}}^k) + F(q_{\text{old}}^k) - g^{\delta}\|_G^2 + \frac{1}{\beta_k} \|q^{\dagger} - q_0\|_Q^2 \,. \end{aligned}$$

2.2 Convergence rates

To prove convergence rates we will consider Hilbert space source conditions

$$\exists s \in Q \text{ s.t. } q^{\dagger} - q_0 = f(F'(q^{\dagger})^* F'(q^{\dagger}))s.$$
(41)

with $f \colon [0,\infty) \to [0,\infty)$ satisfying

$$f(0) = 0, \quad f^2 \text{ strictly monotonically increasing,} \phi \text{ convex, where } \phi := (f^2)^{-1} \text{ i.e., } \phi^{-1}(\lambda) = f^2(\lambda) \text{ and}$$
(42)
$$\Theta : \lambda \mapsto f(\lambda)\sqrt{\lambda} \text{ strictly monotonically increasing.}$$

Examples of functions f satisfying (42) are Hölder type $f(\lambda) = \lambda^{\nu}, \nu > 0$ or logarithmic type $f(\lambda) = \ln(\frac{1}{\lambda})^{-p}, \lambda \in (0, 1/e], p > 0$ functions.

If (41) holds, then we have by Jensen's inequality

$$\begin{aligned} |\langle q^{\dagger} - q_{0}, q - q^{\dagger} \rangle_{Q}| &= |\langle s, f(F'(q^{\dagger})^{*}F'(q^{\dagger}))(q - q^{\dagger}) \rangle_{Q}| \\ &\leq \|s\|_{Q} \|q - q^{\dagger}\|_{Q} f\left(\frac{\|F'(q^{\dagger})(q - q^{\dagger})\|_{G}^{2}}{\|q - q^{\dagger}\|_{Q}^{2}}\right). \end{aligned}$$
(43)

Using estimate (33) from the proof of Theorem 1, as well as the definition of the stopping index according to the discrepancy principle, we can therefore make use of Theorem 1 in [15] to obtain

Theorem 2. Let the conditions of Theorem 1 and additionally the source condition (41) for some function f with (42) be fullfiled.

Then there exists a $\overline{\delta} > 0$ and a constant $\overline{C} > 0$ independent of δ such that for all $\delta \in (0, \overline{\delta}]$

$$\|q_h^{k_*} - q^{\dagger}\|_Q^2 \le \frac{\bar{C}^2 \delta^2}{\Theta^{-1}\left(\frac{\bar{C}}{2\|s\|_Q}\delta\right)} = 4\|s\|^2 f^2(\Theta^{-1}(\frac{\bar{C}}{2\|s\|_Q}\delta))$$
(44)

where $\Theta(\lambda) \coloneqq f(\lambda) \sqrt{\lambda}$.

Proof. The assertion follows from Theorem 1 in [15] using the estimate (33) and

$$||F(q_h^{k_*}) - g^{\delta}||_G \leq \sqrt{I_{3,h}^{k_*} + \eta_3^{k_*}} \leq \sqrt{1 + c_1} \sqrt{I_{3,h}^{k_*}} \leq \sqrt{1 + c_1} \tau \delta.$$

Remark 6. If f satisfies the condition

$$t \mapsto \frac{f(t)}{\sqrt{t}}$$
 monotonically decreasing, (45)

then for all C > 0 the inequality

$$f(\Theta^{-1}(Ct)) \le \max\{\sqrt{C}, 1\} f(\Theta^{-1}(t)) \quad (t \ge 0)$$
 (46)

holds, which implies that we can conclude from (44) the optimal rates

$$\|q_h^{k_*} - q^{\dagger}\|_Q \le C\left(f^2(\Theta^{-1}(\delta))\right) = C\left(\frac{\delta^2}{\Theta^{-1}(\delta)}\right).$$
(47)

The restriction (45) corresponds to the typical saturation phenomenon of Tikhonov regularization in combination with the discrepancy principle, see e.g., [9].

2.3 Computation of the error estimators

The computation of the error estimators η_1^k , η_2^k , η_3^k and η_4^k is done similarly to [10]. The only difference lies in the fact that in I_1^k we have three variables subject to discretization, namely q, u_{old} and w instead of only two (q and u) as usual, which leads to the following error estimators. In this section we omit the iteration index k for simplicity.

2.3.1 Error estimator for I_1

Since the dependence on β is not important for error estimation, we neglect β as argument and consider

$$I_1(q, u_{\text{old}}, w) = \|C'(u_{\text{old}})(w) + C(u_{\text{old}}) - g^{\delta}\|_G^2 + \frac{1}{\beta} \|q - q_0\|_Q^2$$

and define the Lagrange functional

$$\begin{split} L(q, u_{\text{old}}, w, v, v_{\text{old}}) &\coloneqq I_1(q, u_{\text{old}}, w) \\ &\quad + A'_u(q_{\text{old}}, u_{\text{old}})(w)(v) + A'_q(q_{\text{old}}, u_{\text{old}})(q - q_{\text{old}})(v) \\ &\quad + A(q_{\text{old}}, u_{\text{old}})(v_{\text{old}}) - f(v_{\text{old}}) \,. \end{split}$$

Proposition 1. Let $X = Q \times V \times V \times W \times W$ and $X_h = Q_h \times V_h \times V_h \times W_h$. Let $x = (q, u_{old}, w, v, v_{old}) \in X$ be a stationary point of L, i.e.

$$x_h \in X_h$$
: $L'(x)(dx) = 0 \quad \forall dx \in X$

and let $x_h = (q_h, u_{\text{old},h}, w_h, v_h, v_{\text{old},h}) \in X_h$ be a discrete stationary point of L, i.e.

$$L'(x_h)(dx) = 0 \qquad \forall dx \in X_h .$$
(48)

Then there holds

$$I_1(q, u_{\text{old}}, w) - I_1(q_h, u_{\text{old}, h}, w_h) = \frac{1}{2}L'(x_h)(x - \tilde{x}_h) + R,$$

for an arbitrary $\tilde{x}_h \in X_h$ and

$$R = \frac{1}{2} \int_0^1 L'''(x_s e_x)(e_x, e_x, e_x)s(s-1) \ ds$$

with $e_x \coloneqq x - x_h$.

Proof. cf. [10] and [2].

Explicitly such stationary points can be computed by solving the equations

$$u_{\text{old}} \in V: \qquad A(q_{\text{old}}, u_{\text{old}})(dv_{\text{old}}) = f(dv_{\text{old}}) \quad \forall dv_{\text{old}} \in W$$

$$\tag{49}$$

$$w \in V: \qquad A'_u(q_{\text{old}}, u_{\text{old}})(w)(dv) = -A'_q(q_{\text{old}}, u_{\text{old}})(q - q_{\text{old}})(dv) \quad \forall dv \in W$$
(50)

$$v \in W: \qquad A'_u(q_{\text{old}}, u_{\text{old}})(dw)(v) = -I'_{1,w}(q, u_{\text{old}}, w)(dw) \quad \forall dw \in V$$

$$\tag{51}$$

$$v_{\rm old} \in W: \qquad A'_u(q_{\rm old}, u_{\rm old})(du)(v_{\rm old}) = -I'_{1, u_{\rm old}}(q, u_{\rm old}, w)(du) \tag{52}$$

$$-A_{uu}^{\prime\prime}(q_{\text{old}}, u_{\text{old}})(w, du)(v)$$
(53)

$$-A_{qu}^{\prime\prime}(q_{\text{old}}, u_{\text{old}})(q - q_{\text{old}}, du)(v) \quad \forall du \in V \quad (54)$$

$$q \in Q: \qquad I'_{1,q}(q, u_{\text{old}}, w)(dq) = -A'_q(q_{\text{old}}, u_{\text{old}})(dq)(v) \quad \forall dq \in Q.$$
(55)

and their discrete counterparts.

Obviously, we do not actually compute continuous stationary points, but (as in [10]) we choose $\tilde{x}_h = i_h x$ with a suitable interpolation operator $i_h \colon X \to X_h$ and approximate the interpolation error using an operator $\pi_h \colon X_h \to \tilde{X}_h$ with $\tilde{X}_h \neq X_h$, such that $x - \pi_h x_h$ has a better local asymptotical behavior than $x - i_h x$. Then the error estimator η_1 for I_1 can be computed as

$$I_1 - I_{1,h} = I_1(q, u_{\text{old}}, w) - I_1(q_h, u_{\text{old},h}, w_h) \approx \frac{1}{2}L'(x_h)(\pi_h x_h - x_h) = \eta_1$$

(cf. [2]).

Remark 7. Please note that the equations (48) /(49)-(55) are solved anyway in the process of solving the optimization problem (7)-(9).

2.3.2 Error estimator for I_2

We consider

$$I_2(u_{\text{old}}, w) = \|C'(u_{\text{old}})(w) + C(u_{\text{old}}) - g^{\delta}\|_{G}^2$$

and for $x_1 \coloneqq (q_1, u_{\text{old},1}, w_1, v_1, v_{\text{old}}^1) \in X$ we define the Lagrange functional

$$M(x, x_1) \coloneqq I_2(u_{\text{old}}, w) + L'(x)(x_1).$$

Then there holds a similar result to Proposition 1 for the difference $I(u_{\text{old}}, w) - I(u_{\text{old},h}, w_h)$ for stationary points $y = (x, x_1) \in X \times X$ and $y_h = (x_h, x_h^1) \in X_h \times X_h$ of M (cf. [3]). Such a discrete stationary point y_h can be computed by solving the equations (48)/(49)-(55) and

$$x_{h}^{1} \in X_{h}: \qquad L''(x_{h})(x_{h}^{1}, dx) = -I'_{2,w}(u_{\text{old},h}, w_{h})(dw) - I'_{2,u_{\text{old}}}(u_{\text{old},h}, w_{h})(du_{\text{old}}) \qquad \forall dx \in X_{h}$$
(56)

(where $dx = (dq, du_{old}, dw, dv, dv_{old})$). The error estimator η_2 for I_2 can then be computed by

$$I_2 - I_{2,h} = I_2(u_{\text{old}}, w) - I_2(u_{\text{old},h}, w_h) \approx \frac{1}{2}M'(y_h)(\pi_h y_h - y_h) = \eta_2.$$

Remark 8. To avoid the computation of second order information in (56) we would like to refer to [3], where (56) is replaced by an approximate equation of first order.

2.3.3 Error estimator for I_3

For I_3 we again proceed similarly to the sections 2.3.1 and 2.3.2, i.e. we consider

$$I_3(u_{\text{old}}) = ||C(u_{\text{old}}) - g^{\delta}||_G^2$$

and define the Lagrangian

$$N(x, x_2) \coloneqq I_3(u_{\text{old}}) + L'(x)(x_2).$$

As there holds again a similar results to Proposition 1, we compute a discrete stationary point $\chi_h = (x_h, x_h^2) \in X_h \times X_h$ of N by solving the equations (48)/(49)-(55) and

$$x_h^2 \in X_h$$
: $L''(x_h)(x_h^2, dx) = -I'_3(u_{\text{old},h})(du_{\text{old}}) \quad \forall dx \in X_h$, (57)

and compute the error estimator for I_3 as

$$I_3 - I_{3,h} = I_3(u_{\text{old}}) - I_3(u_{\text{old},h}) \approx \frac{1}{2} N'(\chi_h)(\pi_h \chi_h - \chi_h) = \eta_3.$$

2.3.4 Error estimator for I_4

Different to the other error estimates, the bound on the error in I_4 only appears in connection with the very weak assumption $\eta_4^k \to 0$ as $k \to \infty$, which may be satisfied in practice without refining explicitly with respect to η_4 , but simply, by refining with respect to the other error estimators η_1, η_2 , and especially η_3 . Another way to make sure that $\eta_4^k \to 0$ as $k \to \infty$, is, of course, to refine globally every now and then, although this is admittedly, not a very efficient solution.

If one doesn't want to rely on such practically motivated speculations and actually wants to compute an error estimator for I_4 , one has to include the decoupled constraint

$$A(q,u)(v) = f(v) \qquad \forall v \in W$$

in the definition of the Lagrangian L in subsection 2.3.1. In that case we redefine the Lagrange functional L in subsection 2.3.1 as

$$\begin{split} L(q, u_{\text{old}}, w, v, v_{\text{old}}, u, z) &\coloneqq I_1(q, u_{\text{old}}, w) \\ &+ A'_u(q_{\text{old}}, u_{\text{old}})(w)(v) + A'_q(q_{\text{old}}, u_{\text{old}})(q - q_{\text{old}})(v) \\ &+ A(q_{\text{old}}, u_{\text{old}})(v_{\text{old}}) - f(v_{\text{old}}) \\ &+ A(q, u)(z) - f(z) \,. \end{split}$$

and the spaces $X := Q \times V \times V \times W \times W \times V \times W$ and $X_h := Q_h \times V_h \times V_h \times W_h \times W_h \times V_h \times W_h$. Then we consider

$$I_4(u) \coloneqq \|C(u) - g^{\delta}\|_G^2$$

and define the auxiliary Lagrange functional

$$K(x, x_3) \coloneqq I_4(u) + L'(x)(x_3)$$

for $x, x_3 \in X$. Then again (as in the subsections 2.3.1, 2.3.2 and 2.3.3) we could estimate the difference $I_4(u) - I_4(u_h)$ by computing a discrete stationary point $\xi_h = (x_h, x_h^3)$ of K, that means we would solve he equations (48)/(49)-(55) and

$$x_h^3 \in X_h$$
: $L''(x_h)(x_h^3, dx) = -I'_4(u_h)(du) \forall dx = (dq, du, dz) \in X_h$,

with $X_h := Q_h \times V_h \times V_h \times W_h \times W_h \times V_h \times W_h$ and compute the error estimator η_4 for I_4 by

$$I_4 - I_{4,h} = I_4(u) - I_4(u_h) \approx \frac{1}{2} K'(\xi_h)(\pi_h \xi_h - \xi_h) = \eta_4.$$

Algorithm 2.4

As mentioned and justified in Subsection 2.3.4, we neglect η_4^k and the condition $\eta_2^k \to 0$ and $\eta_3^k \to 0$ as $k \to \infty$ from (31) in the following algorithm.

In order to verify the condition (30) more easily, we split (30) into

$$\eta_1^k \le c_4 I_{3,h}^k \quad \text{with} \quad c_4 = \frac{1}{2} \left(\frac{\tilde{\theta}}{\ell} - 2 \left(c_{tc}^2 + \frac{(1+c_{tc})^2}{\tau^2} \right) \right)$$
(58)

and

$$\eta_3^k \le c_5 I_{3,h}^k \quad \text{with} \quad c_5 = \frac{1}{4c_{tc}^2} \left(\frac{\tilde{\theta}}{\ell} - 2\left(c_{tc}^2 + \frac{(1+c_{tc})^2}{\tau^2} \right) \right) \,.$$
 (59)

Additionally we combine the inequality in (31), the first inequality in (32) and (59), since there holds

$$I_{3,h}^k \le I_3^k + \eta_3^k$$
 and $I_{4,h}^{k-1} \ge I_4^{k-1} - \eta_4^{k-1}$,

such that the condition

$$\eta_3^k + (1+c_3)\eta_4^{k-1} \le (1+c_3)I_4^{k-1} - I_3^k + r^k \tag{60}$$

implies the first inequality in (32). As mentioned in Remark 3, I_3^k and I_4^{k-1} and $I_{3,h}^k$ and $I_{4,h}^{k-1}$ only differ in the discretization level, which motivates the assumption that for small h, we have $I_3^k \approx I_4^{k-1}$ and $\eta_4^{k-1} \approx \eta_3^k$, such that instead of (60) we check whether

$$\eta_3^k \le \frac{c_3}{2(1+c_3)} I_{3,h}^k + \frac{r^k}{2(1+c_3)} \,. \tag{61}$$

Thus, as a combination of the inequality in (31), (61) and (59), we formulate

$$\eta_3^k \le \min\left\{c_1, c_5, \frac{c_3}{2(1+c_3)}\right\} I_{3,h}^k.$$
(62)

Algorithm 1. Reduced form of discretized IRGNM

- 1: Choose τ , τ_{β} , $\tilde{\tau}_{\beta}$, $\tilde{\underline{\theta}}$, $\tilde{\overline{\theta}}$ such that $0 < \tilde{\underline{\theta}} \leq \tilde{\overline{\theta}} < 1$ and (29) holds, $\tilde{\theta} = (\tilde{\underline{\theta}} + \tilde{\overline{\theta}})/2$ and $\max\{1, \tilde{\tau}_{\beta}\} < \tau_{\beta} \leq \tau$ and choose the constants c_1 , c_2 and c_3 , such that the second part of (32) is fulfilled.
- 2: Choose a discretization h = h₀ and starting value q_h⁰ = q_{h0}⁰ (not necessarily coinciding with q₀ in the regularization term) and set q_{0ld}⁰ = q_{h0}⁰.
 3: Determine u_{old}⁰ = u_{old,h0}⁰, I_{3,h}⁰ = I_{3,h0}⁰ and q₃⁰ = q_{3,h0}⁰ by applying Algorithm 2 with m = 0
- (and $h = h_0$).
- 4: Set $h_0^1 = h_0$.
- 5: while (62) is violated do

- Refine grids according to the error estimator η_3^0 , such that we obtain a finer discretiza-6: tion h_0^1 .
- Determine $u_{\text{old}}^0 = u_{\text{old},h_0^1}^0$, $I_{3,h}^0 = I_{3,h_0^1}^0$ and $\eta_3^0 = \eta_{3,h_0^1}^0$ by applying Algorithm 2 with $\tilde{7}$: $h = h_0^1$ and m = 0.
- 8: Set k = 0 and $h = h_0^1$ (possibly different from h_0). 9: while $I_{3,h}^k \ge \tau^2 \delta^2$ do 10: Set $h = h_k^1$.
- 10:
- With q_{old}^k , u_{old}^k fixed, apply Algorithm 3 starting with the current mesh $h(=h_k^1)$ to obtain 11: a regularization parameter β_k and a possibly different discretization h_k^2 such that (25) holds and the corresponding $w_h^k = w_{h^2}^k$, $q_h^k = q_{h^2}^k$.
- 12: Set $h = h_k^2$.
- Evaluate error estimator $\eta_1^k = \eta_1^k(h_k^2)$. 13:
- Set $h_k^3 = h_k^2$. 14:
- while (58) is violated do 15:
- Refine grids according to the error estimator η_1^k , such that we obtain a finer dis-16: cretization h_k^3 .
- Set $h = h_k^3$. 17:

18: With
$$q_{\text{old}}^k$$
 and u_{old}^k fixed, determine $q_h^k = q_{h^3}^k$ and $w_h^k = w_{h^3}^k$ by solving (63)

Determine $u_h^k = u_{h_k^3}^k \in V_h = V_{h_k^3}$ by solving 19:

$$A(q_h^k, u_h^k)(v) = f(v) \qquad \forall v \in W_h.$$

- 20:
- Set $q_{\text{old}}^{k+1} = q_h^k (= q_{h_k^3}^k)$ and $u_{\text{old}}^{k+1} (= u_{\text{old},h_{k+1}^3}^{k+1}) = u_h^k (= u_{h_k^3}^k)$. Evaluate $I_{3,h}^{k+1} = I_{3,h_k^3}^{k+1}$ according to (21) and the error estimator $\eta_3^{k+1} = \eta_3^{k+1}(h_k^3)$. Set 21: $h_{k}^{4} = h_{k}^{3}$.
- while (62) is violated do 22:
- Refine grid according to the error estimator η_3^{k+1} , such that we obtain a finer dis-23:
- cretization h_k^4 . Determine $u_{\text{old}}^{k+1} = u_{\text{old},h_{k+1}^4}^{k+1}$, $I_{3,h}^{k+1} = I_{3,h_k^4}^{k+1}$ and $\eta_3^{k+1} = \eta_{3,h_k^4}^{k+1}$ by applying Algorithm 2 24: with m = k + 1 and $h = h_k^4$.
- Set $h_{k+1}^1 = h_k^4$ (i.e. use the current mesh as a starting mesh for the next iteration) 25:

26: Set
$$k = k + 1$$

Algorithm 2. Evaluation of I_3^m

1: Determine

$$u_{\text{old}}^m \in V_h$$
: $A(q_{\text{old}}^m, u_{\text{old}}^m)(v) = f(v)$ $\forall v \in W_h$.

- 2: Evaluate $I_{3,h}^m$ according to (21).
- 3: Evaluate error estimator η_3^m .

Algorithm 3. Inexact Newton method for the determination of a regularization parameter for the IRGNM subproblem from [10]

- 1: Set $\delta_{\beta} = \sqrt{\tilde{\theta} I_{3,h}^{k-1}} / \tau_{\beta}$.
- 2: Compute a Lagrange triple $x_h = (q_h, w_h, z_h)$ to

$$\min_{(q,w)\in Q_h\times V_h} \|C'(u_{\text{old}}^k)(w) + C(u_{\text{old}}^k) - g^{\delta}\|_G^2 + \frac{1}{\beta_k} \|q - q_0\|_Q^2$$
(63)

s.t.
$$A'_u(q^k_{\text{old}}, u^k_{\text{old}})(w, v) + A'_q(q^k_{\text{old}}, u^k_{\text{old}})(q - q^k_{\text{old}})(v) + A(q^k_{\text{old}}, u^k_{\text{old}})(v) - f(v) = 0 \qquad \forall v \in W_h$$

Evaluate $i_k = -I^k - ||C'(u^k_k)|(w_k)| + C(u^k_k) - a^{\delta}||^2$

3: Evaluate $i_h = I_{h,2}^{\kappa} = \|C'(u_{\text{old}}^{\kappa})(w_h) + C(u_{\text{old}}^{\kappa}) - g^{\circ}\|_{G}^{z}$.

- 4: while $i_h > (\tau_{\beta}^2 + \frac{\tilde{\tau}_{\beta}^2}{2})\delta_{\beta}^2$ do 5: Evaluate i'_h (cf. [10]).
- Evaluate error estimator for $i(\beta) = I(w(\beta))$ with $I: w \mapsto I_2(u_{\text{old}}^k, w)$ (cf. [10]). 6:
- Evaluate error estimator for $i'(\beta) = \frac{d}{d\beta}I(w(\beta))$ (cf. [10]). while accuracy requirements (cf. [10]) are violated **do** $\tilde{7}$:
- 8:
- Refine with respect to the corresponding error estimator. 9:
- Compute a Lagrange triple $x_h = (q_h, w_h, z_h)$ to (63). 10:
- Evaluate $i_h = I_{h,2}^k = \|C'(u_{\text{old}}^k)(w_h) + C(u_{\text{old}}^k) g^{\delta}\|_G^2$ 11:

Evaluate i'_h (cf. [10]). 12:

Evaluate error estimator for $i(\beta) = I(w(\beta))$ with $I: w \mapsto I_2(u_{\text{old}}^k, w)$ (cf. [10]). Evaluate error estimator for $i'(\beta) = \frac{d}{d\beta}I(w(\beta))$ (cf. [10]). 13:

- 14:
- Update β according to an inexact Newton method (cf. [10]) $\beta \leftarrow \beta \frac{i_h}{i'}$. 15:
- Compute a Lagrange triple $x_h = (q_h, w_h, z_h)$ to (63). Evaluate $i_h = I_{h,2}^k = \|C'(u_{\text{old}}^k)(w_h) + C(u_{\text{old}}^k) g^{\delta}\|_G^2$. 16:
- 17:

Remark 9. Algorithm 3 corresponds to the algorithm from [10] with the following replacements:

in [10]	here
q	$q-q_0$
T	$F'(q_{ m old}^k)$
g^{δ}	$g^{\delta} - F(q_{\text{old}}^k) + F'(q_{\text{old}}^k)(q_{\text{old}}^k)$
$\tau^2 \delta^2$	$ ilde{ heta}I^k_{3,h}$
$(\tau - \tilde{\tau})^2 \delta^2$	$\frac{ ilde{ heta}}{ ilde{ heta}}I^k_{3,h}$
$(\tau+\tilde{\tau})^2\delta^2$	$\frac{\tilde{\overline{ heta}}}{\tilde{\overline{ heta}}}I^k_{3,h}$

With respect to loops and the solution of PDEs and optimization problems, the algorithm has the following form. (We do not display the refinement loops on lines 5, 15, 22 of Algorithm 1 and on line 8 of Algorithm 3 but only the iteration loops.)

Algorithm 4. Loops in reduced form of discretized IRGNM

- 1: while · · · (Newton iteration) do
- Apply algorithm from [10], i.e. 2.
- 3: while \cdots (Iteration for β_k) do
- Solve linear-quadratic optimization problem (i.e. solve linear PDE). 4:
- Update β and refine eventually. 5:

Solve nonlinear PDE. 6:

In contrast with the nonlinear Tikhonov method

$$\min_{q \in Q} \|F(q) - g^{\delta}\|_{G}^{2} + \frac{1}{\beta} \|q - q_{0}\|_{Q}^{2}$$

investigated in [15] (cf. algorithm 5 below), we have one additional loop, but we only have to solve a linear-quadratic optimization problem instead of a nonlinear problem. On the other hand, we still have to solve (at least) one nonlinear PDE in each outer loop. For this reason we doubt whether algorithm 1 pays off with respect to computation time as compared to the method in [15]. Therefore we do not implement this algorithm, but consider more efficient modifications in [16] (part II of this paper).

Algorithm 5. Loops in Inexact Newton Method (for nonlinear problems)

- 1: while \cdots (Iteration for β) do
- Solve nonlinear optimization problem (i.e. solve nonlinear PDE). 2.
- 3: Update β and refine eventually.

2.5Extension to more general data misfit and regularization terms

Motivated by the increasing use of nonquadratic, non-Hilbert space misfit and regularization terms for modelling, e.g., sparsity of the solution, or non-Gaussian data noise (cf., e.g., [19, 6] for Tikhonov regularization, and [13] for the IRGNM), we now extend our results to a more general setting. To this purpose we consider a more general version of (16):

$$\min_{q \in Q} \mathcal{T}_{\beta}(q) \coloneqq \mathcal{S}(F'(q_{\text{old}}^k)(q - q_{\text{old}}^k) + F(q_{\text{old}}^k), g^{\delta}) + \frac{1}{\beta} \mathcal{R}(q)$$
(64)

with quantities of interest (cf. (10))

$$I_1^k := \mathcal{S}(F'(q_{\text{old}}^k)(q^k - q_{\text{old}}^k) + F(q_{\text{old}}^k), g^\delta) + \frac{1}{\beta} \mathcal{R}(q^k)$$

$$I_2^k := \mathcal{S}(F'(q_{\text{old}}^k)(q^k - q_{\text{old}}^k) + F(q_{\text{old}}^k), g^\delta)$$

$$I_3^k := \mathcal{S}(F(q_{\text{old}}^k), g^\delta)$$

$$I_4^k := \mathcal{S}(F(q^k), g^\delta)$$
(65)

and its discrete counterparts (cf. (19))

$$\min_{q \in Q_h} \mathcal{T}_{\beta,h}(q) \coloneqq \mathcal{S}(F'_h(q^k_{\text{old}})(q - q^k_{\text{old}}) + F_h(q^k_{\text{old}}), g^\delta) + \frac{1}{\beta} \mathcal{R}(q)$$
(66)

with

$$I_{1,h}^{k} \coloneqq S(F_{h_{k}}'(q_{\text{old}}^{k})(q_{h_{k}}^{k} - q_{\text{old}}^{k}) + F_{h_{k}}(q_{\text{old}}^{k}), g^{\delta}) + \frac{1}{\beta} \mathcal{R}(q_{h_{k}}^{k})$$

$$I_{2,h}^{k} \coloneqq S(F_{h_{k}}'(q_{\text{old}}^{k})(q_{h_{k}}^{k} - q_{\text{old}}^{k}) + F_{h_{k}}(q_{\text{old}}^{k}), g^{\delta})$$

$$I_{3,h}^{k} \coloneqq S(F_{h_{k}}(q_{\text{old}}^{k}), g^{\delta})$$

$$I_{4,h}^{k} \coloneqq S(F_{h_{k}}(q_{h_{k}}^{k}), g^{\delta})$$
(67)

(cf. (21)).

The data misfit and regularization functionals $\mathcal S$ and $\mathcal R$ should satisfy

Assumption 1. Let $S: G \times G \to \mathbb{R}$ and $\mathcal{R}: Q \to \mathbb{R}$ have the following properties:

- 1. The mapping $y \mapsto \mathcal{S}(y, g^{\delta})$ is convex.
- 2. S is symmetric, i.e. $S(y, \tilde{y}) = S(\tilde{y}, y)$ for all $y, \tilde{y} \in G$.
- 3. S is positive definite, i.e. $S(y, \tilde{y}) \ge 0$ and S(y, y) = 0 for all $y, \tilde{y} \in G$.
- 4. For all $y, \tilde{y}, \hat{y} \in G$ there exists a constant $c_{\mathcal{S}}$ such that $\mathcal{S}(y, \tilde{y}) \leq c_{\mathcal{S}}(\mathcal{S}(y, \hat{y}) + \mathcal{S}(\hat{y}, \tilde{y}))$.
- 5. The regularization operator \mathcal{R} is proper (i.e. the domain of \mathcal{R} is non-empty) and convex.

where the domain of an operator $\mathcal{R} \colon M \to \mathbb{R}$ should be understood as

$$\mathcal{D}(\mathcal{R}) \coloneqq \{ m \in M | \ \mathcal{R}(m) \neq \infty \}.$$

Remark 10. In fact, it suffices to require S(y,y) = 0 only for y = g, i.e. for the exact data in Item 3 in Assumption 1, but since Item 3 is a more "natural" assumption in terms of general operator properties, we stick with the stronger assumption Item 3.

We refer once more to [13] where convergence and convergence rates for the IRGNM have already been established in an even more general (continuous) setting and mention that we here consider a somewhat simpler situation with stronger assumptions on S, \mathcal{R} , since our main intention is to demonstrate transferrability of the adaptive discretization concept. Moreover note, that we rely on a different choice of the regularization parameter here. The results obtained here will allow us to easily establish convergence rates results for an exact penalty formulation of an all-at-once formulation of the IRGNM in [16] (part II of this paper).

Although we will, again, restrict ourselves to Hilbert spaces in the next sections, at this point we discuss convergence in a Banach space setting to emphasize the generality of the subsequent results. To this purpose we introduce the *Bregman distance*

$$D_{\mathcal{R}}^{\xi}(q,\overline{q}) \coloneqq \mathcal{R}(q) - \mathcal{R}(\overline{q}) - \langle \xi, q - \overline{q} \rangle_{Q^*,Q}$$
(68)

with some $\xi \in \partial \mathcal{R}(q) \subset Q^*$, which coincides with $\frac{1}{2} ||q - q^{\dagger}||_Q^2$ for $\mathcal{R}(q) = \frac{1}{2} ||q - q_0||_Q^2$ and $\xi = q^{\dagger} - q_0$ in a Hilbert space Q.

Well-definedness (i.e. for every $g^{\delta} \in G$ and $\beta_k > 0$ there exists a solution $q_{h_k}^k$ to (66)) and stable dependence on the data (i.e. for every fixed $\beta_k > 0$ the solution $q_k^{h_k}$ depends continuously on g^{δ}) can be shown under the following assumptions (cf., e.g., Assumption 1.32 in [19] or Remark 2.1 in [13])

Assumption 2. 1. Q and G are Banach spaces, with which there are associated topologies τ_Q and τ_G , which are weaker than the norm topologies.

- 2. The mapping $y \mapsto \mathcal{S}(y, g^{\delta})$ is sequentially lower semi-continuous with respect to τ_G .
- 3. $F'(q_{\text{old}}^k): Q \to G$ is continuous with respect to the topologies τ_Q and τ_G .
- 4. $\mathcal{R}: Q \to [0, +\infty]$ is proper, convex and τ_Q -lower semi continuous.
- 5. $\mathcal{D} \coloneqq \mathcal{D}(F) \cap \mathcal{D}(\mathcal{R}) \neq \emptyset$ is closed with respect to τ_Q .

6. For every C > 0 the set

$$\mathcal{C}(C) \coloneqq \{ q \in \mathcal{D} \colon \mathcal{R}(q) \le C \} \,, \tag{69}$$

is τ_Q -sequentially compact in the following sense: every sequence $(q_n)_{n \in \mathbb{N}}$ in $\mathcal{C}(C)$ has a subsequence, which is convergent in Q with respect to the τ_Q -topology.

For well-definedness of the a posteriori chosen regularization parameter β_k we refer to Lemma 1 and Theorem 3 in [18].

Remark 11. For Hilbert spaces Q and G and the choice $S(y, \tilde{y}) \coloneqq \frac{1}{2} ||y - \tilde{y}||_G^2$ and $\mathcal{R}(q) \coloneqq \frac{1}{2} ||q - q_0||_Q^2$ Assumption 1 and Assumption 2 are obviously fulfilled. As for examples in a real Banach spaces setting, we refer to [18, 14, 19].

Consistently the conditions (27) and (28) on F are generalized to the following two assumptions.

Assumption 3. Let the reduced forward operator F be continuous with respect to τ_Q, τ_G and satisfy

$$(q_n \stackrel{'Q}{\rightarrow} q \land \mathcal{S}(F(q_n), g) \to 0 \Rightarrow (q \in \mathcal{D}(F) \land F(q) = g)$$

for all $(q_n)_{n \in \mathbb{N}} \subseteq Q$.

Assumption 4. Let the generalized tangential cone condition

$$\mathcal{S}(F(q), F(\bar{q}) + F'(q)(q - \bar{q})) \le c_{tc}^2 \mathcal{S}(F(q), F(\bar{q}))$$

hold for all $q, \bar{q} \in Q$ in a neighborhood of q_0 for some $0 < c_{tc} < 1$.

Moreover, the source condition (41) is replaced by Assumption 5.

Assumption 5. Let the multiplicative variational inequality

$$|\langle \xi, q - q^{\dagger} \rangle_{Q^*, Q}| \le c D_{\mathcal{R}}^{\xi}(q, q^{\dagger})^{1/2} f\left(\frac{\mathcal{S}(F(q), F(q^{\dagger}))}{D_{\mathcal{R}}^{\xi}(q, q^{\dagger})}\right)$$

for all $q \in \mathcal{D}(F)$ hold.

Based on this groundwork, we can now formulate a convergence theorem similar to Theorem 1:

Theorem 3. Let Assumption 2 be satisfied, let $q^{\dagger} \in \mathcal{B}_{\rho}(q_0)$ be a solution to (1) and let F be continuous and satisfy Assumption 3, Assumption 4 with c_{tc} sufficiently small. Let, further, $\tau > 0$ be chosen sufficiently large such that

$$c_{\mathcal{S}}\left(c_{\mathcal{S}}c_{tc}^{2} + \frac{1 + c_{\mathcal{S}}c_{tc}^{2}}{\tau^{2}}\right) < \underline{\tilde{\theta}} \quad and \quad 0 < \frac{c_{\mathcal{S}}\overline{\tilde{\theta}} + c_{\mathcal{S}}^{2}c_{tc}^{2}}{1 - c_{\mathcal{S}}^{2}c_{tc}^{2}} < 1$$
(70)

and let

$$S(g, g^{\delta}) \le \delta^2 \,. \tag{71}$$

Finally, let for the discretization error with respect to the quantities of interest (65), (67) estimates (24) hold, where η_1^{ϵ} , η_2^{ϵ} , η_3^{ϵ} , η_4^{κ} are selected such that

$$\eta_1^k + c_S^2 c_{tc}^2 \eta_3^k \le \left(\frac{\tilde{\theta}}{\ell} - c_S \left(c_S c_{tc}^2 + \frac{1 + c_S c_{tc}^2}{\tau^2}\right)\right) I_{3,h}^k \tag{72}$$

as well as (31), the first part of (32) and

$$(1+c_3)\frac{c_s\bar{\theta} + c_s^2c_{t_c}^2}{1 - c_s^2c_{t_c}^2} \le c_2 < 1$$
(73)

hold for some constants $c_1, c_2, c_3 > 0$, and a sequence $r^k \to 0$ as $k \to \infty$, where (73) is possible due to the right inequality in (70).

Then with β_k and $h = h_k$ fulfilling (25) and k_* selected according to (26) there holds

(i) For any solution $q^{\dagger} \in \mathcal{B}_{\rho}(q_0)$ of (1)

$$\mathcal{R}(q_h^k) \le \mathcal{R}(q^\dagger) \quad \forall k < k_* \,, \tag{74}$$

- (ii) k_* is finite,
- (iii) $q_{\text{old}}^{k_*} = q_{h_{k_*-1}}^{k_*-1}$ converges (weakly) subsequentially to a solution of (1) as $\delta \to 0$ in the sense that it has a τ_Q convergent subsequence and each τ_Q convergent subsequence converges to a solution of (1). If the solution q^{\dagger} to (1) is unique, then $q_{\text{old}}^{k_*}$ converges with respect to τ_Q to q^{\dagger} as $\delta \to 0$.

Proof. The proof basically follows the lines of the proof of Theorem 1, where we have to replace the specific fitting and regularization terms by S and \mathcal{R} :

(i): For all $k < k_*$ and any solution q^{\dagger} of (1) we have by (24) and minimality of q^k

$$I_{1,h}^{k} \leq I_{1}^{k} + \eta_{1}^{k} \leq \mathcal{S}(F'(q_{\text{old}}^{k})(q^{\dagger} - q_{\text{old}}^{k}) + F(q_{\text{old}}^{k}), g^{\delta}) + \frac{1}{\beta_{k}}\mathcal{R}(q^{\dagger}) + \eta_{1}^{k}.$$
 (75)

In here, according to (71), (26) and Assumption 4, as well as the inequality $(a + b)^2 \leq 2a^2 + 2b^2$ for arbitrary $a, b \in \mathbb{R}$ we can estimate as follows

$$S(F'(q_{\text{old}}^{k})(q^{\dagger} - q_{\text{old}}^{k}) + F(q_{\text{old}}^{k}), g^{\delta}) \leq c_{\mathcal{S}} \left(\mathcal{S}(g, F'(q_{\text{old}}^{k})(q^{\dagger} - q_{\text{old}}^{k}) + F(q_{\text{old}}^{k})) + \delta^{2} \right)$$

$$\leq c_{\mathcal{S}} \left(c_{tc}^{2} \mathcal{S}(g, F(q_{\text{old}}^{k})) + \delta^{2} \right)$$

$$\leq c_{\mathcal{S}} \left(c_{tc}^{2} (c_{\mathcal{S}} (\mathcal{S}(g^{\delta}, F(q_{\text{old}}^{k})) + \delta^{2}) + \delta^{2} \right)$$

$$\leq c_{\mathcal{S}}^{2} c_{tc}^{2} I_{3}^{k} + c_{\mathcal{S}} (1 + c_{\mathcal{S}} c_{tc}^{2}) \delta^{2}$$

$$\leq c_{\mathcal{S}}^{2} c_{tc}^{2} (I_{3,h}^{k} + \eta_{3}^{k}) + c_{\mathcal{S}} (1 + c_{\mathcal{S}} c_{tc}^{2}) \frac{I_{3,h}^{k}}{\tau^{2}}$$

$$\leq c_{\mathcal{S}} \left(c_{\mathcal{S}} c_{tc}^{2} + \frac{1 + c_{\mathcal{S}} c_{tc}^{2}}{\tau^{2}} \right) I_{3,h}^{k} + c_{\mathcal{S}}^{2} c_{tc}^{2} \eta_{3}^{k} . \quad (76)$$

On the other hand, from (25) and the fact that $I_{1,h}^k = I_{2,h}^k + \frac{1}{\beta_k} \mathcal{R}(q_h^k)$ (cf. (23)) there follows that

$$I_{1,h}^{k} = I_{2,h}^{k} + \frac{1}{\beta_{k}} \mathcal{R}(q_{h}^{k}) \ge \underline{\tilde{\theta}} I_{3,h}^{k} + \frac{1}{\beta_{k}} \mathcal{R}(q_{h}^{k}), \qquad (77)$$

which together with the previous inequality and (72) gives

$$\begin{split} \underline{\tilde{\theta}}I_{3,h}^{k} &+ \frac{1}{\beta_{k}}\mathcal{R}(q_{h}^{k}) \leq I_{1}^{k} + \eta_{1}^{k} \\ &\leq \mathcal{S}(F'(q_{\text{old}}^{k})(q^{\dagger} - q_{\text{old}}^{k}) + F(q_{\text{old}}^{k}), g^{\delta}) + \frac{1}{\beta_{k}}\mathcal{R}(q^{\dagger}) + \eta_{1}^{k} \\ &\leq c_{\mathcal{S}}\left(c_{\mathcal{S}}c_{tc}^{2} + \frac{1 + c_{\mathcal{S}}c_{tc}^{2}}{\tau^{2}}\right)I_{3,h}^{k} + c_{\mathcal{S}}^{2}c_{tc}^{2}\eta_{3}^{k} + \frac{1}{\beta_{k}}\mathcal{R}(q^{\dagger}) + \eta_{1}^{k} \\ &\leq \underline{\tilde{\theta}}I_{3,h}^{k} + \frac{1}{\beta_{k}}\mathcal{R}(q^{\dagger})\,, \end{split}$$

which implies (74).

(ii): Furthermore, for all $k < k_*$ we have by the triangle inequality as well as Assumption 4 and (25)

$$\begin{aligned}
I_{4}^{k} &= \mathcal{S}(F(q^{k}), g^{\delta}) \\
&\leq c_{\mathcal{S}} \left(\mathcal{S}(F'(q_{\text{old}}^{k})(q^{k} - q_{\text{old}}^{k}) + F(q_{\text{old}}^{k}), g^{\delta}) + \mathcal{S}(F'(q_{\text{old}}^{k})(q^{k} - q_{\text{old}}^{k}) + F(q_{\text{old}}^{k}), F(q^{k})) \right) \\
&\leq c_{\mathcal{S}} \left(I_{2}^{k} + c_{tc}^{2} \mathcal{S}(F(q^{k}), F(q_{\text{old}}^{k})) \right) \\
&\leq c_{\mathcal{S}} \left(I_{2,h}^{k} + \eta_{2}^{k} \right) + c_{\mathcal{S}}^{2} c_{tc}^{2} \left(\mathcal{S}(F(q^{k}), g^{\delta}) + \mathcal{S}(F(q_{\text{old}}^{k}), g^{\delta}) \right) \\
&= c_{\mathcal{S}} \left(\overline{\tilde{\theta}} I_{3,h}^{k} + \eta_{2}^{k} \right) + c_{\mathcal{S}}^{2} c_{tc}^{2} \left(I_{4}^{k} + I_{3}^{k} \right) ,
\end{aligned}$$
(78)

which implies

$$I_{4}^{k} \leq \frac{1}{1 - c_{\mathcal{S}}^{2} c_{tc}^{2}} \left(c_{\mathcal{S}} \overline{\theta} I_{3,h}^{k} + c_{\mathcal{S}} \eta_{2}^{k} + c_{\mathcal{S}}^{2} c_{tc}^{2} I_{3}^{k} \right) \,.$$

With (24) and (32) we can further deduce

$$\begin{split} I_{4,h}^{k} &\leq \frac{1}{1 - c_{\mathcal{S}}^{2} c_{tc}^{2}} \left((c_{\mathcal{S}} \tilde{\overline{\theta}} + c_{\mathcal{S}}^{2} c_{tc}^{2}) I_{3,h}^{k} + c_{\mathcal{S}} \eta_{2}^{k} + c_{\mathcal{S}}^{2} c_{tc}^{2} \eta_{3}^{k} \right) + \eta_{4}^{k} \\ &\leq \frac{c_{\mathcal{S}} \tilde{\overline{\theta}} + c_{\mathcal{S}}^{2} c_{tc}^{2}}{1 - c_{\mathcal{S}}^{2} c_{tc}^{2}} (1 + c_{3}) I_{4,h}^{k-1} + \frac{1}{1 - c_{\mathcal{S}}^{2} c_{tc}^{2}} \left(r^{k} + c_{\mathcal{S}} \eta_{2}^{k} + c_{\mathcal{S}}^{2} c_{tc}^{2} \eta_{3}^{k} \right) + \eta_{4}^{k} \\ &\leq c_{2} I_{4,h}^{k-1} + \frac{1}{1 - c_{\mathcal{S}}^{2} c_{tc}^{2}} \left(r^{k} + c_{\mathcal{S}} \eta_{2}^{k} + c_{\mathcal{S}}^{2} c_{tc}^{2} \eta_{3}^{k} \right) + \eta_{4}^{k} \,. \end{split}$$

With the notation

$$a^{i} \coloneqq \frac{1}{1 - c_{\mathcal{S}}^{2} c_{tc}^{2}} \left(r^{i} + c_{\mathcal{S}} \eta_{2}^{i} + c_{\mathcal{S}}^{2} c_{tc}^{2} \eta_{3}^{i} \right) + \eta_{4}^{i} \quad \forall i \in \{1, 2, \dots, k\}$$
(79)

there follows recursively

$$I_{4,h}^{k} \le c_{2}^{k} I_{4,h}^{0} + \sum_{j=0}^{k-1} c_{2}^{j} a^{k-j} .$$

$$(80)$$

Note that by the second part of (31), the second part of (32) and the fact that $r^k \to 0$ as $k \to \infty$ (by definition of r^k), we have $c_2^k I_{4,h}^0 + \sum_{j=0}^{k-1} c_2^j a^{k-j} \to 0$ as $k \to \infty$. So, if the discrepancy principle never got active (i.e., $k_* = \infty$), the sequence $(I_{4,h}^k)_{k \in \mathbb{N}}$ and therewith by assumption (31) also $(I_{3,h}^k)_{k \in \mathbb{N}}$ would be bounded by a sequence tending to zero as $k \to \infty$, which implies that $I_{3,h}^k$ would fall below $\tau^2 \delta^2$ for k sufficiently large, thus yielding a contradiction. Hence the stopping index $k_* < \infty$ is well-defined and finite.

(iii): With (71), (24), (31) and definition of k_* , we have

$$S(F(q_{\text{old}}^{k_*}), g) \le c_S \left(S(F(q_{\text{old}}^{k_*}), g^{\delta}) + \delta^2 \right) \le c_S \left(I_{3,h}^{k_*} + \eta_3^k + \delta^2 \right) \le c_S \left((1+c_1)I_{3,h}^{k_*} + \delta^2 \right) \le c_S \left((1+c_1)\tau^2 + 1 \right) \delta^2 \to 0$$
(81)

as $\delta \to 0$. By (i), (ii) and (69) in assumption 2 $q_{\text{old}}^{k_*} = q_{h_{k_*-1}}^{k_*-1}$ has a τ_Q convergent subsequence $(q_{\text{old}}^{k_*(\delta_l)})_{l \in \mathbb{N}}$ and due to Assumption 3 and (81), the limit of every τ_Q convergent subsequence is a solution to F(q) = g.

It is readily checked that (like in the case of \mathcal{R} , \mathcal{S} being defined by squared Hilbert space norms as considered in Theorem 1 of [15]) any approximation \tilde{q} of a solution q^{\dagger} of F(q) = gwith $||g - g^{\delta}|| \leq \delta$ such that

$$\mathcal{R}(\tilde{q}) \leq \mathcal{R}(q^{\dagger}) \text{ and } \mathcal{S}(F(\tilde{q}), g^{\delta}) \leq \hat{\tau}^2 \delta^2$$

with $\hat{\tau}$ independent of δ , as well as the variational inequality (5) holds, satisfies the rate estimate

$$D^{\xi}_{\mathcal{R}}(\tilde{q}, q^{\dagger}) \leq \frac{C^2 \delta^2}{\Theta^{-1}\left(\frac{\bar{C}}{c}\delta\right)} = c^2 f^2(\Theta^{-1}(\frac{\bar{C}}{c}\delta))$$

with $\bar{C}^2 = c_S(\hat{\tau}^2 + 1).$

Hence we directly obtain from (74) and the definition of k_* according to (26) the following rates result.

Theorem 4. Let the conditions of Theorem 3 and additionally the variational inequality (5) for some function f with (42) be fullfiled.

Then there exists a $\bar{\delta} > 0$ and a constant $\bar{C} > 0$ independent of δ such that for all $\delta \in (0, \bar{\delta}]$

$$D_{\mathcal{R}}^{\xi}(q_h^{k_*}, q^{\dagger}) \le \frac{\bar{C}^2 \delta^2}{\Theta^{-1}\left(\frac{\bar{C}}{c}\delta\right)} = c^2 f^2(\Theta^{-1}(\frac{\bar{C}}{c}\delta)), \qquad (82)$$

where $\Theta(\lambda) \coloneqq f(\lambda) \sqrt{\lambda}$.

Proof. The proof can be done analogously to the proof of Theorem 1 in [15] with the same replacements as in the proof of Theorem 3. \Box

3 Conclusions and Remarks

In this paper we consider the iteratively regularized Gauss-Newton method and its adaptive discretization by means of goal oriented error estimators. Our aim is to recover convergence as in the continuous setting for discretized hence approximate computations. The key result is that control of a small number (four) real valued quantities per Newton step suffices to guarantee convergence and convergence rates. While we have studied the problem in a reduced form here, using the parameter-to-solution map, the related paper [16](part II of this paper) develops and studies all-at-once formulations. Numerical tests are provided in part II of this paper [16].

4 Acknowledgments

The authors wish to thank Boris Vexler for fruitful discussions, for his support, and for his valuable suggestions after a careful reading of this manuscript.

Moreover, we gratefully acknowledge financial support by the German Science Foundation (DFG) within the grant KA 1778/5-1 and VE 368/2-1 "Adaptive Discretization Methods for the Regularization of Inverse Problems".

References

- A.B. BAKUSHINSKY AND M. KOKURIN: Iterative Methods for Approximate solution of Inverse Problems. Kluwer Academic Publishers, Dordrecht, 2004.
- [2] R. BECKER, R. RANNACHER, An Optimal Control Approach to a-Posteriori Error Estimation Acta Numerica 2011 ed. A Iseries, (Cambridge University Press), 1–102.
- [3] R. BECKER, B. VEXLER, A posteriori error estimation for finite element discretizations of parameter identification problems *Numer. Math.* 96 (2004), 435–59
- [4] B. BLASCHKE(-KALTENBACHER), A. NEUBAUER, O. SCHERZER, On convergence rates for the iteratively regularized Gauß–Newton method, *IMA J. Numer. Anal.* 17 (1997), 421–436.
- [5] H.-G. BOCK, Randwertproblemmethoden zur Parameteridentifizierung in Systemen nichtlinearer Differentialgleichungen, Bonner Mathematische Schriften 183, Bonn, 1987.
- [6] J. FLEMMING: Theory and examples of variational regularisation with non-metric fitting functionals, *Journal of Inverse and Ill-Posed Problems*, 18(6), 2010.
- [7] M. BURGER AND W. MÜHLHUBER: Iterative regularization of parameter identification problems by sequential quadratic programming methods, *Inverse Problems* 18 (2002) 943–969.
- [8] M. BURGER AND W. MÜHLHUBER: Numerical approximation of an SQP-type method for parameter identification, SIAM J. Numer. Anal. 40 (2002), 1775–1797.
- [9] H.W. ENGL, M. HANKE, A. NEUBAUER, Regularization of Inverse Problems, Kluwer, Dordrecht, 1996.
- [10] A. Griesbaum, B. Kaltenbacher, B. Vexler: Efficient computation of the Tikhonov regularization parameter by goal-oriented adaptive discretization, *Inverse Problems* 24 (2008),
- [11] M. HANKE, A regularization Levenberg-Marquardt scheme, with applications to inverse groundwater filtration problems, *Inverse Problems* 13 (1997), 79–95.
- [12] T. HOHAGE, Iterative Methods in Inverse Obstacle Scattering: Regularization Theory of Linear and Nonlinear Exponentially Ill-Posed Problems, PhD thesis, University of Linz, 1999.
- [13] T. HOHAGE AND F. WERNER: Iteratively regularized Newton methods with general data misfit functionals and applications to Poisson data, *Numerische Mathematik* 123 (2013), 745-779.
- [14] B. KALTENBACHER, B. HOFMANN: Convergence rates for the Iteratively Regularized Gauss-Newton method in Banach spaces, *Inverse Problems* 26 (2010) 035007.

- [15] B. KALTENBACHER, A. KIRCHNER, B. VEXLER: Adaptive discretizations for the choice of a Tikhonov regularization parameter in nonlinear inverse problems, *Inverse Problems* 27 (2011) 125008.
- [16] B. KALTENBACHER, A. KIRCHNER, B. VEXLER: Goal oriented adaptivity in the IRGNM for parameter identification in PDEs II: all-at once formulations, submitted.
- [17] B. KALTENBACHER, A. NEUBAUER, O. SCHERZER: Iterative Regularization Methods for Nonlinear Ill-Posed Problems. Walter de Gruyter, Berlin – New York, 2008.
- [18] B. KALTENBACHER, F. SCHÖPFER, AND T. SCHUSTER: Convergence of some iterative methods for the regularization of nonlinear ill-posed problems in Banach spaces. *Inverse Problems* 25 (2009), 065003 (19pp). DOI:10.1088/0266-5611/25/6/065003.
- [19] C. PÖSCHL. *Tikhonov Regularization with General Residual Term.* PhD thesis, Universität Innsbruck, October 2008.
- [20] A. RIEDER, On the regularization of nonlinear ill-posed problems via inexact Newton iterations, *Inverse Problems* 15 (1999), 309–327.