## G-quadruplexes and mRNA localization

Valentina Agoni valentina.agoni@unipv.it

G-quadruplexes represent a novelty for molecular biology. Their role inside the cell remains mysterious.

We investigate a possible correlation with mRNA localization.

In particular, we hypothesize that Gquadruplexes influence fluid dynamics.

Localization of mRNA molecules within the cytoplasm recently emerged as a prevalent mechanism for cell polarization, thus underlying developmental processes.

[1] reports 5 adventages for a cell of transporting mRNAs rather than proteins:

(i) transport costs are reduced, as several protein molecules can be translated from a single RNA

molecule

(ii) transporting mRNAs can prevent proteins from acting ectopically

(iii) localized translation can facilitate incorporation of proteins into macromolecular complexes [2]

(iv) nascent proteins may have properties distinct from pre-existing copies, by virtue of post-translational modifications or through chaperone-aided folding pathways [3]

(v) a major advantage of mRNA targeting is that it allows fine-tuning of gene expression in both space and time [4] [5].

Until now three distinct mechanisms have been identified underlying mRNA localization: localized protection from degradation, diffusion-coupled local entrapment and directed transport along a polarized cytoskeleton.

But what can control mRNA localization? At the same time the role of fluid dynamics in developmental biology has been acknowledged [6]. Human conception, indeed fertilization in general, takes place in a fluid.

Plant roots direct their growth in response to gravity, light, and mechanical stimuli. Local changes in hormone concentration have been shown to be part of the response mechanisms that result in those tropic growth.

First of all it is very interesting to note that gravistimulation and mechanical stimulation act on the same transcripts [7].

Furthermore [8] indicates that salt-induced altered gravitropism of root growth is mediated by ion disequilibrium.

We know that objects with the same mass do fall at the same speed in the vacuum and hit the ground at the same time, but air resistance hinders this fact. For example, if you drop a tennis ball and a feather at the same time on the Earth, obviously the ball falls faster.

In addition we have to consider parasitic drag. Parasitic drag is caused by moving a solid object through a fluid medium (in the case of hydrodynamics, more specifically, a liquid medium). Parasitic drag is made up of many components, the most prominent being form drag. Skin friction and interference drag are also major components of parasitic drag. The shape of the object is the most important factor in form drag or pressure drag: bodies with a larger apparent cross-section will have a higher drag than thinner bodies. The fluid we are considering is cytoplasm while the objects of our analysis are mRNAs.

Cytoplasmic streaming, also termed cyclosis, is a persistent circulation of the fluid contents of cells and we assume it to flow in every direction [6]. However what happens if we consider a very short instant of time the flow goes in one direction or an higher gravity system? Well, we talk about the tennis ball and the feather in the presence of gravity, now we also have to consider that the shape of the object is the most important factor in form drag or pressure drag: bodies with a larger apparent cross-section will have a higher drag than thinner bodies.

In [7] root apices from 7-day-old etiolated *Arabidopsis* seedlings were harvested and analyzed for relative changes in transcript

levels in response to gravistimulation and mechanical stimulation using the *Arabidopsis* 

ATH1 GeneChip (Affymetrix) in a time course experiment. The analysis of whole genome microarrays on the differential regulation of transcript abundance reveals changes during the first hour after gravity [7]. Four-stranded G-quadruplex nucleic acid structures are G-rich DNA and RNA sequences that can fold into non-canonical secondary structures [9].

G-quadruplexes are coded by different sequences (Figure 1) but we identify a common motif in the AGGG sequence.

## Sequence

```
5'-GG(TTAGGG)<sub>4</sub>TTAG-3'
5'-GG(TTAGGG)<sub>4</sub>TTAG-3'/3'-C(AATCCC)<sub>4</sub>AAT-5'
5'-GGCATAGTGCGTGGGCG-3'
5'-TGAGGGTGGGTAGGGTGGGTAA-3'
5'-AGGGAGGGCGCTGGGAGGAGGGG-3'
5'-CGGCGGGGCGCGAGGGGGGGGGGGGGGG
5'-GGTTGGTGGGTTGG-3'
5'-CAGUACAGAUCUGUACUG-3'
5'-(TTAGGG)<sub>3</sub>-3'
```

**Figure 1.** Oligonucleotides used in enzymelinked immunosorbent assays. Modified from [9].

We considered the sequences of some

mRNAs shown to coherently increase or decrease in response to gravity and movement [7] analyzing the frequency of AGGG sequences.

It emerges that increasing-mRNAs present a significantly lower AGGG content respect to decreasing-mRNAs.

This can be reasonable if we think that increasing gravity the number of Gquadruplexes in transcripts should be lower (on average). This is the reason why we performed a one tailed-test.

Nevertheless the two populations of transcripts are heterogeneous because also in these conditions there should be mRNAs at different levels in the cellular cytoplasm.

In consideration of all these punctualizations and due to the limited number of samples we performed a Mann-Whitney test for not-Gaussian populations.The results are shown in Table 1.

In conclusion mRNAs could arrange in the cell according to their weight and shape also in terms of presence and number of G-quadruplexes.

But is this the major phenomenon regulating mRNA localization?

NCBI Reference Sequence	AGGG percentage	NCBI Reference Sequence	AGGG percentage
Sequences increasing with gravity		Sequences decreasing with gravity	
NM_179725.1	0.041701418	NM_117622.2	0.160578081
DQ105587.1	0.07199424	AY150484.1	0.244498778
AF332444.1	0.258397933	AY093066.1	0.155884645
NM_117342.2	0.066934404	BT004357.1	0.384615385
BT002780.1	0.092936803	AY096456.1	0.666666667
NM_104261.2	0.145772595	BT003121.1	0.225988701
BT010322.1	0.174216028	AY051075.1	0.199071002
AY045865.1	0.186046512	BT003667.1	1.089324619
BT012546.1	0.228310502	BT025023.1	0.366300366
AK317670.1	0.291828794	AY142594.1	0.185070944
NM_116393.1	0.260416667	AY081574.1	0.257069409
BT020422.1	0.229885057	AF361097.1	0.336021505
average	0.170703413		0.355924175
stdev	0.085971011		0.270824472
Z-ratio			-2.0496
U-value			36

## The critical value of U at $p \le 0.05$ is 42. Therefore, the result is significant at $p \le 0.05$ .

 Table 1. The AGGG/sequence length percentages relative to genes responsive to gravity and mechanical stimulation [7] are reported.

[1] L. A. Mingle, N. N. Okuhama, J. Shi, R. H. Singer, J. Condeelis, and G. Liu, "Localization of all seven messenger RNAs for the actin-polymerization nucleator Arp2/3 complex in the protrusions of fibroblasts," *J. Cell Sci.*, Vol. 118, 2005, pp. 2425-2433.

[2] A. C. Lin, and C. E. Holt, "Local translation and directional steering in axons," *EMBO J.*, Vol. 26, 2007, pp.3729-3736.

[3] C.Medioni, K. Mowry and F.Besse, "Principles and roles of mRNA localization in animal development", *Development. Sep.*,Vol. 139, 2012, pp.3263-76.

[4] G. Baj, E. Leone, M. V. Chao and E. Tongiorgi, "Spatial segregation of BDNF transcripts enables BDNF to differentially shape distinct dendritic compartments," *Proc. Natl. Acad. Sci. USA*, Vol. 108, 2011, pp.16813-16818.

[5] F. Besse and A. Ephrussi, Translational control of localized mRNAs: restricting protein synthesis in space and time," *Nat. Rev. Mol. Cell Biol.*, Vol. 9, 2008, pp.971-980.

[6] J.H. Cartwright, O. Piro, I. Tuval, "Fluid dynamics in developmental biology: moving fluids that shape ontogeny," *HFSP J.*, Vol. 3, 2009, pp.77-93.

[7] J.M. Kimbrough, C.S. Brown, Winter, H. Sederoff, "Co-expression and hormonal regulation of genes throughout gravity and mechanical stimulation in the *Arabidopsis* root apex," *Gravitational and Space Biology*, Vol. 18, 2005, pp.117–119.

[8] Sun F, Zhang W, Hu H, Li B, Wang Y, Zhao Y, Li K, Liu M, Li X. "Salt modulates gravity signaling pathway to regulate growth direction of primary roots in Arabidopsis," *Plant Physiol.* Vol.146, N° 1, 2008, pp.178-88.

[9] G. Biffi, D. Tannahill, J. McCafferty and S. Balasubramanian, "Quantitative visualization of DNA G-quadruplex structures in human cells," *Nature Chemistry*, Vol. 5 2013, pp.182-186.

[10] E. Louie, J. Ott and J. Majewski, "Nucleotide Frequency Variation Across Human Genes," *Genome Res.*, Vol. 13, N° 12, 2003, pp.2594–2601.

[11] J. Ramchatesingh, A.M. Zahler, K.M. Neugebauer, M.B. Roth and T.A. Cooper, "A subset of SR proteins activates splicing of the cardiac troponin T alternative exon by direct interactions with an exonic enhancer," *Mol. Cell. Biol.*, Vol. 15, 1995, pp. 4898-4907.

[12] A.J. McCullough and S.M. Berget, "G triplets located throughout a class of small vertebrate introns enforce intron borders and regulate splice site selection," *Mol. Cell. Biol.*, Vol. 17, 1997, pp. 4562-4571.

[13] A.J. McCullough and M.A. Schuler, "Intronic and exonic sequences modulate 5' splice site selection in plant nuclei," *Nucleic Acids Res.*, Vol. 25, 1997 pp.1071-1077.