

Modified (G'/G) -expansion method for solving nonlinear partial differential equations

Mahouton Norbert Hounkonnou* and Rolland Finangnon Kanfon†

*International Chair in Mathematical Physics and Applications
(ICMPA-UNESCO Chair), University of Abomey-Calavi,
072B.P.50, Cotonou, Rep. of Benin*

E-mails: *norbert.hounkonnou@cipma.uac.bj, †rolland.kanfon@cipma.uac.bj

Abstract

This paper addresses a modified (G'/G) -expansion method to obtain new classes of solutions to nonlinear partial differential equations (NPDEs). The cases of Burgers, KdV and Kadomtsev-Petviashvili NPDEs are exhaustively studied. Relevant graphical representations are shown in each case.

1 Introduction

Over the second half of the past century, there was a flurry of activities regarding the investigation, with a solid mathematical background and a wide range of physical observations and practical applications, of nonlinearities in complex phenomena governing our universe. Indeed, nonlinear equations, widely used to describe these complex phenomena, pervade many branches of mathematics and physics, including fluid mechanics, plasma physics, quantum electrodynamics, solid-state physics, nuclear and atomic physics, plasma waves, biology, and so on. See, for instance, [1, 2, 3] (and references therein) for more details from both theoretical and experimental viewpoints. The knowledge of exact solutions to these equations constitutes certainly one of the keys for better understanding the nonlinearities. This explains the search for suitable analytical methods for solving nonlinear equations. The most popular ones are today the variational iteration method [4], [5], inverse scattering method [6], Hirota bi-linear form [7], Painlevé analysis [8], direct algebraic method [9], tanh-function method [10], [11], [12], sine-cosine method [13], Darboux transformation [14, 15], homotopy analysis method, (G'/G) -expansion method [16], [17] and their extensions.

This paper addresses a modified (G'/G) -expansion method to obtain new classes of solutions to nonlinear partial differential equations (NPDEs). This approach consists, first, in linearizing the NPDE, and secondly, in integrating the resulting partial differential equation (PDE) with a non-zero integration constant dependent constant. Then, the (G'/G) -expansion method is applied to deduce the solutions of the nonlinear PDE.

The paper is organized as follows. In Section (2) we describe the (G'/G) -method used to solve the NPDEs. We show that any solution ϕ of the NPDE can be expanded as polynomials depending on the new variable G'/G , where G' is the derivative of G with respect to the coordinate variables. The general form of ϕ is also given. In Section (3) we explicitly solve the Burgers partial differential

equation, the Kadomtsev-Petviashvili and KdV equations. Relevant graphical representations are exhibited in each case. Section (4) is devoted to concluding remarks.

2 Description of the method

The nonlinear partial differential equations (NPDEs), although difficult to solve, are object of intensive study in recent literature as shown in [1]-[25] and references therein. In this section, we consider the following types of NPDEs:

$$Q(u, u_t, u_x, u_{xx}, \dots) = 0, \quad x, t \in \mathbb{R}, \quad (1)$$

$$P(u, u_t, u_x, u_y, u_{tt}, u_{xx}, u_{yy}, u_{tx}, u_{ty}, u_{xy}, \dots) = 0, \quad (2)$$

$$u_t = \frac{\partial u}{\partial t}, u_x = \frac{\partial u}{\partial x}, u_{xx} = \frac{\partial^2 u}{\partial x^2}, u_{tx} = \frac{\partial^2 u}{\partial t \partial x}, \dots; \quad x, y, t \in \mathbb{R}, \quad (3)$$

where $u = u(x, t) \in C^\infty(\mathbb{R}^2)$ and $v = v(x, y, t) \in C^\infty(\mathbb{R}^3)$ are solutions of (1) and (2), respectively. Making the transformations

$$u(x, t) = u(\nu) \quad \text{and} \quad v(x, y, t) = v(\eta) \quad (4)$$

with

$$\nu = x - \omega t, \quad \eta = kx + \alpha y + \omega' t, \quad (5)$$

where ω, ω', k and $\alpha \in \mathbb{R}$, the equations (1) and (2) can be rewritten as

$$H(u, u', u'', \dots) = 0, \quad (6)$$

where $u(\xi) = u(\nu)$ for equation (1) and $u(\xi) = v(\eta)$ for equation (2).

Assume that equation (6) is integrable with respect to ξ without cancelling the integrating constants. Then introduce the following transformation [10]

$$u(\xi) = \phi(\xi) + c_1, \quad c_1 \in \mathbb{R}. \quad (7)$$

Substituting (7) into (6) and setting the constant part equal to zero in the resulting nonlinear ODE in ϕ and assuming that the function ϕ and its derivatives have the following asymptotic values:

$$\phi(\xi) \rightarrow \phi_\pm \quad \text{as} \quad \xi \rightarrow \pm\infty, \quad (8)$$

for $n \geq 1$

$$\phi^{(n)} \rightarrow 0 \quad \text{as} \quad \xi \rightarrow \pm\infty, \quad (9)$$

and that ϕ_{\pm} satisfies the algebraic equation in ϕ [10], we get the value of c_1 .

The (G'/G) -expanding method consists in setting

$$\phi(\xi) = \alpha_m \left(\frac{G'}{G} \right)^m + \alpha_{m-1} \left(\frac{G'}{G} \right)^{m-1} + \dots \quad (10)$$

where $\alpha_m \neq 0$ and $G = G(\xi)$ satisfies the second order linear ordinary differential equation (LODE) in the form [22, 19]:

$$G'' + \lambda G' + \mu G = 0. \quad (11)$$

The parameter m can be found by substituting along with equation (11) into equation (6) and considering a homogeneous balance between the highest order derivative and highest order nonlinear term in equation (6), where $k, \alpha, \omega, a_0, a_1, \dots, a_m$ are to be determined. Substituting (7) and (10) into (4) yields a set of algebraic equations for $k, \alpha, a_0, a_1, \dots, a_m$, vanishing all coefficients of $(G'/G)^i$. From these relations, $k, \alpha, \omega, a_0, a_1, \dots, a_m$ can be obtained. Having determined these parameters, knowing that m is a positive integer in most cases, and using (7) and (10) we obtain analytical solutions $u(x, t)$ and $v(x, y, t)$ of (1) and (2), respectively.

The solutions of the equation (11) are given as follows:

- If $\lambda^2 - 4\mu > 0$,

$$G(\xi) = e^{-\frac{\lambda}{2}\xi} \left(k_1 \cosh \left(\frac{\sqrt{\lambda^2 - 4\mu}}{2} \xi \right) + k_2 \sinh \left(\frac{\sqrt{\lambda^2 - 4\mu}}{2} \xi \right) \right) \quad (12)$$

and

$$\frac{G'}{G} = -\frac{\lambda}{2} + \frac{\sqrt{\lambda^2 - 4\mu}}{2} f(\xi; \lambda, \mu). \quad (13)$$

where

$$f(\xi; \lambda, \mu) = \frac{k_1 \sinh \left(\frac{\sqrt{\lambda^2 - 4\mu}}{2} \xi \right) + k_2 \cosh \left(\frac{\sqrt{\lambda^2 - 4\mu}}{2} \xi \right)}{k_1 \cosh \left(\frac{\sqrt{\lambda^2 - 4\mu}}{2} \xi \right) + k_2 \sinh \left(\frac{\sqrt{\lambda^2 - 4\mu}}{2} \xi \right)} \quad (14)$$

- If $\lambda^2 - 4\mu < 0$,

$$G(\xi) = e^{-\frac{\lambda}{2}\xi} \left(k_1 \cos \left(\frac{\sqrt{4\mu - \lambda^2}}{2} \xi \right) + k_2 \sin \left(\frac{\sqrt{4\mu - \lambda^2}}{2} \xi \right) \right) \quad (15)$$

and

$$\frac{G'}{G} = -\frac{\lambda}{2} + \frac{\sqrt{4\mu - \lambda^2}}{2} g(\xi; \lambda, \mu). \quad (16)$$

where

$$g(\xi; \lambda, \mu) = \frac{-k_1 \sin \left(\frac{\sqrt{4\mu - \lambda^2}}{2} \xi \right) + k_2 \cos \left(\frac{\sqrt{4\mu - \lambda^2}}{2} \xi \right)}{k_1 \cos \left(\frac{\sqrt{4\mu - \lambda^2}}{2} \xi \right) + k_2 \sin \left(\frac{\sqrt{4\mu - \lambda^2}}{2} \xi \right)}, \quad (17)$$

- If $\lambda^2 - 4\mu = 0$,

$$G(\xi) = (k_1\xi + k_2)e^{-\frac{\lambda}{2}\xi} \quad (18)$$

and

$$\frac{G'}{G} = -\frac{\lambda}{2} + \frac{k_1}{k_1\xi + k_2}. \quad (19)$$

The equation (10) can be re-expressed as

$$\phi(\xi) = \sum_{k=1}^m \alpha_k \left(\frac{G'(\xi)}{G(\xi)} \right)^k + \alpha_0, \quad \alpha_0 \in \mathbb{R}, \quad \alpha_k \in \mathbb{R} \quad \forall k > 0, \quad (20)$$

giving

$$\phi'(\xi) = -\sum_{k=1}^m k\alpha_k \left\{ \mu \left(\frac{G'}{G} \right)^{k-1} + \lambda \left(\frac{G'}{G} \right)^k + \left(\frac{G'}{G} \right)^{k+1} \right\} \quad (21)$$

and

$$\begin{aligned} \phi''(\xi) = & \sum_{k=1}^m k\alpha_k \left\{ \mu^2(k-1) \left(\frac{G'}{G} \right)^{k-2} + \mu\lambda(2k-1) \left(\frac{G'}{G} \right)^{k-1} \right. \\ & \left. + k(\lambda^2 + 2\mu) \left(\frac{G'}{G} \right)^k + \lambda(2k+1) \left(\frac{G'}{G} \right)^{k+1} + (k+1) \left(\frac{G'}{G} \right)^{k+2} \right\}. \end{aligned} \quad (22)$$

3 Application to NPDEs describing physical phenomena

In this section, we obtain new classes of solutions to some relevant NPDEs such as the Burgers equations [10], KdV equation [10] and Kadomtsev-Petviashvili equation (also called KP equation in the literature) [21]. In each case the most relevant graphical representations are shown with an appropriate choice of parameters.

3.1 Burgers partial differential equation

Let us consider the nonlinear Burgers equation in two-dimensional space-time given by the following relation :

$$u_t + \alpha uu_x + \beta u_{xx} = 0. \quad (23)$$

Using (4), the equation (23) takes the form

$$M(u, u', u'') = -\omega u' + \alpha uu' + \beta u'' = 0. \quad (24)$$

Equation (24) can be integrated to give the integral form

$$\int M(u, u', u'') d\xi = -\omega u + \frac{1}{2}\alpha u^2 + \beta u' + c_1 C = 0, \quad c_1 C \in \mathbb{R}. \quad (25)$$

Suppose that $u = \phi + c_1$. Then the equation (25) can be transformed into the form

$$\alpha c_1 \phi - \omega \phi + \frac{1}{2} \alpha \phi^2 + \beta \phi' + c_1 \left(\frac{1}{2} \alpha c_1 + C - \omega \right) = 0. \quad (26)$$

Using the equations (8) and (9), the functions ϕ_{\pm} satisfy the relation

$$\alpha c_1 \phi_{\pm} - \omega \phi_{\pm} + \frac{1}{2} \alpha \phi_{\pm}^2 = 0, \quad (27)$$

with the condition

$$c_1 \left(\frac{1}{2} \alpha c_1 + C - \omega \right) = 0. \quad (28)$$

Two cases, deduced from the equation (28), deserve investigation in order to find explicit solutions to the equation (23).

Proposition 3.1 (Case 1) $c_1 = 0$ leads to the two following situations:

(i) $\xi = x - (2C - 2\beta)t$: We have

$$U_{11} = \begin{cases} \frac{2\beta}{\alpha} + \frac{2\beta}{\alpha} \left[-\frac{\lambda}{2} + \frac{\sqrt{\lambda^2 - 4\mu}}{2} f(\xi; \lambda, \mu) \right], & \text{if } \lambda^2 - 4\mu > 0, \\ \frac{2\beta}{\alpha} + \frac{2\beta}{\alpha} \left[-\frac{\lambda}{2} + \frac{\sqrt{4\mu - \lambda^2}}{2} g(\xi; \lambda, \mu) \right], & \text{if } \lambda^2 - 4\mu < 0 \\ \frac{2\beta}{\alpha} + \frac{2\beta}{\alpha} \left(-\frac{\lambda}{2} + \frac{k_1}{k_1 \xi + k_2} \right), & \text{if } \lambda^2 - 4\mu = 0, \end{cases} \quad (29)$$

(ii) $\xi = x - (2C - \beta\mu)t$: We get

$$U_{12} = \begin{cases} \frac{2\beta\mu}{\alpha} + \frac{2\beta}{\alpha} \left[-\frac{\lambda}{2} + \frac{\sqrt{\lambda^2 - 4\mu}}{2} f(\xi; \lambda, \mu) \right], & \text{if } \lambda^2 - 4\mu > 0, \\ \frac{2\beta\mu}{\alpha} + \frac{2\beta}{\alpha} \left[-\frac{\lambda}{2} + \frac{\sqrt{4\mu - \lambda^2}}{2} g(\xi; \lambda, \mu) \right], & \text{if } \lambda^2 - 4\mu < 0 \\ \frac{2\beta\mu}{\alpha} + \frac{2\beta}{\alpha} \left[-\frac{\lambda}{2} + \frac{k_1}{k_1 \xi + k_2} \right], & \text{if } \lambda^2 - 4\mu = 0, \end{cases} \quad (30)$$

Proof of relations (29) and (30).

Equation (26) becomes

$$-\omega \phi + \frac{1}{2} \alpha \phi^2 + \beta \phi' = 0. \quad (31)$$

Using (10) and balancing ϕ^2 into ϕ' , we get

$$2m = m + 1 \quad \text{so,} \quad m = 1. \quad (32)$$

Finally

$$\phi(\xi) = \alpha_1 \left(\frac{G'}{G} \right) + \alpha_0. \quad (33)$$




Figure 1: Solution U_{11} for $\alpha = 2, \beta = 1, \mu = 4, c_1 = 2, c_2 = 2, C = 2$




Figure 2: Solution U_{12} for $\alpha = 2, \beta = 1, \mu = 5, c_1 = 2, c_2 = 4, C = 2$

Substituting (33) and (21) (for $m = 1$) in (31), we obtain the following expression

$$\begin{aligned}
 & -\omega\alpha_1\left(\frac{G'}{G}\right) - \omega\alpha_0 + \frac{1}{2}\alpha\alpha_0^2 + \alpha\alpha_0\alpha_1\left(\frac{G'}{G}\right) + \frac{1}{2}\alpha\alpha_1^2\left(\frac{G'}{G}\right)^2 \\
 & \quad + \beta\alpha_0 - \beta\alpha_1\mu - \beta\alpha_1\mu\left(\frac{G'}{G}\right) - \beta\alpha_1\left(\frac{G'}{G}\right)^2 = 0.
 \end{aligned} \tag{34}$$

Vanishing the polynomial coefficients yields the system

$$\begin{cases}
 -\omega\alpha_0 + \frac{1}{2}\alpha\alpha_0^2 + \beta\alpha_0 - \beta\alpha_1\mu & = 0 \\
 -\omega\alpha_1 + \alpha\alpha_0\alpha_1 - \beta\alpha_1\mu & = 0 \\
 \frac{1}{2}\alpha\alpha_1^2 - \beta\alpha_1 & = 0
 \end{cases} \tag{35}$$

whose solutions are given by

$$\begin{cases}
 \omega & = 2C - 2\beta \\
 \alpha_0 & = \frac{2\beta}{\alpha} \\
 \alpha_1 & = \frac{2\beta}{\alpha},
 \end{cases} \quad \text{or} \quad \begin{cases}
 \omega & = 2C - \beta\mu \\
 \alpha_0 & = \frac{2\beta\mu}{\alpha} \\
 \alpha_1 & = \frac{2\beta}{\alpha}.
 \end{cases} \tag{36}$$

The relations (29) and (30) are therefore well satisfied. ■

Proposition 3.2 (Case 2) $c_1 = \frac{2}{\alpha}(\omega - C)$ furnishes two situations:

(i) $\xi = x - (2C - 2\beta + \beta\mu)t$

$$U_{13} = \begin{cases}
 \frac{-2\beta - \beta\lambda + 2C + 2\beta\mu}{\alpha} + \frac{\beta\sqrt{\lambda^2 - 4\mu}}{\alpha} f(\xi; \lambda, \mu), & \text{if } \lambda^2 - 4\mu > 0, \\
 \frac{-2\beta - \beta\lambda + 2C + 2\beta\mu}{\alpha} + \frac{\beta\sqrt{4\mu - \lambda^2}}{\alpha} g(\xi; \lambda, \mu), & \text{if } \lambda^2 - 4\mu < 0, \\
 \frac{-2\beta + 2C - \beta\lambda + 2\beta\mu}{\alpha} + \frac{2\beta}{\alpha} \left(\frac{k_1}{k_1\xi + k_2} \right), & \text{if } \lambda^2 - 4\mu = 0.
 \end{cases} \tag{37}$$

(ii) $\xi = x - (2C - \beta\mu)t$:

$$U_{14} = \begin{cases} \frac{-\beta\lambda+2C}{\alpha} + \frac{\beta\sqrt{\lambda^2-4\mu}}{\alpha} f(\xi; \lambda, \mu), & \text{if } \lambda^2 - 4\mu > 0, \\ \frac{-\beta\lambda+2C}{\alpha} + \frac{\beta\sqrt{4\mu-\lambda^2}}{\alpha} g(\xi; \lambda, \mu), & \text{if } \lambda^2 - 4\mu < 0, \\ \frac{-\beta\lambda+2C}{\alpha} + \frac{2\beta}{\alpha} \left(\frac{k_1}{k_1\xi+k_2} \right) & \text{if } \lambda^2 - 4\mu = 0. \end{cases} \quad (38)$$




Figure 3: Solution U_{13} for $\alpha = 2$, $\beta = 1$, $\lambda = 5$, $\mu = 4$, $c_1 = 2$, $c_2 = 4$, $C = 2$




Figure 4: Solution U_{14} for $\alpha = 2$, $\beta = 1$, $\lambda = 4$, $\mu = 5$, $c_1 = 2$, $c_2 = 4$, $C = 2$

Proof of relations (37) and (38).

The equation (26) becomes

$$\omega\phi - 2C\phi + \frac{1}{2}\alpha\phi^2 + \beta\phi' = 0. \quad (39)$$

Applying (G'/G) -expansion method on (39), using (10) and balancing ϕ^2 into ϕ' , we get $m = 1$ and

$$\phi(\xi) = \alpha_1 \left(\frac{G'}{G} \right) + \alpha_0. \quad (40)$$

Using (40), the relation (39) is re-expressed as a polynomial in (G'/G) and we obtain

$$\begin{aligned} & \omega\alpha_1 \left(\frac{G'}{G} \right) + \omega\alpha_0 - 2C\alpha_1 \left(\frac{G'}{G} \right) - 2C\alpha_0 + \frac{1}{2}\alpha\alpha_0^2 + \alpha\alpha_0\alpha_1 \left(\frac{G'}{G} \right) \\ & + \frac{1}{2}\alpha\alpha_1^2 \left(\frac{G'}{G} \right)^2 + \beta\alpha_0 - \beta\alpha_1\mu - \beta\alpha_1\mu \left(\frac{G'}{G} \right)^2 - \beta\alpha_1 \left(\frac{G'}{G} \right)^2 = 0. \end{aligned} \quad (41)$$

Vanishing the polynomial coefficients provides the system

$$\begin{cases} \omega\alpha_0 - 2C\alpha_0 + \frac{1}{2}\alpha\alpha_0^2 + \beta\alpha_0 - \beta\alpha_1\mu & = 0 \\ \omega\alpha_1 - 2C\alpha_1 + \alpha\alpha_0\alpha_1 - \beta\alpha_1\mu & = 0 \\ \frac{1}{2}\alpha\alpha_1^2 - \beta\alpha_1 & = 0 \end{cases} \quad (42)$$

which can be solved to give the solutions

$$\begin{cases} \omega & = 2C - 2\beta + \beta\mu \\ \alpha_0 & = \frac{2\beta}{\alpha} \\ \alpha_1 & = \frac{2\beta}{\alpha}, \end{cases} \quad \text{or} \quad \begin{cases} \omega & = 2C - \beta\mu \\ \alpha_0 & = \frac{2\beta\mu}{\alpha} \\ \alpha_1 & = \frac{2\beta}{\alpha}. \end{cases} \quad (43)$$

Similarly as in the previous case, the relations (37) and (38) are well satisfied. ■

3.2 KdV nonlinear equation

We consider the following KdV equation:

$$u_t + \alpha u u_x + \gamma u_{xxx} = 0. \quad (44)$$

To solve it we use the transformation (4) to reduce the equation (44) into an ordinary differential equation (ODE):

$$-\omega u + \frac{1}{2}\alpha u^2 + \gamma u'' + c_1 C = 0, \quad (45)$$

where $c_1 C$ is constant. Combining (7) and (45), we get

$$(\alpha c_1 - \omega)\phi + \frac{1}{2}\alpha\phi^2 + \gamma\phi'' + c_1\left(\frac{1}{2}\alpha c_1 + C - \omega\right) = 0. \quad (46)$$

The solutions of (44) are also given in two different cases.

Proposition 3.3 (Case 1) $c_1 = 0$ leads to two situations:

(i) If $\xi = x - \gamma(4\mu - \lambda^2)t$, we get

$$U_{21} = \begin{cases} \frac{-4\gamma\mu + \gamma\lambda^2}{\alpha} - \frac{3\gamma(\lambda^2 - 4\mu)}{\alpha} f^2(\xi; \lambda; \mu), & \text{if } \lambda^2 - 4\mu > 0, \\ \frac{-4\gamma\mu + \gamma\lambda^2}{\alpha} - \frac{3\gamma(\lambda^2 - 4\mu)}{\alpha} g^2(\xi; \lambda; \mu), & \text{if } \lambda^2 - 4\mu < 0, \\ \frac{-4\gamma\mu + 6\gamma\mu\lambda - 5\gamma\lambda^2}{\alpha} + \frac{12\gamma(\lambda - \mu)}{\alpha} \left(\frac{k_1}{k_1\xi + k_2}\right) - \frac{12\gamma}{\alpha} \left(\frac{k_1}{k_1\xi + k_2}\right)^2, & \text{if } \lambda^2 - 4\mu = 0. \end{cases} \quad (47)$$

(ii) If $\xi = x + \gamma(4\mu - \lambda^2)t$, we have:

$$U_{22} = \begin{cases} \frac{6\gamma(\lambda - \mu)\sqrt{\lambda^2 - 4\mu}}{\alpha} f(\xi; \lambda; \mu) - \frac{3\gamma(\lambda^2 - 4\mu)}{\alpha} f^2(\xi; \lambda; \mu) \\ + \frac{-12\gamma\mu + 6\gamma\mu\lambda - 3\gamma\lambda^2}{\alpha}, & \text{if } \lambda^2 - 4\mu > 0, \\ \frac{6\gamma(\lambda - \mu)\sqrt{\lambda^2 - 4\mu}}{\alpha} g(\xi; \lambda; \mu) - \frac{3\gamma(\lambda^2 - 4\mu)}{\alpha} g^2(\xi; \lambda; \mu) \\ + \frac{-12\gamma\mu + 6\gamma\mu\lambda - 3\gamma\lambda^2}{\alpha}, & \text{if } \lambda^2 - 4\mu < 0, \\ \frac{-12\gamma\mu + 6\gamma\mu\lambda - 3\gamma\lambda^2}{\alpha} + \frac{12\gamma(\lambda - \mu)}{\alpha} \left(\frac{k_1}{k_1\xi + k_2}\right) \\ - \frac{12\gamma}{\alpha} \left(\frac{k_1}{k_1\xi + k_2}\right)^2, & \text{if } \lambda^2 - 4\mu = 0. \end{cases} \quad (48)$$

Proof of relations (47) and (48).

The equation (46) is reduced to

$$-\omega\phi + \frac{1}{2}\alpha\phi^2 + \gamma\phi'' = 0. \quad (49)$$

Taking into account (10), and balancing ϕ^2 into ϕ'' , we get $m = 2$ and arrive at the following system:

$$\begin{cases} -\omega\alpha_0 + \frac{1}{2}\alpha\alpha_0^2 + 2\gamma\alpha_2\mu^2 + \gamma\alpha_1\lambda\mu & = 0 \\ -\omega\alpha_1 + \alpha\alpha_0\alpha_1 + 6\gamma\alpha_2\lambda\mu + 2\gamma\alpha_1\mu + \alpha_1\mu + \alpha_1\lambda^2\gamma & = 0 \\ -\omega\alpha_2 + \alpha\alpha_0\alpha_2 + \frac{1}{2}\alpha\alpha_1^2 + 4\gamma\alpha_2\lambda^2 + 3\alpha_1\gamma\lambda + 8\alpha_2\mu\gamma & = 0 \\ \alpha\alpha_1\alpha_2 + 10\gamma\alpha_2\lambda + 2\alpha_1\gamma & = 0 \\ \frac{1}{2}\alpha\alpha_2 + 6\alpha_2\gamma & = 0. \end{cases} \quad (50)$$




Figure 5: Solution U_{21} for $\alpha = 2$, $\gamma = 1$, $\lambda = 5$, $\mu = 4$, $c_1 = 2$, $c_2 = 4$, $C = 5$




Figure 6: Solution of U_{22} for $\alpha = 2$, $\gamma = 1$, $\lambda = 4$, $\mu = 5$, $c_1 = 2$, $c_2 = 2$, $C = 4$

which can be solved to give

$$\begin{cases} \omega = \gamma(4\mu - \lambda^2) \\ \alpha_0 = \frac{-2\gamma(2\mu + \lambda^2)}{\alpha} \\ \alpha_1 = -\frac{12\gamma\lambda}{\alpha} \\ \alpha_2 = -\frac{12\gamma}{\alpha} \end{cases} \quad \text{or} \quad \begin{cases} \omega = -\gamma(4\mu - \lambda^2) \\ \alpha_0 = \frac{-12\gamma\mu}{\alpha} \\ \alpha_1 = -\frac{12\gamma\lambda}{\alpha} \\ \alpha_2 = -\frac{12\gamma}{\alpha} \end{cases} \quad (51)$$

The relations (47) and (48) then hold. ■

Proposition 3.4 (Case 2) $c_1 = \frac{2\omega - 2C}{\alpha}$ amounts to the solutions of (44) in the two following situations:

(i) $\xi = x - (2C + 4\gamma\mu - \lambda^2\gamma)t$ gives

$$U_{23} = \begin{cases} \frac{-4\gamma\mu + \gamma\lambda^2 + 2C}{\alpha} - \frac{3\gamma(\lambda^2 - 4\mu)}{\alpha} f^2(\xi; \lambda; \mu), & \text{if } \lambda^2 - 4\mu > 0, \\ \frac{-4\gamma\mu + \gamma\lambda^2 + 2C}{\alpha} - \frac{3\gamma(\lambda^2 - 4\mu)}{\alpha} g^2(\xi; \lambda; \mu), & \text{if } \lambda^2 - 4\mu < 0 \text{ and} \\ \frac{-4\gamma\mu + \gamma\lambda^2 + 2C}{\alpha} - \frac{12\gamma}{\alpha} \left(\frac{k_1}{k_1\xi + k_2} \right)^2, & \text{if } \lambda^2 - 4\mu = 0. \end{cases} \quad (52)$$

(ii) $\xi = x - (2C + 4\gamma\mu + \lambda^2\gamma)t$ yields

$$U_{24} = \begin{cases} \frac{-12\gamma\mu + 3\gamma\lambda^2 + 2C}{\alpha} - \frac{3\gamma(\lambda^2 - 4\mu)}{\alpha} f^2(\xi; \lambda; \mu), & \text{if } \lambda^2 - 4\mu > 0, \\ \frac{-12\gamma\mu + 3\gamma\lambda^2 + 2C}{\alpha} - \frac{3\gamma(\lambda^2 - 4\mu)}{\alpha} g^2(\xi; \lambda; \mu), & \text{if } \lambda^2 - 4\mu < 0, \\ \frac{-12\gamma\mu + 3\gamma\lambda^2 + 2C}{\alpha} - \frac{12\gamma}{\alpha} \left(\frac{k_1}{k_1\xi + k_2} \right)^2, & \text{if } \lambda^2 - 4\mu = 0. \end{cases} \quad (53)$$

Proof of relations (52) and (53).

The equation (46) can be re-expressed as

$$(\omega - 2C)\phi + \frac{1}{2}\alpha\phi^2 + \gamma\phi'' = 0. \quad (54)$$




Figure 7: Solution of U_{23} for $\alpha = 2$, $\gamma = 5$, $\lambda = 1$, $\mu = 1/4$, $c_1 = 2$, $c_2 = 4$, $C = 1$, $\beta = 1$




Figure 8: Soliton solution of U_{24} for $\alpha = 1/2$, $\gamma = 1$, $\lambda = 2$, $\mu = 1/2$, $c_1 = 1$, $c_2 = 0.3$, $C = 0.1$

Using the equation (10) and balancing ϕ^2 into ϕ'' imply that $m = 2$ and we arrive at the following system:

$$\begin{cases} \omega\alpha_0 - 2C\alpha_0 + \frac{1}{2}\alpha\alpha_0^2 + 2\gamma\alpha_2\mu^2 + \gamma\alpha_1\lambda\mu^2 & = 0 \\ \omega\alpha_1 - 2C\alpha_1 + \alpha\alpha_0\alpha_1 + 6\gamma\alpha_2\lambda\mu + 2\gamma\alpha_1\mu + \alpha_1\mu + \alpha_1\lambda^2\gamma & = 0 \\ \omega\alpha_2 - 2C\alpha_2 + \alpha\alpha_0\alpha_2 + \frac{1}{2}\alpha\alpha_1^2 + 4\gamma\alpha_2\lambda^2 + 3\alpha_1\gamma\lambda + 8\alpha_2\mu\gamma & = 0 \\ \alpha\alpha_1\alpha_2 + 10\gamma\alpha_2\lambda + 2\alpha_1\gamma & = 0 \\ \frac{1}{2}\alpha\alpha_2^2 + 6\alpha_2\gamma & = 0 \end{cases} \quad (55)$$

whose solutions are given by

$$\begin{cases} \omega & = 2C + 4\gamma\mu - \lambda^2\gamma \\ \alpha_0 & = -\frac{12\gamma\mu}{\alpha} \\ \alpha_1 & = -\frac{12\gamma\lambda}{\alpha} \\ \alpha_2 & = -\frac{12\gamma}{\alpha} \end{cases} \quad \text{or} \quad \begin{cases} \omega & = 2C - 4\gamma\mu + \lambda^2\gamma \\ \alpha_0 & = \frac{2\gamma(2\mu + \lambda^2)}{\alpha} \\ \alpha_1 & = -\frac{12\gamma\lambda}{\alpha} \\ \alpha_2 & = -\frac{12\gamma}{\alpha} \end{cases} \quad (56)$$

Therefore we simply get relations (52) and (53). ■

3.3 Kadomtsev-Petviashvili nonlinear equation and its solutions

This subsection is devoted to the resolution of the Kadomtsev-Petviashvili NPDE in three dimensions given by

$$(u_t + 6uu_x + u_{xxx})_x + 3\sigma^2 u_{yy} = 0, \quad \sigma^2 = \pm 1. \quad (57)$$

Using the transformation (4), the equation (57) takes the form

$$N(u, u', u'', u''') = k\omega u'' + 6k^2(u'u)' + k^4 u^{(4)} + 3\sigma^2 \alpha^2 u'' = 0. \quad (58)$$

Integrating this latter equation, we get the following nonlinear ODE

$$\int N(u, u', u'', u''') d\xi = (k\omega + 3\sigma^2 \alpha^2)u' + 6k^2 u'u + k^4 u''' + c_1 C = 0, \quad (59)$$

where c_1C is a constant to be determined. Substituting (7) into (59), we simply get

$$(k\omega + 3\sigma^2\alpha^2 + 6c_1k^2)\phi + 3k^2\phi^2 + k^4\phi'' + c_1(k\omega + 3\sigma^2\alpha^2 + 3k^2c_1 + C) = 0. \quad (60)$$

The equation (57) can be now solved in different cases to give the next results:

Proposition 3.5 (Case 1) $c_1 = 0$. *The solutions of (57) are:*

(i) *If $\xi = x - (4k^4\mu - 3\sigma^2\alpha^2 - k^4\lambda^2)t/k$:*

$$U_{31} = \begin{cases} -2k^2\mu + \frac{1}{2}k^2\lambda^2 - \frac{k^2(\lambda^2-4\mu)}{2}f(\xi; \lambda, \mu)^2, & \text{if } \lambda^2 - 4\mu > 0, \\ -2k^2\mu + \frac{1}{2}k^2\lambda^2 - \frac{k^2(4\mu-\lambda^2)}{2}g(\xi; \lambda, \mu)^2, & \text{if } \lambda^2 - 4\mu < 0, \\ -2k^2\mu + \frac{1}{2}k^2\lambda^2 - 2k^2\left(\frac{c_1}{c_1\xi+C_2}\right)^2, & \text{if } \lambda^2 - 4\mu = 0. \end{cases} \quad (61)$$

(ii) *If $\xi = x - (4k^4\mu - 3\sigma^2\alpha^2 - k^4\lambda^2)t/k$:*

$$U_{32} = \begin{cases} -\frac{2}{3}k^2\mu + \frac{1}{6}k^2\lambda^2 - \frac{k^2(\lambda^2-4\mu)}{2}f(\xi; \lambda, \mu)^2, & \text{if } \lambda^2 - 4\mu > 0, \\ -\frac{2}{3}k^2\mu + \frac{1}{6}k^2\lambda^2 - \frac{k^2(4\mu-\lambda^2)}{2}g(\xi; \lambda, \mu)^2, & \text{if } \lambda^2 - 4\mu < 0, \\ -\frac{2}{3}k^2\mu + \frac{1}{6}k^2\lambda^2 - 2k^2\left(\frac{c_1}{c_1\xi+C_2}\right)^2, & \text{if } \lambda^2 - 4\mu = 0. \end{cases} \quad (62)$$

Proof of relations (61) and (62).

The equation (60) takes the form

$$(k\omega + 3\sigma^2\alpha^2)\phi + 3k^2\phi^2 + k^4\phi'' = 0. \quad (63)$$

Let us consider (10) and balance ϕ^2 into ϕ'' to get $m = 2$. Then we arrive at the following system

$$\begin{cases} 3k^2\alpha_2^2 + 6k^4\alpha_2 & = 0 \\ 6k^4\alpha\alpha_2 + 10k^4\alpha_2\lambda + 2k^4\alpha_1 & = 0 \\ 3\sigma^2\alpha^2\alpha_2 + k\omega\alpha_2 + 3k^2\alpha_1^2 + 3k^4\alpha_1\lambda + 8k^4\alpha_2\mu + 6k^2\alpha_0\alpha_1\alpha_2 + 4k^4\alpha_2\lambda^2 & = 0 \\ k\omega\alpha_1 + 2k^4\alpha_1\mu + \alpha k^2\alpha_0\alpha_1 + 3\sigma^2\alpha^2\alpha_1 + 6k^4\alpha_2\lambda\mu + k^4\alpha_1\lambda^2 & = 0 \\ k\omega\alpha_0 + k^4\alpha_1\lambda\mu + 3k^2\alpha_0^2 + 3\sigma^2\alpha^2\alpha_0 + 2k^4\alpha_2\mu^2 & = 0 \end{cases} \quad (64)$$

giving the solutions

$$\begin{cases} \omega = \frac{4k^4\mu - 3\sigma^2\alpha^2 - k^4\lambda^2}{k} \\ \alpha_0 = -2k^2\mu \\ \alpha_1 = -2k^2\mu \\ \alpha_2 = -2k^2 \end{cases} \quad \text{or} \quad \begin{cases} \omega = -\frac{4k^4\mu + 3\sigma^2\alpha^2 - k^4\lambda^2}{k} \\ \alpha_0 = -\frac{2}{3}k^2\mu - \frac{1}{3}k^2\lambda^2 \\ \alpha_1 = -2k^2\lambda \\ \alpha_2 = -2k^2. \end{cases} \quad (65)$$

There result the relations (61) and (62). ■

Proposition 3.6 (Case 2) $c_1 = \frac{-C-3\sigma^2\alpha^2-k^2\omega}{3}$. *It leads to the solutions of (57) in two situations:*

(i) $\xi = kx + \alpha y - (4k^4\mu - 2k^2C - k^4\lambda^2 + 6k^2\sigma^2\alpha^2 - 3\sigma^2\alpha^2)t/k(-1 + 2k^3)$ provides

$$U_{33} = \begin{cases} \frac{-12k^2\mu + 3k^2\lambda^2 - 2C - 6\sigma^2\alpha^2 - 2k^2\omega}{6} - \frac{k^2(\lambda^2 - 4\mu)}{2} f(\xi; \lambda, \mu)^2, \\ \text{if } \lambda^2 - 4\mu > 0, \\ \frac{-12k^2\mu + 3k^2\lambda^2 - 2C - 6\sigma^2\alpha^2 - 2k^2\omega}{6} - \frac{k^2(4\mu - \lambda^2)}{2} g(\xi; \lambda, \mu)^2 \\ \text{if } \lambda^2 - 4\mu < 0, \\ \frac{-12k^2\mu + 3k^2\lambda^2 - 2C - 6\sigma^2\alpha^2 - 2k^2\omega}{6} - 2k^2 \left(\frac{c_1}{c_1\xi + C_2} \right)^2 \\ \text{if } \lambda^2 - 4\mu = 0 \end{cases} \quad (66)$$

(ii) $\xi = kx + \alpha y - (-4k^4\mu + 2k^2C + k^4\lambda^2 + 6k^2\sigma^2\alpha^2 - 3\sigma^2\alpha^2)t/k(-1 + 2k^3)$ gives

$$U_{34} = \begin{cases} \frac{-4k^2\mu + k^2\lambda^2 - 2C - 6\sigma^2\alpha^2 - 2k^2\omega}{6} - \frac{k^2(\lambda^2 - 4\mu)}{2} f(\xi; \lambda, \mu)^2, \\ \text{if } \lambda^2 - 4\mu > 0, \\ \frac{-4k^2\mu + k^2\lambda^2 - 2C - 6\sigma^2\alpha^2 - 2k^2\omega}{6} - \frac{k^2(4\mu - \lambda^2)}{2} g(\xi; \lambda, \mu)^2, \\ \text{if } \lambda^2 - 4\mu < 0 \\ \frac{-4k^2\mu + k^2\lambda^2 - 2C - 6\sigma^2\alpha^2 - 2k^2\omega}{6} - 2k^2 \left(\frac{c_1}{c_1\xi + C_2} \right)^2 \\ \text{if } \lambda^2 - 4\mu = 0. \end{cases} \quad (67)$$

Proof of (66) and (67).

The equation (60) can be rewritten as

$$(k\omega + 3\sigma^2\alpha^2 - 2k^2C - 6k^2\sigma^2\alpha^2 - 2k^4\omega)\phi + 3k^2\phi^2 + k^4\phi'' = 0. \quad (68)$$

Considering (10) and balancing ϕ^2 into ϕ'' , we get $m = 2$ and obtain the following system

$$\begin{cases} 3k^2C_2^2 + 6k^4\alpha_2 & = 0 \\ 6k^2\alpha_1\alpha_2 + 10k^4\alpha_2\lambda + 2k^4\alpha_1 & = 0 \\ -6k^2\sigma^2\alpha^2\alpha_2 - 2k^2\omega\alpha_2 + 6k^2\alpha_0\alpha_2 - 2k^2C\alpha_2 + 3k^4\alpha_1\lambda \\ + 3k^2\alpha_1^2 + k\omega\alpha_2 + 8k^4\alpha_2\mu + 3\sigma^2\alpha^2\alpha_2 + 4k^4\alpha_2\lambda^2 & = 0 \\ 2k^4\alpha_1\mu - 2k^4\omega\alpha_1 + 2k^4\alpha_1\lambda^2 + k\omega\alpha_1 - 2k^2C\alpha_1 + 6k^4\alpha_2 \\ - 6k^2\sigma^2\alpha^2\alpha_1 + 6k^2\alpha_0\alpha_1 + 3\sigma^2\alpha^2\alpha_1 & = 0 \\ k\omega\alpha_0 - 2k^2C\alpha_0 - 6k^2\sigma^2\alpha^2\alpha_0 + 3k^2\alpha_0^2 \\ + 3\sigma^2\alpha^2\alpha_0 + 2k^4\alpha_2\mu^2 - 2k^4\omega\alpha_0 + k^4\alpha_1\lambda\mu & = 0 \end{cases} \quad (69)$$

Its solutions are given by

$$\begin{cases} \omega = -\frac{2k^2 + 6k^2\sigma^2\alpha^2 + 4k^4\mu - 3\sigma^2\alpha^2 - k^4\lambda^2}{k(-1 + 2k^3)} \\ \alpha_0 = -2k^2\mu \\ \alpha_1 = -2k^2\lambda \\ \alpha_2 = -2k^2 \end{cases} \quad (70)$$

or

$$\begin{cases} \omega = \frac{-4k^4\mu + 2k^2C + k^4\lambda^2 + 6k^2\sigma^2\alpha^2 - 3\sigma^2\alpha^2}{k(-1 + 2k^3)} \\ \alpha_0 = -\frac{2}{3}k^2\mu - \frac{1}{3}k^2\lambda^2 \\ \alpha_1 = -2k^2\lambda \\ \alpha_2 = -2k^2 \end{cases} \quad (71)$$

validating the relations (66) and (67). ■

4 Conclusion

In this work we have modified the (G'/G) -expansion method to find new classes of solutions to known important NPDEs, in addition to solutions obtained by the usual (G'/G) -expansion method. The most relevant graphical representations have been drawn to show the pertinence of the obtained analytical solutions.

Acknowledgment

RK thanks the University of Abomey-Calavi and the Ministry for High Education and Scientific Research of Benin for financial support. This work is partially supported by the ICTP through the OEA-ICMPA-Prj-15. The ICMPA is in partnership with the Daniel Iagolnitzer Foundation (DIF), France.

References

- [1] C. L. Zheng, Localized Coherent Structures with Chaotic and Fractal Behaviors in a $(2 + 1)$ -Dimensional Modified Dispersive Water-Wave System, *Communication in Theoretical Physics* Vol. **39** (2003), pp.261-266.
- [2] J. H. He, Nonlinear Science as a Fluctuating Research Frontier, *Chaos, Soliton and Fractals* **41** (2009), 2533-2537.
- [3] L. Debnath, *Partial Differential Equations for Scientist and Engineers*, (Balkema, Rotterdam, 2002).
- [4] J. H. He, G. C. Wu and F. Austin, The Variational Iteration Method Which Should be Followed, *Nonlinear Science Letters A* **1** (1) (2010), 1-30.
- [5] M.-Din, S. T., M. A. Noor and K. I. Noor, Variational Iteration Method for Solving Fierl-Petviashvili Equation Using He's Polynomials and Pade Approximates *World Applied Sciences Journal* **6**(9) (2009), 1298-1303.
- [6] M. J. Ablowitz and H. Segur, *Solitons and Inverse Scattering Transform*, (SIAM, Philadelphia, 1981).
- [7] R. Hirota, *The Direct Method in Soliton Theory*, (Cambridge University Press, Cambridge, 2004).
- [8] R. Conte, *Painlevé Property*, (Springer, Berlin, 1999).
- [9] W. Hereman, P. P. Banerjee, A. Korpel, G. Assanto, A. Van Immerzele and A. Meerpoel, Exact Solitary Wave Solutions of Nonlinear Evolution and Wave Equations Using a Direct Algebraic Method, *Journal of Physics A: Mathematical and General*, Vol. **9**, No. 5 (1986), pp.607-628 .

- [10] N. H. Abdel-All, M. A. Abdel-Razek and A. K. Seddeek, Expanding the Tanh-Function Method for Solving Nonlinear Equation, *Assiut University, Assiut, Egypt*, (2011).
- [11] W. Malfliet and W. Hereman, The Tanh Method: I. Exact Solutions of Nonlinear Evolution and Wave Equations, *Physica Scripta*, Vol. **54** (1996), pp. 563-568.
- [12] A. A. Soliman, The Modified Extended Tanh-Function Method for Solving Burgers-Type Equations, *Physica A: Statistical Mechanics and its Applications*, Vol. **361**, No.2 (2006), pp. 394-404.
- [13] C. T. Yan, A simple Transformation for Nonlinear Waves, *Physics Letters A* Vol. **224**, No.1-2 (1996), pp.77-84.
- [14] R. Guo, B. Tian, L. Wang, F. Qi and Y. Zhan, Darboux Transformation and Soliton Solutions for a System Describing Ultrashort Pulse Propagation in a Multicomponent Nonlinear Medium, *Physica Scripta*, **81**:025002 (2010).
- [15] Y. Zhao, N -Fold Darboux Transformation for a Nonlinear Evolution Equation, *Applied Mathematics*, Vol. **3** No.8 (2012), pp. 943-948.
- [16] İ. E. İnan, U. Yavuz, B. Kiliç, Traveling Wave Solutions of the RLW-Burgers Equation and Potential Kdv Equation by Using the (G'/G) -Expansion Method, *Journal of Arts and Sciences Sayt* **12** Aralık (2009).
- [17] A. Roozi. and A. G. Mahmeiani, The (G'/G) -Expansion Method for $(2+1)$ -Dimensional Kadomtsev-Petviashvili Equation, *World Applied Sciences Journal*, **13**(10) (2011), pp. 2231-2234.
- [18] E. Fan, Extended tanh-Function Method and Its Applications to Nonlinear Equations, *Physics Letters A*, Vol.**277** No.4-5 (2000), pp. 212-218.
- [19] H. Naher, F. A. Abdullah and M. A. Akbar, New Travelling Wave Solutions of the Higher Dimensional Nonlinear Evolution Equation by the Improved (G'/G) -Expansion Method, *World applied Sciences Journal* **161** (2012), pp. 11-21.
- [20] J. H. He, Variational Iteration Method-Some Recent Result and New Interpretations, *Journal of Computational and Applied Mathematics*, Vol. **207** No.1 (2007), pp. 3-17.
- [21] M. J. Ablowitz and P. A. Clarkson, *Solitons, Nonlinear Evolution Equations and Inverse Scattering*, (Cambridge: Cambridge University Press, 1991).
- [22] H. Naher, F. A. Abdullah and M. A. Akbar, The (G'/G) -Expansion Method for Abundant Travelling Wave Solutions of Caudrey-Dodd-Gibbon Equation *Math Prob. Eng.* (2011), 11pages.
- [23] S. A. El-Wakil, S. K. El-Labany, M. A. Zahran and R. Sabry, Modified Extended Tanh-Function Method for Solving Nonlinear Partial Differential Equations, *Physics Letters A*, **299** (2002), pp. 179-188.
- [24] M. Wang, X. Li and J. Zhang, The (G'/G) -Expansion Method and Travelling Wave Solutions of Nonlinear Evolution Equations in Mathematical Physics, *Phys. Lett. A*, **372** (2008), pp. 417-423.

- [25] Y. T. Gao and B. Tian, Generalized Tanh Method with Symbolic Computation and Generalized Shallow Water Wave Equation, *Computers Mathematics With Applications*, Vol. **33** No. 4 (1997), pp. 115-118.