

On the Leray-Hopf Extension Condition for the Steady-State Navier–Stokes Problem in Multiply-Connected Bounded Domains

Giovanni P. Galdi

November 9, 2018

Abstract

Employing the approach of A. Takeshita [Pacific J. Math., **157** (1993), 151–158], we give an elementary proof of the invalidity of the Leray-Hopf Extension Condition for certain multiply connected bounded domains of \mathbb{R}^n , $n = 2, 3$, whenever the flow through the different components of the boundary is non-zero. Our proof is alternative to and, to an extent, more direct than the recent one proposed by J.G. Heywood [J. Math. Fluid Mech. **13** (2011), 449–457].

Keywords. Navier–Stokes equations, Non-homogeneous boundary conditions, Multiply connected domains

MSC (2000) 35Q30, 35Q35, 30E25

1 Introduction

Let Ω be a bounded domain of \mathbb{R}^n , $n = 2, 3$. As is well known, the Leray-Hopf Extension Condition is related to the solvability of the following Navier–Stokes equations

$$\left. \begin{aligned} \nu \Delta \mathbf{v} &= \mathbf{v} \cdot \nabla \mathbf{v} + \nabla p + \mathbf{f} \\ \operatorname{div} \mathbf{v} &= 0 \end{aligned} \right\} \text{ in } \Omega, \quad (1)$$

under prescribed non-homogeneous boundary conditions

$$\mathbf{v} = \mathbf{v}_* \quad \text{at } \partial\Omega. \quad (2)$$

Here, as customary, \mathbf{v} , p and $\nu > 0$ denote velocity and pressure fields, and kinematic viscosity of the liquid, respectively, while \mathbf{f} is representative of a body force possibly acting on it. Moreover, \mathbf{v}_* is a given distribution of velocity at the boundary $\partial\Omega$, which, by $(1)_1$ and the Gauss theorem must satisfy the compatibility condition

$$\sum_{k=1}^N \int_{\Gamma_k} \mathbf{v}_* \cdot \mathbf{n} := \sum_{k=1}^N \Phi_k = 0, \quad (3)$$

where Γ_k , $k = 1, \dots, N$, are the connected components of $\partial\Omega$, and \mathbf{n} is its unit outer normal. From the physical viewpoint, Φ_k is the (mass) flow-rate through the portion Γ_k of the boundary. To fix the ideas, we assume that Γ_i , $i = 1, 2, \dots, N - 1$, are all surrounded by Γ_N and lie outside of each other.

The existence of a (weak, in principle) solution to the problem (1)–(3) is readily established (e.g., by Galerkin method or by Leray-Schauder theory), provided we are able to show (formally, at least) that the velocity field of the searched solution satisfies the a priori bound

$$\|\nabla \mathbf{v}\|_2 \leq C, \quad (4)$$

where $\|\cdot\|_2$ is the $L^2(\Omega)$ -norm,¹ and C , here and in the following, denotes a constant depending at most on Ω , \mathbf{f} , \mathbf{v}_* and ν ; see [7, Chapter IX] for details.

One way of attempting to prove (4) is to extend the boundary data \mathbf{v}_* to Ω by a solenoidal function \mathbf{V} , and introduce the new velocity field $\mathbf{u} := \mathbf{V} - \mathbf{v}$. Clearly, \mathbf{v} satisfies a bound of the type (4) if and only if \mathbf{u} does. Now, writing (1) in terms of \mathbf{u} , dot-multiplying both sides of the resulting equation by \mathbf{u} , integrating by parts over Ω and using the fact that \mathbf{u} is solenoidal and that vanishes at $\partial\Omega$, we formally show the following relation:

$$\nu \|\nabla \mathbf{u}\|_2^2 = -(\mathbf{u} \cdot \nabla \mathbf{V}, \mathbf{u}) - (\mathbf{V} \cdot \nabla \mathbf{V}, \mathbf{u}) - \nu(\nabla \mathbf{V}, \nabla \mathbf{u}) + (\mathbf{f}, \mathbf{u}), \quad (5)$$

where we have adopted the standard notation²

$$(\mathbf{a}, \mathbf{b}) = \int_{\Omega} a_i b_i, \quad \mathbf{a}, \mathbf{b} \in \mathbb{R}^n; \quad (\mathbf{A}, \mathbf{B}) = \int_{\Omega} A_{ij} B_{ij}, \quad \mathbf{A}, \mathbf{B} \in \mathbb{R}^{n^2}.$$

Thus, assuming $\mathbf{V} \in W^{1,2}(\Omega)$,³ and using Cauchy–Schwartz and classical embedding inequalities, from (5) we show

$$\nu \|\nabla \mathbf{u}\|_2^2 \leq -(\mathbf{u} \cdot \nabla \mathbf{V}, \mathbf{u}) + C. \quad (6)$$

Since both terms in (6) are quadratic in \mathbf{u} , from this relation it is not clear how to get a bound on $\nabla \mathbf{u}$ of the type (4), unless we make the obvious assumption that the viscosity ν is “sufficiently large” compared to the magnitude of $\nabla \mathbf{V}$, or equivalently, of \mathbf{v}_* in suitable trace norm. However, such a restriction can be avoided whenever \mathbf{v}_* obeys the *Leray–Hopf Extension Condition* [11, p. 38], [9, p. 772], namely, *for any $\varepsilon > 0$, there exists a solenoidal extension, $\mathbf{V}_\varepsilon \in W^{1,2}(\Omega)$,⁴ of \mathbf{v}_* such that*

$$-(\mathbf{u} \cdot \nabla \mathbf{V}_\varepsilon, \mathbf{u}) \leq \varepsilon \|\nabla \mathbf{u}\|_2^2, \quad (\text{EC})$$

for all solenoidal vector functions $\mathbf{u} \in W_0^{1,2}(\Omega)$.⁵ It is then obvious that the validity of (EC) along with (6) furnishes the desired uniform bound for \mathbf{u} , without imposing any restriction on the magnitude of $\nu > 0$.

¹We employ standard notation for Lebesgue, Sobolev and trace spaces; see e.g. [1].

²Summation convention over repeated indices applies.

³This condition on \mathbf{V} is certainly satisfied if $\mathbf{v}_* \in W^{1/2,2}(\partial\Omega)$ and Ω is Lipschitz.

⁴See footnote 3.

⁵Notice that (EC) is *weaker* than the so-called “Leray Inequality”, the latter consisting in replacing the left-hand side of (EC) with $|(\mathbf{u} \cdot \nabla \mathbf{V}_\varepsilon, \mathbf{u})|$. The validity of Leray’s Inequality is originally studied, and disproved under certain conditions, in [15] and, more recently, in [3].

The validity of (EC) has been investigated by many authors, beginning with the cited pioneering works of J. Leray and E. Hopf, who showed that (EC) certainly holds provided \mathbf{v}_* satisfies a condition *stronger* than (3), namely, that $\Phi_k = 0$, for each $k = 1, \dots, N$. More recently, a proof of (EC) under the *general* assumption (3) was given in the two-dimensional case by L.I. Sazonov [14], and, independently, by H. Fujita [5], provided, however Ω , \mathbf{v}_* , and \mathbf{u} satisfy suitable *symmetry* hypotheses; see also [12, 4]. As a result, existence to problem (1)–(3) follows on condition that also \mathbf{f} is prescribed in an appropriate class of symmetric functions. The method of Fujita was successively extended by V.V. Pukhnachev [13] to cover the three-dimensional case, again under appropriate symmetry assumptions.

The fact that (EC) may not be true unless some restrictions are imposed, was already clear after the work of A. Takeshita [15, Section 3] and the present author [6, pp. 22–23], where it was shown that even in the simplest case when Ω is an annulus \mathcal{A} , (EC) fails in general. More precisely, denoting by Γ_2 and Γ_1 the outer and inner concentric circles bounding \mathcal{A} , one proves that (EC) cannot hold at least when the flow-rate, $\Phi := \Phi_2 = -\Phi_1$ through Γ_2 is *strictly negative* (inflow condition). A similar result remains valid also in the case when Ω is a spherical shell, as stated in [15, p. 157] and clearly worked out in [3].

The counterexamples mentioned above require $\Phi < 0$. The case $\Phi > 0$ (outflow condition) presented, presumably, more difficulty and, as a result, the question of whether (EC) holds under the latter assumption on the flow-rate remained apparently open for several years.⁶ Quite recently, in [8], J.G. Heywood finally provided very interesting ideas on how to show the invalidity of (EC) also for the case $\Phi > 0$. This is achieved by using appropriate functions \mathbf{u} in (EC), that he names “U-tube test functions.”

Objective of this note is to give a direct and elementary proof of the invalidity of (EC) when Ω is an annulus (see Section 2) or a spherical shell (see Section 3), and $\Phi > 0$. Our proof uses Takeshita’s approach –which allows us to replace in (EC) the extension \mathbf{V}_ε with its integral average over all possible rotations– in conjunction with an appropriate choice of the function \mathbf{u} .

It should be emphasized that once the result is established for these special domains, the invalidity of (EC) can be extended to more general domains, even multiply connected, whenever for each “interior” connected component Γ_i of $\partial\Omega$, $i = 1, 2, \dots, N - 1$, there is a circumference (spherical surface) completely contained in Ω , and that surrounds only Γ_i . Actually, combining the results of [3] with ours, no restrictions need to be imposed on the sign of Φ_k , provided, of course, (3) is satisfied. This generalization can be obtained by following exactly the same argument of [3, Corollary 1], and it is stated in Theorem 1 in Section 4.

We wish to end this introductory section with a final observation. A different way of proving the a priori estimate (4), again suggested by J. Leray [11, p. 28 and ff], is to use a contradiction argument. By this argument one shows that (4) is true (and so existence to (1)–(2) is proved under the general condition (3))

⁶In this regard, see [8, Section 8].

provided the following requirements on the pair (\mathbf{w}, π) (in a suitable function class) are incompatible

$$\left. \begin{aligned} \mathbf{w} \cdot \nabla \mathbf{w} &= \nabla \pi \\ \operatorname{div} \mathbf{w} &= 0 \end{aligned} \right\} \text{ in } \Omega, \quad (7)$$

$$\begin{aligned} \mathbf{w} &= \mathbf{0} \text{ on } \partial\Omega, \\ -(\mathbf{w} \cdot \nabla \mathbf{V}, \mathbf{w}) &= \nu. \end{aligned}$$

Here (\mathbf{w}, π) are limits (in appropriate topology) of certain normalized sequences of solutions to (1)–(2), while \mathbf{V} is a given extension of the boundary data \mathbf{v}_* . It is then interesting to notice that if (7) has a solution, then (EC) cannot be true. In fact, writing $\mathbf{V}_\varepsilon = \mathbf{V} + (\mathbf{V}_\varepsilon - \mathbf{V})$, from (7)_{1,4} we find

$$-(\mathbf{w} \cdot \nabla \mathbf{V}_\varepsilon, \mathbf{w}) = \nu + (\mathbf{w} \cdot \nabla \mathbf{w}, \mathbf{V}_\varepsilon - \mathbf{V}) = \nu + (\nabla \pi, \mathbf{V}_\varepsilon - \mathbf{V}) = \nu$$

where, in the last step, we have used that $\mathbf{V}_\varepsilon - \mathbf{V}$ is solenoidal and vanishes at $\partial\Omega$. Consequently, admitting (EC) would imply $\nu \leq \varepsilon \|\nabla \mathbf{w}\|_2^2$ for all $\varepsilon > 0$, namely, $\nu = 0$. These considerations suggest that the contradiction argument could be a weaker requirement than the validity of (EC), and that it might lead to the proof of the a priori estimate (4) under more general assumptions than those of symmetry requested by the use of (EC). This fact was already hinted by C.J. Amick [2], but only recently was it fully confirmed by M.V. Korobkov, K. Pileckas, and R. Russo [10] who showed that (7) are indeed incompatible when Ω is a doubly-connected, two-dimensional (Lipschitz) domain, under the sole assumption that the flow-rate satisfies the inflow condition.

2 The Case Ω an Annulus

We follow and specialize the approach of [15]. Let $\Omega := \{x \in \mathbb{R}^2 : R_1 < |x| < R_2\}$, $R_1 > 0$, $\Gamma_i := \{x \in \mathbb{R}^2 : |x| = R_i\}$, $i = 1, 2$. Moreover, set

$$\Phi := \int_{\Gamma_2} \mathbf{v}_* \cdot \mathbf{n} = - \int_{\Gamma_1} \mathbf{v}_* \cdot \mathbf{n},$$

and assume $\Phi > 0$. We want to show that the validity of (EC) then leads to a contradiction. For $\mathbf{x} \in \Omega$, we put

$$\mathbf{y} = \mathcal{R}_\varphi \cdot \mathbf{x} \quad (8)$$

with $\mathcal{R}_\varphi \in SO(2)$ rotation matrix of angle $\varphi \in [0, 2\pi]$, and define the average of \mathbf{V}_ε :

$$\mathcal{A}(\mathbf{V}_\varepsilon)(\mathbf{y}) := \frac{1}{2\pi} \int_0^{2\pi} \mathcal{R}_\varphi \cdot \mathbf{V}_\varepsilon(\mathcal{R}_\varphi^\top \cdot \mathbf{y}) d\varphi,$$

where $\mathbf{V}_\varepsilon \in W^{1,2}(\Omega)$ is a solenoidal extension of \mathbf{v}_* for which (EC) is supposed to hold, and $^\top$ denotes transpose. Taking into account the properties of \mathbf{V}_ε ,

and the proper orthogonality of \mathcal{R}_φ , one at once shows that

$$\begin{aligned} \operatorname{div} \mathcal{A}(\mathbf{V}_\varepsilon)(\mathbf{y}) &= 0 \quad \mathbf{y} \in \Omega, \\ \int_{\Gamma_2} \mathcal{A}(\mathbf{V}_\varepsilon) \cdot \mathbf{n} &= - \int_{\Gamma_1} \mathcal{A}(\mathbf{V}_\varepsilon) \cdot \mathbf{n} = \Phi. \end{aligned} \quad (9)$$

Furthermore, by construction, $\mathcal{A}(\mathbf{V}_\varepsilon)$ is invariant under rotation. Therefore, observing that, denoted by (r, θ) a system of polar coordinates with the origin at $x = 0$, the corresponding base vectors $\{\mathbf{e}_r, \mathbf{e}_\theta\}$ are both invariant, we infer

$$\mathcal{A}(\mathbf{V}_\varepsilon) = v_1(r)\mathbf{e}_r + v(r)\mathbf{e}_\theta.$$

However, $\mathcal{A}(\mathbf{V}_\varepsilon)$ must satisfy (9), so that we conclude

$$\mathcal{A}(\mathbf{V}_\varepsilon) = \frac{1}{2\pi} \frac{\Phi}{r} \mathbf{e}_r + v(r)\mathbf{e}_\theta. \quad (10)$$

It is now straightforward to prove that since \mathbf{V}_ε satisfies (EC), also $\mathcal{A}(\mathbf{V}_\varepsilon)$ does. In fact, following [15], by Fubini theorem and (8), for all solenoidal $\mathbf{u} \in W_0^{1,2}(\Omega)$ we have

$$\begin{aligned} \int_{\Omega} \mathbf{u} \cdot \nabla_y (\mathcal{A}(\mathbf{V}_\varepsilon)) \cdot \mathbf{u} \, dy &= \frac{1}{2\pi} \int_0^{2\pi} \left(\int_{\Omega} \mathbf{u} \cdot \nabla_y (\mathcal{R}_\varphi \cdot \mathbf{V}_\varepsilon (\mathcal{R}_\varphi^\top \cdot \mathbf{y})) \cdot \mathbf{u} \, dy \right) d\varphi \\ &= \frac{1}{2\pi} \int_0^{2\pi} \left(\int_{\Omega} (\mathcal{R}_\varphi^\top \cdot \mathbf{u}) \cdot \nabla_x \mathbf{V}_\varepsilon \cdot (\mathcal{R}_\varphi^\top \cdot \mathbf{u}) \, dx \right) d\varphi \end{aligned} \quad (11)$$

Clearly, $\mathcal{R}_\varphi^\top \cdot \mathbf{u} \in W_0^{1,2}(\Omega)$ and is solenoidal, and $\|\nabla(\mathcal{R}_\varphi^\top \cdot \mathbf{u})\|_2 = \|\nabla \mathbf{u}\|_2$, so that from (EC) we deduce

$$-\frac{1}{2\pi} \int_0^{2\pi} \left(\int_{\Omega} (\mathcal{R}_\varphi^\top \cdot \mathbf{u}) \cdot \nabla_x \mathbf{V}_\varepsilon \cdot (\mathcal{R}_\varphi^\top \cdot \mathbf{u}) \, dx \right) d\varphi \leq \frac{\varepsilon}{2\pi} \int_0^{2\pi} \|\nabla \mathbf{u}\|_2^2 d\varphi = \varepsilon \|\nabla \mathbf{u}\|_2^2.$$

Combining the latter with (11) we thus obtain the desired inequality, namely,

$$-\int_{\Omega} \mathbf{u} \cdot \nabla (\mathcal{A}(\mathbf{V}_\varepsilon)) \cdot \mathbf{u} \, dy \leq \varepsilon \|\nabla \mathbf{u}\|_2^2, \quad (12)$$

for all solenoidal $\mathbf{u} \in W_0^{1,2}(\Omega)$. Denoting by $\mathcal{D}[\mathcal{A}(\mathbf{V}_\varepsilon)]$ the symmetric part of $\nabla \mathcal{A}(\mathbf{V}_\varepsilon)$, we show, on the one hand,

$$\int_{\Omega} \mathbf{u} \cdot \nabla (\mathcal{A}(\mathbf{V}_\varepsilon)) \cdot \mathbf{u} \, dy = \int_{\Omega} \mathbf{u} \cdot \mathcal{D}[\mathcal{A}(\mathbf{V}_\varepsilon)] \cdot \mathbf{u} \, dy$$

and, on the other hand, from (10),

$$\mathcal{D}[\mathcal{A}(\mathbf{V}_\varepsilon)] = \begin{pmatrix} -\frac{\Phi}{\pi} \frac{1}{r^2} & v'(r) - \frac{1}{r}v(r) \\ v'(r) - \frac{1}{r}v(r) & \frac{\Phi}{\pi} \frac{1}{r^2} \end{pmatrix}.$$

As a result, (12) becomes

$$\frac{\Phi}{\pi} \int_{\Omega} \frac{u_r^2 - u_{\theta}^2}{r^2} + \int_{\Omega} (v'(r) - \frac{1}{r}v(r))u_r u_{\theta} \leq \varepsilon \|\nabla \mathbf{u}\|_2^2, \quad (13)$$

for all solenoidal $\mathbf{u} \in W_0^{1,2}(\Omega)$, and where u_r and u_{θ} denote the polar components of \mathbf{u} . We now choose $\mathbf{u} = (u_r, u_{\theta})$ where

$$\begin{aligned} u_r &= \frac{m}{r} \frac{R_2 - R_1}{2\pi} \left\{ \cos \left[\frac{2\pi(r - R_1)}{R_2 - R_1} \right] - 1 \right\} \cos(m\theta) := U(r)[m \cos(m\theta)] \\ u_{\theta} &= \sin \left[\frac{2\pi(r - R_1)}{R_2 - R_1} \right] \sin(m\theta) := W(r)[\sin(m\theta)], \end{aligned} \quad (14)$$

with m an integer that will be specified further on. It is obvious that $\mathbf{u} \in W_0^{1,2}(\Omega)$, as well as, by taking into account that $(rU)' = -W$, that $\operatorname{div} \mathbf{u} = 0$ in Ω . Moreover, by a direct computation we show (with $\rho = R_2/R_1$)

$$\int_{\Omega} \frac{u_r^2 - u_{\theta}^2}{r^2} = m^2 F_1(\rho) - F_2(\rho), \quad (15)$$

where

$$\begin{aligned} F_1(\rho) &:= \frac{(\rho - 1)^3}{4\pi} \int_0^1 \frac{1}{[(\rho - 1)z + 1]^3} \{\cos(2\pi z) - 1\}^2 dz, \\ F_2(\rho) &:= \pi(\rho - 1) \int_0^1 \frac{1}{(\rho - 1)z + 1} \sin^2(2\pi z) dz. \end{aligned}$$

Furthermore, setting $G(r) := (v'(r) - \frac{1}{r}v(r))U(r)W(r)$, we get

$$\int_{\Omega} (v'(r) - \frac{1}{r}v(r))u_r u_{\theta} = m \int_{R_1}^{R_2} rG(r) dr \int_0^{2\pi} \sin(m\theta) \cos(m\theta) d\theta = 0. \quad (16)$$

Thus, by fixing m sufficiently large so that $\kappa := m^2 F_1(\rho) - F_2(\rho) > 0$, from (13)–(16) we conclude

$$\kappa \Phi \leq \varepsilon \|\nabla \mathbf{u}\|_2^2,$$

which, by the arbitrariness of $\varepsilon > 0$, and the assumption $\Phi > 0$ furnishes a contradiction. As originally showed by A. Takeshita [15], a similar result also holds if $\Phi < 0$. Actually, it is enough to choose in (13) instead of the field (14), the following one

$$u_r \equiv 0, \quad u_{\theta} = f(r)$$

with $f(r)$ any sufficiently smooth function satisfying $f(R_1) = f(R_2) = 0$, in which case the left-hand side of (13) becomes $-2\Phi \int_{R_1}^{R_2} r f^2(r) dr$.

3 The case Ω a Spherical Shell

In this case $\Omega := \{x \in \mathbb{R}^3 : R_1 < |x| < R_2\}$, $R_1 > 0$, $\Gamma_i := \{x \in \mathbb{R}^3 : |x| = R_i\}$, $i = 1, 2$,

$$\Phi := \int_{\Gamma_2} \mathbf{v}_* \cdot \mathbf{n} = - \int_{\Gamma_1} \mathbf{v}_* \cdot \mathbf{n},$$

and assume $\Phi > 0$. Again following the strategy of [15], the proof, in its first part is basically the same as in the case of the two-dimensional annulus. The only change being that the generic rotation matrix is now an element $\mathcal{R}_{\alpha_1\alpha_2\alpha_3} \in SO(3)$ characterized by the Euler angles α_i , $i = 1, 2, 3$. Consequently, (8) takes the form $\mathbf{y} = \mathcal{R}_{\alpha_1\alpha_2\alpha_3} \cdot \mathbf{x}$, and the average $\mathcal{A}(\mathbf{V}_\varepsilon)$ becomes

$$\mathcal{A}(\mathbf{V}_\varepsilon)(\mathbf{y}) = \frac{1}{8\pi^2} \int_0^{2\pi} \int_0^{2\pi} \int_0^\pi \mathcal{R}_{\alpha_1\alpha_2\alpha_3} \cdot \mathbf{V}_\varepsilon(\mathcal{R}_{\alpha_1\alpha_2\alpha_3}^\top \cdot \mathbf{y}) \sin \alpha_3 \, d\alpha_1 d\alpha_2 d\alpha_3.$$

Again, by the properties of \mathbf{V}_ε and the proper orthogonality of the rotation we show that the average satisfies (9). Moreover, the invariance of $\mathcal{A}(\mathbf{V}_\varepsilon)$ under the action of $SO(3)$ along with (9) implies that

$$\mathcal{A}(\mathbf{V}_\varepsilon) = \frac{\Phi}{4\pi r^2} \mathbf{e}_r,$$

where $\{\mathbf{e}_r, \mathbf{e}_\chi, \mathbf{e}_\theta\}$ is the base of a system of spherical coordinates (r, χ, θ) with the origin at $x = 0$; see [15, p. 157] for details. Next, proceeding verbatim as in Section 2, we prove that $\mathcal{A}(\mathbf{V}_\varepsilon)$ must satisfy (12), which by taking into account that this time

$$\{\mathcal{D}[\mathcal{A}(\mathbf{V}_\varepsilon)]\}_{ij} = \frac{\Phi}{4\pi r^3} \left(-3 \frac{x_i x_j}{r^2} + \delta_{ij} \right),$$

is equivalent to the following

$$\frac{\Phi}{4\pi} \int_\Omega \frac{2u_r^2 - u_\chi^2 - u_\theta^2}{r^3} \leq \varepsilon \|\nabla \mathbf{u}\|_2^2, \quad (17)$$

for all solenoidal $\mathbf{u} \in W_0^{1,2}(\Omega)$. In order to show that (17) leads to a contradiction, we choose

$$u_r = m \tilde{U}(r) \cos(m\theta) \sin \chi, \quad u_\chi \equiv 0, \quad u_\theta = \frac{1}{r} W(r) \sin(m\theta) \sin^2 \chi, \quad (18)$$

where $\tilde{U} = U/r$, and the functions U and W are defined in (14). It is easily proved that the vector \mathbf{u} with components given in (18) is solenoidal and is in $W_0^{1,2}(\Omega)$, and therefore can be replaced in (17). Since by a direct calculation we show that

$$\int_\Omega \frac{2u_r^2 - u_\chi^2 - u_\theta^2}{r^3} = m^2 G_1(\rho) - G_2(\rho)$$

where G_i , $i = 1, 2$, are *positive* functions of $\rho = R_2/R_1$, taking m sufficiently large and using the arbitrariness of ε , we show that (17) is incompatible with the assumption $\Phi > 0$. One can show the incompatibility of (17) also with the alternative assumption $\Phi < 0$, by using in (17) an appropriate function \mathbf{u} different from (18). This has been shown in [3] by the choice $\mathbf{u} = f(r) \sin \chi \mathbf{e}_\varphi$, with f sufficiently smooth and satisfying $f(R_1) = f(R_2) = 0$, in which case the left-hand side of (17) becomes $-\frac{2\Phi}{3} \int_{R_1}^{R_2} r^2 f(r) dr$.

4 The case Ω Multiply-Connected

We now assume that $\Omega \subset \mathbb{R}^n$, $n = 2, 3$, is a multiply-connected Lipschitz domain of the type defined in the Introduction. Furthermore, following [3], we suppose that for each connected component Γ_i of $\partial\Omega$, $i = 1, 2, \dots, N - 1$, there is a circumference (spherical surface) completely contained in Ω , and surrounding only the component Γ_i .

Combining the results of the previous two sections with Remarks 1 and 2, and the argument of [3, Theorem 1 and Corollary 1], we can show the following general result, whose proof follows exactly the same lines of [3, Corollary 1], and, consequently, will be omitted.

Theorem 4.1 *Let $\Omega \subset \mathbb{R}^n$, $n = 2, 3$, be a bounded domain satisfying the assumptions mentioned above. Moreover, let $\mathbf{v}_* \in W^{1/2,2}(\partial\Omega)$ obey the compatibility condition (3). Then the Leray-Hopf Extension Condition holds for \mathbf{v}_* (if and) only if $\Phi_k = 0$ for all $k = 1, 2, \dots, N$.*

Acknowledgements. Work partially supported by the NSF Grant DMS-1311983

References

- [1] R.A. Adams, and J.J. Fournier, Sobolev spaces. 2nd ed. *Pure and Applied Mathematics*, **140**, Elsevier/Academic Press, Amsterdam (2003)
- [2] C.J. Amick, Existence of Solutions to the Nonhomogeneous Steady Navier-Stokes Equations, *Indiana Univ. Math. J.*, **33**, 817–830 (1984)
- [3] R. Farwig, H. Kozono, and T. Yanagisawa, Leray’s Inequality in General Multi-Connected Domains in \mathbb{R}^n , *Math. Ann.*, **354**, 137–145 (2012)
- [4] R. Farwig, and H. Morimoto, Leray’s Inequality for Fluid Flow in Symmetric Multi-connected Two-dimensional Domains, *Tokyo J. of Math.*, **35**, 63–70 (2012)
- [5] H. Fujita, On Stationary Solutions to Navier–Stokes Equations in Symmetric Plane Domains under General Outflow Conditions, *Proceedings of International Conference on Navier-Stokes Equations, Theory and Numerical Methods*, June 1997, Varenna Italy, Pitman Research Notes in Mathematics, **388**, 16–30 (1998)
- [6] G.P. Galdi, An Introduction to the Mathematical Theory of the Navier-Stokes Equations, Vol. 2, Springer-Verlag, New York (1994)
- [7] G.P. Galdi, An Introduction to the Mathematical Theory of the Navier–Stokes Equations: Steady-State Problems, 2nd edition, Springer-Verlag, New York (2011)

- [8] J.G. Heywood, On the Impossibility, in Some Cases, of the Leray–Hopf Condition for Energy Estimates, *J. Math. Fluid Mech.*, **13**, 449–457 (2011)
- [9] E. Hopf, Ein allgemeiner Endlichkeitssatz der Hydrodynamik, *Math Annalen*, **117**, 764–775 (1941)
- [10] M.V. Korobkov, K. Pileckas, and R. Russo, On the Flux Problem in the Theory of Steady Navier–Stokes Equations with Nonhomogeneous Boundary Conditions, *Arch. Rational Mech. Anal.*, **207**, 185–213 (2013)
- [11] J. Leray, Etude de Diverses Équations Intégrales non Linéaires et de Quelques Problèmes que Pose l’ Hydrodynamique. *J. Math. Pures Appl.* **12**, 1–82 (1933)
- [12] H. Morimoto, A Remark on the Existence of 2-D Steady Navier–Stokes Flow in Bounded Symmetric Domain under General Outflow Condition, *J. Math. Fluid Mech.*, **9**, 411–418 (2007)
- [13] V.V. Pukhnachev, Viscous Flows in Domains with a Multiply Connected Boundary, *New directions in mathematical fluid mechanics, Adv. Math. Fluid Mech.*, 333–348, Birkhäuser Verlag, Basel, (2010)
- [14] L.I. Sazonov, On the Existence of a Stationary Symmetric Solution of the Two-dimensional Fluid Flow Problem, *Mathematical Notes*, **54**, 1280–1283 (1993)
- [15] A. Takeshita, A Remark on Leray’s Inequality, *Pacific J. Math.*, **157**, 151–158 (1993)