# Non-identifiability of identity coefficients at biallelic loci

Miklós Csűrös\*

May 7, 2018

#### Abstract

Shared genealogies introduce allele dependencies in diploid genotypes, as alleles within an individual or between different individuals will likely match when they originate from a recent common ancestor. At a locus shared by a pair of diploid individuals, there are nine combinatorially distinct modes of identity-by-descent (IBD), capturing all possible combinations of coancestry and inbreeding. A distribution over the IBD modes is described by the nine associated probabilities, known as (Jacquard's) identity coefficients. The genetic relatedness between two individuals can be succinctly characterized by the identity coefficients corresponding to the joint genealogy. The identity coefficients (together with allele frequencies) determine the distribution of joint genotypes at a locus. At a locus with two possible alleles, identity coefficients are not identifiable because different coefficients can generate the same genotype distribution.

We analyze precisely how different IBD modes combine into identical genotype distributions at diallelic loci. In particular, we describe IBD mode mixtures that result in identical genotype distributions at all allele frequencies, implying the non-identifiability of the identity coefficients from independent loci. Our analysis yields an exhaustive characterization of relatedness statistics that are always identifiable. Importantly, we show that identifiable relatedness statistics include the kinship coefficient (probability that a random pair of alleles are identical by descent between individuals) and inbreeding-related measures, which can thus be estimated from genotype distributions at independent loci.

#### 1 Introduction

Non-random mating histories, selection, finite population sizes, and many other causes create dependencies between alleles in a diploid population. Because of joint genealogies, alleles may match within or across genotypes for being unmodified copies of a common ancestral state. Such alleles are said to be

<sup>\*</sup>Department of Computer Science and Operations Research, University of Montréal; E-mail: csuros@iro.umontreal.ca.

IBD mode	A's genotype	B's genotype	identity coefficient	IBD mode	A's genotype	B's genotype	identity coefficient
A x x B x x	<i>x</i> / <i>x</i>	<i>x</i> / <i>x</i>	$\Delta_1$	A <u>x</u> x B <b>y</b> y	<i>x</i> / <i>x</i>	y/y	$\Delta_2$
A X X B X Y	<i>x</i> / <i>x</i>	<i>x</i> / <i>y</i>	$\Delta_3$	A x x B (y) (2)	<i>x</i> / <i>x</i>	y/z	$\Delta_4$
А <u>х</u> <u>у</u> В <u>х</u> х	<i>x</i> / <i>y</i>	<i>x</i> / <i>x</i>	$\Delta_5$	A 🗶 Z B (Y)-(Y)	<i>x</i> / <i>z</i>	y/y	$\Delta_6$
A x y B x z	x/y	x/z	$\Delta_8$	A <u>x</u> <u>y</u> B <u>z</u> <u>w</u>	x/y	z/w	$\Delta_9$
Α <u>×</u> γ Β × γ	x/y	x/y	$\Delta_7$				

Figure 1: Identity modes for a diploid genotype pair. Thick red lines mark identity by descent. Alleles  $x, y, \ldots$  observed in the genotypes may be equal. Probabilities for different modes are denoted by the identity coefficients  $\Delta_i$ .

*identical by descent* [9, 6]. Combinatorially distinct partitionings of identicalby-descent (IBD) alleles for a pair of unordered diploid genotypes are called *identity modes* [6]. The two individuals' joint pedigree defines the possible inheritance histories for the four alleles, which combine into a distribution over the identity modes. Every identity mode generates its own probability distribution over the joint genotypes at a locus, and the observable genotypic distribution is the mixture of the mode-specific distributions. The probabilities of the identity modes, or *identity coefficients*, characterize thus the individuals' genetic relatedness succinctly.

The identity coefficients can be computed for any known pairwise genealogy [3, 7]. Hypothetical pedigrees can be thus assessed by comparing implied genotype distributions with empirical ones [14, 10]. But can identity coefficients be directly inferred from genotype distributions without genealogies? The answer depends on the number of alleles. There are nine identity modes for a pair of diploid individuals (Figure 1), which define eight independent identity coefficients; the ninth one is implied since the coefficients sum to one. At loci with only two alleles, nine genotype pairs are possible, but because of redundancy, there are not enough many different genotype pairs to make certain inference possible: more than one set of coefficients generate the same joint genotype distribution. Genotype distributions at loci with three or more alleles, however, convey enough information in principle to identify a single set of identity coefficients that produce it. Among molecular markers, multiallelic microsatellite loci provide in consequence high discriminatory power for a detailed characteriza-

Mode	0/0:0/0	1/1:1/1	1/1:0/1	0/1:1/1	0/1:0/1	1/1:0/0	0/0:1/1	0/1:0/0	0/0:0/1
1	q	p	0	0	0	0	0	0	0
2	$q^2$	$p^2$	0	0	0	pq	pq	0	0
3	$q^2$	$p^2$	pq	0	0	0	0	0	pq
4	$q^3$	$p^3$	$2p^2q$	0	0	$pq^2$	$p^2q$	0	$2pq^2$
5	$q^2$	$p^2$	0	pq	0	0	0	pq	0
6	$q^3$	$p^3$	0	$2p^2q$	0	$p^2q$	$pq^2$	$2pq^2$	0
7	$q^2$	$p^2$	0	0	2pq	0	0	0	0
8	$q^3$	$p^3$	$p^2q$	$p^2q$	pq(p+q)	0	0	$pq^2$	$pq^2$
9	$q^4$	$p^4$	$2p^3q$	$2p^3q$	$4p^2q^2$	$p^2q^2$	$p^2q^2$	$2pq^3$	$2pq^3$

 Table 1: Distribution of biallelic genotypes by identity mode

Genotypic probabilities for every mode are given by assuming that alleles are chosen independently for each IBD group, with probability p for allele 1 (minor allele) and with probability q for allele 0 (major allele). Genotypes are unordered (1/0 and 0/1 are considered equivalent).

tion of genetic relatedness, but diallelic single-nucleotide and insertion-deletion have restricted utility [17]. Here, we scrutinize the inherent ambiguity of relatedness in diallelic genotypes. Specifically, our aim is to find what aspects of coancestry result in non-identifiability and to characterize statistical measures of the identity mode distribution that can be consistently estimated from joint genotype frequencies.

#### 2 Theory

#### 2.1 Identity coefficients and biallelic genotype distributions

*Identity by descent* [9] encapsulates the dependence between diploid genotypes due to shared parentage. Two alleles are identical by descent (IBD) if they originate from a common ancestral allele without modification. Equivalence relations for four alleles of two diploid genotypes take one of nine combinatorially distinct forms [5, 6, 7], or *identity modes*, as illustrated in Figure 1.

The individuals' joint pedigree determines the possible identity modes and their associated frequencies, specified by the vector of coefficients  $\Delta_i: i = 1, \ldots, 9$  using the notation of Jacquard [6]. For instance, children of the same parents from non-overlapping lineages inherit two IBD alleles with probability  $\Delta_7 = \frac{1}{4}$ , one IBD set from either parent with probability  $\Delta_8 = \frac{1}{2}$ , and four independent alleles with probability  $\Delta_9 = \frac{1}{4}$ .

Suppose that the locus has two alleles, and alleles 1 and 0 (minor and major) occur with frequencies p and q, respectively in both individuals. Every mode generates its own conditional distribution of joint genotypes. In mode 8, the individuals are 0/1 heterozygotes simultaneously with probability  $pq^2 + p^2q$  since either the IBD alleles are the mutants, or two mutant alleles are chosen independently. In contrast, if all four alleles are sampled independently (iden-

tity mode 9) then the joint genotype 0/1 : 0/1 occurs with probability  $4p^2q^2$ , accounting for two minor and two major alleles in 4 possible orderings. Table 1 lists the complete set of genotypic probabilities. Denote the distribution of joint genotypes by

$$\mathbf{f} = (f_{0000}, f_{1111}, f_{1101}, f_{0111}, f_{0101}, f_{1100}, f_{0011}, f_{0100}, f_{0001}).$$

Table 1 corresponds to the system of equations

$$f_{0000} = q\Delta_1 + q^2 (\Delta_2 + \Delta_3 + \Delta_5 + \Delta_7) + q^3 (\Delta_8 + \Delta_4 + \Delta_6) + q^4 \Delta_9$$
  

$$f_{1111} = p\Delta_1 + p^2 (\Delta_2 + \Delta_3 + \Delta_5 + \Delta_7) + p^3 (\Delta_4 + \Delta_6 + \Delta_8) + p^4 \Delta_9$$
  

$$f_{1101} = pq (\Delta_3 + p(\Delta_8 + 2\Delta_4 + 2p\Delta_9))$$
  

$$f_{0111} = pq (\Delta_5 + p(\Delta_8 + 2\Delta_6 + 2p\Delta_9))$$
  

$$f_{0101} = pq (\Delta_2 + q\Delta_8 + 4pq\Delta_9)$$
  

$$f_{0011} = pq (\Delta_2 + q\Delta_4 + p\Delta_6 + pq\Delta_9)$$
  

$$f_{0100} = pq (\Delta_5 + q\Delta_8 + 2q\Delta_6 + 2q^2\Delta_9)$$
  

$$f_{0001} = pq (\Delta_3 + q\Delta_8 + 2q\Delta_4 + 2q^2\Delta_9).$$
  
(1)

In matrix form,

$$\mathbf{f} = \mathbf{F} \cdot \mathbf{\Delta},\tag{2}$$

and Table 1 gives the transpose of **F**. Note that the matrix structure guarantees  $\sum_i \Delta_i = 1$  when  $f_{0000} + f_{1111} + \ldots + f_{0001} = 1$  since the all-1 row vector  $\mathbf{e} = \begin{pmatrix} 1 & 1 & \cdots & 1 \end{pmatrix}$  is a left eigenvector:

$$1 = \mathbf{e} \cdot \mathbf{f} = \mathbf{e} \cdot \mathbf{F} \cdot \mathbf{\Delta} = \mathbf{e} \cdot \mathbf{\Delta}.$$

The matrix  $\mathbf{F}$  projects the vector of identity coefficients  $\Delta$  to the vector of genotype probabilities  $\mathbf{f}$ . Consequently, identity coefficients can be inferred from the biallelic genotype distribution if and only if the matrix  $\mathbf{F}$  is invertible. The matrix rows are, however, linearly dependent.

**Claim 1** When p + q = 1, dependencies between genotype probabilities include the following two.

$$f_{1101} + 2f_{1100} + f_{0100} = f_{0111} + 2f_{0011} + f_{0001};$$
(3)

$$p = f_{1111} + \frac{3}{4} (f_{1101} + f_{0111})$$
(4)

$$+\frac{1}{2}(f_{0101}+f_{1100}+f_{0011})+\frac{1}{4}(f_{0100}+f_{0001}).$$

Theorem 2 below characterizes the set of identity coefficients that lead to the same distribution over joint biallelic genotypes. **Theorem 2** Suppose that p + q = 1. If  $\Delta_i$ : i = 1, ..., 9 satisfy (2) then so do the following coefficients, for all choices of  $\xi, \eta \in \mathbb{R}$ .

$$\begin{aligned}
\Delta_1' &= \Delta_1 - \eta p q & \Delta_2' &= \Delta_2 + \xi - \eta p q \\
\Delta_3' &= \Delta_3 + 2\eta p q & \Delta_4' &= \Delta_4 - \xi \\
\Delta_5' &= \Delta_5 + 2\eta p q & \Delta_6' &= \Delta_6 - \xi \\
\Delta_7' &= \Delta_7 - \xi + \eta (1 - 2pq) \\
\Delta_8' &= \Delta_8 + 2\xi - 2\eta & \Delta_9' &= \Delta_9 + \eta.
\end{aligned}$$
(5)

Starting from an arbitrary particular solution set  $\Delta_i$ , Equation (5) generates all vector solutions to (2).

Values of  $(\xi, \eta)$  for which Eq. (5) produces a proper distribution are precisely those where  $\Delta'_i \geq 0$  for all *i*:

$$\eta \leq \frac{\Delta_1}{pq} \qquad \eta \leq \frac{\Delta_2}{pq} + \frac{\xi}{pq} \qquad \eta \leq \frac{\Delta_8}{2} + \xi$$
  
$$\eta \geq -\frac{\Delta_3}{2pq} \qquad \eta \geq -\frac{\Delta_5}{2pq} \qquad \eta \geq -\frac{\Delta_7}{1-2pq} + \frac{\xi}{1-2pq} \qquad \eta \geq -\Delta_9 \qquad (6)$$
  
$$\xi \leq \Delta_4 \qquad \xi \leq \Delta_6$$

Figure 2 illustrates the solution area of (6) for the identity coefficients of a real-life example (Queen Victoria and Prince Albert, who shared multiple common ancestors within seven generations — Appendix for complete family tree).

#### 2.2 Identifiable relatedness parameters

Despite multiple solutions, some aspects of the identity coefficients can be ascertained from the genotype distribution. In particular, if a linear combination stays the same for all sets of identity coefficients from (5), then it is computable from the biallelic genotype distribution. Theorem 3 formalizes our argument.

**Definition 1** A function of the identity distribution  $\theta(\Delta)$  is called a linear relatedness parameter if and only if it can be written as a linear combination

$$\theta(\mathbf{\Delta}) = \sum_{i=1}^{9} a_i \Delta_i,\tag{7}$$

where  $a_i$  are constants. In particular,  $a_i$  may not depend on the allele frequency p.

**Theorem 3** A linear relatedness parameter  $\theta$  is identifiable from the joint genotype distribution only if

$$a_{2} + 2a_{8} = a_{4} + a_{6} + a_{7};$$
  

$$a_{7} + a_{9} = 2a_{8};$$
 and (8)  

$$2a_{3} + 2a_{5} = a_{1} + a_{2} + 2a_{7}.$$



Identity coefficient	$\times 2^{-20}$	decimal value	ambiguity range	
			$\min$	max
$\Delta_1$	34	$3.24 \cdot 10^{-5}$	0	$1.78 \cdot 10^{-4}$
$\Delta_2$	1	$9.54 \cdot 10^{-7}$	0	$7.58 \cdot 10^{-4}$
$\Delta_3$	324	$3.09 \cdot 10^{-4}$	$1.85 \cdot 10^{-5}$	$3.74 \cdot 10^{-4}$
$\Delta_4$	665	$6.34 \cdot 10^{-4}$	0	$7.80 \cdot 10^{-4}$
$\Delta_5$	3140	$2.99 \cdot 10^{-3}$	$2.70 \cdot 10^{-3}$	$3.06 \cdot 10^{-3}$
$\Delta_6$	9113	$8.69 \cdot 10^{-3}$	$8.06 \cdot 10^{-3}$	$8.84 \cdot 10^{-4}$
$\Delta_7$	5087	$4.85 \cdot 10^{-3}$	0	$5.94 \cdot 10^{-3}$
$\Delta_8$	278698	0.266	0.263	0.276
$\Delta_9$	751514	0.717	0.711	0.718
Relatedness parameter				
$\theta_1$ (kinship)	73984	0.071		
$\theta_{2A}$ (Victoria's inbreeding)	1024	$9.77 \cdot 10^{-3}$		
$\theta_{2B}$ (Albert's inbreeding)	12288	0.0117		
$\theta_3$ (two-out-of-three IBD)	152824	0.146		
$\theta_{3:3}$ (three-out-of-three IBD)	1766	$1.68 \cdot 10^{-3}$		
$\theta_4 = (\Delta_4 - \Delta_6)/2$	-8448	$-8.06 \cdot 10^{-3}$		

Figure 2: Null space identity coefficients. The intersection of the constraints from Equation (6) defines the convex polygonal area within which  $(\xi, \eta)$  values plugged into Eq. (5) yield valid identity mode distributions that generate the same genotypic distribution. The example is based on the joint parentage of Queen Victoria of the United Kingdom (1819–1901) and her spouse Prince Albert of Saxe-Coburg and Gotha (1819–1861), for which the relevant parameters are listed below the plot. The illustration assumes pq = 0.02746... reflecting typical allele frequency moments in humans (from the 1000 Genomes project). Extremal values of possible  $\Delta_i$  listed under "ambiguity range" are attained in the corners of the unshaded area.

**Theorem 4** The following linear relatedness parameters are identifiable from the biallelic genotype distribution.

$$\theta_0 = \sum_{i=1}^9 \Delta_i \tag{9a}$$

$$\theta_1 = \Delta_1 + \frac{1}{2} (\Delta_3 + \Delta_5 + \Delta_7) + \frac{1}{4} \Delta_8 \qquad (kinship \ coefficient) \qquad (9b)$$

 $\theta_{2A} = \Delta_1 + \Delta_2 + \Delta_3 + \Delta_4 \qquad (A's inbreeding) \qquad (9c)$ 

$$\theta_{2B} = \Delta_1 + \Delta_2 + \Delta_5 + \Delta_6 \qquad (B's inbreeding) \qquad (9d)$$

$$\theta_3 = \Delta_1 + \Delta_2 + \Delta_3 + \Delta_5 + \Delta_7 \tag{9e}$$

$$+\frac{1}{2}(\Delta_4 + \Delta_6 + \Delta_8)$$
  
$$\theta_4 = \frac{1}{2}(\Delta_4 - \Delta_6) \tag{9f}$$

All other identifiable parameters are linear combinations of  $\theta_i$  in Equations (9).

Theorem 3 shows that, in general, the identity coefficients  $\Delta_i$  are not identifiable separately. In particular, probabilities for various inbred modes  $(\Delta_3, \Delta_4, \Delta_5, \Delta_6)$  are not identifiable, only their differences  $(\Delta_4 - \Delta_6 = 2\theta_4 \text{ and } \Delta_3 - \Delta_5 = \theta_{2A} - \theta_{2B} - 2\theta_4)$ .

The identifiable parameters of Theorem 4 include the usual measures of inbreeding  $(\theta_{2*})$  and coancestry  $(\theta_1)$  generalized to inbred parents [5], as well as the trivial  $\sum_i \Delta_i$ . The parameter  $\theta_3$  is the probability that there is at least one pair of IBD alleles among three randomly selected ones. A simpler three-gene characterization of inbred coancestry is

$$\theta_{3:3} = \Delta_1 + \frac{1}{2} \left( \Delta_3 + \Delta_5 \right) = \theta_1 - \frac{1}{2} \theta_3 + \frac{1}{4} \left( \theta_{2A} + \theta_{2B} \right), \tag{10}$$

which is the probability that three randomly chosen alleles are simultaneously identical by descent. By Theorem 3,  $\theta_{3:3}$  is identifiable, and (10) shows how to write it as a linear combination of identifiable parameters from Theorem 4.

Linear relatedness parameters, defined as linear combinations of identity coefficients, can be written as linear combinations of genotypic probabilities in the linear-algebraic framework of Equation (1). For the archetypical parameters of Theorem 4, we consider the following expressions.

$$\tau_{1A} = \frac{f_{1101} + f_{0100}}{2} + \frac{f_{0101}}{4} + f_{1100} \qquad \tau_{1B} = \frac{f_{0111} + f_{0001}}{2} + \frac{f_{0101}}{4} + f_{0011}$$
  

$$\tau_{1} = \frac{\tau_{1A} + \tau_{1B}}{2}$$
  

$$\tau_{2A} = \frac{f_{0111} + f_{0101} + f_{0100}}{2} \qquad \tau_{2B} = \frac{f_{1101} + f_{0101} + f_{0001}}{2}$$
  

$$\tau_{3} = \frac{(f_{0100} - f_{0111}) + (f_{0001} - f_{1101})}{4} \qquad \tau_{4} = \frac{f_{1100} - f_{0011}}{2}$$
(11)

Due to the linear dependencies, multiple equivalent formulas exist that relate the genotype distribution and any specific parameter. For instance, the intermediate quantities  $\tau_{1A}$ ,  $\tau_{1B}$  and  $\tau_1$ , which weigh genotypic probabilities differently, are equal by Equation (3).

#### 2.3 Relatedness estimation from independent loci

We examine two applications of estimating pairwise relatedness from observed genotypes at independent diallelic loci.

Application I: relatedness between two individuals In this application [11, 8], n independent sites are genotyped in two genomes. Minor-allele frequencies  $p_i$  follow population-wide background frequencies, and apply to both genomes equally. Our aim is to characterize the IBD mode distribution implied by the joint genealogy, using the observed genotypes.

Application II: background relatedness in a structured population We sample a set of n random pairs from a population at the same locus, and we would like to infer the background structure of relatedness [1, 16], described as the population-wide distribution of IBD modes. Accordingly, the same minorallele frequency  $p_i \equiv p$  applies to all pairs i = 1, 2, ..., n.

Theorem 2 suggests that, identity coefficients cannot always be inferred from observed genotypes in either application. We formalize our argument using the following abstraction. Suppose that minor-allele frequencies  $p_i: i = 1, ..., n$  apply to n independent biallelic loci sampled by random pairwise genotypes  $X_i \in$  $\{0/0, 0/1, 1/1\}^2: i = 1, 2, ..., n$ . The pairs have identical IBD mode distribution  $\Delta$ . As in Eq. (2), the joint genotype distribution  $\mathbf{f}^{(i)}$  of  $X_i$  is related to the identity coefficients by a matrix  $\mathbf{F}^{(i)}$  through  $\mathbf{f}^{(i)} = \mathbf{F}^{(i)} \cdot \Delta$ . The matrix entries are given by (1), substituting  $p \leftarrow p_i$  and  $q \leftarrow 1 - p_i$ .

**Theorem 5** Let  $T_n(X_1, X_2, ..., X_n)$  be an estimator of the identity coefficients  $\Delta$ . If  $\min{\{\Delta_4, \Delta_6, \Delta_7\}} > 0$  or  $\min{\{\Delta_2, \Delta_8\}} > 0$  or both, then  $T_n$  may not converge (in probability) to  $\Delta$  as  $n \to \infty$  for any sequence of minor-allele frequencies  $p_i: i = 1, 2, ...$ 

In other words, identity coefficients cannot be estimated consistently in general. Theorem 4 suggests, however, that a maximal set of linear relatedness

So,

parameters can be inferred consistently. As an illustration, we prove the consistency of simple moment-based estimators using observed genotype frequencies. Define the genotype counts  $n_{0000}, n_{1111}, n_{1101}, \ldots, n_{0001}$ , and the *empirical genotype frequencies*  $\hat{f}_x = n_x/n$ . Written with indicator variables  $\{X_i = x\}$  for  $x = 0000, 1111, \ldots, 0001$ ,

$$\hat{f}_x = \frac{n_x}{n} = \frac{\sum_{i=1}^n \{X_i = x\}}{n}.$$
(13)

Plugged into Equation (11), we get the sample statistics

$$\begin{aligned} \hat{\tau}_{1A} &= \frac{\hat{f}_{1101} + \hat{f}_{0100}}{2} + \frac{\hat{f}_{0101}}{4} + \hat{f}_{1100} \\ \hat{\tau}_{1B} &= \frac{\hat{f}_{0111} + \hat{f}_{0001}}{2} + \frac{\hat{f}_{0101}}{4} + \hat{f}_{0011} \qquad \hat{\tau}_{1} = \frac{\hat{\tau}_{1A} + \hat{\tau}_{1B}}{2} \\ \hat{\tau}_{2A} &= \frac{\hat{f}_{0111} + \hat{f}_{0101} + \hat{f}_{0100}}{2} \qquad \hat{\tau}_{2B} = \frac{\hat{f}_{1101} + \hat{f}_{0101} + \hat{f}_{0001}}{2} \\ \hat{\tau}_{3} &= \frac{(\hat{f}_{0100} - \hat{f}_{0111}) + (\hat{f}_{0001} - \hat{f}_{1101})}{4} \qquad \hat{\tau}_{4} = \frac{\hat{f}_{1100} - \hat{f}_{0011}}{2}. \end{aligned}$$
(14)

Write the moments for the minor-allele frequency (MAF) distribution as

$$\bar{\mu} = \frac{1}{n} \sum_{i=1}^{n} p_i \quad \bar{\mu}_2 = \frac{1}{n} \sum_{i=1}^{n} p_i^2 \quad \bar{\mu}_3 = \frac{1}{n} \sum_{i=1}^{n} p_i^3 \quad \bar{\mu}_4 = \frac{1}{n} \sum_{i=1}^{n} p_i^4,$$

and the average centralized moments as

$$\bar{\nu}_2 = \frac{1}{n} \sum_{i=1}^n \left( p_i (1-p_i)^2 + (1-p_i) p_i^2 \right) = \bar{\mu} - \mu_2$$
$$\bar{\nu}_3 = \frac{1}{n} \sum_{i=1}^n \left( p_i (1-p_i)^3 - (1-p_i) p_i^3 \right) = \bar{\mu} - 3\bar{\mu}_2 + 2\bar{\mu}_3$$

Finally, define the relatedness estimators

$$\hat{\theta}_{1} = 1 - \frac{\hat{\tau}_{1}}{\bar{\nu}_{2}} \quad \hat{\theta}_{2A} = 1 - \frac{\hat{\tau}_{2A}}{\bar{\nu}_{2}} \quad \hat{\theta}_{2B} = 1 - \frac{\hat{\tau}_{2B}}{\bar{\nu}_{2}}$$

$$\hat{\theta}_{3} = 1 - \frac{\hat{\tau}_{3}}{\bar{\nu}_{3}} \quad \hat{\theta}_{4} = \frac{\hat{\tau}_{4}}{\bar{\nu}_{3}}.$$
(15)

**Theorem 6** Suppose that there exist constants  $\nu_2, \nu_3$  such that in Eq. (15),  $\bar{\nu}_2 \xrightarrow{P} \nu_2$  (converges in probability to  $\nu_2$ ) and  $\bar{\nu}_3 \xrightarrow{P} \nu_3$  as  $n \to \infty$ . If  $\nu_2, \nu_3 \neq 0$ , then  $\hat{\theta}_y \xrightarrow{P} \theta_y$  for all parameters y = 1, 2A, 2B, 3, 4 from Theorem 4.

The theorem implies that one can consistently estimate five non-trivial parameters of the background IBD structure in a population (Application II) if the sampled locus has minor-allele frequency  $p \neq 0, 1, 1/2$ . In Application I, our moment-based estimators are consistent when, for instance, minor-allele frequencies are independent and identically distributed, with non-zero variance (i.e., sites are mostly segregating) and kurtosis (excluding symmetric MAF distributions around 1/2).

## 3 Discussion

Identity coefficients [6, 5] encapsulate the allele dependencies within the joint genotype distribution for a pair of diploid individuals If the individuals are not inbred, only three of the coefficients may be positive  $(\Delta_7, \Delta_8, \Delta_9)$ , corresponding to Cotterman's k-gene coefficients [14] for the individuals sharing k = 0, 1, or 2 alleles between them. The three coefficients can be retrieved from sampled genotypes using well-established methods relying on likelihood maximization [14, 10] or allele frequency moments [11, 8].

It may be of interest to estimate higher-order identity coefficients simultaneously. In particular, all nine IBD modes may occur if the individuals have inbred coancestries [5], or come from a structured population [16]. Biallelic genotypes, however, do not convey enough information about the generic IBD structure, since different identity coefficients can generate the same joint genotype distribution. Theorem 2 scrutinizes the inherent ambiguity about the identity coefficients, describing the linear subspace in which all solutions are found. One particular source of the ambiguity (corresponding to the null vector  $\mathbf{z}_1$ in (16) is that symmetric mixtures of simultaneous inbreeding and coancestry (modes  $\Delta_7 - \Delta_4 - \Delta_6$  vs.  $\Delta_2 - \Delta_8 - \Delta_8$  in Figure 1) manifest identically in the genotype distribution. Importantly, these equivalent solutions (varying only the  $\xi$ coordinate) remain equivalent for any minor-allele frequency. The uncertainties about the identity coefficients are within the same magnitude as the inbreeding levels (Eq. (5)) when both individuals are inbred  $(\Delta_4, \Delta_6 > 0)$  and share multiple ancestors ( $\Delta_7 > 0$ ). Indeed, the real-life example of Figure 2 shows that the subtle details of coancestry can be irretrievable from the genotype distribution.

Consistent estimation is thus impossible since even as the number of independent sampled loci n goes to infinity and genotype frequencies concentrate around their true probabilities, the identity coefficients stay ambiguous regardless of the estimation method used (Theorem 5). The decomposition of the solution space (Theorem 3) shows the aspects of the IBD structure that can instead be inferred from biallelic genotypes. Specifically, Theorem 4 lists five non-trivial relatedness parameters, deconvolving the IBD structure to the maximum degree that is attainable. Principal aspects of genetic relatedness, quantified by the coefficients of kinship and inbreeding, are identifiable. Other identifiable attributes are probabilities for three-gene joint IBD and the asymmetry of inbreeding modes with and without simultaneous coancestry. In contrast, parameters that do not weigh the identity coefficients properly (Eq. (8)) are not identifiable from the biallelic genotypes. Ill-defined relatedness parameters include the probabilities of separate identity modes (e.g., the probability  $\Delta_1$  of fourfold IBD), the fraternity coefficient  $(\Delta_1 + \Delta_7)$  and other generalizations of Cotterman's k-gene coefficients.

In the case of human genomes, the limits of inferring relatedness are set by linkage and finite genome size, and not identifiability [12]. The mean length of a segment with the same particular history involving m meioses decreases linearly with m. In human whole genome sequences, IBD segments of length  $0.4 \,\mathrm{cM}$  can be demarcated [13] with confidence by high-coverage sequencing. The detection of shared ancestry is thus constrained by the fact that descendants inherit a common ancestor's allele simultaneously with exponentially small probability in the number of meioses separating them  $(2^{-m+1})$ . As Browning and Browning [2] point out, fifth cousins (m = 12) simultaneously inherit 1/2048 of their genome on expectation from the shared great-great-great-great-great grandfather or greatgreat-great-great grandmother each, which amounts to about 1.5 cM in an entire human genome, while the average IBD segment length is 8.3 cM. Then, by Markov's inequality, there is at least one IBD segment between the two cousins' genomes with probability at most  $\frac{2 \times 1.5}{8.3} = 0.35...$  Identity modes with more than two IBD alleles  $(\Delta_1, \Delta_2, \Delta_3, \Delta_5, \Delta_7)$  usually involve even more distant ancestries and are, thus, almost certainly undetectable [15]. For example, if both individuals are children of fourth cousins (as the royal cousins here), the simultaneous inbreeding mode  $\Delta_2$  appears in segments of average length 4.2 cM, covering  $1/2^{20}$  of their genome (about 0.003 cM); so, no such segments are seen in at least 99.93% of the cases.

The ambiguity of identity coefficients (outlined by Theorem 2) complements well-known results on equivalent pedigrees [4, 12]. In the absence of linkage information, the non-identifiable mode combinations represent the theoretical limits of dissecting the IBD structure. By our results (Theorem 4), only two more distribution parameters can be inferred in addition to the usual two-gene coefficients for coancestry and inbreeding: one for three-gene IBD, and another measuring asymmetry in inbreeding modes.

#### Acknowledgments

I am grateful for valuable commentary on earlier versions of the manuscript by Damian Labuda and anonymous reviewers. This research project was partially funded by the author's individual discovery grant from the *Natural Sciences and Engineering Research Council of Canada*.

## References

- Amy D. Anderson and Bruce S. Weir. A maximum-likelihood method for the estimation of pairwise relatedness in structured populations. *Genetics*, 176:421–440, 2007.
- [2] Sharon R. Browning and Brian L. Browning. Identity by descent between distant relatives: Detection and applications. *Annual Review of Genetics*, 46:617–633, 2012.
- [3] C. Clark Cockerham. Higher order probability functions of identity of alleles by descent. *Genetics*, 69:235–246, 1971.
- [4] Kevin P. Donnelly. The probability that related individuals share some section of genome identical by descent. *Theoretical Population Biology*, 23:34–63, 1983.
- [5] Dewey L. Harris. Genotypic covariances between inbred relatives. *Genetics*, 50:1319–1348, 1964.
- [6] Albert Jacquard. Genetics of Human Populations. Springer, New York, 1974.
- [7] Kenneth Lange. Mathematical and Statistical Methods for Genetic Analysis. Springer, New York, 1997.
- [8] Michael Lynch and Kermit Ritland. Estimation of pairwise relatedness with molecular markers. *Genetics*, 152:1753–1766, 1999.
- [9] G. Malécot. The Mathematics of Relationship. W. H. Freeman, San Francisco, 1969.
- [10] Brook G. Milligan. Maximum-likelihood estimation of relatedness. Genetics, 163:1153–1167, 2003.
- [11] Kermit Ritland. Estimators for pairwise relatedness and individual inbreeding coefficients. *Genetical Research*, 67:175–185, 1996.
- [12] Øivind Skare, Nuala Sheehan, and Thoire Egeland. identification of distant family relationships. *Bioinformatics*, 25(18):2376–2382, 2009.
- [13] Shu-Yi Su, Jay Kasberger, Sergio Baranzini, William Byerley, Wilson Liao, et al. Detection of identity by descent using next-generation whole genome sequencing data. *BMC Bioinformatics*, 13:121, 2012.
- [14] E. A. Thompson. The estimation of pairwise relationships. Annals of Human Genetics, 39:173–188, 1975.
- [15] E. A. Thompson. The IBD process along four chromosomes. *Theoretical Population Biology*, 73(3):369–373, 2008.

- [16] Jinliang Wang. Unbiased relatedness estimation in structured populations. Genetics, 187:887–901, 2011.
- [17] Bruce S. Weir, Amy D. Anderson, and Amanda B. Hepler. Genetic relatedness analysis: modern data and new challenges. *Nature Reviews Genetics*, 7:771–780, 2006.

# A Linear algebra for genotypic probabilities and identity coefficients

Proof of Claim 1. The equalities can be seen by inspecting the rows of  $\mathbf{F}$ , but considering allele counts gives a more straightforward proof. Consider the expected number  $\omega$  of minor ('1') alleles in the random joint genotype. Since it is the expectation for the sum of four indicator variables,  $\mathbb{E}\omega = 4p$ . Alternatively, by summing over the possible joint genotypes,  $\mathbb{E}\omega = 4f_{1111} + 3(f_{1101} + f_{0111}) + 2(f_{0101} + f_{1100} + f_{0011}) + (f_{0100} + f_{0001})$ , and Equation (4) follows after dividing by 4. Now consider the expected number of minor alleles in the A's and B's genotype separately. Clearly, both equal 2p. Counting by joint genotypes:

$$\underbrace{2(f_{1111} + f_{1101} + f_{1100}) + (f_{0111} + f_{0101} + f_{0100})}_{\text{expected count in A}} = \underbrace{2(f_{1111} + f_{0111} + f_{0011}) + (f_{1101} + f_{0101} + f_{0001})}_{\text{expected count in B}}.$$

After elimination of common terms, Equation (3) follows.

*Proof of Theorem 2.* The null space of  $\mathbf{F}$  is spanned by the vectors

$$\mathbf{z}_{1} = \begin{pmatrix} 0 \\ 1 \\ 0 \\ -1 \\ 0 \\ -1 \\ -1 \\ 2 \\ 0 \end{pmatrix} \qquad \mathbf{z}_{2} = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 1 \\ -2 \\ 1 \end{pmatrix} + pq \begin{pmatrix} -1 \\ -1 \\ 2 \\ 0 \\ 2 \\ 0 \\ -2 \\ 0 \\ 0 \end{pmatrix} = \mathbf{z}_{2}^{(1)} + pq\mathbf{z}_{2}^{(2)} \qquad (16)$$

It is straightforward to verify that  $\mathbf{F}\mathbf{z}_1 = \mathbf{F}\mathbf{z}_2 = \mathbf{0}$ , the null vector. Hence,

$$\mathbf{F} \cdot (\mathbf{\Delta} + \xi \mathbf{z}_1 + \eta \mathbf{z}_2) = \mathbf{F} \cdot \mathbf{\Delta} = \mathbf{f}$$

for all choices of  $\xi, \eta$ . The rank of the  $9 \times 9$  matrix **F** is 7 (established by Gaussian elimination), and therefore no other solutions exist.

Proof of Theorem 3. In order to be identifiable,  $\theta(\Delta)$  must remain the same for all distributions satisfying (2). The vector of coefficients  $(a_i: i = 1, \ldots, 9)$ then has to be orthogonal to the null space of **F**. The identities of (8) express the orthogonality with the vectors  $\mathbf{z}_1$ ,  $\mathbf{z}_2^{(1)}$  and  $\mathbf{z}_2^{(2)}$ : the latter two are used separately since orthogonality must be maintained for all p.

Proof of Theorem 4. By Theorem 3, identifiable parameters satisfy three independent linear equations. The theorem lists a maximal set of 6 linearly independent parameters.  $\hfill \Box$ 

#### Genotype counts and MAF moments $\mathbf{B}$

By Eq. (6), a proper distribution  $\Delta'$  is generated in Proof of Theorem 5. Eq. (5) with  $\eta = 0$  and all  $\xi$  satisfying

$$-\min\{\Delta_2, \Delta_8/2\} \le \xi \le \min\{\Delta_4, \Delta_6, \Delta_7\}.$$

If  $\xi \neq 0$ , then  $\Delta' \neq \Delta$ , yet they generate the same joint genotype distributions  $\mathbf{f}^{(i)}$  with all minor-allele frequency sequences  $(p_i: i = 1, 2, ...)$ . 

Proof of Theorem 6. Define the expectations  $\bar{f}_x = \mathbb{E}\hat{f}_x$ . Since  $\hat{f}_x$  is an average of indicator random variables in (13),

$$\hat{f}_x \xrightarrow{\mathrm{P}} \bar{f}_x \tag{17}$$

as  $n \to \infty$ . (Hoeffding's inequality gives  $\mathbb{P}\left\{\left|\hat{f}_x - \bar{f}_x\right| \ge \epsilon\right\} \le 2\exp(-2n\epsilon^2)$ , thus  $\lim_{n \to \infty} \mathbb{P}\left\{ \left| \hat{f}_x - \bar{f}_x \right| \ge \epsilon \right\} = 0 \text{ for all } \epsilon > 0.$ By averaging (1) across the *n* sites,

$$\bar{\mathbf{f}} = \frac{\sum_{i=1}^{n} \mathbf{f}^{(i)}}{n} = \frac{\sum_{i=1}^{n} \mathbf{F}^{(i)}}{n} \cdot \mathbf{\Delta} = \bar{\mathbf{F}} \cdot \mathbf{\Delta}.$$

The entries of  $\bar{\mathbf{F}}$  link expected genotype frequencies  $\bar{f}$  to the common identity coefficients  $\Delta_i$ :  $i = 1, \ldots, 9$ , as follows in Eqs. (18a–18i).

$$\bar{f}_{0000} = \frac{\mathbb{E}n_{0000}}{n} = \frac{(1-\bar{\mu})\Delta_1 + (1-2\bar{\mu}+\bar{\mu}_2)(\Delta_2+\Delta_3+\Delta_5+\Delta_7)}{+(1-3\bar{\mu}+3\bar{\mu}_2-\bar{\mu}_3)(\Delta_8+\Delta_4+\Delta_6)} + (1-4\bar{\mu}+6\bar{\mu}_2-4\bar{\mu}_3+\bar{\mu}_4)\Delta_9$$
(18a)

$$\bar{f}_{1111} = \frac{\mathbb{E}n_{1111}}{n} = -\bar{\mu}\Delta_1 + \bar{\mu}_2(\Delta_2 + \Delta_3 + \Delta_5 + \Delta_7) + \bar{\mu}_3(\Delta_4 + \Delta_6 + \Delta_8) + \bar{\mu}_4\Delta_9$$
(18b)

$$\bar{f}_{1101} = \frac{\mathbb{E}n_{1101}}{n} = (\bar{\mu} - \bar{\mu}_2)\Delta_3 + (\bar{\mu}_2 - \bar{\mu}_3)(\Delta_8 + 2\Delta_4) + 2(\bar{\mu}_3 - \bar{\mu}_4)\Delta_9$$
(18c)

$$\bar{f}_{0111} = \frac{\mathbb{E}n_{0111}}{n} = (\bar{\mu} - \bar{\mu}_2)\Delta_5 + (\bar{\mu}_2 - \bar{\mu}_3)(\Delta_8 + 2\Delta_6) + 2(\bar{\mu}_3 - \bar{\mu}_4)\Delta_9$$
(18d)

$$\bar{f}_{0101} = \frac{\mathbb{E}n_{0101}}{n} = 2(\bar{\mu} - \bar{\mu}_2)\Delta_7 + (\bar{\mu} - 2\bar{\mu}_2 + \bar{\mu}_3)\Delta_8$$

$$+4(\bar{\mu}_2 - 2\bar{\mu}_3 + \bar{\mu}_4)\Delta_9$$
(18e)

$$\bar{f}_{1100} = \frac{\mathbb{E}n_{1100}}{n} = (\bar{\mu} - \bar{\mu}_2)\Delta_2 + (\bar{\mu} - 2\bar{\mu}_2 + \bar{\mu}_3)\Delta_4 + (\bar{\mu}_2 - \bar{\mu}_3)\Delta_6 \qquad (18f) + (\bar{\mu}_2 - 2\bar{\mu}_3 + \bar{\mu}_4)\Delta_9$$

$$\bar{f}_{0011} = \frac{\mathbb{E}n_{0011}}{n} = (\bar{\mu} - \bar{\mu}_2)\Delta_2 + (\bar{\mu} - 2\bar{\mu}_2 + \bar{\mu}_3)\Delta_6 + (\bar{\mu}_2 - \bar{\mu}_3)\Delta_4 \qquad (18g) + (\bar{\mu}_2 - 2\bar{\mu}_3 + \bar{\mu}_4)\Delta_9$$

$$\bar{f}_{0100} = \frac{\mathbb{E}n_{0100}}{n} = \frac{(\bar{\mu} - \bar{\mu}_2)\Delta_5 + (\bar{\mu} - 2\bar{\mu}_2 + \bar{\mu}_3)(\Delta_8 + 2\Delta_6)}{+2(\bar{\mu} - 3\bar{\mu}_2 + 3\bar{\mu}_3 - \bar{\mu}_4)\Delta_9}$$
(18h)

$$\bar{f}_{0001} = \frac{\mathbb{E}n_{0001}}{n} = (\bar{\mu} - \bar{\mu}_2)\Delta_3 + (\bar{\mu} - 2\bar{\mu}_2 + \bar{\mu}_3)(\Delta_8 + 2\Delta_4) + 2(\bar{\mu} - 3\bar{\mu}_2 + 3\bar{\mu}_3 - \bar{\mu}_4)\Delta_9.$$
(18i)

In a similar vein, define the expected sample statistics  $\bar{\tau}_y = \mathbb{E}\hat{\tau}_y$  for y = 1, 2A, 2B, 3, 4. It follows from (14) and (11) that each  $\bar{\tau}_y$  can be written as a linear combination of the expected genotype counts  $\bar{f}_x$ ; i.e., Equation (11) holds after substituting  $\tau_y \leftarrow \bar{\tau}_y$  and  $f_x \leftarrow \bar{f}_x$  appropriately. Consequently, (18) and (17) imply that

$$\hat{\tau}_{1} \xrightarrow{P} (\bar{\mu} - \bar{\mu}_{2})(1 - \theta_{1}) 
\hat{\tau}_{2A} \xrightarrow{P} (\bar{\mu} - \bar{\mu}_{2})(1 - \theta_{2A}) 
\hat{\tau}_{2B} \xrightarrow{P} (\bar{\mu} - \bar{\mu}_{2})(1 - \theta_{2B}) 
\hat{\tau}_{3} \xrightarrow{P} (\bar{\mu} - 3\bar{\mu}_{2} + 2\bar{\mu}_{3})(1 - \theta_{3}) 
\hat{\tau}_{4} \xrightarrow{P} (\bar{\mu} - 3\bar{\mu}_{2} + 2\bar{\mu}_{3})\theta_{4}.$$
(19)

Since

$$\bar{\mu} - \bar{\mu}_2 \xrightarrow{\mathrm{p}} \nu_2$$
 and  $(\bar{\mu} - 3\bar{\mu}_2 + 2\bar{\mu}_3) \xrightarrow{\mathrm{p}} \nu_3$ 

and the convergence in the denominators of  $\hat{\theta}_y$  is towards non-zero constants  $(\nu_2 \text{ or } \nu_3)$ , Eq. (19) implies  $\hat{\theta}_y \xrightarrow{\mathrm{P}} \theta_y$  for all  $y = 1, 2\mathrm{A}, 2\mathrm{B}, 3, 4$ .  $\Box$ 



Figure 3: Joint ancestry of Queen Victoria (V1819) and Prince Albert (A1819). Rectangles denote men; ovals denote women. Nodes are placed by the indicated time scale, according to birth year.

# C Family tree of Queen Victoria and Prince Albert

Queen Victoria of the United Kingdom (1819–1901) and her spouse Prince Albert of Saxe-Coburg and Gotha (1819–1861) descended from prominent ducal families of the Holy Roman Empire. Figure 3 shows their family tree covering 200 years, pruned back to founders. Table 2 lists the 49 members of the tree. Genealogical records were culled using peerage.com, and Wikipedia's English and German editions.

Table 2: Members of the family tree for Victoria and Albert. Identifiers are formed by initial and birth year; founders are marked by boldface.

Identifier	Lineage		Name	Title
	Victoria	Albert		
E1601	₽ď	ço™	Ernest I	Duke of Saxe-Gotha and Altenburg
E1619	₽ď	çơ	Elisabeth Sophie	of Saxe-Altenburg
A1633	Ŷ	çoĩ	Anthony Ulrich	Duke of Brunswick-Wolfenbttel
E1634	Ŷ	çơ	Elisabeth Juliane	of Schleswig-Holstein-Sønderborg-Nordborg
F1646	ਾ	Ŷ	Frederick I	Duke of Saxe-Gotha-Altenburg
M1647		Ŷ	Marie Hedwig	of Hesse-Darmstadt
M1648	ੋ	Ŷ	Magdalena Sibylle	of Saxe-Weissenfels
B1649		Ŷ	Bernhard I	Duke of Saxe-Meiningen
E1658		Ŷ	Elisabeth Eleonore	of Brunswick-Wolfenbüttel
J1658	Ŷ	çoĩ	John Ernest IV	Duke of Saxe-Coburg-Saalfeld
C1664	ę	çoĩ	Charlotte-Johanna	of Waldeck-Wildungen
C1671	Ŷ		Christine Louise	of Oettingen-Oettingen

Table 2. Family tree for Victoria and Albert

Identifier	Lineage		Name	Title
	Victoria	Albert		
L1671	Ŷ	്	Louis Rudolph	Duke of Brunswick-Lüneburg
E1672		Ŷ	Ernst Ludwig I	Duke of Saxe-Meiningen
D1674		Ŷ	Dorothea Marie	of Saxe-Gotha-Altenburg
F1676	്	Ŷ	Frederick II	Duke of Saxe-Gotha-Altenburg
M1679	്	Ŷ	Magdalene Augusta	of Anhalt-Zerbst
F1680	Ŷ	ď	Ferdinand-Albert II	Duke of Brunswick-Wolfenbüttel
A1687		Ŷ	Anton Ulrich	Duke of Saxe-Meiningen
A1696	Ŷ	ď	Antoinette	of Brunswick-Wolfenbüttel
F1697	Ŷ	₽ď	Francis Josias	Duke of Saxe-Coburg-Saalfeld
F1699		Ŷ	Frederick III	Duke of Saxe-Gotha-Altenburg
A1700	Ŷ	₽ď	Anna Sophie	of Schwarzburg-Rudolstadt
J1704		Ŷ	John August	of Saxe-Gotha-Altenburg
F1707	്		Frederick Louis	Prince of Wales
L1710		Ŷ	Luise Dorothea	of Saxe-Meiningen
A1719	്		Augusta	of Saxe-Gotha-Altenburg
E1724	Ŷ	്	Ernest Frederick	Duke of Saxe-Coburg-Saalfeld
S1724	Ŷ	്	Sophie Antoinette	of Brunswick-Wolfenbüttel
L1725		Ŷ	Louis	Duke of Mecklenburg-Schwerin
L1726		Ŷ	Louise	of Reuss-Schleiz
C1730		Ŷ	Charlotte Amalie	of Hesse-Phillipstal
C1731		Ŷ	Charlotte Sophie	of Saxe-Coburg-Saalfeld
G1738	്		George III	King of the United Kingdom
C1744	്		Charlotte	of Mecklenburg-Strelizt
E1745		Ŷ	Ernest II	Duke of Saxe-Gotha-Altenburg
F1750	Ŷ	ď	Francis	Duke of Saxe-Coburg-Saalfeld
C1751		Ŷ	Charlotte	of Saxe-Meiningen
F1756		Ŷ	Frederick Francis I	Grand-Duke of Mecklenburg-Schwerin
L1756		Ŷ	Louise	of Saxe-Gotha-Altenburg
A1757	Ŷ	്	Augusta	of Reuss-Ebersdorf
E1767	്		Edward	Duke of Kent and Strathearn
A1772		Ŷ	Augustus	Duke of Saxe-Gotha-Altenburg
L1779		Ŷ	Louise-Charlotte	of Mecklenburg-Schwerin
E1784		്	Ernest I	Duke of Saxe-Coburg and Gotha
V1786	Ŷ		Victoria	of Saxe-Coburg-Saalfeld
L1800		Ŷ	Louise	of Saxe-Gotha-Altenburg
A1819			Albert	of Saxe-Coburg and Gotha
V1819			Victoria	Queen of the United Kingdom