

# Comment on “Exact Analytic Second Virial Coefficient for the Lennard-Jones Fluid” (arXiv:0909.3326v1)

Isidro Cachadiña Gutiérrez \*

*Departamento de Física Aplicada  
Universidad de Extremadura  
Avda. de Elvas, s/n  
06006 Badajoz (Spain)*

January 31, 2020

### Abstract

The asymptotic expansion for  $T \rightarrow 0$  from Byung Chan Eu of  $B_2(T) = -16\sqrt{2\pi}v_0e^{\varepsilon\beta}(\varepsilon\beta)^{3/2}\left[1 + \frac{19}{16\varepsilon\beta} + \frac{105}{512(\varepsilon\beta)^2} \dots\right]$  is wrong. The correct expression is  $B_2(T) = -2\sqrt{2\pi}v_0e^{\varepsilon\beta}(\varepsilon\beta)^{-1/2}\left[1 + \frac{15}{16\varepsilon\beta} + \frac{945}{512(\varepsilon\beta)^2} + \dots\right]$ .

Byung Chan Eu gave an exact analytical solution for the second virial coefficient valid for the entire range of temperature which reads as:

$$-\frac{B_2(x)}{v_0\sqrt{2}x^{1/4}} = 4\Gamma\left(\frac{3}{4}\right)\left[6xM\left(\frac{7}{4}, \frac{3}{2}, x\right) - (1+4x)M\left(\frac{3}{4}, \frac{1}{2}, x\right)\right] + \left[2\Gamma\left(\frac{1}{4}\right)\sqrt{x}\left[\frac{10}{3}xM\left(\frac{9}{4}, \frac{5}{2}, x\right) + (1-4x)M\left(\frac{5}{4}, \frac{3}{2}, x\right)\right]\right]. \quad (1)$$

where  $x = \varepsilon\beta = \varepsilon/kT$ ,  $v_0 = \pi\sigma^3/6$ , and  $M(a, b, x)$ , is a confluent hypergeometric function of Kummer [2]:

$$M(a, b, x) = \sum_{n=0}^{\infty} \frac{(a)_n x^n}{(b)_n n!}, \quad (2)$$

where  $(a)_n$  is the Pochhammer symbol [2]

$$\begin{aligned} (a)_0 &= 1, \\ (a)_n &= a(a+1)(a+2)\dots(a+n-1) \quad (n \geq 1). \end{aligned} \quad (3)$$

The asymptotic behaviour for  $T \rightarrow 0$  can be determined using the following asymptotic expansion of  $M(a, b, x)$  for real  $x$  [2] when  $x \rightarrow \infty$ :

$$M(a, b, x) = \frac{\Gamma(b)}{\Gamma(a)} e^x x^{a-b} \left[ \sum_{n=0}^{m-1} \frac{(b-a)_n (1-a)_n}{n!} x^{-n} + O(x^{-m}) \right]. \quad (4)$$

---

\*Electronic address: [icacha@unex.es](mailto:icacha@unex.es)

Byung Chan Eu claimed that the following expression

$$B_2^{Chan}(T) = -16\sqrt{2\pi}v_0e^{\varepsilon\beta}(\varepsilon\beta)^{3/2}\left[1 + \frac{19}{16\varepsilon\beta} + \frac{105}{512(\varepsilon\beta)^2} + \dots\right] \quad (5)$$

is the asymptotic behaviour of the second virial coefficient for  $T \rightarrow 0$ .

We can see easily that this expansion is wrong by calculating

$$\left|\frac{B_2^{Chan}(x)}{B_2(x)} - 1\right|, \quad (6)$$

and seeing that the above expression doesn't go to zero for larger values of  $x$ . When this function is plotted (as it is done in Fig. 1) one can see that ratio between  $|B_2^{Chan}(x)/B_2(x) - 1|$  grows as  $x^2$  and therefore  $B_2^{Chan}(x)$  is not the asymptotic expansion of  $B_2(x)$ .




Figure 1: Absolute relative deviation of the asymptotic expansion proposed by B. Chan Eu for the second virial coefficient. This curve have to tend to zero for larger values of  $x$ , but on the contrary, it grows as  $x^2$ .

With the aid of the Mathematica program [3] the asymptotic expansion can be easily calculated (see Appendix 1) to give:

$$B_2(x) = -2\sqrt{2\pi}v_0e^{\varepsilon\beta}(\varepsilon\beta)^{-1/2}\left[1 + \frac{15}{16x} + \frac{945}{512x^2} + \frac{45045}{8192x^3} + \dots\right]. \quad (7)$$

The factor  $x^2$  that can be seen in Fig. 1 is due the ratio of  $x^{3/2}/x^{-1/2}$  between the expression here deduced and the one from Byung Chan Eu.

In order to check the convergence Eq. (7) I plotted in Fig. 2 this expansion taking different number of terms in the polynomial of  $x^{-1}$ . Thus, calling:

$$B_2(x, n) = -2\sqrt{2\pi}v_0e^{\varepsilon\beta}(\varepsilon\beta)^{-1/2}\sum_{i=0}^n \frac{a_i}{x^i} \quad (8)$$

where:

$$\begin{aligned}
 a_0 &= 1 \\
 a_1 &= 15/16 \\
 a_2 &= 945/512 \\
 a_3 &= 45045/8192 \\
 a_4 &= 11486475/524288 \\
 a_5 &= 916620705/8388608 \\
 a_6 &= 175685635125/268435456 \\
 a_7 &= 19651693186125/4294967296 \\
 a_8 &= 20103682129405875/549755813888
 \end{aligned} \tag{9}$$

we can check how many terms are needed to achieve the desired accuracy. For example, taking  $n = 6$  the expansion of Eq. (8) gives 6 exact digits when  $x > 25$ .




Figure 2: Absolute relative deviation of the asymptotic expansion,  $B_2(x, n)$  (Eq. (8)), as a function of the number of terms  $n$  included.

## Appendix

This is the Mathematica [3] notebook I used to calculate the asymptotic expansions.

```
In[1]:=
```

```
(* True value of  $-B2/v0$  *)
```

```
B2[x_] := Sqrt[2]*x^(1/4)*(4*Gamma[3/4]*(6*x*Hypergeometric1F1[7/4, 3/2, x] -
(1 + 4*x)*Hypergeometric1F1[3/4, 1/2, x]) + 2*Gamma[1/4]*
Sqrt[x]*(10/3*x*Hypergeometric1F1[9/4, 5/2, x] + (1 - 4*x)*
Hypergeometric1F1[5/4, 3/2, x]))
```

```
(* Check if  $TB^{-1} = 0.2925$  *)
```

```
In[2]:= FindRoot[B2[x] == 0, {x, 0.3}]
```

```
Out[2]= {x -> 0.292575}
```

In[3]:= (\* Asymptotic expansion of Krummer functions up to n \*)

M[a\_, b\_, x\_, n\_] :=  
Gamma[b]/Gamma[a]\*Exp[x]\*x^(a - b)\*  
Sum[Pochhammer[b - a, k]\*Pochhammer[1 - a, k]\*x^-k/k!, {k, 0, n}]

In[5]:= (\* The value of  $-B2/v0$  using the asymptotic expansion up to term n \*)

F[x\_, n\_] := Sqrt[2]\*x^(1/4)\*(4\*  
Gamma[3/4]\*(6\*x\*M[7/4, 3/2, x, n] - (1 + 4\*x)\*M[3/4, 1/2, x, n]) +  
2\*Gamma[1/4]\*Sqrt[x]\*(10/3\*x\*M[9/4, 5/2, x, n]  
+ (1 - 4\*x)\*M[5/4, 3/2, x, n]))

In[6]:= (\* The two first approximations of  $F[x, n]$  are zero \*)

FullSimplify[F[x, 0]]

Out[6]= 0

In[7]:= FullSimplify[F[x, 1]]

Out[7]= 0

In[8]:= Ap2[x\_] = FullSimplify[F[x, 2]]

Out[8]= (2 E^x Sqrt[2 \[Pi]])/Sqrt[x]

In[9]:= Apn[x\_, n\_] :=  
Expand[FullSimplify[F[x, n]/F[x, 2]]]\*FullSimplify[F[x, 2]]

In[10]:= Apn[x, 7]

Out[10]= (2 E^x Sqrt[  
2 \[Pi]] (1 + 916620705/(8388608 x^5) + 11486475/(524288 x^4) +  
45045/(8192 x^3) + 945/(512 x^2) + 15/(16 x)))/Sqrt[x]

In[11]:= CoefficientList[Expand[FullSimplify[F[x, 10]/F[x, 2]]], 1/x]

Out[11]= {1, 15/16, 945/512, 45045/8192, 11486475/524288, \  
916620705/8388608, 175685635125/268435456, 19651693186125/4294967296, \  
20103682129405875/549755813888}

## References

- [1] Byung Chan Eu, arXiv:0909.3326v1 (2009)
- [2] M. Abramowitz, I.A. Stegun, Handbook of Mathematical Functions, Dover Pub. Inc, Washington, D.C, 1964.
- [3] Wolfram Research, Inc., Mathematica, Version 8.0, Champaign, IL (2010).